Genome-wide association mapping of correlated traits in cassava: dry matter and total carotenoid content

Ismail Y. Rabbi[1]*, Lovina I. Udoh[1], Marnin Wolfe[2], Elizabeth Y. Parkes[1], Melaku A. Gedil[1], Alfred Dixon[1], Punna Ramu[3], Jean-Luc Jannink[2,4], Peter Kulakow[1].

1. International Institute of Tropical Agriculture (IITA), PMB 5320 Ibadan, Oyo, Nigeria
2. Department of Plant Breeding and Genetics, Cornell University, Ithaca, NY, USA
3. Institute of Genomic Diversity, Cornell University, Ithaca, NY, USA.
4. USDA-ARS, R.W. Holley Center for Agriculture and Health, Ithaca, NY, USA

* Corresponding author, email: I.Rabbi@cgiar.org

ABSTRACT

Cassava (*Manihot esculenta* (L.) Crantz) is a starchy root crop cultivated in the tropics for fresh consumption and commercial processing. Dry matter content and micronutrient density, particularly of provitamin A – traits that are negatively correlated – are among the primary selection objectives in cassava breeding. This study aimed at identifying genetic markers associated with these traits and uncovering the potential underlying cause of their negative correlation – whether linkage and/or pleiotropy. A genome-wide association mapping using 672 clones genotyped at 72,279 SNP loci was carried out. Root yellowness was used indirectly to assess variation in carotenoid content. Two major loci for root yellowness was identified on chromosome 1 at positions 24.1 and 30.5 Mbp. A single locus for dry matter content that co-located with the 24.1 Mbp peak for carotenoid content was identified. Haplotypes at these loci explained a large proportion of the phenotypic variability. Evidence of mega-base-scale linkage disequilibrium around the major loci of the two traits and detection of the major dry matter locus in independent analysis for the white- and yellow-root subpopulations suggests that physical linkage rather that pleiotropy is more likely to be the cause of the negative correlation between the target traits. Moreover, candidate genes for carotenoid (*phytoene synthase*) and starch biosynthesis (*UDP-glucose pyrophosphorylase* and *sucrose synthase*) occurred in the vicinity of the identified locus at 24.1 Mbp. These findings elucidate on the genetic architecture of carotenoids and dry matter in cassava and provides an opportunity to accelerate genetic improvement of these traits.

CORE IDEAS
- Cassava, a starchy root crop, is a major source of dietary calories in the tropics.
- Most varieties consumed are poor in micronutrients, including pro-vitamin A.
- These two traits are governed by few major loci on chromosome one.
- Genetic linkage, rather than pleiotropy, is the most likely cause of their negative correlation.

INTRODUCTION

Cassava (*Manihot esculenta* (L.) Crantz) is one of the most important food and feed crops in the tropics and Africa accounts for more than half of the total world-wide production of

270.3 million tonnes (http://faostat3.fao.org/, accessed 26.03.2016). Because of its
remarkable tolerance to drought (El-Sharkawy, 1993), its ability to grow in poor soils (Cock,
1982), and its perennial nature which allows it to be harvested as and when required, this
heterozygous and clonally propagated species plays a particularly important role in food
security for millions of small-holder farmers in developing countries. Moreover, cassava is
increasingly being cultivated for commercial processing to convert its storage roots into
dehydrated chips, flour and starch (Balagopalan, 2002). Dry matter content, of which a large
proportion is starch, is therefore a primary factor that defines adoption of new cassava
varieties by farmers and the market value of harvested roots (Okechukwu and Dixon, 2008).
As a result, breeding of improved varieties with high dry matter content is one of the
primary objectives of cassava genetic improvement programs in the world.
58
Another important target trait for cassava improvement in developing countries is
biofortification for micronutrients (Pfeiffer and McClafferty, 2007; Saltzman et al., 2013).
Most varieties grown and consumed throughout the world have white storage roots with
negligible amounts of micronutrients in general, and provitamin A in particular (Welsch et
al., 2010). Dietary diversification and breeding of farmer-preferred improved varieties with
higher nutritional density are complementary approaches used in addressing potential
micronutrient deficiency associated with consumption of cassava as the sole staple food
(Sayre et al., 2011). The crop's gene-pool exhibits considerable natural variation for storage
root carotenoids that can be tapped for breeding of biofortified varieties, with some
breeding populations reported to accumulate as much as 25.8 µg/g fresh root weight
(Ceballos et al., 2013; Sánchez et al., 2014).
70
Despite availability of natural genetic diversity in the global germplasm that is relevant to
breeding for increased dry matter and total carotenoid contents, improving these traits
through phenotype-based recurrent selections is a lengthy process, due to the breeding
complexities associated with the species including an annual cropping cycle of 12 to 24
months and low multiplication rate of planting materials. Understanding the genetic basis of
variation in these traits is essential for increasing their selection efficiency and the rate of
genetic gain. More importantly, several studies using diverse germplasm have reported that
dry matter and carotenoid content are negatively correlated, with r values ranging from -0.1
to -0.5 (Marín Colorado et al., 2009; Akinwale et al., 2010; Esuma et al., 2012; Ceballos et al.,
2013; Njoku et al., 2015). Despite its significant implication in breeding, the genetic basis of
this correlation – whether it is due to genetic linkage or pleiotropy – is not understood.
82
Several mapping studies using either Bulk Segregant Analysis (BSA) or Quantitative Trait Loci
(QTL) mapping of S1 or F1 populations have been reported separately for dry matter and
carotenoid content (Balyejusa Kizito et al., 2007; Marín Colorado et al., 2009; Welsch et al.,
2010; Morillo C et al., 2013; Njoku et al., 2014). The mapping resolution from single-cross
experimental populations is expected to be limited due to the use of sparse genetic maps
and the limited number of recombination events observed (Hamblin et al., 2011). Moreover,
QTLs from such bi-parental populations may not provide insight into the tremendous
genetic and phenotypic variation of the larger gene pool (Zhao et al., 2011). The increased
availability of genomic resources for cassava, including the chromosome-scale reference
genome and integrated linkage map (Prochnik et al., 2012; International Cassava Genetic
Map Consortium (ICGMC), 2014) and high-density genotyping using next-generation

sequencing (Rabbi et al., 2014a; b) makes it possible to use genome-wide association
(GWAS) mapping to dissect the phenotypic diversity of cassava germplasm with respect to
dry matter and carotenoid content. GWAS, which takes advantage of natural linkage
disequilibrium (LD) generated by ancestral mutation, drift, and recombination events in
diverse germplasm, offers the possibility to overcome the shortcomings of traditional bi-
parental QTL mapping. These advantages mean GWAS is able to reveal a broader spectrum
of trait-linked allelic variation and thus may provide the most useful markers for marker-
assisted selection (MAS). Indeed, GWAS has already been applied in other crops such as
maize to study the genetic architecture of carotenoid accumulation (Harjes et al., 2008;
Owens et al., 2014; Suwarno et al., 2015). In cassava, Esuma et al., (2016) carried out a
GWAS study using a panel of partial inbreds (S1 and S2 generation) produced from eight
clones. Using this limited number of parents, they reported a single genomic region on
Chromosome 1 that underlies the variation in total carotenoid content. However, no joint
association analysis examining carotenoids and dry matter content has hitherto been
reported. Here, we present the results of a GWAS using a collection of more than 650
cassava clones representing diverse African germplasm genotyped at high-density using
genotyping-by-sequencing (Elshire et al., 2011). The population was phenotyped in two
locations for three consecutive field seasons. The results of this study will be used to
develop efficient strategies to breed for high dry matter and provitamin A content varieties.
113
114 **METHODS**
115
Germplasm: The present work was carried out using the Tropical Manihot Selection (TMS)
cultivars developed at the International Institute of Tropical Agriculture (IITA) in Nigeria. This
population, also known as the Genetic Gain collection, consist of more than 650 advanced
breeding lines and key landraces selected over four decades from 1970 (Okechukwu and
Dixon, 2008; Ly et al., 2013). The pedigree of the collection is mainly composed of crosses
between germplasm from West Africa and early introductions of CMD-tolerant lines arising
from interspecific hybridization between *Manihot glaziovii* and cultivated cassava at the
Amani station in Tanzania (Hahn et al., 1980). The collection also includes hybrid germplasm
from Latin America (Wolfe et al., 2016).
125
Locations and experimental design: The Genetic Gain population was planted using an
incomplete block design with two checks per block and single row of either 5 or 10 plants
spaced at $1m^2$. Data used for this study was collected in the 2012-2013, 2013-2014, and
2014-2015 field seasons in Ibadan (7.40° N, 3.90° E) and Ubiaja (6.66° N, 6.38° E). The trials
are usually planted in June, at the onset of the raining season in South West Nigeria and
harvested in June of the following year.
132
Assessment of dry matter content and yellow color intensity of storage roots: Dry matter
content was assessed using the oven-drying method. Eight fully developed roots were
randomly selected from each plot, peeled, chipped and thoroughly mixed. For each sample,
100g was weighed and oven-dried for 48 hours at 104°C till constant weight was achieved.
The samples were then re-weighed and the dry matter content was expressed as the
percentage of dry weight relative to fresh weight.
139
Because of the well-established linear relationship between intensity of yellow color and

carotenoid content in cassava storage roots (Pearson's coefficient, r, ranges from 0.81 to
0.89), we used root yellowness as an indirect measure of carotenoid content (Iglesias et al.,
1997; Chávez et al., 2005; Marín Colorado et al., 2009; Akinwale et al., 2010; Sánchez et al.,
2014). The relative difference among clones in the Genetic Gain population was assessed
using two complementary methods. The first was a visual gradation of yellow color using a
standard color-chart starting from one (white) to 7 (deep yellow). Due to the potential
subjectivity inherent in visual color scores, we complemented the color-chart method
through the use of a Minolta CR-410® chromameter. Approximately 100g of grated samples
from freshly peeled roots were placed in transparent Nasco Whirl-Pak® sampling bags and
four chromameter measurements taken in different sections of the bag. We chose the
commission internationale de l'éclairage (CIELAB) method that records color values in a
three-dimensional color space, where the L* coordinate corresponds to a lightness
coordinate, and the a* coordinate corresponds either to red (positive values) or to green
(negative values). Of importance to this study was the b* coordinate, whose positive values
represents yellow while the negative values represent blue. The illuminant used was D65
and calibration was done each day with a white ceramic.

SNP genotyping: DNA was extracted as described in Rabbi et al. (2014) and Genotyping-by-
sequencing was carried out as described by (Elshire et al., 2011). DNAs from the Genetic
Gain individuals were digested individually with *ApeK*I, a methylation sensitive restriction
enzyme that recognizes a five base-pair sequence (GCWGC, where W is either A or T). The
GBS sequencing libraries consisting of 95-plex DNA samples each were prepared by ligating
the digested DNA to unique sample identifier barcodes (nucleotide adapters) followed by
standard PCR. Sequencing was performed using Illumina HiSeq2500. The sequenced reads
from different genotypes were de-convoluted using their unique barcodes and aligned to
version 6.0 of the cassava reference genome (www.phytozome.org/cassava) with the
Bowtie 2 (Langmead and Salzberg, 2012). SNPs were discovered using the GBS pipeline
Version 2 implemented in TASSEL software (Glaubitz et al., 2014) and converted to dosage
format (0 = homozygous reference, 1 = heterozygous, 2 = homozygous non-reference
alleles). Missing data were filtered as described in (Wolfe et al., 2016) and imputed with the
glmnet algorithm in R (http://cran.r-project.org/web/packages/glmnet/index.html) (Wong
et al., 2014).

Phenotypic data analysis:
The phenotypic data across two locations and three years was collapsed to single best linear
unbiased predictor (BLUP) values for each clone by fitting the following mixed linear model
with the *lme4* package in R:
$y_{lij} = \mu + c_l + \beta_i + c_l * \beta_i + \varepsilon_{lij}$
Here, y_{lij} represents raw phenotypic observations, μ is the grand mean, c_l is a random
effects term for clone with $c_l \sim N(0, \sigma_I^2)$, β_i is a fixed effect for the combination of location
and year harvested, $C_l * \beta_i$ is a random effect for genotype-by-environment variance, and
ε_{lij} is the residual variance, assumed to be random and distributed $N(0, \sigma_e^2)$. Broad-sense
heritability for dry matter content and yellow color intensity was calculated according to (Ly
et al., 2013). Genetic correlation among traits was also calculated from BLUP values.

Population structure and Genome-Wide Association Analyses: Inherent population
structure and cryptic relatedness can lead to spurious associations in GWAS (Astle and

Balding, 2009). To control for these confounding factors, three standard GWAS models were
compared: a simple one-way ANOVA model with no correction (naïve model); a general
linear model (GLM) with the first five PCs of the SNP matrix as covariates (GLM + 5PCs); and
a mixed-linear model (MLM) using the five PCs and marker-estimated kinship matrix (Yu et
al., 2006). The models correcting for kinship and 5 PCs had the lowest inflation-factors as
determined from quartile-quartile (QQ) plots and therefore the lowest false-discovery rate
(**Supplementary Figure 1**). The association analyses were implemented in TASSEL (Bradbury
et al., 2007; Zhang et al., 2010). Association test P-values were considered significant when
more extreme than the Bonferroni threshold (with experiment-wise type I error rate of
0.05).
198
199 The patterns and extent of linkage disequilibrium (LD) in a population not only determines
200 the obtainable resolution in association mapping studies (Hamblin et al., 2011) but also has
201 strong implication in the interpretation of association peaks. Therefore the level of LD decay
202 and the local patterns of LD along each chromosome were determined by calculating intra-
203 chromosomal pairwise squared correlation (r^2) using PLINK (Purcell et al., 2007).
204
205 **RESULTS**
206 **SNP genotyping:** A total of 72,279 genome-wide SNP markers were called for the 672
207 genetic gain individuals after filtering for minor allele frequency threshold of 0.005. The
208 high-density coverage of SNPs resulted in an average of 4015 markers per chromosome,
209 ranging from 3101 on chromosome 16 to 5880 on chromosome 1.
210
Phenotypic variability: We investigated the phenotypic variation in dry matter content as
well as carotenoid-based intensity of yellow root color using a visual color chart and
chromameter. The dry matter content varied widely in the Genetic Gain population ranging
from 8.4% to 45% (average 28.6%, Table 1). About two-thirds of the evaluated clones have
white storage roots while the remaining showed a range of yellow color suggesting varying
levels of carotenoid content. On average, the visual score was 1.7 and ranged from 1 (white)
to 7 (yellow). The average chromameter measure of yellow color intensity (b* value) was
20.8 and ranged from 11.1 (white) to 40.8 (yellow). Dry matter was approximately normally
distributed while chromameter b* values showed a bimodal distribution (**Figure
1**) in which the first peak (b* values from 10 to 20) is associated with the white clones while
the second peak (b* values from 20 to 40) is associated with the variations among the
yellow clones.
Broad-sense heritability was high for root yellow color (H^2 = 0.87 and 0.82 for color chart
and chromameter b* values, respectively) but moderate for dry matter content (H^2 = 0.51).
These values are within the range of heritabilities reported previously for these traits
(Balyejusa Kizito et al., 2007; Ceballos et al., 2013). The relative importance of genotype-by-
environment variance (V_{GxE}) compared to genotype (V_G) variance was measured by the ratio
V_{GxE}/V_G. For all traits, the genetic variance component was larger than the genotype-by-
environment interaction variance. The interaction is minimal for the yellow color
measurements (0.054 and 0.173 for color chart and chromameter b* values, respectively).
For dry matter content, we observed a slightly higher interaction ratio of 0.214.
232
The BLUPs for dry matter content and gradation of yellow color were negatively correlated
in our germplasm collection (Pearson's correlation coefficient, r = -0.59; P-value < 0.0001),

indicating that clones with higher carotenoid content are more likely to have low dry matter
content (**Figure 2**) which confirms previous findings in cassava (Marín Colorado et al., 2009;
Akinwale et al., 2010; Esuma et al., 2012; Njoku et al., 2015). On the other hand, we found a
positive association between dry matter content and color lightness (chromameter L* value,
r = 0.60, P-value < 0.0001). The two measures of yellow color (i.e. color chart and
chromameter b* axis) were strongly correlated (*r* = 0.96, P-value < 0.0001).
241
242 **Population structure:** Analysis of population structure in 672 accessions genotyped across
243 72,279 SNPs using PCA detected subtle genetic differentiation in the genetic gain collection,
244 with the first 10 PCs explaining about 23 % of the genetic variation. The first two principal
245 components, which accounted for 8% of the genetic variation, revealed genetic
246 differentiation between white and yellow-root clones (**Figure 3**).
247
248 **Linkage disequilibrium:** Several regions of extensive mega-base-scale LD were discovered in
249 chromosomes 1, 4 and 10 as well as smaller regions in other chromosomes (**Figure 4**).
250 Excluding results from chromosomes with large LD blocks (i.e. chromosomes 1, 4 and 10),
251 we found that on average, LD drops almost to background levels (r^2 < 0.1) at around 2 Mb in
252 the Genetic Gain population (**Figure 5**).
253
Population-wide GWAS:
Variation in carotenoid content estimated by root yellowness: The MLM-based GWAS
analysis for yellow color in the storage root parenchyma using both the color chart and the
chromameter-based methods uncovered the same major association regions occurring at
24.1 and 30.5 Mbp of chromosome 1 (**Figure 6**). This is not surprising given the high
correlation between the two color assessment methods. The first major peak is tagged by
marker S1_24121306 for visual gradation of color ($-\log_{10}$(p-value) of 21.8) and marker
S1_24159585 ($-\log_{10}$(p-value) of 18.8) for chromameter b* value (Table 2). The second peak
was tagged by the same marker, S1_30543382, for both measures of color intensity and
occurred 6.5 Mbp away from the first peak ($-\log_{10}$(p-value) of 10.54 and 10.78, for color
chart and chromameter, respectively). All other SNPs between the two major regions were
not significant at the Bonferroni significance threshold (P = 6.92e-07) (**Figure 7**). The LD
between SNPs S1_24121306 and S1_30543382 was 0.3, suggesting moderate non-random
segregation of alleles of the two markers in these regions.
268
Dry matter content: Genetic variation in dry matter content was found to be associated
with a major locus occurring at 24.1 Mbp region of chromosome 1 and tagged by marker
S1_24121306 ($-\log10$(p-value) of 11.73). Importantly, this locus for dry matter content co-
locates with one of the two peaks found to be associated with carotenoid content (**Figure
7**). The genomic co-location of the major loci for dry matter content and root yellowness
suggests either a strong physical linkage between the genes underlying these important
traits or a pleiotropic effect. Distinguishing between these two possible causes is important
in cassava improvement efforts that target both traits. We therefore attempted to unravel
the genetic cause of the observed association by: (i) exploring the underlying linkage
disequilibrium patterns in the QTL regions on chromosome 1; (ii) carrying out independent
association analysis for dry matter content within the white root and yellow root
subpopulations; and (iii) searching for plausible biological explanation by identifying
candidate genes for both traits in the target region.

282
Exploration of the LD landscape along Chromosome 1 uncovered a mega-base-scale region
of low recombination extending from 22Mb to 33 Mb surrounding the association peaks for
dry matter content and yellow color (**Figure 7**). This region was recently shown to coincide
with a large *Manihot glaziovii* introgression segment (Bredeson et al., 2016) that traces back
to early breeding for resistance to cassava mosaic and cassava brown streak viruses in the
1930's (Hahn et al., 1980). Clustering of the Genetic Gain population based on identity-by-
descent relationship (i.e. a measure of how many alleles at any marker in each of the two
samples came from the same ancestral chromosomes) calculated using only markers from
this extensive LD region (2150 SNPs from markers S1_21567540 to S1_34950326) revealed
at least two major groups of accessions (**Supplementary Figure 2**), indicating presence of
few major haplotypes associated with the LD blocks.
294
GWAS for dry matter content in white root and yellow root subpopulation: If the
phenotypic association between dry matter and carotenoid contents and the colocation of
their association signals (~ 24.1 Mbp region) is largely caused by physical linkage rather than
by pleiotropy, the major dry matter locus should be detectable in both white root and
yellow root germplasm when analyzed independently. We therefore split the Genetic Gain
dataset into white root (n=210) and yellow root (n=427) subpopulations and repeated the
GWAS analysis. Clones that were at the borderline between yellow-root and white-root
were excluded from these analyses. To mitigate the loss of power as a result of double-
fitting markers in the MLM model both as a fixed effect tested for association and as a
random effect as part of the kinship (Lippert et al., 2011; Listgarten et al., 2012), the MLM
analysis was carried out using a kinship matrix calculated excluding markers from
chromosome 1.
307
We recovered the major dry matter content association signal in both the white root and
yellow-root subpopulation (**Figure 8**). Though coinciding with the locus identified in the
population-wide GWAS, the association signal in the white subpopulation was much
broader, extending from 24 to 33 Mbp and generally overlaps with the broad LD region of
the chromosome 1 (**Figure 8**). On the contrary, association signal for the yellow
subpopulation was relatively narrow. Survey of the underlying LD pattern in the same
chromosome region for the yellow subpopulation showed a recombination spot.
315
Selection sweep associated with breeding for yellow-root varieties: To determine whether
the breeding for carotenoid content trait in the Genetic Gain germplasm resulted in a
selection sweep around the major QTL region, we quantified genome-wide nucleotide
variation in the yellow root subpopulation (n = 210) and the non-yellow subpopulation (n =
427). A sliding-window scan of expected heterozygosity (π) and Tajima's D detected a ~ 6
Mb region with decrease in nucleotide diversity in the yellow compared to white-root
subpopulation around the first major carotenoid locus site (~ 24.1 Mbp) relative to its
chromosomal neighborhood (**Figure 9**).
324
Proportion of variance explained by markers QTL haplotypes.
To determine predictive ability of the discovered loci for yellow color intensity and dry
matter content, we carried out a multiple linear regression analysis using the *lm* function in
R and considered the top markers for these traits as independent variables and the traits

measurements as the response variables. A model considering the two major peaks
associated with gradation of yellow color as assessed using a color chart (S1_24121306 and
S1_30543382) returned an adjusted squared correlation (R^2) of 0.81. For the measure of
continuous variation in intensity of yellow color using chromameter (b* value), the adjusted
R^2 from same genomic regions (S1_24159585 and S1_30543382) was 0.70 while that for dry
matter content was moderate ($R^2 = 0.37$). This finding suggests that the major loci on
Chromosome 1 would be useful in Marker Assisted Selection breeding in cassava. Single or
joint allelic substitution effects at the associated loci with respect to chromameter b* value,
color chart and dry matter content is shown in **Figure 10**.
338
Candidate genes: The first of the two genomic regions associated color intensity (tagged by
SNP S1_24159585) was found ~ 4.5 Kbp away from phytoene synthase 2 (*PSY*2) in the
cassava version 6 reference genome. The *PSY*2 enzyme, named Manes.01G124200 and
located at 24,155,070 bp, is involved in the first dedicated step of the carotenoid
biosynthesis pathway in cassava roots which converts geranylgeranyl diphosphate to
phytoene (Welsch et al., 2010). Presence of the null versus the functional *PSY*2 allele is
responsible for the qualitative color difference between the white and the yellow roots,
respectively (Welsch et al., 2010; Rabbi et al., 2014a). Our study suggests that allelic
variation associated with increases in enzyme activity could contribute to deeper yellow by
increasing the flux into the pathway. No known candidate genes were found in the vicinity
of the second significant association signal on chromosome 1 occurring at 30.5 Mbp.
350
For dry matter content, we found two particular genes that are pivotal in central carbon
metabolism in the vicinity of top SNP linked to that trait. The first is UDP-glucose
pyrophosphorylase (named Manes.01G123000 in the cassava reference genome). This gene
which occurs at 24.06 Mbp region, plays a key role in carbohydrate metabolism, and is
strongly associated with the yield production both in grains and root crops (Smith, 2008;
Zeeman et al., 2010). UDP-glucose pyrophosphorylase was recently found to be up-
regulated during bulking of cassava storage roots (Yang et al., 2011; Wang et al., 2016). The
second key carbohydrate metabolism gene was sucrose synthase (named
Manes.01G123800), which occurred in 24.14 Mbp region. Finding of these potential
candidate genes for carotenoid and carbohydrate biosynthesis strongly favors the possibility
that the association between these two traits is caused by physical linkage rather than
pleiotropy. This hypothesis warrants further investigation.
363
DISCUSSION AND CONCLUSION
The present study revealed that the genetic architecture for dry matter content and
intensity of yellow color resulting from carotenoid accumulation in cassava roots is
governed by few major loci on chromosome 1 and explains the large repeatability estimates,
particularly for yellow color. These findings expand on those from previous genetic mapping
efforts for dry matter and carotenoid content. Using a candidate gene mapping approach,
Welsch et al. (2010) reported that a SNP mutation in the *PSY2* gene, leading to amino-acid
substitution, differentiates white and yellow cassava storage roots. Similarly, a bi-parental
QTL mapping study that used two clones from the Genetic Gain collection (TMS-I961089A
and TMEB117) also uncovered a single QTL peak whose confidence interval encompassed
the same PSY2 gene (Rabbi et al., 2014a). The F1 progenies from the TMS-I961089A x
TMEB117 population, also genotyped using the GBS method, segregated at an

approximately 1:1 ratio for white versus light-yellow roots, suggesting that the yellow-root
parent was heterozygous for the functional allele at the PSY2 locus. Kizito et al. (2007)
reported a QTL for dry matter content in a bi-parental population genotyped using SSR
markers that also corresponds to this region on chromosome 1. More recently, Esuma et
al., (2016) reported a single genomic region on Chromosome 1 underlies the variation in
total carotenoid content in eight S1 and S2 partially inbred families. This peak, around 24.66
Mbp, is close to our first locus tagged by SNP S1_24121306. However that study did not look
at genetic architecture for dry matter content.
384
While the amount of total carotenoids in the Genetic Gain collection was not directly
estimated, previous studies of diverse cassava germplasm have consistently reported a
strong linear relationship between yellow color and carotenoid content (Pearson's
coefficient, r, ranging from 0.81 to 0.89) (Iglesias et al., 1997; Chávez et al., 2005; Marín
Colorado et al., 2009; Akinwale et al., 2010; Sánchez et al., 2014). Hence the results
obtained here should be useful for breeding efforts targeting breeding for improved
carotenoid content. Nevertheless, we propose to quantify total carotenoids and its
constituents as a future study to corroborate the current findings.
393
Given the importance of dry matter content in cassava, and the fact that we found a single
genomic region associated with this trait, further studies are warranted to fine-map and
validate the identity of the causal locus. To do this effectively would require different
populations that are lacking the wild introgression segments in chromosome 1. This will lead
to reduced LD and allow higher mapping resolution. Additionally, special crosses such as
nested-association mapping population design (Yu et al., 2008) using strategically selected
sets of parents will reduce the confounding effect of population structure. Given our marker
density and sample size, this study is sufficiently powered to find large effect alleles that are
common in the studied germplasm. To detect more QTLs of small effects will require a
larger association panel genotyped at higher density.
404
The use of a broad cassava diversity panel in GWAS not only provides the foundation to map
genomic regions associated with natural variation in dry matter and carotenoid content but
also allows us to unravel the genetic cause of the negative correlation between these traits,
that is, pleiotropy versus genetic linkage. In the context of breeding to simultaneously
increase carotenoid and dry matter content, the observed negative association between
these traits in our germplasm is undesirable. Several lines of investigation pointed to a
possibility of genetic linkage rather than pleiotropy to be the cause of the observed
association. Firstly, the genomic region harboring the QTLs for yellow color and dry matter
content was found to occur in chromosomal segments that exhibits low overall
recombination in this region compared to the genome-wide patterns. Recent work by
Bredeson et al. (2016) has shown that this chromosome 1 region harbors a large *M. glaziovii*
introgression that commonly occurs in the Genetic Gain collection. Secondly, independent
association analysis for dry matter content on the white and the yellow subpopulations
detected the same association signal although the QTL in the white subpopulation was
broader suggesting that the favorable alleles were located in non-recombining haplotype.
Thirdly, strong candidate genes for dry matter (UDP-glucose pyrophosphorylase and
sucrose synthase) and carotenoid content (phytoene synthase) were found in the vicinity
of the major association region (24.1 Mbp) of chromosome 1. Presence of these genes hints

at possibly distinct biological causes of the observed associations with the two traits. These
hypotheses need to be tested through functional genetics studies at these candidate genes.
Taken together, these findings suggest that the phenotypic correlation between dry matter
and carotenoid content is mainly caused by physical linkage of loci underlying these traits.
Moreover, Ortiz et al. (2011) found a fairly large positive correlation (r = 0.62) between
these traits. It is therefore possible that the nature of association (whether positive or
negative) is dependent on the allelic status at the linked dry matter and carotenoid
biosynthesis genes. We also detected a reduction of expected heterozygosity (π) around the
major gene region in the yellow versus white sub population. This suggests that the genetic
base for sources of favorable alleles with respect to carotenoid biosynthesis at this locus is
narrow, possibly arising from a single haplotype, which could be linked in *cis* to low-dry
matter alleles in the dry matter locus. Alternatively, balancing selection of the *M. glaziovii*
introgression in the white cassava sub population might be the cause of the higher levels of
heterozygosity relative to the yellow sub population.
437
Although cassava is a predominantly outcrossing species, its clonally propagated nature
means that modern varieties have undergone relatively few recombination cycles compared
to seed crops. Most accessions in the Genetic Gain collection are not far removed from
founder clones. Accordingly, the extent of LD in this study (~ 2 Mbp) is much greater than
the LD in maize (< 10 Kb) (Yan et al., 2009) as well as in grape (< 10 Kb) (Myles et al., 2011),
another clonal species. Moreover, the overall recombination pattern is far from
homogeneous owing to the persistent introgressions of *M. glaziovii* chromosomal segments
that are legacies of the historical breeding program in East Africa (Hahn et al., 1980;
Jennings, 1994). From these results, it is expected that the mapping resolution will vary
widely across the cassava genome depending mainly on whether a locus-of-interest occurs
in or outside the large-LD blocks.
449
This study presents a significant progress toward dissecting the genetic architecture of two
key breeding goal traits in cassava. The major loci associated with carotenoid content
variation and a single locus associated with dry matter content represents markers that will
be useful for marker-assisted selection in these traits. Although the results of the present
study suggests genetic linkage is more likely to be responsible for the negative correlation
between the studied traits, there is need for further investigations to confirm or reject this
hypothesis. For example, will dry matter content be increased by knocking out the PSY2
gene using gene-silencing methods (Lu et al., 2003; Burch-Smith et al., 2004; Fofana et al.,
2004)?. Alternately, could the activation of PSY2 in clones with high dry matter content and
lacking in carotenoids using gene-editing technologies like CRISPR-CAS9 (Hsu et al., 2014;
Sander and Joung, 2014) lead to not only carotenoid production and accumulation but also
lowering of dry matter content?
462

Acknowledgements:
Oluwafemi Alaba and Ruth Uwugiaren for DNA processing; Staff of the Cassava Breeding
Unit of IITA for conducting field trials; Sharon Mitchell, Charlotte Acharya of Cornell's
Genomic Diversity Facility; Lukas Mueller and Guillaume Bauchet for SNP processing and
imputation. This work was supported through HarvestPlus project, The CGIAR Research
Programme on Roots, Tubers, and Bananas (CRP-RTB) and The Next Generation Cassava
Breeding grant OPP1048542 from Bill and Melinda Gates Foundation and the United

Kingdom Department for International Development.

References

Akinwale, M.G.. b, R.D.. Aladesanwa, B.O.. Akinyele, A.G.O.. Dixon, and A.C.. Odiyi. 2010. Inheritance of β-carotene in cassava (Manihot esculenta crantza). Int. J. Genet. Mol. Biol. 2(10): 198–201.

Astle, W., and D. Balding. 2009. Population Structure and Cryptic Relatedness in Genetic Association Studies. Stat. Sci. 24(4): 451–471.

Balagopalan, C. 2002. Cassava: biology, production and utilization (RJ Hillocks and JM Thresh, Eds.). CABI, Wallingford.

Balyejusa Kizito, E., A.-C. Rönnberg-Wästljung, T. Egwang, U. Gullberg, M. Fregene, and A. Westerbergh. 2007. Quantitative trait loci controlling cyanogenic glucoside and dry matter content in cassava (Manihot esculenta Crantz) roots. Hereditas 144(4): 129–136.

Bradbury, P.J., Z. Zhang, D.E. Kroon, T.M. Casstevens, Y. Ramdoss, and E.S. Buckler. 2007. TASSEL: software for association mapping of complex traits in diverse samples. Bioinformatics 23(19): 2633–2635.

Bredeson, J. V, J.B. Lyons, S.E. Prochnik, G.A. Wu, C.M. Ha, E. Edsinger-Gonzales, J. Grimwood, J. Schmutz, I.Y. Rabbi, C. Egesi, P. Nauluvula, V. Lebot, J. Ndunguru, G. Mkamilo, R.S. Bart, T.L. Setter, R.M. Gleadow, P. Kulakow, M.E. Ferguson, S. Rounsley, and D.S. Rokhsar. 2016. Sequencing wild and cultivated cassava and related species reveals extensive interspecific hybridization and genetic diversity. Nat. Biotechnol. 34(5): 562–570.

Burch-Smith, T.M., J.C. Anderson, G.B. Martin, and S.P. Dinesh-Kumar. 2004. Applications and advantages of virus-induced gene silencing for gene function studies in plants. Plant J. 39(5): 734–746.

Ceballos, H., N. Morante, T. Sánchez, D. Ortiz, I. Aragón, A.L. Chávez, M. Pizarro, F. Calle, and D. Dufour. 2013. Rapid Cycling Recurrent Selection for Increased Carotenoids Content in Cassava Roots. Crop Sci. 53(6): 2342.

Chávez, a. L., T. Sánchez, G. Jaramillo, J.M. Bedoya, J. Echeverry, E. a. Bolaños, H. Ceballos, and C. a. Iglesias. 2005. Variation of quality traits in cassava roots evaluated in landraces and improved clones. Euphytica 143(1–2): 125–133.

Cock, J.H. 1982. Cassava: a basic energy source in the tropics. Science 218(4574): 755–762.

El-Sharkawy, M.A. 1993. Drought-tolerant cassava for Africa, Asia, and Latin America. Bioscience 43(7): 441–451.

Elshire, R.J., J.C. Glaubitz, Q. Sun, J. a Poland, K. Kawamoto, E.S. Buckler, and S.E. Mitchell. 2011. A robust, simple genotyping-by-sequencing (GBS) approach for high diversity species. PLoS One 6(5): e19379.

Esuma, W., L. Herselman, M.T. Labuschagne, P. Ramu, F. Lu, Y. Baguma, E.S. Buckler, and R.S. Kawuki. 2016. Genome-wide association mapping of provitamin A carotenoid content in cassava. Euphytica.

Esuma, W., P. Rubaihayo, A. Pariyo, R. Kawuki, B. Wanjala, I. Nzuki, J.J. Harvey, and Y. Baguma. 2012. Genetic Diversity of Provitamin A Cassava in Uganda. J. Plant Stud. 1(1): 60–71.

Fofana, I.B.F., A. Sangaré, R. Collier, C. Taylor, and C.M. Fauquet. 2004. A geminivirus-induced gene silencing system for gene function validation in cassava. Plant Mol. Biol. 56(4): 613–24.

Glaubitz, J.C., T.M. Casstevens, F. Lu, J. Harriman, R.J. Elshire, Q. Sun, and E.S. Buckler. 2014.
TASSEL-GBS: A high capacity genotyping by sequencing analysis pipeline. PLoS One 9(2).
Hahn, S.K., E.R. Terry, and K. Leuschner. 1980. Breeding cassava for resistance to cassava
mosaic disease. Euphytica 29(3): 673–683.
Hamblin, M.T., E.S. Buckler, and J.-L. Jannink. 2011. Population genetics of genomics-based
crop improvement methods. Trends Genet. 27(3): 98–106.
Harjes, C.E., T.R. Rocheford, L. Bai, T.P. Brutnell, C.B. Kandianis, S.G. Sowinski, A.E. Stapleton,
R. Vallabhaneni, M. Williams, E.T. Wurtzel, J. Yan, and E.S. Buckler. 2008. Natural
genetic variation in lycopene epsilon cyclase tapped for maize biofortification. Science
319: 330–333.
Hsu, P.D., E.S. Lander, and F. Zhang. 2014. Development and applications of CRISPR-Cas9 for
genome engineering. Cell 157(6): 1262–1278.
Iglesias, C., J. Mayer, L. Chavez, and F. Calle. 1997. Genetic potential and stability of
carotene content in cassava roots. Euphytica 94(3): 367–373.
International Cassava Genetic Map Consortium (ICGMC). 2014. High-resolution linkage map
and chromosome-scale genome assembly for cassava (Manihot esculenta Crantz) from
10 populations. G3 (Bethesda). 5(1): 133–44.
Jennings, D.L. 1994. Breeding for resistance to African cassava mosaic geminivirus in East
Africa. Trop. Sci. 34(1): 110–122.
Langmead, B., and S.L. Salzberg. 2012. Fast gapped-read alignment with Bowtie 2. Nat
Methods 9(4): 357–359.
Lippert, C., J. Listgarten, Y. Liu, C.M. Kadie, R.I. Davidson, and D. Heckerman. 2011. FaST
linear mixed models for genome-wide association studies. Nat. Methods 8(10): 833–
837.
Listgarten, J., C. Lippert, C.M. Kadie, R.I. Davidson, E. Eskin, and D. Heckerman. 2012.
Improved linear mixed models for genome-wide association studies. Nat. Methods
9(6): 525–526.
Lu, R., A.M. Martin-Hernandez, J.R. Peart, I. Malcuit, and D.C. Baulcombe. 2003. Virus-
induced gene silencing in plants. Methods 30(4): 296–303.
Ly, D., M. Hamblin, I. Rabbi, G. Melaku, M. Bakare, H.G. Gauch, R. Okechukwu, A.G.O. Dixon,
P. Kulakow, and J.-L. Jannink. 2013. Relatedness and Genotype × Environment
Interaction Affect Prediction Accuracies in Genomic Selection: A Study in Cassava. Crop
Sci. 53(4): 1312–1325.
Marín Colorado, J.A., H. Ramírez, and M. Fregene. 2009. Genetic mapping and QTL analysis
for carotenes in a S1 population of cassava. Acta Agron. Univ. Nac. Colomb. 58(1): 15–
21.
Morillo C, A.C., Y. Morillo C, and H. Ceballos L. 2013. Identification of QTLs for carotene
content in the genome of cassava (Manihot esculenta Crantz) and S1 population
validation. Acta Agronómica, Univ. Nac. Colomb. 62(3): 196–206.
Myles, S., A.R. Boyko, C.L. Owens, P.J. Brown, F. Grassi, M.K. Aradhya, B. Prins, A. Reynolds,
J.-M. Chia, D. Ware, C.D. Bustamante, and E.S. Buckler. 2011. Genetic structure and
domestication history of the grape. Proc. Natl. Acad. Sci. U. S. A. 108(9): 3530–3535.
Njoku, D.N., V.E. Gracen, S.K. Offei, I.K. Asante, E.Y. Danquah, C.N. Egesi, and E. Okogbenin.
2014. Molecular marke r analysis of F1 progenies and their parents for carotenoids i
nheritance in African cassava (Manihot esculenta Crantz). African J. Biotechnol. 13(40):
3999–4007.
Njoku, D.N., V.E. Gracen, S.K. Offei, I.K. Asante, C.N. Egesi, P. Kulakow, and H. Ceballos.

2015. Parent-offspring regression analysis for total carotenoids and some agronomic traits in cassava. Euphytica 206(3): 657–666.

Okechukwu, R.U., and a. G.O. Dixon. 2008. Genetic Gains from 30 Years of Cassava Breeding in Nigeria for Storage Root Yield and Disease Resistance in Elite Cassava Genotypes. J. Crop Improv. 22(2): 181–208.

Ortiz, D., T. Sánchez, N. Morante, H. Ceballos, H. Pachón, M.C. Duque, A.L. Chávez, and A.F. Escobar. 2011. Sampling strategies for proper quantification of carotenoid content in cassava breeding. J. Plant Breed. Crop Sci. 3(1): 14–23.

Owens, B.F., A.E. Lipka, M. Magallanes-Lundback, T. Tiede, C.H. Diepenbrock, C.B. Kandianis, E. Kim, J. Cepela, M. Mateos-Hernandez, C.R. Buell, E.S. Buckler, D. DellaPenna, M. a Gore, and T. Rocheford. 2014. A Foundation for Provitamin A Biofortification of Maize: Genome-Wide Association and Genomic Prediction Models of Carotenoid Levels. Genetics 198(4): 1699–1716.

Pfeiffer, W.H., and B. McClafferty. 2007. HarvestPlus: Breeding Crops for Better Nutrition. Crop Sci. 47(Supplement_3): S-88.

Prochnik, S., P.R. Marri, B. Desany, P.D. Rabinowicz, C. Kodira, M. Mohiuddin, F. Rodriguez, C. Fauquet, J. Tohme, T. Harkins, D.S. Rokhsar, and S. Rounsley. 2012. The Cassava Genome: Current Progress, Future Directions. Trop. Plant Biol. 5(1): 88–94.

Purcell, S., B. Neale, K. Todd-Brown, L. Thomas, M. a R. Ferreira, D. Bender, J. Maller, P. Sklar, P.I.W. de Bakker, M.J. Daly, and P.C. Sham. 2007. PLINK: a tool set for whole-genome association and population-based linkage analyses. Am. J. Hum. Genet. 81(September): 559–575.

Rabbi, I., M. Hamblin, M. Gedil, P. Kulakow, M. Ferguson, A.S. Ikpan, D. Ly, and J.L. Jannink. 2014a. Genetic mapping using genotyping-by-sequencing in the clonally propagated cassava. Crop Sci. 54(4): 1384–1396.

Rabbi, I.Y., M.T. Hamblin, P.L. Kumar, M. a Gedil, A.S. Ikpan, J.-L. Jannink, and P. a Kulakow. 2014b. High-resolution mapping of resistance to cassava mosaic geminiviruses in cassava using genotyping-by-sequencing and its implications for breeding. Virus Res. 186: 87–96.

Saltzman, A., E. Birol, H.E. Bouis, E. Boy, F.F. De Moura, Y. Islam, and W.H. Pfeiffer. 2013. Biofortification : Progress toward a more nourishing future. Glob. Food Sec. 2(1): 9–17.

Sánchez, T., H. Ceballos, D. Dufour, D. Ortiz, N. Morante, F. Calle, T. Zum Felde, M. Domínguez, and F. Davrieux. 2014. Prediction of carotenoids, cyanide and dry matter contents in fresh cassava root using NIRS and Hunter color techniques. Food Chem. 151: 444–451.

Sander, J.D., and J.K. Joung. 2014. CRISPR-Cas systems for editing, regulating and targeting genomes. Nat. Biotechnol. 32(4): 347–55.

Sayre, R., J.R. Beeching, E.B. Cahoon, C. Egesi, C. Fauquet, J. Fellman, M. Fregene, W. Gruissem, S. Mallowa, M. Manary, B. Maziya-Dixon, A. Mbanaso, D.P. Schachtman, D. Siritunga, N. Taylor, H. Vanderschuren, and P. Zhang. 2011. The BioCassava plus program: biofortification of cassava for sub-Saharan Africa. Annu. Rev. Plant Biol. 62: 251–72.

Smith, A.M. 2008. Prospects for increasing starch and sucrose yields for bioethanol production. Plant J. 54(4): 546–558.

Suwarno, W.B., K. V Pixley, N. Palacios-Rojas, S.M. Kaeppler, and R. Babu. 2015. Genome-wide association analysis reveals new targets for carotenoid biofortification in maize. Theor. Appl. Genet. 128(5): 851–864.

Wang, X., L. Chang, Z. Tong, D. Wang, Q. Yin, D. Wang, X. Jin, Q. Yang, L. Wang, Y. Sun, Q.
Huang, A. Guo, and M. Peng. 2016. Proteomics Profiling Reveals Carbohydrate
Metabolic Enzymes and 14-3-3 Proteins Play Important Roles for Starch Accumulation
during Cassava Root Tuberization. Sci. Rep. 6(January): 19643.
Welsch, R., J. Arango, C. Bär, B. Salazar, S. Al-Babili, J. Beltrán, P. Chavarriaga, H. Ceballos, J.
Tohme, and P. Beyer. 2010. Provitamin A accumulation in cassava (Manihot esculenta)
roots driven by a single nucleotide polymorphism in a phytoene synthase gene. Plant
Cell 22(10): 3348–56.
Wolfe, M.D., I.Y. Rabbi, C. Egesi, M. Hamblin, R. Kawuki, P. Kulakow, R. Lozano, D.P. Del
Carpio, P. Ramu, and J.-L. Jannink. 2016. Genome-Wide Association and Prediction
Reveals Genetic Architecture of Cassava Mosaic Disease Resistance and Prospects for
Rapid Genetic Improvement. Plant Genome 9(2): 1–248.
Wong, W.W.L., J. Griesman, and Z.Z. Feng. 2014. Imputing genotypes using regularized
generalized linear regression models. Stat. Appl. Genet. Mol. Biol. 13(5).
Yan, J., T. Shah, M.L. Warburton, E.S. Buckler, M.D. McMullen, and J. Crouch. 2009. Genetic
characterization and linkage disequilibrium estimation of a global maize collection
using SNP markers. PLoS One 4(12): e8451.
Yang, J., D. An, and P. Zhang. 2011. Expression Profiling of Cassava Storage Roots Reveals an
Active Process of Glycolysis/Gluconeogenesis. J. Integr. Plant Biol. 53(3): 193–211.
Yu, J., J.B. Holland, M.D. Mcmullen, and E.S. Buckler. 2008. Genetic Design and Statistical
Power of Nested Association Mapping in Maize. 551(January): 539–551.
Yu, J., G. Pressoir, W.H. Briggs, I. Vroh Bi, M. Yamasaki, J.F. Doebley, M.D. McMullen, B.S.
Gaut, D.M. Nielsen, J.B. Holland, S. Kresovich, and E.S. Buckler. 2006. A unified mixed-
model method for association mapping that accounts for multiple levels of relatedness.
Nat. Genet. 38(2): 203–208.
Zeeman, S.C., J. Kossmann, and A.M. Smith. 2010. Starch: its metabolism, evolution, and
biotechnological modification in plants. Annu. Rev. Plant Biol. 61: 209–234.
Zhang, Z., E. Ersoz, C.-Q. Lai, R.J. Todhunter, H.K. Tiwari, M. a Gore, P.J. Bradbury, J. Yu, D.K.
Arnett, J.M. Ordovas, and E.S. Buckler. 2010. Mixed linear model approach adapted for
genome-wide association studies. Nat. Genet. 42(4): 355–60.
Zhao, K., C.-W. Tung, G.C. Eizenga, M.H. Wright, M.L. Ali, A.H. Price, G.J. Norton, M.R. Islam,
A. Reynolds, J. Mezey, A.M. McClung, C.D. Bustamante, and S.R. McCouch. 2011.
Genome-wide association mapping reveals a rich genetic architecture of complex traits
in Oryza sativa. Nat. Commun. 2: 467.
645
646

Tables

Table 1. Summary of phenotype variation, variance components (±se) and broad-sense heritability (H^2) for dry matter, color chart and Chromameter CIELAB readings.

Trait	DM (%)	TC-CHART	L*	a*	b*
Minimum	8.4	1.0	69.5	-3.0	11.1
Average	28.6	1.7	84.4	-0.3	20.8
Maximum	45.4	7.0	90.2	4.6	40.8
N	3232	4237	1360	1360	1360
V_G	16.52 (4.06)	1.11 (1.05)	5.22 (2.29)	0.86 (0.92)	16.52 (4.06)
V_{GxE}	3.53 (1.88)	0.06 (0.24)	0.77 (0.88)	0.16 (0.41)	2.87 (1.69)
V_E	1.91 (1.38)	0.01 (0.11)	0.53 (0.73)	0.08 (0.28)	1.11 (1.05)
Residual	10.27 (3.20)	0.17 (0.41)	3.26 (1.80)	0.30 (0.55)	2.71 (1.65)
H^2	0.51	0.82	0.53	0.61	0.87

Table 2. Summary of significant associations between selected traits and SNP markers from the MLM analysis. Only results in the major loci from Chromosome 1 are shown.

Trait	SNP	Chr	Position (bp)	P-value	Candidate genes and mid-position (bp)
Color chart	S1_24121306	1	24,121,306	1.74E-22	Phytoene synthase (Manes.01G124200; 24,155,070 bp)
Color chart	S1_30543382	1	30,543,382	2.91E-11	NA
Chromameter b*	S1_24159585	1	24,159,585	1.79E-19	Phytoene synthase (Manes.01G124200; 24,155,070 bp)
Chromameter b*	S1_30543382	1	30,543,382	1.66E-11	NA
Dry matter	S1_24121306	1	24,121,306	1.86E-12	UDP-glucose pyrophosphorylase (Manes.01G123000; 24,061,652 bp); sucrose synthase (Manes.01G123800; 24,142,314 bp)

Chr = Chromosome (version 6 of cassava reference genome);

658 **Figures**

659
660 **Figure 1.** Distribution of phenotype for TCHART, Dry matter content, and chromameter (b*).
661
662
663

664
Figure 2. Relationship between dry matter content and root yellowness BLUPs (expressed as
b* value of chromameter measurement). Different symbols denote the genotype at marker
S1_24121306 that is associated with both dry matter content and root color intensity.

Figure 3. Population structure of the Genetic Gain collection. (A) PCA bi-plot of the first two

axes; (B) Neighbor-joining dendrogram calculated from pairwise IBS distance. Yellow color

highlights accessions with yellow roots.

Figure 4. Local pattern of linkage disequilibrium (r^2) along each of the 18 cassava

chromosomes. Note the large LD blocks in chromosomes 1, 4 and 10. SNPs are arrayed

according to their order, and not their physical position.

678
679 **Figure 5.** A Moving-average based LD decay profile in the Genetic Gain population. (A) all
680 chromosomes except 1, 4 and 10; (B) chromosomes 1, 4 and 10.
681

Figure 6. Genome-wide association results. Manhattan and Quantile-quantile plot of the MLM model for: root yellowness estimated using (A) chromameter b* value; and (B) color chart method; and (C) dry matter content. The red horizontal line indicates the genome-wide significance threshold.

Figure 7. GWAS results for chromosome 1. Manhattan and Quantile-quantile plot of the MLM model for: (A) root yellowness as measured by chromameter b* value; (B) color chart; and (C) dry matter content. Note the common peak at ~ 24.1 Mbp region for the three traits. Red horizontal line indicates the genome-wide significance threshold. The SNPs are colored according to their degree of linkage disequilibrium (r^2) with the leading variant (i.e. top SNP for the first peak at 24.1 Mbp). The vertical blue lines in (A) and (B) denote the position of the carotenoid biosynthesis gene, phytoene synthase (24,155,070 bp), and those

on (C) denotes the positions of the UDP-glucose pyrophosphorylase (24,061,652 bp) and
sucrose synthase (24,142,314 bp).

Figure 8. Manhattan plots of the MLM analysis of the yellow root (top) and white root
subpopulations (bottom). Below each is an LD heatmap showing pairwise squared
correlation of alleles between markers along chromosome 1. Note the large number of SNPs
showing significant association with dry matter in the white subpopulation compared to
that of the yellow subpopulation. Red horizontal line indicates the genome-wide
significance threshold. The vertical blue lines are same as in Figure 7.

706
707 **Figure 9.** Selection sweep associated with positive selection for provitamin A trait varieties
708 in the genetic gain population. Red dashed line indicates the position of SNP S1_24121306.
709

Figure 10. Effect of the most significantly associated markers on the BLUPs for yellow color
measured by: (A) Chromameter (b* value) and (B) color chart in the Genetic Gain
population. The boxplots show the effects of the most significantly associated SNPs at first
and second peaks (above) and the two-locus haplotypes (below) on chromosomes 1. (C)
Effect of the most significantly associated markers on the BLUPs for dry matter content.
Alleles are coded as 0 = homozygous reference genome; 1 = heterozygous and 2 =
homozygous non-reference genome. The dashed line represents the population mean of the
BLUPs. The numbers in parenthesis below genotypic categories refer to the number of
accessions for each genotype.
720
721

722
723
724 **Supplementary Figure 1.** Quantile–quantile plots for P-values obtained from simple GLM,
725 GLM+5PCs and MLM model for color chart, chromameter b* value and dry matter content.
726

727
728 **Supplementary Figure 2.** Heatmap of identity-by-descent relationship using SNPs from large
729 LD block in chromosome 1 around the major QTL region.
730