Ecological drivers of jellyfish blooms – the complex life history

of a 'well-known' medusa (*Aurelia aurita*)

J. Goldstein[*, a, b], U.K. Steiner[b, c]

[a]Marine Biological Research Centre, University of Southern Denmark, Hindsholmvej 11, 5300

Kerteminde, Denmark

[b]Department of Biology, University of Southern Denmark, Campusvej 55, 5230 Odense,

Denmark

[c]Center on Population Dynamics, University of Southern Denmark, Campusvej 55, 5230 Odense,

Denmark

[*]Corresponding author: jgoldstein@biology.sdu.dk

Running head Drivers of jellyfish blooms

Summary

1. Jellyfish blooms are conspicuous demographic events with significant ecological

and socio-economic impact. Despite worldwide concern about an increased

frequency and intensity of jellyfish mass occurrences, predicting their booms and

busts remains challenging.

2. Forecasting how jellyfish populations may respond to environmental change

requires taking into account their complex life histories. Metagenic life cycles,

which include a benthic polyp stage, can boost jellyfish mass occurrences via

asexual recruitment of pelagic medusae.

3. Here we present stage-structured matrix population models with monthly,

individual-based demographic rates of all life stages of the moon jellyfish *Aurelia*

aurita (sensu stricto) for low and high food conditions. We investigate the life

stage-dynamics of such complex populations to illustrate how changes in medusa

density depend on non-medusa stage dynamics.

4. We show that increased food availability can be an important ecological driver of

jellyfish mass occurrence, as it can temporarily shift the population structure from

polyp- to medusa-dominated. Projecting populations for a winter warming

scenario additionally enhanced the booms and busts of jellyfish blooms.

5. We identify demographic key variables that control the intensity and frequency of

jellyfish blooms in response to environmental drivers such as habitat

eutrophication and climate change. By contributing to an improved understanding

of mass occurrence phenomena, our findings provide perspective for future

management of ecosystem health.

Key-words biodemography, environmental change, jellyfish blooms, matrix model,

population structure

Introduction

The biodiversity, productivity and resilience of marine ecosystems worldwide are

threatened by an imbalance in the ecological parameters regulating population outbreaks.

Mass occurrences of micro- and macroalgae, cnidarians, ctenophores, molluscs,

echinoderms, tunicates and fish seem to appear with increased frequency and intensity,

covering larger geographical areas (Breithaupt, 2003; Lucas & Dawson, 2014; Maji,

Bhattacharyya, & Pal 2016). Despite mass occurrences and mass mortality events being

widespread – with major implications for marine ecosystem health – predictions have

remained challenging due to an incomplete concept of underlying demographic processes

(Fey et al., 2015). Jellyfish blooms are a prominent example for mass occurrences in

which the population density of one life stage, the pelagic medusa, shows distinct peaks

over broad temporal and regional scales (Lucas, Graham, & Widmer 2012). These

demographic events (cf. Lucas & Dawson, 2014) may have lasting ecological, economic

and social implications. Jellyfish blooms can trigger trophic cascades such as collapse of

pelagic food web structures due to their severe predatory impact on a broad spectrum of

zooplankton species and fish larvae (Richardson, Bakun, Hays, & Gibbons 2009;

Riisgård, Andersen, & Hoffmann 2012; Schneider & Behrends, 1994, 1998). Further,

jellyfish mass occurrences can interfere with human activities such as tourism, fisheries,

aquaculture and power production (Purcell, 2012; Purcell, Uye, & Lo 2007). Jellyfish

populations, i.e. aggregations of the medusa stage, appear to be increasing in numerous

coastal ecosystems around the world (Brotz, Cheung, Kleisner, Pakhomov, & Pauly

2012). Available evidence suggests that jellyfish outbreaks, i.e. exceptional, sudden

increases in biomass (Lucas & Dawson, 2014), may be favored by anthropogenic

influences, in particular overfishing, eutrophication, habitat modification, species

translocation and climate change (Purcell, 2012; Richardson, Bakun, Hays, & Gibbons

2009).

Bloom-forming jellyfish species are primarily found within the cnidarian taxon

Scyphozoa and can feature complex life histories (Dawson & Hamner, 2009; Hamner &

Dawson, 2009; Lucas & Dawson, 2014). Their metagenic life cycles typically include a

pelagic, sexually reproducing medusa and a benthic, asexually reproducing polyp stage

(Lucas, Graham, & Widmer 2012). Such life cycles are characterized by high stage-

specific plasticity to changing environmental conditions (e.g. Gröndahl, 1989; Hamner &

Jenssen, 1974; Liu, Lo, Purcell, & Chang 2009). Although the development of medusa

populations is directly linked to survival rates of several precedent life stages, i.e. planula

larvae, polyps and ephyrae (Lucas, 2001), information on the demographic rates of

bloom-forming jellyfish has remained scarce (Xie, Fan, Wang, & Chen 2015).

Quantifying demographic rates of all stages in the life cycle of species showing mass

occurrence is thus fundamental to identifying the environmental variables causing

changes in population dynamics and structure.

Here, we explore the demographic structure and environmental drivers underlying

mass occurrence of the moon jellyfish *Aurelia aurita* (sensu stricto). We quantify

demographic rates of all life stages of *A. aurita* (larvae, polyps, ephyrae and medusae),

i.e. survival, stage-transitions and fecundity, based on experimental and field data from

two temperate Danish water systems with contrasting prey availability. Using these

demographic rates, we parameterize stage-structured population models to project the life

cycle of jellyfish populations for low and high food levels. Since warm temperature has

been associated with increased numbers of medusae for most temperate scyphozoan

species (Purcell, 2005), we further predict the impact of a winter warming scenario on the

demographic dynamics of *A. aurita*. Our results provide novel insights into the

development of metagenic jellyfish populations under increased food levels and increased

winter temperatures. Such information is essential to assess the boosting potential of

environmental drivers on species showing mass occurrence and provides a milestone for

identifying and managing demographic irregularities in marine ecosystems.

Materials and Methods

Study organism

The moon jellyfish *Aurelia* (Scyphozoa, Cnidaria) belongs to an ubiquitous, well-studied

genus with at least 16 species, most of which seem to be distributed in specific coastal

and shelf sea regions around the world (Dawson, Gupta, & England 2005). The life

history, i.e. individual stage-trajectory, of *Aurelia aurita*, sensu stricto, is characterized by

a complex, metagenic life cycle which involves sexually reproducing medusae and

asexually reproducing polyps (Figure 1a). Medusae of *A. aurita* are pelagic filter feeders

which differentiate into distinct male and female individuals upon sexual maturation

(Lucas 2001). After fertilization of oocytes, female medusae produce large numbers of

lecithotrophic (yolk-feeding) planula larvae (Ishii & Takagi, 2003), which spend 12 h to 1

week in the water column (Lucas, 2001) before they settle on suitable hard substrate and

metamorphose into polyps (Holst & Jarms, 2007; Keen, 1987). The inconspicuous

benthic polyp stage of *A. aurita* can occur in densities between 60 000 and 400 000 ind.

m^{-2} (Lucas, 2001) and reproduces asexually by several modes, including i) the production

of well protected chitin-covered resting stages called podocysts (Arai, 2009), ii) budding

of new polyps (Ishii & Watanabe, 2003), and iii) the release of multiple immature

medusae, i.e. ephyrae, through transverse fission (strobilation). After asexual

propagation, polyps remain in the polyp stage, while the development of pelagic ephyrae

into sexually mature medusae closes the life cycle (Lucas, Graham, & Widmer 2012).

Data collection

We cultivated planula larvae (L), polyps (P), ephyrae (E) and medusae (M) of *A. aurita* in

the laboratories of the Marine Biological Research Centre, Kerteminde (Denmark)

between June 2013 and November 2014. We set up separate experimental series for each

life stage using filtered (38 µm) seawater at constant salinity (20 PSU) and with water

temperature following seasonal variability of the local temperate climate zone (5-22 °C).

We fed all life stages, except the planula larvae, with low (*L*) or high (*H*) amounts

(corresponding to a ratio ~1:10) of two-day old *Artemia salina* nauplii and regularly

exchanged the seawater in our cultivation containers. We adjusted the fed biomass of *A.*

salina (2.55 µg C ind.$^{-1}$; Vanhaecke & Sorgeloos, 1980) to the size-specific energy

demands of each life stage (Table 1).

We obtained planula larvae from female medusae that were collected in two

contrasting Danish water systems – the fjord system Kerteminde Fjord/Kertinge Nor and

the major water system Limfjorden. Zooplankton prey biomasses are in a range <1−100

µg C L^{-1} in Kerteminde Fjord/Kertinge Nor (Nielsen, Pedersen, & Riisgård 1997) and

>10−1000 µg C L^{-1} in Limfjorden, respectively (Møller & Riisgård, 2007). Directly after

the release of planulae, we transferred the larvae individually to multi-well seawater

containers (200 µL). During daily counts, we determined the number of living larvae (l_{LL},

ind. day^{-1}) and the number of larvae that had completed metamorphosis into polyps (l_{LP},

ind. day^{-1}) until all planulae had metamorphosed into polyps or had died.

We cultivated polyps of *A. aurita* from planulae, released by female medusae from

Kertinge Nor, which we initially added to small seawater containers (50 mL) with

floating polystyrene substrate plates to facilitate settlement. After 24 h, we removed all

but one newly settled polyp per substrate plate and replaced the seawater in the

containers. We subsequently raised and cultivated the individual polyps under low or

high food conditions (Table 1; cf. Kamiyama, 2011). To maintain the initial polyp

cohorts, we instantly removed any buds of the "mother" polyps from substrate plates.

During the whole experimental period of 12 months, no excystment of polyps from

podocysts was observed. Each week we counted the number of live polyps (l_{PP}, ind.

week^{-1}) as well as the number of buds (N_{PP}, ind. week^{-1}) and ephyrae per polyp (N_{PE}, ind.

week^{-1}).

Ephyrae which were released by polyps under low and high food conditions were

individually transferred to seawater flasks (250 mL) with constant water movement and

air supply, and continuously reared under either low or high food conditions. Food levels

for ephyrae corresponded to the minimum energy demands for maintenance of a

'standard' ephyra with 8 mm inter-rhopalia diameter (Båmstedt, Lane, & Martinussen

1999; Bengtson, Leger, & Sorgeloos 1991; Eckert, Randall, & Augustine 1988; Frandsen

& Riisgård, 1997), or the 10-fold carbon load, respectively (Table 1). Each week we

counted the number of surviving ephyrae (l_{EE}, ind. week^{-1}) and the number of ephyrae

that had completed transition to medusa (l_{EM}, ind. week^{-1}); the latter we defined by the

presence of fully developed intermediate lappets and the final closure of adradial clefts.

We performed these counts until all ephyrae had either developed into medusae or had

died.

We continuously tracked all newly transitioned medusae and additional sexually

mature females collected in Kertinge Nor, in individual aquaria (1 L) with constant air

supply. We kept the medusae under low or high food levels which resembled complete

starvation or the 10-fold energy demands of a 'standard' 50 mm umbrella diameter

medusa (Table 1; Olesen, Frandsen, & Riisgård 1994; Goldstein & Riisgård, 2016). We

recorded the survival of medusae (l_{MM}, ind. day^{-1}) daily until all individuals in the low

food treatment had died; at that time all medusae in the high food treatment were still

alive. We estimated the daily release of planula larvae by female medusae during the

reproductive period (N_{ML}, ind. day^{-1}) using the field observation-based relationship

$N_{\mathrm{ML}} = 0.087 \times 160.8 \times e^{0.029d}$ (Goldstein & Riisgård, 2016), where average umbrella

diameters (d, mm) of 40 mm and 200 mm reflected low and high food conditions.

Estimating stage-specific demographic rates

We used the experimental data to parameterize age-stage-structured matrix population

models. For these models, we used discrete time steps of one month to estimate survival,

stage transitions and fecundity of the stages L, P, E and M between age x and age $x+1$.

Individuals in each life stage i have certain monthly probabilities to survive in their

current stage (t_{ii}), to transition to another stage j (t_{ij}) and to produce offspring of stage j

(F_{ij}). Each stage has its unique survival rate (t_{LL}, t_{PP}, t_{EE}, t_{MM}, month^{-1}) and stage

transitions are possible from larva to polyp (t_{LP}, month^{-1}) or from ephyra to medusa (t_{EM},

month^{-1}). Fecundity includes asexual reproduction of polyps by budding (F_{PP}, ind. ind.$^{-1}$

month^{-1}) or release of ephyrae (F_{PE}, ind. ind.$^{-1}$ month^{-1}), and sexual reproduction of

medusae by the release of planula larvae (F_{ML}, ind. ind.$^{-1}$ month^{-1}; Figure 1b).

Details on estimating monthly stage-specific demographic rates of *A. aurita* are

available from the Supporting Information (Appendix S1 & S2). We first estimated

monthly transition probabilities and fecundities for each life-stage i under low and high

food conditions (Appendix S1) based on count ratios from our empirical data with either

daily (L, M) or weekly resolution (P, E). For converting the daily release of planula

larvae into monthly fecundity of medusae, we assumed a sex ratio of 1:1. Since transition

probabilities and fecundities between subsequent time steps depend on survivorship

within each monthly time step, we assumed that individuals that died during a given

month, had died halfway during this month, i.e. they survived throughout half the month

on average (Appendix S2). This assumption is needed as, for instance, a medusa might

still release larvae before its death within a given month. Further, some ephyrae

completed transition to the medusa stage within less than a month after they were

released by a polyp. We therefore corrected the monthly release of ephyrae by polyps not

only for daily survivorship of ephyrae, but also for the transition probability to the

medusa stage within a given month.

Monthly stage-structured matrix population model

Our matrix population model describes the transitions of each life stage taking place

during each projection interval, as defined by monthly stage-specific demographic rates

for each of the 12 month of a year. Age is determined by month, and the final stages are

P, E and M. Since larval transition to the polyp stage was restricted to days or hours ($t_{LP} =$

0 for incompetent larvae with $t_{LL} > 0$), we linked the monthly transition from larva to

polyp stage to fecundity of medusae ($F_{ML} \times t_{LP}$) to build the basic structure of an

irreducible monthly stage-structured matrix model (Figure 1c). Using our empirical

datasets on all demographic rates of *A. aurita* under low and high food conditions,

respectively, we constructed monthly stage-structured submatrices of the following

structure:

$$
A_x = \begin{pmatrix} t_{PP} + F_{PP} & 0 & F_{ML} \times t_{LP} \\ F_{PE} & t_{EE} & 0 \\ F_{PE} \times t_{EM} & t_{EM} & t_{MM} \end{pmatrix}
$$

225   ,

226   where *x* denotes the age in months. Based on 12 monthly submatrices A_x (3 × 3) which

227   reflect seasonal life cycle dynamics of temperate *A. aurita* populations (Appendix S3:

228 Figures S1 & S2), we constructed yearly population matrices A (36×36; time step = 1

229 month) for projections over a 10-year interval (120 iterations):

$$
A = \begin{pmatrix}
\cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & A_{12} \\
A_1 & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & A_2 & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & A_3 & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & A_4 & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & \cdot & A_5 & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & \cdot & \cdot & A_6 & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & \cdot & \cdot & \cdot & A_7 & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & A_8 & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & A_9 & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & A_{10} & \cdot & \cdot \\
\cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & A_{11} & \cdot
\end{pmatrix}
$$

.

In some months, we observed no survivorship of ephyrae or medusae, which would lead

to reducible matrices (Caswell, 2001); in these cases, we included a small transition

probability $c = 1 \times 10^{-10}$ as placeholder (Appendix S3: Figures S1 & S2). Our results

were robust to this assumption.

Adjusting matrix parameters to environmental conditions

We constructed yearly projection matrix models for the jellyfish population under low

and high food conditions. Since our transition probabilities from larva to polyp were

unrealistically high due to the sheltered laboratory environment (as an important

microzooplankton component, planula larvae are highly subject to predation under

natural conditions), we calibrated the larva to polyp transition (t_{LP}) to a level that resulted

in a stationary population (i.e. characterized by constant population size over time) under

low food conditions. We applied the same level of calibration as for the low food

condition to the projection matrix model for high food conditions, leaving us with a

rapidly growing population.

We parameterized additional projection matrix models to explore the potential effects

of a winter warming scenario under low and high food conditions. A predicted winter

warming trend will likely increase present water temperatures from 5 to 10 °C and thus

benefit the production of *A. aurita* ephyrae by extended strobilation periods and enhanced

ephyra release per polyp (Holst, 2012). Our winter warming scenario included 1.8-fold

increased ephyra production (i.e. $F_{PE} \times 1.8$) during strobilation periods that we extended

by one month, i.e. mean F_{PE} of proceeding and subsequent month. We additionally

adjusted corresponding probabilities for ephyra survival and transition from ephyra to

medusa to t_{EE} and t_{EM} of subsequent month, respectively (Appendix S3: Figures S1 &

S2). With these modified matrices, we investigated the demographic effects of a

predicted winter warming trend on *A. aurita* populations under low and high food

conditions.

Computing demographic parameters and population projection

We analyzed our yearly population projection matrices using R version 3.1.3 (R Core

Team, 2015) and compared the demographic dynamics of *A. aurita* under low and high

food levels, and in combination with a winter warming trend. We used R packages

'popbio' (Stubben & Milligan, 2007) and 'popdemo' (Stott, Hodgson, & Townley 2012)

to compute population growth rate λ (dominant eigenvalue), stable stage distribution

(right eigenvector **w** corresponding to the dominant eigenvalue; i.e. proportions of

individuals in each stage remain constant on the long-term), reproductive value *R* (left

eigenvector **v** corresponding to the dominant eigenvalue; i.e. expected lifetime

reproduction of an individual in stage i), the net reproductive rate R_0 (i.e. average number

of offspring by which an individual will be replaced during its lifetime), the generation

time T (i.e. average age at which an individual has produced the average number of

offspring, R_0), the life expectancy e_0 (i.e. mean time to death), and the sensitivity of λ to

perturbations of the matrix elements (only elements >0 are considered). We further

performed a fixed life-table response experiment (LTRE) analysis (Caswell, 1996;

Tuljapurkar & Caswell, 1997) for pairwise comparison of the sensitivity of λ to changes

in stage-specific transitions under high versus low food conditions.

277   **Results**

Demographic dynamics of jellyfish under low and high food conditions

The metagenic life cycle of *A. aurita* (Figure 1a) was reflected by cyclic (imprimitive)

structure of our matrix model, expressing annual peak and pit dynamics in projected

population density (Figure 2a). Relative to the calibrated stationary stable population with

$\lambda = 1$ month^{-1} under low food conditions, the calibrated population growth rate for 10-

fold increased food availability indicated ($\lambda^{12} =$) 346-fold yearly increase in population

size (Table 2). Such differences in growth rate due to changes in food availability provide

high leverage for population booms and busts. Similar to low food levels, the projection

matrix under high food availability showed cyclic annual structure with typical peak and

pit dynamics in projected population density, but life cycle amplitudes were much

increased (Figure 2b).

Under low food conditions, the overall population density (taking into account all life

stages) increased to a peak during spring and summer and thereafter decreased to low

densities during autumn and winter. Peak density and dynamics under high food levels

differed significantly compared to low food conditions, showing two distinct maxima in

population density, one during spring and one during autumn, each followed by a

subsequent minimum in summer and winter (Figures 2a–b). The convergence time to

stable stage (i.e. the damping ratio ρ) of our modelled *A. aurita* population was ~5 years

under low food conditions and reduced by a factor 21, for the high food population

(Table 2). The stable stage distribution under low food availability was vastly dominated

by polyps (0.94) that survived throughout the year. Ephyrae and medusae accounted for

comparatively small fractions of the population (0.04 and 0.02, respectively) and showed

typical seasonal patterns. Ephyrae were released during April/May and survived into

September. Only few ephyrae transitioned to the medusa stage under low food

conditions, so that medusae were present from June to December and contributed only

marginally to restocking polyp densities by the release of planula larvae during autumn

and winter (Figure 2c). Under high food supply, less polyps (0.76) dominated the overall

A. aurita population and the stable stage distribution was characterized by a 10-fold

increased medusa fraction (0.2) compared to low food conditions (Table 2). Compared to

a year-round standing stock of polyps under low food conditions, the stable stage

proportion of polyps under high food levels showed a peak in autumn after being

restocked by the release of planula larvae, and subsequently decreased throughout the

year. Ephyrae were released in March/April, and the medusa fraction dominated the *A.*

aurita population from April to July under high food levels (Figure 2d).

The reproductive value was low for ephyrae (0.03) under low food conditions, and

sexual reproduction of medusae contributed only about half as much (0.42) to future

generations as the asexually reproducing polyp stage (1; note scaling to 1 for polyps in

January). Under high food conditions, the reproductive value of ephyrae and medusae

increased by a factor 8.1 and 21.6, respectively (Table 2). A net reproductive rate of 1

indicated that each polyp is replaced by 1 new polyp at the end of its life under low food

conditions which resulted in long generation times of ~16.7 years. High food levels

increased the net reproductive rate to 57 polyps replacing each polyp at the end of its life

and boosted the average time between two consecutive generations to 8 months. Life

expectancy of polyps, estimated as the mean time to death, was 33 years under low food

conditions, while ephyrae and medusae revealed much shorter life expectancies of 2 and

3 months, respectively. In consensus with a more pronounced impact of the medusa stage

on population structure under high food levels, life expectancy of polyps decreased to 7

years, while the length of ephyra and especially medusa lives extended to 3 and 8 months,

respectively (Table 2).

Sensitivity analysis generally highlighted a strong influence of survival rates and less

pronounced effects of fecundity on the growth of jellyfish populations. Under low food

conditions, population growth was most sensitive to changes in polyp survival ($t_{PP}+F_{PP}$),

expressed by constant sensitivity 0.08 (relative maximum value in the sensitivity matrix)

throughout the months of the year (Figure 3a). Comparatively, λ showed a much lower

sensitivity of 0.02 to changes in ephyra production (F_{PE}) from March to May. The overall

sensitivity of λ to perturbations in the projection matrix elements increased under high

compared to low food levels. Under high food conditions, changes in polyp survival

($t_{PP}+F_{PP}$) affected λ particularly from October to April, with sensitivities of 0.06 to 0.13.

In contrast to low food levels however, λ was most sensitive to changes in medusa

survival (t_{MM}) under high food levels, with sensitivity remaining high, ranging within

0.07 to 0.16 from April to October, and reaching a maximum of 0.32 from June to July

(Figure 3b). We further detected sensitivities of λ in the range 0.02 to 0.03 for

perturbations in ephyra production (F_{PE}) from March to May under high food conditions.

Differences in the sensitivity of λ between food treatments were expressed by the results

of our life table response experiment (LTRE) analysis. High food availability led to 7.7 to

12.6 % increased sensitivity for the transition probabilities from polyp to medusa stage

(i.e. $F_{PE}+t_{EM}$) from March to May. Further, increased food conditions enhanced the

sensitivity of λ for medusa survival (i.e. t_{MM}) during April by 7.7 % and for polyp

production via sexual reproduction of medusae (i.e. $F_{ML}+t_{LP}$) from August to December

by 5.0 to 6.8 % (Figure 3c).

Demographic dynamics of jellyfish in a winter warming scenario

For our winter warming scenario with extended strobilation periods and increased ephyra

release, population growth rates of *A. aurita* increased to 1.01 and 1.73 per month under

low and high food conditions, corresponding to 1.1- and 729-fold yearly increases in

population size (i.e., relative increases by 6.2 % and 110.9 %), respectively. Winter

warming led to stronger variability in inter-annual population structure in both food

treatments, expressed by more distinct peaks and pits in population density (Figures

2a–b). Under low food levels, the stable population structure of *A. aurita* consisted of 2.7

% more ephyrae, 7.9 % more medusae, and 10.6 % less polyps per year as a consequence

of winter warming (Figure 2c). Under high food levels, increased winter temperatures

resulted in a stable stage distribution with 17.4 % more medusae, while the proportions of

ephyrae and polyps decreased by 1.7 % and 15.7 %, respectively (Table 2). An earlier

onset of strobilation periods in the scenario and fast transition of ephyrae caused

significant proportions of medusae to be already present from February (Figure 2d).

Winter warming caused substantial changes in reproductive rates and generation times

of the projected jellyfish populations; these changes were particularly pronounced under

low food conditions. The reproductive value of ephyrae in the low food population was

257-fold increased, while the relative contribution to the number of offspring by medusae

decreased to near zero. Winter warming in combination with high food levels decreased

reproductive values of both ephyrae and medusae by a factor 1.5 and 1.2, respectively

(Table 2). Increased winter temperatures raised the net reproductive rate by 75.1 % under

low food conditions and by 42.0 % under high food conditions. The winter warming

scenario shortened the generation time by 7.4 years under low food conditions, while

generation times were shortened by only 9 days under high food conditions. Winter

warming increased the life expectancy of ephyrae by 12 and 18 days, respectively,

whereas life expectancy of polyps and medusae remained unchanged under both low and

high food levels (Table 2).

Discussion

We reveal the demographic dynamics and life history shifts underlying jellyfish mass

occurrence and show that variable intensity and frequency is subject to specific

ecological triggers. Our matrix projections quantify how the booms and busts

characterizing jellyfish blooms are enhanced by naturally occurring variations in the food

regime. We suggest the observed temporal shift from polyp- to medusa-dominated

populations in response to increased food availability as a key mechanism causing

jellyfish outbreaks, which are likely boosted in intensity and frequency as a consequence

of the combined effects with a predicted winter warming. Our findings support previous

studies which have pointed out metagenic life cycles as an ultimate driver of boom and

bust population dynamics (Brotz, Cheung, Kleisner, Pakhomov, & Pauly 2012; Hamner

& Dawson, 2009; Lucas & Dawson, 2014, their Figure 2.4b; Pitt, Welsh, & Condon

2009). Considering jellyfish outbreaks as highly sensitive indicators for ocean

degradation (Schrope, 2012), the present insights may provide fundamental knowledge

regarding the health of marine ecosystems worldwide.

Food availability as a key driver of jellyfish blooms

Our findings suggest food availability as a major constraint for the intensity and

frequency of jellyfish blooms, confirming previous observations of food availability as a

key variable in limiting the development of ephyrae (Fu, Shibata, Makabe, Ikeda, & Uye

2014) and the survivorship and fecundity of medusae (Goldstein & Riisgård, 2016). Yet,

to our knowledge, we present the first comprehensive quantitative model of how

increased food availability can trigger demographic changes in a scyphozoan jellyfish

population. Such changes include enhanced boom and bust dynamics and shorter

generation times. We show that complex life cycles can facilitate jellyfish blooms and

emphasize the key role of benthic polyps in ensuring long-term survival (Boero et al.,

2008; Lucas, Graham, & Widmer 2012), particularly when food is scarce. Life-history

trade-offs such as extended polyp life-spans along with reduced rates of asexual

reproduction under low food levels (Stearns, 1976) may explain high plasticity of

jellyfish populations to changing environmental conditions. In the moon jellyfish,

increased food levels are associated with boosted population growth rates and dramatic

life history shifts, as expressed by longer-lived medusae with enhanced release of planula

larvae, increased ephyra production, as well as faster and more successful development of

ephyrae into medusae. Our study provides insight into the possible consequences of

habitat eutrophication by taking into account increased food availability in context with

jellyfish blooms, as investigated food levels are motivated by eutrophic to hypereutrophic

conditions in our study areas Kerteminde Fjord/Kertinge Nor and Limfjorden (cf.

Riisgård, Jensen, & Rask 2008, their Table 16.1).

The availability of food resources further drives a population density-dependent

mechanism controlling individual size (Lucas, 2001) and thus, biomass, which plays a

major role for the ecological impact of *A. aurita* (Riisgård, Barth-Jensen, & Madsen

2010; Schneider & Behrends, 1994). Size-specific traits should hence be taken into

account for future population models (cf. Bordehore et al., 2015) to quantify and evaluate

the development of jellyfish blooms from a food web perspective (cf. Schnedler-Meyer,

Kiørbøe, & Mariani 2018). Although the temporal and spatial patterns of abundance in

each life stage may vary according to jellyfish species (Lucas, Graham, & Widmer 2012;

Scorrano, Aglieri, Boero, Dawson, & Piraino 2016) and even on a population-level

(Dawson et al., 2015), our observations on the moon jellyfish are likely not unique. Field

evidence of a strong correlation between medusa density and prey abundance for two

distinct *A. aurita* populations (Dawson et al., 2015, their Figure 4d) implies that described

general trends in population stage-structure and dynamics in response to variable food

supply seem broadly applicable across the *Aurelia* species complex and other metagenic

jellyfish species.

Winter warming and jellyfish blooms

Our findings support that increased water temperature can boost the booms and busts of

jellyfish blooms (Richardson, Bakun, Hays, & Gibbons 2009; Xie, Fan, Wang, & Chen

2015). Based on recent climate analyses, rises in global ocean surface temperature by 1 to

2 °C can be expected within the next 100 years (Belkin, 2009). Our results indicate

pronounced effects of this hypothetical warming period on the development of temperate

A. aurita populations in combination with regionally observed increases in food level.

Due to seasonal changes in population composition and structure, winter warming may

cause a general trend towards more medusae and less polyps under increased food

conditions. In regions with limited food availability, winter warming can further increase

the net reproductive rate of temperate jellyfish populations. Due to enhanced strobilation

periods and increased numbers of released ephyrae per polyp (Holst, 2012; Purcell,

2005), warmer winter temperatures may shorten the generation time under low food

conditions drastically. Climate warming has previously been pointed out to benefit the

population size of temperate jellyfish species, but not tropical or boreal species (Purcell et

al., 2012). Further taking into account that a cold trigger is necessary to stimulate

strobilation in many scyphozoans (Holst, 2012; Lotan, Fine, & Ben-Hillel 1994) and that

more frequent and long-lasting heatwave events may negatively affect the population

dynamics of jellyfish (Chi, Mueller-Navarra, Hylander, Sommer, & Javidpour 2018), we

suggest winter warming may promote the blooming potential of certain temperate

jellyfish species but seems to be a less powerful driver when compared to increased food

availability.

Jellyfish life histories in a changing environment

While our matrix population model projections are based on changes in prey abundance

and the temperature regime, several additional ecological parameters including intra- and

inter-specific competition, predation, substrate availability, hydrodynamic currents and

environmental conditions are considered important constraints for the development of

jellyfish blooms (Arai, 2005; Johnson, Perry, & Burke 2001; Lucas, Graham, & Widmer

2012). Indicated by the results of our life table response experiment analysis for high

versus low food conditions, a high natural mortality risk is especially likely for the

planktonic planula stage (cf. Lucas, 2001). A transition probability of 0.001 month^{-1} from

larva to polyp was found during calibration of our presented matrix models for low and

high food conditions and matches well previous estimates for *A. aurita* (Xie, Fan, Wang,

& Chen 2015). Our resulting stable population under low food levels (λ = 1 month^{-1}) is

comparable to the local *A. aurita* jellyfish population in the semi-enclosed fjord system

Kerteminde Fjord/Kertinge Nor, which has remained unchanged over the last 25 years

(Goldstein & Riisgård, 2016; Olesen, Frandsen, & Riisgård 1994; Riisgård, Barth-Jensen,

& Madsen 2010). The rapid growth potential we show for our calibrated jellyfish

population under constantly high food (optimum) conditions however is in contrast to

rather low densities of *A. aurita* medusae observed in open coastal waters (Goldstein &

Riisgård, 2016; Möller, 1980) or in Limfjorden (Hansson, Moeslund, Kiørboe, &

Riisgård 2005; Møller & Riisgård, 2007), which implies additional factors regulating

polyp, ephyra and medusa density in less protected ecosystems. According to our matrix

projection for high food levels, a pronounced increase in polyp mortality to 0.3 day^{-1} (cf.

Xie, Fan, Wang, & Chen 2015, Table 1 therein; i.e. a survivorship of 2.3×10^{-5} month^{-1})

would decrease population growth rates only marginally from 1.63 month^{-1} to 1.44

month^{-1}. In contrast, reduced medusa survival to 0.1 month^{-1} (corresponding to a

mortality of 0.07 day^{-1}; Xie, Fan, Wang, & Chen 2015) or alternatively, a lowered

transition probability from ephyra to medusa stage to 0.0001 month^{-1}, would result in

remarkably decelerated population growth rates of $\lambda = 1.10$ month^{-1} and 1.09 month^{-1},

respectively. Above-mentioned estimates and the results of our life table response

experiment analysis suggest that besides successful metamorphosis of planula larvae,

transition from ephyra to medusa stage and survival of medusae are the most critical

parameters for the development of *A. aurita* populations in exposed coastal regions.

Removal of jellyfish and reduction of their prey availability have previously been

considered as possible management strategies for keeping box jellyfish populations in

check (Bordehore et al., 2015) which, however, may have unpredictable consequences for

the sensitive balance of marine food webs. Jellyfish constitute major parts of the diet of

some fish species (Arai, 2005; Hamilton, 2016) and can in turn be important competitors

for zooplankton prey with forage fish (Purcell, 2012). For that reason, the present work

indicates devastating effects of overfishing in context with jellyfish blooms beyond its

implications for eutrophication and climate change.

Conclusions

Our study highlights that food availability can drive and specifically shape the occurrence

of jellyfish blooms. We suggest that for most temperate scyphozoans predicted changes

in water temperature may cause less dramatic increases in jellyfish density than already

established food concentrations in several eutrophic regions around the world. The role of

natural competitors and predators in controlling jellyfish populations needs further

exploration, but our findings might serve as a promising starting point. We conclude that

especially the combination of key environmental drivers, such as habitat eutrophication,

overfishing and climate change, can promote jellyfish blooms severely. Our findings

emphasize the fundamental importance of an integrative, quantitative view on jellyfish

life histories. We believe such perspective, comprising species-, life stage- and age-

specific traits, is an inevitable prerequisite in providing visions for future management of

mass occurrences in marine ecosystems.

Authors' contributions

510   US and JG conceived the ideas and designed the methodology, JG collected the data and

511   led the writing of the manuscript; JG and US analyzed the data. Both authors contributed

512   critically to the drafts and gave final approval for publication.

Acknowledgements

We thank Kim Lundgreen for technical assistance and Daniel A. Levitis, Hans Ulrik

Riisgård, Hal Caswell, Iain Stott, Cesar Bordehore, Per Andersen, Owen Jones and Cathy

H. Lucas for helpful comments and discussions. We are grateful to the reviewers for

providing valuable feedback on the manuscript. This study was financially supported by

the Max-Planck Society (Max-Planck Odense Center on the Biodemography of Aging,

Denmark). The authors declare no conflict of interest.

Data Accessibility

523   A preprint version of this article is available from bioRxiv https://doi.org/10.1101/102814

524   (Goldstein & Steiner, 2019). Data will be made publicly accessible on the Dryad Digital

525   Repository at the time of publication.

References

Arai, M. N. (2005). Predation on pelagic coelenterates: a review. Journal of the Marine

Biological Association of the United Kingdom, 85(3), 523–536. doi:

10.1017/S0025315405011458

Arai, M. N. (2009). The potential importance of podocysts to the formation of

scyphozoan blooms: a review. In J. E. Purcell & K. A. Pitt (Eds.), *Jellyfish blooms:*

causes, consequences, and recent advances (pp. 241–246). Netherlands: Springer.

doi: 10.1007/s10750-008-9588-5

Båmstedt, U., Lane, J., & Martinussen, M. B. (1999). Bioenergetics of ephyra larvae of

the scyphozoan jellyfish *Aurelia aurita* in relation to temperature and salinity. Marine

Biology, 135(1), 89–98. doi: 10.1007/s002270050605

Belkin, I. M. (2009). Rapid warming of large marine ecosystems. Progress in

Oceanography, 81(1), 207–213. doi: 10.1016/j.pocean.2009.04.011

540 Bengtson, D. A., Leger, P., & Sorgeloos, P. (1991). Use of *Artemia* as a food source for

541   aquaculture. In R. A. Browne, P. Sorgeloos & C. A. N. Trotman (Eds.), *Artemia*

542   *Biology* (pp. 255–285). Florida, USA: CRC Press.

543 Boero, F., Bouillon, J., Gravili, C., Miglietta, M. P., Parsons, T., & Piraino, S. (2008).

544   Gelatinous plankton: irregularities rule the world (sometimes). Marine Ecology

545   Progress Series, 356, 299–310. doi: 10.3354/meps07368

546 Bordehore, C., Fuentes, V. L., Segarra, J. G., Acevedo, M., Canepa, A., & Raventós, J.

547   (2015). Use of an inverse method for time series to estimate the dynamics of and

548   management strategies for the box jellyfish *Carybdea marsupialis*. PLoS ONE, 10(9),

549   e0137272. doi: 10.1371/journal.pone.0137272

550 Breithaupt, H. (2003). Aliens on the shores. EMBO reports, 4(6), 547–550. doi:

551   10.1038/sj.embor.embor877

552 Brotz, L., Cheung, W. W., Kleisner, K., Pakhomov, E., & Pauly, D. (2012). Increasing

553   jellyfish populations: trends in large marine ecosystems. Hydrobiologia, 690(1), 3–20.

554   doi: 10.1007/s10750-012-1039-7

555 Caswell, H. (1996). Analysis of life table response experiments II. Alternative

556   parameterizations for size-and stage-structured models. Ecological Modelling, 88(1-

557   3), 73–82. doi: 10.1016/0304-3800(95)00070-4

558 Caswell, H. (2001). Matrix population models: construction, analysis, and interpretation

559   (2nd ed.). Massachusetts, USA: Sinauer Associates.

560 Chi, X., Mueller-Navarra, D. C., Hylander, S., Sommer, U., & Javidpour, J. (2018). Food

561   quality matters: Interplay among food quality, food quantity and temperature affecting

life history traits of *Aurelia aurita* (Cnidaria: Scyphozoa) polyps. Science of the Total

Environment, 656, 1280−1288. doi: 10.1016/j.scitotenv.2018.11.469

Dawson, M. N., & Hamner, W. M. (2009). A character-based analysis of the evolution of

jellyfish blooms: adaptation and exaptation. Hydrobiologia, 616(1), 193−215. doi:

10.1007/s10750-008-9591-x

Dawson, M. N., Gupta, A. S., & England, M. H. (2005). Coupled biophysical global

ocean model and molecular genetic analyses identify multiple introductions of

cryptogenic species. Proceedings of the National Academy of Sciences, 102(34),

11968−11973. doi: 10.1073/pnas.0503811102

Dawson, M. N., Cieciel, K., Decker, M. B., Hays, G. C., Lucas, C. H., & Pitt, K. A.

(2015). Population-level perspectives on global change: genetic and demographic

analyses indicate various scales, timing, and causes of scyphozoan jellyfish blooms.

Biological Invasions, 17(3), 851−867. doi: 10.1007/s10530-014-0732-z

Eckert, R., Randall, D., & Augustine, G. (1988). Animal energetics and temperature

relations. In R. Eckert & G. Augustine (Eds.), *Animal physiology, mechanisms and*

adaptions (pp. 555−605). Los Angeles, USA: WH Freeman and Company.

Fey, S. B., Siepielski, A. M., Nusslé, S., Cervantes-Yoshida, K., Hwan, J. L., Huber, E.

R., ... Carlson, S. M. (2015). Recent shifts in the occurrence, cause, and magnitude of

animal mass mortality events. Proceedings of the National Academy of Sciences,

112(4), 1083−1088. doi: 10.1073/pnas.1414894112

Frandsen, K. T., & Riisgård, H. U. (1997). Size dependent respiration and growth of

jellyfish, *Aurelia aurita*. Sarsia, 82(4), 307−312. doi:

10.1080/00364827.1997.10413659

Fu, Z., Shibata, M., Makabe, R., Ikeda, H., & Uye, S.-I. (2014). Body size reduction

under starvation, and the point of no return, in ephyrae of the moon jellyfish *Aurelia*

aurita. Marine Ecology Progress Series, 510, 255−263. doi: 10.3354/meps10799

Goldstein, J., & Riisgård, H. U. (2016). Population dynamics and factors controlling

somatic degrowth of the common jellyfish, *Aurelia aurita*, in a temperate semi-

enclosed cove (Kertinge Nor, Denmark). Marine Biology, 163(2), 33−44. doi:

10.1007/s00227-015-2802-x

Goldstein, J., & Steiner, U. K. (2019). Preprint version: Ecological drivers of jellyfish

blooms - the complex life history of a cosmopolitan medusa (*Aurelia aurita*). bioRxiv.

doi: 10.1101/102814.

Gröndahl, F. (1989). Evidence of gregarious settlement of planula larvae of the

scyphozoan *Aurelia aurita*: An experimental study. Marine Ecology Progress Series,

56, 119−125.

Hamilton, G. (2016). The secret lives of jellyfish. Nature News, 531(7595), 432−434.

doi: 10.1038/531432a

Hamner, W. M., & Dawson, M. N. (2009). A review and synthesis on the systematics and

evolution of jellyfish blooms: advantageous aggregations and adaptive assemblages.

Hydrobiologia, 616(1), 161−191. doi: 10.1007/s10750-008-9620-9

Hamner, W. M., & Jenssen, R. M. (1974). Growth, degrowth, and irreversible cell

differentiation in *Aurelia aurita*. Amercian Zoologist, 14(2), 833−849. doi:

10.1093/icb/14.2.833

Hansson, L. J., Moeslund, O., Kiørboe, T., & Riisgård, H. U. (2005). Clearance rates of

jellyfish and their potential predation impact on zooplankton and fish larvae in a

neritic ecosystem (Limfjorden, Denmark). Marine Ecology Progress Series, 304,

117–131. doi: 10.3354/meps304117

Holst, S. (2012). Effects of climate warming on strobilation and ephyra production of

North Sea scyphozoan jellyfish. Hydrobiologia, 690(1), 127–140. doi:

10.1007/s10750-012-1043-y

Holst, S., & Jarms, G. (2007). Substrate choice and settlement preferences of planula

larvae of five Scyphozoa (Cnidaria) from German Bight, North Sea. Marine Biology,

151(3), 863–871. doi: 10.1007/s00227-006-0530-y

Ishii, H., & Takagi, A. (2003). Development time of planula larvae on the oral arms of

the scyphomedusa *Aurelia aurita*. Journal of Plankton Research, 25(11), 1447–1450.

doi: 10.1093/plankt/fbg094

Ishii, H., & Watanabe, T. (2003). Experimental study of growth and asexual reproduction

in *Aurelia aurita* polyps. Sessile Organisms, 20(2), 69–73. doi: 10.4282/sosj.20.69

Johnson, D. R., Perry, H. M., & Burke, W. D. (2001). Developing jellyfish strategy

hypotheses using circulation models. Hydrobiologia, 451(1), 213–221.

Kamiyama, T. (2011). Planktonic ciliates as a food source for the scyphozoan *Aurelia*

aurita (s.l.): Feeding activity and assimilation of the polyp stage. Journal of

Experimental Marine Biology and Ecology, 407(2), 207–215. doi:

10.1016/j.jembe.2011.06.016

Keen, S. L. (1987). Recruitment of *Aurelia aurita* (Cnidaria: Scyphozoa) larvae is

position-dependent, and independent of conspecific density, within a settling surface.

Marine Ecology Progress Series, 38, 151–160.

Liu, W. C., Lo, W. T., Purcell, J. E., & Chang, H. H. (2009). Effects of temperature and

light intensity on asexual reproduction of the scyphozoan, *Aurelia aurita* (L.) in

Taiwan. Hydrobiologia, 616(1), 247−258. doi: 10.1007/s10750-008-9597-4

Lotan, A., Fine, M., & Ben-Hillel, R. (1994). Synchronization of the life cycle and

dispersal pattern of the tropical invader scyphomedusan *Rhopilema nomadica* is

temperature dependent. Marine Ecology Progress Series, 109, 59−65.

Lucas, C. H. (2001). Reproduction and life history strategies of the common jellyfish,

Aurelia aurita, in relation to its ambient environment. Hydrobiologia, 451(1-3),

229−246.

Lucas, C. H., & Dawson, M. N., (2014). What are jellyfishes and thaliaceans and why do

they bloom? In K. A. Pitt & C. H. Lucas (Eds.) *Jellyfish blooms* (pp. 9−44).

Netherlands: Springer. doi: 10.1007/978-94-007-7015-7_2

Lucas, C. H., Graham, W. M., & Widmer, C. (2012). Jellyfish Life Histories: Role of

Polyps in Forming and Maintaining Scyphomedusa Populations. Advances in Marine

Biology, 63, 133−196.

Maji, B., Bhattacharyya, J., & Pal, S. (2016). Potential effects of invasive *Pterois volitans*

in coral reefs. Letters in Biomathematics, 3(1), 119−139. doi:

10.1080/23737867.2016.1213146

Möller, H. (1980). Population dynamics of *Aurelia aurita* medusae in Kiel Bight,

Germany (FRG). Marine Biology, 60(2-3), 123−128. doi: 10.1007/BF00389155

Møller, L. F., & Riisgård, H. U. (2007). Population dynamics, growth and predation

impact of the common jellyfish *Aurelia aurita* and two hydromedusae (*Sarsia*

tubulosa and *Aequorea vitrina*) in Limfjorden (Denmark). Marine Ecology Progress

Series, 346, 153−165. doi: 10.3354/meps06960

Nielsen, A. S., Pedersen, A. W., & Riisgård, H. U. (1997). Implications of density driven

currents for interaction between jellyfish (*Aurelia aurita*) and zooplankton in a Danish

fjord. Sarsia, 82(4), 297−305. doi: 10.1080/00364827.1997.10413658

Olesen, N. J., Frandsen, K., & Riisgård, H. U. (1994). Population dynamics, growth and

energetics of jellyfish, *Aurelia aurita*, in a shallow fjord. Marine Ecology Progress

Series, 105, 9−18.

Pitt, K. A., Welsh, D. T., & Condon, R. H. (2009). Influence of jellyfish blooms on

carbon, nitrogen and phosphorus cycling and plankton production. Hydrobiologia,

616(1), 133−149. doi: 10.1007/s10750-008-9584-9

Purcell, J. E. (2005). Climate effects on formation of jellyfish and ctenophore blooms: a

review. Journal of the Marine Biological Association of the United Kingdom, 85(3),

461−476. doi: 10.1017/S0025315405011409

Purcell, J. E. (2012). Jellyfish and ctenophore blooms coincide with human proliferations

and environmental perturbations. Annual Review of Marine Science, 4, 209−235. doi:

10.1146/annurev-marine-120709-142751

Purcell, J. E., Uye, S.-I., & Lo, W. T. (2007). Anthropogenic causes of jellyfish blooms

and their direct consequences for humans: a review. Marine Ecology Progress Series,

350, 153−174. doi: 10.3354/meps07093

Purcell, J. E., Atienza, D., Fuentes, V. L., Olariaga, A., Tilves, U., Colahan, C., & Gili J.

M. (2012). Temperature effects on asexual reproduction rates of scyphozoan species

from the northwest Mediterranean Sea. Hydrobiologia, 690(1), 169−180. doi:

10.1007/s10750-012-1047-7

R Core Team (2015). R: A language and environment for statistical computing. R

Foundation for Statistical Computing, Vienna, Austria. http://www.r-project.org

Richardson, A. J., Bakun, A., Hays, G. C., & Gibbons, M. J. (2009). The jellyfish joyride:

causes, consequences and management responses to a more gelatinous future. Trends

in Ecology & Evolution, 24(6), 312−322. doi: 10.1016/j.tree.2009.01.010

Riisgård, H. U., Jensen, M. H., & Rask, N. (2008). Odense Fjord and Kerteminde

Fjord/Kertinge Nor. In U. Schiewer (Ed.), *Ecology of Baltic Coastal Waters* (pp.

361−394). Germany: Springer.

Riisgård, H. U., Barth-Jensen, C., & Madsen, C. V. (2010). High abundance of the

jellyfish *Aurelia aurita* excludes the invasive ctenophore *Mnemiopsis leidyi* to

establish in a shallow cove (Kertinge Nor, Denmark). Aquatic Invasions, 5(4),

347−356. doi: 10.3391/ai.2010.5.4

Riisgård, H. U., Andersen, P., & Hoffmann, E. (2012). From fish to jellyfish in the

eutrophicated Limfjorden (Denmark). Estuaries and Coasts, 35(3), 701−713. doi:

10.1007/s12237-012-9480-4

Scorrano, S., Aglieri, G., Boero, F., Dawson, M.N., & Piraino, S. (2016). Unmasking

Aurelia species in the Mediterranean Sea: an integrative morphometric and molecular

approach. Zoological Journal of the Linnean Society, 180(2), 243−267. doi:

10.1111/zoj.12494

Schneider, G., & Behrends, G. (1994). Population dynamics and the trophic role of

Aurelia aurita medusae in the Kiel Bight and western Baltic. ICES Journal of Marine

Science, 51(4), 359−367.

Schneider, G., & Behrends, G. (1998). Top-down control in a neritic plankton system by

Aurelia aurita medusae – a summary. Ophelia, 48(2), 71−82. doi:

10.1080/00785236.1998.10428677

Schnedler-Meyer, N. A., Kiørboe, T., & Mariani, P. (2018). Boom and Bust: Life

History, Environmental Noise, and the (un)Predictability of Jellyfish Blooms.

Frontiers in Marine Science, 5, 257. doi: 10.3389/fmars.2018.00257

Schrope, M. (2012). Marine ecology: attack of the blobs. Nature, 482(7383), 20−21.

Stearns, S. C. (1976). Life-history tactics: a review of the ideas. The Quarterly Review of

Biology, 51(1), 3−47. doi: 10.1086/409052

Stott, I., Hodgson, D. J., & Townley, S. (2012). popdemo: an R package for population

demography using projection matrix analysis. Methods in Ecology and Evolution,

3(5), 797−802. doi: 10.1111/j.2041-210X.2012.00222.x

Stubben, C. J., & Milligan, B. G. (2007). Estimating and Analyzing Demographic Models

Using the popbio Package in R. Journal of Statistical Software, 22(11), 1−23. doi:

10.18637/jss.v022.i11

Tuljapurkar, S., & Caswell, H. (1997). Structured-population models in marine,

terrestrial, and freshwater systems. New York, NY: Chapman & Hall.

Vanhaecke, P., & Sorgeloos, P. (1980). International Study on *Artemia*. IV. The

biometrics of *Artemia* strains from different geographical origin. The brine shrimp

Artemia, 3, 393−405.

Xie, C., Fan, M., Wang, X., & Chen, M. (2015) Dynamic model for life history of

scyphozoa. PLoS ONE, 10(6), e0130669. doi: 10.1371/journal.pone.0130669

Supporting Information

Additional Supporting Information is available online in the supporting information tab

for this article.

Tables

Table 1. Experimental conditions for life stages of the jellyfish *Aurelia aurita* (L: planula

larva, P: polyp, E: ephyra, M: medusa) under low (*L*) and high (*H*) food conditions. l_0:

initial number of individuals per treatment, t_0, Δt: start and duration of experiment, *c*:

food concentration based on the carbon content of available prey organisms (*Artemia*

salina).

Food treatment	Life stage	l_0 (ind.)	t_0 (month)	Δt (d)	c (μg C L^{-1} d^{-1})
L	L	960	Sept	52	0
	P	100	Aug	364	1
	E	33	Mar	153	10
	M	16	Apr	296	-
H	L	960	Jun	48	0
	P	100	Aug	364	10
	E	33	Jan	158	100
	M	33	Feb	391	>1000

731 Table 2. Demographic parameters of yearly stage-structured projection matrices (P: polyp, E: ephyra, M: medusa) for a local *Aurelia*

732 *aurita* population under low (*L*) and high food conditions (*H*) at present water temperatures (PTW) and for a predicted winter warming

733 trend (WW).

Demographic parameter	*L*		*H*	
	PWT	WW	PWT	WW
Scaled population growth rate λ (month^{-1})	1.00	1.01	1.63	1.73
Damping ratio ρ (months)	59	59.2	3	3.2
Stable stage proportion (P, E, M)	0.94, 0.04, 0.02	0.84, 0.12, 0.04	0.76, 0.04, 0.20	0.60, 0.02, 0.38
Scaled reproductive value R (ind.) (P, E, M)	1, 0.03, 0.42	1, 8.53, 3×10^{-7}	1, 0.27, 9.13	1, 0.18, 7.43
Net reproductive rate R_0 (ind.)	1.0	1.8	57.0	80.9
Generation time T (months)	200.0	111.8	8.3	8.0
Life expectancy e_0 (months) (P, E, M)	394.5, 1.7, 3.1	394.5, 2.1, 3.1	86.7, 2.9, 8.0	86.7, 3.5, 8.0

Figures

Figure 1. Life history of the moon jellyfish *A. aurita* (Scyphozoa, Cnidaria). (a)

Metagenic life cycle with alternation of generations between the sexually reproducing

medusa and the asexually reproducing polyp stage. (b) Life cycle graph for *A. aurita*,

including transition probability *t*, fecundity *F* (solid arrows) and mortality † (broken

arrows) of all life stages, i.e. planula larva (L), polyp (P), ephyra (E) and medusa (M). (c)

Reduced life cycle graph for polyps, ephyrae and medusae considering fast (daily)

transition from larva to polyp stage.

Figure 2. Population matrix model for *A. aurita* under low (upper panels) and high (lower panels) food conditions (~1:10) at present water temperatures (PWT; black) and in a winter warming scenario (WW; grey). (a−b) Matrix projection over a 10-year period, standardized to the long-term effects of population growth rates λ = 1 month^{-1} and λ = 1.63 month^{-1}, respectively. (c−d) Stable stage distribution (i.e. relative, constant long-term proportions) of polyps (P; solid lines), ephyrae (E; dotted lines) and medusae (M; broken lines).

Figure 3. Sensitivity of the population growth rate λ of *A. aurita* to changes in monthly stage-

specific transitions (1: $t_{PP}+F_{PP}$, 2: F_{PE}, 3: $F_{PE}×t_{EM}$, 4: t_{EE}, 5: t_{EM}, 6: $F_{ML}×t_{LP}$, 7: t_{MM}) under (a) low

and (b) high food conditions. (c) Life-table response experiment (LTRE) analysis for sensitivity

of λ to perturbations in the population projection matrix elements when comparing high to low

food conditions (~10:1).