Ecological drivers of jellyfish blooms – the complex life history

of a 'well-known' medusa (*Aurelia aurita*)

8   J. Goldstein[* a, b], U.K. Steiner [b, c]

10   [a]Marine Biological Research Centre, University of Southern Denmark, Hindsholmvej 11, 5300

11   Kerteminde, Denmark

12   [b]Department of Biology, University of Southern Denmark, Campusvej 55, 5230 Odense,

13   Denmark

14   [c]Center on Population Dynamics, University of Southern Denmark, Campusvej 55, 5230 Odense,

15   Denmark

17   [*]Corresponding author: jgoldstein@biology.sdu.dk

21   **Running head** Drivers of jellyfish blooms

Summary

1. Jellyfish blooms are conspicuous demographic events with significant ecological and socio-economic impact. Despite worldwide concern about an increased frequency and intensity of such mass occurrences, predicting their booms and busts remains challenging.

2. Forecasting how jellyfish populations may respond to environmental change requires taking into account their complex life histories. Metagenic life cycles, which include a benthic polyp stage, can boost jellyfish mass occurrences via asexual recruitment of pelagic medusae.

3. Here we present stage-structured matrix population models with monthly, individual-based demographic rates of all life stages of the moon jellyfish *Aurelia aurita* (sensu stricto). We investigate the life stage-dynamics of these complex populations under low and high food conditions to illustrate how changes in medusa density depend on non-medusa stage dynamics.

4. We show that increased food availability can be an important ecological driver of jellyfish mass occurrences, as it can temporarily shift the population structure from polyp- to medusa-dominated. Projecting populations for a winter warming scenario additionally enhanced the booms and busts of jellyfish blooms.

5. We identify demographic key variables that control the intensity and frequency of

jellyfish blooms in response to environmental drivers such as habitat

eutrophication and climate change. By contributing to an improved understanding

of mass occurrence phenomena, our findings provide perspective for future

management of ecosystem health.

Key-words biodemography, environmental change, jellyfish blooms, matrix model,

population structure

Introduction

The biodiversity, productivity and resilience of marine ecosystems worldwide are

threatened by an imbalance in the ecological parameters regulating population outbreaks.

Mass occurrences of micro- and macroalgae, cnidarians, ctenophores, molluscs,

echinoderms, tunicates and fish appear with increasing frequency and intensity at a global

scale in recent decades (Breithaupt, 2003; Lucas & Dawson, 2014; Maji, Bhattacharyya,

& Pal 2016). Despite mass occurrences and mass mortality events are widespread – and

can provide major implications for marine ecosystem health – predictions have remained

challenging due to an incomplete understanding of underlying demographic processes

(Fey et al., 2015). Jellyfish blooms are a prominent example for mass occurrences in

which the population density of one life stage, the pelagic medusa, shows distinct peaks

over broad temporal and regional scales (Lucas, Graham, & Widmer 2012). These

demographic events may have lasting ecological, economic and social implications (cf.

Lucas & Dawson, 2014). Jellyfish blooms can trigger trophic cascades such as collapse of

pelagic food webs due to their severe predatory impact on a broad spectrum of

zooplankton species and fish larvae (Richardson, Bakun, Hays, & Gibbons 2009;

Riisgård, Andersen, & Hoffmann 2012; Schneider & Behrends, 1994, 1998). Further,

jellyfish mass occurrences can interfere with human activities including tourism,

fisheries, aquaculture and power production (Purcell, 2012; Purcell, Uye, & Lo 2007).

Jellyfish populations, i.e. local aggregations of the medusa stage, seem to be increasing in

numerous coastal ecosystems around the world (Brotz, Cheung, Kleisner, Pakhomov, &

Pauly 2012). Available evidence suggests that jellyfish outbreaks which are defined as

exceptional, sudden or 'unnatural' increases in biomass (Lucas & Dawson, 2014), may be

favored by anthropogenic influences such as overfishing, eutrophication, habitat

modification, species translocation and climate change (Purcell, 2012; Richardson,

Bakun, Hays, & Gibbons 2009).

Bloom-forming jellyfish species are primarily found within the cnidarian taxon

Scyphozoa and can feature complex life histories (Dawson & Hamner, 2009; Hamner &

Dawson, 2009; Lucas & Dawson, 2014). Their metagenic life cycles typically include a

pelagic, sexually reproducing medusa and a benthic, asexually reproducing polyp stage

(Lucas, Graham, & Widmer 2012). Such life cycles are characterized by high stage-

specific plasticity to changing environmental conditions (e.g. Gröndahl, 1989; Hamner &

Jenssen, 1974; Liu, Lo, Purcell, & Chang 2009). Although the development of medusa

populations is directly linked to survival rates of several precedent life stages, i.e. planula

larvae, polyps and ephyrae (Lucas, 2001), information on the demographic rates of

bloom-forming jellyfish has remained scarce (Xie, Fan, Wang, & Chen 2015).

Quantifying demographic rates of all stages in the life cycle of species showing mass

occurrence is thus fundamental to identifying the environmental variables causing

changes in population dynamics and structure.

Here, we explore the demographic structure and environmental drivers underlying

mass occurrence of the moon jellyfish *Aurelia aurita* (sensu stricto). We quantify

demographic rates of all life stages of *A. aurita* (larvae, polyps, ephyrae and medusae),

i.e. survival, stage-transitions and fecundity, based on experimental and field data from

two temperate Danish water systems with contrasting prey availability. Using these

demographic rates, we parameterize stage-structured population models to project the life

cycle of jellyfish populations for low and high food levels (i.e. zooplankton prey

biomasses). Since warm temperature has been associated with increased numbers of

medusae for most temperate scyphozoan species (Purcell, 2005), we further predict the

impact of a winter warming scenario on the population dynamics of *A. aurita* by

perturbing demographic rates. Our results provide novel insights into the development of

metagenic jellyfish populations under increased food levels and increased winter

temperatures. Such information is essential to assess the boosting potential of

environmental drivers on species showing mass occurrence and provides a milestone for

identifying and managing demographic irregularities in marine ecosystems.

Materials and Methods

Study organism

The moon jellyfish *Aurelia* (Scyphozoa, Cnidaria) belongs to an ubiquitous, well-studied

genus with at least 16 species, most of which are distributed in specific coastal and shelf

sea regions around the world (Dawson, Gupta, & England 2005). The life history, i.e.

individual stage-trajectory, of *Aurelia aurita*, sensu stricto, is characterized by a complex,

metagenic life cycle which involves sexually reproducing medusae and asexually

reproducing polyps (Figure 1a). Medusae of *A. aurita* are pelagic filter feeders which

differentiate into distinct male and female individuals upon sexual maturation (Lucas

2001). After fertilization of oocytes, female medusae produce large numbers of

lecithotrophic (yolk-feeding) planula larvae (Ishii & Takagi, 2003), which spend 12 h to 1

week in the water column (Lucas, 2001) before they settle on suitable hard substrate and

metamorphose into polyps (Holst & Jarms, 2007; Keen, 1987). The inconspicuous

benthic polyp stage of *A. aurita* can occur in densities between 60,000 and 400,000 ind.

m^{-2} (Lucas, 2001) and reproduces asexually by several modes, including i) the production

of well protected chitin-covered resting stages called podocysts (Arai, 2009), ii) budding

of new polyps (Ishii & Watanabe, 2003), and iii) the release of multiple immature

medusae, i.e. ephyrae, through transverse fission (strobilation). After asexual

propagation, polyps remain in the polyp stage, while the development of pelagic ephyrae

into sexually mature medusae closes the life cycle (Lucas, Graham, & Widmer 2012).

Data collection

We cultivated planula larvae (L), polyps (P), ephyrae (E) and medusae (M) of *A. aurita* in

the laboratories of the Marine Biological Research Centre, Kerteminde (Denmark)

between June 2013 and November 2014 (Table 1). We set up separate experimental

series for each life stage using filtered (38 µm) seawater at constant salinity (20 PSU) and

with water temperature following seasonal variability of the local temperate climate zone

(5-22 °C). We fed all life stages, except the planula larvae, with low (L) or high (H)

amounts (corresponding to a ratio ~1:10) of two-day old *Artemia salina* nauplii and

regularly exchanged the seawater in our cultivation containers. We adjusted the fed

biomass of *A. salina* (2.55 µg C ind.$^{-1}$; Vanhaecke & Sorgeloos, 1980) to the size-specific

energy demands of each life stage. In the following, we describe in more detail the origin

and experimental setup for each investigated life stage.

We obtained planula larvae from female medusae that were collected in two

contrasting Danish water systems – the fjord system Kerteminde Fjord/Kertinge Nor and

the major water system Limfjorden. Zooplankton prey biomasses are in a range $<1-100$

µg C L^{-1} in Kerteminde Fjord/Kertinge Nor (Nielsen, Pedersen, & Riisgård 1997) and

$>10-1000$ µg C L^{-1} in Limfjorden, respectively (Møller & Riisgård, 2007). Directly after

the release of planulae, we transferred the larvae individually to multi-well seawater

containers (200 µL). During daily counts, we determined the number of living larvae (l_{LL},

ind. day^{-1}) and the number of larvae that had completed metamorphosis into polyps (l_{LP},

ind. day^{-1}) until all planulae had metamorphosed into polyps or had died.

We cultivated polyps of *A. aurita* from planulae, released by female medusae from

Kertinge Nor, which we initially added to small seawater containers (50 mL) with

floating polystyrene substrate plates to facilitate settlement. After 24 h, we removed all

but one newly settled polyp per substrate plate and replaced the seawater in the

containers. We subsequently raised and cultivated the individual polyps under low or

high food conditions (Table 1; cf. Kamiyama, 2011). To maintain the initial polyp

cohorts, we instantly removed any buds of the "mother" polyps from substrate plates.

During the whole experimental period of 12 months, no excystment of polyps from

podocysts was observed. Each week we counted the number of live polyps (l_{PP}, ind.

week^{-1}) as well as the number of buds (N_{PP}, ind. week^{-1}) and ephyrae per polyp (N_{PE}, ind.

week^{-1}).

Ephyrae which were released by polyps under low and high food conditions were

individually transferred to seawater flasks (250 mL) with constant water movement and

air supply, and continuously reared under either low or high food conditions. Food levels

for ephyrae corresponded to the minimum energy demands for maintenance of a

'standard' ephyra with 8 mm inter-rhopalia diameter (Båmstedt, Lane, & Martinussen

1999; Bengtson, Leger, & Sorgeloos 1991; Eckert, Randall, & Augustine 1988; Frandsen

& Riisgård, 1997), or the 10-fold carbon load, respectively (Table 1). Each week we

counted the number of surviving ephyrae (l_{EE}, ind. week^{-1}) and the number of ephyrae

that had completed transition to medusa (l_{EM}, ind. week^{-1}); the latter we defined by the

presence of fully developed intermediate lappets and the final closure of adradial clefts.

We performed these counts until all ephyrae had either developed into medusae or had

died.

We continuously tracked all newly transitioned medusae and additional sexually

mature females collected in Kertinge Nor, in individual aquaria (1 L) with constant air

supply. We kept the medusae under low or high food levels which resembled complete

starvation or the 10-fold energy demands of a 'standard' 50 mm umbrella diameter

medusa (Table 1; Olesen, Frandsen, & Riisgård 1994; Goldstein & Riisgård, 2016). We

recorded the survival of medusae (l_{MM}, ind. day^{-1}) daily until all individuals in the low

food treatment had died; at that time all medusae in the high food treatment were still

alive. We estimated the daily release of planula larvae by female medusae during the

reproductive period (N_{ML}, ind. day^{-1}) using the field observation-based relationship

$N_{ML} = 0.087 \times 160.8 \times \mathrm{e}^{0.029d}$ (Goldstein & Riisgård, 2016), where average umbrella

diameters (d, mm) of 40 mm and 200 mm reflected low and high food conditions.

Estimating stage-specific demographic rates

We used the experimental data to estimate survival rates, stage transition rates and

fecundity of the stages L, P, E and M between age x and age x+1 for discrete time steps of

one month. Individuals in each life stage i have certain monthly probabilities to survive in

their current stage (t_{ii}), to transition to another stage j (t_{ij}) and to produce offspring of

stage j (F_{ij}). Each stage has its unique survival rate (t_{LL}, t_{PP}, t_{EE}, t_{MM}, month^{-1}) and stage

transitions are possible from larva to polyp (t_{LP}, month^{-1}) or from ephyra to medusa (t_{EM},

month^{-1}). Fecundity includes asexual reproduction of polyps by budding (F_{PP}, ind. ind.$^{-1}$

month^{-1}) or release of ephyrae (F_{PE}, ind. ind.$^{-1}$ month^{-1}), and sexual reproduction of

medusae by the release of planula larvae (F_{ML}, ind. ind.$^{-1}$ month^{-1}; Figure 1b). Details on

estimating monthly stage-specific demographic rates of *A. aurita* are available from the

Supporting Information (Appendix S1 & S2).

201   We first estimated monthly transition probabilities and fecundities for each life-stage i

under low and high food conditions (Appendix S1) based on count ratios from our

empirical data with either daily (L, M) or weekly resolution (P, E). For converting the

daily release of planula larvae into monthly fecundity of medusae, we assumed a sex ratio

of 1:1. Since transition probabilities and fecundities between subsequent time steps

depend on survivorship within each monthly time step, we assumed that individuals that

died during a given month, had died halfway during this month, i.e. they survived

throughout half the month on average (Appendix S2). This assumption is needed as, for

instance, a medusa might still release larvae before its death within a given month.

Further, some ephyrae completed transition to the medusa stage within less than a month

after they were released by a polyp. We therefore corrected the monthly release of

ephyrae by polyps not only for daily survivorship of ephyrae, but also for the transition

probability to the medusa stage within a given month.

Stage-structured matrix population models

Our matrix population models describe the transitions of each life stage taking place

during each projection interval, as defined by monthly stage-specific demographic rates

for each of the 12 month of a year. Age is determined by month, and the final stages are

P, E and M. Since larval transition to the polyp stage was restricted to days or hours ($t_{LP} =$

0 for incompetent larvae with $t_{LL} > 0$), we linked the monthly transition from larva to

polyp stage to fecundity of medusae ($F_{ML} \times t_{LP}$) to build the basic structure of an

irreducible monthly stage-structured matrix model (Figure 1c). Using our empirical

datasets on all demographic rates of *A. aurita* under low and high food conditions,

respectively, we constructed monthly stage-structured submatrices of the following

structure:

$$
\mathbf{A}_x = \begin{pmatrix} t_{PP} + F_{PP} & 0 & F_{ML} \times t_{LP} \\ F_{PE} & t_{EE} & 0 \\ F_{PE} \times t_{EM} & t_{EM} & t_{MM} \end{pmatrix},
$$

227   where x denotes the age in months. Based on 12 monthly submatrices A_x (3×3) which

228   reflect seasonal life cycle dynamics of temperate *A. aurita* populations (Appendix S3:

229   Figures S1 & S2), we constructed yearly population matrices A (36×36; time step = 1

230   month) for monthly projections over a 10-year interval (120 iterations):

$$A = \begin{pmatrix} \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & A_{12} \\ A_1 & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & A_2 & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & A_3 & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & A_4 & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & A_5 & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & A_6 & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & A_7 & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & A_8 & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & A_9 & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & A_{10} & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & A_{11} & \cdot \end{pmatrix}$$

.

In some months, we observed no survivorship of ephyrae or medusae, which would lead

to reducible matrices (Caswell, 2001); in these cases, we included a small transition

probability $c = 1 \times 10^{-10}$ as placeholder (Appendix S3: Figures S1 & S2). Our results

were robust to this assumption.

Adjusting matrix parameters to environmental conditions

We constructed yearly projection matrix models for the jellyfish population under low

and high food conditions. Since our transition probabilities from larva to polyp were

unrealistically high due to the sheltered laboratory environment (as an important

microzooplankton component, planula larvae are subject to a high mortality risk under

natural conditions), we calibrated the larva to polyp transition to a level ($t_{LP} = 0.001$

month^{-1}) that resulted in a stationary population (i.e. characterized by constant population

size over time) under low food conditions. We used the same t_{LP} as for the calibrated low

food condition to scale the projection matrix model for high food conditions, leaving us

with a rapidly growing population.

We parameterized additional projection matrix models to explore the potential effects

of a winter warming scenario under low and high food conditions. A simulated increase

in present water temperatures from 5 to 10 °C is predicted to benefit the production of *A.*

aurita ephyrae in terms of extended strobilation periods and enhanced ephyra release per

polyp (Holst, 2012). Our winter warming scenario included 1.8-fold increased ephyra

production (i.e. $F_{PE}\times1.8$) during strobilation periods that we extended by one month, i.e.

mean F_{PE} of proceeding and subsequent month. We additionally adjusted corresponding

probabilities for ephyra survival and transition from ephyra to medusa to t_{EE} and t_{EM} of

subsequent month, respectively (Appendix S3: Figures S1 & S2). With these modified

matrices, we investigated the demographic effects of a predicted winter warming trend on

A. aurita populations under low and high food conditions.

Computing demographic parameters and population projection

We analyzed our yearly population projection matrices using R version 3.1.3 (R Core

Team, 2015) and compared the demographic dynamics of *A. aurita* under low and high

food levels, and in combination with a winter warming trend. We used R packages

'popbio' (Stubben & Milligan, 2007) and 'popdemo' (Stott, Hodgson, & Townley 2012)

to compute the following population-level metrics: population growth rate λ (dominant

eigenvalue), stable stage distribution (right eigenvector **w** corresponding to the dominant

eigenvalue; i.e. proportions of individuals in each stage remain constant on the long-

term), reproductive value R (left eigenvector $\mathbf{v}$ corresponding to the dominant eigenvalue;

i.e. expected lifetime reproduction of an individual in stage i), net reproductive rate R_0

(i.e. average number of offspring by which an individual will be replaced during its

lifetime), generation time T (i.e. average age at which an individual has produced the

average number of offspring, R_0), life expectancy e_0 (i.e. mean time to death), and

sensitivity of λ to perturbations of the matrix elements (only elements >0 are considered).

We further performed a fixed life table response experiment (LTRE) analysis (Caswell,

1996; Tuljapurkar & Caswell, 1997) for pairwise comparison of the sensitivity of λ to

changes in stage-specific transitions under high versus low food conditions.

Results

Demographic dynamics of jellyfish under low and high food conditions

The metagenic life cycle of *A. aurita* (Figure 1a) was reflected by the cyclic (imprimitive)

structure of our matrix model, expressing annual peak and pit dynamics in projected

population density (Figure 2a). Relative to the calibrated stationary stable population with

$\lambda = 1$ month^{-1} under low food conditions, the scaled population growth rate for 10-fold

increased food availability was $\lambda = 1.63$ month^{-1}, corresponding to an approximated

increase in population size by a factor ($\lambda^{12} =$) 346 year^{-1} (Table 2). Such differences in

growth rate due to changes in food availability provide high leverage for population

booms and busts. Similar to low food levels, the projection matrix under high food

availability showed cyclic annual structure with typical peak and pit dynamics in

projected population density, but life cycle amplitudes were much increased (Figure 2b).

Under low food conditions, the overall population density (taking into account all life

stages) increased to a peak during spring and summer and thereafter decreased to low

densities during autumn and winter. Within-year dynamics under high food levels

differed significantly compared to low food conditions, showing two distinct maxima in

population density, one during spring and one during autumn, each followed by a

subsequent minimum in summer and winter (Figures 2a–b). The convergence time to

stable stage (i.e. the damping ratio ρ) of our modelled *A. aurita* population was ~5 years

under low food conditions and reduced by a factor 21, for the high food population

(Table 2).

The stable stage distribution under low food availability was vastly dominated by

polyps (0.94) that survived throughout the year. Ephyrae and medusae accounted for

comparatively small fractions of the population (0.04 and 0.02, respectively) and showed

typical seasonal patterns. Ephyrae were released during April/May and survived into

September. Only few ephyrae transitioned to the medusa stage under low food

conditions, so that medusae were present from June to December and contributed only

marginally to restocking polyp densities by the release of planula larvae during autumn

and winter (Figure 2c). Under high food supply, less polyps (0.76) dominated the overall

A. aurita population and the stable stage distribution was characterized by a 10-fold

increased medusa fraction (0.2) compared to low food conditions (Table 2). Compared to

a year-round standing stock of polyps under low food conditions, the stable stage

proportion of polyps under high food levels showed a peak in autumn after being

restocked by the release of planula larvae, and subsequently decreased throughout the

year. Ephyrae were released in March/April, and the medusa fraction dominated the *A.*

aurita population from April to July under high food levels (Figure 2d).

The reproductive value was low for ephyrae (0.03) under low food conditions, and

sexual reproduction of medusae contributed only about half as much (0.42) to future

generations as the asexually reproducing polyp stage (1; note scaling to 1 for polyps in

January). Under high food conditions, the reproductive value of ephyrae and medusae

increased by a factor 8.1 and 21.6, respectively (Table 2). A net reproductive rate of 1

indicated that each polyp is replaced by 1 new polyp at the end of its life under low food

conditions which resulted in long generation times of ~16.7 years. High food levels

increased the net reproductive rate to 57 polyps replacing each polyp at the end of its life

and boosted the average time between two consecutive generations to 8 months. Life

expectancy of polyps, estimated as the mean time to death, was 33 years under low food

conditions, while ephyrae and medusae revealed much shorter life expectancies of 2 and

3 months, respectively. In consensus with a more pronounced impact of the medusa stage

on population structure under high food levels, life expectancy of polyps decreased to 7

years, while the length of ephyra and especially medusa lives extended to 3 and 8 months,

respectively (Table 2).

Sensitivity of demographic rates to increased food availability

Sensitivity analysis generally highlighted a strong influence of survival rates and less

pronounced effects of fecundity on the growth of jellyfish populations. Under low food

conditions, population growth was most sensitive to changes in polyp survival ($t_{PP}+F_{PP}$),

expressed by constant sensitivity 0.08 (relative maximum value in the sensitivity matrix)

throughout the months of the year (Figure 3a). Comparatively, λ showed a much lower

sensitivity of 0.02 to changes in ephyra production (F_{PE}) from March to May. The overall

sensitivity of λ to perturbations in the projection matrix elements increased under high

compared to low food levels. Under high food conditions, changes in polyp survival

($t_{PP}+F_{PP}$) affected λ particularly from October to April, with sensitivities of 0.06 to 0.13.

In contrast to low food levels, however, λ was most sensitive to changes in medusa

survival (t_{MM}) under high food levels. Sensitivity to t_{MM} remained high throughout the

year, ranging from 0.07 to 0.16 from April to October, and reaching a maximum of 0.32

from June to July (Figure 3b). We further detected sensitivities of λ in the range 0.02 to

0.03 for perturbations in ephyra production (F_{PE}) from March to May under high food

conditions. Differences in the sensitivity of λ between food treatments were also evident

from the results of our life table response experiment (LTRE) analysis. High food

availability led to 7.7 to 12.6 % increased sensitivity for the transition probabilities from

polyp to medusa stage (i.e. $F_{PE}+t_{EM}$) from March to May. Further, increased food

conditions enhanced the sensitivity of λ for medusa survival (i.e. t_{MM}) during April by 7.7

% and for polyp production via sexual reproduction of medusae (i.e. $F_{ML}+t_{LP}$) from

August to December by 5.0 to 6.8 % (Figure 3c).

Demographic dynamics of jellyfish in a winter warming scenario

For our winter warming scenario with extended strobilation periods and increased ephyra

release, population growth rates of *A. aurita* increased to 1.01 and 1.73 month^{-1} under

low and high food conditions, corresponding to 1.1- and 729-fold yearly increases in

population size (i.e., relative increases by 6.2 % and 110.9 %), respectively. Winter

warming led to stronger variability in inter-annual population structure in both food

treatments, expressed by more distinct peaks and pits in population density (Figures

2a–b). Under low food levels, the stable population structure of *A. aurita* consisted of 2.7

% more ephyrae, 7.9 % more medusae, and 10.6 % less polyps per year as a consequence

of winter warming (Figure 2c). Under high food levels, increased winter temperatures

resulted in a stable stage distribution with 17.4 % more medusae, while the proportions of

ephyrae and polyps decreased by 1.7 % and 15.7 %, respectively (Table 2). An earlier

onset of strobilation periods in the scenario and fast transition of ephyrae caused

significant proportions of medusae to be already present from February (Figure 2d).

Winter warming caused substantial changes in reproductive rates and generation times

of the projected jellyfish populations; these changes were particularly pronounced under

low food conditions. The reproductive value of ephyrae in the low food population was

257-fold increased, while the relative contribution to the number of offspring by medusae

decreased to near zero. Winter warming in combination with high food levels decreased

reproductive values of both ephyrae and medusae by a factor 1.5 and 1.2, respectively

(Table 2). Increased winter temperatures raised the net reproductive rate by 75.1 % under

low food conditions and by 42.0 % under high food conditions. The winter warming

scenario shortened the generation time by 7.4 years under low food conditions, while

generation times were shortened by only 9 days under high food conditions. Winter

warming increased the life expectancy of ephyrae by 12 and 18 days, respectively,

whereas life expectancy of polyps and medusae remained unchanged under both low and

high food levels (Table 2).

Discussion

We reveal the demographic dynamics and life history shifts underlying jellyfish mass occurrence and show that variable intensity and frequency is subject to specific ecological triggers. Our matrix projections quantify how jellyfish blooms are enhanced by naturally occurring variations in the food regime. We suggest the observed temporal shift from polyp- to medusa-dominated populations in response to increased food availability as a key mechanism causing jellyfish outbreaks, which can be boosted by the combined effects with a predicted winter warming. Our findings support previous studies which have pointed out metagenic life cycles as an ultimate driver of boom and bust population dynamics (Brotz, Cheung, Kleisner, Pakhomov, & Pauly 2012; Hamner & Dawson, 2009; Lucas & Dawson, 2014, their Figure 2.4b; Pitt, Welsh, & Condon 2009). Considering jellyfish outbreaks as highly sensitive indicators for ocean degradation (Schrope, 2012), the present insights may provide fundamental knowledge regarding the health of marine ecosystems worldwide.

Food availability as a key driver of jellyfish blooms

Our study provides insight into probable consequences of habitat eutrophication in context with jellyfish blooms, as investigated food levels mirror eutrophic to hypereutrophic conditions prevailing in several Danish coastal regions (Riisgård, Jensen, & Rask 2008, their Table 16.1). Resulting demographic dynamics suggest that food availability is a major constraint for the intensity and frequency of jellyfish blooms, confirming previous observations of food availability as a key variable in limiting the development of ephyrae (Fu, Shibata, Makabe, Ikeda, & Uye 2014) and the survivorship

and fecundity of medusae (Goldstein & Riisgård, 2016). Yet, to our knowledge, we

present the first comprehensive quantitative model of how increased food availability can

trigger demographic changes in a scyphozoan jellyfish population. Such changes include

enhanced boom and bust dynamics and shorter generation times. We show that complex

life cycles can facilitate jellyfish blooms and emphasize the key role of benthic polyps in

ensuring long-term survival (Boero et al., 2008; Lucas, Graham, & Widmer 2012),

particularly when food is scarce. Life-history trade-offs such as extended polyp life-spans

along with reduced rates of asexual reproduction under low food levels (Stearns, 1976)

may explain high plasticity of jellyfish populations to changing environmental conditions.

In the moon jellyfish, increased food levels are associated with boosted population

growth rates and dramatic life history shifts, as expressed by longer-lived medusae with

enhanced release of planula larvae, increased ephyra production, as well as faster and

more successful development of ephyrae into medusae.

The availability of food resources further drives a population density-dependent

mechanism controlling individual size (Lucas, 2001) and thus, biomass, which plays a

major role for the ecological impact of *A. aurita* (Riisgård, Barth-Jensen, & Madsen

2010; Schneider & Behrends, 1994). Size-specific traits should hence be taken into

account for future population models (cf. Bordehore et al., 2015) to quantify and evaluate

the development of jellyfish blooms from a food web perspective (cf. Schnedler-Meyer,

Kiørbøe, & Mariani 2018). Although the temporal and spatial patterns of abundance in

each life stage may vary according to jellyfish species (Lucas, Graham, & Widmer 2012;

Scorrano, Aglieri, Boero, Dawson, & Piraino 2016) and even populations of the same

species (Dawson et al., 2015), our observations on the moon jellyfish are likely

describing general trends. Field evidence of a strong positive correlation between medusa

density and prey abundance for two distinct *A. aurita* populations (Dawson et al., 2015,

their Figure 4d) implies that observed population outbreaks in response to increased food

supply are common across the *Aurelia* species complex and other metagenic jellyfish

species.

Winter warming and jellyfish blooms

Our projected winter warming scenario indicates pronounced effects of an increase in

water temperatures from 5 to 10 °C on the development of temperate *A. aurita*

populations in combination with regionally observed increases in food level. Based on

recent climate analyses, rises in global ocean surface temperature by 1 to 2 °C can be

expected within the next 100 years, with a particularly strong winter warming trend in

semi-enclosed ecosystems such as the North and Baltic Seas (Belkin, 2009; Mackenzie &

Schiedeck, 2007). As reflected by doubled yearly population growth rates under high

food conditions in response to the simulated increase in winter temperatures, our findings

support that increased water temperature can boost the future booms and busts of jellyfish

blooms (cf. Richardson, Bakun, Hays, & Gibbons 2009; Xie, Fan, Wang, & Chen 2015).

According to our population matrix model, underlying seasonal changes in population

composition and structure due to winter warming include a general trend towards more

medusae and less polyps under increased food conditions. In regions with limited food

availability, winter warming can further increase the net reproductive rate of temperate

jellyfish populations. Due to enhanced strobilation periods and increased numbers of

released ephyrae per polyp (Holst, 2012; Purcell, 2005) as simulated in our matrix model

projections, warmer winter temperatures may shorten the generation time under low food

conditions drastically.

Climate warming has previously been pointed out to benefit the population size of

temperate jellyfish species, but not tropical or boreal species (Purcell et al., 2012).

Further, a cold trigger is necessary to stimulate strobilation in many scyphozoans (Holst,

2012; Lotan, Fine, & Ben-Hillel 1994), while heatwave events can negatively affect the

population dynamics of jellyfish (Chi, Mueller-Navarra, Hylander, Sommer, & Javidpour

2018). Taking into account these physiological constraints, we suggest winter warming

may promote the blooming potential of certain temperate jellyfish species but seems to be

a less powerful driver when compared to increased food availability.

Jellyfish life histories in a changing environment

Beyond revealing the demographic key variables that control population growth of *A.*

aurita in response to habitat eutrophication and climate warming, our matrix model

projections provide general implications for the development of jellyfish blooms with

regard to environmental change. In the following, we address how other potential drivers

of jellyfish blooms which have not been quantified in the present study can overrule

important ecological parameters that constrain the demography of metagenic jellyfish.

Such natural limitations include intra- and inter-specific competition, predation, substrate

availability, hydrodynamic currents and the effects of salinity, hypoxia, pH, pollution,

light and sedimentation (Arai, 2005; Johnson, Perry, & Burke 2001; Lucas, Graham, &

Widmer 2012).

Our findings emphasize the crucial role of predation by benthic fauna, spatial

competition and availability of settlement substrate for transition from the planktonic

planula to the benthic polyp stage under natural conditions (Lucas, 2001; Lucas, Graham,

& Widmer 2012). We estimated a transition probability of 0.001 month^{-1} from larva to

polyp, which is in agreement with previous estimates for *A. aurita* (Xie, Fan, Wang, &

Chen 2015), when scaling to a stable *A. aurita* population under low food levels ($\lambda = 1$

month^{-1}). The local *A. aurita* jellyfish population in the semi-enclosed fjord system

Kerteminde Fjord/Kertinge Nor, where zooplankton biomasses are comparable to our low

food conditions, has remained unchanged over the last 25 years (Goldstein & Riisgård,

2016; Olesen, Frandsen, & Riisgård 1994; Riisgård, Barth-Jensen, & Madsen 2010) and

may, similar to our projected *A. aurita* population under low food conditions, mainly

depend on environmental parameters that determine larval transition to the benthic polyp

stage.

The rapid growth potential we show for our calibrated jellyfish population under

constantly high food (optimum) conditions however stands in contrast to rather low

densities of *A. aurita* medusae observed in open coastal waters (Goldstein & Riisgård,

2016; Möller, 1980) or in Limfjorden (Hansson, Moeslund, Kiørboe, & Riisgård 2005;

Møller & Riisgård, 2007), which implies additional factors regulating the demographic

dynamics of jellyfish in less protected ecosystems. The results of our life table response

experiment analysis suggest that besides successful metamorphosis of planula larvae,

transition from ephyra to medusa and survival of medusae are the most critical

parameters for the development of *A. aurita* populations in exposed coastal regions.

According to our matrix projection for high food levels, reduced medusa survival ($t_{MM} =$

0.1 month^{-1}), as a possible consequence of natural predation by other cnidarians,

ctenophores or fish (Arai, 2005), or alternatively, a lowered transition probability from

ephyra to medusa stage (t_{EM} = 0.0001 month^{-1}), would remarkably decelerate the

population growth rate (from λ =1.63 month^{-1} to 1.10 month^{-1} and 1.09 month^{-1},

respectively).

Since jellyfish constitute major parts of the diet of some fish species (Arai, 2005;

Hamilton, 2016) and can in turn be important competitors for zooplankton prey with

forage fish (Purcell, 2012), these findings indicate dramatic effects of large-scale

alterations in ecosystem dynamics by overfishing on the demography of the pelagic

medusa and ephyra stages. Lack of competition and predation by fish may further create

protected niches for nonindigenous jellyfish species that are inter-regionally transported

via ballast water exchange (Richardson, Bakun, Hays, & Gibbons 2009). Man-made

shoreline modifications contribute decisively to the availability of hard substrate for

benthic polyps (Purcell, 2012) and are, according to our findings, likely to enhance the

impact of changing food web structures and climate conditions on jellyfish blooms. For

these reasons, we expect that future jellyfish populations will further exploit their growth

potential in the context of environmental change. Removal of jellyfish and reduction of

their prey availability have previously been considered as possible management strategies

(Bordehore et al., 2015) which, however, could have unpredictable consequences for the

sensitive balance of marine food webs. A decisive step forward will be to further explore

jellyfish life histories in stochastic environments, taking into account specific traits that

facilitate survival of metagenic species under unfavorable conditions, such as production

of resting stages (podocysts and planulocysts), regeneration and reverse development

(Piraino, De Vito, Schmich, Bouillon, & Boero 2004).

Conclusions

Our study highlights that food availability can drive and specifically shape the occurrence

of jellyfish blooms. We suggest that for most temperate scyphozoans predicted changes

in water temperature may cause less dramatic increases in jellyfish density than already

established food concentrations in several eutrophic regions around the world. The role of

natural competitors and predators in controlling jellyfish populations needs further

exploration, but our findings might serve as a promising starting point. We conclude that

especially the combination of key environmental drivers, such as habitat eutrophication

and climate change, may severely promote jellyfish blooms. Our findings emphasize the

fundamental importance of studying jellyfish life histories using quantitative approaches.

Such approaches, which take into account species-, life stage- and age-specific variation

in demographic rates, are highly valuable tools for informing future management of mass

occurrences in marine ecosystems.

Authors' contributions

US and JG conceived the ideas and designed the methodology, JG collected the data and

led the writing of the manuscript; JG and US analyzed the data. Both authors contributed

critically to the drafts and gave final approval for publication.

Acknowledgements

We thank Kim Lundgreen for technical assistance and Daniel A. Levitis, Hans Ulrik

Riisgård, Hal Caswell, Iain Stott, Cesar Bordehore, Per Andersen, Owen Jones, Matthew

Spencer and Cathy H. Lucas for helpful comments and discussions. We are grateful to the

reviewers for providing valuable feedback on the manuscript. This study was financially

supported by the Max-Planck Society (Max-Planck Odense Center on the

Biodemography of Aging, Denmark). The authors declare no conflict of interest.

Data Accessibility

547   A preprint version of this article is available from bioRxiv https://doi.org/10.1101/102814

548   (Goldstein & Steiner, 2019). Data will be made publicly accessible on the Dryad Digital

549   Repository at the time of publication.

References

Arai, M. N. (2005). Predation on pelagic coelenterates: a review. Journal of the Marine

Biological Association of the United Kingdom, 85(3), 523–536. doi:

10.1017/S0025315405011458

Arai, M. N. (2009). The potential importance of podocysts to the formation of

scyphozoan blooms: a review. In J. E. Purcell & K. A. Pitt (Eds.), *Jellyfish blooms:*

causes, consequences, and recent advances (pp. 241–246). Netherlands: Springer.

doi: 10.1007/s10750-008-9588-5

Båmstedt, U., Lane, J., & Martinussen, M. B. (1999). Bioenergetics of ephyra larvae of

the scyphozoan jellyfish *Aurelia aurita* in relation to temperature and salinity. Marine

Biology, 135(1), 89–98. doi: 10.1007/s002270050605

Belkin, I. M. (2009). Rapid warming of large marine ecosystems. Progress in

Oceanography, 81(1), 207–213. doi: 10.1016/j.pocean.2009.04.011

Bengtson, D. A., Leger, P., & Sorgeloos, P. (1991). Use of *Artemia* as a food source for

aquaculture. In R. A. Browne, P. Sorgeloos & C. A. N. Trotman (Eds.), *Artemia*

Biology (pp. 255–285). Florida, USA: CRC Press.

Boero, F., Bouillon, J., Gravili, C., Miglietta, M. P., Parsons, T., & Piraino, S. (2008).

Gelatinous plankton: irregularities rule the world (sometimes). Marine Ecology

Progress Series, 356, 299–310. doi: 10.3354/meps07368

Bordehore, C., Fuentes, V. L., Segarra, J. G., Acevedo, M., Canepa, A., & Raventós, J.

(2015). Use of an inverse method for time series to estimate the dynamics of and

management strategies for the box jellyfish *Carybdea marsupialis*. PLoS ONE, 10(9),

e0137272. doi: 10.1371/journal.pone.0137272

Breithaupt, H. (2003). Aliens on the shores. EMBO reports, 4(6), 547–550. doi:

10.1038/sj.embor.embor877

Brotz, L., Cheung, W. W., Kleisner, K., Pakhomov, E., & Pauly, D. (2012). Increasing

jellyfish populations: trends in large marine ecosystems. Hydrobiologia, 690(1), 3–20.

doi: 10.1007/s10750-012-1039-7

Caswell, H. (1996). Analysis of life table response experiments II. Alternative

parameterizations for size-and stage-structured models. Ecological Modelling, 88(1-

3), 73–82. doi: 10.1016/0304-3800(95)00070-4

Caswell, H. (2001). Matrix population models: construction, analysis, and interpretation

(2nd ed.). Massachusetts, USA: Sinauer Associates.

Chi, X., Mueller-Navarra, D. C., Hylander, S., Sommer, U., & Javidpour, J. (2018). Food quality matters: Interplay among food quality, food quantity and temperature affecting life history traits of *Aurelia aurita* (Cnidaria: Scyphozoa) polyps. Science of the Total Environment, 656, 1280−1288. doi: 10.1016/j.scitotenv.2018.11.469

Dawson, M. N., & Hamner, W. M. (2009). A character-based analysis of the evolution of jellyfish blooms: adaptation and exaptation. Hydrobiologia, 616(1), 193−215. doi: 10.1007/s10750-008-9591-x

Dawson, M. N., Gupta, A. S., & England, M. H. (2005). Coupled biophysical global ocean model and molecular genetic analyses identify multiple introductions of cryptogenic species. Proceedings of the National Academy of Sciences, 102(34), 11968−11973. doi: 10.1073/pnas.0503811102

Dawson, M. N., Cieciel, K., Decker, M. B., Hays, G. C., Lucas, C. H., & Pitt, K. A. (2015). Population-level perspectives on global change: genetic and demographic analyses indicate various scales, timing, and causes of scyphozoan jellyfish blooms. Biological Invasions, 17(3), 851−867. doi: 10.1007/s10530-014-0732-z

Eckert, R., Randall, D., & Augustine, G. (1988). Animal energetics and temperature relations. In R. Eckert & G. Augustine (Eds.), *Animal physiology, mechanisms and adaptions* (pp. 555−605). Los Angeles, USA: WH Freeman and Company.

Fey, S. B., Siepielski, A. M., Nusslé, S., Cervantes-Yoshida, K., Hwan, J. L., Huber, E. R., ... Carlson, S. M. (2015). Recent shifts in the occurrence, cause, and magnitude of animal mass mortality events. Proceedings of the National Academy of Sciences, 112(4), 1083−1088. doi: 10.1073/pnas.1414894112

Frandsen, K. T., & Riisgård, H. U. (1997). Size dependent respiration and growth of

jellyfish, *Aurelia aurita*. Sarsia, 82(4), 307−312. doi:

10.1080/00364827.1997.10413659

Fu, Z., Shibata, M., Makabe, R., Ikeda, H., & Uye, S.-I. (2014). Body size reduction

under starvation, and the point of no return, in ephyrae of the moon jellyfish *Aurelia*

aurita. Marine Ecology Progress Series, 510, 255−263. doi: 10.3354/meps10799

Goldstein, J., & Riisgård, H. U. (2016). Population dynamics and factors controlling

somatic degrowth of the common jellyfish, *Aurelia aurita*, in a temperate semi-

enclosed cove (Kertinge Nor, Denmark). Marine Biology, 163(2), 33−44. doi:

10.1007/s00227-015-2802-x

Goldstein, J., & Steiner, U. K. (2019). Preprint version: Ecological drivers of jellyfish

blooms - the complex life history of a cosmopolitan medusa (*Aurelia aurita*). bioRxiv.

doi: 10.1101/102814.

Gröndahl, F. (1989). Evidence of gregarious settlement of planula larvae of the

scyphozoan *Aurelia aurita*: An experimental study. Marine Ecology Progress Series,

56, 119−125.

Hamilton, G. (2016). The secret lives of jellyfish. Nature News, 531(7595), 432−434.

doi: 10.1038/531432a

Hamner, W. M., & Dawson, M. N. (2009). A review and synthesis on the systematics and

evolution of jellyfish blooms: advantageous aggregations and adaptive assemblages.

Hydrobiologia, 616(1), 161−191. doi: 10.1007/s10750-008-9620-9

Hamner, W. M., & Jenssen, R. M. (1974). Growth, degrowth, and irreversible cell

differentiation in *Aurelia aurita*. Amercian Zoologist, 14(2), 833−849. doi:

10.1093/icb/14.2.833

Hansson, L. J., Moeslund, O., Kiørboe, T., & Riisgård, H. U. (2005). Clearance rates of

jellyfish and their potential predation impact on zooplankton and fish larvae in a

neritic ecosystem (Limfjorden, Denmark). Marine Ecology Progress Series, 304,

117−131. doi: 10.3354/meps304117

Holst, S. (2012). Effects of climate warming on strobilation and ephyra production of

North Sea scyphozoan jellyfish. Hydrobiologia, 690(1), 127−140. doi:

10.1007/s10750-012-1043-y

Holst, S., & Jarms, G. (2007). Substrate choice and settlement preferences of planula

larvae of five Scyphozoa (Cnidaria) from German Bight, North Sea. Marine Biology,

151(3), 863−871. doi: 10.1007/s00227-006-0530-y

Ishii, H., & Takagi, A. (2003). Development time of planula larvae on the oral arms of

the scyphomedusa *Aurelia aurita*. Journal of Plankton Research, 25(11), 1447−1450.

doi: 10.1093/plankt/fbg094

Ishii, H., & Watanabe, T. (2003). Experimental study of growth and asexual reproduction

in *Aurelia aurita* polyps. Sessile Organisms, 20(2), 69−73. doi: 10.4282/sosj.20.69

Johnson, D. R., Perry, H. M., & Burke, W. D. (2001). Developing jellyfish strategy

hypotheses using circulation models. Hydrobiologia, 451(1), 213−221.

Kamiyama, T. (2011). Planktonic ciliates as a food source for the scyphozoan *Aurelia*

aurita (s.l.): Feeding activity and assimilation of the polyp stage. Journal of

Experimental Marine Biology and Ecology, 407(2), 207−215. doi:

10.1016/j.jembe.2011.06.016

Keen, S. L. (1987). Recruitment of *Aurelia aurita* (Cnidaria: Scyphozoa) larvae is

position-dependent, and independent of conspecific density, within a settling surface.

Marine Ecology Progress Series, 38, 151−160.

Liu, W. C., Lo, W. T., Purcell, J. E., & Chang, H. H. (2009). Effects of temperature and

light intensity on asexual reproduction of the scyphozoan, *Aurelia aurita* (L.) in

Taiwan. Hydrobiologia, 616(1), 247−258. doi: 10.1007/s10750-008-9597-4

Lotan, A., Fine, M., & Ben-Hillel, R. (1994). Synchronization of the life cycle and

dispersal pattern of the tropical invader scyphomedusan *Rhopilema nomadica* is

temperature dependent. Marine Ecology Progress Series, 109, 59−65.

Lucas, C. H. (2001). Reproduction and life history strategies of the common jellyfish,

Aurelia aurita, in relation to its ambient environment. Hydrobiologia, 451(1-3),

229−246.

Lucas, C. H., & Dawson, M. N. (2014). What are jellyfishes and thaliaceans and why do

they bloom? In K. A. Pitt & C. H. Lucas (Eds.) *Jellyfish blooms* (pp. 9−44).

Netherlands: Springer. doi: 10.1007/978-94-007-7015-7_2

Lucas, C. H., Graham, W. M., & Widmer, C. (2012). Jellyfish Life Histories: Role of

Polyps in Forming and Maintaining Scyphomedusa Populations. Advances in Marine

Biology, 63, 133−196.

MacKenzie, B. R., & Schiedek, D. (2007). Daily ocean monitoring since the 1860s shows

record warming of northern European seas. Global Change Biology, 13(7),

1335−1347. doi: 10.1111/j.1365-2486.2007.01360.x

Maji, B., Bhattacharyya, J., & Pal, S. (2016). Potential effects of invasive *Pterois volitans*

in coral reefs. Letters in Biomathematics, 3(1), 119−139. doi:

10.1080/23737867.2016.1213146

Möller, H. (1980). Population dynamics of *Aurelia aurita* medusae in Kiel Bight,

Germany (FRG). Marine Biology, 60(2-3), 123−128. doi: 10.1007/BF00389155

Møller, L. F., & Riisgård, H. U. (2007). Population dynamics, growth and predation

impact of the common jellyfish *Aurelia aurita* and two hydromedusae (*Sarsia*

tubulosa and *Aequorea vitrina*) in Limfjorden (Denmark). Marine Ecology Progress

Series, 346, 153−165. doi: 10.3354/meps06960

Nielsen, A. S., Pedersen, A. W., & Riisgård, H. U. (1997). Implications of density driven

currents for interaction between jellyfish (*Aurelia aurita*) and zooplankton in a Danish

fjord. Sarsia, 82(4), 297−305. doi: 10.1080/00364827.1997.10413658

Olesen, N. J., Frandsen, K., & Riisgård, H. U. (1994). Population dynamics, growth and

energetics of jellyfish, *Aurelia aurita*, in a shallow fjord. Marine Ecology Progress

Series, 105, 9−18.

Piraino, S., De Vito, D., Schmich, J., Bouillon, J., & Boero, F. (2004). Reverse

development in Cnidaria. Canadian Journal of Zoology, 82(11), 1748−1754. doi:

10.1139/z04-174

Pitt, K. A., Welsh, D. T., & Condon, R. H. (2009). Influence of jellyfish blooms on

carbon, nitrogen and phosphorus cycling and plankton production. Hydrobiologia,

616(1), 133−149. doi: 10.1007/s10750-008-9584-9

Purcell, J. E. (2005). Climate effects on formation of jellyfish and ctenophore blooms: a

review. Journal of the Marine Biological Association of the United Kingdom, 85(3),

461–476. doi: 10.1017/S0025315405011409

Purcell, J. E. (2012). Jellyfish and ctenophore blooms coincide with human proliferations

and environmental perturbations. Annual Review of Marine Science, 4, 209–235. doi:

10.1146/annurev-marine-120709-142751

Purcell, J. E., Uye, S.-I., & Lo, W. T. (2007). Anthropogenic causes of jellyfish blooms

and their direct consequences for humans: a review. Marine Ecology Progress Series,

350, 153–174. doi: 10.3354/meps07093

Purcell, J. E., Atienza, D., Fuentes, V. L., Olariaga, A., Tilves, U., Colahan, C., & Gili J.

M. (2012). Temperature effects on asexual reproduction rates of scyphozoan species

from the northwest Mediterranean Sea. Hydrobiologia, 690(1), 169–180. doi:

10.1007/s10750-012-1047-7

R Core Team (2015). R: A language and environment for statistical computing. R

Foundation for Statistical Computing, Vienna, Austria. http://www.r-project.org

Richardson, A. J., Bakun, A., Hays, G. C., & Gibbons, M. J. (2009). The jellyfish joyride:

causes, consequences and management responses to a more gelatinous future. Trends

in Ecology & Evolution, 24(6), 312–322. doi: 10.1016/j.tree.2009.01.010

Riisgård, H. U., Jensen, M. H., & Rask, N. (2008). Odense Fjord and Kerteminde

Fjord/Kertinge Nor. In U. Schiewer (Ed.), *Ecology of Baltic Coastal Waters* (pp.

361–394). Germany: Springer.

Riisgård, H. U., Barth-Jensen, C., & Madsen, C. V. (2010). High abundance of the

jellyfish *Aurelia aurita* excludes the invasive ctenophore *Mnemiopsis leidyi* to

establish in a shallow cove (Kertinge Nor, Denmark). Aquatic Invasions, 5(4),

347−356. doi: 10.3391/ai.2010.5.4

Riisgård, H. U., Andersen, P., & Hoffmann, E. (2012). From fish to jellyfish in the

eutrophicated Limfjorden (Denmark). Estuaries and Coasts, 35(3), 701−713. doi:

10.1007/s12237-012-9480-4

Scorrano, S., Aglieri, G., Boero, F., Dawson, M.N., & Piraino, S. (2016). Unmasking

Aurelia species in the Mediterranean Sea: an integrative morphometric and molecular

approach. Zoological Journal of the Linnean Society, 180(2), 243−267. doi:

10.1111/zoj.12494

Schneider, G., & Behrends, G. (1994). Population dynamics and the trophic role of

Aurelia aurita medusae in the Kiel Bight and western Baltic. ICES Journal of Marine

Science, 51(4), 359−367.

Schneider, G., & Behrends, G. (1998). Top-down control in a neritic plankton system by

Aurelia aurita medusae – a summary. Ophelia, 48(2), 71−82. doi:

10.1080/00785236.1998.10428677

Schnedler-Meyer, N. A., Kiørboe, T., & Mariani, P. (2018). Boom and Bust: Life

History, Environmental Noise, and the (un)Predictability of Jellyfish Blooms.

Frontiers in Marine Science, 5, 257. doi: 10.3389/fmars.2018.00257

Schrope, M. (2012). Marine ecology: attack of the blobs. Nature, 482(7383), 20−21.

Stearns, S. C. (1976). Life-history tactics: a review of the ideas. The Quarterly Review of

Biology, 51(1), 3−47. doi: 10.1086/409052

Stott, I., Hodgson, D. J., & Townley, S. (2012). popdemo: an R package for population

demography using projection matrix analysis. Methods in Ecology and Evolution,

3(5), 797–802. doi: 10.1111/j.2041-210X.2012.00222.x

Stubben, C. J., & Milligan, B. G. (2007). Estimating and Analyzing Demographic Models

Using the popbio Package in R. Journal of Statistical Software, 22(11), 1–23. doi:

10.18637/jss.v022.i11

Tuljapurkar, S., & Caswell, H. (1997). Structured-population models in marine,

terrestrial, and freshwater systems. New York, NY: Chapman & Hall.

Vanhaecke, P., & Sorgeloos, P. (1980). International Study on *Artemia*. IV. The

biometrics of *Artemia* strains from different geographical origin. The brine shrimp

Artemia, 3, 393–405.

Xie, C., Fan, M., Wang, X., & Chen, M. (2015) Dynamic model for life history of

scyphozoa. PLoS ONE, 10(6), e0130669. doi: 10.1371/journal.pone.0130669

Supporting Information

Additional Supporting Information is available online in the supporting information tab

for this article.

 Tables

755 Table 1. Experimental conditions for life stages of the jellyfish *Aurelia aurita* (L: planula

756 larva, P: polyp, E: ephyra, M: medusa) under low (*L*) and high (*H*) food conditions. l_0:

757 initial number of individuals per treatment, t_0, Δt: start and duration of experiment, c:

758 food concentration based on the carbon content of available prey organisms (*Artemia*

759 *salina*).

Food treatment	Life stage	l_0 (ind.)	t_0 (month)	Δt (d)	c (μg C L^{-1} d^{-1})
L	L	960	Sept	52	0
	P	100	Aug	364	1
	E	33	Mar	153	10
	M	16	Apr	296	-
H	L	960	Jun	48	0
	P	100	Aug	364	10
	E	33	Jan	158	100
	M	33	Feb	391	>1000

761 Table 2. Demographic parameters of yearly stage-structured projection matrices (P: polyp, E: ephyra, M: medusa) for a local *Aurelia*

762 *aurita* population under low (*L*) and high food conditions (*H*) at present water temperatures (PWT) and for a predicted winter warming

763 trend (WW).

Demographic parameter	L		H	
	PWT	WW	PWT	WW
Scaled population growth rate λ (month^{-1})	1.00	1.01	1.63	1.73
Damping ratio ρ (months)	58.6	59.2	2.8	3.2
Stable stage proportion (P, E, M)	0.94, 0.04, 0.02	0.84, 0.12, 0.04	0.76, 0.04, 0.20	0.60, 0.02, 0.38
Scaled reproductive value R (ind.) (P, E, M)	1, 0.03, 0.42	1, 8.53, 3×10^{-7}	1, 0.27, 9.13	1, 0.18, 7.43
Net reproductive rate R_0 (ind.)	1.0	1.8	57.0	80.9
Generation time T (months)	200.0	111.8	8.3	8.0
Life expectancy e_0 (months) (P, E, M)	394.5, 1.7, 3.1	394.5, 2.1, 3.1	86.7, 2.9, 8.0	86.7, 3.5, 8.0

Figures

Figure 1. Life history of the moon jellyfish *A. aurita* (Scyphozoa, Cnidaria). (a)

Metagenic life cycle with alternation of generations between the sexually reproducing

medusa and the asexually reproducing polyp stage. (b) Life cycle graph for *A. aurita*,

including transition probability t, fecundity F (solid arrows) and mortality † (broken

arrows) of all life stages, i.e. planula larva (L), polyp (P), ephyra (E) and medusa (M). (c)

Reduced life cycle graph for polyps, ephyrae and medusae considering fast (daily)

transition from larva to polyp stage.

Figure 2. Population matrix model for *A. aurita* under low (upper panels) and high (lower panels) food conditions (~1:10) at present water temperatures (PWT) and in a winter warming scenario (WW). (a–b) Matrix projection over a 10-year period, standardized to the long-term effects of population growth rates λ = 1 month^{-1} (low food availability) and λ = 1.63 month^{-1} (high food availability), respectively. (c–d) Stable stage distribution (i.e. relative, constant long-term proportions) of polyps (P; solid lines), ephyrae (E; dotted lines) and medusae (M; broken lines).

Figure 3. Sensitivity of the population growth rate λ of *A. aurita* to changes in monthly stage-specific transitions (1: $t_{PP}+F_{PP}$, 2: F_{PE}, 3: $F_{PE}×t_{EM}$, 4: t_{EE}, 5: t_{EM}, 6: $F_{ML}×t_{LP}$, 7: t_{MM}) under (a) low and (b) high food conditions. (c) Life table response experiment (LTRE) analysis for sensitivity of λ to perturbations in the population projection matrix elements when comparing high to low food conditions (~10:1).