Unifying Population and Landscape Ecology with Spatial Capture-recapture

J. Andrew Royle, USGS Patuxent Wildlife Research Center
Angela K. Fuller, USGS, Dept. of Natural Resources, Cornell University
Christopher Sutherland, Dept. of Environmental Conservation, UMASS-Amherst

Running title: spatial capture-recapture

Keywords: Density, landscape connectivity, landscape ecology, non-invasive sampling, population ecology, spatial capture-recapture, resource selection, spatial point process.

Article type: Reviews and Syntheses

Statement of authorship: JAR, AKF and CS contributed equally to the development of the structure, content and writing of this manuscript.

Word count (abstract/main text/box1/box2/box3/box4): 172/5814/267/152/252/250

Reference count: 93

Figures: 1

Tables: 0

Boxes: 4

Corresponding author information: J. Andrew Royle; USGS Patuxent Wildlife Research Center; 12100 Beech Forest Rd; Laurel MD 20708.

39
40
41
42
Abstract
Spatial heterogeneity in the environment induces variation in population demographic rates and
dispersal patterns, which result in spatio-temporal variation in density and gene flow. Unfortunately,
applying theory to learn about the role of spatial structure on populations has been hindered by the
lack of mechanistic spatial models and inability to make precise observations of population structure.
Spatial capture-recapture (SCR) represents an individual-based analytic framework for overcoming this
fundamental obstacle that has limited the utility of ecological theory. SCR methods make explicit use of
spatial encounter information on individuals in order to model density and other spatial aspects of
animal population structure, and have been widely adopted in the last decade. We review the
historical context and emerging developments in SCR models that enable the integration of explicit
ecological hypotheses about landscape connectivity, movement, resource selection, and spatial
variation in density, directly with individual encounter history data obtained by new technologies (e.g.,
camera trapping, non-invasive DNA sampling). We describe ways in which SCR methods stand to
revolutionize the study of animal population ecology.
57

58   *Keywords*
59   Density, landscape connectivity, landscape ecology, non-invasive sampling, population ecology, spatial
60   capture-recapture, resource selection, spatial point process.

INTRODUCTION

Understanding factors influencing natural variation in population size and structure, demographic

rates, and movement has long been a central research focus for population ecologists. Despite well-

developed theories over the last half century demonstrating the importance of spatial structure in

shaping spatio-temporal population dynamics (e.g., Huffaker 1958, Hanski 1999, Elner et al. 2001), the

field of population ecology remains, by and large, unconcerned about within-population spatial

processes and their effects on populations. Ecologists routinely study such processes as how

individuals use space within their home range, how they perceive connectivity of the landscape,

interact with other individuals of the same or other species and how survival or recruitment might be

impacted by landscape heterogeneity. However, the population level implications of these processes

are not widely studied. Instead the focus is at the individual level, often by studying only a few

individuals, with no accounting for how those individuals are sampled from the population. Extending

inferences from the individual to the population is not straightforward and in some cases not even

possible without a formal statement of a population model linking the sample to the true state, and a

description of the sampling process.

Much of what drives the spatial ecological processes that give rise to spatio-temporal

population dynamics is the structure and configuration of the landscape (Turner et al. 2001). In fact,

linking landscape structure to ecological processes is the primary focus of landscape ecology. This focus

on how spatial structure influences ecosystem composition, structure, and function (Turner et al. 2001)

by definition, avoids any assumption about spatial homogeneity. When related to animal populations,

the tendency in landscape ecology is to focus on movement processes, specifically landscape

connectivity (Taylor et al. 1993) and resource selection functions (Chetkiewicz et al. 2009, McLoughlin

et al. 2010) rather than demographic rates and population state variables that are of interest in

population ecology. Population and landscape ecology offer alternative, yet equally important

approaches for understanding spatio-temporal dynamics, yet a consistent theory or quantitative

framework linking spatial landscape structure and population ecology does not yet exist.

The ecological theory underpinning landscape and population ecology is well-developed

(Tilman & Karieva 1997; Hanski & Ovaskainen 2000, Hanski 2001, Turner et al. 2001, Kot 2001, Williams

et al. 2002; Gets & Saltz 2008; Allen & Singh 2016), but testing theoretical models and predictions

about spatial ecology in practice is both logistically and statistically challenging. One major impediment

is the lack of general mechanistic spatial models that can be applied to empirical data; this precludes

rigorous testing of theoretical predictions. Spatial point process models (Illian et al. 2008) provide a

natural framework for characterizing the spatial structure of populations assumed to be static and that

can be observed with a high degree of accuracy. However, point process models have not been widely

adopted in practical field studies of population ecology where individuals cannot be enumerated easily.

In practice, populations distributed widely in space must be studied by observing a sample of

individuals, sometimes only a very small fraction, at only a few time points and at only a few locations.

In some cases, individuals can be continuously monitored (e.g., by telemetry), but in general it is not

possible to observe the status of animals perfectly – either their demographic status, their location, or

even whether or not they are alive. This is one of the key considerations that has motivated the

development and widespread adoption of capture-recapture methods which are now ubiquitous in

ecology (Williams et al. 2002; Cooch & White 2006).

For decades, capture-recapture methods have been the cornerstone of ecological statistics as

applied to population biology (Nichols 1992, Williams et al. 2002). At their core, capture-recapture

models are the canonical class of models for "individual encounter history" data. These data are

obtained by capturing or encountering individuals (e.g., using camera traps, acoustic sampling, non-

invasive genetic sampling, or direct physical capture), marking them, and observing them over time.

Capture-recapture methods have had a profound influence on the study and understanding of

demography in wild populations (Karanth et al. 2006, Pradel et al. 2010), in advancing ecological theory

(Cooch et al. 2002), and informing modern conservation and wildlife management practices (Nichols

and MacKenzie 2004). While capture-recapture has become the standard sampling and analytical

framework for the study of population processes (Williams, Nichols & Conroy 2002) it has advanced

independent of and remained unconnected to the spatial structure of the population or the landscape

within which populations exist. Furthermore, capture-recapture does not invoke any spatially explicit

biological processes and thus is distinctly non-spatial, accounting neither for the inherent spatial

nature of the sampling nor of the spatial distribution of individual encounters. This precludes the study

of many important spatial processes and/or the emerging within-population spatial structure that is

arguably as important as demographic rates in population ecology. Recently developed *spatial* capture-

recapture (SCR) methods (Royle et al. 2014) couple a spatio-temporal point process with a spatially

explicit observation model which resolves these important criticisms and offers a significant advance in

our ability to quantify and study spatial processes using encounter history data. Spatial capture-

recapture represents an extension of classical capture-recapture and allows for both the spatial

organization of sampling devices and the spatial information that is inherent in essentially all studies of

animal populations, i.e., *spatial encounter histories*.

Although a relatively recent advance in the field of statistical ecology (Efford 2004), the past

decade has seen an explosive growth in SCR methodological development and applications (Box 1).

Spatial capture-recapture provides a quantitative framework that links ecological processes at the

individual and population levels. SCR promises the integration of models (hypotheses) of within-

population dynamics with "population level" parameters and dynamics. Thus, SCR has proven to be

more than simply an extension of a technique, but has emerged as a flexible framework that allows

ecologists to test hypotheses about a wide range of ecological theories including landscape and

network connectivity (Sutherland et al. 2015, Fuller et al. 2015), demography (Ergon & Gardner 2013,

Whittington and Sawaya 2015, Munoz et al, 2016), resource selection (Royle et al. 2013b, Proffitt et al.

2015), and movement and dispersal (Borchers et al. 2014, Lagrange et al. 2014, Schaub & Royle 2015,

Royle et al. 2016). While classical capture-recapture methods focus on population level quantities, SCR

models allow for the "downscaling" of population structure from coarse summaries (spatial and/or

demographic) into finer-scale components by the use of a spatially explicit individual-based point

process model. By connecting population level attributes to individual level attributes that are spatially

realistic, SCR unifies the fundamental concepts of population and landscape ecology, relating spatial

encounters of individuals to explicit descriptions of spatial structure and of how space is sampled (Box

2).

In this review, we describe the basic elements of spatial capture-recapture and how SCR

methods advance spatial population ecology by providing a unified framework that integrates

important concepts and elements of population ecology and landscape ecology. As such, the

framework allows for the study of density, movement, resource selection, landscape connectivity, and

other spatial population processes using ordinary encounter history data. Finally, we discuss new

directions in the study of animal populations that are made possible by the existence of spatially

explicit capture-recapture methods.

THE ELEMENTS OF SPATIAL CAPTURE-RECAPTURE

Traditional capture-recapture (CR) models were largely motivated by a formal statistical sampling view

of how individuals are encountered by sampling, with little or no direct consideration given to the

fundamental spatial nature of the sampling. As a result, traditional CR models represent, in essence,

"fish bowl" sampling – that is, a system that is devoid of any meaningful spatial context. This leads

immediately to several important technical concerns that arise in the application of traditional CR to

the study of animal populations which are necessarily spatially explicit.

One important deficiency with classical closed population models is the inability to directly

estimate animal *density* (D), arguably *the* state variable of interest in the vast majority of animal

monitoring studies (Krebs 1985, Turchin 1998). This is because, in almost all practical field applications,

it is not possible to precisely define the effective area sampled by a set of trapping devices due to

movement of individuals into and out of the region within which sampling occurs (Dice 1938; Hayne

1949; Wilson and Anderson 1985a,b). Secondly, the probability of encountering an individual is

necessarily heterogeneous among those individuals exposed to sampling. For example, individuals on

the periphery of a trapping grid should have lower probability of capture than individuals with home

ranges on the interior of the trapping grid (Figure 1). This heterogeneity in encounter probability has

long been known to induce negative bias in estimates of abundance (N), and hence density (K. Ullas

Karanth & Nichols, 1998; Otis, Burnham, White, & Anderson, 1978), and was one of the factors that

motivated the development of SCR methods (Efford, 2004). These (and other) technical limitations of

the non-spatial CR framework arise directly as a result of a lack of spatial explicitness. On the other

hand, SCR integrates models that describe the spatially explicit nature of sampling, how individuals are

distributed, and how they use space.

SCR models assume that a population of N individuals is sampled and that each individual has

associated with it a spatial location which represents its activity centre which can be expressed by its

X and Y coordinates as $\mathbf{s}_i = [s_{i,X}, s_{i,Y}]$. The collection of activity centres $\mathbf{s}_1, \ldots, \mathbf{s}_N$ can be thought of

as the realization of a statistical point process (Illian et al. 2008), a class of probability models for

characterizing the spatial pattern and distribution of points. This is perhaps the key innovation of

spatial capture-recapture because it is this model that connects observations to much of the ecological

theory that can be addressed by SCR. To formalize the point process model it is necessary to describe

the probability distribution function of the point locations. The simplest possible point process model

is to assume that each of the N point locations are distributed uniformly in space (the "uniformity

assumption"):

$$\mathbf{s}_i \sim \mathrm{Uniform}(S) \qquad\qquad (1)$$

where S is an explicit spatial region within which sampling of individuals occurs, and for which

inferences about density will be made. Formally this is referred to as the *state-space* of the point

process and is an essential component of a probabilistic characterization of potential activity centres,

which are equivalent to individuals in the SCR framework. One important distinction to be made

between SCR and classical CR methods is that the state-space S is an explicit component of the SCR

model. The state-space induces an explicit model of heterogeneous detection probabilities which may

affect inferences about density and, hence, population size.

The introduction of this statistical point process – that is, the association of a spatial coordinate

with each individual in the population – leads naturally to two distinct conceptually important and

powerful modifications of the classical capture-recapture framework which are at the core of the SCR

method: First, we can formulate a spatial model for the probability that an individual is captured in

each sample location or trap x_j for $j = 1, 2, \ldots, J$, conditional on its activity centre rather than simply

whether an individual was captured at all in a sample occasion, as is the case in traditional CR.

Acknowledging the spatial structure of the traps means observations can be spatially indexed (Box 2,

top panel) so encounter histories describe *who* (*i*), *when* (*k*) and importantly *where* (*j*) individuals were

encountered, i.e. $y_{i,j,k}$. Often, these observations are assumed to be Bernoulli outcomes:

$$y_{i,j,k} \sim \mathrm{Bernoulli}(p_{i,j,k}), \qquad\qquad (2)$$

where $p_{i,j,k}$ is the probability of encountering individual i in trap j , and occasion k . It is this model

which links the observations (spatially indexed encounters) to the underlying latent point process

describing biological pattern and process. At a minimum the encounter probability model depends on

the distance between the trap location (x_j) and the individual's activity centre (s_i) such as the half-

normal encounter model (Box 2, middle panel):

$$p_{i,j,k} = p_{0,j,k} \times e^{-(1/2\sigma^2)\mathrm{d}(x_j, s_i)^2}. \tag{3}$$

where p_0 is the baseline encounter probability, the probability of encountering an individual at its'

activity centre, the parameter σ describes the rate at which detection probability declines as a

function of distance, and $\mathrm{d}(x_j, s_i)$ is the Euclidean distance between trap j and the activity centre of

individual i (Box 2, middle panel). In a spatial capture-recapture analysis, the parameters to be

estimated are p_0, σ and population size N or density D. We note that the parameter σ

accommodates individual heterogeneity in detectability but, unlike classical models of heterogeneity

(Otis et al. 1978; Dorazio & Royle 2003) the parameter represents an explicit source of heterogeneity,

due to the distance between individual activity or home range centres and traps.

The uniformity assumption yields what is usually referred to as a homogenous point process

model, although very general models of the point process are possible. For example, when spatially

referenced covariates, say $z(\mathbf{s})$, can be identified that result in spatially heterogeneous density

surfaces (Box 2, lower panel), then a standard inhomogeneous point process model posits that

$$\Pr(\mathbf{s}) \propto exp(\beta z(\mathbf{s})), \tag{4}$$

where the parameter β corresponds to an explicit hypothesis: "does density depend on the covariate

z ?"

Integrating the point process model with the CR sampling framework leads naturally to a direct

focus on inference about parameters of the underlying point process, instead of the abstract quantity

N which is devoid of spatial context. Under the SCR modelling framework, the goal is to estimate the

number of individuals (or activity centres) within any region of the *state-space* S. For example, we may

estimate density D, the number of activity centres per unit area of S, or produce predictions of the

number of points in any formal subset of S, or functions of the entire set of N points which might be

used to test for spatial randomness, clustering mechanisms (Reich & Gardner, 2014) or other point

process assumptions.

For purposes of statistical estimation and inference, the activity centres are regarded as latent

variables (i.e., as in classical random effects or mixed models, Laird & Ware 1982). The point process

model is then precisely equivalent to the random effects distribution or prior distribution. The resulting

model is amenable to analyses by classical methods of statistical inference such as based on marginal

likelihood (Borchers & Efford 2008), in which the latent variables are removed from the likelihood by

integration, or Bayesian analysis by Markov chain Monte Carlo (MCMC; Royle & Young 2008), in which

the activity centres are explicitly estimated along with other unknown parameters and random

variables.

SCR models are now routinely applied to many taxa, across a wide variety of systems using a

range of sampling methodologies (Box 1). However, the utility of the model reaches far beyond simply

estimating density and includes the investigation of important questions about population and

landscape ecology which we describe next. Moreover, an enormous number of extensions to SCR

models can be accommodated, both to the structure of the ecological processes and the types of

observation method, including acoustic sampling (Dawson & Efford, 2009), sampling continuous space

instead of using traps (Royle, Kéry, & Guélat, 2011; Royle & Young, 2008), sampling continuous time

instead of having discrete sampling intervals (Borchers et al. 2014) and modeling population dynamics

such as survival and recruitment using individual level or state-space formulations of classical Jolly-

Seber and Cormack-Jolly Seber models (Gimenez et al. 2007; Gardner et al. 2010). We discuss some of

these extensions below.

Figure 1. Left: Two home ranges of individuals (gray circles) juxtaposed with a spatial sampling grid

(traps) illustrating the variable exposure to trapping based on home range location.   Right: the implied

distribution of individual encounter probability for a population exposed to sampling by a regular grid

(taken from Royle et al. 2014, ch. 5).

SCR – A DECADE OF DEVELOPMENT AND APPLICATION

As SCR methods were first appearing more than 10 years ago, the motivation for their development

and use was exclusively as a technical device for resolving specific technical limitations of ordinary

capture-recapture (Fig. 1). More generally, SCR methods have proved to be a flexible framework for

making ecological processes explicit in models of individual encounter history data, and for studying

spatial processes such as individual movement, resource selection, space usage, landscape

connectivity, population dynamics, spatial distribution, density and inter- and intra-specific

interactions. Historically, researchers studied these questions independently, using ostensibly

unrelated study designs and statistical procedures.

SCR for resource selection

SCR models provide a coherent framework for modeling both 2nd and 3rd order resource selection

(Johnson 1980; Box 3). Second order resource selection is defined as the processes by which

individuals select the location of their home range within a particular landscape. SCR models

parameterize an explicit representation of this selection process in the specification of the latent point

process model (individual activity centres $\mathbf{s}_1,\ldots,\mathbf{s}_N$). While typical applications involve a relatively

simple homogeneous point process model in which activity centers are distributed independently and

uniformly over the state-space S , the SCR framework accommodates inhomogeneous point process

models in which the density of activity centers varies as a function of explicit covariates or flexible

spatial response surface models (Borchers & Kidney 2014) that affect density. Inhomogeneous point

process models show great promise for testing explicit hypotheses about 2^{nd} order selection,

understanding mechanisms that influence species density distribution, and developing conservation

and management strategies with explicit abundance- or density-based objectives (Kendall et al. 2015,

Proffitt et al. 2015, Sun et al. 2015, Linden et al. 2016).

Third order resource selection – that is, selection that occurs within an individual's home range

– can be modeled explicitly in SCR models by accommodating habitat structure in the vicinity of

sampling (or trap locations) as a covariate that affects the probability of encounter (Royle et al. 2013b).

Traditionally, resource selection was studied exclusively by telemetry, and more recently GPS, and

because of the high cost are often based on a small sample of individuals observed many times.

Conversely, SCR methods may produce a sample of many more individuals, and direct information

about population level resource selection from spatial encounter data. However, individuals in the

population often do not need to be physically captured to obtain this information by SCR methods

(e.g., by camera trapping). Thus, SCR provides an alternative, efficient, and cost effective framework

for studying the important population process of resource selection from individual encounter history

data for species that classical telemetry may not be viable and that is based on a larger sample of the

population.

SCR for modeling movement and dispersal

The direct linkage of the SCR encounter probability model to movement of an individual within its

home range is one of the basic concepts of SCR (Royle & Young 2008; Borchers et al. 2014). However,

one of the key assumptions of most SCR methods to date is that the latent point locations which

represent the individual activity centers are static variables. In a sense this is a manifestation of type of

population "closure"; individuals are allowed to move around in space, but their expected location is

assumed not to change over the course of the study. Recent attention has been given to modifying the

underlying state point process model to accommodate temporal dynamics such as dispersal or

transience (Schaub & Royle 2014; Ergon & Gardner 2014). These models formally allow for the

estimation of survival probability that is free of biasing effects of dispersal whereas, classically, only

"apparent survival" has been estimated from standard capture recapture data (Schaub et al. 2004).

Even in populations where mortality or recruitment are absent, including a dynamic spatial process to

account for dispersal and transience is possible by coupling a latent movement model with a spatial

model of the encounter process (Royle et al. 2016). For example, to modify the point process model to

allow for an individual's activity center to shift from time $t-1$ to time t we might accommodate this

with a simple Markovian movement model where the difference between successive activity centers

has variance τ^2 :

$$\mathbf{s}_{i,t} \mid \mathbf{s}_{i,t-1} \sim Normal(\mathbf{s}_{i,t-1}, \tau^2 \mathbf{I})$$

Thus, SCR has a characterization as a state-space (Patterson et al. 2008) or hidden Markov model

(HMM; Langrock et al. 2012) with specific forms of observation model governed by spatial sampling

and an underlying latent process model that describes movement of individuals on the landscape. As a

result, SCR offers a general framework for the formal study of dispersal, transience, and other types of

movement from individual encounter history data.

Modeling landscape connectivity

One of the core elements of SCR is the model for encounter probability which we described above as a

function of Euclidean distance between activity centers and sample locations (Box 2). However, the

Euclidean distance assumption implies a simplistic model of space usage – that individual home ranges

are symmetric and stationary. In practice, we expect individual home ranges to be influenced by local

landscape characteristics and structure. One approach for accommodating this landscape structure-

induced asymmetry in space use in SCR models is the relaxation of the Euclidean distance assumption.

This is achieved using an alternative distance metric that is related to the landscape through

which distance is being measured, thus allowing the degree of asymmetry to be estimated using a

model that relates the observed spatial pattern of observations explicitly to the measurable landscape

characteristics. For example, Royle et al. (2013a) suggested using least-cost path distance with the

exception that, rather than being defined *a priori* based on opinion as is customary (Zeller et al. 2012),

the resistance parameters are estimated using standard likelihood methods based on spatial encounter

histories (Box 4). This idea was extended by Sutherland et al. (2015) for highly structured dendritic

networks, an extreme yet intuitive conceptual setting for investigating the utility of this asymmetric

space use model. The major development is that the model of asymmetric space use can be used to

jointly estimate density and resistance parameters which are typically defined a priori based on

opinion (Zeller, McGarigal & Whitely 2012), yielding ecologically interesting and realistic individual

home range geometries which can be scaled up to the landscape level based on direct estimates of the

landscape structure-space use relationship (Box 4). What results is the important notion that ordinary

encounter history data that is extensively collected in ecological studies with relative ease can now be

used to formally characterize landscape connectivity within a framework of statistical inference.

Sutherland et al. (2015) defined several intuitive measures of landscape connectivity based on such

upscaling based on the SCR encounter probability model to the landscape scale. The asymmetric space

use model described here and in Box 4 in general, can be extended to multiple landscape

characteristics as would be done with any log-linear regression model, and requires only that the

landscape covariates are defined at the pixel level. While current applications have focused on river

networks (Fuller et al. 2016), the approach should be highly relevant for any species for which one or

more landscape features act to impede or facilitate movement (Morin et al. *in press*) (e.g., extreme

topographic variation, well-defined networks of roads and trails used by a species, Box 4). Moreover,

SCR offers a formal model-based solution for investigating the strength of landscape interactions,

avoiding the need to arbitrarily prescribe resistance values. The possibility exists to consider other non-

Euclidean distance metrics such as circuit distance (McRae et al. 2006) or flexible deformations of

geographic space (Sampson & Guttorp 1992).

FUTURE DIRECTIONS OF SPATIAL CAPTURE-RECAPTURE

The relevance of SCR methods is expanding rapidly because these techniques allow ecologists to

explicitly test hypotheses about the mechanisms that drive ecological phenomena as diverse as habitat

selection, persistence of rare species, community assembly, invasion, and genetic diversity. The

developments described above represent significant contributions to applied population ecology

despite their relative infancy, and we believe the potential for SCR in ecology has not yet been fully

realized. We highlight specific and potentially fruitful development areas for SCR that have the

potential to make further contributions with regard to wildlife population sampling, and/or developing

and testing ecological hypotheses.

Landscape management and corridor design

It is possible to use SCR with individual encounter history data to inform landscape management

decisions such as corridor and reserve design. Because SCR models provide spatially explicit within-

population information about density, they provide objective inferences about where the population is

distributed in space and why. Therefore, SCR can serve as an empirical framework for characterizing

the utility of landscapes to populations. In particular, when combined with explicit models of

connectivity (previous section), spatially explicit metrics which integrate information about both

density and connectivity (Sutherland et al. 2015; Fuller et al. 2016; Morin et al. 2016) can be estimated,

thus providing information about quality of the landscape for maintaining connectivity and also for

maintaining source populations of important species.

Corridors are increasingly used as conservation tools, designed to facilitate movement of

individuals between habitat patches, or between two nodes or habitat blocks separated by some

distance (e.g., two protected areas) with the ultimate goal of maintaining landscape connectivity. In

the most general sense, corridor design involves defining a resistance value (i.e., resistance of the

landscape to animal movement) of each pixel in the landscape as a function of pixel characteristics,

and then subsequently selecting the lowest cost pixels, typically evaluated by estimating the

cumulative cost of moving from one area to another. The resistance of a landscape is approximated by

a 'cost' value, representing how difficult it is for an individual to move through a landscape. High

quality habits are more permeable to movement and infer lower ecological costs (i.e., they provide

increased survival and reproduction) relative to lower quality habitat. Resistance values are most often

based on subjective expert opinion or data from previously published studies (Zeller et al. 2012). These

user defined resistance models have been tested based on limited inference from few radiomarked

individuals (Driezen et al. 2007). However, examples exist of deriving resistance values from

occurrence probability from occupancy models (Walpole et al. 2012) or using a variety of different

threshold values based on the most traversable habitat from radio-marked individuals (Poor et al.

2012). Of importance is that these applications fail to utilize the information from animal movements

to directly estimate landscape resistance values.

We are aware of only one application of using capture-recapture data for formal inference

about landscape resistance for a species (Fuller et al. 2016), which was based on the Ecological

distance SCR model (Royle et al. 2013a, Sutherland et al. 2015). Further, corridor conservation has

been devoid of explicit consideration of local population density. SCR models allow for the

simultaneous estimation of the two processes that are most critical to the conservation of spatially-

structured populations, density and connectivity. Morin et al. (in press) derived a model-based

estimator of landscape connectivity (i.e., density-weighted connectivity) that estimates both the spatial

distribution and connectivity of individuals across a landscape. Spatial optimization approaches that

maximize density-weighted connectivity would identify areas on the landscape that support the

highest number of individuals and best landscape connectivity and would therefore have the greatest

potential for application in corridor conservation and landscape management.

Modeling spatial interactions

The latent point process describing the spatial distribution of individuals is a central component of SCR

methods. Parameterization of this point process allows encounter history data to be used to develop

models that explicitly address theories related to competition, including territoriality (Reich & Gardner

2014) and maintenance of coexisting species and species diversity. Extending the point process model

to account for dependencies among multiple species simultaneously occupying a landscape may

provide an analytic framework for the empirical study of inter and intra-specific competition and

landscape level spatial structure in species assemblages. This should have enormous relevance in

understanding host-pathogen and disease in natural systems where transmission depends on local

interactions of individuals and local density. Where individuals live and who they interact with are

fundamental elements contributing to the dynamics of disease and pathogen systems.

Acoustic sampling

Acoustic sampling is emerging as a promising technology for sampling vocal species such as birds,

anurans, marine mammals, and primates, and the application of these methods is increasing rapidly

(Marques et al. 2009; Blumstein et al. 2011). Information on signal strength and/or direction gives

imperfect information about the source of the vocalization although statistically pinpointing the source

has been recognized as being analogous to inference about the activity center in SCR methods, and

therefore SCR has been adapted to accommodate data obtained by acoustic sampling methods

(Dawson & Efford 2009; Efford et al. 2009; Borchers et al. 2015; Stevenson et al. 2015; Kidney et al.

2016). It seems probable that these technologies will become the *de facto* sampling method in bird

population studies due to the increasing affordability of the technology.

Uncertain identity

Given the widespread adoption of non-invasive sampling technologies, which may only yield partial

information on the identity of individual samples, it will become important to accommodate

uncertainty in individual identity in to studies of animal populations that use individual encounter

histories. There has been considerable attention paid to the problem of uncertain identity in capture-

recapture (Link et al. (2010), Bonner & Holmberg (2013), McClintock et al. (2013)). However, such

methods have developed in the context of classical capture-recapture methods which ignore the

spatial information inherent in most animal population sampling studies. On the other hand, for most

populations we should expect that the spatial location of samples should be informative about the

uncertain identity of those samples (Chandler & Royle 2013, Chandler & Clark 2014, Royle 2015,

Augustine et al. 2016). That is, all other things being equal, spatial samples that are in close spatial

proximity to one another should more likely be of the same individual than samples that are far apart.

Thus, dealing effectively with an uncertain identity of an individual is fundamentally a spatial problem

for which SCR offers a solution.

Methods of accommodating uncertain identity and partially marked populations are promising

avenues for the formal integration of citizen science data collection with population ecology studies

based on capture-recapture. SCR facilitates the use of citizen science in studies of population ecology

because citizen science schemes naturally produce abundant information about individual location,

potentially useful in spatial mark-resight and similar SCR models. Thus, involving citizens in data

collection will produce large quantities of confirmations of species and their locations, but potentially

no individual identity of the observations.

CONCLUSIONS

Two technological advances have influenced the present and future of animal population ecology in a

way that we believe is more profound than any advance in quantitative ecology since the invention of

computers. First is the development of new technologies for obtaining spatial encounter information

on individuals (Box 1). These technologies have revolutionized applied population ecology.

Simultaneous to the development of these new field techniques has been the increasing spatialization

of ecological process models seen in the advancing fields of landscape ecology and metapopulation

ecology, along with the increasing utilization of statistical point process models throughout population

ecology. SCR lies at the convergence of these two technological advances, combining a spatially explicit

observation model that describes data collected using new technologies such as noninvasive genetics

or camera trapping, with spatially explicit models describing how individuals are distributed across a

landscape.

Estimating abundance or population size is one of the most important problems in applied

ecology permitting the evaluation of sophisticated questions in population dynamics (Krebs 1984,

Williams et al. 2002, Sutherland et al. 2013) and providing necessary information for the conservation

and management of important species (Karanth & Nichols, 2002). SCR has become the standard

method for obtaining such information for many species, and is now routinely used to estimate

abundance of populations of conservation concern including species such as tigers (Royle et al. 2009),

grizzly bears (Efford & Mowat 2014; Kendall et al. 2015), wolverines (Magoun et al. 2011, see Box 1)

and jaguars (Sollmann et al. 2011). These and many other species are extremely difficult to capture and

so non-invasive sampling combined with SCR methods are perfectly suited to study these species.

Moreover, many populations exist in such low densities that obtaining sufficient sample sizes of

individuals can be challenging, and thus making the most efficient use of all data, and in particular,

spatial recaptures which are discarded by ordinary capture-recapture, is critically important.

While SCR methods were developed originally as a tool for inference about animal density from

capture-recapture data (Efford, 2004) motivated by the need to address specific technical limitations

of ordinary capture recapture methods (Fig. 1), they have proven to be more than simply an extension

of technique. Spatial capture recapture has profoundly affected the manner in which capture-

recapture is used in studies of animal populations because they allow testing of explicit hypotheses of

core elements of population and landscape ecology by formally integrating technical descriptions of

these processes with encounter history data obtained by sampling animal populations. SCR models

include an explicit model of density and thus relationships can be modeled between density and

landscape features or other population attributes. For example, SCR models can be used to test

hypotheses related to density dependence in animal populations, such as the relationship between

density and home range size (Efford et al. 2015). In addition, understanding movement of individuals

over the landscape is a key objective throughout ecology, and SCR enables the formal integration of

explicit movement, dispersal, and survival models with models of density and other population

characteristics (Ergon & Gardner 2015; Schaub & Royle 2015). Because SCR models are spatially

explicit, we believe it is possible to consider explicit modeling of population dynamic rates as a function

of local density.   Finally, landscape connectivity is a fundamental element of landscape ecology and

explicit models of connectivity can be integrated directly with models of spatial encounter history data

within the SCR framework to provide population-level estimates of connectivity parameters.

Sutherland et al. (2015) and Fuller et al. (2016) develop SCR models in highly structured landscapes and

demonstrate formal inference for an explicit model of landscape connectivity and resistance,

estimated from individual encounter history data from a capture-recapture study of mink.

SCR is not only revolutionizing how population and landscape ecology questions can be

addressed but also in the way we observe populations. For example, use of scat dogs to sample space

using unstructured area searches is extremely practical for studying many species and this method has

grown rapidly in recent years. When data are obtained in this manner, it is imperative that the spatial

structure of sampling be accounted for and SCR accommodates this by explicitly using GPS search lines

in place of trap locations. Acoustic sampling (Dawson & Efford 2009) is a promising new technology for

sampling birds and many other species. However, without a spatially explicit model that describes both

the sampling and underlying process, it is not possible to connect observed acoustic encounter data to

meaningful biological parameters such as population density. Finally, SCR has potential as the

framework for integrating individual encounter data with inexpensive, broad scale auxiliary data such

as from occupancy studies (Chandler & Clark 2014; Whittingham & Chandler *in press*) and potentially

even citizen science programs.

At the core of science is the notion of testing theories by confronting models with data. At the

level of the population this has been recognized as a promise of capture-recapture for several decades

(Nichols 1992), but the use of capture-recapture has not been widely used to address questions related

to within-population spatial structure and population dynamics. However, SCR achieves this promise,

by integrating a formal spatial model describing how individuals are distributed over a landscape, with

a formal spatial model for how the population is sampled. SCR enables testing explicit spatial

mechanisms and processes and improves understanding of spatial ecology from individual encounter

history data. SCR incorporates elements of population structure and dynamics and explicit spatial and

landscape structure to provide a quantitative framework that unifies population and landscape

ecology.

REFERENCES

Allen, A.M. & Singh, N.J. (2016). Linking movement ecology with wildlife management and
conservation. *Front. Ecol. Evol.*, 12, Jan 2016.

Augustine, B., Royle, J.A., Kelly, M., Satter, C., Alonso, R., Boydston, E. & Crooks, K. (2016). Spatial
capture-recapture with partial identity: an application to camera traps. bioRxiv, p.056804.

Blumstein, D. T., Mennill, D. J., Clemins, P., Girod, L., Yao, K., Patricelli, G., *et al.* (2011). Acoustic
monitoring in terrestrial environments using microphone arrays: applications, technological
considerations and prospectus. *J. Appl. Ecol.*, 48, 758–767.

Bonner, S. & Holmberg, J. (2013). Mark-recapture with multiple, non-invasive marks. *Biometrics*, 69,
766-775.

Borchers, D.L. & Efford, M.G. (2008). Spatially explicit maximum likelihood methods for capture-
recapture studies. *Biometrics*, 64, 377-385.

Borchers, D. L. & Kidney, D. (2014). Flexible density surface estimation for spatially explicit capture-
recapture surveys Technical Report, University of St Andrews.

Borchers, D., Distiller, G., Foster, R., Harmsen, B. & Milazzo, L. (2014). Continuous-time spatially
explicit capture–recapture models, with an application to a jaguar camera-trap survey. *Methods Ecol.*
Evol., 5, 656-665.

Borchers, D.L., Stevenson, B.C., Kidney, D., Thomas, L. & Marques, T.A. (2015). A unifying model for
capture–recapture and distance sampling surveys of wildlife populations. *J. Am. Stat. Assoc.*, 110, 195-
204.

Chandler, R.B. & Royle, J.A. (2013). Spatially explicit models for inference about density in unmarked
or partially marked populations. *Ann. Appl. Stat.*, 7, 936-954.

Chandler, R.B. & Clark, J.D. (2014). Spatially explicit integrated population models. *Methods Ecol.*
Evol., 5, 1351-1360.

Chetkiewicz C. B. & Boyce, M.S. (2009). Use of resource selection functions to identify conservation
corridors. *J. Appl. Ecol.*, 46, 1036–1047.

Cooch, E.G., Cam, E. & Link, W. (2002). Occam's shadow: Levels of analysis in evolutionary ecology--
where to next? *J. Appl. Stat.*, 29, 19-48.

Cooch, E. & White, G. (2006). Program MARK: a gentle introduction. Available in pdf format for free
download at http://www. phidot. org/software/mark/docs/book.

Converse, S.J., Block, W.M. & White, G.C. (2006). Small mammal population and habitat responses to
forest thinning and prescribed fire. *Forest Ecol. Manag.*, 228, 263-273.

Dawson, D.K. & Efford, M.G. (2009). Bird population density estimated from acoustic signals. *J. Appl.*
Ecol., 46, 1201-1209.

Dice, L.R. (1938). Some census methods for mammals. *J. Wildl. Manage.*, 2, 119-130.

Dorazio, R.M. & Royle, J.A. (2003). Mixture models for estimating the size of a closed population when
capture rates vary among individuals. *Biometrics*, 59, 351-364.

Driezen K., Adriaensen F., Rondini C., Doncaster C.P. & Matthysen E. (2007). Evaluating least-cost
model predictions with empirical dispersal data: a case-study using radiotracking data of hedgehogs
(Erinaceus europaeus). *Ecol. Model.*, 209, 314–322.

Efford, M. G. (2004). Density estimation in live-trapping studies. *Oikos*, 106, 598-610.

Ellner, S.P., McCauley, E., Kendall, B.E., Briggs, C.J., Hosseini, P.R., Wood, S.N., *et al.* (2001).
Habitat structure and population persistence in an experimental community. *Nature*, 412, 538-543.
591
592 Efford, M. G., Dawson, D.K. & Borchers, D.L. (2009). Population density estimated from locations of
593 individuals on a passive detector array. *Ecology*, 90, 2676-2682.
594
595 Efford, M.G. & Mowat, G. (2014). Compensatory heterogeneity in spatially explicit capture–recapture
596 data. *Ecology*, 95, 1341-1348.
597
598 Efford, M.G., Dawson, D.K., Jhala, Y.V. & Qureshi, Q. (2015). Density-dependent home-range size
599 revealed by spatially explicit capture–recapture. *Ecography*, 39, 676-688.
600
601 Ergon, T. & Gardner, B. (2013). Separating mortality and emigration: modelling space use, dispersal
602 and survival with robust-design spatial capture–recapture data. *Methods Ecol. Evol.*, 5, 1327-1336.
603
604 Fuller, A.K., Sutherland, C.S., Royle, J.A. & Hare, M.P. (2015). Estimating population density and
605 connectivity of American mink using spatial capture-recapture. *Ecol. Appl.*, 26, 1125-1135.
606
607 Gardner, B., Reppucci, J., Lucherini, M. & Royle, J.A. (2010). Spatially explicit inference for open
608 populations: estimating demographic parameters from camera-trap studies. *Ecology*, 91, 3376-3383.
609
610 Getz, W.M. & Saltz, D. (2008). A framework for generating and analyzing movement paths on
611 ecological landscapes. *Proc. Natl. Acad. Sci. U.S.A.*, 105, 19066-19071.
612
613 Gopalaswamy, A.M., Royle, J.A., Hines, J.E., Singh, P., Jathanna, D., Kumar, N. & Karanth, K.U.,
614 2012. Program SPACECAP: software for estimating animal density using spatially explicit capture–
615 recapture models. *Methods Ecol. Evol.*, *3*,1067-1072.
616
617 Gimenez, O., Rossi, V., Choquet, R., Dehais, C., Doris, B., Varella, H., *et al.* (2007). State-space
618 modelling of data on marked individuals. *Ecol. Modell.*, 206, 431-438.
619
620 Gustafson, E.J. & Gardner, R.H. (1996). The effect of landscape heterogeneity on the probability of
621 patch colonization. *Ecology*, 77, 94-107.
622
623 Hanski, I. (1999). *Metapopulation Ecology*. Oxford University Press, 266 pp.
624
625 Hayne, D.W. (1949). Calculation of size of home range. *J. Mammal.*, 30, 1-18.
626
Huffaker, C.B. (1958). Experimental studies on predation: dispersion factors and predator–prey
oscillations. *Hilgardia*, 27, 343–383.

Illian, J., Penttinen, A., Stoyan, H. & Stoyan, D. (2008). *Statistical Analysis and Modelling of Spatial*
Point Patterns (Vol. 70). John Wiley & Sons.

Johnson, D.H. (1980). The comparison of usage and availability measurements for evaluating resource
preference. *Ecology*, *61,* 65-71.

Karanth, K.U. & Nichols, J.D. (1998). Estimation of tiger densities in India using photographic captures
and recaptures. *Ecology*, 79, 2852-2862.

Kāranta, K.U. & Nichols, J.D. eds. (2002). *Monitoring Tigers and Their Prey: a Manual for*
Researchers, Managers, and Conservationists in Tropical Asia. Centre for Wildlife Studies.

Karanth, K.U., Nichols, J.D., Kumar, N. & Hines, J.E. (2006). Assessing tiger population dynamics
using photographic capture–recapture sampling. *Ecology*, 87, 2925-2937.

Kendall, K.C., Macleod, A.C., Boyd, K.L., Boulanger, J., Royle, J.A., Kasworm, W.F., *et al.* (2016).
Density, distribution, and genetic structure of grizzly bears in the Cabinet-Yaak Ecosystem. *J. Wildl.*
Manage., 80, 314-331.

Kidney, D., Rawson, B.M., Borchers, D.L., Stevenson, B.C., Marques, T.A. & Thomas, L. (2016). An
efficient acoustic density estimation method with human detectors applied to gibbons in Cambodia.
PLOS ONE, 11, p.e0155066.

Kot, M. (2001). *Elements of Mathematical Ecology*. Cambridge University Press, 453 pp.

Krebs, C. J. 1985. *Ecology*, 3rd ed. Harper & Row.

Lagrange, P., Pradel, R., Bélisle, M. & Gimenez, O. (2014). Estimating dispersal among numerous sites
using capture–recapture data. *Ecology*, 95, 2316-2323.

Laird, N.M. & Ware, J.H. (1982). Random-effects models for longitudinal data. *Biometrics*, 38, 963-
974.

Langrock, R., King, R., Matthiopoulos, J., Thomas, L., Fortin, D. & Morales, J.M. (2012). Flexible and
practical modeling of animal telemetry data: hidden Markov models and extensions. *Ecology*, 93, 2336-
2342.

Linden, D.W., Fuller, A.K., Royle, J.A. & Hare, M.P. (2016). Estimating the occupancy-density
relationship for a low density carnivore. *J. Appl. Ecol.* (in press).

Link, W.A., Yoshizaki, J., Bailey, L.L. & Pollock, K.H. (2010). Uncovering a latent multinomial:
analysis of mark–recapture data with misidentification. *Biometrics*, 66, 178-185.

MacArthur, R.H. & MacArthur, A.T. (1974). On the use of mist nets for population studies of birds.
Proc. Natl. Acad. Sci. U.S.A., 71, 3230-3233.

Magoun, A.J., Long, C.D., Schwartz, M.K., Pilgrim, K.L., Lowell, R.E. & Valkenburg, P. (2011).
Integrating motion-detection cameras and hair snags for wolverine identification. *J. Wildl. Manag.*, 75,
731-739.

Marques, T.A., Thomas, L., Ward, J., DiMarzio, N. & Tyack, P.L. (2009). Estimating cetacean
population density using fixed passive acoustic sensors: an example with Blainville's beaked whales. *J.
Acoust. Soc. Am.*, 125, 1982-1994.

McLoughlin, P.D., Morris, D.W., Fortin, D., Vander Wal, E. & Contasti, A.L. (2010). Considering
ecological dynamics in resource selection functions. *J. An. Ecol.*, 79, 4-12.

McClintock, B.T., Conn, P.B., Alonso, R.S. & Crooks, K.R. (2013). Integrated modeling of bilateral
photo-identification data in mark–recapture analyses. *Ecology*, 94, 1464-1471.

McRae, B.H. (2006). Isolation by resistance. *Evolution*, 60, 1551-1561.

Morin, D.J., Fuller, A.K., Royle, J.A. & Sutherland, C. (2016). Model-based estimators of density and
connectivity to inform conservation of spatially-structured populations. *Ecosphere* (in press)

Muñoz, D.J., Miller, D.A., Sutherland, C. & Grant, E.H.C. (2016). Using spatial capture-recapture to
elucidate population processes and space-use in herpetological studies. *J. Herpetol.* (in press)

Nichols, J.D. (1992). Capture-recapture models. *BioScience*, 42, 94-102.

Nichols, J.D., Pollock, K.H. & Hines, J.E. (1984). The use of a robust capture-recapture design in small
mammal population studies: a field example with Microtus pennsylvanicus. *Acta Theriol.*, 29, 357-365.

Nichols, J.D. and MacKenzie, D.I., 2004. Abundance estimation and conservation biology. *An. Biod.
Cons.*, 27, 437-439.

Otis, D.L., Burnham, K.P., White, G.C. & Anderson, D.R. (1978). Statistical inference from capture
data on closed animal populations. *Wildl. Monogr.*, 62, 3-135.

Ovaskainen, O. (2004). Habitat-specific movement parameters estimated using mark-recapture data and
a diffusion model. *Ecology*, *85*, 242-257.

Patterson, T.A., Thomas, L., Wilcox, C., Ovaskainen, O. & Matthiopoulos, J. (2008). State–space
models of individual animal movement. *Trends Ecol. Evol. 23*, 87-94.

Poor, E.E., Loucks, C., Jakes, A. & Urban, D.L. (2012). Comparing habitat suitability and connectivity
modeling methods for conserving pronghorn migrations. *PLOS ONE*, 7, p.e49390.

Pradel, R., Choquet, R., Lima, M.A., Merritt, J. & Crespin, L. (2010). Estimating population growth rate
from capture–recapture data in presence of capture heterogeneity. *J. Agric. Biol. Environ. Stat.*, 15, 248-
258.

Proffitt, K.M., Goldberg, J.F., Hebblewhite, M., Russell, R.E., Jimenez, B.S., Robinson, H.S., *et al.*
(2015). Integrating resource selection into spatial capture-recapture models for large carnivores.
Ecosphere, 6, 1-15.

Reich, B.J. & Gardner, B. (2014). A spatial capture-recapture model for territorial species.
Environmetrics, 25, 630-637.

Royle, J.A. (2015). Spatial capture-recapture with partial identity. *arXiv preprint arXiv:1503.06873*.

Royle, J.A. & Young, K.V. (2008). A hierarchical model for spatial capture-recapture data. *Ecology*, 89,
2281-2289.

Royle, J.A., Chandler, R.B., Sollmann R. & Gardner, B. (2014). *Spatial Capture-Recapture*. Academic
Press/Elsevier. 612 pages.

Royle, J.A., Chandler, R.B., Gazenski, K.D & Graves, T.A. (2013a). Spatial capture-recapture models
for jointly estimating population density and landscape connectivity. *Ecology*, 94, 287-294.

Royle, J.A., Chandler, R.B., Sun C.C. & Fuller, A.K. (2013b). Integrating resource selection information
with spatial capture-recapture. *Methods Ecol. Evol.*, 3, 545-554.

Royle, J.A., Kery, M. & Guelat, J. (2011). Spatial capture-recapture models for search-encounter data.
Methods Ecol. Evol., 2, 602-611.

Royle, J.A., Fuller, A.K. & Sutherland, C. (2016). Spatial capture–recapture models allowing Markovian
transience or dispersal. *Popul. Ecol.*, 58, 53-62.
748

749 Sampson, P.D. & Guttorp, P. (1992). Nonparametric estimation of nonstationary spatial covariance
750 structure. *J. Am. Stat. Assoc.*, 87, 108-119.
751
752 Schaub, M., Gimenez, O., Schmidt, B.R. & Pradel, R. (2004). Estimating survival and temporary
753 emigration in the multistate capture-recapture framework. *Ecology*, 85, 2107-2113.
754
755 Schaub, M. & Royle, J.A. (2014). Estimating true instead of apparent survival using spatial Cormack-
756 Jolly–Seber models. *Methods Ecol. Evol.*, 5, 1316-1326.
757
758 Schwartz, M.K., Luikart, G. & Waples, R.S. (2007). Genetic monitoring as a promising tool for
759 conservation and management. *Trends Ecol. Evol.*, 22, 25-33.
760
761 Sollmann, R., Furtado, M.M., Gardner, B., Hofer, H., Jácomo, A.T., Tôrres, N.M. & Silveira, L. (2011).
762 Improving density estimates for elusive carnivores: accounting for sex-specific detection and
763 movements using spatial capture–recapture models for jaguars in central Brazil. *Biol. Cons.*, 144, 1017-
764 1024.
765
766 Sollmann, R., B. Gardner, A.W. Parsons, J.J. Stocking, B.T. McClintock, T.R. Simons, & O'Connell,
767 A.F. (2013). A spatial mark-resight model augmented with telemetry data. *Ecology*, 94, 553-559.
768
769 Stevenson, B.C., Borchers, D.L., Altwegg, R., Swift, R.J., Gillespie, D.M. & Measey, G.J. (2015). A
770 general framework for animal density estimation from acoustic detections across a fixed microphone
771 array. *Methods Ecol. Evol.*, 6, 38-48.
772
773 Sun, C.C., Fuller, A.K. & Royle, J.A. (2014). Trap configuration and spacing influences parameter
774 estimates in spatial capture-recapture models. *PLOS ONE*, 9, e88025.
775
776 Sutherland, C., Fuller, A.K. & Royle, J.A. (2015). Modelling non-Euclidean movement and landscape
777 connectivity in highly structured ecological networks. *Methods Ecol. Evol.*, 6, 69-177.
778
779 Sutherland, W. J., Freckleton, R. P., Godfray, H. C. J., Beissinger, S. R., Benton, T., Cameron, D. D., *et
780 al.* (2013), Identification of 100 fundamental ecological questions. *J. Ecol.*, 101, 58–67.
781
782 Taberlet, P. & Luikart, G. (1999). Non-invasive genetic sampling and individual identification. *Biol. J.
783 Linn. Soc.*, 68, 41-55.
784
785 Taylor, P.D., Fahrig, L., Henein, K. & Merriam, G. (1993). Connectivity is a vital element of landscape
786 structure. *Oikos*, 68, 571-572.
787

788 Tilman, D. & Kareiva, P.M. (1997). Spatial ecology: the role of space in population dynamics and
789 interspecific interactions (Vol. 30). Princeton University Press.
790
791 Turchin, P. (1998). *Quantitative Analysis of Movement: Measuring and Modeling Population*
792 *Redistribution in Animals and Plants* (Vol. 1). Sunderland: Sinauer Associates.
793
794 Turner, M.G., Gardner, R.H. & O'neill, R.V. (2001). *Landscape Ecology in Theory and Practice*. New
795 York: Springer.
796
797 Walpole A.A., Bowman J., Murray D.L., & Wilson P.J. (2012). Functional connectivity of lynx at their
798 southern range periphery in Ontario, Canada. *Land. Ecol.*, 27, 761–773
799
800 Whittington J. & Sawaya, M.A. (2015). A comparison of grizzly bear demographic parameters
801 estimated from non-spatial and spatial open population capture-recapture models. *PLOS ONE*, 10,
802 e0134446.
803
804 Whittington, J., Hebblewhite, M. & Chandler, R.B. (2017). Generalized spatial mark-resight models
805 with an application to grizzly bears. *J. Appl. Ecol.*, (in press).
806
807 Williams, B.K., Nichols, J.D. & Conroy, M.J. (2002). *Analysis and Management of Animal Populations*.
808 Academic Press.
809
810 Wilson, K.R. & Anderson, D.R. (1985a). Evaluation of a nested grid approach for estimating density. *J.*
811 *Wildl. Manag.*, 49, 675-678.
812
813 Wilson, K.R. & Anderson, D.R. (1985b). Evaluation of a density estimator based on a trapping web and
814 distance sampling theory. *Ecology*, 66, 1185-1194.
815
816 Zeller, K.A., McGarigal, K. & Whiteley, A.R. (2012). Estimating landscape resistance to movement: a
817 review. *Land. Ecol.*, 27, 777-797.

<div style="border: 1px solid black; padding: 10px;">

Box 1 – New technologies for generating spatial encounter data

The advent of new field-based methodologies for individual identification allows researchers to collect spatial encounter information on individual animals without the need for physically capturing and marking individuals. Additionally, many of the methods are amenable to citizen science approaches (Dickinson et al. 2010) whereby non-professional scientists are engaged in the collection of data (e.g., camera traps, hair snares), providing increased spatial extent of sampling.

Camera Traps: With improvements in camera technology, there are many commercially available cameras (a) with superior digital technology that provide still photographs and videos to capture species that are elusive and otherwise difficult to capture. Individuals can be identified from photographs for species that possess distinctive natural marks (e.g., Andean bears (b), tigers (c), wolverines (d), bobcats, jaguars, snow leopards, and others).

a

b

c

d

Non-Invasive Genetic Sampling (NGS): NGS allows for the identification of individuals without direct observations via the extraction of DNA from samples. Genetic data can be collected from scat, hair, feathers, shed skin, saliva, and urine. Two common methods of obtaining genetic samples are by using devices that snag hair (i.e., hair snares) (a) and scat detection dogs (b). These methods have been employed on marine and terrestrial mammals (e.g., right whales, black bear, fisher, American mink).

a) Hair Sampling   *b) Scat sampling*

Bioacoustics: Spatially separated microphones or hydrophones can be used to detect species that produce sounds for biological purposes such as defending territories, social calling, and mate attracting. Recent advances in bioacoustics technologies and signal detection and recognition algorithms of spectrographs (left) permit the collection of sounds from species such as mammals, birds, and marine mammals.

</div>

<div style="border:1px solid black">

Box 2 – Core elements of spatial capture-recapture

Spatial encounter history data: Classical capture-recapture summarize spatial data and records only *when* each individual is encountered (a). In practice, data are reduced from a richer 3-d data structure - a record of *when* and *where* each individual was captured (b). Such spatial pattern data are informative about spatial population processes.

a) Non-spatial   b) Spatial

Encounter probability model: SCR models describe encounter probability as a function of the distance between a sample location and *s*, the individual's activity center (the half-normal form is shown to the left). The spatial scale parameter σ accommodates individual heterogeneity in detection due to the juxtaposition of individuals with detectors.

Spatially explicit point process model: Encounter histories are modelled conditional on a latent point process describing the spatial distribution of individuals. The null model of uniformity (a) is typically applied and robust to violations. More realistic models allow individuals to be distributed explicitly according to some covariate (b).

a)   b)

</div>

Box 3 – Modeling resource selection with SCR

Resource selection is a multi-scale process (Johnson 1980), determining the range of a species (1st order selection), the distribution of individuals within their range (2nd order), and the use of habitat by an individual within its home range or territory (3rd order). SCR methods allow for explicit modeling of both 2nd and 3rd order resource selection from encounter history data produced from standard capture-recapture methods such as camera trapping, scat and hair sampling for DNA and live trapping.

2nd order resource selection is the process that governs the placement or location of individual activity centers. SCR models accommodate explicit models for the probability distribution of activity centers s_i , referred to as inhomogeneous point process models, in which the intensity function depends on landscape or habitat structure:

$$log\big(D(s)\big) = \beta_0 \; + \textstyle\sum \beta_k \, X_{sk}$$

At right, mapped tiger density (from Gopalaswamy et al. 2012).

3rd order resource selection affects the SCR encounter probability model (Royle et al. 2013). For some spatially explicit covariate $z(x)$, the probability of encounter can be modeled as a function of both distance and measured covariate with parameter to be estimated:

$$p(x|s) \propto \exp\big(\alpha * z(x) - \; dist(x,s\,)\big).$$

This corresponds to the kernel of standard resource selection models, providing a framework for formal integration of capture-recapture data with data from telemetry studies. At right, estimated encounter probability surface for a black bear population if a trap were placed at a pixel relative to the encounter probability at a pixel of average elevation (from Royle et al. 2013).

Box 4: Estimating landscape connectivity

Landscape structure influences local movements such that, during monitoring, the pattern of individual observations around its activity center is likely to deviate from the assumption of a circular home range. Understanding patterns of space use, and thus estimating encounter rates without bias, requires that the structure of ecological landscapes is explicitly accounted for. Using a least cost path approach, SCR allows the estimation of one or more resistance parameters, δ, characterizing how movement is influenced by landscape structure. Euclidean distance in Eqn. 3 and Box 2 is replaced by *ecological* distance, d_{ecol}, the length of the least cost path between two points (v_0 and v_T):

$$d_{ecol}(v_0, v_T) = \min_{\mathcal{L}_1,\dots,\mathcal{L}_W} \sum_{a=0}^{T} \text{cost}(v_a, v_{a+1}) \times d_{euc}(v_a, v_{a+1}),$$

where $\mathcal{L}_w(v_0, v_T) = \{v_0, \dots, v_T\}$ denotes any path consisting of T adjacent lines connecting adjacent pixels, and $\text{cost}(v_a, v_{a+1})$ is a cost function which is a log linear function of the average pixel-specific covariate values:

$$\log\big(\text{cost}(v_a, v_{a+1})\big) = \delta \frac{z(v_a) + z(v_{a+1})}{2}.$$

Estimation of the resistance parameter provides a direct measure of the strength of species–landscape interactions which has important implications. First, this is a model for asymmetric space use that simultaneously relaxes assumptions of symmetry and stationarity, allowing *home range geometry* to vary depending on location and local landscape structure. Secondly, for a known landscape, the probability of use for any pixel on the landscape can be computed given an individual's location, i.e., individual local connectivity, and it follows therefore that the composite local connectivity surfaces for any collection of activity centers provides a *model-based landscape connectivity* surface informed by the estimate of the species-landscape parameter.

841