Temperature-driven selection on metabolic traits increases the strength of an algal-grazer interaction in naturally warmed streams

Authors
C. -Elisa Schaum[1,4]*, Bio244 Students[1],Richard ffrench-Constant[2], Chris Lowe[1,2], Jón S. Ólafsson[3], Daniel Padfield[1] & Gabriel Yvon-Durocher[1]*

Author affiliations
1 Environment and Sustainability Institute, University of Exeter, Penryn, Cornwall TR10 9EZ, UK

2 Centre for Ecology and Conservation, College of Life and Environmental Sciences, University of Exeter, Penryn, Cornwall, TR10 9FE, U.K.

3. Marine and Freshwater Research Institute, Árleyni 22, 112 Reykjavik, Iceland.

4. Institute for Hydrobiology and Fisheries, Section Biological Oceanography, University of Hamburg, Hamburg, 22767, Germany

* Corresponding authors: bav0352@uni-hamburg.de, g.yvon-durocher@exeter.ac.uk,

Author Contributions:
GYD conceived the experiment. GYD and ES supervised the experiment, analysed data, and wrote the manuscript. All authors contributed to later versions of the manuscript.

Data accessibility statement: All data will be made available as supporting information should the manuscript be accepted. R-Code for analysis will be available from the authors on request.

Type of Article: Primary Research Article
This file includes the supporting information at the end of the file
Number of words in: Abstract: 220;
Main Text without legends and references: 5060
Number of Figures: 3
List: Figure 1: Map and experimental set-up. 2. Thermal reaction norms of grazer metabolic rates. 3. Long and short-term effects of elevated temperature on interaction strength and metabolic rates.
Number of Tables:2
List: 1. Physical and chemical characteristics of geothermal streams. 2. Parameter estimates and output of gnls models.
Number of cited references: 70
Correspondence during submission to: Elisa Schaum bav0352@uni-hamburg.de or
Gabriel Yvon Durocher g.yvon-durocher@exeter.ac.uk

ABSTRACT

Trophic interactions are important determinants of the structure and functioning of

ecosystems. As the metabolism and consumption rates of ectotherms increase sharply with

temperature, there are major concerns that global warming will increase the strength of

trophic interactions, destabilizing food webs, and altering ecosystem structure and function.

We used geothermally warmed streams that span a ~10ºC temperature gradient to investigate

the interplay between temperature-driven selection on traits related to metabolism and

resource acquisition, and the interaction strength between the keystone gastropod grazer, the

wandering snail *Radix balthica,* and a common algal resource. Populations from a warm

stream (~28ºC) had higher maximal metabolic rates and optimal temperatures than their

counterparts from a cold stream (~17ºC). We found that metabolic rates of the population

originating from a warmer stream were higher across all measurement temperatures. A

reciprocal transplant experiment demonstrated that the interaction strengths between the

grazer and its algal resource were highest for both populations when transplanted into the

warm stream. In line with the thermal dependence of respiration, interaction strengths of

grazers from the warm stream were always higher than those of grazers from the cold stream.

These findings suggest that warming can increase the strength of algal-grazer interactions

through the thermodynamic effects of higher temperatures on physiological rates as well as

through correlated increases in *per capita* metabolism and consumption.

Keywords: Consumer-resource interactions, global warming, metabolism, thermal

adaptation, interaction strength

INTRODUCTION

The strength of consumer-resource interactions (e.g. the effect of a consumer on the population density of its prey) plays a critical role in shaping the stability of food webs (May, 1973; Paine, 1980; McCann *et al.*, 1998; Otto *et al.*, 2007). Grazing is an important class of consumer-resource interaction, determining the flux of energy and materials from autotrophs to heterotrophs. There are currently major concerns that global warming will increase the impact of grazers on algal or plant communities because the ingestion and respiration rates of heterotrophs tend to be more sensitive to rising temperatures than rates of photosynthesis and growth in autotrophs (O'Connor, 2009; Gilbert *et al.*, 2014; West & Post, 2016). Stronger interactions have the potential to destabilise food webs and consequently, warming induced increases in interaction strengths could have fundamental implications for ecosystem structure and function. For example, elevated grazing rates in aquatic ecosystems, driven by the mismatch in thermal sensitivity between autotrophs and heterotrophs, are a key driver of projected declines in aquatic primary production over the 21^{st} century in models of ocean biogeochemistry (Laufkötter *et al.*, 2015).

The effects of temperature on metabolic rates and traits associated with consumer-resource interactions (e.g. attack rates, handling times) often follow characteristic unimodal thermal response curves, in which rates increase exponentially to an optimum and decline rapidly thereafter (Dell *et al.*, 2011; Englund *et al.*, 2011; Rall *et al.*, 2012; Dell *et al.*, 2014; Gilbert *et al.*, 2014). Integrating thermal responses for metabolism and interaction-traits with dynamical models of consumer-resource interactions offers a promising framework for predicting food web responses to global warming (Vasseur & McCann, 2005; Shurin *et al.*, 2012; Binzer *et al.*, 2015). However, thermal response curves are often flexible, and can shift when organisms are exposed to novel thermal environments, both via phenotypic plasticity and adaptive evolution (Angilletta *et al.*, 2003; Kingsolver *et al.*, 2004; Deutsch *et al.*, 2008;

Kingsolver & Huey, 2008). Consequently, plasticity and evolution have the potential to

modulate the effects of rising temperatures on the strength of species interactions (Sentis *et*

al., 2015). For example, if metabolic rates are down-regulated after long-term exposure to

higher temperatures (Addo-Bediako *et al.*, 2000), then compensatory metabolic responses to

warming could mitigate predicted increases in consumer-resource interaction strength. How

these long-term responses to warming affect rates of metabolism and in turn, the strength of

consumer-resource interactions, are largely unknown, limiting our ability to predict how

trophic interactions will change in response to warming in the long-term.

There is evidence from studies across naturally occurring thermal gradients over large

spatial scales, that local thermal adaptation can play an important role in shaping the strength

of species interactions (Barton, 2011; De Block *et al.*, 2012). While these studies provide

important insights into how consumer-resource interactions are shaped by evolution across

thermal gradients (Fukami & Wardle, 2005), their usefulness for understanding responses to

rapid climate warming might be limited, because other factors, such as day length, light

intensity and precipitation, tend to be confounded with temperature along such broad scale

spatial gradients. Furthermore, the timescales over which local adaptation has occurred in

such broad scale studies could be much longer than the rapid evolutionary change required to

keep pace with climate warming (Loarie *et al.*, 2009; Hoffmann & Sgrò, 2011). Here, we

investigate how temperature-driven selection on traits that determine the thermal responses of

metabolism and resource acquisition affect the strength of a keystone grazing interaction (the

gastropod *Radix balthica,* which grazes algal biofilms in streams) in naturally warmed

Icelandic geothermal streams spanning a gradient of 11ºC. Critically, temperature is the main

abiotic factor that varies among streams in the catchment and is not correlated with pH,

conductivity or inorganic nutrient concentrations (see Table 1). These streams are thought to

have been subject to geothermal heating for at least the last century (OGorman *et al.*, 2012).

This system therefore provides the opportunity to investigate how long-term differences in

temperature between otherwise similar sites shape the expression of metabolic traits and the

subsequent impact of any temperature-driven selection on species interactions in a natural

system. Specifically, we ask: can long-term differences in temperature drive selection for

metabolic traits that attenuate the direct effects of warming on the strength of consumer-

resource interactions?

121
122   **METHODS**
123
Study site

The streams are located North of the Hveragerði valley, in the south east of the Hengil high

temperature geothermal field, Iceland (N64° 0' 2.944" W21° 11' 17.451") and consist of a

catchment of 11 streams spanning a temperature gradient of approximately 20 ºC (see Figure

1 and SI Figure 1). Two streams, stream 5 (17.5 ºC ± 4.5 ºC, hereafter 'cold stream') and

stream 11A (28.3 ºC ± 1.3 ºC, hereafter 'warm stream', see Table 1 for a comparison of other

chemical and physical parameters), were chosen for experiments due to their close proximity

to each other, the large temperature differential and similar abundances of the keystone

grazer, *Radix balthica*. The grazer plays an important functional role in geothermal stream

ecosystems, where grazer biomass as well as grazing rates are strongly influenced by

temperature (OGorman *et al.*, 2012). The two streams are similar in all other measured

physical and chemical characteristics but differ in average temperature by 11 ºC (see Table

1), and hence present an opportunity to investigate how the effects long-term differences in

temperature shape consumer-resource interactions.

Grazer metabolism

To quantify whether the different thermal regimes in the two adjacent streams resulted in

divergence in metabolic traits of *R. balthica* we measured the acute responses of respiration

to a broad gradient in temperature. We collected 33 individuals of similar weight and length

from each stream, which were cleaned from any algal debris to avoid carry-over of a food

source into the tank or subsequent respiratory measurements on the oxygen electrode. The

snails were kept overnight in aerated tanks at the average stream temperature of origin and in

the absence of a food source to minimise any potential effects of differences in food quantity

or quality between streams. Respiration was quantified as the rate of oxygen consumption in

a Clark-Type oxygen electrode, measured between 4 – 44 ºC in 4 ºC increments (11

temperatures in total). At each temperature, respiration was measured for 3 individuals, and a

different set of individuals was measured at each temperature (i.e. each animal was only

subjected to a single assay). Individuals were allowed 15 minutes at the assay temperature

prior to the measurements. The subsequent thermal responses of respiration were quantified

using a modification of the Sharpe-Schoolfield equation (see (Schoolfield *et al.*, 1981) and

(Sharpe & DeMichele, 1977)for the original equation):

$$\ln\big(b(T)\big) = E_a\left(\frac{1}{kT_c} - \frac{1}{kT}\right) + \ln\big(b(T_c)\big) + \alpha\ln(M_i) - \ln\left(1 + e^{E_h\left(\frac{1}{kT_h} - \frac{1}{kT}\right)}\right) \qquad (1)$$

where $b(T)$, is the *per capita* metabolic rate (μmol O_2 L^{-1} h^{-1}) at temperature T in Kelvin (K),

k is Boltzmann's constant (8.62×10^{-5} eV K^{-1}), E_a is an apparent activation energy (in eV) for

the metabolic process, ln ($b(T_c)$) is the rate of metabolism normalised to an arbitrary

reference temperature, $T_c = 18$ ºC, where no low or high temperature inactivation is

experienced. M_i is the mass (g) of an individual i, α is the allometric scaling exponent that

characterises the power-law relation of mass and metabolic rate (Brown *et al.*, 2004). E_h

characterizes temperature-induced inactivation of enzyme kinetics above T_h where half the

enzymes are rendered non-functional. Differentiating equation (1) and solving for the global

maxima yields an expression for the optimum temperature

$$T_{opt} = \frac{E_h T_h}{E_h + kT_h\ln\left(\frac{E_h}{E_a} - 1\right)} \qquad (2)$$

Equation (1) differs from the Sharpe-Schoolfield equation (Sharpe & DeMichele, 1977;

Schoolfield *et al.*, 1981) in a number of ways. First, we account for the power law relation

between body mass and metabolic rate, M^{α} (Brown *et al.*, 2004). Second, we exclude

parameters from Eq. (1) used to characterize low-temperature inactivation due to insufficient

data to quantify this phenomenon in our analysis. Third, rather than characterize temperature

effects below T_{opt} using the Eyring (1935) relation, $\left(\frac{T}{T_c}\right) e^{E_a\left(\frac{1}{kT_c}-\frac{1}{kT}\right)}$, we instead use the

simpler Boltzmann factor, $e^{E_a\left(\frac{1}{kT_c}-\frac{1}{kT}\right)}$. This simplification enables an explicit solution for T_{opt}

(Eq. 2) and facilitates more direct comparison with previous work on the temperature

dependence of metabolism using metabolic theory e.g. (Allen *et al.*,2005, Gillooly, 2001;

Brown *et al.*, 2004; Van M Savage *et al.*, 2015).

The parameters, ln $b(T_c)$, α, E_a, E_h, T_h, and T_{opt}, in Eqs. (1) & (2) represent traits

characterising the metabolic thermal response that we expect to be under selection in *R.*

balthica inhabiting the hot and cold streams. We tested for differences in each of the

parameters between the populations of *R. balthica* by fitting the respiration data to Eq. (1)

using generalised non-linear least squares regression (within the 'gnls' function in the 'nlme'

package for R, package version 3.1-128) and including 'origin' as a two level factor (i.e.

'cold' and 'warm' stream). We tested for differences between populations for each parameter

by sequentially removing the effect of 'origin' on each parameter and comparing the Akaike

information criterion for small sample sizes (AICc) for all possible models (see SI Table 1

and SI Table 2) using the 'aictab' and 'modavg' functions from the AICcmodavg package

(package version 2.1-0). The model chosen for further exploration was that with the lowest

(AICc) value. Model averaging was carried out when models fell within 2 AICc units of each

other, and the conditional averages of the parameters were used for curve fitting and

interpretation (see also Table 2). The relative importance of the fixed factors in the averaged

model was determined using the sum of their relative weights.

Reciprocal transplant experiment

We carried out a reciprocal transplant experiment to determine how long-term differences in

temperature and the resultant impacts on metabolic traits affect the strength of algal-grazer

interactions. We achieved this by placing snails from each population in microcosms

consisting of a tissue culture flask on which diatom biofilms had been established. Diatoms

of the genera, *Acnanthes, Nitzschia, Navicula*, and *Gomphonema* are common in streams

across the Hengill volcanic area (Gudmundsdottir *et al.* 2013)and were ordered from culture

collections (Culture collection of algae and protozoa and Sciento) and grown in the

laboratory in mixed assemblages to yield common resource for testing the effects of

temperature and local adaptation on grazing . The diatom assemblages were inoculated into

Corning plastic translucent flasks (maximum volume 1L) with 20 mL COMBO medium

(Kilham *et al.*, 1998), and brought to a salinity of 5-10 (equivalent to approximately 5-10 g

salts/kg water) to match the slightly elevated salinity and conductivity found in these thermal

stream environments (Gudmundsdottir et al. 2013). The flasks were turned onto their sides to

allow for a larger area of biofilm growth on the base (~ 60 cm^2 in total per flask) and the algal

communities were left to grow for 14 days prior to the experiment. After 14 days, all flasks

had substantial biofilm development on the base and were used as microcosms for the *in situ*

reciprocal transplant experiment. Analysis of control flasks (no grazer) showed that growth of

the diatom lawn *per se* did not differ significantly for flasks placed in hot or cold streams (SI

Figure 2, one-way ANOVA $F_{1,10} = 1.28$, $P = 0.26$). Thus, any changes to the biofilm biomass

in the experiment can be attributed to the per capita effects of the grazer.

The experiment consisted of 3 treatments (each with 6 replicate microcosms placed in

each of the 2 streams): (i) a control microcosm in which a biofilm was present and no *R.*

balthica were added, (ii) an 'origin' treatment in which *R. balthica* that were resident in the

stream were added to microcosms, and (iii) a 'transplanted' treatment in which *R. balthica*

that were from the adjacent stream were added to microcosms. *R. balthica* individuals were

collected from the 2 streams the day before the experiment and were starved for 24h in the

laboratory in aerated tanks at the average temperature of the stream of origin. There was no

significant difference in average snail weight between the two streams (see SI Figure 3, one-

way ANOVA: $F_{1,408} = 0.15$, $P = 0.7$). Microcosms were assembled by adding 3 snails of

similar body dimensions (0.35 ± 0.03 g of *R. balthica* weight reported as blotted fresh weight

throughout) and 100 mL of 0.4 µm filtered water from the stream in which the microcosm

was to be placed. This resulted in a grazer density of 5 individuals m^{-2}, which was

comparable to the average *in situ* density in the streams (see SI Figure 4, no significant

difference in in situ density between the two-streams: one-way ANOVA: $F_{1,66}$, $P = 0.54$).

This design was preferred to a set-up with each microcosm holding a single grazer, which

attempt to exclude the effects of mutual interference on feeding behaviour e.g. (Skalski &

Gilliam, 2001; Rall *et al.*, 2009; Lang *et al.*, 2011; Vucic-Pestic *et al.*, 2011), because (i) the

experimental densities are representative of natural conditions; and (ii) the consumption rates

of a single individual were insufficient to detect a significant change in algal biomass. The

microcosms were submerged in each stream and the snails were left to graze for 48 hours.

We observed no grazer mortality over the experimental period.

Interaction strength

At the end of the experiment, algal biomass in each of the microcosms was quantified via

methanol chlorophyll extraction modified from (Holm-Hansen & Riemann, 1978). Here, the

walls of the microcosms were scrubbed until all biofilm particles were in suspension. The

solution was filtered onto a 0.4µm GF/F filter, which was then ground in methanol for 5

minutes. The samples were centrifuged at 3500 rpm for 15 minutes and the absorbance of the

supernatant was measured at 632nm, 665nm, and 750nm. Total chlorophyll content in μg

mL^{-1} was then calculated as described in Holm-Hansen & Riemann (1978). The *per capita*

interaction strength in each microcosm was then estimated by calculating the dynamic index

(DI, see also (Berlow *et al.*, 2004)for a technically similar set-up):

$$DI = \frac{\ln\left(\frac{N}{D}\right)}{Yt} \qquad (3)$$

where DI is the dynamic index (g C^{-1} h^{-1}), N is total chlorophyll (sum of Chl a + Chl c)

content of control, D total chlorophyll in the grazed microcosm, Y is the grazer biomass (g

C), and t is time in hours. Snail blotted wet weight was converted to carbon mass (in grams)

using conversion factors that assume dry weight to be 7.5% of the blotted wet weight

(Ricciardi & Bourget, 1998) and a carbon content of 22% dry weight (Burgmer *et al.*, 2010).

We carried out two analyses using the data from the reciprocal transplant experiment.

The first analysis, used a generalised linear model (GLM), with 'interaction strength' as the

response variable and 'origin' ('cold' or 'warm' stream) and 'transplant temperature' (17.5

and 28.3 ºC) as potentially interacting factors. We used this analysis to determine (i) whether

interaction strengths differed between snails that originated from the warm or cold streams

(e.g. a main effect of 'origin'); (ii) whether interaction strengths were temperature dependent

(e.g. a main effect of 'temperature'); and (iii) whether the temperature dependence of

interaction strength differed between the snails from the cold and warm streams (e.g.

interaction between 'origin' and 'temperature').

The design of the reciprocal transplant experiment also enabled us to disentangle

short-term temperature responses attributable to acclimation (e.g. responses to the

temperature in the 'transplanted' stream) from those reflecting processes operating over

longer, time scales (e.g. adaptation to the stream of 'origin'). Note that these 'long-term'

effects, which we call 'adaptation', could reflect strict genetic microevolution (e.g. resulting

in divergent genotypes among populations) or they could represent non-genetic effects of the

different temperature regimes that manifest over ontogenetic development, but are

nevertheless adaptive (Bonduriansky *et al.*, 2011) . In the second GLM we included

'interaction strength' as the response variable and 'timescale' ('short' or 'long') and

'transplant temperature' (17.5 and 28.3 ºC) as potentially interacting factors. Here, 'short-

term' temperature responses were characterised as the change in interaction strength between

the stream of origin and the transplant stream. By contrast, the 'long-term' temperature

response was characterised as the change in interaction strength comparing measurements

made only when the snails were in their stream of origin. For better comparison of the

steepness of the respiration reaction norms, we re-express the transplant temperature data as

Boltzmann temperatures $\left(\frac{1}{kT_c} - \frac{1}{kT}\right)$ so that the coefficients of the model yield activation

energies in units of eV (see Eq. (1)). In this analysis, a significant interaction between

'transplant temperature' and 'timescale' would demonstrate that the temperature dependence

of interaction strength differs between the 'short-term' (E_{short}, change in interaction strength

between the stream of origin and the transplant stream, see also Fig. 3), and 'long-term'

(E_{long}, i.e. change in interaction strength comparing measurements made only when the snails

were in their stream of origin, see also Fig. 3). We assume that E_{short} captures rapid

physiological plasticity (e.g. acclimation) in interaction strength in response to a change in

temperature and E_{long} captures processes operating over longer timescales – e.g. genetic

microevolution and non-genetic developmental effects. Consequently, the component of the

temperature sensitivity attributable to 'adaptation' (recognising that this might be genetically

and/or developmentally determined) is given by $E_{adapt} = E_{long} - E_{short}$.

RESULTS

Metabolic thermal response curves

The allometric scaling coefficient, α, and the apparent activation energy, E_a, were consistent

between the populations of *R. balthica* from the cold and warm streams (see Table 2 for

model comparison and estimated parameter values). The temperature normalised rate of

respiration, ln $b(T_c)$, and T_h (the temperature at which respiration was 50% inactivated) were

both higher in the population of *R. balthica* from the warm stream. Because the optimum

temperature, T_{opt}, depends strongly on T_h (see Eq. (2)), T_{opt} was higher in *R. balthica* from

the warmer stream (T_{opt} warm = 38.25 ± 0.6 ℃; T_{opt} cold = 33.05 ± 1.5 ℃). As ln $b(T_c)$ and

T_{opt} were both higher, the warm populations of *R. balthica* had elevated metabolic rates

across the full range of measurement temperatures (Fig. 2).

Local adaptation of interaction strength

Interaction strength increased with elevated transplant temperature for the populations of *R.*

balthica from both the warm and the cold streams (Fig. 3A; main effect of 'transplant

temperature' GLM $t_{1,21}$ = 2.56; $P < 0.01$). Furthermore, interaction strengths were

consistently higher for the populations of *R. balthica* from the warm stream in both transplant

temperatures (Fig. 3A; GLM main effect of 'origin' $t_{1,121}$ = 2.92; $P < 0.005$). These findings

are consistent with the higher respiration rates observed in the warm population (Fig. 2) and

highlight the association between metabolism and interaction strength.

Disentangling the short- and long-term effects of warming on interaction strengths

Our experimental design enabled us to compare temperature sensitivities that capture short-

term thermal acclimation (e.g. changes in interaction strength in response to the reciprocal

transplant) as well as the long-term temperature sensitivity, which also includes effects of

local adaptation (e.g. changes in rates between warm and cold populations quantified in the

stream of origin). We found that interaction strength increased with temperature in both the

short- and the long-term (Fig. 3). However, the magnitude of the temperature response was

significantly larger in the long-term (Fig. 3; interaction between 'transplant temperature' and

'timescale' on interaction strength; $GLM_{1,18} = -2.91$; $p < 0.05$), where, the average E_{short} was

0.46 eV, while E_{long} was significantly higher at 0.99 eV. This divergence between the short-

and long-term temperature sensitivities implies a non-trivial contribution of adaptation in

amplifying the effects of temperature on interaction strength *in situ*, with the contribution of

E_{adapt} of 0.51 and 0.53 eV in the cold and warm adapted populations respectively.

DISCUSSION

Understanding how global warming will affect the strength of consumer-resource interactions

and the stability of aquatic food webs is a fundamental challenge in evolutionary ecology that

requires insight on the short-term effects of temperature on metabolism and interaction traits

and they are modulated by evolutionary and developmental processes over longer time scales.

There is evidence from terrestrial (Rall *et al.*, 2009; Barton, 2011; Vucic-Pestic *et al.*, 2011;

Brose *et al.*, 2012), freshwater (Kratina *et al.*, 2012) and marine ecosystems (Sanford, 1999),

that warming is likely to increase the strength of consumer-resource interactions, at least in

the short-term, owing to the exponential effects of temperature on the consumption rates of

mobile ectothermic consumers (Dell *et al.*, 2014; Gilbert *et al.*, 2014). What is less clear

however, is how long-term responses to rising temperatures will modulate the direct effects

of warming on species interactions. Space-for-time substitutions across broad spatial scales

indicate that local adaptation to different thermal regimes can play an important role in

shaping species interactions, often compensating for the direct effects of temperature on

interaction traits (Barton, 2011; De Block *et al.*, 2012). Here, we build on this work by

investigating the effects of temperature and local adaptation on the interaction between the

gastropod grazer, *R. balthica*, and its algal resource. Our study contributes novel insights in a

number of ways. First, we explore patterns of local adaptation over a relatively small spatial

scale (m as opposed to km). The two streams in our experiment are separated by

approximately 500 m but differ in temperature by 11 ºC. Because dispersal, gene flow and

genetic divergence among populations in this species are strongly related to geographic

distance (Johansson *et al.*, 2016), our study over a relatively small spatial scale, provides

insight into how metabolic and resource acquisition traits in closely related natural

populations have divergeed in response to warming and is therefore directly relevant for

understanding the effects of rapid climate change (Richter-Boix *et al.*, 2010; Keller *et al.*,

2013; Merilä & Hendry, 2014). Second, we quantified the effects of temperature on both

metabolic and consumption rates to determine how temperature-driven selection on key traits

shape the effects of long-term warming on the strength of consumer-resource interactions.

We found significant variation in the thermal response curves for respiration between

the populations of *R. balthica* from the warm and cold streams. The optimum temperature

(T_{opt}) for respiration was higher in the warm population (i.e. metabolic rates peaked at higher

temperatures). Furthermore, the inactivation energy (E_h) was lower in the warm population,

indicating that declines in the rate of respiration after the optimum (i.e. at high temperatures)

were less pronounced than in grazers from the cold stream, where metabolic rates peaked at

lower temperatures and declined markedly at temperatures above T_{opt}. These divergences

indicate that the different thermal regimes in these streams have selected for different

metabolic traits in warm and cold populations of *R. balthica*. Whilst the higher T_{opt} and lower

E_h in the warm population were in line with expectations assuming local thermal adaptation,

we found no evidence that metabolic performance at high temperature was traded-off against

performance at low temperature. Instead, metabolic rates were higher for *R. balthica* from the

warm stream across all measurement temperatures. These results are in broad agreement with

the "hotter is better" hypothesis, which proposes that maximal performance of organisms

with higher optimal temperatures should be greater than those with lower optimum

temperatures because of the thermodynamic constraints imposed by high temperatures on

enzyme kinetics (Huey & Kingsolver, 1993; Kingsolver *et al.*, 2004; Angilletta *et al.*, 2010).

Indeed maximal respiration rates in the population from the warm stream were greater than

those from the cool (ln(R) warm stream: 3.39 ± 0.14 µmol O_2 L^{-1} h^{-1}, and cool stream: $2.54 \pm$

0.26 µmol O_2 L^{-1} h^{-1}, both $\pm$ 1.s.e.m.). The lower E_h, (i.e. the steepness of the decline of the

thermal reaction norm past the optimum)*,* and higher ln $b(T_c)$, i.e. the rate of respiration

normalised to 18 ºC, in the warm population also meant that the thermal response curve for *R.*

balthica from the warm stream was broader. In agreement with previous work (e.g. on

bacteriophages, (Knies *et al.*, 2009), our data for the gastropod *R. balthica* indicate that

adaptation to higher temperatures resulted in both greater maximal metabolic performance

and a broader metabolic thermal reaction norm.

The general patterns observed in the metabolic traits were also reflected in the effects

of temperature on interaction strength. Interaction strength was higher for individuals placed

in the warm stream, irrespective of their stream of origin. These findings suggest that

elevated temperatures increase consumption rates though the effects of temperature on

respiratory physiology, but local adaptation to warmer environments also results in a

correlated increase in metabolism and interaction strength at low temperature. This may have

important wider implications for the effects of warming on the structure, functioning and

stability of aquatic food webs (Rall *et al.*, 2009; O'Connor *et al.*, 2011; Vucic-Pestic *et al.*,

2011; Dell *et al.*, 2014; Fussmann *et al.*, 2014; Gilbert *et al.*, 2014). If long-term responses to

increasing temperature give rise to higher maximal rates of metabolism and consumption as

well as elevating rates at lower temperatures, then the effects of warming on the strength of

consumer-resource interactions in the long-term could be greater than previously anticipated

(Gilbert *et al.*, 2014). Indeed, work on experimental warming of aquatic ecosystems has

shown that increases in the strength of top-down control can have profound effects on

community structure and ecosystem processes (Burgmer & Hillebrand, 2011; Kratina *et al.*,

2012; Yvon-Durocher *et al.*, 2015). Elevated grazing rates at warmer temperatures can have a

wide range of impacts in aquatic systems, with evidence for both increases (Yvon-Durocher

et al., 2015) and decreases (Burgmer & Hillebrand, 2011) in algal species richness, biomass

and productivity.

In our experiments, the thermal sensitivities of metabolic rates were much larger than

those of interaction strengths in the short-term (e.g. 0.96 and 0.45 eV respectively), in line

with findings in other invertebrate systems (Rall *et al.*, 2009; Vucic-Pestic *et al.*, 2011;

Fussmann *et al.*, 2014). These findings suggest that rates of grazing and metabolism were

clearly linked, but became decoupled when individuals experience rapid changes in

temperature that depart substantially from those in their local environment. In the short-term,

if increases in metabolic demands with temperature are greater than those of consumption

rates (as found here), then less energy will be transferred from the resource to the consumer,

i.e. more is lost through respiration, see also (Rall *et al.*, 2009). If such imbalances are

maintained over long periods of time then starvation of the consumer can ultimately result in

a decline in top-down control on the resource (Fussmann *et al.*, 2014; Binzer *et al.*, 2015).

However, when consumers' feeding rates are more sensitive to temperature than metabolic

rates, interaction strengths can become amplified in warmer environments, leading to faster

resource depletion and eventually driving either the resource or the consumer to extinction

(Vasseur & McCann, 2005). Long-term effects of temperature on interaction strengths have

so far only been explored using food web models, parameterised using temperature

sensitivities derived from short-term experiments (Vasseur & McCann, 2005; Rall *et al.*,

2012; Fussmann *et al.*, 2014). Consequently, such analyses don't capture the evolutionary

and developmental effects which can modulate the short-term effects of temperature on *per*

capita rates. Our results highlight substantial differences between the short- and long-term

effects of temperature on interaction strength; implying that longer term processes plays an

important role in maintaining the balance between metabolic and consumption rates.

418   We quantified the short- and long-term effects of temperature in the reciprocal

transplant experiment. The short-term temperature response (E_{short}) captures the effects of

physiological plasticity over the 48h experiment. Conversely, the long-term response (E_{long})

also accounts for processes operating over longer timescales, including genetic micro

evolution and non-genetic developmental effects of temperature. In our experiment, E_{long} was

higher than E_{short}, implying a significant role for long-term proceses in shaping the effects of

temperature on *in situ* interaction strengths. Notably, the higher E_{long} was driven both by

elevated grazing rates in the warm populations in the warm stream and lower rates in the cold

populations in the cold stream. These results diverge from our expectations based on the

metabolic cold adaptation hypothesis (Addo-Bediako *et al.*, 2000) which would predict

populations from warmer environments should dampen the acute effects of temperature on

metabolic rates. On the contrary, our results suggest that adaptation to warming amplified the

effects of temperature on metabolic as well as grazing rates. The lower interaction strengths

in the population of *R. balthica* from the colder stream highlight unexpected long-term

effects of temperature on species interactions. Maintenance of lower than anticipated grazing

rates in the cold stream could be selected for since lower grazing rates might result in greater

food chain stability and/or stoichiometric homeostasis (Cross *et al.*, 2005; 2014) under the

prevailing temperature regime. Thus, understanding the impacts of environmental change on

the strength of consumer-resource interactions over timescales that are relevant to the rate of

climate change (e.g. gradual warming over decades) will require an appreciation both of the

direct effects of rising temperatures on species interactions and the reciprocal feedback

between ecological and evolutionary dynamics (Fussmann *et al.*, 2007; Gravel *et al.*, 2010;

Loeuille, 2010; Urban, 2013; Barraclough, 2015)

Conclusions

We used a natural geothermal temperature gradient to investigate how warming influences

the strength of algal-grazer interactions via the direct effects of temperature on metabolism

and consumption, and indirect feedbacks through adaptation. Metabolic rates and interaction

strength increased with temperature in the same way for both the warm and cold populations

of *R. balthica*, suggesting that rapid changes in temperature have a consistent effect on

interactions between mobile consumers and sessile resources, mediated by the effects of

temperature on consumer metabolic rates (Dell *et al.*, 2014). However, the warm populations

had higher metabolic and grazing rates across all measurement temperatures compared to

their colder counterparts. These findings are consistent with the 'hotter is better and broader'

hypothesis (Huey & Kingsolver, 1993; Knies *et al.*, 2009; Angilletta *et al.*, 2010) (e.g.

adaptation to warming gives rise to higher maximal metabolic rates and broader thermal

reaction norms). In consequence, our results suggest that warming could increase the strength

of algal-grazer interactions, which are often 'keystone' interactions in aquatic systems, both

via the thermodynamic effects of higher temperatures on enzyme kinetics and through

correlated increases in *per capita* metabolism and consumption as organisms adapt to warmer

temperatures.

Conflict of interest

The authors declare no conflict of interest

Acknowledgments

The authors thank Eoin O'Gorman and for comments on an earlier version of this manuscript.

This study was funded by a Leverhulme Trust research grant (RPG-2013-335), and an ERC

starting grant (ERC-StG 677278) awarded to GYD; and the University of Exeter.

References

Addo-Bediako A, Chown SL, Gaston KJ (2000) Thermal tolerance, climatic variability, and
latitude. *Proceedings of the Royal Society B: Biological Sciences*, **267**, 739–745.
Allen, A. P., J. F. Gillooly, and J. H. Brown. 2005. Linking the global carbon cycle to
individual metabolism. Functional Ecology 19:202–213.
Angilletta MJ Jr, Huey RB, Frazier MR (2010) Thermodynamic Effects on Organismal
Performance: Is Hotter Better? *Physiological and Biochemical Zoology*, **83**, 197–206.
Angilletta MJ Jr, Wilson RS, Navas CA, James RS (2003) Tradeoffs and the evolution of
thermal reaction norms. *Trends in Ecology & Evolution*, **18**, 234–240.
Barraclough TG (2015) How Do Species Interactions Affect Evolutionary Dynamics Across
Whole Communities? *Annual Review of Ecology, Evolution, and Systematics*, **46**, 25–48.
Barton BT (2011) Local adaptation to temperature conserves top-down control in a grassland
food web. *Proceedings of the Royal Society B: Biological Sciences*, **278**, 3102–3107.
Berlow EL, Neutel AM, Cohen JE et al. (2004) Interaction strengths in food webs: issues and
opportunities. *Journal of Animal Ecology*, **73**, 585–598.
Binzer A, Guill C, Rall BC, Brose U (2015) Interactive effects of warming, eutrophication
and size structure: impacts on biodiversity and food-web structure. *Global Change*
Biology, **22**, 220–227.
Bonduriansky R, Crean AJ, Day T (2011) The implications of nongenetic inheritance for
evolution in changing environments. *Evolutionary Applications*, **5**, 192–201.
Brose U, Dunne JA, Montoya JM, Petchey OL, Schneider FD, Jacob U (2012) Climate
change in size-structured ecosystems. *Philosophical Transactions of the Royal Society B:*
Biological Sciences, **367**, 2903–2912.
Brown JH, Gillooly JF, Allen AP, Van M Savage, West GB (2004) Toward A Metabolic
Theory Of Ecology. *Ecology*, **85**, 1771–1789.
Burgmer T, Hillebrand H (2011) Temperature mean and variance alter phytoplankton
biomass and biodiversity in a long-term microcosm experiment. *Oikos*, **120**, 922–933.
Burgmer T, Reiss J, Wickham SA, Hillebrand H (2010) Effects of snail grazers and light on
the benthic microbial food web in periphyton communities. *Aquatic Microbial Ecology*,
61, 163–178.
Cross WF, Benstead JP, Frost PC, Thomas SA (2005) Ecological stoichiometry in freshwater
benthic systems: recent progress and perspectives. *Freshwater Biology*, **50**, 1895–1912.
Cross WF, Hood JM, Benstead JP, Huryn AD, Nelson D (2014) Interactions between
temperature and nutrients across levels of ecological organization. *Global Change*
Biology, **21**, 1025–1040.
De Block M, Pauwels K, Van Den Broeck M, De Meester L, Stoks R (2012) Local genetic
adaptation generates latitude-specific effects of warming on predator-prey interactions.
Global Change Biology, **19**, 689–696.
Dell AI, Pawar S, Savage VM (2014) Temperature dependence of trophic interactions are
driven by asymmetry of species responses and foraging strategy. *J Anim Ecol*, **83**, 70–84.
Dell AI, Pawar S, Savage VM (2011) Systematic variation in the temperature dependence of
physiological and ecological traits. *Proceedings of the National Academy of Sciences*,

108, 10591–10596.

Deutsch CA, Tewksbury JJ, Huey RB, Sheldon KS, Ghalambor CK, Haak DC, Martin PR
(2008) Impacts of climate warming on terrestrial ectotherms across latitude. *Proceedings*
of the National Academy of Sciences, **105**, 6668–6672.

Englund G, Öhlund G, Hein CL, Diehl S (2011) Temperature dependence of the functional
response. *Ecology Letters*, **14**, 914–921.

Fukami T, Wardle DA (2005) Long-term ecological dynamics: reciprocal insights from
natural and anthropogenic gradients. *Proceedings of the Royal Society B: Biological*
Sciences, **272**, 2105–2115.

Fussmann GF, Loreau M, Abrams PA (2007) Eco-evolutionary dynamics of communities
and ecosystems. *Functional Ecology*, **21**, 465–477.

Fussmann KE, Schwarzmüller F, Brose U, Jousset A, Rall BC (2014) Ecological stability in
response to warming. *Nature Climate Change*, **4**, 206–210.

Gilbert B, Tunney TD, McCann KS et al. (2014) A bioenergetic framework for the
temperature dependence of trophic interactions (ed Wootton T). *Ecology Letters*, **17**,
902–914.

Gillooly JF (2001) Effects of Size and Temperature on Metabolic Rate. *Science*, **293**, 2248–
2251.

Gravel D, Bell T, Barbera C, Bouvier T, Pommier T, Venail P, Mouquet N (2010)
Experimental niche evolution alters the strength of the diversity-productivity
relationship. *Nature*, **469**, 89–92.

Gudmundsdottir, R., S. Palsson, E. R. Hannesdottir, J. S. Olafsson, G. M. Gislason, and B.
Moss. 2013. Diatoms as indicators: The influences of experimental nitrogen enrichment
on diatom assemblages in sub-Arctic streams. Ecological Indicators 32:74–81.

Hoffmann AA, Sgrò CM (2011) Climate change and evolutionary adaptation. *Nature*, **470**,
479–485.

Holm-Hansen O, Riemann B (1978) Chlorophyll a Determination: Improvements in
Methodology. *Oikos*, **30**, 438.

Huey RB, Kingsolver JG (1993) Evolution of Resistance to High Temperature in Ectotherms.
The American Naturalist, **142**, S21–S46.

Johansson MP, Quintela M, Laurila A (2016) Genetic divergence and isolation by thermal
environment in geothermal populations of an aquatic invertebrate. *Journal of*
Evolutionary Biology, **29(9)**:1701-12

Keller I, Alexander JM, Holderegger R, Edwards PJ (2013) Widespread phenotypic and
genetic divergence along altitudinal gradients in animals. *Journal of Evolutionary*
Biology, **26**, 2527–2543.

Kilham SS, Kreeger DA, Lynn SG, Goulden CE, Herrera L (1998) COMBO: a defined
freshwater culture medium for algae and zooplankton. *Hydrobiologia*, **377**, 147–159.

Kingsolver JG, Huey RB (2008) Size, temperature, and fitness: three rules. *Evolutionary*
Ecology Research, **10**, 251–268.

Kingsolver JG, Ragland GJ, Shlichta JG (2004) Quantitative genetics of continuous reaction
norms: thermal sensitivity of caterpillar growth rates. *Evolution*.

Knies JL, Kingsolver JG, Burch CL (2009) Hotter Is Better and Broader: Thermal Sensitivity
of Fitness in a Population of Bacteriophages. *The American Naturalist*, **173**, 419–430.

Kratina P, Greig HS, Thompson PL (2012) Warming modifies trophic cascades and
eutrophication in experimental freshwater communities. *Ecology*, **93**, 1421–1430.

Lang B, Rall BC, Brose U (2011) Warming effects on consumption and intraspecific
interference competition depend on predator metabolism. *Journal of Animal Ecology*, **81**,
516–523.

Laufkötter C, Vogt M, Gruber N et al. (2015) Drivers and uncertainties of future global

marine primary production in marine ecosystem models. *Biogeosciences*, **12**, 6955–6984.

Loarie SR, Duffy PB, Hamilton H, Asner GP, Field CB, Ackerly DD (2009) The velocity of climate change. *Nature*, **462**, 1052–1055.

Loeuille N (2010) Influence of evolution on the stability of ecological communities. *Ecology Letters*, **13**, 1536–1545.

May RM (1973) Qualitative stability in model ecosystems. *Ecology*, **54**, 638–641.

McCann K, Hastings A, Huxel GR (1998) Weak trophic interactions and the balance of nature. *Nature*, **395**, 794–798.

Merilä J, Hendry AP (2014) Climate change, adaptation, and phenotypic plasticity: the problem and the evidence. *Evolutionary Applications*, **7**, 1–14.

O'Connor MI (2009) Warming strengthens an herbivore–plant interaction. *Ecology*, **90**, 388–398.

O'Connor MI, Gilbert B, Brown CJ (2011) Theoretical predictions for how temperature affects the dynamics of interacting herbivores and plants. *The American Naturalist*, **178**, 626–638.

OGorman EJ, Pichler DE, Adams G et al. (2012) *Impacts of Warming on the Structure and Functioning of Aquatic Communities: Individual- to Ecosystem-Level Responses*, 1st edn, Vol. 47. Elsevier Ltd., 96 p.

Otto SB, Rall BC, BROSE U (2007) Allometric degree distributions facilitate food-web stability. *Nature*, **450**, 1226–1229.

Paine RT (1980) Food webs: linkage, interaction strength and community infrastructure. *Journal of Animal Ecology*, **49**, 666.

Vucic-Pestic O, Ehnes RB, Rall BC (2011) Warming up the system: higher predator feeding rates but lower energetic efficiencies. *Global Change Biology, 17, 1301–1310*

Rall BC, Brose U, Hartvig M, Kalinkat G, Schwarzmuller F, Vucic-Pestic O, Petchey OL (2012) Universal temperature and body-mass scaling of feeding rates. *Philosophical Transactions of the Royal Society B: Biological Sciences*, **367**, 2923–2934.

Rall BC, Vucic-Pestic O, Ehnes RB, Emmerson M, Brose U (2009) Temperature, predator-prey interaction strength and population stability. *Global Change Biology*, **16**, 2145–2157.

Ricciardi A, Bourget E (1998) Weight-to-weight conversion factors for marine benthic macroinvertebrates. *Marine Ecology Progress Series*, **171**, 245–251.

Richter-Boix A, Teplitsky C, Rogell B, Laurila A (2010) Local selection modifies phenotypic divergence among Rana temporariapopulations in the presence of gene flow. *Molecular Ecology*, **19**, 716–731.

Sanford E (1999) Regulation of keystone predation by small changes in ocean temperature. *Science*, **283**, 2095–2097.

Schoolfield RM, Sharpe PJH, Magnuson CE (1981) Non-linear regression of biological temperature-dependent rate models based on absolute reaction-rate theory. *Journal of Theoretical Biology*, **88**, 719–731.

Sentis A, Morisson J, Boukal DS (2015) Thermal acclimation modulates the impacts of temperature and enrichment on trophic interaction strengths and population dynamics. *Global Change Biology*, **21**, 3290–3298.

Sharpe PJ, DeMichele DW (1977) Reaction kinetics of poikilotherm development. *Journal of Theoretical Biology*, **64**, 649–670.

Shurin JB, Clasen JL, Greig HS, Kratina P, Thompson PL (2012) Warming shifts top-down and bottom-up control of pond food web structure and function. *Philosophical Transactions of the Royal Society B: Biological Sciences*, **367**, 3008–3017.

Skalski GT, Gilliam JF (2001) Functional Responses With Predator Interference: Viable Alternatives To The Holling Type II Model . *Ecology*, **82**, 3083.

Urban MC (2013) Evolution mediates the effects of apex predation on aquatic food webs.
Proceedings of the Royal Society B: Biological Sciences, **280**, 20130859–20130869.
Van M Savage, Gillooly JF, Brown JH, West GB, Charnov EL (2015) Effects of Body Size
and Temperature on Population Growth. *The American Naturalist*, **163**, 429–441.
Vasseur DA, McCann KS (2005) A mechanistic approach for modeling temperature-
dependent consumer-resource dynamics. *The American Naturalist*, **166**, 184–198.
West DC, Post DM (2016) Impacts of warming revealed by linking resource growth rates
with consumer functional responses (ed Behmer S). *Journal of Animal Ecology*, **85**, 671–
680.
Yvon-Durocher G, Allen AP, Cellamare M et al. (2015) Five Years of Experimental
Warming Increases the Biodiversity and Productivity of Phytoplankton (ed Levin SA).
PLoS Biology, **13**, e1002324–22.
622

Tables

Table 1. Physical and chemical characteristics of the streams. Temperature data were
collected over a 3-day period. All other parameters were collected on the first day of the day
of the experiment. Temperature data are displayed as means ± 1SD. All other data were
originally collected for correlation with temperature across the catchment area (all 11
streams), so that replication was on the level of stream identity.
630

Parameter	Stream 5	Stream 11
Average temperature (^{o}C, 5 days)	17.5 ± 4.5	28.3 ± 1.3
pH	7.63	7.17
Conductivity	273.6	235.7
NO_2	0.22	0.24
NO_3	0.57	0.29
NH_4	0.17	0.19
PO_4	0.27	0.35

Table 2. Parameter estimates and output from the best fitting gnls model to the thermal
response curves of respiration rates. Differences in treatments are given in **bold.** Parameter
estimates are taken from the averaged generalised linear models along with their standard
errors (± 1 s.e.m). C = cold stream. W = warm stream. See Supporting Information for details
on model selection and information on AICsc scores for all possible models. Here, the model
average of the conditional average output for the four best models (within 2 AICc units of
each other) is displayed.
641

Non-linear mixed model output for respiration rates (R)		
Treatment effect on	Estimate	± 1 s.e.m.
E_a	C: 0.96	0.05
ln $R(T_c)$	**C: 1.77**	**0.15**
	W: 2.04	**0.12**
E_h	**C: 5.01**	**0.97**
	W: 3.16	**0.96**
T_h [K]	**C: 307.16 (34.01)**	**0.94**
(°C)	**W: 314.15**	
	(41.00)	**1.69**
		0.78
α	0.36	0.03

FIGURE 1 Map and experimental set-up. **Top panel**: The catchment area, with streams used in this experiment indicated by black (for the colder stream 5 with 17.5 ºC ± 4.5 ºC) and red (for the warmer stream 11A with 28.3 ºC ± 1.3 ºC) snail icons. **Lower panel**: Schematic overview of experimental set-up for the grazing experiment.

646
FIGURE 2 Thermal response curves for respiration. Thermal response curves of respiration rates in μmol
O2 L^{-1} h^{-1} as a function of increasing temperature for populations of the snail *Radix balthica* from the cold
(black) and warm (red) stream. Lines are derived from fitting a modified Sharpe-Schoolfied equation (see
methods) to the rate data. Snails from the warm stream have higher temperature normalised metabolic rates (ln
$R(T_c)$, with T_c=18ºC) at all measurement temperatures and have higher optimal temperatures (T_{opt}), than snails
from the cold stream. The inactivation energy (E_h) is lower in snails from the warm stream, resulting in a curve
that is both broader and elevated in comparison to the thermal response curve of respiration for snails from the
cold stream.
655

FIGURE 3 Long-term and short-term effects of stream temperature on interaction strength Long-term
and short-term effects of temperature in interaction strength measured via the dynamic index in units of
chlorophyll consumed per hour. Populations originating from the warm stream have stronger interaction
strength indices in all environments and the highest dynamic index overall was found for snails from the warm
stream in their original environment. Interaction strength increased with temperature both in the short-term
(E_{short}, dashed blue lines) and in the long-term (E_{long}, solid blue line), with E_{long} significantly greater than E_{short}.
663

Supporting Information

for

Temperature-driven selection on metabolic traits increases the strength of an algal-grazer interaction in naturally warmed streams

Authors

C. -Elisa Schaum[1]*, Richard ffrench-Constant[2], Chris Lowe[1,2], Jón S. Ólafsson[3], Daniel Padfield[1] & Gabriel Yvon-Durocher[1]*

Author affiliations

1 Environment and Sustainability Institute, University of Exeter, Penryn, Cornwall TR10 9EZ, UK

2 Centre for Ecology and Conservation, College of Life and Environmental Sciences, University of Exeter, Penryn, Cornwall, TR10 9FE, U.K.

3. Marine and Freshwater Research Institute, Árleyni 22, 112 Reykjavik, Iceland.

* Corresponding authors: g.yvon-durocher@exeter.ac.uk, bav0352@uni-hamburg.de

List of SI Items:
SI Figure 1: The catchment area and experimental set-up.
SI Figure 2: Chlorophyll content in control microcosms.
SI Figure 3: Snail weight for 205 randomly selected snails from each stream
SI Figure 4: Grazer density (snails/m$^{2)}$) in stream 5 (17.5ºC) and stream 11 (28.3ºC)

SI Table 1: AICc comparison details for model simplification of the gnls models fitted to the thermal response data for respiration rates.

692
693
694 **Figure S1: the catchment area and experimental set-up.** A: View of representative stream
695 in the Hveragerði catchment area. B and C: Microcosms fixed in stream 5 (Microcosm
696 pictures courtesy of B. Flello).
697

702 **Figure S2: Chlorophyll content in control microcosms.** Chlorophyll content at the end of
703 the experiment in control microcosms did not differ significantly between cold and warm
704 streams (one-way ANOVA $F_{1,10}$=1.28, p=0.26) implying growth and mortality of the biofilms
705 was consistent in both treatments.
706

707
708

709
710
711 **Figure S3: Snail weight for 205 randomly selected snails from each stream.** Snail wet
712 weight did not vary significantly between the cold (black) and warm (red) streams (one-way
713 ANOVA: $F_{1,408} = 0.15$, p = 0.7).

714
715 **Figure S4: *In-situ* grazer density in the experimental streams.** Snails were counted in 34
716 randomly distributed 1m^2 quadrats. There was no significant difference in snail abundance
717 between streams (one-way ANOVA: $F_{1,66}$, p = 0.54). The mean abundance of 5.5
718 individuals per square meter was comparable to the densities established in the experimetal
719 microcosms. Snails used for the experiments were sampled randomly from these patches.
720

Table S1. AICc comparison details for model simplification of the gnls models fitted to the thermal response data for respiration rates.
Equation 1 was used to account for effects of mass (alpha in the model) on metabolic rate, and fitted to data for respiration (ln R) to characterize acute thermal responses. Model selection was carried out by fist fitting the most complex model to the data and then calculating AICc scores for all possible models with all possible treatment effects, and comparing the models through the aictab function within the AICmodav package (version 2.1-0) in R. The most parsimonious model included differences between warm and cold populations in ln $R(T_c)$, T_h, and E_h, while E_a and α were consistent between populations. For the four models with a difference in AICc of less than two, models were averaged using the modelavg function in the same package. **Lower part of table: Model averaged parameter estimates, with 95% confidence intervals (95%CI) and relative importance as sum of AICc based relative weights (SW), for models where Δ AICc <2.**

Model name	Model details	K	AICC	Dellta AICC	AICc weights	cumulative weights	LogLik
r.gnls19	**alpha+Ea+Eh~ 1, ln.b0+Th~1 + StreamT**	**8**	**84.98**	**0**	**0.26**	**0.26**	**-33.13**
r.gnls7	**ln.b0 +Ea~ 1, alpha+ Eh + Th ~ 1 + StreamT**	**9**	**86.15**	**1.17**	**0.14**	**0.4**	**-32.34**
r.gnls10	**Ea +alpha~ 1,ln.b0+ Eh + Th ~ 1 + StreamT**	**9**	**86.37**	**1.39**	**0.13**	**0.53**	**-32.45**
r.gnls22	**ln.b0+alpha+Ea+Eh~ 1, Th~1 + StreamT**	**7**	**86.51**	**1.53**	**0.12**	**0.65**	**-35.22**
r.gnls13	alpha+Eh~ 1,ln.b0+ Ea+ Th ~ 1 + StreamT	9	87.68	2.7	0.07	0.72	-33.11
r.gnls9	alpha+ Ea + Eh ~ 1 + StreamT,ln.b0 +Th~ 1	9	88.24	3.26	0.05	0.77	-33.39
r.gnls16	ln.b0+alpha+Ea~ 1,Eh+Th~ 1 + StreamT	8	88.81	3.83	0.04	0.8	-35.05
r.gnls1	Ea ~ 1, ln.b0+alpha + Eh + Th ~ 1 + StreamT	10	88.97	3.99	0.04	0.84	-32.33
r.gnls2	ln.b0 ~ 1, alpha +Ea+ Eh + Th ~ 1 + StreamT	10	89.01	4.03	0.03	0.87	-32.35
r.gnls17	ln.b0+alpha+Eh~ 1,Ea+Th~ 1 + StreamT	8	89.12	4.14	0.03	0.91	-35.2
r.gnls3	alpha ~ 1, ln.b0+Ea+ Eh + Th ~ 1 + StreamT	10	89.21	4.23	0.03	0.94	-32.45
r.gnls4	Eh ~ 1, ln.b0+alpha+Ea + Th ~ 1 + StreamT	10	90.02	5.04	0.02	0.96	-32.85
r.gnls5	ln.b0+alpha+Ea+Eh ~ 1 + StreamT,Th ~ 1	10	91.04	6.06	0.01	0.97	-33.36
r.gnls12	ln.b0+ alpha+ Eh ~ 1 + StreamT,Ea +Th~ 1	9	91.24	6.25	0.01	0.98	-34.89
r.gnls6	ln.b0 +alpha~ 1, Ea+ Eh + Th ~ 1 + StreamT	9	91.6	6.61	0.01	0.99	-35.07
r.gnls	Ful model	11	91.97	6.99	0.01	1	-32.34
r.gnls23	Eh~1 + StreamT,ln.b0+alpha+Ea+Th~ 1	7	99.12	14.14	0	1	-41.52

r.gnls18	Ea+Eh~ 1 + StreamT,ln.b0+alpha+Th~ 1	8	102.36	17.38	0	1	-41.82
r.gnls20	ln.b0+Ea~1 + StreamT,alpha+Eh+Th~ 1	8	105.68	20.69	0	1	-43.48
r.gnls8	ln.b0 +Eh~ 1, alpha+ Ea + Th ~ 1 + StreamT	9	106.1	21.12	0	1	-42.32
r.gnls15	ln.b0+ alpha+Ea ~ 1 + StreamT,Eh+Th~ 1	9	107.84	22.86	0	1	-43.19
r.gnls26	Ea~1 + StreamT,ln.b0+alpha+Eh+Th~ 1	7	113.1	28.12	0	1	-48.51
r.gnls25	alpha~1 + StreamT,ln.b0+Ea+Eh+Th~ 1	7	115.95	30.97	0	1	-49.94
r.gnls24	ln.b0~1 + StreamT,alpha+Ea+Eh+Th~ 1	7	116.79	31.81	0	1	-50.36
r.gnls21	ln.b0+alpha~1 + StreamT,Ea+Eh+Th~ 1	8	118.31	33.33	0	1	-49.8
r.gnls27	ln.b0+alpha+Ea+Eh+Th~ 1	6	131.06	46.08	0	1	-58.77
r.gnls11	Ea +Eh~ 1,ln.b0+ alpha+ Th ~ 1 + StreamT	no convergence					
r.gnls14	ln.b0+ Ea+ Eh ~ 1 + StreamT,alpha+Th~ 1	no convergence					

For AICc <2 (bold in upper part of table)							
Parameter	**lnR(Tc)**		**Alpha**	**Ea**	**Eh**		**Th [K]**	
Stream temperature (ºC)	17.5	28.3	No treatment effect	No treatment effect	17.5	28.3	17.5	28.3
SW (sum of weights)	0.6		0.22	0	0.42		1.00	
Parameter Estimate	1.77	2.04	0.36	0.96	5.01	3.16	307.16	314.15
95% CI [lower,upper]	[0.98-2.77]	[1.09-3.28]	[0.34-0.76]	[0.85-1.07]	[3.07-6.96]	[2.96-5.11]	[305.74-309.99]	[309.04-319.13]

732
733