Double maternal effect: duplicated nucleoplasmin 2 genes, *npm2a* and *npm2b*, are

shared by fish and tetrapods, and have distinct and essential roles in early

embryogenesis

Caroline T. Cheung[1, #], Jérémy Pasquier[1, #], Aurélien Bouleau[1], Thao-Vi Nguyen[1], Franck

Chesnel[2], Yann Guiguen[1], and Julien Bobe[1]*

[1]INRA LPGP UR1037, Campus de Beaulieu, 35042 Rennes, FRANCE.

[2]CNRS/ UMR6290, Université de Rennes 1, 35000 Rennes, FRANCE.

* Corresponding author E-mail: julien.bobe@inra.fr

Authors contributed equally to this work: #

Short Title: *npm2* genes have maternal effect in vertebrates

Abstract

Nucleoplasmin 2 (*npm2*) is an essential maternal-effect gene that mediates early embryonic events through its function as a histone chaperone that remodels chromatin. Here we report the existence of two *npm2* (*npm2a* and *npm2b*) genes in zebrafish. We examined the evolution of *npm2a* and *npm2b* in a variety of vertebrates, their potential phylogenetic relationships, and their biological functions using knockout models via the CRISPR/cas9 system. We demonstrated that the two *npm2* duplicates exist in a wide range of vertebrates, including sharks, ray-finned fish, amphibians, and sauropsids, while *npm2a* was lost in Coelacanth and mammals, as well as some specific teleost lineages. Using phylogeny and synteny analyses, we traced their origins to the early stages of vertebrate evolution. Our findings suggested that *npm2a* and *npm2b* resulted from an ancient local gene duplication, and their functions diverged although key protein domains were conserved. We then investigated their functions by examining their tissue distribution in a wide variety of species and found that they shared ovarian-specific expression, a key feature of maternal-effect genes. We also showed that both *npm2a* and *npm2b* are maternally-inherited transcripts in vertebrates. Moreover, we used zebrafish knockouts to demonstrate that *npm2a* and *npm2b* play essential, but distinct, roles in early embryogenesis. *npm2a* functions very early during embryogenesis, at or immediately after fertilization, while *npm2b* is involved in processes leading up to or during zygotic genome activation. These novel findings will broaden our knowledge on the evolutionary diversity of maternal-effect genes and underlying mechanisms that contribute to vertebrate reproductive success.

Author Summary

The protein and transcript of the *npm2* gene have been previously demonstrated as maternal contributions to embryos of several vertebrates. Recently, two *npm2* genes, denoted here as *npm2a* and *npm2b*, were discovered in zebrafish. This study was conducted to explore the evolutionary origin and changes that occurred that culminated in their current functions. We found that an ancient local duplication of the ancestral *npm2* gene created the current two forms, and while most vertebrates retained both genes, notably, mammals and certain species of fish lost *npm2a* and, albeit rarely, both *npm2a* and *npm2b*. Our functional analyses showed that *npm2a* and *npm2b* have diverse but essential functions during embryogenesis, as *npm2a* mutants failed to undergo development at the earliest stage while *npm2b* mutants developed, although abnormally, until the zygotic genome activation stage after which their development was arrested followed subsequently by death. Our study is the first to clearly demonstrate the evolution, diversification, and functional analyses of the *npm2* genes, which are essential maternal factors that are required for proper embryonic development and survival.

Introduction

In animals and plants, early embryonic development relies strictly on maternal products until maternal-to-zygotic transition (MZT) during which zygotic genome activation (ZGA) occurs [1]. Maternal-effect genes are those that are transcribed from the maternal genome and whose products, which include transcripts, proteins, and other biomolecules, are deposited into the oocytes during their production in order to coordinate embryonic development before ZGA [2]. MZT is a key step that is needed firstly for clearance of maternal components, and secondly to activate zygotic gene expression and to allow subsequent embryonic development. Among the maternally-inherited transcripts that play

important roles during early development, some genes were demonstrated to regulate zygotic

program activation such as *nanog*, *pou5f1*, and *soxb1* in zebrafish (*Danio rerio*) [3].

Henceforth, all gene and protein nomenclature will be based on that of zebrafish regardless of

species for simplification purposes. Those factors were shown to participate in the regulation

of "first wave" zygotic genes and among them, mir-430, a conserved microRNA that has

been shown to be involved in clearance of maternal mRNAs in zebrafish [4][5], as well as

mir-427, its orthologue in Xenopus [6]. microRNAs may also have maternal effects by

functioning in a similar manner to protein-coding genes, and recent revelations showed that

there is a subset of microRNAs that are predominantly expressed in the fish ovary and may

function in oogenesis and early embryogenesis [7]. Another gene, *nucleoplasmin 2* (*npm2*),

belongs to the family of nucleoplasmins/nucleophosmins that is maternally-inherited at both

protein and mRNA levels, whereby both play important roles in early development [8].

Historically, this protein was identified and defined as a nuclear chaperone in Xenopus

[9,10]. While the protein has been shown to be the most abundant nuclear protein in the

Xenopus oocyte [11] and to play a crucial role at fertilization due to its role in sperm

chromatin decondensation [12,13], its maternally-inherited mRNA has been recently

demonstrated to be translated as a *de novo* synthesized protein that could play a crucial role

during ZGA in zebrafish [8]. Further, *npm2* is one of the first identified maternal-effect genes

in mouse whereby its deficiency results in developmental defects and eventual embryonic

mortality [14].

The *npm2* gene belongs to the *npm* gene family that encompasses four members,

npm1, npm2, npm3, and npm4. The diversity of the Npm family has been shown to result

from the two rounds of whole genome duplication (WGD) that occurred in early vertebrates

(vertebrate genome duplication 1 and 2, or VGD1 & VGD2, respectively) [15,16]. Former

evolutionary studies clearly provided a phylogenic model of this family; VGD1 produced two

genes, *npm3/2* and *npm4/1*, from an ancestral *npm* gene and the following WGD, VGD2,

further created the current four *npm* types with subsequent loss of *npm4* from mammals, but

retained in most fish species [17–20]. Recently, two *npm2* genes were automatically

annotated in the zebrafish genome, i.e. *npm2a* (ENSDARG00000076391) and *npm2b*

(previously known as *npm2*, ENSDARG00000053963). As the teleost ancestor experienced

an extra WGD event (TGD, or teleost-specific genome duplication) [21], doubling of genes

and other types of genomic rearrangement may be present in teleost species compared with

other vertebrates. Moreover, a fourth round of duplication occurred more recently in

salmonids (salmonid-specific genome duplication or SaGD) [22,23], leading to further

possible doubling of genes and other genomic rearrangements. As multiple evolutionary

events (impact of TGD, local duplication, teleost-specific duplication, etc.) could have led to

the *npm2a*/*npm2b* diversity in zebrafish, which in turn may have significant impact on

vertebrate reproduction, we investigated the evolution of *npm2a* and *npm2b* in a wide range

of vertebrate species, their potential phylogenetic relationship, as well as their biological

functions using transgenic zebrafish models created by the CRISPR/cas9 system in order to

broaden our knowledge on the evolutionary diversity of maternal-effect genes and the

underlying mechanisms that contribute to reproductive success in vertebrates.

Results and Discussion

As previously shown, *npm1*, *npm2*, *npm3* and *npm4* genes are thought to have

originated from the first two rounds of WGD, VGD1 and VGD2, which occurred early on in

vertebrate evolution [17–20]. However, two *npm2* genes have been automatically annotated

in the zebrafish genome, *i.e. npm2a* and *npm2b*. In order to verify if these two *npm2* genes

are paralogous to each other and to determine their origination, we used a Blast search

approach in various public databases to retrieve a multitude of sequences that could be related

to *npm2* genes. All retrieved sequences are compiled in Supplemental Table S1.

Diversity of npm2a and npm2b in vertebrates

<u>Phylogenetic analysis</u>

In order to verify that the retrieved protein sequences (Supplemental Table S1) were

homologous to zebrafish Npm2a and Npm2b, a phylogenetic analysis on Npm2 was

performed. Based on the alignment of 76 vertebrate Npm2-related sequences, and using

vertebrate Npm1 and Npm3 amino acid sequences as out-groups, a phylogenetic tree was

generated (Fig. 1). As shown in Fig. 1, vertebrate Npm2-related sequences clustered into two

clades, Npm2a and Npm2b, which were supported by significant bootstrap values, 84.2% and

78.2%, respectively.

There were 28 sequences that clustered in the Npm2a clade, which encompassed

sequences belonging to species from various vertebrate groups, including chondrichthyans

(such as dogfish Npm2a), ray-finned fish, and sarcopterygians (lobe-finned fish and

tetrapods). The sequences belonging to ray-finned fish included gar and bowfin Npm2a, as

well as teleost sequences such as zebrafish, northern pike, and rainbow trout Npm2a. In

contrast, no Npm2a sequence was identified in neoteleostei (medaka, European perch, and

Atlantic cod). The sequences belonging to sarcopterygians included amphibian (Xenopus)

Npm2a, as well as sauropsid sequences such as Chinese alligator, chicken, penguin, and ibis

Npm2a, although no Npm2a sequence could be identified in mammals. These results

provided the first evidence for the presence of orthologs to zebrafish Npm2a across

evolutionary divergent vertebrate groups (i.e. sauropsids, amphibians, and fish).

The Npm2b clade included 48 of the retrieved sequences. Like the Npm2a clade, the

Npm2b clade also encompassed sequences belonging to species from all of the main

vertebrate groups, including chondrichthyans (such as dogfish Npm2b), ray-finned fish, and

sarcopterygians. Ray-finned fish sequences included gar and bowfin Npm2b in addition to

teleost sequences such as zebrafish and northern pike Npm2b. Our analysis also demonstrated

the existence of two paralogous Npm2b proteins (*i.e.* Npm2b1 and Npm2b2) in all of the

investigated salmonid species (rainbow trout, brown trout, and brook trout). In contrast, no

Npm2b sequence could be identified in any neoteleostean species, including medaka, cod,

perch, fugu, tetraodon, and stickleback. The sequences belonging to sarcopterygians included

amphibian sequences such as Xenopus Npm2b, sauropsid sequences such as Chinese

alligator, ostrich, penguin and chicken Npm2b, as well as mammalian sequences such as

human and mouse Npm2. These results demonstrated, for the first time, that proteins

previously reported as Npm2 (Xenopus, cattle, mouse, human, zebrafish) are orthologous to

Npm2b in all investigated vertebrate species, and should therefore now be referred to as

Npm2b. In addition, with the presence of Npm2b1 and Npm2b2 in all investigated salmonid

species, we provided for the first time evidence of the existence of two Npm2b protein forms

in vertebrate species.

The existence of the two Npm2 clades indicated that Npm2a and Npm2b are likely

paralogous to each other. In addition, comparison of Npm2a and Npm2b amino acid

sequences in the species that harbor both revealed that they share between 30.2% and 46.4%

homology, depending on the species (Supplemental Table S2). The low sequence identity is

consistent with an ancient duplication event that gave rise to *npm2a* and *npm2*b genes.

However, neither the topology of the Npm2 phylogenetic tree nor the comparison between

Npm2a and Npm2b could indicate the kind of duplication event that had occurred. In

contrast, the high sequence identity shared by Npm2b1 and Npm2b2 in salmonids (between

75.1% and 86%) suggested a more recent duplication event. This observation in all of the

investigated salmonid species is consistent with the hypothesis that *npm2b1* and *npm2b2*

genes likely resulted from the SaGD.

<u>Synteny analysis</u>

In order to further understand the origin of the *npm2a* and *npm2b* genes in vertebrates,

we performed a synteny analysis of their neighboring genes in representative vertebrate

genomes. We focused our study on two mammals (human and mouse), two sauropsids

(Chinese alligator and chicken), two amphibians (*Xenopus tropicalis* and *laevi*s L), one basal

sarcopterygian (coelacanth), one basal actinopterygian (spotted gar), and two teleosts

(zebrafish and tetraodon) (Fig. 2). Analysis was also performed on the *Xenopus laevi*s S

subgenome, but since the results were very similar to that of *Xenopus laevi*s L subgenome,

we only showed the latter's data [24].

The human, mouse, Chinese alligator, *Xenopus tropicalis* and *laevis*, coelacanth,

spotted gar, and zebrafish *npm2b* genes are located in genomic regions containing common

loci, including *dok2, xpo7, fgf17, dmtn, lgi3, bmp1, sorbs3, pdlim2, gpr124, prlhr, got1l1,*

adrb3, and *nkx6-3*. Together with the phylogenetic analysis, this indicates that *npm2b* genes

investigated here are orthologous. Synteny analysis revealed the absence of the *npm2b* gene

in tetraodon although the above-mentioned neighbouring genes are present in its genome

(Fig. 2 and Supplemental Table S3).

The Chinese alligator, chicken, *Xenopus tropicalis* and *laevis*, spotted gar, and

zebrafish *npm2a* genes are located in genomic regions containing the same loci as *npm2b*

conserved regions (Fig. 2). Indeed, Chinese alligator and spotted gar *npm2a* and *npm2b* genes

are located in the vicinity of each other on scaffold 98.1 and the linkage group LG1,

respectively. The presence of *npm2a* and *npm2b* genes in the same genomic region in

representative species of sarcopterygians (Chinese alligator and *Xenopus tropicalis* and

laevis) and actinopterygians (spotted gar) strongly suggested that *npm2a* and *npm2b* genes

could have resulted from a unique local duplication of an ancestral *npm2* gene.

Evolutionary history of npm2 genes in vertebrates

The presence/absence of *npm2a* and *npm2b* in the current vertebrate phyla and species

is summarized in Fig.3, and we also propose an evolutionary scenario for the diversification

of the *npm2* genes across vertebrate evolution.

In this study, we demonstrated that *npm2a* and *npm2b* may be paralogous genes

present in the different vertebrate groups, chondrichthyans (Fig.1), sarcopterygians, and

actinopterygians (Figs. 1 and 2), which strongly suggested that the *npm2* genes originated

from a duplication event prior to the divergence of chondrichthyans and osteichthyans. Since

the four *npm* family members (*npm1*, *npm2*, *npm3*, and *npm4*) are thought to be produced

from the first two rounds of WGD (VGD1 & VGD2) that occurred early on in vertebrate

evolution, we can thus hypothesize that the duplication event that generated *npm2a* and

npm2b took place after VGD2, but before emergence of chondrichthyans and osteichthyans,

between 450 and 500 million years ago (Mya) [25]. The findings from our synteny analysis

demonstrated that in representative species of actinopterygians (spotted gar) and

sarcopterygians (*Xenopus tropicalis* and *laevis* and Chinese alligator), *npm2a* and *npm2b*

genes are at two distinct loci located on the same chromosomic region (Fig. 2). These results

strongly suggested that *npm2a* and *npm2b* genes arose from local gene duplication rather than

a whole genome (or chromosome) duplication event (Fig. 3).

The teleost ancestor experienced an extra WGD event (TGD) [21], but in all

investigated teleosts, we observed a maximum of one *npm2a* ortholog and one *npm2b*

ortholog (except in salmonids) (Fig. 1). In addition, *npm2a* and *npm2b* are located on two

TGD ohnologous regions on chromosomes 8 and 10, respectively (Fig. 2), in zebrafish,

which is consistent with the loss of one of the *npm2a* duplicates from one region and the loss

of one *npm2b* duplicate from the corresponding ohnologous region. Thus, *npm2* diversity is

most likely due to the early loss of one of the two *npm2a* and *npm2b* TGD ohnologs (Fig. 3).

This observation in zebrafish strengthened the hypothesis that TGD did not impact the *npm2a*

and *npm2b* diversity in teleosts. In addition, the lack of *npm2a* and *npm2b* in neoteleostei

species, such as tetraodon, suggested that additional gene losses occurred early in the

evolutionary history of this group (Fig. 2).

In salmonids, we identified two *npm2b* paralogs, i.e. *npm2b1* and *npm2b2*, in all

investigated salmonid species (Fig. 1). Considering the high sequence identity shared by

npm2b1 and *npm2b2*, it is strongly hypothesized that these duplicates originated from SaGD.

In contrast, we identified only one *npm2a* gene in all investigated salmonids suggesting that

SaGD did not have any impact on the current salmonid *npm2a* diversity mostly due to an

early loss of the SaGD ohnolog of this gene (Fig. 3).

In addition to the early gene losses after WGD, various other independent and

phylum-specific gene losses may have contributed to shape the current diversity of *npm2a*

and *npm2b* in vertebrates. In fact, although both *npm2a* and *npm2b* have been globally

conserved in sarcopterygians and actinopterygians, some phyla in each group lack at least one

of the genes. In sarcopterygians, *npm2a* is conserved in amphibians such as *Xenopus*

tropicalis and *laevis*, and in sauropsids such as Chinese alligator and chicken (Figs. 1-3). In

contrast, we did not find any *npm2a* genes in the Coelacanth genome (Figs. 1-3), which could

be due to the lower assembly quality of the concerned genomic region (see Supplemental

Table S3), and in the mammalian genomes including human and mouse (Figs. 1-3), whose

absence may have been due to the loss of this gene in the common ancestor of mammals. On

the other hand, *npm2b* is conserved in most investigated species. In actinopterygians, *npm2a*

and *npm2b* are both conserved in all investigated species except neoteleostei in which neither

npm2a nor *npm2b* is found. In contrast, one *npm2a* and two *npm2b*, i.e. *npm2b1* and *npm2b2*,

are found in salmonids, representing to date the most important diversity of *npm2* genes in

vertebrates (Figs. 1-3). Considering the essentialness of the *npm2* gene in embryonic

development, its lack in some neoteleosteon species, such as Atlantic cod and medaka, raises

the question on how evolution can cope with its loss. Further analyses of data from the

Phylofish database [26] (data not shown) showed that *npm3* had the strongest homology to

npm2 and was predominantly expressed in the ovaries in medaka and cod which suggest that

it could potentially compensate for *npm2* deficiency. In contrast, *npm3* does not show an

ovarian-predominant expression in spotted gar and zebrafish, thus suggesting that its strong

ovarian expression could be restricted to neoteleostei.

Evolution of npm2a and npm2b coding sequences

To examine the evolutionary adaptation of the *npm2* proteins, estimation of the ratio

of substitution rates (dN/dS) between the paralogous *npm2a* and *npm2b* CDS for all of the

species harbouring multiple *npm2* paralogs was performed (Supplemental Table S4). For all

investigated species, the dN/dS values were well below 1, indicating that whilst *npm2a* and

npm2b genes diverged approximately 500 Mya, they have remained under strong purifying

selection and evolutionary pressure, which may have tended to conserve their distinct protein

structures and functions. Thus, in this study, we found that Npm2a and Npm2b may have

diverged from an ancient duplication and currently share low sequence identity, suggesting

that they could have thus evolved with different roles, which were likely conserved through

strong negative selection.

Expression domains of Npm2a and Npm2b in vertebrates

In order to investigate further the potential functions of Npm2a and Npm2b, we first

explored the tissue distributions of both transcripts using two different approaches, qPCR and

RNA-Seq, the latter of which was obtained from the Phylofish online database [26]. In

zebrafish and *Xenopus tropicalis*, we observed that *npm2a* and *npm2b* were both

predominantly expressed in the ovary, and to a lesser extent, the muscle, as well as in the

zebrafish gills (Fig. 4A and 4B). We also demonstrated that *npm2a* and *npm2b* were

predominantly expressed in the ovary of bowfin, elephantnose fish, panga, European eel,

sweetfish, and northern pike (Fig. 4C to 4H). In the investigated salmonid species, brook

trout and brown trout, *npm2a*, *npm2b1*, and *npm2b2* were also predominantly expressed in

the ovary (Fig. 4I and 4J). In addition, *npm2* transcripts were expressed at a very low level in

the testis of European eel, northern pike, and salmonid species (Fig. 4F, 4H, 4I, and 4J). In

teleosts, *npm2a* mRNA levels globally tended to be lower than *npm2b* (or *npm2b1* and

npm2b2 for salmonids) (Fig. 4A and 4D-J). The clear ovarian-specific expression profiles of

npm2a and *npm2b* transcripts were conserved in all investigated species, from teleosts to

tetrapods. This is consistent with the *npm2b* mRNA profiles reported in the literature for

mouse [14], cattle [27], *Xenopus tropicalis* [28], and zebrafish [8]. Despite the long passage

of time since their divergence (approximately 500 Mya) [25] as demonstrated above, *npm2a*

and *npm2b* appeared to have conserved their ovarian-specific expression profiles, which

suggested that they both have roles in female reproduction and/or embryonic development.

Embryonic expression of npm2a and npm2b

Thus, in order to delve deeper into their functions in reproduction and embryogenesis,

we investigated *npm2a* and *npm2b* mRNA expression during oogenesis and early

development in zebrafish. During zebrafish oogenesis, both *npm2a* and *npm2b* transcripts

were found at high levels in oocytes (Fig. 5A), and despite gradual decreases in the levels of

both *npm2* during oogenesis, they still can be detected at reasonable amounts in the

unfertilized egg (i.e. metaphase 2 oocyte) (Fig. 5B). Thereafter, *npm2a* and *npm2b* transcript

levels progressively decreased during embryonic development after fertilization before

reaching very low levels at 24 hours post-fertilization (hpf) (Fig. 5B). This suggests that both

mRNAs are strictly maternal (*i.e.* not re-expressed by the zygote), which is consistent with

previous studies on zebrafish *npm2a* and *npm2b* transcripts [8,29]. Their expression profiles

are typical features of maternally-inherited mRNAs, which highly suggested that the novel

npm2a is also a maternal-effect gene.

Peptidic domains of npm2 paralogs

We further examined the functions of the Npm2 proteins by analysis of their protein

domains. To date, only the role of *npm2b* has been investigated in *Xenopus tropicalis*

[12,30,31], mouse [14,32], humans [33–35], cattle [27], and zebrafish [8]. As previously

demonstrated, *npm2b* is a maternal-effect gene whose transcripts are accumulated in the

growing oocyte and maternally-inherited by the zygote, where it functions as a histone

chaperone to decondense sperm DNA as well as reorganize chromatin, thus, it has also been

suggested to contribute to ZGA as well [8,14,27,28]. Npm2b is thought to be activated by

various post-translational modifications and homo-pentamerisation [36], and subsequently

interacts with chromatin by exchanging sperm-specific basic proteins with histones via its

core domain and acidic tracts A1, A2 and A3 [34,37–42]. However, no data is available on

the structure and function of Npm2a. Using a predictive approach, we identified the presence

of the Npm core domain, acidic tract A1, and acidic tract A2 in the various investigated

Npm2 sequences. The nucleoplasmin family is defined by the presence of an Npm core

domain, which enables oligomerization of Npm proteins, in all of its members [38,43]. We

were able to predict the presence of this domain in all investigated Npm2a and Npm2b

sequences (Supplemental Fig. S1), suggesting that Npm2a proteins could also form homo- or

hetero-polymers. The acidic tract domains were demonstrated to facilitate histone binding by

increasing the recognition and affinity for different histones [40,44]. Acidic tract A1 was

demonstrated to be absent from most of the Npm2b investigated so far, except *Xenopus*

tropicalis Npm2b, and acidic tract A2 was predicted to be in all investigated Npm2b apart

from guinea pig Npm2b and American whitefish Npm2b1 (Supplemental Fig. S1). This

strongly suggested that the histone and basic protein binding activity of Npm2b is mediated

predominantly by acidic tract A2. On the other hand, in all investigated Npm2a proteins, we

predicted an acidic tract A1 except in that of dogfish, zebrafish, and allis shad (Supplemental

Fig. S1). In contrast, only half of the investigated Npm2a proteins harbored an acidic tract

A2, which is additionally shorter than the one present in Npm2b proteins (Supplemental Fig.

S1). Our results demonstrated that all Npm2a proteins, except those of Allis shad and

dogfish, possess acidic tracts, which could potentially mediate histone and basic protein

interactions.

Functional analysis of npm2a and npm2b in zebrafish

Lastly, we performed functional analysis of these two Npm2 proteins by genetic

knockout using the CRISPR/cas9 system. One-cell staged embryos were injected with the

CRISPR/cas9 guides that targeted either *npm2a* or *npm2b* and allowed to grow to adulthood.

Mosaic founder mutant females (F0) were identified by fin clip genotyping and subsequently

mated with wild-type (WT) males, and embryonic development of the F1 fertilized eggs was

recorded. Since the mutagenesis efficiency of the CRISPR/cas9 system was very high, as

previously described [45,46], the *npm2* genes were sufficiently knocked-out even in the

transgenic mosaic F0 females. This was evidenced by the substantially lower transcript levels

of *npm2a* and *npm2b* in the F1 embryos as compared to those from control WT pairings

(Fig.6A). Thus, the phenotypes of *npm2a* (n=3) and *npm2b* (n=4) mutants could be observed

even in the F0 generation. Since none of the mutated genes were transmissible to future

generations neither through the male nor the female, therefore, all of our observations were

obtained from the F0 generation. We observed that most of the embryos from the *npm2b*

mutant females underwent cellular division during the very early stages of development (1-3

hpf) despite a considerable number of embryos with abnormal morphology (24.00±7.84%

versus 0% in controls) (Fig.6B), which included smaller size, enlarged yolk to membrane

ratio, and small yolk to membrane ratio. The diameter of the embryo is demonstrated by the

red dotted lines at oblong and germ ring stages, which reveal the extremely reduced size of

npm2a embryos (Fig.6C). Notably, around two thirds of the embryos showed abnormal cell

division, even in those with normal morphology, that culminated in developmental arrest at

around 4 hpf following which the cells stopped dividing and appeared to regress. Thus, the

developmental success at 4 hpf (oblong/sphere stage) was 38.09±7.31% vs. 85.09±4.88% in

controls (Fig.6B and 6C [right column]). By the 24 somite stage (24 hpf), these embryos were

all dead while the remaining embryos that showed normal development and normal cell

division continued to progress normally. This finding corresponded to our previous results,

which showed that *npm2b*-deficient embryos targeted by morpholino arrested at 4 hpf and

eventually died [8].

On the other hand, *npm2a* mutant-derived embryos had a very low developmental

success even at a very early stage of growth (1 hpf) (17.62±5.06% vs. 85.09±4.88% in

controls) as defined by a complete lack of cell division, and a large population of them

(36.36±6.98% vs. 0% in controls) had an abnormal morphology (Fig.6B and 6C [middle

columns]). The abnormal morphology of these embryos was similar to that found in *npm2b*-

deficient embryos. The F1 embryos that did not undergo any cell division at 1 hpf continued

to display a complete lack of development at 64-cell, oblong, dome, germ ring, and finally

somite stages (Fig.6B [middle columns]). Similar to the *npm2b* mutant-derived embryos, the

npm2a-deficient embryos were all dead by 24 hpf while the remaining embryos that showed

normal development and normal cell division continued to progress normally. This novel

finding showed for the first time that *npm2a* is essential for early development of embryos,

and is therefore a crucial maternal-effect gene. Further, we demonstrated that while *npm2a*

and *npm2b* share similar tissue distribution, as both are found specifically in the ovaries and

early stage embryos, their roles are distinct and essential to embryogenesis, and one could not

compensate for the other. Notably, *npm2a* probably plays a major role at or immediately after

fertilization since the embryos are arrested before the first mitotic division, which is in line

with a role in sperm DNA decondensation since lack of proper processing of the paternal-

derived DNA would lead to developmental arrest at the earliest stage and embryonic death. In

contrast, *npm2b* appears to function at a later stage or other factors can replace its role in the

early stages as the *npm2b*-deficient embryos are capable of dividing until around 4 hpf, which

corresponds to previous reports [8,14]. These findings are in line with those demonstrated for

other maternal-effect genes in mammals, such as *mater*, *floped*, *filia*, *tle6* [47], and *bcas2*

[48] as well as in zebrafish, including *fue* [49] and *cellular atoll* [50], all of which function to

moderate the early events of fertilization and embryogenesis and their ablation leads to arrest

and mortality at the earliest stages of embryogenesis.

In consequence, the dominancy of the mutant allele combined with the strong

maternal effect of these genes, which led to massive early embryonic mortality such that none

of the mutant carriers survived, very likely contributed to the lack of germline transmission of

the mutations despite reported heritability rates of 10-30% in CRISPR/cas9-targeted zebrafish

from previous studies [45,46]. This was supported by detection of the mutant allele via PCR

genotyping in the genome of F1 *npm2a*- and *npm2b*-deficient embryos that were

developmentally-arrested at the germ ring stage, but not in the surviving animals at 24 hpf

(Supplemental Fig.S2).

Conclusions

Since two annotated genes corresponding to *npm2*, which is known as an essential

maternal-effect gene in mammals and amphibians, were recently found in zebrafish, we set

out to investigate their evolution and function. In this study, we demonstrated that the two

duplicates of *npm2*, *npm2a* and *npm2b*, exist in a wide range of vertebrate species, including

ray-finned fish, amphibians, and bird. We also found that the mammalian *npm2* gene is in

fact an ortholog of *npm2b*. Using phylogeny and synteny analyses, we traced the origins of

those two duplicates to the early stages of vertebrate evolution. Our findings indicate that

npm2a and *npm2b* genes resulted from a local gene duplication that may have occurred

between VGD2 and the divergence of ray-finned fish and tetrapods (~450-500 Mya). This

ancient origin is in line by the low sequence identity between the two *npm2* genes, although

the main protein domains remain conserved. Both genes exhibit a strict ovarian expression

and corresponding transcripts are maternally-inherited in fish and amphibians. Moreover, we

demonstrated and confirmed by CRISPR/cas9-directed genetic knockout of *npm2a* and

npm2b that they are both maternal-effect genes that play essential, but distinct, roles in early

embryogenesis. *npm2a* probably plays a major role at or immediately after fertilization, and

most likely in chromatin sperm decondensation. In contrast, *npm2b* appears to function at a

later stage and could participate in ZGA. Our findings will help us gain further insight into

the evolutionary diversity of maternal-effect genes and understand the underlying

mechanisms that contribute to reproductive success in vertebrates.

Material and Methods

Genomic databases

The following genomic data were extracted and investigated from the ENSEMBL genomic

database (http://www.ensembl.org/index.html): human, *Homo sapiens*; mouse, *Mus*

musculus; chicken, *Gallus gallus*; Xenopus, *Xenopus tropicalis*; coelacanth, *Latimeria*

chalumnae; spotted gar, *Lepisosteus oculatus*; zebrafish, *Danio rerio*; and tetraodon,

Tetraodon nigroviridis. The Chinese alligator (*Alligator sinensis*) genome was extracted and

investigated from the NCBI genomic database (http://www.ncbi.nlm.nih.gov/genome/22419).

The *Xenopus laevis* L and S genomes were analyzed from the Xenbase database

(www.xenbase.org).

Transcriptomic databases

The following actinopterygian transcriptomes were retrieved and investigated from the

Phylofish database (http://phylofish.sigenae.org/index.html): bowfin, *Amia calva*; spotted gar,

Lepisosteus oculatus; elephantnose fish, *Gnathonemus petersi*; arowana, *Osteoglossum*

bicirrhosum; butterflyfish, *Pantodon buchholzi*; European eel, *Anguilla anguilla*; rainbow

trout, *Oncorhynchus mykiss*; allis shad, *Alosa alosa*; zebrafish, *Danio rerio*; panga,

Pangasius hypophthalmus; northern pike, *Esox lucius*; grayling, *Thymallus thymallus*;

Atlantic cod, *Gadhus morua*; medaka, *Oryzias latipes*; European perch, *Perca fluviatilis*;

brown trout, *Salmo trutta*; European whitefish, *Coregonus lavaretus*; brook trout, *Salvelinus*

fontinalis; Astyanax, *Astyanax mexicanus*; lake whitefish, *Coregonus clupeaformis*; eastern

mudminnow, *Umbra pygmae*, and sweetfish, *Plecoglossus altivelis*.

Gene predictions

TBLASTN search

Genomic data were analyzed using the TBLASTN algorithm (search sensitivity: near exact

match short) on the ENSEMBL website or the NCBI browser for the Chinese alligator

genome. The TBLASTN algorithm on the SIGENAE platform was used on the

transcriptomic data.

<u>Predictions of *npm2* genes</u>

The peptidic sequences of zebrafish Npm2a and Npm2b were used as query in TBLASTN

search to identify the open reading frame (ORF) encoding *npm2* genes in the various

investigated genomes and transcriptomes.

Phylogenetic analysis

Amino acid sequences of 75 Npm2, 3 Npm3, and 3 Npm1 proteins were first aligned using

ClustalW. The JTT (Jones, Taylor, and Thornton) protein substitution matrix of the resulting

alignments was determined using ProTest software. Phylogenetic analysis of Npm proteins

was performed using the Maximum Likelihood method (MEGA 5.1 software) with 1,000

bootstrap replicates.

Synteny analyses

Synteny maps of the conserved genomic regions in human, mouse, chicken, Xenopus,

coelacanth, spotted gar, zebrafish, and tetraodon were produced using PhyloView on the

Genomicus v75.01 website ([http://www.genomicus.biologie.ens.fr/genomicus-75.01/cgi-](http://www.genomicus.biologie.ens.fr/genomicus-75.01/cgi-bin/search.pl)

[bin/search.pl](http://www.genomicus.biologie.ens.fr/genomicus-75.01/cgi-bin/search.pl)). Synteny analysis of the Chinese alligator conserved genomic regions was

performed using TBLASTN searches in the corresponding genomic database. For each gene,

the peptidic sequences of human and chicken were used as query, as far as they were

referenced in the databases.

RNA-seq

RNA-seq data were deposited into Sequence Read Archive (SRA) of NCBI under accession

references SRP044781-84, SRP045138, SRP045098-103, and SRP045140-146. The

construction of sequencing libraries, data capture and processing, sequence assembly,

mapping, and interpretation of read counts were all performed as previously reported with

some modifications [26]. In order to study the transcript expression patterns and levels of

npm2 for each actinopterygian species presenting 2 or 3 *npm2* genes, we mapped the double

stranded RNA-seq reads to the corresponding *npm2* CDS using BWA-Bowtie with stringent

mapping parameters (maximum number of allowed mismatches –aln 2). Mapped reads were

counted using the idxstat command in SAMtools, with a minimum alignment quality value (–

q 30) to discard ambiguous mapping reads. For each species, the number of mapped reads

was then normalized for each *npm2* gene across the 11 tissues using RPKM normalization.

Quantitative real-time PCR (QPCR)

For each sample, total RNA was extracted using Tri-Reagent (Molecular Research Center,

Cincinnati, OH) according to the manufacturer's instructions. Reverse transcription (RT) was

performed using 1 μg of RNA from each sample with M-MLV reverse transcriptase and

random hexamers (Promega, Madison, WI). Briefly, RNA and dNTP were denatured for

6 min at 70°C and then chilled on ice for 5 min before addition of the RT reagents. RT was

performed at 37°C for 1 h and 15 min followed by a 15-min incubation step at 70°C. Control

reactions were run without reverse transcriptase and used as negative control in the QPCR

study. For each studied tissue, cDNA originating from three individual fish were pooled and

subsequently used for QPCR. Follicular oocytes at different stages of oogenesis were

obtained from at least 3 different wildtype animals, and pooled embryos were obtained from

at least 3 clutches from each individual mutant. QPCR experiments were performed with the

Fast-SYBR GREEN fluorophore kit (Applied Biosystems, Foster City, CA) as per the

manufacturer's instructions using 200 nM of each primer in order to keep PCR efficiency

between 90% and 100%, and an Applied Biosystems StepOne Plus instrument. RT products,

including control reactions, were diluted 1/25, and 4 μl of each sample were used for each

PCR. All QPCR were performed in triplicate. The relative abundance of target cDNA was

calculated from a standard curve of serially diluted pooled cDNA and normalized to 18S, β-

actin, β-2 microglobulin, and EF1α transcripts. The primer sequences can be found in

Supplemental Table S5.

Comparison of npm2a and npm2b peptidic sequences

For the species harbouring Npm2a and Npm2b, or Npm2a, Npm2b1, and Npm2b2, we

compared the corresponding protein sequences using EMBOSS Matcher tool on the EBI

website (http://www.ebi.ac.uk/Tools/psa/emboss_matcher/). Results of the pair-wise

comparisons are presented as identity percentages in Supplemental Table S2. We also

estimated the synonymous substitution rates (dS) and non-synonymous substitution rates

(dN) between the paralogous Npm2a and Npm2b coding sequences (CDS) using JCoDA

v1.14 software (http://www.tcnj.edu/~nayaklab/jcoda). The alignment options were set to

"clustal", and the Yang and Nielsen dN/dS substitution models were used.

CrispR-cas9 genetic knockout

CRISPR/cas9 guide RNA (gRNA) were designed using the ZiFiT online software and were

made against 3 targets within each gene to generate large genomic deletions, ranging from

250-1600 base pairs, that span exons which allow the formation of non-functional proteins.

Nucleotide sequences containing the gRNA were ordered, annealed together, and cloned into

the DR274 plasmid. *In vitro* transcription of the gRNA from the T7 initiation site was

performed using the Maxiscript T7 kit (Applied Biosystems), and their purity and integrity

were assessed using the Agilent RNA 6000 Nano Assay kit and 2100 Bioanalyzer (Agilent

Technologies, Santa Clara, CA). Zebrafish embryos at the one-cell stage were micro-injected

with approximately 30-40 pg of each CRISPR/cas9 guide along with 8-9 nM of purified cas9

protein (a generous gift from Dr. Anne de Cian from the National Museum of Natural History

in Paris, France). The embryos were allowed to grow to adulthood, and genotyped using fin

clip and PCR that detected the deleted regions. The PCR bands of the mutants were then sent

for sequencing to verify the deletion. Once confirmed, the mutant females were mated with

wildtype males to produce F1 embryos, whose phenotypes were subsequently recorded.

Images were captured with a Carl Zeiss microscope (Jena, Germany) and ToupCam camera

(ToupTek, Hangzhou, China).

Genotyping by PCR

Fin clips were harvested from animals under anesthesia (0.1% phenoxyethanol) and lysed

with 5% chelex containing 100 μg of proteinase K at 55^{o}C for 2 hrs and then 99^{o}C for 10

minutes. The extracted DNA was subjected to PCR using Advantage2 system (Clontech,

Mountain View, CA) for *npm2b* and Jumpstart Taq polymerase (Sigma-Aldrich, St. Louis,

MO) for *npm2a*. The primers are listed in Supplemental Table S5.

Statistical Analysis

Comparison of two groups was performed using the GraphPad Prism statistical software (La

Jolla, CA), and either the Student's t-test or Mann-Whitney U-test was conducted depending

on the normality of the groups based on the Anderson-Darling test.

Acknowledgements

This work was supported by ANR grants PHYLOFISH (ANR-10-GENM-017) and

Maternal Legacy (ANR-13-BSV7-0015) to JB. Authors would like to thank Dr. Yann Audic

for providing Xenopus tropicalis animals and Ms. Amélie Patinote for zebrafish rearing and

egg production.

References

1.   Baroux C, Autran D, Gillmor CS, Grimanelli D, Grossniklaus U. The maternal to

zygotic transition in animals and plants. Cold Spring Harb Symp Quant Biol. 2008;73:

89–100.

2.   Lindeman RE, Pelegri F. Vertebrate maternal-effect genes: Insights into fertilization,

early cleavage divisions, and germ cell determinant localization from studies in the

zebrafish. Mol Reprod Dev. 2010;77: 299–313.

3.   Lee MT, Bonneau AR, Takacs CM, Bazzini AA, DiVito KR, Fleming ES, et al.

Nanog, Pou5f1 and SoxB1 activate zygotic gene expression during the maternal-to-

zygotic transition. Nature. 2013;503: 360–364.

4.   Giraldez AJ, Mishima Y, Rihel J, Grocock RJ, Van Dongen S, Inoue K, et al.

Zebrafish MiR-430 promotes deadenylation and clearance of maternal mRNAs.

Science. 2006;312: 75–79.

5.   Giraldez AJ. microRNAs, the cell's Nepenthe: clearing the past during the maternal-

to-zygotic transition and cellular reprogramming. Curr Opin Genet Dev. 2010; 20:369–

375.

6.   Lund E, Liu M, Hartley RS, Sheets MD, Dahlberg JE. Deadenylation of maternal

mRNAs mediated by miR-427 in Xenopus laevis embryos. Rna. 2009;15: 2351–2363.

7.   Bouchareb A, Le Cam A, Montfort J, Gay S, Nguyen T, Bobe J, et al. Genome-wide

identification of novel ovarian-predominant miRNAs: new insights from the medaka

(Oryzias latipes). Sci Rep. 2017;7: 40241.

8.   Bouleau A, Desvignes T, Traverso JM, Nguyen T, Chesnel F, Fauvel C, et al.

Maternally inherited npm2 mRNA is crucial for egg developmental competence in

zebrafish. Biol Reprod. 2014;91: 43.

9.   Laskey RA, Honda BM, Mills AD, Finch JT. Nucleosomes are assembled by an acidic

protein which binds histones and transfers them to DNA. Nature. 1978;275: 416–420.

10.   Laskey RA, Earnshaw WC. Nucleosome assembly. Nature. 1980;286: 763–767.

11.   Mills AD, Laskey RA, Black P, De Robertis EM. An acidic protein which assembles

nucleosomes in vitro is the most abundant protein in Xenopus oocyte nuclei. J Mol

Biol. 1980;139: 561–568.

12. Philpott A, Leno GH, Laskey RA. Sperm decondensation in Xenopus egg cytoplasm is

mediated by nucleoplasmin. Cell. 1991;65: 569–578.

13. Philpott A, Leno GH. Nucleoplasmin remodels sperm chromatin in Xenopus egg

extracts. Cell. 1992;69: 759–767.

14. Burns KH, Viveiros MM, Ren Y, Wang P, DeMayo FJ, Frail DE, et al. Roles of

NPM2 in chromatin and nucleolar organization in oocytes and embryos. Science.

2003;300: 633–636.

15. Dehal P, Boore JL. Two rounds of whole genome duplication in the ancestral

vertebrate. PLoS Biol. 2005;3: e314.

16. Van de Peer Y, Maere S, Meyer A. The evolutionary significance of ancient genome

duplications. Nat Rev Genet. 2009;10: 725–732.

17. Eirin-Lopez JM, Frehlick LJ, Ausio J. Long-term evolution and functional

diversification in the members of the nucleophosmin/nucleoplasmin family of nuclear

chaperones. Genetics. 2006;173: 1835–1850.

18. Frehlick LJ, Eirin-Lopez JM, Jeffery ED, Hunt DF, Ausio J. The characterization of

amphibian nucleoplasmins yields new insight into their role in sperm chromatin

remodeling. BMC Genomics. 2006;7: 99.

19. Wotton KR, Weierud FK, Dietrich S, Lewis KE. Comparative genomics of Lbx loci

reveals conservation of identical Lbx ohnologs in bony vertebrates. BMC Evol Biol.

2008;8: 171.

20. Jovelin R, Yan YL, He X, Catchen J, Amores A, Canestro C, et al. Evolution of

developmental regulation in the vertebrate FgfD subfamily. J Exp Zool B Mol Dev

Evol. 2010;314: 33–56.

21.   Glasauer SM, Neuhauss SC. Whole-genome duplication in teleost fishes and its

evolutionary consequences. Mol Genet Genomics. 2014;289: 1045–1060.

22.   Volff JN. Genome evolution and biodiversity in teleost fish. Heredity (Edinb).

2005;94: 280–294.

23.   Berthelot C, Brunet F, Chalopin D, Juanchich A, Bernard M, Noel B, et al. The

rainbow trout genome provides novel insights into evolution after whole-genome

duplication in vertebrates. Nat Commun. 2014;5: 3657.

24.   Session AM, Uno Y, Kwon T, Chapman JA, Toyoda A, Takahashi S, et al. Genome

evolution in the allotetraploid frog Xenopus laevis. Nature. 2016;538: 336–343.

25.   Near TJ, Eytan RI, Dornburg A, Kuhn KL, Moore JA, Davis MP, et al. Resolution of

ray-finned fish phylogeny and timing of diversification. Proc Natl Acad Sci U S A.

2012;109: 13698–13703.

26.   Pasquier J, Cabau C, Nguyen T, Jouanno E, Severac D, Braasch I, et al. Gene

evolution and gene expression after whole genome duplication in fish: the PhyloFish

database. BMC Genomics. 2016;17: 368.

27.   Lingenfelter BM, Tripurani SK, Tejomurtula J, Smith GW, Yao J. Molecular cloning

and expression of bovine nucleoplasmin 2 (NPM2): a maternal effect gene regulated

by miR-181a. Reprod Biol Endocrinol. 2011;9: 40.

28.   Burglin TR, Mattaj IW, Newmeyer DD, Zeller R, De Robertis EM. Cloning of

nucleoplasmin from Xenopus laevis oocytes and analysis of its developmental

expression. Genes Dev. 1987;1: 97–107.

29.   Harvey SA, Sealy I, Kettleborough R, Fenyes F, White R, Stemple D, et al.

Identification of the zebrafish maternal and paternal transcriptomes. Development.

2013;140: 2703–2710.

30.   Laskey RA, Mills AD, Philpott A, Leno GH, Dilworth SM, Dingwall C. The role of

nucleoplasmin in chromatin assembly and disassembly. Philos Trans R Soc Lond B

Biol Sci. 1993;339: 263–269.

31.   Prado A, Ramos I, Frehlick LJ, Muga A, Ausio J. Nucleoplasmin: a nuclear chaperone.

Biochem Cell Biol. 2004;82: 437–445.

32.   De La Fuente R, Viveiros MM, Burns KH, Adashi EY, Matzuk MM, Eppig JJ. Major

chromatin remodeling in the germinal vesicle (GV) of mammalian oocytes is

dispensable for global transcriptional silencing but required for centromeric

heterochromatin function. Dev Biol. 2004;275: 447–458.

33.   Lee HH, Kim HS, Kang JY, Lee BI, Ha JY, Yoon HJ, et al. Crystal structure of human

nucleophosmin-core reveals plasticity of the pentamer-pentamer interface. Proteins.

2007;69: 672–678.

34.   Platonova O, Akey I V, Head JF, Akey CW. Crystal structure and function of human

nucleoplasmin (npm2): a histone chaperone in oocytes and embryos. Biochemistry.

2011;50: 8078–8089.

35.   Okuwaki M, Sumi A, Hisaoka M, Saotome-Nakamura A, Akashi S, Nishimura Y, et

al. Function of homo- and hetero-oligomers of human nucleoplasmin/nucleophosmin

family proteins NPM1, NPM2 and NPM3 during sperm chromatin remodeling.

Nucleic Acids Res. 2012;40: 4861–4878.

36.   Onikubo T, Nicklay JJ, Xing L, Warren C, Anson B, Wang WL, et al.

Developmentally Regulated Post-translational Modification of Nucleoplasmin

Controls Histone Sequestration and Deposition. Cell Rep. 2015. pii:S2211-

1247(15)00196-5. doi: 10.1016/j.celrep.2015.02.038.

37.   Dingwall C, Dilworth SM, Black SJ, Kearsey SE, Cox LS, Laskey RA. Nucleoplasmin

cDNA sequence reveals polyglutamic acid tracts and a cluster of sequences

homologous to putative nuclear localization signals. Embo J. 1987;6: 69–74.

38.   Dutta S, Akey I V, Dingwall C, Hartman KL, Laue T, Nolte RT, et al. The crystal

634   structure of nucleoplasmin-core: implications for histone binding and nucleosome

635   assembly. Mol Cell. 2001;8: 841–853.

636   39.   Banuelos S, Hierro A, Arizmendi JM, Montoya G, Prado A, Muga A. Activation

637   mechanism of the nuclear chaperone nucleoplasmin: role of the core domain. J Mol

638   Biol. 2003;334: 585–593.

639   40.   Salvany L, Chiva M, Arnan C, Ausio J, Subirana JA, Saperas N. Mutation of the small

640   acidic tract A1 drastically reduces nucleoplasmin activity. FEBS Lett. 2004;576: 353–

641   357.

642   41.   Taneva SG, Banuelos S, Falces J, Arregi I, Muga A, Konarev P V, et al. A mechanism

643   for histone chaperoning activity of nucleoplasmin: thermodynamic and structural

644   models. J Mol Biol. 2009;393: 448–463.

645   42.   Fernandez-Rivero N, Franco A, Velazquez-Campoy A, Alonso E, Muga A, Prado A. A

646   Quantitative Characterization of Nucleoplasmin/Histone Complexes Reveals

647   Chaperone Versatility. Sci Rep. 2016;6: 32114.

648   43.   Namboodiri VM, Akey I V, Schmidt-Zachmann MS, Head JF, Akey CW. The

649   structure and function of Xenopus NO38-core, a histone chaperone in the nucleolus.

650   Structure. 2004;12: 2149–2160.

651   44.   Ramos I, Fernandez-Rivero N, Arranz R, Aloria K, Finn R, Arizmendi JM, et al. The

652   intrinsically disordered distal face of nucleoplasmin recognizes distinct

653   oligomerization states of histones. Nucleic Acids Res. 2014;42: 1311–1325.

654   45.   Auer TO, Duroure K, De Cian A, Concordet JP, Del Bene F. Highly efficient

655   CRISPR/Cas9-mediated knock-in in zebrafish by homology-independent DNA repair.

656   Genome Res. 2014;24: 142–153.

46.   Gagnon JA, Valen E, Thyme SB, Huang P, Akhmetova L, Pauli A, et al. Efficient

mutagenesis by Cas9 protein-mediated oligonucleotide insertion and large-scale

assessment of single-guide RNAs. PLoS One. 2014;9: e98186.

47.   Li L, Baibakov B, Dean J. A subcortical maternal complex essential for

preimplantation mouse embryogenesis. Dev Cell. 2008;15: 416–425.

48.   Xu Q, Wang F, Xiang Y, Zhang X, Zhao ZA, Gao Z, et al. Maternal BCAS2 protects

genomic integrity in mouse early embryonic development. Development. 2015;142:

3943–3953.

49.   Dekens MP, Pelegri FJ, Maischein HM, Nusslein-Volhard C. The maternal-effect gene

futile cycle is essential for pronuclear congression and mitotic spindle assembly in the

zebrafish zygote. Development. 2003;130: 3907–3916.

50.   Yabe T, Ge X, Pelegri F. The zebrafish maternal-effect gene cellular atoll encodes the

centriolar component sas-6 and defects in its paternal function promote whole genome

duplication. Dev Biol. 2007;312: 44–60.

51.   Kimmel CB, Ballard WW, Kimmel SR, Ullmann B, Schilling TF. Stages of embryonic

development of the zebrafish. Dev Dyn. 1995;203: 253–310.

Figure legends

Figure 1: Consensus phylogenetic tree of Npm2 proteins.

This phylogenetic tree was constructed based on the amino acid sequences of Npm2 proteins

(for the references of each sequence see Supplemental Table S1) using the Neighbour Joining

method with 1,000 bootstrap replicates. The number shown at each branch node indicates the

bootstrap value (%). The tree was rooted using Npm3 and Npm1 sequences. The Npm2a

sequences are in blue, the Npm2b sequences are in red, the salmonid Npm2b1 sequences are

in purple, and the salmonid Npm2b2 sequences are in pink.

Figure 2: Conserved genomic synteny of *npm2* genes.

Genomic synteny maps comparing the orthologs of *npm2a, npm2b*, and their neighbouring

genes. *npm2* genes are named as *npm2a* and *npm2b* (formerly known as *npm2*). The other

genes were named after their human orthologs according to the Human Genome Naming

Consortium (HGNC). Orthologs of each gene are shown in the same color. The direction of

arrows indicates the gene orientation, with the ID of the genomic segment indicated above

and the position of the gene (in 10^{-6} base pairs) indicated below. The full gene names and

detailed genomic locations are given in Supplemental Table S3.

Figure 3: Current status and proposed evolutionary history of *npm2* genes among

gnathostomes.

The names of the current representative species of each phylum are given at the end of the

final branches, together with a red and/or blue X to denote the *npm2* genes they possess

(*npm2a*=red, *npm2b*=blue). The black X upon an *npm2* gene symbol indicates a gene loss.

LGD: local gene duplication; TGD: teleost-specific whole genome duplication; SaGD:

salmonid-specific whole genome duplication.

Figure 4: Tissue distribution of *npm2a* and *npm2b* in different species.

A) Tissue expression analysis by QPCR of *npm2a* and *npm2b* mRNAs in zebrafish and

Xenopus. **(B)** Expression level is expressed as a percentage of the expression in the ovary for

the most expressed gene. Data were normalized using *18S* expression. **(C-J)** Tissue

expression level by RNA-Seq of *npm2a* and *npm2b* mRNAs in different fish species. mRNA

levels are expressed in read per kilobase per million reads (RPKM). In salmonids (**I** and **J**),

the two 4R isoforms of *npm2b* are *npm2b1* and *npm2b2*. Br, brain ; Gi, gills ; Lu, lung ; In,

intestine ; Li, liver ; Mu, muscle ; He, heart ; Bo, bone ; Ki, kidney ; Ov, ovary ; and Te,

testis.

Figure 5: *npm2a* and *npm2b* expression during oogenesis and early development.

QPCR analysis of **(A)** *npm2a* and **(B)** *npm2b* expression during oogenesis and early

embryonic development in zebrafish. Data were normalized using luciferase, and relative

expression was based on *npm2b* expression at the indicated stage. UF, unfertilized egg; hpf,

hours post-fertilization.

Figure 6: CRISPR/cas9 knockout of *npm2a* and *npm2b* in zebrafish.

(A) Relative expression level of *npm2a* and *npm2b* transcripts by QPCR in the fertilized

zebrafish eggs from crosses between *npm2a* or *npm2b* mutant female and wildtype (WT)

male, respectively. **(B)** Developmental success as measured by the proportion of fertilized

eggs that underwent normal cell division and reached normal developmental milestones

based on Kimmel et al. [51] from crosses between WT animals (control), and *npm2a* or

npm2b mutant female and WT male at 1 and 4 hours post-fertilization (hpf). Shaded grey

columns denote normal morphology and black columns denote abnormal morphology of the

fertilized eggs at 1 hpf. **(C)** Representative images demonstrating development of fertilized

eggs from crosses between control animals (left panels), and *npm2a* (middle panels) or

npm2b (right panels) mutant female and WT male at 64-cell, oblong, dome, germ ring, and

24-somite stages according to Kimmel et al [51]. N=3 for *npm2a* mutant and N=4 for *npm2b*

mutant. All assessments were performed from at least 3 clutches from each mutant. Scale

bars denote 500 nm. Red dotted lines define the diameter of the embryo.   ** $p<0.01$,

***$p<0.001$, ****$p<<0.0001$

Supplemental Figure S1: Npm2a and Npm2b protein domain conservation.

Presence of the Npm core (red color bar) and acidic tract (blue color bar) domains in *npm2a*

and *npm2b* types in various vertebrate species.

Supplemental Figure S2: Genotyping by PCR for *npm2a* and *npm2b* in F1 embryos.

npm2a and *npm2b* knockout animals were crossed with wildtype (WT) zebrafish, and

individual F1 embryos were harvested at 6 hours post-fertilization and subjected to

genotyping by PCR with gene-specific primers. The wildtype PCR product for *npm2a* was

1634 base pairs (bp) and the CRISPR/cas9-targeted knockout mutant band was 240 bp while

the WT *npm2b* band was 813 bp and the CRISPR/cas9-targeted knockout mutant band was

560 bp. The yellow asterisks denote the embryos carrying the mutant allele.

Supplementary Data

Supplemental Table S1: list of Npm2 protein sequences used in the phylogenetic analysis.

Supplemental Table S2: Npm2a and Npm2b protein sequence comparisons.

Supplemental Table S3: list of the genes from the conserved syntenic region of *npm2* genes.

Supplemental Table S4: Estimation of the ratio of substitution rates (dN/dS) in *npm2a* and

npm2b genes.

Supplemental Table S5: QPCR and PCR primers.

Supplemental Fig.S1: Npm2a and Npm2b protein domain conservation.

Supplemental Fig.S2: Genotyping of F1 embryos from *npm2a* and *npm2b* mutants.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Supplemental Figure S1

Supplemental Figures

S2