Group-level progressive alterations in brain connectivity patterns revealed by diffusion-tensor brain networks across severity stages in Alzheimer's disease

J. Rasero[1,2*], C. Alonso-Montes[3*], I. Diez[2*], L. Olabarrieta-Landa[4], L. Remaki[3], I. Escudero[2,5], B.
Mateos[2,5], P. Bonifazi[2,6], M. Fernandez[2,7], J.C. Arango-Lasprilla[2,6], S. Stramaglia[1,3], J.M.**
Cortes[2,6,8], for the Alzheimer's Disease Neuroimaging Initiative[8].**

[1]Dipartamento di Fisica, Universita degli Studi di Bari and INFN. Bari, Italy
[2]Biocruces Health Research Institute. Barakaldo, Spain
[3]BCAM - Basque Center for Applied Mathematics. Bilbao, Spain
[4]Department of Education and Psychology, University of Deusto. Bilbao, Spain
[5]Radiology Service. Cruces University Hospital. Barakaldo, Spain
[6]IKERBASQUE: The Basque Foundation for Science. Bilbao, Spain
[7]Neurology Service. Cruces University Hospital. Barakaldo, Spain
[8]Department of Cell Biology and Histology. University of the Basque Country. Leioa, Spain
13

[8]Data used in the preparation of this article were obtained from the Alzheimer's Disease
Neuroimaging Initiative (ADNI) database (adni.loni.ucla.edu). As such, the investigators within the
ADNI contributed to the design and implementation of ADNI and/or provided data but did not
participate in the analysis or writing of this report. A complete listing of ADNI investigators is
available at
http://adni.loni.usc.edu/wpcontent/uploads/how_to_apply/ADNI_Acknowledgement_List.pdf

* Equal first-author contribution

** Equal last-author contribution

Keywords: Diffusion-Tensor Imaging; Brain networks; Alzheimer's disease, Severity progression

Abstract. Alzheimer's disease (AD) is a chronically progressive neurodegenerative disease highly correlated to aging. Whether AD originates by targeting a localized brain area and propagates to the rest of the brain across disease-severity progression is a question with an unknown answer. Here, we aim to provide an answer to this question at the group-level by looking at differences in diffusion-tensor brain networks. In particular, making use of data from Alzheimer's Disease Neuroimaging Initiative (ADNI), four different groups were defined (all of them matched by age, sex and education level): G_1 ($N_1=36$, healthy control subjects, Control), G_2 ($N_2=36$, early mild cognitive impairment, EMCI), G_3 ($N_3=36$, late mild cognitive impairment, LMCI) and G_4 ($N_4=36$, AD). Diffusion-tensor brain networks were compared across three disease stages: stage I (Control vs EMCI), stage II (Control vs LMCI) and stage III (Control vs AD). The group comparison was performed using the multivariate distance matrix regression analysis, a technique that was born in genomics and was recently proposed to handle brain functional networks, but here applied to diffusion-tensor data. The results were three-fold: First, no significant differences were found in stage I. Second, significant differences were found in stage II in the connectivity pattern of a subnetwork strongly associated to memory function (including part of the hippocampus, amygdala, entorhinal cortex, fusiform gyrus, inferior and middle temporal gyrus, parahippocampal gyrus and temporal pole). Third, a widespread disconnection across the entire AD brain was found in stage III, affecting more strongly the same memory subnetwork appearing in stage II, plus the other new subnetworks, including the default mode network, medial visual network, frontoparietal regions and striatum. Our results are consistent with a scenario where progressive alterations of connectivity arise as the disease severity increases and provide the brain areas possibly involved in such a degenerative process. Further studies applying the same strategy to longitudinal data are needed to fully confirm this scenario.

1.Introduction

Alzheimer's disease (AD), the most common form of dementia, is a chronically progressive

neurodegenerative disease highly correlated to aging; indeed, although the prevalence of clinically

manifested AD is about 2% at the age of 65 years, it increases to 30% at the age of 85 years (Wimo et

al. 1997).

AD is characterized by an accumulation of beta-amyloid plaques and neurofibrillary tangles

composed of tau amyloid fibrils (Hardy 2006) associated with synapse loss and neurodegeneration

leading to long-term memory impairment and other cognitive problems. To date, there is no treatment

known to slow down the progression of this disorder.

The initial AD pathology develops many years before the cognitive and functional impairments are

evident. Different terms have been used to describe this disease-starting condition, including pre-

dementia and prodromal AD and, more often, MCI (mild cognitive impairment). The concept of MCI

varied over the past two decades and has been classified into different broad categories depending on

memory performance and the number of impaired cognitive functions (Mueller *et al.* 2005).

An accurate prediction for the conversion from MCI to AD can help to clinicians to evaluate AD risk

pre-symptomatically, initiate treatments at early stage, and monitor their effectiveness (Cheng *et al.*

2015, Li *et al.* 2014). However, such a prediction is challenging, as the MCI group is highly

heterogeneous and only a few patients convert to AD, a rate of about 8% to 15% convert per year

69(Ritter *et al.* 2015, Mitchell and Shiri-Feshki 2009). However, the amnestic subtype of MCI is more 70prevalent than the non-amnestic MCI (Petersen *et al.* 2010), and has an annual conversion rate higher 71of about 30% to 40% (Schmidtke and Hermeneit 2008, Rozzini *et al.* 2007, Geslani *et al.* 2005).

73This study aims to search for neuroimaging biomarkers that can account for differences with respect 74to a healthy control population from the early to the final stages of AD. Multitude of different 75neuroimaging studies has addressed the conversion from MCI to AD, see (Zhang *et al.* 2014) and 76references therein. In relation to structural magnetic resonance imaging (MRI), it was shown that the 77hippocampus volume and the volume from other subcortical structures at MCI were well correlated 78to a worse progression to AD, with accuracy of about 65% in the prediction from MCI to AD (Teipel 79*et al.* 2015).

81Rather than assuming that specific brain regions are affected in AD, a blind approach using multiple 82regions of interest has been shown to achieve a better predictive accuracy (of about 80%) of the 83conversion from MCI to AD (Westman *et al.* 2011, Eskildsen *et al.* 2013, Liu *et al.* 2013). The use of 84tensor diffusion MRI in combination with structural MRI has provided better results as compared to 85only structural MRI, showing that white-matter integrity of the fornix, cingulum, and 86parahippocampal gyrus provided accuracy varying from 80% to even 95% (Wee *et al.* 2013, Mielke 87*et al.* 2012, Douaud *et al.* 2013).

89Initiatives like the Alzheimer's Disease Neuroimaging Initiative (ADNI) provide important resources 90to study AD to the research community (including demographic data, imaging datasets, cognitive

91 tests, etc.), pushing forward multimodal studies correlating different imaging modalities to 92 neuropsychological functioning. Interestingly, ADNI also allows the possibility of studying variations 93 in the images at a group level across disease's progression, as brain images are categorized in 94 different groups ranging from Control to AD, with two intermediate stages, early and late mild 95 cognitive impairment, EMCI and LMCI, respectively. Importantly, although EMCI and LMCI 96 patients have memory impairment (Medina *et al.* 2006), the conversion rate to AD is only between 8- 97 15% per year (Mitchell and Shiri-Feshki 2009), making this group have a special relevance in the 98 development of novel imaging techniques that could correlate with disease progression.

100 Despite extensive research shedding light into the MCI to AD conversion, the precise mechanisms 101 and clinical variables responsible for such progression are poorly understood, mainly due to the lack 102 of time-resolved longitudinal studies in large populations. Taking into consideration previous work 103 (Khedher *et al.* 2015, Douaud *et al.* 2011, Bosch *et al.* 2012, Liu *et al.* 2013, Acosta-Cabronero *et al.* 104 2012, Preti *et al.* 2012), the present study focus on the variations of brain networks across AD 105 progression at a group level. It is hypothesized that if in the transition from Control to MCI the 106 connectivity pattern of some subnetworks are altered, in further disease stages the alterations of the 107 same subnetworks will coexist together with alterations of new different subnetworks in the AD 108 brain, in a manner that connectivity alterations will finally extend to the rest of the brain.

110 2. Material and Methods

111 2.1 Ethics

112 The present study made use of ADNI data previously collected in 50 different institutions.

113 Participants provided informed consent before recruitment and data collection started. In addition,

114 participants filled questionnaires approved by each participating site's Institutional Review Board

115 (IRB). The complete list of ADNI sites' IRBs can be found using the following link:

116 http://adni.loni.ucla.edu/about/data-statistics/.

118 2.2 Alzheimer's Disease Neuroimaging Initiative (ADNI)

Diffusion tensor imaging (DTI) data was used in this paper from ADNI database

http://adni.loni.usc.edu. ADNI was launched in 2003 by the Nat. Inst. on Aging (NIA), the Nat. Inst.

Biomedical Imaging and Bioengineering (NIBIB), the Food and Drug Administration (FDA), private

pharmaceutical companies and non-profit organizations, as a $60 million, 5-year public-private

partnership. ADNI's main goal has been to test whether serial MRI, positron emission tomography

(PET), other biological markers, and clinical and neuropsychological assessment can be combined to

measure the progression of MCI and early AD. Determination of sensitive and specific markers of

very early AD progression is intended to aid researchers and clinicians to develop new treatments and

monitor their effectiveness, as well as to lessen the time and cost of clinical trials. The Principal

Investigator of this initiative is Michael W. Weiner, MD, VA Medical Center and Univ. California –

San Francisco. ADNI subjects have been recruited from over 50 sites across the U.S. and Canada.

Currently, around 1500 adults were recruited in the different ADNI initiatives, ages 55 to 90,

consisting of cognitively normal older (NC), early/late MCI (EMCI/LMCI), significant memory

concern (SMC) and early AD (AD) individuals. The follow up duration of each group is specified in

the protocols for ADNI-1, ADNI-2 and ADNI-GO, see further information in www.adni-info.org .

135 2.3 Demographic Data

A total number of N=144 subjects were used in this study (Table S1). This number was chosen in

order to get the biggest four groups as possible (Control, EMCI, LMCI and AD), balanced by size,

age and sex. DTI images were selected and downloaded from ADNI database, belonging to four

different groups: Control (N_1=36), EMCI (N_2=36), LMCI (N_3=36) and AD (N_4=36). Age and sex

were balanced across groups (Table 1), respectively, using a t-test and chi-squared test. In addition, it

is important to remark that the "years of education" variable was already controlled by the ADNI

group classification, for details see Inclusion criteria in page 31 of https://adni.loni.usc.edu/wp-

content/uploads/2008/07/**adni2-procedures-manual**.pdf

145 2.4 ADNI group classification

The group labels Control, EMCI, LMCI and AD are based on several test scores, such as the Logical

Memory II subscale (LMIIS) from the Wechsler Memory Scale, the Mini-Mental State Examination

(MMSE) and the Clinical Dementia Rating (CDR), as well as National Institute of Neurological and

Communicative Disorders and Stroke and the AD and Related Disorders Association

(NINCDS/ADRDA) criteria in AD cases. In the procedures manual each of the criteria are cited

(http://adni.loni.usc.edu/wp-content/uploads/2008/07/adni2-procedures-manual.pdf).

153Control subjects are free of memory complaints (beyond normal ageing), verified by a study partner.

154EMCI, LMCI and AD must have a subjective memory concern as reported by the subject, study

155partner, or clinician. Details of specific groups are given in Table 2.

1572.5 Group-level stages for AD progression

158AD progression was defined by three different stages: stage I (control vs EMCI), stage II (control vs

159LMCI) and stage III (control vs AD). Further details are given in Figure 1.

1612.6 DTI acquisitions

162All subjects in this study had the same ADNI imaging protocol, explained in

163http://adni.loni.usc.edu/methods/documents/mri-protocols/ and consisting in whole-brain MRI 3T

164scanners and Diffusion Weighted Images (DWI) images of the axial DTI series. The DTI images

165were acquired using spin echo pulse sequence echo-planar-imaging (SE-EPI) with the following

166parameters: TR = 9050.0 ms; TE set to minimum (values ranging from 60 ms till 69 ms); 59 slices

167with thickness of 2.7 mm with no gap among slices; 128x128 matrix with a FOV of 35.0 cm; with

168matrix pixels 256x256x2714 and voxel size 1.36x1.36x2.7 mm^3, flip angle = 90º. A diffusion gradient

169was applied along 41 non-collinear directions with a b value of 1000 s/mm2. Additionally, one set of

170images was acquired with no diffusion weighting (b= 0 s/mm2).

1722.7 Diffusion tensor brain networks

173Diffusion tensor brain networks were built following a similar methodology to previous work

174(Marinazzo *et al.* 2014, Diez *et al.* 2015, Alonso-Montes *et al.* 2015, Amor *et al.* 2015, Diez *et al.*

1752017) using FSL (FMRIB Software Library v5.0) and the Diffusion Toolkit. First, all the selected

176images were downloaded in DICOM and transformed to Nifti format for further analysis. Next, an

177eddy current correction was applied to overcome the artifacts produced by variation in the gradient

178field directions, together with the artifacts produced by head movements. Next, using the corrected

179data, a local fitting of the diffusion tensor was applied to compute the diffusion tensor model for each

180voxel. Next, a Fiber Assignment by Continuous Tracking (FACT) algorithm was applied (Mori *et al.*

1811999). Next, a transformation from the Montreal Neurological Institute (MNI) space to the

182individual-subject diffusion space was computed and applied to the brain hierarchical atlas (BHA)

183with M=20 modules, which was shown in (Diez *et al.* 2015) to have the best correspondence

184between functional and structural modules. This atlas developed by the authors is available to

185download at http://www.nitrc.org/projects/biocr_hcatlas/. This allowed building 20 x 20 structural

186connectivity (SC) matrices, each per subject, by counting the number of white matter streamlines

187connecting all module pairs. Thus, the element matrix *(i,j)* of SC is given by the streamlines number

188between modules *i* and *j*. As a result, SC is a symmetric matrix, where the connectivity from *i* to *j* is

189equal to the one from *j* to *i*.

1912.8 Labelling of anatomical regions

192The anatomical representation of the initial 2,514 brain regions existing in BHA was identified by

193using the Automated Anatomical Labelling (AAL) brain atlas (Tzourio-Mazoyer *et al.* 2002).

194Therefore, the anatomical identification of the brain regions used in this work followed the labels

195existing in the AAL atlas.

198 **2.9 Cross-group analysis: Multivariate Distance Matrix Regression**

199 The cross-group analysis has been performed using the Multivariate Distance Matrix Regression
200 (MDMR) approach proposed in (Shehzad *et al.* 2014). Connectome-wide association studies are
201 usually performed by means of mass-univariate statistical analyses, in which the association between
202 a phenotypic variable (e.g., the score in a neuropsychological test) with each entry of the brain
203 connectivity matrix is tested across subjects. Such analysis, however, exhibits two main pitfalls: First
204 even at the level of region of interest (ROI) and thus choosing much less regions as voxels, the
205 number of statistical tests entailed is large (Milham 2012), which increases the potential for false
206 positives. On the other hand, studying each brain connectivity matrix entry separately, concurrent
207 contributions from other entries are necessarily ignored (Cole *et al.* 2010). In multivariate methods,
208 instead, the simultaneous contribution of entire sets of brain connectivity entries to a phenotypic
209 variable is evaluated, in a manner that it better captures the concurrent global changes and reduces
210 the number of false positives.

212 A multivariate distance regression was applied and the variation of distance in connectivity patterns
213 between groups as a response of the Alzheimer's progression as compared to the Control state was
214 tested. For a fixed brain module i, the distance between connectivity patterns of module i to the rest
215 of the brain was calculated per pair of subjects (u,v) --by calculating Pearson correlation between
216 connectivity vectors of subject pairs--, thus leading to a distance matrix in the subject space for each
217 module i investigated. In particular, the following formula was calculated

$\quad d_{uv}^{i} = \sqrt{2 \cdot \left(1 - r_{uv}\right)}$ \hfill (Eq. 1)

where $\quad r_{uv}\quad$ is the Pearson correlation between connectivity patterns of *i* for subjects *u* and *v*. After

repeating the same procedure for all subjects, as many distance matrices as partition modules

$\left(i = 1, \ldots, 20\right)\quad$ were obtained. Next, MDMR was applied to perform cross-group analysis as

implemented in R (McArtor 2016).

It is important to emphasize that MDMR does not look to how individual modules are locally

organized or connected, but to the integration connectivity pattern between those segregated modules

to the rest of the brain. Therefore, when group differences were found on a MDMR given module,

the connectivity alterations from that module suggests an significant affect to the rest of the brain.

MDMR yielded a pseudo-F estimator (analogous to that F-estimator in standard ANOVA analysis),

which addresses significance of disease strength due to between-group variation as compared to

within-group variations (McArdle and Anderson 2001). To compare between groups when the

regressor variable is categorical (*i.e.* the group label), given a distance matrix, one can calculate the

total sum of squares as

$\quad SS_{T} = \dfrac{1}{N} \sum\limits_{u=1} \sum\limits_{v=1+1} d_{uv}^{2}\quad$, \hfill (Eq. 2)

with $\quad N\quad$ being the total number of subjects. Notice that, from here on, we will consider

$\quad d_{uv} \equiv d_{uv}^{i}$. Thus, we got a different $\quad SS_{T}\quad$ for each module *i*. Similarly, the within-group sum of

squares can be written as

$$SS_W = \sum \frac{1}{n_g} \sum_{u=1} \sum_{v=u+1} d_{uv}^2 \varepsilon_{uv}^g \quad ,$$ (Eq. 3)

where n_g is the number of subjects per group and ε_{uv}^g a variable equal to 1 if subjects u and v

belong to group g and 0 otherwise. The between-group variation is simply $SS_B = SS_T - SS_W$,

which leads to a pseudo-F statistic that reads

$$F = \frac{SS_B / (m-1)}{SS_W / (N-m)}$$ (Eq. 4)

where m is the number of groups. As it was acknowledged in (Zapala and Schork 2006), the pseudo-F

statistic is not distributed like the usual Fisher's F-distribution under the null hypothesis. Accordingly,

we randomly shuffled the subject indices and computed the pseudo-F statistic for each time. A p-

value is computed by counting those pseudo F-statistic values from permuted data greater than that

from the original data respect to the total number of performed permutations.

Finally, we controlled for type I errors due to the 20 independent statistical performed tests by false

discovery rate corrections (Benjamini and Hochberg 1995). Corrected whole-brain connectivity

patterns of modules are the ones related to AD progression at the different stages. A schematic

overview of the method can be found in Figure 2.

3. Results

Results are summarized in Table 3 and modules involved in the disease progression at the group level

are shown in Figure 3. See also Table S2 for examples of the different terms participating in the

statistical test.

259 3.1 Stage I: Control vs EMCI

260 A total number of 36 images per each group were selected to perform group comparison. No
261 significant differences were found in terms of module connectivity patterns to the whole brain.

263 3.2 Stage II: Control vs LMCI

264 A total number of 36 images per each group were selected to perform group comparison. Significant
265 differences were found for the connectivity between the <u>module 18</u> and the rest of the brain
266 (p=0.007). As detailed in (27), the module 18 of the brain hierarchical atlas incorporated part of the
267 hippocampus, amygdala, entorhinal cortex, fusiform gyrus, inferior temporal gyrus, middle temporal
268 gyrus, parahippocampal gyrus and temporal pole.

270 3.3 Stage III: Control vs AD

A total of 36 images per group were selected to perform group comparison. At this stage, significant
different connectivity patterns were found in multiple modules existing in BHA:

<u>Module 1</u> (p=0.023); including part of the posterior cingulate.

<u>Module 2</u> (p=0.049); including part of the putamen, anterior cingulate, rostral pars of the middle
frontal gyrus, superior parietal gyrus, supramarginal gyrus, insula, inferior parietal gyrus, precentral
gyrus and superior frontal gyrus.

277Module 3 (p=0.049); part of the paracentral lobe, precentral gyrus, postcentral gyrus, precuneus,

278superior frontal gyrus, superior parietal gyrus, superior temporal gyrus, supramarginal gyrus and

279insula.

280Module 4 (p=0.031); part of the cuneus, lateral occipital sulcus, lingual gyrus, pericalcarine cortex

281and precuneus.

282Module 8 (p=0.031); part of the caudate nucleus and putamen.

283Module 12 (p=0.031); part of the inferior parietal gyrus, inferior temporal gyrus, lateral frontal

284orbital gyrus, pars orbitalis, pars triangularis, rostral pars of the middle frontal gyrus, superior frontal

285gyrus, caudate nucleus and anterior cingulate.

286Module 14 (p=0.006); part of the thalamus, hippocampus, amygdala, putamen, ventral

287diencephalon, banks of the superior temporal sulcus, parahippocampal gyrus, superior temporal

288gyrus, insula, middle temporal gyrus and temporal pole.

289Module 15 (p=0.031); part of the thalamus, putamen, pallidum, brainstem, hippocampus, amygdala,

290accumbens nucleus, ventral diencephalon, orbital gyrus and insula.

291Module 16 (p=0.031); part of the cerebellum, banks of the superior temporal sulcus, inferior parietal

292gyrus, cingulate isthmus, middle temporal gyrus, precuneus and superior temporal gyrus.

293Module 18 (p=0.002); see previous 3.2 section for the anatomical description, but notice a reduction

294in p value from 0.007 (Control vs LMCI) to 0.002 (Control vs AD).

2963.4 Common affected modules between stages

297Connectivity pattern of <u>module 18</u> to the rest of the brain was found at stage II (p=0.007) and at stage

298III (p=0.002), indicating that the further the disease progresses, the greater the connectivity of

299module 18 is altered to the rest of the brain.

301**4. Discussion**

302The aim of the present study was to identify differences in brain connectivity patterns between a

303control group and three pathological groups by disease severity. For this purpose, diffusion tensor

304brain networks were built allowing determining connectivity differences at three consecutive severity

305stages: stage I (Control vs EMCI), stage II (Control vs LMCI) and stage III (Control vs AD).

307The results showed an absence of significant changes in connectivity patterns in stage I, that is,

308between patients with early mild cognitive impairment and healthy individuals. The MDMR analysis

309we have applied finds group differences in the connectivity patterns from different modules to the

310rest of the brain. Therefore, when observing early mild cognitive impairment, our analysis allows for

311some possible structural damages to locally occur. This study has shown that even if local alterations

312exist, they are not capable of producing global inter-module network reorganization/redistribution

313detectable by the MDMR analysis.

315Significant differences were found by the MDMR method in stage II in relation to a network

316involved with memory (module 18), which includes the hippocampus, amygdala, entorhinal cortex,

317fusiform gyrus, inferior temporal gyrus, mean temporal gyrus, parahippocampal gyrus and the

318temporal pole. Strikingly, the change in module 18 connectivity becomes more evident in stage III

319(i.e., in patients with AD), and *memory* alterations coexist with alterations in a multitude of different
320modules (modules 1-4, 8, 12, 14-16 and 18), which encompass the default mode network, the
321sensory-motor network, the medial visual network, frontoparietal regions and subcortical networks
322(including part of the hippocampus, amygdala and putamen).

324The brain connectivity alterations found in this study in stage II might be related to the appearance of
325several cognitive manifestations, which are typical of AD. For example, many studies have
326determined the main cognitive impairment in the preclinical phase of AD is episodic memory
327(Almkvist, 1996, Arnaiz et al., 2003; Albert et al., 2001; Bäckman et al., 2004, 2005; Grober et al.,
3282008), in which hippocampus; entorhinal cortex and amygdala are involved. Following this line of
329results, research has found that alterations in the temporal-medial lobe have an affect before AD is
330even clinically diagnosed (Almkvist, 1996; Bäckman et al., 2004, 2005; Small et al., 1999; Estévez-
331González et al., 2003; Small et al., 2003). Moreover, research has also shown that the initial neuronal
332lesions in AD begin in the entorhinal region (included in module 18, therefore, in agreement with our
333results) with the accumulation of neurofibrillary tangles and neuritic plaques (Gómez-Isla et al.,
3341996).

336Although alterations of the episodic memory are considered the most critical ones at the preclinical
337phase of AD (Small, et al., 2003; Storandt, 2008) and tasks that measure episodic memory have been
338shown to be particularly effective at identifying people at risk for developing AD (Elias et al., 2000;
339Tierney et al., 1996), studies have shown that people with mild cognitive impairment who have
340altered (in addition to episodic memory) other cognitive areas such as verbal ability (Apostolova et
341al., 2008; Arnaiz et al et al., 2003; Bäckman et al., 2004, 2005; Joubert et al., 2010), executive

342functions (Albert et al., 2001; Bäckman et al., 2004, 2005; Dickerson et al., 2007; Grober et al., 2008;

343Storandt, 2008; Blacker et al., 2007; Rapp et al., 2005), perceptual speed (Bäckman et al., 2005),

344Visuo-spatial / visuoperceptive skills (Almkvist, 1996, Arnaiz et al., 2003; Bäckman et al., 2004,

3452005; Alegret et al., 2009), attention (Bäckman et al., 2005; Rapp et al., 2005), etc. are more likely to

346convert to AD than those with only memory impairment (Bozoki et al., 2001). As indicated by

347Bäckman et al. (2004, 2005), a number of different areas in addition to the ones in the temporal-

348medial lobe are altered prior to the diagnosis of AD (such as the anterior cingulate, temporal sulcus,

349posterior cingulate, temporoparietal regions, frontal regions and precuneus). This may explain why

350studies attempting to find cognitive markers of the AD preclinical stage find alterations in other

351cognitive functions apart from episodic memory.

353As the disease progresses, not only the disconnection pattern of module 18 becomes more evident

354(increasing the distance between AD and controls, Table 3), but such significant changes extend to

355other brain regions. For example, areas of the hippocampus affected by module 14 are well known to

356suffer a very severe cognitive degeneration, a fact also confirmed by functional connectivity studies

357(Zhou *et al.* 2008). The results also indicate a significant connectivity change with temporal medial

358areas, as revealed by module 16, as shown in Tract Based Spatial Statistics at (Stricker *et al.* 2009,

359Acosta-Cabronero *et al.* 2010, Salat *et al.* 2010). Similarly to the results of this study, authors of (He

360*et al.* 2007) demonstrated, through a combined structural and functional analysis, changes in

361connectivity between the lingual and cuneus, by using only structural connectivity data.

363The results of the present study indicate a significant change in the connectivity from the entire brain

364to the areas provided by module 4, mainly associated to visual function. A decrease in virtual

365capacity in AD is well known, especially in those areas involving movement blindness, depth

366perception, color perception and contrast sensitivity (Whittaker *et al.* 2002). Again, this damage

367expansion to other brain regions also agrees with the extent and worsening of cognitive aspects (e.g.,

368memory, attention, language; Weintraub *et al.* 2012) and neurobehavioral problems (e.g. personality

369changes, anxiety, depression, agitation, hallucinations; Chung & Cummings, 2000, Bassiony et al.,

3702000, Senanarong et al., 2003) of patients with AD.

372Previous studies have analyzed the connectivity differences from tensor diffusion networks in AD

373and have found significant alterations in the inferior longitudinal fasciculus for patients at risk of AD

374(Smith *et al.* 2010), which could correspond to LMCI. Similarly, a voxel-based analysis in (Honea *et*

375*al.* 2009) showed a significant decrease in FA for fibers connecting the parahippocampal gyrus. In

376addition, patients diagnosed in the early stages of AD (corresponding to early or late mild cognitive

377impairment in this study) had a significant reduction in white matter in the upper longitudinal

378fasciculus, which also connects part of module 18 in the brain hierarchical atlas with the frontal lobe

379(Rose *et al.* 2000). The authors (Hanyu *et al.* 1998) found significant changes in apparent diffusion

380coefficients and diffusion anisotropy in patients with recent progressive cognitive impairment,

381suggesting an early decrease in temporal fiber density, a region included in the module 18, therefore

382in concordance to our results.

384A different comparison between pathological groups

385By defining disease progression across three stages, I (control vs EMCI), II (control vs LMCI) and III

386(control vs AD), we have found progressive variations in connectivity patterns that start in a module

387clearly associated to memory function (including part of the hippocampus, amygdala, entorhinal

388cortex, fusiform gyrus, inferior and middle temporal gyrus, parahippocampal gyrus and temporal

389pole) and later on, alterations are found widespread along the entire brain. Therefore, it is important

390to emphasize that we have defined disease progression by comparing each pathological group with

391respect to the control group. A different possibility for assessing connectivity variations is to perform

392comparisons between pathological groups, i.e., EMCI vs LMCI, LMCI vs AD, EMCI vs AD. For the

393two comparisons EMCI vs LMCI and LMCI vs AD, none of the module showed differences in

394connectivity patterns   (Table S3). However, the EMCI vs AD comparison showed differences in

395modules 2,3,4,14 and 16.

397The reason why our strategy of defining disease progression with respect to the control group found

398differences in module 18 at the beginning of the progression is due to the fact that the within-group

399distance contribution of the control group is smaller than the corresponding one in any of the

400pathological groups. In particular, we calculated the sum of distances squared (defined in Eq. 1)

401between pairs of subjects of connectivity between module 18 and the rest of the brain and obtained

402values of 62 (control), 76 (EMCI), 83 (LMCI) and 82 (AD). In other words, the tensor-diffusion

403connectivity values of module 18 are more homogeneous between subjects within the control group

404as compared to subjects within any other pathological group, what makes our strategy to successfully

405detect differences in the connectivity pattern of module 18 at the early stages of disease progression.

407<u>Implications</u>

408In recent years a great deal of emphasis has been placed on early AD detection (Albert et al., 2001);

409from looking for pharmacological or non-pharmacological treatments to help delay the age of onset

410disease and to slow down the clinical disease progression. Similar to other studies, these results

411provide (by looking to diffusion tensor brain networks) that the earliest detection in connectivity

412patterns affecting globally the rest of the brain starts in a network mainly encompassing memory

413function.

415On the other hand, identifying brain connectivity patterns in patients who have not yet developed AD

416might shed some light in determining how these connectivity patterns evolve as time goes on. In

417addition, it will be possible to associate connectivity patterns with clinical patient's variations existing

418at each disease stage. This might help better understand the relationship between deterioration in

419brain functioning and clinical patient's characteristics.

421Limitations

422The results of the present study should be interpreted in light of the following limitations. First, it is a

423cross-sectional study with different groups of patients in each experimental group and with a small

424sample size, so future studies should try to extend to bigger cohorts and follow the same group of

425people over time as the disease progresses. Second, the patients included in the study have a probable

426AD, which means that the definitive diagnosis of AD can only be performed post-mortem (Fearing et

427al., 2007). The use of patients with familiar AD could help to know in depth the evolution of the

428disease and the changes in cerebral connectivity from many years back to its onset. Third, there are a

429number of risk factors associated with the decline of mild cognitive impairment which can affect

brain connectivity such as advanced diabetes, symptomatology depressive disorder, hypertension,

hypotension, obesity, history of traumatic brain injury and APOE genotype, that have not been taken

into account in this study. Future studies should take into account the possible influence of these

variables on the processes of cerebral connectivity.

<u>Summary</u>

In conclusion, the results obtained from this study applying a multivariate method to diffusion tensor

connectivity networks across AD severity progression, are in line with the evolution of AD from both

the neuropathological and neuropsychological points of view. That is, first alterations occur in the

connectivity of regions of the middle temporal lobe (hippocampus and entorhinal), which coincides

with the first symptoms of altered episodic memory in the preclinical stage and in mild cognitive

impairment. As the disease progresses, the brain damage and its disconnection of these regions

become more evident and expands to other areas, which coincides with the expansion and/or

worsening of other cognitive functions and neurobehavioral aspects seen in the individuals with AD.

Future developments will deal with the application of the same methodology to longitudinal data, a

mandatory step to confirm our results.

Author Contributions

JR, CAM and ID had equal first-author contribution; JR, CAM and ID analyzed the data and made

the figures; LOL and JCAL connected results to cognitive deficits in AD; LR, IE, BM, PB, MF,

JCAL, SS and JMC designed the research; all the authors wrote the manuscript and agreed in its

submission; SS and JMC had equal last author contributions

Acknowledgements

SS acknowledges financial support from Bizkaia Talent and European Commission through

COFUND with the research project BRAhMS – Brain Aura Mathematical Simulation– (AYD-000-

285). JR acknowledges financial support from the Minister of

Education, Language Policy and Culture (Basque Government) under

Doctoral Research Staff Improvement Programme. JMC and JCAL acknowledge financial support

from Ikerbasque (The Basque Foundation for Science) and from Ministerio Economia, Industria y

Competitividad (Spain) and FEDER (grant DPI2016-79874-R). PB acknowledges financial support

from Ikerbasque and from the Ministerio Economia, Industria y Competitividad (Spain) and FEDER

(grant SAF2015-69484-R).

Data collection and sharing for this project was funded by the Alzheimer's Disease Neuroimaging Ini-

tiative (ADNI) National Institutes of Health grant U01 AG024904. ADNI is funded by the National

Institute on Aging, the National Institute of Biomedical Imaging and Bioengineering, and through

generous contributions from the following: Abbott, AstraZeneca, AB, Amorfix, Bayer Schering

Pharma AG, Bio- clinica Inc., Biogen Idec, Bristol-Myers Squibb, Eisai Global Clinical Develop-

ment, Elan Corporation, Genentech, GE Healthcare, Innogenetics, IXICO, Janssen Alzheimer Im-

munotherapy, Johnson and Johnson, Eli Lilly and Co., Medpace, Inc., Merck and Co., Inc., Meso

Scale Diagnostic, & LLC, Novartis AG, Pfizer Inc, F. Hoffman-La Roche, Servier, Synarc, Inc., and

Takeda Pharmaceuticals, as well as non-profit partners the Alzheimer's Association and Alzheimer's

Drug Discovery Foundation, with participation from the U.S. Food and Drug Administration. Private

sector contributions to ADNI are facilitated by the Foundation for the National Institutes of Health

(www.fnih.org). The grantee organization is the Northern California Institute for Research and Edu-

cation, and the study is coordinated by the Alzheimer's Disease Cooperative Study at the University

of California, San Diego. DATA are disseminated by the Laboratory for Neuro Imaging at the Uni-

versity of California, Los Angeles. This research was also supported by NIH grants P30AG01012 9,

K01 AG0305 14, and the Dana Foundation.

References

Acosta-Cabronero, J., Alley, S., Williams, G. B., Pengas, G. & Nestor, P. J. Diffusion Tensor Metrics
as Biomarkers in Alzheimer's Disease. PLOS ONE 7, e49072 (2012).

Acosta-Cabronero, J., Williams, G. B., Pengas, G. & Nestor, P. J. Absolute diffusivities define the
landscape of white matter degeneration in Alzheimer's disease. Brain 133, 529–539 (2010).

Albert, M. S., Moss, M. B., Tanzi, R., & Jones, K. (2001). Preclinical prediction of AD using
neuropsychological tests. Journal of the International Neuropsychological Society, 7(05), 631-639.

Alegret, M., Boada-Rovira, M., Vinyes-Junqué, G., Valero, S., Espinosa, A., Hernández, I., ... &
Tárraga, L. (2009). Detection of visuoperceptual deficits in preclinical and mild Alzheimer's disease.
Journal of clinical and experimental neuropsychology, 31(7), 860-867.

Almkvist, O. (1996). Neuropsychological features of early Alzheimer's disease: preclinical and
clinical stages. Acta Neurologica Scandinavica, 94(S165), 63-71.

Alonso-Montes, C. et al. Lagged and instantaneous dynamical influences related to brain structural
connectivity. Front. Psychol. 6, (2015).

Amor, T. A. et al. Extreme brain events: Higher-order statistics of brain resting activity and its
relation with structural connectivity. EPL Europhys. Lett. 111, 68007 (2015).

Apostolova, L. G., Lu, P., Rogers, S., Dutton, R. A., Hayashi, K. M., Toga, A. W., ... & Thompson, P.
M. (2008). 3D mapping of language networks in clinical and pre-clinical Alzheimer's disease. Brain
and language, 104(1), 33-41.

Arnáiz, E., & Almkvist, O. (2003). Neuropsychological features of mild cognitive impairment and
preclinical Alzheimer's disease. Acta Neurologica Scandinavica, 107(s179), 34-41.

Bassiony, M. M., Steinberg, M. S., Warren, A., Rosenblatt, A., Baker, A. S., & Lyketsos, C. G.
(2000). Delusions and hallucinations in Alzheimer's disease: prevalence and clinical correlates.
International journal of geriatric psychiatry, 15(2), 99-107.

Benjamini, Y. & Hochberg, Y. Controlling the False Discovery Rate: A Practical and Powerful
Approach to Multiple Testing. J. R. Stat. Soc. Ser. B Methodol. 57, 289–300 (1995).

Blacker, D., Lee, H., Muzikansky, A., Martin, E. C., Tanzi, R., McArdle, J. J., ... & Albert, M. (2007).
Neuropsychological measures in normal individuals that predict subsequent cognitive decline.
Archives of neurology, 64(6), 862-871.

Bosch, B. et al. Multiple DTI index analysis in normal aging, amnestic MCI and AD. Relationship
with neuropsychological performance. Neurobiol. Aging 33, 61–74 (2012).

Bozoki, A., Giordani, B., Heidebrink, J. L., Berent, S., & Foster, N. L. (2001). Mild cognitive
impairments predict dementia in nondemented elderly patients with memory loss. Archives of
Neurology, 58, 411–416.

Bäckman, L., Jones, S., Berger, A. K., Laukka, E. J., & Small, B. (2004). Multiple cognitive deficits
during the transition to Alzheimer's disease. Journal of internal medicine, 256(3), 195-204.

Bäckman, L., Jones, S., Berger, A. K., Laukka, E. J., & Small, B. J. (2005). Cognitive impairment in
preclinical Alzheimer's disease: a meta-analysis. Neuropsychology, 19(4), 520.

519Cerhan, J. H., Ivnik, R. J., Smith, G. E., Machulda, M. M., Boeve, B. F., Knopman, D. S., ... &
520Tangalos, E. G. (2007). Alzheimer's disease patients' cognitive status and course years prior to
521symptom recognition. Aging, Neuropsychology, and Cognition, 14(3), 227-235.

522Cheng, B. et al. Multimodal manifold-regularized transfer learning for MCI conversion prediction.
523Brain Imaging Behav. 9, 913–926 (2015).

524Chung, J. A., & Cummings, J. L. (2000). Neurobehavioral and neuropsychiatric symptoms in
525Alzheimer's disease. Neurologic clinics, 18(4), 829-846.

526Cole D.M., Smith S.M., Beckmann C.F. Advances and Pitfalls in the Analysis and Interpretation of
527Resting-State FMRI Data. Frontiers in Systems Neuroscience 4, 8 (2010).

528Daniel B. McArtor, D. B. M. MDMR: Multivariate Distance Matrix Regression. (2016).

529Dickerson, B. C., Sperling, R. A., Hyman, B. T., Albert, M. S., & Blacker, D. (2007). Clinical
530prediction of Alzheimer disease dementia across the spectrum of mild cognitive impairment.
531Archives of General Psychiatry, 64(12), 1443-1450.

532Diez, I. et al. A novel brain partition highlights the modular skeleton shared by structure and function.
533Sci. Rep. 5, 10532 (2015).

534Diez, I. et al. Enhanced pre-frontal functional-structural networks to support postural control deficits
535after traumatic brain injury in a pediatric population. Network Neuroscience 1–56 (2017).

536Douaud, G. et al. Brain Microstructure Reveals Early Abnormalities more than Two Years prior to
537Clinical Progression from Mild Cognitive Impairment to Alzheimer's Disease. J. Neurosci. 33, 2147–
5382155 (2013).

Douaud, G. et al. DTI measures in crossing-fibre areas: increased diffusion anisotropy reveals early

white matter alteration in MCI and mild Alzheimer's disease. NeuroImage 55, 880–890 (2011).

Elias, M. F., Beiser, A., Wolf, P. A., Au, R., White, R. F., & D'Agostino, R. B. (2000). The preclinical

phase of Alzheimer's disease: A 22-year prospective study of the Framingham cohort. Archives of

Neurology, 57, 808–813.

Eskildsen, S. F. et al. Prediction of Alzheimer's disease in subjects with mild cognitive impairment

from the ADNI cohort using patterns of cortical thinning. NeuroImage 65, 511–521 (2013).

Estevez Gónzalez, A., Kulisevsky, J., Boltes, A., Otermin, P., García-Sánchez, C. Rey. Verbal

Learning test is a useful tool for differential diagnosis in the preclinical phase of Alzheimer's disease:

comparison with mild cognitive impairment and normal aging. Int J Geriatr Psychiatry 2003; 18 (11):

1021-8.

Fearing, M.A., Bigler, E.D., Norton, M., Tschanz, J.A., Hulette, C., Leslie, C., Welsh-Bohmer, K., &

Cache County Investigators (2007). Autopsy-confirmed Alzheimer's disease versus clinically

diagnosed Alzheimer's disease in the Cache County Study on Memory and Aging: a comparison of

quantitative MRI and neuropsychological findings. Journal of Clinical and Experimental

Neuropsychology, 29(5): 553-560

Geslani, D. M., Tierney, M. C., Herrmann, N. & Szalai, J. P. Mild cognitive impairment: an

operational definition and its conversion rate to Alzheimer's disease. Dement. Geriatr. Cogn. Disord.

19, 383–389 (2005).

Gomez-Isla, T., Price, J., McKeel, D., Morris, J., Growdon, J., & Hyman, B. (1996). Profound loss of

layer II entorhinal cortex neurons occurs in very mild Alzheimer's disease. Journal of Neuroscience,

16, 4491– 4500.

Grober, E., Hall, C. B., Lipton, R. B., Zonderman, A. B., Resnick, S. M., & Kawas, C. (2008).

Memory impairment, executive dysfunction, and intellectual decline in preclinical Alzheimer's

disease. Journal of the International Neuropsychological Society, 14(02), 266-278.

Hanyu, H. et al. Diffusion-weighted MR imaging of the hippocampus and temporal white matter in

Alzheimer's disease. J. Neurol. Sci. 156, 195–200 (1998).

Hardy, J. Alzheimer's disease: the amyloid cascade hypothesis: an update and reappraisal. J.

Alzheimers Dis. JAD 9, 151–153 (2006).

Honea, R. A., Vidoni, E., Harsha, A. & Burns, J. M. Impact of APOE on the Healthy Aging Brain: A

Voxel-Based MRI and DTI Study. J. Alzheimers Dis. JAD 18, 553–564 (2009).

Joubert, S., Brambati, S. M., Ansado, J., Barbeau, E. J., Felician, O., Didic, M., ... & Kergoat, M. J.

(2010). The cognitive and neural expression of semantic memory impairment in mild cognitive

impairment and early Alzheimer's disease. Neuropsychologia, 48(4), 978-988.

Khedher, L., Ramírez, J., Górriz, J. M., Brahim, A. & Segovia, F. Early diagnosis of Alzheimer´s

disease based on partial least squares, principal component analysis and support vector machine

using segmented MRI images. Neurocomputing 151, Part 1, 139–150 (2015).

Li, H. et al. Hierarchical Interactions Model for Predicting Mild Cognitive Impairment (MCI) to

Alzheimer's Disease (AD) Conversion. PLOS ONE 9, e82450 (2014).

Liu, J. et al. White Matter Changes in Patients with Amnestic Mild Cognitive Impairment Detected

by Diffusion Tensor Imaging. PLOS ONE 8, e59440 (2013).

Liu, Y. et al. Predicting AD Conversion: Comparison between Prodromal AD Guidelines and

Computer Assisted PredictAD Tool. PLOS ONE 8, e55246 (2013).

Marinazzo, D. et al. Information transfer and criticality in the Ising model on the human connectome.

PLoS ONE 9, e93616 (2014).

McArdle, B. H. & Anderson, M. J. Fitting Multivariate Models to Community Data: A Comment on

Distance-Based Redundancy Analysis. Ecology 82, 290–297 (2001).

Medina, D. et al. White matter changes in mild cognitive impairment and AD: A diffusion tensor

imaging study. Neurobiol. Aging 27, 663–672 (2006).

Mielke, M. M. et al. Fornix integrity and hippocampal volume predict memory decline and

progression to Alzheimer's disease. Alzheimers Dement. J. Alzheimers Assoc. 8, 105–113 (2012).

Milham M.P. Open Neuroscience Solutions for the Connectome-wide Association Era. Neuron. 73,

214–218 (2012).

Mitchell, A. J. & Shiri-Feshki, M. Rate of progression of mild cognitive impairment to dementia –

meta-analysis of 41 robust inception cohort studies. Acta Psychiatr. Scand. 119, 252–265 (2009).

Mori, S., Crain, B. J., Chacko, V. P. & van Zijl, P. C. Three-dimensional tracking of axonal

projections in the brain by magnetic resonance imaging. Ann. Neurol. 45, 265–269 (1999).

Mueller, S. G. et al. The Alzheimer's disease neuroimaging initiative. Neuroimaging Clin. N. Am. 15,

869–877, xi–xii (2005).

Petersen, R. C. et al. Prevalence of mild cognitive impairment is higher in men The Mayo Clinic

Study of Aging. Neurology 75, 889–897 (2010).

Preti, M. G. et al. Assessing Corpus Callosum Changes in Alzheimer's Disease: Comparison between

Tract-Based Spatial Statistics and Atlas-Based Tractography. PLOS ONE 7, e35856 (2012).

602Rapp, M. A., & Reischies, F. M. (2005). Attention and executive control predict Alzheimer disease in
603late life: results from the Berlin Aging Study (BASE). The American Journal of Geriatric Psychiatry,
60413(2), 134-141.

605Ritter, K. et al. Multimodal prediction of conversion to Alzheimer's disease based on incomplete
606biomarkers∗. Alzheimers Dement. Diagn. Assess. Dis. Monit. 1, 206–215 (2015).

607Rose, S. et al. Loss of connectivity in Alzheimer's disease: an evaluation of white matter tract
608integrity with colour coded MR diffusion tensor imaging. J. Neurol. Neurosurg. Psychiatry 69, 528–
609530 (2000).

610Rozzini, L. et al. Conversion of amnestic Mild Cognitive Impairment to dementia of Alzheimer type
611is independent to memory deterioration. Int. J. Geriatr. Psychiatry 22, 1217–1222 (2007).

612Salat, D. H. et al. White matter pathology isolates the hippocampal formation in Alzheimer's disease.
613Neurobiol. Aging 31, 244–256 (2010).

614Schmidtke, K. & Hermeneit, S. High rate of conversion to Alzheimer's disease in a cohort of
615amnestic MCI patients. Int. Psychogeriatr. 20, 96–108 (2008).

616Senanarong, V., Cummings, J. L., Fairbanks, L., Mega, M., Masterman, D. M., O'connor, S. M., &
617Strickland, T. L. (2003). Agitation in Alzheimer's disease is a manifestation of frontal lobe
618dysfunction. Dementia and geriatric cognitive disorders, 17(1-2), 14-20.

619Shehzad, Z. et al. A multivariate distance-based analytic framework for connectome-wide association
620studies. NeuroImage 93, Part 1, 74–94 (2014).

621Small, B. J., Mobly, J. L., Laukka, E. J., Jones, S., & Bäckman, L. (2003). Cognitive deficits in
622preclinical Alzheimer's disease. Acta Neurologica Scandinavica, 107(s179), 29-33.

Small, S. A., Perara, G., DeLaPaz, R., Mayeux, R., & Stern, Y. (1999). Differential regional

dysfunction of the hippocampal formation among elderly with memory decline and Alzheimer's

disease. Annals of Neurology, 45, 466–472.

Smith, C. D. et al. White matter diffusion alterations in normal women at risk of Alzheimer's disease.

Neurobiol. Aging 31, 1122–1131 (2010).

Storandt, M. (2008). Cognitive deficits in the early stages of Alzheimer's disease. Current Directions

in Psychological Science, 17(3), 198-202.

Stricker, N. H. et al. Decreased white matter integrity in late-myelinating fiber pathways in

Alzheimer's disease supports retrogenesis. NeuroImage 45, 10–16 (2009).

Teipel, S. et al. Multimodal imaging in Alzheimer's disease: validity and usefulness for early

detection. Lancet Neurol. 14, 1037–1053 (2015).

Tierney, M. C., Szalai, J. P., Snow, W. G., & Fisher, R. H. (1996). The prediction of Alzheimer

disease: The role of patient and informant perceptions of cognitive deficits. Archives of Neurology,

53, 423–427

Tzourio-Mazoyer, N. et al. Automated anatomical labeling of activations in SPM using a

macroscopic anatomical parcellation of the MNI MRI single-subject brain. Neuroimage 15, 273–289

(2002).

Wee, C.-Y., Yap, P.-T., Shen, D. & Alzheimer's Disease Neuroimaging Initiative. Prediction of

Alzheimer's disease and mild cognitive impairment using cortical morphological patterns. Hum.

Brain Mapp. 34, 3411–3425 (2013).

Westman, E. et al. Sensitivity and Specificity of Medial Temporal Lobe Visual Ratings and

Multivariate Regional MRI Classification in Alzheimer's Disease. PLOS ONE 6, e22506 (2011).

Whittaker, K. W., Burdon, M. A. & Shah, P. Visual field loss and Alzheimer's disease. Eye 16, 206–

208 (2002).

Wimo, A., Ljunggren, G. & Winblad, B. Costs of dementia and dementia care: a review. Int. J.

Geriatr. Psychiatry 12, 841–856 (1997).

Y. et al. Regional coherence changes in the early stages of Alzheimer's disease: A combined

structural and resting-state functional MRI study. NeuroImage 35, 488–500 (2007).

Zapala, M. A. & Schork, N. J. Multivariate regression analysis of distance matrices for testing

associations between gene expression patterns and related variables. Proc. Natl. Acad. Sci. U. S. A.

103, 19430–19435 (2006).

Zhang, S. et al. in Cochrane Database of Systematic Reviews (John Wiley & Sons, Ltd, 2014).

Zhou, Y. et al. Abnormal connectivity in the posterior cingulate and hippocampus in early

Alzheimer's disease and mild cognitive impairment. Alzheimers Dement. 4, 265–270 (2008).

List of Tables

Table 1: t-test and Chi^2 test across groups

EMCI: Early mild cognitive impairment; LMCI=Late mild cognitive impairment; AD= Alzheimer
disease.

	Control vs EMCI		Control vs LMCI		Control vs AD	
	test value	*p-value*	*test value*	*p-value*	*test value*	*p-value*
Age (t-test)	0.0349	0.9722	0.5539	0.5814	0.2071	0.8365
Sex (Chi^2 test)	0.2338	0.6287	0.2338	0.6287	0.2338	0.6287

666 **Table 2: Further information about ADNI group classification.**

667 EMCI: Early mild cognitive impairment; LMCI=Late mild cognitive impairment; AD= Alzheimer
668 disease; LMIIS=Logical Memory II subscale; MMSE= Mini Mental State Examination; CDR=
669 Clinical Dementia Rating.

	Control	EMCI	LMCI	AD

Progressive brain disconnection in AD

LMIIS (maximum of 25 points)				
Education ≥16 years	≥ 9	[9-11]	≤ 8	≤ 8
Education [8-15] years	≥ 5	[5-9]	≤ 4	≤ 4
Education [0-7] years	≥ 3	[3-6]	≤ 2	≤ 2
MMSE (Maximum of 30 points)	[24-30]	[24-30]	[24-30]	[20-26]
CDR	0	0.5	0.5	0.5 or 1
Memory Box Score (subpart of CDR)	0	at least 0.5	at least 0.5	NA

672 **Table 3: p-values associated to each module from the brain hierarchical atlas.**

673 EMCI: Early mild cognitive impairment; LMCI=Late mild cognitive impairment; AD= Alzheimer

674 disease; * 0.01<p<0.05; ** 0.005 < p <0.01: *** p<0.005.

676 Connectivity alterations start in module 18 (marked in black and underlined), and in later stages grow

677 (increasing significance) and extend to a multitude of different other modules.

Module	Control vs EMCI	Control vs LMCI	Control vs AD
1	0.956	0.753	0.023*
2	0.956	0.466	0.049*
3	0.956	0.441	0.049*
4	0.880	0.532	0.031*
5	0.859	0.689	0.973
6	0.859	0.438	0.546
7	0.956	0.900	0.503
8	0.859	0.449	0.031*
9	0.859	0.600	0.591
10	0.956	0.900	0.627
11	0.956	0.438	0.759

12	0.956	0.466	0.031*
13	0.859	0.600	0.531
14	0.956	0.438	0.006**
15	0.956	0.753	0.031*
16	0.956	0.986	0.031*
17	0.890	0.898	0.546
<u>18</u>	**<u>0.399</u>**	**<u>0.007</u>****	**<u>0.002</u>*****
19	0.956	0.438	0.109
20	0.956	0.986	0.972

List of Captions

Figure 1: Methodological sketch. Alzheimer's disease progression is addressed across three stages.

Four groups of 36 subjects each at different stages of AD (Control, Early and Late MCI, Alzheimer)

following the ADNI classification criterion. All groups have been balanced with respect to age, sex

and years of education. Brain connectivity patterns and its relation with disease progression are

accomplished by comparing the control group with the rest of groups, i.e. Control vs EMCI (stage I),

Control vs LMCI (stage II) and Control vs AD (stage III).

Figure 2: Multivariate distance matrix regression analysis to find differences in brain

connectivity patterns across severity progression of AD. In a first step (*Image preprocessing* in a

red box), brain images are preprocessed by using standard techniques, mainly eddy current and head

motion corrections). Next, diffusion tensor reconstructions are built that allows calculating the

tractography for each subject (further details in Methods). In the next step (*Distance matrix*

calculation in a green box), first the streamline number connectivity matrix is obtained (here,

represented by λ), one per subject, corresponding to 20×20 entries of values given by α .

Second, the connectivity patterns of subjects for a given module are used to construct the distance

matrix in the subject space by means of Pearson correlation coefficients. Once the distance matrix for

a given module is calculated (here, we highlight in red the first row that corresponds to the first

module), we test in the third step (*Multivariate regression* in a blue box) whether the variability in

distance between different groups is statistically related with disease, for which we compare the

observed results with a simulated distribution given by N permutations of the labels. We repeat this

operation for every module. We finally apply the fourth step (*False discovery rate corrections* in a

black box) to correct for multiple comparisons.

Figure 3: Pseudo F-statistic brain maps across the severity progression of AD. Brain

disconnection as disease progresses is quantitatively addressed by looking at the Pseudo F-statistic

values of the modules. At first stages (Control vs EMC, top), fibers deterioration is not sufficient to

yield significant changes in modules connectivity patterns. In the following stage (Control vs LMCI -

middle), the connectivity pattern of module 18, which involves parts of the hippocampus, entorhinal

cortex, amygdala and other memory-related areas, disconnects statistically with respect to control (p-

val = 0.007). Such connectivity differences are widely spread to the rest of the brain at the final stage

(Control vs AD, bottom).

I. IMAGE PREPROCESSING

Eddy current and motion correction → Diffusion Tensor Reconstruction →

II. DISTANCE MATRIX CALCULATION

Streamline number   **Streamline 1st region**

Subject 1
$$\lambda_1^{(20\times20)} = \begin{pmatrix} \alpha_{1,1} & \alpha_{1,2} & \cdots & \alpha_{1,20} \\ \alpha_{2,1} & \alpha_{2,2} & \cdots & \alpha_{2,20} \\ \vdots & \vdots & \ddots & \vdots \\ \alpha_{20,1} & \alpha_{20,2} & \cdots & \alpha_{20,20} \end{pmatrix}$$

Subject 2
$$\lambda_2^{(20\times20)} = \begin{pmatrix} \alpha_{1,1} & \alpha_{1,2} & \cdots & \alpha_{1,20} \\ \alpha_{2,1} & \alpha_{2,2} & \cdots & \alpha_{2,20} \\ \vdots & \vdots & \ddots & \vdots \\ \alpha_{20,1} & \alpha_{20,2} & \cdots & \alpha_{20,20} \end{pmatrix}$$

GROUP1
____
GROUP2

Subject 71
$$\lambda_{71}^{(20\times20)} = \begin{pmatrix} \alpha_{1,1} & \alpha_{1,2} & \cdots & \alpha_{1,20} \\ \alpha_{2,1} & \alpha_{2,2} & \cdots & \alpha_{2,20} \\ \vdots & \vdots & \ddots & \vdots \\ \alpha_{20,1} & \alpha_{20,2} & \cdots & \alpha_{20,20} \end{pmatrix}$$

Subject 72
$$\lambda_{72}^{(20\times20)} = \begin{pmatrix} \alpha_{1,1} & \alpha_{1,2} & \cdots & \alpha_{1,20} \\ \alpha_{2,1} & \alpha_{2,2} & \cdots & \alpha_{2,20} \\ \vdots & \vdots & \ddots & \vdots \\ \alpha_{20,1} & \alpha_{20,2} & \cdots & \alpha_{20,20} \end{pmatrix}$$

$$d_{ij} = \sqrt{2(1-r_{ij})} \rightarrow D(72\times72) = \begin{pmatrix} d_{1,1} & d_{1,2} & \cdots & d_{1,72} \\ d_{2,1} & d_{2,2} & \cdots & d_{2,72} \\ \vdots & \vdots & \ddots & \vdots \\ d_{72,1} & d_{72,2} & \cdots & d_{72,72} \end{pmatrix}$$

Distance Matrix

REPEAT WITH ALL MODULES

III. MULTIVARIATE REGRESSION

Pseudo F-stat measured:

$$F = \frac{SS_A/(m-1)}{SS_W/(N-m)}$$

Recalculate F from permuted data

$$p_{val} = \frac{\#F_{permuted} > F}{N_{permutations}}$$

IV. FALSE DISCOVERY RATE CORRECTIONS

$$FDR = < \frac{v}{v+s} > = < \frac{v}{R} >$$

control control
vs vs
EMCI

control
vs
LMCI

No significant modules

Module 18

pseudo-F statistic

0.08

0.03

Module 18

Module 1

Control vs AD

Module 2

Module 12

Module 3

Module 14

Module 4

Module 15

Module 8

Module 16

Table S1: ADNI subjects within each group

EMCI: Early mild cognitive impairment; LMCI=Late mild cognitive impairment; AD= Alzheimer disease; M=Male, F=Female.

Control			EMCI			LMCI			AD		
SubjectId	Sex	Age	SubjectId	Sex	Age	SubjectId	Sex	Age	SubjectId	Sex	Age
003_S_4119	M	79	003_S_2374	F	81	003_S_0908	F	70	003_S_4373	F	71
003_S_4839	M	66	007_S_2394	M	69	003_S_4354	M	76	003_S_5165	M	79
007_S_4488	M	73	016_S_4575	F	62	016_S_4584	F	78	003_S_5187	F	62
007_S_4516	M	72	021_S_2077	M	81	016_S_4646	F	61	005_S_4707	M	68
007_S_4620	M	77	021_S_2100	F	88	016_S_4902	F	75	005_S_4910	F	82
016_S_4121	M	89	021_S_2125	F	78	021_S_4402	F	73	005_S_5038	M	82

021_S_4558	F	71	021_S_2142	F	83	021_S_4633	F	73	005_S_5119	F	77
029_S_4279	M	84	021_S_4419	F	65	021_S_4857	M	68	007_S_4568	F	71
029_S_4290	M	74	021_S_4659	M	86	027_S_4729	F	78	007_S_4911	M	75
029_S_4384	M	62	021_S_4744	F	73	027_S_4757	F	63	007_S_5196	F	73
029_S_4385	F	68	029_S_2370	F	64	027_S_4804	M	80	016_S_4591	F	66
029_S_4585	M	66	029_S_2395	M	73	027_S_4869	M	77	016_S_4887	M	75
029_S_4652	M	79	029_S_4327	M	83	027_S_4873	M	83	016_S_4963	F	72
057_S_0934	F	77	029_S_5135	M	77	027_S_4936	M	78	016_S_5057	M	75
094_S_4234	M	70	094_S_2201	F	64	027_S_4943	M	76	016_S_5251	F	66
094_S_4459	F	68	094_S_2216	M	69	027_S_4955	M	72	021_S_4718	M	79
094_S_446	F	67	094_S_223	M	69	052_S_462	M	69	021_S_492	M	77

0			8			6			4		
094_S_4503	F	72	094_S_2367	M	75	052_S_4807	F	72	027_S_4801	M	78
094_S_4649	M	66	094_S_4434	M	68	052_S_4945	M	57	027_S_4802	M	83
098_S_4002	F	74	098_S_2052	M	74	057_S_4888	M	75	027_S_4938	M	71
098_S_4003	F	72	098_S_2071	M	85	057_S_4909	F	78	027_S_4962	F	80
098_S_4018	M	76	099_S_4205	F	84	094_S_4162	F	71	027_S_4964	M	81
098_S_4050	M	77	099_S_4498	F	80	094_S_4295	F	70	052_S_5062	F	71
098_S_4275	M	73	109_S_2110	F	68	094_S_4630	F	66	094_S_4089	M	74
098_S_4506	M	72	109_S_2111	M	72.	109_S_4471	M	73	094_S_4737	F	74
099_S_4076	F	75	109_S_2200	F	76	109_S_4531	M	74	098_S_4201	F	64
127_S_4148	M	73	109_S_4380	M	72	126_S_4458	F	76	098_S_4215	M	82

127_S_4198	F	78	109_S_4455	M	64	126_S_4507	M	78	109_S_4378	M	80
127_S_4604	M	65	109_S_4594	M	62	126_S_4675	M	80	126_S_4494	M	71
127_S_4645	F	76	126_S_2360	M	64	126_S_4712	M	74	127_S_4749	F	78
127_S_4843	F	73	126_S_4891	M	60	126_S_4743	M	70	127_S_4992	F	64
129_S_0778	M	80	127_S_4301	M	75	126_S_4896	M	68	127_S_5028	M	62
129_S_4369	M	70	127_S_4624	F	78	127_S_4197	M	79	127_S_5056	M	85
129_S_4371	M	70	127_S_4765	M	76	127_S_4210	M	64	127_S_5058	M	62
129_S_4396	F	81	129_S_2347	M	73	127_S_4240	M	71	127_S_5067	M	81
131_S_0123	M	81	129_S_4220	F	73	129_S_4287	F	73	127_S_5095	M	66

Table S2: Examples of pseudo F-statistics, between-group and within-group sum of squares. Three different situations: Node 10, that does not provide any significant change in pattern connectivity; Node 16, significantly different in stage III; and Node 18, with pattern connectivity significantly different in stages II and III.

EMCI: Early mild cognitive impairment; LMCI=Late mild cognitive impairment; AD= Alzheimer disease

Module	Control vs EMCI			Control vs LMCI			Control vs AD		
	F	*SSA*	*SSW*	*F*	*SSA*	*SSW*	*F*	*SSA*	*SSW*
10	0.527	0.008	1.028	0.630	0.008	0.940	0.894	0.012	0.941
16	0.380	0.019	3.457	0.285	0.014	3.404	3.410	0.173	3.550
18	3.057	0.024	0.558	5.854	0.049	0.595	6.018	0.051	0.588

Table S3: Group comparison not involving the healthy control group.

EMCI: Early mild cognitive impairment; LMCI=Late mild cognitive impairment;

*AD= Alzheimer disease * $0.01<p<0.05$; ** $0.005 < p < 0.01$: *** $p<0.005$.*

Module	EMCI vs LMCI	LMCI vs AD	EMCI vs AD
1	0.869	0.089	0.153
2	0.473	0.089	0.049*
3	0.474	0.089	0.049*
4	0.473	0.433	0.049*
5	0.474	0.433	0.352
6	0.736	0.372	0.969
7	0.869	0.395	0.688
8	0.869	0.395	0.055
9	0.473	0.235	0.352
10	0. 473	0.222	0.383
11	0. 869	0.533	0.969
12	0. 473	0.395	0.383
13	0.473	0.089	0.352
14	0.473	0.060	0.035*

15	0.930	0.089	0.092
16	0.869	0.089	0.049*
17	0.869	0.410	0.905
18	0.736	0.698	0.623
19	0.474	0.089	0.383
20	0.787	0.698	0.969