**Sex chromosome evolution, heterochiasmy and physiological QTL in the**

**salmonid Brook Charr *Salvelinus fontinalis***

Ben J. G. Sutherland[*], Ciro Rico[†‡], Céline Audet[§] and Louis Bernatchez[*]

* Institut de Biologie Intégrative et des Systèmes (IBIS), Université Laval, Québec, QC, Canada G1V
0A6

† School of Marine Studies, Molecular Diagnostics Laboratory, The University of the South Pacific,
Laucala Campus, Suva, Fiji

‡ Department of Wetland Ecology, Estación Biológica de Doñana (EBD-CSIC), c/Américo Vespucio s/n,
41092 Sevilla, Spain

§ Institut des Sciences de la Mer de Rimouski, Université du Québec à Rimouski, Rimouski, QC, Canada
G5L 3A1

**Running title:** Sex, heterochiasmy and QTL in Charr

**Keywords:** heterochiasmy; salmon; sex chromosomes; QTL; whole genome duplication

**Author for Correspondence**:

Ben Sutherland
Institut de Biologie Intégrative et des Systèmes (IBIS)
Université Laval, Québec, QC, Canada
G1V 0A6
Email: ben.sutherland.1@ulaval.ca (BJGS)

## **ABSTRACT**

Whole genome duplication can have large impacts on genome evolution. However, much remains

unknown about these impacts, such as the mechanisms of coping with a duplicated sex determination

system, which may result in increased sex determination mechanism diversity. Sexual conflict (i.e. alleles

having different optimums in each sex) can result in sequestration of genes into non-recombining sex

chromosomes. Development of sex chromosomes may involve heterochiasmy (i.e. sex-specific

recombination rate), which is also poorly understood. Family Salmonidae is a model system for these

phenomena, having undergone autotetraploidization and subsequent rediploidization in most of the

genome at the base of the lineage. The salmonid master sex determining gene is known, and many species

have non-homologous sex chromosomes, putatively due to transposition of this gene. In this study, we

identify the sex chromosome of Brook Charr *Salvelinus fontinalis* and compare sex chromosome

identities across the lineage (eight species, four genera). Although non-homology is frequent, homologous

sex chromosomes and other consistencies are present in distantly related species, indicating probable

convergence on specific sex and neo-sex chromosomes. We also characterize strong heterochiasmy with

2.7-fold more crossovers in maternal than paternal haplotypes with paternal crossovers biased to

chromosome ends. Y chromosome crossovers are restricted to a single end of the chromosome, and this

chromosome contains a large interspecific inversion, although its status between males and females

remains unknown. Finally, we identify QTL for 21 unique growth, reproductive and stress-related

phenotypes to improve knowledge of the genetic architecture of these traits important to aquaculture and

evolution.

## INTRODUCTION

Characterizing the genetic architecture of ecologically-relevant phenotypes is essential for organism and genome evolution research (Rogers and Bernatchez 2007; Gagnaire *et al.* 2013) and selective breeding (Yáñez *et al.* 2014). Regions of the genome associated with specific traits can be identified by quantitative trait loci (QTL) analysis (Mackay 2001) or genome-wide association studies (GWAS; Bush and Moore 2012). Mapped traits can include morphological, behavioral, physiological or molecular phenotypes, but must show sufficient heritable variation. Power to detect QTL is determined by the number of individuals in the study (Henning *et al.* 2014), the effect size of the QTL, allele frequencies (Mackay *et al.* 2009) and the degree of polygenic control of the trait (Rockman 2012; Ashton *et al.* 2016). QTL mapping precision depends on recombination frequency (Mackay *et al.* 2009) as well as map density, although QTL are often only in linkage with causative mutations, which are rarely identified (Slate 2005). Trait genetic architecture can differ between families or populations (e.g. Santure *et al.* 2015) but parallelism and shared QTL can be identified (e.g. Laporte *et al.* 2015; Larson *et al.* 2015). It is therefore valuable to analyze multiple crosses to understand the broader implications of a QTL (e.g. Hecht *et al.* 2012; Palti *et al.* 2015; Lv *et al.* 2016), to more accurately determine the amount of variation explained by the QTL (Slate 2005), and to identify QTL that are not dependent on specific genetic backgrounds, which is particularly valuable for marker-assisted selection (Lv *et al.* 2016).

Advances in massively parallel sequencing (MPS) technology and comparative genomics have benefited QTL and association studies in several ways. MPS greatly increases marker density (Catchen *et al.* 2011; Ashton *et al.* 2016), provides markers with flanking sequence that can be aligned against reference genomes for integrating across species (Sutherland *et al.* 2016) or for identifying genes near QTL to inform on potential drivers underlying a trait (e.g. McKinney *et al.* 2016; Johnston *et al.* 2016). When causative mutations are not known, detecting orthologous QTL in other species can provide further evidence for a region or gene being related to a trait (Mackay 2001; Larson *et al.* 2015). As an example, QTL for recombination rate in mammalian model and non-model systems occur near the same genes (Johnston *et al.* 2016). Comparative genomics clearly has an important role in identifying drivers of trait variation.

Genetic architecture can be strongly affected by sexually antagonistic selection and sex determination. Sexually antagonistic alleles (i.e. alleles that benefit sexes differently) produce genetic conflict (Mackay 2001; Charlesworth *et al.* 2005), which can be resolved by the sequestration of alleles in non-recombining sex chromosomes. As an example of the effect this can have on genome architecture, the *Drosophila* Y chromosome is made almost exclusively of genes that have migrated from other chromosomes, presumably due to their specific benefit to males (Carvalho 2002). An additional benefit

occurs by constant sex-specific selection occurring for alleles on the Y (or W) chromosome, as these are
always only in the heterogametic sex (Lahn *et al.* 2001). However, the lack of recombination between the
sex chromosomes can also result in Y degeneration due to accumulation of mutations that are not able to
be purged through recombination with X (Charlesworth 1991). A different resolution to genetic conflict
involves sex-dependent dominance, whereby allelic dominance depends on the sex of the individual,
which need not be on the sex chromosome (Barson *et al.* 2015). Much remains to be understood about
resolving these conflicts.

Genome evolution can also be affected by large mutational forces, such as polyploidization
events including whole genome duplication (WGD) (Ohno 1970), which may disrupt sex determination
systems (Davidson *et al.* 2009). Although details of this disruption remain generally unknown, some
hypotheses have been proposed involving the independent segregation of duplicated sex determining
chromosomes or imbalances in gene dosages when X inactivation occurs (Muller 1925; Orr 1990;
Davidson *et al.* 2009). Highly diverse sex determination systems are observed in teleosts (Marshall
Graves and Peichel 2010), which may have been influenced by the teleost-specific WGD due to a post-
WGD adoption of numerous different sex determination mechanisms (Mank and Avise 2009). Evolution
of sex determination post-WGD may occur through the mutational disruption of one duplicated portion of
the existing system, or by the development of a new system (Davidson *et al.* 2009), which can thus result
in the evolution of new sex chromosomes.

Sex chromosome evolution may be facilitated by differences in recombination rates between the
sexes (i.e. heterochiasmy) (Charlesworth *et al.* 2005). The evolution of heterochiasmy remains under
investigation, although several explanations have been proposed (Lenormand and Dutheil 2005;
Brandvain and Coop 2012; Lenormand *et al.* 2016). First, sexes can experience different extents of
selection at haploid stages (Lenormand 2003), and heterochiasmy permits retention of epistatically-
interacting alleles within a haplotype specifically within the sex experiencing more haploid selection
(Lenormand and Dutheil 2005). Second, physical meiotic differences may play a role; female meiosis
occurs with a long delay, and chiasma (i.e. locations where crossovers occur) stabilize chromatids during
this process (Lenormand 2003; Lenormand *et al.* 2016). Third, recombination protects from meiotic drive,
to which the sexes have different susceptibilities (Brandvain and Coop 2012). Other hypotheses have also
been proposed (Trivers 1998; Lenormand 2003). However, in general it is unclear which of the above
explanations have the largest influence, and thus the relationships between WGD, heterochiasmy and sex
chromosome evolution require further study.

Salmonids (Family Salmonidae) are an ideal system to study genetic architecture and sex
determination post-WGD (Davidson *et al.* 2010). The salmonid genome remains in a residually tetraploid
state, where some chromosomal telomeric regions continue recombining between homeologous

chromosomes and others have rediploidized (Allendorf and Thorgaard 1984; Allendorf *et al.* 2015; May

and Delany 2015; Lien *et al.* 2016). Salmonid sex determination is genetically controlled (Davidson *et al.*

2009) by a truncated gene from the *interferon-response factor* transcription factor family, *sdY* (sexually

dimorphic on the Y-chromosome; Yano *et al.* 2012a). *sdY* may be a salmonid innovation as it has not yet

been identified in the non-duplicated sister group for the salmonid WGD, Northern Pike *Esox lucius*

(Yano *et al.* 2012b). Male genome-specific conservation of *sdY* occurs in more than ten salmonid species,

but some exceptions exist, including the Lake Whitefish *Coregonus clupeaformis* and European

Whitefish *C. lavaretus* (Yano *et al.* 2012b), and some Atlantic Salmon *Salmo salar* and Sockeye Salmon

*Oncorhynchus nerka* individuals (Eisbrenner *et al.* 2013; Larson *et al.* 2016). Sex chromosomes are not

homologous among many salmonid species, potentially due to transposition of *sdY* between chromosomes

(Woram *et al.* 2003). Additional evidence for transposition includes repetitive flanking regions with

putative transposable elements (Brunelli *et al.* 2008; Lubieniecki *et al.* 2015) and sequence conservation

that abruptly stops outside of the sex determination cassette (Faber-Hammond *et al.* 2015). This

transposition to different chromosomes may be delaying Y degeneration (Yano *et al.* 2012b; Lubieniecki

*et al.* 2015). In general, the salmonids are at an early stage of sex chromosome evolution (Phillips and

Ihssen 1985; Yano *et al.* 2012b) where sex chromosomes are homomorphic (Devlin *et al.* 1998; Phillips

and Ráb 2001; Davidson *et al.* 2009). Male salmonids have low recombination rates relative to females

with crossover events primarily occurring at telomeric regions, as observed in Rainbow Trout *O. mykiss*

(Sakamoto *et al.* 2000) and Atlantic Salmon (Moen *et al.* 2004). Heterochiasmy is not viewed in the sister

species of the salmonid WGD Northern Pike (Rondeau *et al.* 2014) and therefore salmonids are a valuable

model to study the evolution and effects of heterochiasmy in relation to sex determination post-WGD.

The combination of characterizing heterochiasmy, sex chromosome identity and the genetic

architecture for reproductive, growth and stress response traits provides much-needed information

regarding the function of the Brook Charr *Salvelinus fontinalis* genome post-duplication. The goals of this

study were to use a high-density genetic map for Brook Charr (Sutherland *et al.* 2016) to (*a*) identify the

sex-linked chromosome; (*b*) quantify heterochiasmy in this mapping family while correcting for probable

genotyping errors; and (*c*) search for growth, stress resistance and reproduction-related QTL. Furthermore,

using the recent characterization of homology to ancestral chromosomes and homeolog identification

among the salmonids (Sutherland *et al.* 2016), we subsequently compare identities of sex chromosomes

and identified QTL across the salmonids to identify consistencies, and discuss the implications of

consistencies as well as the observed heterochiasmy regarding sex chromosome evolution.

## METHODS

**Fish and phenotyping**

Juvenile Brook Charr used in this study were the same individuals used to construct a low-density genetic map and perform QTL analysis for 21 phenotypes (29 including repeated measurements occurring at three time points; Table S1) by Sauvage *et al.* for growth (2012a) and reproductive QTL (2012b). Fish were raised in tanks as previously described until 65-80 g, at which point weight, length and condition factor were measured. These phenotypes were measured on the same fish two and six months after the initial measurements. Growth rate was calculated between the multiple sampling times. At the final sampling, all phenotypes were collected. Stress response was also evaluated at this final sampling through an acute handling stress by reducing water levels, capturing fish without chasing and holding out of water for one minute in order to phenotype the stress response using blood parameters chloride, osmolality and cortisol before and after the stress. After fish had re-acclimatized, they were anaesthetized and killed by decapitation as per regulations of Canadian Council of Animal Protection recommendations and protocols approved by the University Animal Care Committee, as previously reported (Sauvage *et al.* 2012a). The sex of each individual was determined by visual inspection of the gonads as reported by Sauvage *et al.* (2012b).

**Genetic map and quality control of markers and phenotypes**

A recently developed high-density genetic map with 3826 markers was used with genotypes for 192 offspring (Sutherland *et al.* 2016). In brief, genotype data was obtained using the population module of STACKS v.1.32 (Catchen *et al.* 2011), phased in JoinMap v.4.1 (van Ooijen 2006), and imported into R/qtl (Broman *et al.* 2003) using the *read.cross* function with data interpreted as a four-way cross type in the *mapqtl* format (see File S1 for map, genotype and phenotype input files).

All 29 phenotypes (including eight measures at multiple time points) related to blood parameters, growth, growth-related gene expression, reproduction and stress response were used to search for QTL (Table S1). Correlation between phenotypes was evaluated using Pearson correlation in R (R Development Core Team 2017) and a correlation plot was generated using the R package *corrplot* (v.0.77; Wei and Simko 2017). Phenotypes were inspected for normal distribution, and when required, log transformed (Broman and Sen 2009). Outlier phenotype values (>3 SD from the mean) were removed to prevent spurious associations (Broman and Sen 2009), including two individuals each for T1-T2 and T2-T3 growth rates, two individuals for length at T2, four individuals for condition factor at T2, one individual for change in osmolality and one individual for sperm diameter.

Markers present in the map were tested for segregation distortion by chi-square tests for

Mendelian segregation in R/qtl and removed when $p \leq 0.01$ (Broman and Sen 2009). A total of 157

markers with significant segregation distortion were removed, leaving a remainder of 3669 markers.

Proportions of identical genotypes were tested in R/qtl to ensure that there were no mis-labeled samples.

Recombination fraction between marker pairs was estimated using Expectation Maximization algorithm

within *est.rf* in R/qtl. The minimum number of obligate crossover events was calculated per individual

using *count.XO* in R/qtl, and an outlier sample with 1093 crossovers was removed (other samples had

mean and median crossovers of 101 and 83, respectively, before correcting for unlikely double

crossovers).

**Recombination rate**

To characterize heterochiasmy in the mapping family parents, the *plotGenotypes* function of R/qtl was

used to identify positions of crossovers per parental chromosome (total = 84 chromosomes per individual

offspring) and modified to export these positions (see Data Availability section for all code used in the

analysis). Male-specific markers were not included in the original map due to low recombination rate and

poor positioning (Sutherland *et al.* 2016), and therefore to avoid bias of including female-specific but not

male-specific markers, crossovers were evaluated in a map with only markers informative in both sexes

(i.e. *ef* x *eg* and *hk* x *hk*). Furthermore, as recombination rates can be inflated by a genotyping error

appearing to be flanked by two false recombination events (Hackett and Broadfoot 2003; Slate 2008),

which can also occur in RAD-seq data (Andrews *et al.* 2016), an additional correction was made to more

accurately quantify heterochiasmy. Specifically, per individual and per phased haplotype within

individual the number of crossovers within 50 cM of each crossover were counted and crossovers were

only considered when the number of crossovers was odd, suggesting that a true phase change occurred. If

the number was even, these crossovers were not counted as they probably reflect a genotyping error since

crossover interference is expected within the salmonids and double crossovers therefore should not occur.

This is similar to the approach used by Johnston *et al.* (2016) to avoid false double crossovers by only

including crossovers that flank more than a single marker. Subsequently, the cumulative number of

crossovers for fused metacentric and acrocentric chromosome were calculated and cumulatively displayed

in positions as a percentage of the total chromosome length. The corrected crossover counts were used to

calculate the female:male recombination rates of the parents. This was also conducted without cumulating

and displayed on a per chromosome per haplotype basis.

**QTL analysis**

The effect of sex on each phenotype was tested using linear models in R (R Development Core Team 2017). If a marginal effect of sex was found (p ≤ 0.20), sex was included in the model as a covariate for the phenotype to reduce residual variation and improve power to identify the QTL (Broman and Sen 2009). The R/qtl function *scanone* with permutation testing (10,000 permutations; p ≤ 0.05) was used to identify the presence of a single QTL within each linkage group (Broman *et al.* 2003). Chromosome-wide significance was tested in the same way but per chromosome (10,000 permutations; p ≤ 0.01). Confidence interval estimates (95%) for QTL positions were identified using *summary.scanone* calculating LOD support intervals with a 1.5 LOD drop. Sex-specific phenotypes (i.e. sperm diameter and concentration, egg diameter) were tested in only one sex, and therefore had smaller sample sizes. Percent variance explained by the identified QTL was performed using *makeqtl* and *fitqtl* within R/qtl, including all genome and chromosome-wide QTL per trait in the formula (trait ~ $QTL_1 + QTL_2 + QTL_n$), as well as sex as a covariate when required. Phenotypic effects were estimated by calculating the differences between the mean phenotype values among the genotype groups for the marker closest to the identified QTL, including only individuals that were successfully genotyped. For markers that only segregate in one parent (i.e. *nn* x *np*) only two phenotype by genotype averages are given, one for the homozygote and one for the heterozygote offspring. Alternatively, for markers segregating in both parents (i.e. *hk* x *hk* or *ef* x *eg*), three phenotype averages are given, two for the alternate homozygotes and one for the heterozygote in *hk* x *hk* marker types and two for the alternate heterozygotes and one for the homozygote in *ef* x *eg* marker types. Sex-specific averages were calculated when the QTL required sex as a covariate in the model. RAD tags for all alleles and associated QTL results are available in File S2.

To identify the sex chromosome, offspring sex was coded as a binary trait to identify linkage to any of the LG by QTL mapping as described above (Broman and Sen 2009). Furthermore, the effect of a QTL may vary depending on the sex of an individual in a non-additive manner (Broman and Sen 2009) due to genetic variation in sexual dimorphism for the trait (e.g. loci that have a different effect in males and females Mackay 2001). Therefore, QTL by sex interaction effects were inspected per trait by subtracting an additive model (*genotype* and *sex*) from a full model (*genotype*, *sex* and a *sex-by-genotype* interaction term) as described by Broman and Sen (2009) . If the additive model is largely driving the effect, the model with only the interaction effect will not be significant. Significant interaction effects were only considered when the full model was found to be significant (Broman and Sen 2009).

Identities of sex chromosomes of other species were obtained from references listed in Table 1. For Atlantic Salmon, Artieri *et al.* (2006) identify that the sex determining region is on the long (q) arm of chromosome Ssa02, and Lien *et al.* identify that Ssa02q is homeologous to Ssa12q, indicating that the chromosome arm holding the sex determining region corresponds to the ancestral chromosome 9.1

(Sutherland *et al.* 2016). Other species were directly obtained from references in Table 1. Correspondence

between Arctic Charr *S. alpinus* and Brook Charr were identified indirectly through other species shared

between Nugent *et al.* (2016) and Sutherland *et al.* (2016).

**Data availability**

The raw data for this study is available in the NCBI SRA in BioProject PRJNA308100 and accession

SRP068206. All input files used for the analysis are in the supplementary files (File S1) and all code used

to perform analyses is available on Github at the following link:

https://github.com/bensutherland/sfon_pqtl/

253                                    **RESULTS**

**Sex-linked chromosome in Brook Charr**

Sex was highly associated with the majority of Brook Charr (BC) linkage group (LG) BC35, indicating

that this is the sex-linked chromosome in Brook Charr (Figure 1). Linkage across the entire LG until the

LOD score decreases at the distal end can be explained by male salmonid-specific low recombination rate

and male bias of crossovers towards telomeric regions (Sakamoto *et al.* 2000). The drop in LOD suggests

the far end of the chromosome is pseudoautosomal, which even occurs in the highly differentiated

mammalian X/Y chromosomes in a recombinogenic distal region of the Y chromosome (Lahn *et al.* 2001).

Many of the recombination events in BC35 were at a similar section of the LG (~90-110 cM; Figure S2B).

BC35 is an acrocentric chromosome homologous to the Northern Pike chromosome 15.1

(homeolog naming from Sutherland *et al.* 2016). There are no other sex chromosomes in the other

salmonid species with high-density genetic maps available with the chromosome arm containing the sex

determining region homologous to BC35 (Table 1). Arctic Charr has a sex chromosome that is comprised

of a triple fused chromosome (although this may vary across populations) that contains 15.1 in the fusion,

but in Arctic Charr this is not the chromosome arm that holds the sex determining region, which is held

within the arm on the other side of the chromosome AC04p (Nugent *et al.* 2016). Other salmonids have

different sex chromosomes, as shown in Table 1, including Lake Whitefish (3.1; Gagnaire *et al.* 2013),

Atlantic Salmon (9.1-20.2; Artieri *et al.* 2006; Lien *et al.* 2011), Rainbow Trout (14.2; Palti *et al.* 2015),

Coho Salmon *O. kisutch* (3.1; Phillips *et al.* 2005; Kodama *et al.* 2014), Chinook Salmon *O. tshawytscha*

(23.2; Phillips *et al.* 2005; Naish *et al.* 2013; Brieuc *et al.* 2014) or Sockeye Salmon (3.1-19.1; Larson *et*

*al.* 2016). This further refines previous observations of the general lack of homology in the sex

chromosomes of the salmonids (Woram *et al.* 2003). Some information on sex chromosomes identities

across *Salmo, Salvelinus* and *Oncorhynchus* have been previously reported (Phillips 2013) and most of

the results correspond with those here, with the exception of the Brook Charr sex chromosome, which the two studies identify as corresponding to opposite arms of the Arctic charr sex chromosome. This is possibly due to a population polymorphism, but more work would be needed to confirm this.

Considering the importance of inversions to sex chromosome formation through the reduction of recombination between X and Y (Lahn *et al.* 2001; van Doorn and Kirkpatrick 2007; Berset-Brandli *et al.* 2008), it is interesting to note that Brook Charr has a species-specific inversion in BC35 in the female map (15.1; see Figure 5 in Sutherland *et al.* 2016). As is usual for salmonid linkage maps, the male-specific map was not produced as the low recombination frequency resulted in poorly placed male-specific markers (Sutherland *et al.* 2016), and so it is not possible to check whether this inversion is heterozygous within the species, but this will be valuable to investigate in future studies.

**Figure 1.** The acrocentric linkage group BC35 is highly associated with sex in Brook Charr. Due to low recombination in males, high linkage is viewed across the majority of the linkage group.

**Sex-specific recombination rate and positions of crossovers**

Crossovers occurred 2.7-fold more often in the maternal haplotypes (total = 3679) than in the paternal (total = 1368; Figure 2) based on the phased haplotypes of 169 individual offspring (Wu *et al.* 2002; Sutherland *et al.* 2016). The double recombinant correction (see Methods) in the autosomes removed 606 and 682 crossover events due to probable genotyping errors from the dam and sire, respectively, providing a more accurate estimation of the heterochiasmy ratio, although the trends regarding the crossover positions remained similar. Crossovers were biased towards the center of the linkage groups in the dam and towards the external 20% of the linkage groups in the sire (Figure 2). This bias is similar to that observed in Rainbow Trout (Sakamoto *et al.* 2000) although reasons for it remain unknown.

**Table 1.** Salmonid sex chromosomes from high-density genetic maps named with Northern Pike designations (ancestral). The chromosome arm that contains the sex determining region is underlined, and the fusion status of the chromosome and original reference are provided. Ancestral chromosomes are defined by Sutherland *et al*. (2016) and are based on Northern Pike chromosomes from Rondeau *et al*. (2014).

| Common name | Scientific name | Linkage group (sex) | Ancestral | Fused (F) or acrocentric (A) | Evidence type | Citations |
|---|---|---|---|---|---|---|
| Lake Whitefish | *Coregonus clupeaformis* | LW25 | <u>3.1</u> | A | Linkage | (Gagnaire *et al.* 2013) |
| Atlantic Salmon | *Salmo salar* | Ssa02 | <u>9.1</u>-20.2 | F | FISH & Linkage | (Artieri *et al.* 2006; Phillips *et al.* 2009; Lien *et al.* 2011) |
| Arctic Charr | *Salvelinus alpinus* | AC04 | <u>1.2</u>-19.1-15.1 | F | Linkage | (Nugent *et al.* 2016) |
| Brook Charr | *Salvelinus fontinalis* | BC35 | <u>15.1</u> | A | Linkage | (Sutherland *et al.* 2016) and *current paper* |
| Rainbow Trout | *Oncorhynchus mykiss* | OmySex (29) | <u>14.2</u> | A | FISH & Linkage | (Phillips *et al.* 2006; Rexroad *et al.* 2008; Palti *et al.* 2015) |
| Coho Salmon | *O. kisutch* | Co30 | <u>3.1</u> | A | FISH & Linkage | (Phillips *et al.* 2005; Kodama *et al.* 2014) |
| Chinook Salmon | *O. tshawytscha* | Ots17 | <u>23.2</u> | A | FISH & Linkage | (Phillips *et al.* 2005; Naish *et al.* 2013; Brieuc *et al.* 2014) |
| Sockeye Salmon | *O. nerka* | So09 | <u>3.1</u>-19.1* | A | Linkage | (Larson *et al.* 2016) |

Separating chromosomes into fused metacentric (n = 8) and acrocentric chromosomes (n = 34) indicated a higher heterochiasmy ratio in fused metacentric than acrocentric chromosomes (5.6-fold and 2.2-fold, respectively). The male had fewer crossovers per chromosome in the fused metacentrics (mean = 26.5) than acrocentrics (mean = 32.9), even though fused metacentrics are comprised of two acrocentric chromosomes combined and thus are longer. In contrast, the female had approximately twice as many crossovers per fused metacentric chromosome (mean = 148.9) than acrocentric (mean = 70.5). The lower recombination in the paternal fused metacentrics than the paternal acrocentrics is probably due to missing regions of the genetic map that are residually tetraploid that were removed during marker filtering due to quality filtering, as the map was produced using a diploid cross (see Limborg *et al.* 2016). Inspection of

individual chromosomes indicates that the chromosomes expected to still exhibit residual tetraploidy
(Sutherland *et al.* 2016) all show an absence of crossovers in the male relative to the chromosomes
expected to have returned to a diploid state (Figure S2). The missing regions in the 16 (of 50)
chromosome arms expected to be residually tetraploid will result in an inflation of the heterochiasmy ratio,
as male crossovers will be specifically underestimated for these arms. Regardless, most chromosomes are
not residually tetraploid and heterochiasmy can be viewed in these other chromosomes. In summary, the
male has fewer recombination events than the female and the crossovers are biased to the distal portions
of the chromosomes.

The identified sex chromosome had more crossovers than the average male acrocentric
chromosomes (sex = 71, other acrocentrics average = 32.9), and all of the crossover events in the sex
chromosome occurred at one end of the chromosome and almost no crossovers occurred in the rest of the
chromosome (Figure S2). This bias to only a single end of each chromosome in the male map was
consistent throughout all of the chromosomes with crossovers present. Using the positions of centromeres
determined for Chinook Salmon (Brieuc *et al.* 2014), and placing them in the corresponding position on
the Brook Charr map using map correspondence (Sutherland *et al.* 2016), indicates that the end of the
acrocentric chromosomes where the crossovers occur is the opposite end to that containing the probable
centromere (see Figure S2).

**QTL identification: growth, reproduction and stress response**
Genome-wide significant QTL were identified for weight, length, condition factor, specific growth rate,
and liver weight (Table 2; Table S2). A total of 29 QTL were found to be significant at the chromosome-
wide level ($p \leq 0.01$), and these included QTL for phenotypes egg and sperm diameter, change in cortisol,
chloride and osmolality after an acute handling stress, *growth hormone receptor* gene expression and
hematocrit (Table 2). In total, QTL were identified on 14 of the 42 Brook Charr linkage groups (Figure 3).

Several traits showed sexual dimorphism and therefore required sex as a model covariate. These
included weight, length, liver weight, hepatosomatic index, hematocrit, change in osmolality and cortisol
from stressor, resting plasma chloride, hepatic glycogen, *insulin-like growth factor 1* and *igf receptor 1*
(Table S1). Specific growth rate, condition factor, change in chloride, resting plasma osmolality and
glucose, and *growth hormone receptor* gene expression did not show sexual dimorphism. Traits with high
phenotypic correlation included length and weight (r = 0.90 at T1), and liver weight and hepatosomatic
index (r = 0.85; Figure S1). Specific growth rate T1-T2 was negatively correlated with weight at T1 (r = -
0.64), suggesting that larger individuals measured at T1 subsequently grew slower than smaller
individuals. Other traits generally were not as highly correlated (r < 0.35). Even though the phenotypes
were not highly correlated, QTL were identified for condition factor and weight in the same region of

    BC20, and QTL affecting hematocrit and weight (r = 0.24) were found in the same region on BC04

(Figure 3).

**Figure 2**. Maternal and paternal cumulative crossover positions across the chromosomes. The position of
each crossover is expressed as a percent of the total crossover length and cumulated for all crossovers
within each chromosome type, specifically fused metacentric chromosomes (A,C) and acrocentric
chromosomes (B,D) in the maternal and paternal haplotypes, respectively. Maternal haplotypes had 2.7-
fold more crossovers than paternal haplotypes, with the maternal crossovers occurring throughout the
chromosome and the paternal crossovers restricted mainly to the first and/or last 20% of the linkage
groups.

A few specific trait-linkage group combinations had elevated LOD across a large portion of the

LG. This was observed for length and weight on BC03, BC04 and BC05. To determine if this was due to

one specific marker type, the three marker types were each tested independently for QTL (female-specific

*nn* x *np*; informative in both parents *ef* x *eg*; and semi-informative *hk* x *hk*). Interestingly, when including

markers only polymorphic in the female (i.e. *nn* x *np*) these elevated LOD baselines were not observed

(*data not shown*). It is possible that this may be therefore due to an effect originating from paternal alleles,

which have strong linkage across the entire LG. As these QTL explained a substantial amount of variance

for these three traits, the full analysis includes these markers, but the exact locations within the LG of

these QTL cannot be determined without additional families or crosses (markers noted in Figure 3).

A substantial amount of trait variance for length at T2 within this mapping family was explained

by five QTL. Together with the additive sex covariate, this collectively explained 54.5% of the trait

variation. The QTL with broad elevated LOD on BC04 and BC05 (see above) individually explained 5.5

and 8.2% of the variation, respectively. QTL for condition factor varied depending on the sampling time

(T1-T3), where each time point had a QTL at a different LG that explained over 10% of the trait variation

within the time point. However, trait variation was small for this trait and therefore effect sizes of the

QTL were also small (Table 2). QTL for specific growth rate (SGR) were identified on BC03, BC09 and

BC36, with three QTL explaining 26% of the SGR (T2-T3) trait variation (Table 2).

**Figure 3.** All identified QTL plotted on the Brook Charr genetic map. QTL for growth related traits are
shown in blue, reproductive in black, and blood or stress-related in red. QTL with asterisks are at the
genome-wide significance level, and the rest are chromosome-wide. QTL with broad confidence intervals
discussed in the Results are denoted with a positive symbol (+). More details on phenotypes can be found
in Table S1 and on QTL can be found in Table 2 and Table S2.

Reproductive traits were sex-specific and therefore had approximately half of the individuals as
the traits with values in both sexes, and thus had less statistical power, reducing the ability to detect QTL.
Nonetheless, enough power was present in the data to detect chromosome-wide significant QTL for egg
and sperm diameter (Table 2; Figure 3). A QTL for egg diameter was identified at 67 cM of BC05,
explaining 39% of the trait variation and a QTL for sperm diameter was identified at 48 cM of BC24,
explaining 33% of the trait variation. Neither of these reproductive-related traits mapped to the sex-linked
chromosome (BC35). A QTL for *growth hormone receptor* (*ghr*) gene expression was identified at 138
cM of BC24 explaining 24% of the trait variation.

Stress response QTL were identified only at the chromosome-wide significance level. This
included responses of cortisol (114 cM of BC06), chloride (43 cM of BC38), and osmolality (60 cM on
BC40; Table 2; Figure 3). Change in cortisol from acute handling stress was highly dependent on sex; the
identified QTL explained 9% of the trait variation whereas sex explained 43% (total PVE = 58%; Table
2). Females heterozygous at the marker closest to the QTL increased cortisol by 2.3 µg/dL plasma more
than the homozygote, and heterozygous males had 0.65 µg/dL lower than the homozygote. To further
demonstrate the large sex effect, averaging the two genotypes shows that females increased blood cortisol
by 8.6 µg/dL whereas males only increase by 0.43 µg/dL (i.e. 20-fold higher cortisol response in females).
Osmolality change (mmol/kg) was also affected by sex but to a lesser extent than was cortisol. More
specifically, the identified QTL explained 14% of the trait variance and sex explained 14.6%. For this
QTL, both sexes showed suggestive additive effects with the heterozygote having a value in between the
two homozygotes (Table S2). Chloride change (mmol/L) was not sex-dependent, and the identified QTL
on BC38 explained 18% of the trait variation. Chloride reduced in the heterozygote individuals by 2.58
mmol/L whereas it stayed approximately the same in the homozygote (0.3 mmol/L; Table 2). Resting
blood hematocrit was affected by sex, and two QTL were identified at the chromosome-wide level (23
cM on BC04 and 14 cM on BC25). Together with the sex covariate, these explained 43% of the trait
variation, with each QTL explaining approximately 12%. Together, these results indicate the importance
of including sex as covariate in these models. These markers will provide targets for selective breeding.

**Table 2.** Identified QTL in Brook Charr with positions, percent variance explained (PVE) and the effect
of the allelic state on the trait. Sex was included as a covariate when required, and in these cases the
allelic effect is given for both males and females, and the PVE from sex is also given. When sex was not
required as a covariate, the second averages are displayed as NA and the first averages represent both
sexes. The phenotype average for the homozygote common allele (aa avg) is shown for comparison to the
largest effect size (effect ♀ or ♂). LG = linkage group; Pos = cM position; CI = confidence interval; PVE
= percent variance explained; sex.cov = sex included as a covariate. QTL significance is displayed as
genome wide p ≤ 0.01 ***; p ≤ 0.05 **; or chromosome-wide p ≤ 0.01 *.

| Phenotype | LG | Pos | 95% CI | Marker | QTL pval | Tot. PVE | Ind. PVE | aa avg ♀ | effect ♀ | aa avg ♂ | effect ♂ |
|---|---|---|---|---|---|---|---|---|---|---|---|
| Weight (g) T2 | 4 | 28.3 | 16-214 | 7187 | * | 42.8 | 9.5 | 126 | +22.1 | 129.5 | +52.1 |
| | 5 | 198 | 39-262 | 125487 | * | | 8.1 | 121.4 | +27.9 | 145.3 | +30.6 |
| | 20 | 162 | 105-175 | 6352 | * | | 6.2 | 133.9 | -2.2 | 171.2 | -22.8 |
| | | | | sex.cov | | | 9.5 | | | | |
| Length (cm) T2 | 3 | 83 | 19-267 | 90770 | * | 54.5 | 2.0 | 22.4 | -0.3 | 23.6 | +0.4 |
| | 4 | 115 | 16-215 | 66075 | ** | | 5.5 | 21.8 | +0.9 | 23.6 | +0.2 |
| | 5 | 261 | 185-262 | 85980 | *** | | 8.2 | 21.7 | +1.1 | 23.6 | +0.3 |
| | 20 | 169 | 99-175 | 60142 | * | | 4.1 | 21.7 | +0.8 | 22.9 | +1.3 |
| | 34 | 112 | 85-132 | 120757 | * | | 3.1 | 22.5 | -1.1 | 24.0 | +0.7 |
| | | | | sex.cov | | | 13.5 | | | | |
| Cond. Fact. T1 | 16 | 89.9 | 48-105 | 118085 | * | 10.0 | 10.0 | 1.0 | -0.02 | NA | NA |
| Cond. Fact. T2 | 39 | 46.9 | 35-83 | 39977 | * | 10.3 | 10.3 | 1.2 | -0.03 | NA | NA |
| Cond. Fact. T3 | 20 | 150 | 116-175 | 55565 | ** | 12.2 | 12.2 | 1.1 | +0.04 | NA | NA |
| SGR T2-T3 | 3 | 35.7 | 18-85 | 115199 | * | 26.0 | 6.8 | 0.6 | +0.10 | NA | NA |
| | 9 | 139 | 89-189 | 128240 | * | | 5.3 | 0.6 | +0.04 | NA | NA |
| | 36 | 31.4 | 1-80 | 30493 | *** | | 5.3 | 0.6 | -0.09 | NA | NA |
| Egg diameter | 5 | 66.8 | 41.7-185 | 37572 | * | 39.0 | 38.95 | 4.0 | -0.046 | NA | NA |
| Sperm diameter | 24 | 47.8 | 0-59.8 | 202134 | * | 32.6 | 32.59 | NA | NA | 2.9 | -0.001 |
| Δ cortisol | 6 | 114 | 108-135 | 113752 | * | 57.6 | 9.0 | 7.4 | +2.3 | 0.8 | -0.6 |
| | | | | sex.cov | | | 43.0 | | | | |
| Δ chloride | 38 | 43.3 | 25.3-60.9 | 116693 | * | 18.5 | 18.5 | 0.3 | -2.9 | NA | NA |
| Δ osmolality | 40 | 60.3 | 26.3-82.1 | 52306 | * | 31.1 | 14.0 | 15.2 | -3.4 | -5.4 | +4.6 |
| | | | | sex.cov | | | 14.6 | | | | |
| *ghr* | 24 | 138 | 47.8-153 | 141355 | * | 23.8 | 23.8 | -3.9 | +0.2 | NA | NA |
| Hematocrit | 4 | 22.6 | 16.3-161 | 105237 | * | 42.8 | 12.0 | 35.3 | +0.9 | 39.6 | -1.7 |
| | 25 | 139 | 113-159 | 1153 | * | | 12.4 | 35.1 | +0.5 | 38.3 | -0.2 |
| | | | | sex.cov | | | 13.9 | | | | |

**DISCUSSION**

Salmonids are a model system for studying the effects of whole genome duplication (WGD) on genome

evolution, sex determination and speciation. Specifically, the evolution of sex determination after WGD

and its interaction with heterochiasmy remain active areas of research. In this study we identify the sex-

linked chromosome and strong heterochiasmy using the high-density genetic map of Brook Charr. Female

recombination rates were 2.7-fold higher than those in the male, and male recombination was highly

biased to chromosome ends. Using recently established correspondence among salmonid chromosomes,

we show that the Brook Charr sex chromosome is not the same chromosome arm linked to sex in any

other species characterized with high-density genetic maps. However, this chromosome arm (ancestral

15.1) is contained in a fusion within a triple chromosome fused sex chromosome of the congener Arctic

Charr. In other salmonid species, some consistencies in sex chromosomes can be viewed, even as distant

as Lake Whitefish and members of the genus *Oncorhynchus* (discussed below). We additionally evaluate

linkage to 29 reproductive, stress and growth phenotypes (21 not including phenotypes measured at

multiple time points) and identify 29 genome- or chromosome-wide QTL on 14 of 42 linkage groups and

compare these to known QTL in other salmonids. This work provides markers for selective breeding of

Brook Charr as well as insight into the role of heterochiasmy in sex determination and genome evolution

in a post-WGD salmonid genome.

**Sex determination post whole genome duplication**

In species with genetic sex determination, WGD generates multiple copies of sex chromosomes and this

may present challenges to the new lineage (Davidson *et al.* 2009) such as unbalanced gametes and

independent segregation of sex chromosomes (Muller 1925) or disruptions in dosage balance in species

with heteromorphic sex chromosomes (Orr 1990). However, polyploidization also poses other challenges

independent to the duplicated sex chromosome system (Mable 2004; Otto 2007). Nonetheless, ancestral

polyploidization resulting in paleopolyploid lineages has occurred throughout plant and animal evolution,

with some notable examples in vertebrates including the teleost-specific duplication (3R; Taylor *et al.*

2003), the salmonid-specific duplication (4R; Allendorf and Thorgaard 1984), and an allopolyploidization

within the *Xenopus* genus (Session *et al.* 2016). Debate still exists on the effect of polyploidization on

diversification (e.g. Clarke *et al.* 2016). Diversification may involve sex determination systems, as

paleopolyploids may develop a wide variety of sex determination systems, for example in the teleosts

(Mank and Avise 2009). A wide variety of sex chromosomes are used in different teleost groups including

stickleback species (Ross *et al.* 2009; Kirkpatrick 2016) and Medaka, in which *Oryzias latipes* sex

chromosome is LG1 with a probable translocation of the sex gene to a neo-sex chromosome (Kondo 2006)

whereas other members such as those of the Medaka *celebensis* group initially used LG24 as the sex

chromosome and then transitioned to LG10 in several other species (Myosho *et al.* 2015). Ancestral
allotetraploids can also develop new sex determination systems. For example, the African clawed frog
*Xenopus laevis* (ZZ/ZW), has a probable translocated W-specific region that is out of synteny with the
sister group to the polyploidization and with the homeolog of the sex chromosome (Session *et al.* 2016).
However, variable sex determination systems are not unique to paleopolyploid lineages. For example, *X.*
*tropicalis* did not undergo the allotetraploid event but has a unique sex determination system that involves
W, Z, and Y chromosomes, where the presence of a single Y can override the feminizing presence of two
W chromosomes in triploids (Roco *et al.* 2015). The extent of the involvement of polyploidization on the
lability of sex determination systems remains to be determined.

Taxa with high rates of turnover in sex chromosomes have indicated that some chromosomes are
more likely to become sex chromosomes. This is possibly due to favourable gene content, for example
when an autosome contains sexually antagonistic genes it can be repeatedly selected to become a sex
chromosome (Marshall Graves and Peichel 2010). A comparative analysis of teleosts indicates repeated
independent evolution of the same chromosomes as sex chromosomes throughout evolutionary history
(see Table 2 in Marshall Graves and Peichel 2010). Therefore it is not only the master sex determining
gene that can be repeatedly utilized by evolution, but also certain chromosomes due to the gene content,
which can occur over large evolutionary distance. For example, the teleost tongue sole *Cynoglossus*
*semilaevis* and the chicken *Gallus gallus* (both ZZ/ZW) independently evolved sex chromosomes in
homologous chromosomes (Chen *et al.* 2014). Similarly, species from three anuran genera that have
diverged for over 210 million years (*Bufo*, *Hyla* and *Rana* spp.) all have sex-linked markers that map to
the same *X. tropicalis* chromosome containing *doublesex and mab-3 related transcription factor 1*
(DMRT1) and a large region homologous to the avian sex chromosome (Brelsford *et al.* 2013), which the
authors suggest is due to independent evolution to the same sex chromosome across the different genera.
The platypus *Ornithorhynchus anatinus* has five Y and five X chromosomes, all of which are independent
but form a chain at meiosis to co-segregate all together into sperm; this system connects the two sex
determination types, with the most degenerate sex chromosome as homologous to the Z chromosome of
birds and the least degenerate as that homologous to the X chromosome of mammals (Grützner *et al.*
2004; Charlesworth and Charlesworth 2005). Finally, although at least three non-homologous sex
chromosomes exist within *Xenopus*, the sex determining region of *X. borealis* shares orthologous genes to
mammals including humans (Furman and Evans 2016). In summary, repeated, independent evolution of
the same sex chromosome or use of the same set of specific genes for sex determination therefore has
been documented across a variety of animal taxa.

Salmonids have genetically controlled sex determination with XX/XY systems (Thorgaard 1977;
Davidson *et al.* 2009), but putative translocation of the sex determining gene to different autosomes has

resulted in many different sex chromosomes in the different lineages (Woram *et al.* 2003) and even within
the same species (Küttner *et al.* 2011; Eisbrenner *et al.* 2013). However, comparing across the phylogeny
indicates some noteworthy consistencies. First, several species use 3.1 as the sex chromosome, or have
this chromosome fused with the sex chromosome, including the neo-Y of Sockeye Salmon and the sex
chromosome of Coho Salmon (Faber-Hammond *et al.* 2012), as well as the sex-linked LG in Lake
Whitefish (Gagnaire *et al.* 2013), identified as 3.1 during map comparisons (Sutherland *et al.* 2016).
Relative to the variability seen in sex chromosomes in the salmonids, this is a striking consistency
considering that these species have diverged for approximately 50 million years (Crête-Lafrenière *et al.*
2012). This consistency may indicate that either a) 3.1 is an ancestral sex chromosome in the salmonids;
or b) the different species converged on this chromosome independently as it contains a gene complement
that is highly beneficial to be present as a sex chromosome (Marshall Graves and Peichel 2010; Chen *et al.*
2014; Furman and Evans 2016). Second, the Brook Charr sex chromosome (15.1) is fused within the sex
chromosome of Arctic Charr, but is not the same arm as that containing the sex marker in Arctic Charr
(Nugent *et al.* 2016), indicating one of these is a neo-sex chromosome. Furthermore, the middle
chromosome arm fused in the Arctic Charr triple chromosome fusion is 19.1, which is the neo-Y of
Sockeye Salmon (Table 1). These observations provide further evidence for the fusion of specific
chromosomes together that are beneficial for maintaining within the sex chromosome environment.
Finally, intraspecific polymorphism in sex chromosomes occurs in Arctic Charr (Moghadam *et al.* 2007;
Küttner *et al.* 2011), and Icelandic Arctic Charr were identified as having a sex chromosome as one of the
two homeologs AC01 or AC21 instead of AC04, and state that this is homologous to the sex chromosome
of Atlantic Salmon Ssa02 (Küttner *et al.* 2011), which is 9.1 (Sutherland et al 2016), again indicating the
potential re-use of the same chromosome as the sex chromosome.

The presence of both the Y chromosome of Brook Charr and the neo-Y of Sockeye Salmon as
putative neo-Y chromosomes of Arctic Charr is worth further investigation because neo-Y chromosomes
can influence phenotypic divergence and reproductive isolation, as observed in sympatric Threespine
Stickleback *Gasterosteus aculeatus* populations for male courtship displays and hybrid instability,
respectively (Kitano *et al.* 2009; Kitano and Peichel 2011). These consistencies across a phylogeny can
provide insight into speciation. For example, the Threespine Stickleback and Ninespine Stickleback
*Pungitius pungitius* have two different sex chromosomes (LG19 and LG12, respectively), and the
Blackspotted Stickleback *G. wheatlandi* has a fused Y-chromosome made up of these two linkage groups
(Ross *et al.* 2009), to which the authors suggest multiple independent recruitment of LG12 as the sex or
neo-Y chromosome. Other sticklebacks have different sex chromosomes, such as the Brook Stickleback
*Culaea inconstans*, and the Fourspine Stickleback *Apeltes quadracus*, which has a ZW sex determination
system (Ross *et al.* 2009). The salmonid variety and consistencies identified here provide another group

for analyzing sex chromosome differences in relation to gene content and speciation, and in salmonids

also occurs with the salmonid-specific WGD. As more salmonid genomes are characterized, it will

become clearer whether certain sex chromosomes are ancestral or have independently evolved, and

whether there is a favourable gene content within often-viewed sex chromosomes.

In the context of sex chromosome fusions and residual tetraploidy, several additional observations

on the nature of salmonid sex chromosomes can be made from the present analysis (four genera; eight

species; Table 1). First, sex chromosomes with fusions only occur in species specific fusions in the data

here; the three fused sex chromosomes (in Atlantic Salmon, Arctic Charr, and Sockeye Salmon) are

products of species-specific fusions and not in conserved fusions (Sutherland *et al.* 2016). Arctic Charr *S.*

*alpinus* has a sex chromosome that in some individuals involved three fused chromosomes (Nugent *et al.*

2016), and all available evidence suggests this is a species-specific fusion given that these fusions are not

present in the more basally diverging Atlantic Salmon nor the congener Brook Charr (Sutherland *et al.*

2016). Y fusions are the most common of sex chromosome fusions (Pennell *et al.* 2015) and can permit

other sexually antagonistic genes to be linked to the non-recombining regions (Charlesworth and

Charlesworth 1980; Charlesworth *et al.* 2005). Y fusions may also occur due to drift with only slightly

deleterious effects (Kirkpatrick 2016), as males have increased fusion prevalence in general and increased

repeat content (and thus fusion potential) in degenerating Y (Pennell *et al.* 2015). However, since the same

chromosomes that are involved in Y fusions in some species are the sex chromosomes in others (e.g. 15.1

or 19.1, discussed above), it suggests that these fusions could have an adaptive advantage, such as the

movement of an autosome with alleles under sexually antagonistic selection into the Y chromosome

environment, as discussed by Charlesworth and Charlesworth (1980) and Kirkpatrick (2016). When

recombination is low in males (i.e. heterochiasmy), this Y fusion holds an additional chromosome in a

constantly lower recombination environment as it will always be present in males. The use of the same

chromosomes as sex chromosomes and as fusion partners within the salmonids merits further study.

Secondly, chromosomes with regions of residual tetraploidy can become sex chromosomes; two of the

seven identified sex chromosomes (Chinook Ots17 (23.2) and Atlantic Salmon Ssa02q (9.1)) are

chromosomes known to exhibit residual tetraploidy (Brieuc *et al.* 2014; Allendorf *et al.* 2015; Sutherland

*et al.* 2016; Lien *et al.* 2016), therefore suggesting that exhibiting residual tetraploidy does not prevent a

chromosome from becoming a sex chromosome.

Translocation of a sex determining gene to an autosome and the adoption of the autosome as a

new sex chromosome may be possible if the gene moves into linkage with a locus that is under sexually

antagonistic selection (van Doorn and Kirkpatrick 2007). The probability of this adoption is increased

with the occurrence of an inversion in the region which will increase linkage by reducing recombination

(van Doorn and Kirkpatrick 2007), but it requires that the benefit of the new chromosome is greater than

that existing on the original sex chromosome. Interestingly, in the unique sex chromosome of Brook Charr
(15.1), there is a large inversion in relation to the other salmonids (Sutherland *et al.* 2016). However, this
is an interspecific inversion and has not yet been determined whether it is also heterozygous within the
species due to low recombination and resultant challenges of generating male maps. To further
characterize this, genome sequence for both the X and Y chromosomes of Brook Charr will be valuable.

The salmonids, being at an early stage of sex chromosome evolution (Phillips and Ihssen 1985)
provide a good system to study sex chromosome evolution (van Doorn and Kirkpatrick 2007). As we
observed here, reduced recombination occurs consistently in male salmonids, being restricted to the
telomeric region opposite the centromere, resulting in a lack of recombination between X and Y in the
middle of the chromosome. This may facilitate sex chromosome formation, with tight linkage developing
across the entire Y chromosome (Haldane 1922; Nei 1969; Lenormand 2003). Heterochiasmy is not only
restricted to the sex chromosome, but rather occurs throughout the genome, as has been viewed in several
systems with developing sex chromosomes, such as the European tree frog *Hyla arborea* (Berset-Brandli
*et al.* 2008), Medaka (see Kondo *et al.* 2001; Kondo 2006), zebrafish *Danio rerio* (Singer *et al.* 2002), and
the salmonids of genera *Oncorhynchus* (Sakamoto *et al.* 2000), *Salmo* (Moen *et al.* 2004) and *Salvelinus*
(present study). However, heterochiasmy also occurs in systems with fully developed sex chromosomes,
such as humans, where females have ~1.6-fold higher rates than males, which recombine predominantly at
telomeric regions (Broman *et al.* 1998).

Y degeneration can occur from a lack of recombination in sex chromosomes (Charlesworth 1991;
Charlesworth *et al.* 2005), and this can also result in degeneration of fused neo-Y, when present. Neo-Y
degeneration has occurred rapidly in achiasmate male species such as *Drosophila miranda*, having
degenerated after only 1-2 My in the non-recombining state (Steinemann and Steinemann 1998;
Charlesworth and Charlesworth 2005). In species with heterochiasmy, even before large degeneration,
accumulated substitutions can occur throughout a neo-Y and increased sex-biased gene expression occurs
for genes within the neo-Y than the other autosomes, as observed in stickleback (Yoshida *et al.* 2014).
These changes are not only degenerative, migration to the Y, and preservation of male-beneficial genes on
the Y also occurs, as well as dosage compensation and migration of female-beneficial genes to the X
(Bachtrog 2006). Many changes can occur between X and Y when crossovers do not occur throughout the
chromosomes.

In the salmonids, sex chromosome turnover by *sdY* translocation may restart the process of Y
degeneration (Yano *et al.* 2012b). In species with heterochiasmy rather than achiasmy, occasional
crossover between X and Y would also reduce sex chromosome heteromorphism and Y degeneration. This
may be the reason for sex chromosomes remaining homomorphic in green toad species (*Bufo viridis*) all
which have the same sex chromosomes (Stöck *et al.* 2013), and in several members of the *Hyla* genus of

European tree frogs, which also all share the same sex chromosomes (Stöck *et al.* 2011). Regeneration of

Y chromosomes by occasional crossover is termed the 'fountain-of-youth' hypothesis, and is particularly

likely for species with the possibility of sex reversal, as recombination rate is based on phenotypic sex

rather than genetic sex (discussed in Perrin 2009). Some salmonid sex chromosomes are heteromorphic

(Davidson *et al.* 2009) and accumulate repeats (Devlin *et al.* 1998), this may suggest in some species this

regeneration is not occurring. Lack of recombination will be accentuated by inversion accumulation and

other differentiation between sex chromosomes reducing meiotic pairing and crossovers. Sex reversal is

possible in salmonids (Johnstone *et al.* 1978) and has been observed in the wild, for example in Chinook

Salmon (Nagler *et al.* 2001), but the greater extent of this occurring in nature in other salmonids is yet to

be determined. Relative effects of sex chromosome turnovers, occasional X/Y crossovers, and large sex

chromosomal polymorphisms merits further investigation for which the salmonids are a good model

system. The extent of Y or neo-Y degeneration, gene migration, or other aspects of sex chromosome

evolution have not yet been explored comparatively in the salmonids. As these aspects may differ among

species depending on the length of time the chromosome has been used as the Y chromosome, further

investigation into interspecific differences (e.g. 3.1 sex chromosome in both Lake Whitefish and members

of *Oncorhynchus*), or intraspecific differences between populations having different sex chromosomes

(Eisbrenner *et al.* 2013), will be valuable to determine the history of the sex chromosome evolution in the

salmonids.

**QTL mapping, hotspots and consistencies with other species**

Knowledge on the genetic architecture of important traits in the salmonids is improving, for example for

aquaculture-related traits such as disease resistance (Yáñez *et al.* 2014) and stress tolerance (Rexroad *et al.*

2012), and ecologically-relevant traits such as age-at-sea (Barson *et al.* 2015) and body shape evolution

(Laporte *et al.* 2015). In the present study we improve the understanding of genetic architecture of growth,

reproductive and stress-response traits by identifying QTL on 14 of the 42 LGs in the Brook Charr linkage

map (four fused metacentric and 10 acrocentric chromosomes). This improves the previous analysis of

these traits on a low-density map (Sauvage *et al.* 2012a, 2012b) and brings the QTL for these phenotypes

into the context of the more characterized high-density map generated here (e.g. with information on

correspondence of arms with other salmonids, probable residual tetraploidy and centromere positions,

ancestral chromosomes, and identified sex chromosome).

Although correlated phenotypes clustered on the map as expected (e.g. length, weight), no

clustering was observed for blood and stress-related parameters (i.e. hematocrit, change in cortisol,

chloride and osmolality), with each trait having a QTL on a different chromosome. As pleiotropy can

occur with both positive and negative genetic correlations between traits with common underlying biology

(Mackay *et al.* 2009). This is important to consider in marker-assisted selection, to identify QTL useful for

simultaneous selective breeding of multiple traits and to avoid negative correlations between desirable

traits (Lv *et al.* 2016). Mapping multiple correlated traits simultaneously can help define regions (Jiang

and Zeng 1995). However, it can be difficult to determine whether two traits are truly pleiotropic or

whether causal variants for each trait are in in tight linkage, especially when a QTL region is wide

(Mackay *et al.* 2009).

Consistencies in QTL across multiple species can be useful for identifying regions of the genome

with highly conserved roles. Several QTL hotspots have been identified within *Oncorhynchus*, specifically

for thermotolerance, length and weight on So6b (Hecht *et al.* 2012), So7a (except weight; also viewed in

Rainbow Trout and Chinook Salmon), and So11b (see Larson *et al.* 2015). The corresponding Brook

Charr LGs to So6 and So11b (Sutherland *et al.* 2016) did not contain any QTL in the present study, but

the corresponding LG to So7a (BC34) contains a length QTL (Table 2). This further implicates this

chromosome (ancestral 10.2) as having an evolutionary conserved influence on salmonid growth.

Weight and growth are expected to be highly polygenic traits, therefore requiring many

individuals to have sufficient power to identify loci of minor effect (Rockman 2012; Ashton *et al.* 2016).

For example, sample sizes of at least 500 individuals may be required to identify QTL accounting for less

than 5% of the total phenotypic variance (Mackay 2001). This means that often only large effect QTL are

identified, leading to the misconception that these are the norm and to an inflation of the actual percent

variance explained by the QTL (Beavis 1997; Xu 2003). High powered studies can identify more QTL,

such as a recent study in Atlantic Salmon with 1695 offspring and 20 sires, which identified four

chromosomes harboring major effect growth QTL (Tsai *et al.* 2015). Similarly, a study in Common Carp

*Cyprinus carpio* with 522 offspring and eight families identified 10 genome-wide and 28 chromosome-

wide significant QTL for three growth traits, with 30/50 chromosomes containing suggestive QTL (Lv *et*

*al.* 2016). Nonetheless, QTL can be detected with fewer individuals. For example, QTL for polygenic

traits growth rate, behavior and morphology were identified in Lake Whitefish with 102 individuals in the

mapping family (Gagnaire *et al.* 2013; Laporte *et al.* 2015). Furthermore in the present study we identified

QTL for many of the traits with 169 or fewer individuals. Since the effect of a QTL can differ in different

genetic backgrounds due to epistasis (Mackay 2001), it is therefore important to evaluate the effect of

markers in different crosses with different genetic backgrounds to better understand the broader use of the

marker (Lv *et al.* 2016).

The precision of mapping QTL within a family depends on recombination rate (Mackay 2001;

Mackay *et al.* 2009). Therefore the low number of crossovers in male salmonids will reduce the overall

precision of trait mapping. This effect of heterochiasmy has been used by recent salmonid studies to use a

two-stage approach by initially using a sire-based analysis with few markers per chromosome to identify

chromosomes of interest followed by a dam-based analysis to more finely resolve the QTL positions (Tsai

*et al.* 2015). Heterochiasmy is therefore important to consider when designing QTL experiments for
species exhibiting this trait. In the present study, several QTL with very broad regions of elevated LOD
were identified (e.g. for length on BC03, BC04, and BC05), which may be due to low recombination and
paternally associated haplotypes (see Results). In contrast, many of the other identified QTL in this study
have small confidence intervals and high percent variance explained, and therefore will be useful for
selective breeding (Table 2; Figure 3).

Although QTL mapping connects nucleotide sequence with trait variation, it generally ignores
intermediate phenotypes that can be very useful in determining underlying drivers of traits, and the use of
the expression levels of gene transcripts as traits to identify eQTL can provide information on the
intermediate steps to generate a phenotype (Mackay *et al.* 2009). Traits queried in eQTL experiments have
the additional information on gene location in the genome, providing information on cis or trans-eQTL
(Mackay *et al.* 2009). This will be an important next step in determining the underlying causes of the
genotype-phenotype interaction in Brook Charr.

672
**CONCLUSIONS**

The relationships between sex chromosomes, heterochiasmy and polyploidization have important
influences on genome architecture for key biological traits, but much remains unknown about these
interactions. Here we identify the sex-linked chromosome in Brook Charr and compare sex chromosome
identities across the salmonids to investigate consistencies. Although many different chromosomes are
used as sex chromosomes in salmonids, some consistencies can be identified, even in lineages that have
diverged for ~50 million years, *Coregonus* and *Oncorhynchus*. Sex chromosomes that are contained
within fused chromosomes thus far are only observed in species-specific fusions and not in conserved
fusions. Heterochiasmy, or differences in recombination rate between sexes, may play an important role in
the evolution of sex chromosomes. Heterochiasmy is viewed here in the *Salvelinus* genus, and in other
salmonid genera *Oncorhynchus* and *Salmo*, where male recombination is much lower than female, and
crossovers are restricted to telomeric regions. Inversions are also important for sex chromosome evolution,
and the Brook Charr sex chromosome from the female map exhibits a large interspecific inversion,
although the intraspecific polymorphism of this inversion has not yet been determined. Additional analysis
of salmonid genomes is needed to understand the effect of the mobile sex determining gene on phenomena
such as Y degeneration. To improve the characterization of important traits and potential for selective
breeding, we additionally identify 29 QTL across the genome for growth, reproduction, and stress-
response traits, several of which having high PVE and well-refined intervals. Hotspots for multiple traits
were not common, but we identify that an earlier identified hotspot in *Oncorhynchus* also contains a

length QTL in Brook Charr, further indicating the importance of this chromosome region and the value of

identifying orthologous QTL with comparative genomics.

**ACKNOWLEDGEMENTS**

This work was funded by a Fonds de Recherche du Québec Nature et Technologies (FRQNT) research

grant awarded to Céline Audet, Louis Bernatchez and Nadia Aubin-Horth, a grant from the Société de

Recherche et de Développement en Aquaculture Continentale (SORDAC) awarded to Louis Bernatchez

and Céline Audet, and a grant from the Spanish Ministry of Education (Grant PR2010-0601) awarded to

Ciro Rico. Thanks to G. Côté for laboratory assistance, M. Laporte for discussion on QTL and for

comments on the manuscript, M. Lamothe and T. Gosselin for discussion on double recombinants and

genotyping errors in RADseq data and to T. Gosselin for exporting the required files from STACKs for

QTL analysis. During this work, BJGS was supported by an NSERC postdoctoral fellowship, and then an

FRQS postdoctoral fellowship.

**REFERENCES**

Allendorf F. W., Thorgaard G. H., 1984 Tetraploidy and the evolution of salmonid fishes. In:Turner BJ
(Ed.), *Evolutionary genetics of fishes*, Plenum Publishing Corporation, New York, pp. 1–53.
Allendorf F. W., Bassham S., Cresko W. A., Limborg M. T., Seeb L. W., Seeb J. E., 2015 Effects of
crossovers between homeologs on inheritance and population genomics in polyploid-derived
salmonid fishes. J. Hered. **106**: 217–227.
Andrews K. R., Good J. M., Miller M. R., Luikart G., Hohenlohe P. A., 2016 Harnessing the power of
RADseq for ecological and evolutionary genomics. Nat Rev Genet **17**: 81–92.
Artieri C. G., Mitchell L. A., Ng S. H. S., Parisotto S. E., Danzmann R. G., Hoyheim B., Phillips R. B.,
Morasch M., Koop B. F., Davidson W. S., 2006 Identification of the sex-determining locus of
Atlantic Salmon (*Salmo salar*) on chromosome 2. Cytogenet Genome Res **112**: 152–159.
Ashton D. T., Ritchie P. A., Wellenreuther M., 2016 15 years of QTL studies in fish: challenges and
future directions. Mol. Ecol.
Bachtrog D., 2006 A dynamic view of sex chromosome evolution. Current Opinion in Genetics &
Development **16**: 578–585.
Barson N. J., Aykanat T., Hindar K., Baranski M., 2015 Sex-dependent dominance at a single locus
maintains variation in age at maturity in salmon. Nature **528**: 405–408.
Beavis W. D., 1997 QTL Analyses: Power, Precision, and Accuracy. In:Paterson AH (Ed.), *Molecular
Dissection of Complex Traits*, CRC Press, Boca Raton, pp. 145-159.
Berset-Brandli L., Jaquiery J., Broquet T., Ulrich Y., Perrin N., 2008 Extreme heterochiasmy and nascent
sex chromosomes in European tree frogs. Proceedings of the Royal Society B: Biological Sciences
**275**: 1577–1585.
Brandvain Y., Coop G., 2012 Scrambling eggs: Meiotic drive and the evolution of female recombination
rates. Genetics **190**: 709–723.
Brelsford A., Stöck M., Betto-Colliard C., Dubey S., Dufresnes C., Jourdan-Pineau H., Rodrigues N.,

Savary R., Sermier R., Perrin N., 2013 Homologous sex chromosomes in three deeply divergent
anuran species. Evolution **67**: 2434–2440.
Brieuc M. S. O., Brieuc M. S. O., Waters C. D., Waters C. D., Seeb J. E., Seeb J. E., Naish K. A., 2014 A
dense linkage map for Chinook Salmon (*Oncorhynchus tshawytscha*) reveals variable chromosomal
divergence after an ancestral whole genome duplication event. G3 - Genes|Genomes|Genetics **4**: 447–
460.
Broman K. W., Sen S., 2009 *A Guide to QTL Mapping with R/qtl*. Springer, New York.
Broman K. W., Murray J. C., Sheffield V. C., White R. L., Weber J. L., 1998 Comprehensive human
genetic maps: Individual and sex-specific variation in recombination. The American Journal of
Human Genetics **63**: 861–869.
Broman K. W., Wu H., Sen S., Churchill G. A., 2003 R/qtl: QTL mapping in experimental crosses.
Bioinformatics **19**: 889–890.
Brunelli J. P., Wertzler K. J., Sundin K., Thorgaard G. H., 2008 Y-specific sequences and polymorphisms
in Rainbow Trout and Chinook Salmon. Genome **51**: 739–748.
Bush W. S., Moore J. H., 2012 Chapter 11: Genome-Wide Association Studies. PLoS Comput Biol **8**:
e1002822.
Carvalho A., 2002 Origin and evolution of the *Drosophila* Y chromosome. Current Opinion in Genetics &
Development **12**: 664–668.
Catchen J. M., Amores A., Hohenlohe P., Cresko W., Postlethwait J. H., 2011 Stacks: building and
genotyping loci *de novo* from short-read sequences. G3 - Genes|Genomes|Genetics **1**: 171–182.
Charlesworth B., 1991 The evolution of sex chromosomes. Science **251**: 1030–1033.
Charlesworth D., Charlesworth B., 1980 Sex differences in fitness and selection for centric fusions
between sex-chromosomes and autosomes. Genet. Res. **35**: 205–214.
Charlesworth D., Charlesworth B., 2005 Sex chromosomes: Evolution of the weird and wonderful.
Current Biology **15**: R129–R131.
Charlesworth D., Charlesworth B., Marais G., 2005 Steps in the evolution of heteromorphic sex
chromosomes. Heredity **95**: 118–128.
Chen S., Zhang G., Shao C., Huang Q., Liu G., Zhang P., Song W., An N., Chalopin D., Volff J.-N., Hong
Y., Li Q., Sha Z., Zhou H., Xie M., Yu Q., Liu Y., Xiang H., Wang N., Wu K., Yang C., Zhou Q.,
Liao X., Yang L., Hu Q., Zhang J., Meng L., Jin L., Tian Y., Lian J., Yang J., Miao G., Liu S., Liang
Z., Yan F., Li Y., Sun B., Zhang H., Zhang J., Zhu Y., Du M., Zhao Y., Schartl M., Tang Q., Wang J.,
2014 Whole-genome sequence of a flatfish provides insights into ZW sex chromosome evolution and
adaptation to a benthic lifestyle. Nat Genet **46**: 253–260.
Clarke J. T., Lloyd G. T., Friedman M., 2016 Little evidence for enhanced phenotypic evolution in early
teleosts relative to their living fossil sister group. Proc. Natl. Acad. Sci. U.S.A.
Crête-Lafrenière A., Weir L. K., Bernatchez L., 2012 Framing the Salmonidae Family phylogenetic
portrait: a more complete picture from increased taxon sampling. PLoS ONE **7**: e46662.
Davidson W. S., Huang T.-K., Fujiki K., Schalburg von K. R., Koop B. F., 2009 The sex determining loci
and sex chromosomes in the Family Salmonidae. Sexual Development **3**: 78–87.
Davidson W. S., Koop B. F., Jones S. J. M., Iturra P., Vidal R., Maass A., Jonassen I., Lien S., Omholt S.
W., 2010 Sequencing the genome of the Atlantic Salmon (*Salmo salar*). Genome Biol. **11**: 403.
Devlin R. H., Stone G. W., Smailus D. E., 1998 Extensive direct-tandem organization of a long repeat
DNA sequence on the Y chromosome of Chinook Salmon (*Oncorhynchus tshawytscha*). J Mol Evol
**46**: 277–287.
Eisbrenner W. D., Botwright N., Cook M., Davidson E. A., Dominik S., Elliott N. G., Henshall J., Jones S.
L., Kube P. D., Lubieniecki K. P., Peng S., Davidson W. S., 2013 Evidence for multiple sex-
determining loci in Tasmanian Atlantic Salmon (*Salmo salar*). **113**: 86–92.
Faber-Hammond J. J., Phillips R. B., Brown K. H., 2015 Comparative analysis of the shared sex-

determination region (SDR) among salmonid fishes. Genome Biology and Evolution **7**: 1972–1987.

Faber-Hammond J., Phillips R. B., Park L. K., 2012 The Sockeye Salmon neo-Y chromosome is a fusion
between linkage groups orthologous to the Coho Y chromosome and the long arm of Rainbow Trout
chromosome 2. Cytogenet Genome Res **136**: 69–74.

Furman B., Evans B. J., 2016 Sequential turnovers of sex chromosomes in African clawed frogs
(*Xenopus*) suggest some genomic regions are good at sex determination. G3 -
Genes|Genomes|Genetics.

Gagnaire P.-A., Normandeau E., Pavey S. A., Bernatchez L., 2013 Mapping phenotypic, expression and
transmission ratio distortion QTL using RAD markers in the Lake Whitefish (*Coregonus*
*clupeaformis*). Mol. Ecol. **22**: 3036–3048.

Grützner F., Rens W., Tsend-Ayush E., El-Mogharbel N., O'Brien P. C. M., Jones R. C., Ferguson-Smith
M. A., Marshall Graves J. A., 2004 In the platypus a meiotic chain of ten sex chromosomes shares
genes with the bird Z and mammal X chromosomes. Nature **432**: 913–917.

Hackett C. A., Broadfoot L. B., 2003 Effects of genotyping errors, missing values and segregation
distortion in molecular marker data on the construction of linkage maps. **90**: 33–38.

Haldane J., 1922 Sex ratio and unisexual sterility in hybrid animals. Journal of Genetics **12**: 101–109.

Hecht B. C., Thrower F. P., Hale M. C., Miller M. R., Nichols K. M., 2012 Genetic architecture of
migration-related traits in rainbow and steelhead trout, *Oncorhynchus mykiss*. G3 -
Genes|Genomes|Genetics **2**: 1113–1127.

Henning F., Lee H. J., Franchini P., Meyer A., 2014 Genetic mapping of horizontal stripes in Lake
Victoria cichlid fishes: benefits and pitfalls of using RAD markers for dense linkage mapping. Mol.
Ecol. **23**: 5224–5240.

Jiang C., Zeng Z. B., 1995 Multiple trait analysis of genetic mapping for quantitative trait loci. Genetics
**140**: 1111–1127.

Johnston S. E., Berenos C., Slate J., Pemberton J. M., 2016 Conserved genetic architecture underlying
individual recombination rate variation in a wild population of Soay sheep (*Ovis aries*). Genetics **203**:
583–598.

Johnstone R., Simpson T. H., Youngson A. F., 1978 Sex reversal in salmonid culture. Aquaculture **13**:
115–134.

Kirkpatrick M., 2016 The evolution of genome structure by natural and sexual selection. J. Hered **108**: 3–
11.

Kitano J., Peichel C. L., 2011 Turnover of sex chromosomes and speciation in fishes. Environ Biol Fish
**94**: 549–558.

Kitano J., Ross J. A., Mori S., Kume M., Jones F. C., Chan Y. F., Absher D. M., Grimwood J., Schmutz J.,
Myers R. M., Kingsley D. M., Peichel C. L., 2009 A role for a neo-sex chromosome in stickleback
speciation. Nature **461**: 1079–1083.

Kodama M., Brieuc M. S. O., Devlin R. H., Hard J. J., Naish K. A., 2014 Comparative mapping between
Coho Salmon (*Oncorhynchus kisutch*) and three other salmonids suggests a role for chromosomal
rearrangements in the retention of duplicated regions following a whole genome duplication event.
G3 - Genes|Genomes|Genetics **4**: 1717–1730.

Kondo M., 2006 Genomic organization of the sex-determining and adjacent regions of the sex
chromosomes of medaka. Genome Research **16**: 815–826.

Kondo M., Nagao E., Mitani H., Shima A., 2001 Differences in recombination frequencies during female
and male meioses of the sex chromosomes of the medaka, *Oryzias latipes*. Genet Res **78**.

Küttner E., Nilsson J., Skúlason S., Gunnarsson S., Ferguson M. M., Danzmann R. G., Civetta A., 2011
Sex chromosome polymorphisms in Arctic Charr and their evolutionary origins. Genome **54**: 852–
861.

Lahn B. T., Pearson N. M., Jegalian K., 2001 The human Y chromosome, in the light of evolution. Nat

Rev Genet **2**: 207–216.

Laporte M., Rogers S. M., Dion-Côté A.-M., Normandeau E., Gagnaire P.-A., Dalziel A. C., Chebib J., Bernatchez L., 2015 RAD-QTL mapping reveals both genome-level parallelism and different genetic architecture underlying the evolution of body shape in Lake Whitefish (*Coregonus clupeaformis*) species pairs. G3 - Genes|Genomes|Genetics **5**: 1481–1491.

Larson W. A., McKinney G. J., Limborg M. T., Everett M. V., Seeb L. W., Seeb J. E., 2015 Identification of multiple QTL hotspots in Sockeye Salmon (*Oncorhynchus nerka*) using Genotyping-by-Sequencing and a dense linkage map. J. Hered. **107**: 122-133.

Larson W. A., McKinney G. J., Seeb J. E., Seeb L. W., 2016 Identification and characterization of sex-associated loci in Sockeye Salmon using genotyping-by-sequencing and comparison with a sex-determining assay based on the *sdY* gene. J. Hered. **107**: 559-566.

Lenormand T., 2003 The evolution of sex dimorphism in recombination. Genetics **163**: 811–822.

Lenormand T., Dutheil J., 2005 Recombination difference between sexes: a role for haploid selection. Plos Biol **3**: e63.

Lenormand T., Engelstädter J., Johnston S. E., Wijnker E., Haag C. R., 2016 Evolutionary mysteries in meiosis. Phil. Trans. R. Soc. B **371**: 20160001

Lien S., Gidskehaug L., Moen T., Ben J Hayes, Berg P. R., Davidson W. S., Omholt S. W., Kent M. P., 2011 A dense SNP-based linkage map for Atlantic Salmon (*Salmo salar*) reveals extended chromosome homeologies and striking differences in sex-specific recombination patterns. BMC Genomics **12**: 615.

Lien S., Koop B. F., Sandve S. R., Miller J. R., Kent M. P., Nome T., Hvidsten T. R., Leong J. S., Minkley D. R., Zimin A., Grammes F., Grove H., Gjuvsland A., Walenz B., Hermansen R. A., Schalburg von K., Rondeau E. B., Di Genova A., Samy J. K. A., Olav Vik J., Vigeland M. D., Caler L., Grimholt U., Jentoft S., Våge D. I., de Jong P., Moen T., Baranski M., Palti Y., Smith D. R., Yorke J. A., Nederbragt A. J., Tooming-Klunderud A., Jakobsen K. S., Jiang X., Fan D., Hu Y., Liberles D. A., Vidal R., Iturra P., Jones S. J. M., Jonassen I., Maass A., Omholt S. W., Davidson W. S., 2016 The Atlantic Salmon genome provides insights into rediploidization. Nature **533**: 200–205.

Limborg M. T., Seeb L. W., Seeb J. E., 2016 Sorting duplicated loci disentangles complexities of polyploid genomes masked by genotyping by sequencing. Mol. Ecol. **25**: 2117–2129.

Lubieniecki K. P., Lin S., Cabana E. I., Li J., Lai Y. Y. Y., Davidson W. S., 2015 Genomic instability of the sex-determining locus in Atlantic Salmon (*Salmo salar*). G3 - Genes|Genomes|Genetics **5**: 2513–2522.

Lv W., Zheng X., Kuang Y., Cao D., Yan Y., Sun X., 2016 QTL variations for growth-related traits in eight distinct families of Common Carp (*Cyprinus carpio*). BMC Genet **17**: 65.

Mable B. K., 2004 'Why polyploidy is rarer in animals than in plants': myths and mechanisms. Biological Journal of the Linnean Society **82**: 453–466.

Mackay T. F., 2001 The genetic architecture of quantitative traits. Annu. Rev. Genet. **35**: 303–339.

Mackay T. F. C., Stone E. A., Ayroles J. F., 2009 The genetics of quantitative traits: challenges and prospects. Nat Rev Genet **10**: 565–577.

Mank J. E., Avise J. C., 2009 Evolutionary diversity and turn-over of sex determination in teleost fishes. Sex Dev **3**: 60–67.

Marshall Graves J. A., Peichel C. L., 2010 Are homologies in vertebrate sex determination due to shared ancestry or to limited options? Genome Biol. **11**: 205.

May B., Delany M. E., 2015 Meiotic models to explain classical linkage, pseudolinkage, and chromosomal pairing in tetraploid derivative salmonid genomes: II. Wright is still right. J. Hered. **106**: 762–766.

McKinney G. J., Seeb L. W., Larson W. A., Gomez Uchida D., Limborg M. T., Brieuc M. S. O., Everett M. V., Naish K. A., Waples R. K., Seeb J. E., 2016 An integrated linkage map reveals candidate

genes underlying adaptive variation in Chinook Salmon (*Oncorhynchus tshawytscha*). Mol. Ecol.
Resour. **16**: 769–783.

Moen T., Hoyheim B., Munck H., Gomez-Raya L., 2004 A linkage map of Atlantic Salmon (*Salmo salar*)
reveals an uncommonly large difference in recombination rate between the sexes. Anim. Genet. **35**:
81–92.

Moghadam H. K., Ferguson M. M., Danzmann R. G., 2007 Linkage variation at the sex-determining locus
within Fraser strain Arctic charr *Salvelinus alpinus*. Journal of Fish Biology **71**: 294–301.

Muller H. J., 1925 Why polyploidy is rarer in animals than in plants. The American Naturalist **59**: 346–
353.

Myosho T., Takehana Y., Hamaguchi S., 2015 Turnover of sex chromosomes in Celebensis group
Medaka fishes. G3 - Genes|Genomes|Genetics **5**: 2685-2691.

Nagler J. J., Bouma J., Thorgaard G. H., Dauble D. D., 2001 High incidence of a male-specific genetic
marker in phenotypic female Chinook Salmon from the Columbia River. Environ. Health Perspect.
**109**: 67–69.

Naish K. A., Phillips R. B., Brieuc M. S. O., Newton L. R., Elz A. E., Park L. K., 2013 Comparative
genome mapping between Chinook Salmon (*Oncorhynchus tshawytscha*) and Rainbow Trout (*O.*
*mykiss*) based on homologous microsatellite loci. G3 - Genes|Genomes|Genetics **3**: 2281–2288.

Nei M., 1969 Linkage modification and sex difference in recombination. Genetics **63**: 681-699.

Nugent C. M., Easton A. A., Norman J. D., Ferguson M. M., Danzmann R. G., 2016 A SNP based linkage
map of the Arctic Charr (*Salvelinus alpinus*) genome provides insights into the diploidization process
after whole genome duplication. G3 - Genes|Genomes|Genetics.

Ohno S., 1970 *Evolution by Gene Duplication*. Springer Berlin Heidelberg, Berlin, Heidelberg.

Orr H. A., 1990 "Why polyploidy is rarer in animals than in plants" revisited. The American Naturalist
**136**: 759–770.

Otto S. P., 2007 The Evolutionary Consequences of Polyploidy. Cell **131**: 452–462.

Palti Y., Vallejo R. L., Gao G., Liu S., Hernandez A. G., Rexroad C. E. III, Wiens G. D., 2015 Detection
and validation of QTL affecting bacterial cold water disease resistance in Rainbow Trout using
restriction-site associated DNA sequencing. PLoS ONE **10**: e0138435.

Pennell M. W., Kirkpatrick M., Otto S. P., Vamosi J. C., Peichel C. L., Valenzuela N., Kitano J., 2015 Y
fuse? Sex chromosome fusions in fishes and reptiles. PLoS Genet **11**: e1005237.

Perrin N., 2009 Sex reversal: a fountain of youth for sex chromosomes? Evolution **63**: 3043–3049.

Phillips R., Ráb P., 2001 Chromosome evolution in the Salmonidae (Pisces): an update. Biol Rev Camb
Philos Soc **76**: 1–25.

Phillips R. B., 2013 Evolution of the sex chromosomes in salmonid fishes. Cytogenet Genome Res **141**:
177–185.

Phillips R. B., Ihssen P. E., 1985 Identification of sex chromosomes in lake trout (*Salvelinus namaycush*).
Cytogenet. Cell Genet. **39**: 14–18.

Phillips R. B., Keatley K. A., Morasch M. R., Ventura A. B., Lubieniecki K. P., Ben F Koop, Danzmann
R. G., Davidson W. S., 2009 Assignment of Atlantic Salmon (*Salmo salar*) linkage groups to specific
chromosomes: Conservation of large syntenic blocks corresponding to whole chromosome arms in
Rainbow Trout (*Oncorhynchus mykiss*). BMC Genet **10**: 46.

Phillips R. B., Morasch M. R., Park L. K., Naish K. A., Devlin R. H., 2005 Identification of the sex
chromosome pair in Coho Salmon (*Oncorhynchus kisutch*): lack of conservation of the sex linkage
group with Chinook Salmon (*Oncorhynchus tshawytscha*). Cytogenet Genome Res **111**: 166–170.

Phillips R. B., Nichols K. M., DeKoning J. J., Morasch M. R., Keatley K. A., Rexroad C., Gahr S. A.,
Danzmann R. G., Drew R. E., Thorgaard G. H., 2006 Assignment of Rainbow Trout linkage groups
to specific chromosomes. Genetics **174**: 1661–1670.

R Development Core Team, 2017 R: A language and environment for statistical computing.

Rexroad C. E., Palti Y., Gahr S. A., Vallejo R. L., 2008 A second generation genetic map for Rainbow
Trout (*Oncorhynchus mykiss*). BMC Genet **9**: 74.

Rexroad C. E., Vallejo R. L., Liu S., Palti Y., Weber G. M., 2012 QTL affecting stress response to
crowding in a Rainbow Trout broodstock population. BMC Genet **13**: 97.

Rockman M. V., 2012 The QTN program and the alleles that matter for evolution: all that's gold does not
glitter. Evolution **66**: 1–17.

Roco Á. S., Olmstead A. W., Degitz S. J., Amano T., Zimmerman L. B., Bullejos M., 2015 Coexistence of
Y, W, and Z sex chromosomes in *Xenopus tropicalis*. Proceedings of the National Academy of
Sciences **112**: E4752–E4761.

Rogers S. M., Bernatchez L., 2007 The genetic architecture of ecological speciation and the association
with signatures of selection in natural Lake Whitefish (*Coregonus* sp. *Salmonidae*) species pairs.
Molecular Biology and Evolution **24**: 1423–1438.

Rondeau E. B., Minkley D. R., Leong J. S., Messmer A. M., Jantzen J. R., Schalburg von K. R., Lemon C.,
Bird N. H., Koop B. F., 2014 The genome and linkage map of the Northern Pike (*Esox lucius*):
conserved synteny revealed between the salmonid sister group and the Neoteleostei. PLoS ONE **9**:
e102089.

Ross J. A., Urton J. R., Boland J., Shapiro M. D., Peichel C. L., 2009 Turnover of sex chromosomes in the
stickleback fishes (*Gasterosteidae*). PLoS Genet **5**: e1000391–12.

Sakamoto T., Danzmann R. G., Gharbi K., Howard P., Ozaki A., Khoo S. K., Woram R. A., Okamoto N.,
Ferguson M. M., Holm L. E., Guyomard R., Hoyheim B., 2000 A microsatellite linkage map of
Rainbow Trout (*Oncorhynchus mykiss*) characterized by large sex-specific differences in
recombination rates. Genetics **155**: 1331–1345.

Santure A. W., Poissant J., De Cauwer I., Oers K., Robinson M. R., Quinn J. L., Groenen M. A. M.,
Visser M. E., Sheldon B. C., Slate J., 2015 Replicated analysis of the genetic architecture of
quantitative traits in two wild great tit populations. Mol. Ecol. **24**: 6148–6162.

Sauvage C., Vagner M., Derôme N., Audet C., Bernatchez L., 2012a Coding gene single nucleotide
polymorphism mapping and quantitative trait loci detection for physiological reproductive traits in
Brook Charr, *Salvelinus fontinalis*. G3 - Genes|Genomes|Genetics **2**: 379–392.

Sauvage C., Vagner M., Derôme N., Audet C., Bernatchez L., 2012b Coding gene SNP mapping reveals
QTL linked to growth and stress response in Brook Charr (*Salvelinus fontinalis*). G3 -
Genes|Genomes|Genetics **2**: 707–720.

Session A. M., Uno Y., Kwon T., Chapman J. A., Toyoda A., Takahashi S., Fukui A., Hikosaka A.,
Suzuki A., Kondo M., van Heeringen S. J., Quigley I., Heinz S., Ogino H., Ochi H., Hellsten U.,
Lyons J. B., Simakov O., Putnam N., Stites J., Kuroki Y., Tanaka T., Michiue T., Watanabe M.,
Bogdanovic O., Lister R., Georgiou G., Paranjpe S. S., van Kruijsbergen I., Shu S., Carlson J.,
Kinoshita T., Ohta Y., Mawaribuchi S., Jenkins J., Grimwood J., Schmutz J., Mitros T., Mozaffari S.
V., Suzuki Y., Haramoto Y., Yamamoto T. S., Takagi C., Heald R., Miller K., Haudenschild C.,
Kitzman J., Nakayama T., Izutsu Y., Robert J., Fortriede J., Burns K., Lotay V., Karimi K., Yasuoka
Y., Dichmann D. S., Flajnik M. F., Houston D. W., Shendure J., DuPasquier L., Vize P. D., Zorn A.
M., Ito M., Marcotte E. M., Wallingford J. B., Ito Y., Asashima M., Ueno N., Matsuda Y., Veenstra
G. J. C., Fujiyama A., Harland R. M., Taira M., Rokhsar D. S., 2016 Genome evolution in the
allotetraploid frog *Xenopus laevis*. Nature **538**: 336–343.

Singer A., Perlman H., Yan Y., Walker C., Corley-Smith G., Brandhorst B., Postlethwait J., 2002 Sex-
specific recombination rates in zebrafish (*Danio rerio*). Genetics **160**: 649–657.

Slate J., 2005 Quantitative trait locus mapping in natural populations: progress, caveats and future
directions. Mol. Ecol. **14**: 363–379.

Slate J., 2008 Robustness of linkage maps in natural populations: a simulation study. Proceedings of the
Royal Society B: Biological Sciences **275**: 695–702.

Steinemann M., Steinemann S., 1998 Enigma of Y chromosome degeneration: neo-Y and neo-X

chromosomes of *Drosophila miranda* a model for sex chromosome evolution. Genetica **102-103**:
409–420.

Stöck M., Horn A., Grossen C., Lindtke D., Sermier R., Betto-Colliard C., Dufresnes C., Bonjour E.,
Dumas Z., Luquet E., Maddalena T., Sousa H. C., Martinez-Solano I., Perrin N., 2011 Ever-young
sex chromosomes in European tree frogs. Plos Biol **9**: e1001062–9.

Stöck M., Savary R., Betto-Colliard C., Biollay S., Jourdan-Pineau H., Perrin N., 2013 Low rates of X-Y
recombination, not turnovers, account for homomorphic sex chromosomes in several diploid species
of Palearctic green toads (*Bufo viridis* subgroup). Journal of Evolutionary Biology **26**: 674–682.

Sutherland B. J. G., Gosselin T., Normandeau E., Lamothe M., Isabel N., Audet C., Bernatchez L., 2016
Salmonid chromosome evolution as revealed by a novel method for comparing RADseq linkage maps.
Genome Biology and Evolution: **8**: 3600-3617.

Taylor J. S., Braasch I., Frickey T., Meyer A., Van de Peer Y., 2003 Genome duplication, a trait shared by
22,000 species of ray-finned fish. Genome Research **13**: 382–390.

Thorgaard G. H., 1977 Heteromorphic sex chromosomes in male Rainbow Trout. Science **196**: 900–902.

Trivers R., 1998 Sex differences in rates of recombination and sexual selection. In:Michod R, Levin D
(Eds.), *The evolution of sex*,, pp. 270–286.

Tsai H. Y., Hamilton A., Guy D. R., Tinch A. E., Bishop S. C., Houston R. D., 2015 The genetic
architecture of growth and fillet traits in farmed Atlantic Salmon (*Salmo salar*). BMC Genet **16**: 117–
11.

van Doorn G. S., Kirkpatrick M., 2007 Turnover of sex chromosomes induced by sexual conflict. **449**:
909–912.

van Ooijen J. W., 2006 JoinMap4: Software for the calculation of genetic linkage maps in experimental
populations.

Wei T., Simko V., 2017 corrplot: Visualization of a correlation matrix.

Woram R. A., Gharbi K., Sakamoto T., Høyheim B., Holm L.-E., Naish K., McGowan C., Ferguson M.
M., Phillips R. B., Stein J., Guyomard R., Cairney M., Taggart J. B., Powell R., Davidson W.,
Danzmann R. G., 2003 Comparative genome analysis of the primary sex-determining locus in
salmonid fishes. Genome Research **13**: 272–280.

Wu R., Ma C.-X., Painter I., Zeng Z.-B., 2002 Simultaneous maximum likelihood estimation of linkage
and linkage phases in outcrossing species. Theor Popul Biol **61**: 349–363.

Xu S., 2003 Theoretical basis of the Beavis effect. Genetics **165**: 2259–2268.

Yano A., Guyomard R., Nicol B., Jouanno E., Quillet E., Klopp C., Cabau C., Bouchez O., Fostier A.,
Guiguen Y., 2012a An immune-related gene evolved into the master sex-determining gene in
Rainbow Trout, *Oncorhynchus mykiss*. Curr. Biol. **22**: 1423–1428.

Yano A., Nicol B., Jouanno E., Quillet E., Fostier A., Guyomard R., Guiguen Y., 2012b The sexually
dimorphic on the Y-chromosome gene (*sdY*) is a conserved male-specific Y-chromosome sequence in
many salmonids. Evol Appl **6**: 486–496.

Yáñez J. M., Houston R. D., Newman S., 2014 Genetics and genomics of disease resistance in salmonid
species. Front. Genet. **5**: 415.

Yoshida K., Makino T., Yamaguchi K., Shigenobu S., Hasebe M., Kawata M., Kume M., Mori S., Peichel
C. L., Toyoda A., Fujiyama A., Kitano J., 2014 Sex chromosome turnover contributes to genomic
divergence between incipient stickleback species. PLoS Genet **10**: e1004223–16.

1014

 **SUPPLEMENTAL MATERIAL**

**Table S1.** Phenotype average, standard deviation and sample size in males and females. Phenotypes
showing any differences between males and females ($p \leq 0.2$) included sex as a covariate in the model.
Sex-specific phenotypes were only tested within the one sex and therefore had smaller sample sizes.

**Table S2.** Complete QTL table with all identified genome- and chromosome-wide QTLs and associated
values, including marker sequence and SNP, and effect size of different genotypes.

**Figure S1.** Correlation plot of phenotypes used in QTL analysis. Phenotype pairs that do not share any
individuals for correlation are shown with '?'.

**Figure S2.** Heterochiasmy plots for individual chromosomes in the maternal (a) and paternal (b)
haplotypes. Grey boxes above the plot indicate probable residually tetraploid chromosome arms, and
centromere positions transferred from Chinook Salmon are indicated by stars. When Chinook Salmon
chromosomes were not fusions but were metacentrics (n = 2), this is denoted by a '?' to denote the
uncertainty as to the centromere position in Brook Charr.

**File S1.** Required files for running Rqtl analysis (phenotype (.qua), map (.map) and genotype (.loc)). The
map file corresponds to the female map from Sutherland *et al.* 2016. See the Data Availability for code for
performing complete analysis with these files.

**File S2.** Fasta file with RAD-seq tags output using the STACKs population module for alleles from all
individual offspring and parents.