Research Article

SYSTEMATIC   CONSERVATION   PLANNING   FOR   INTRASPECIFIC

GENETIC DIVERSITY.

Ivan   PAZ-VINAS[1,2,3*],   Géraldine   LOOT[1,4],   Virgilio   HERMOSO[6],   Charlotte

VEYSSIERE[1,4],   Nicolas   POULET[7],   Gaël   GRENOUILLET[1,4]   and   Simon

BLANCHET[1,5]

[1] Centre National de la Recherche Scientifique (CNRS), Université Paul Sabatier (UPS),

École Nationale de Formation Agronomique (ENFA); UMR 5174 EDB (Laboratoire

Évolution & Diversité Biologique), 118 route de Narbonne, 31062 Toulouse cedex 4,

France.

[2] Aix-Marseille Université, CNRS, IRD, Avignon Université; UMR 7263 IMBE, Équipe

EGE, Centre Saint-Charles, Case 36, 3 place Victor Hugo, 13331 Marseille cedex 3,

France.

[3] Université de Lyon, CNRS, ENTPE; UMR 5023 LEHNA, Laboratoire d'Écologie des

Hydrosystèmes Naturels et Anthropisés, 6 rue Raphaël Dubois, 69622 Villeurbanne,

France.

[4] Université de Toulouse, UPS; UMR 5174 EDB, 118 route de Narbonne, 31062

Toulouse cedex 4, France.

[5] CNRS, Station d'Écologie Théorique et Expérimentale; UMR 5321, 09200 Moulis,

France.

[6] Centre Tecnologic Forestal de Catalunya, Crta. Sant Llorenc de Monunys, Km 2,

25280, Solsona. Lleida, Spain.

[7] Pôle Écohydraulique ONEMA-IMFT, Office National de l'Eau et des Milieux

Aquatiques, France.

Running head: Conservation of genetic diversity

Keywords: biodiversity conservation, dendritic ecological networks, multi-specific,

freshwater fish, conservation genetics, spatial patterns of biodiversity

*Corresponding author:

32
Ivan PAZ-VINAS
Université Lyon 1
CNRS, ENTPE, UMR 5023 LEHNA
Laboratoire d'Écologie des Hydrosystèmes Naturels et Anthropisés
3, rue Raphaël Dubois - Bât. Darwin C
69622 Villeurbanne Cedex France
Phone: (+33) 04 72 43 26 03
E-mail: ivanpaz23@gmail.com

41
42   **Co-author's e-mails:**

43
Simon Blanchet: simon.blanchet@sete.cnrs.fr
Virgilio Hermoso: virgilio.hermoso@gmail.com
Charlotte Veyssière: veyssiere.charlotte@gmail.com
Nicolas Poulet: nicolas.poulet@onema.fr
Gaël Grenouillet: gael.grenouillet@univ-tlse3.fr
Géraldine Loot: geraldine.loot@univ-tlse3.fr
50

Abstract

Intraspecific diversity informs the demographic and evolutionary histories of

populations, and should hence be a main conservation target. Although approaches exist

for identifying relevant biological units of conservation, attempts to identify priority

conservation areas for intraspecific diversity are scarce, especially within a multi-

specific framework. We used neutral molecular data on six European freshwater

cyprinid fish species (*Squalius cephalus*, *Phoxinus phoxinus, Barbatula barbatula*,

Gobio occitaniae, *Leuciscus burdigalensis* and *Parachondrostoma toxostoma*) sampled

at the riverscape scale (i.e. the Garonne-Dordogne River basin, France) to determine

hot- and cold-spots of genetic diversity, and to identify priority conservation areas using

a systematic conservation planning approach. We demonstrate that systematic

conservation planning is efficient for identifying priority areas representing a predefined

part of the total genetic diversity of a whole landscape. Classical indices of genetic

diversity (allelic richness, genetic uniqueness), but private allelic richness, were poor

predictors for identifying these priority areas. Moreover, we identified weak

congruencies and surrogacies among conservation solutions found for each species,

implying that conservation solutions are highly species-specific. Nonetheless, we

showed that priority areas identified using intraspecific genetic data from multiple

species provide more effective conservation solutions than areas identified for single

species or on the basis of traditional taxonomic criteria.

1. Background

Biodiversity conservation is often addressed by identifying protected areas with

high biodiversity and/or landscape values [1]. Conservation areas are generally

identified as areas with high proportions of endemic, rare or iconic species [2].

Alternatively, conservation planning can be based on the concept of complementarity

between conservation areas [3], and on cost-effectiveness analyses such as systematic

conservation planning methods (hereafter SCP [4]). SCP aims at identifying a number

of complementary and *irreplaceable* sites best representing a predefined proportion of

the biodiversity observed in a region and that should be managed for conservation in

priority at a minimum cost.

In the meantime, there have been major conceptual and empirical leaps

advocating the importance of considering species evolutionary history into conservation

plans [5,6]. For instance, there have been attempts to include information on the

phylogenetic history of species assemblages into SCP approaches to preserve both

species identities and their macro-evolutionary history [7–9]. However, genetic

diversity observed at the *population level* (i.e. within species) has rarely been

considered in SCP [but see 10–13]. *Intraspecific genetic diversity* is a fundamental facet

of biodiversity, as it informs the demographic and colonisation history of populations

(i.e. changes in effective population size over time and space), their level of biological

connectivity, as well as their potential for adapting to environmental changes [14–17].

Conservation geneticists generally use neutral genetic diversity (i.e. part of the genome

not affected by adaptive processes) to identify relevant and significant evolutionary

units within a set of individuals sampled in a landscape [18], mainly by analysing their

levels of genetic differentiation or (dis)continuity [19,20]. For instance, the

"Evolutionary Significant Units" (ESUs) framework has been proposed for representing

different lineages for conservation prioritization [21,22]. By emphasizing

complementarity among areas –rather than differentiation–, SCP appears as a promising

and complementary framework for identifying an optimal set of sites allowing

preserving a pre-defined amount of the total genetic diversity observed within a

landscape.

Most SCP studies that have yet considered intraspecific genetic diversity aimed

at identifying efficient surrogates for preserving genetic diversity [12,13]. For instance,

two recent studies showed that protected areas identified using traditional species

distribution data [13], and environmental and geographic descriptors [12] can be

efficient surrogates in conservation planning for representing intraspecific genetic

diversity. Although these studies shed light on how SCP can help preserving the genetic

diversity of multiple species, only few studies have directly used raw genotypic data

obtained from multiple species as source of information for SCP analyses (see [11] for

the first attempt using data from a single species). Datasets generated using neutral

genetic markers such as microsatellite markers (i.e. alleles presence/absence matrices)

are similar to datasets generally used for SCP (i.e. species presence/absence matrices),

hence making possible applying SCP to genotypic surveys without needing intermediate

modelling procedures or the calculation of summary statistics of genetic diversity (as in

[10,12] for instance) that may fail to capture the whole genetic diversity of populations.

Here, we tested the potential of SCP analyses to identify priority conservation

areas accounting for intraspecific genetic diversity measured for a species assemblage at

the landscape scale (i.e. at the scale of an entire river basin). We first considered a set of

four common and representative freshwater fish species (*Squalius cephalus*, *Gobio*

occitaniae, *Barbatula barbatula* and *Phoxinus phoxinus*) to test the influence of

conservation targets (proportion of the total amount of genetic diversity to be secured by

irreplaceable sites) and analytical strategies (analysing each species independently or all

species pooled) on final conservation solutions (number and identity of irreplaceable

sites). We then included two rare species of particular conservation interest (*Leuciscus*

burdigalensis and *Parachondrostoma toxostoma*) to test the relevance of the SCP

approach in a "real conservation-oriented study". For these two species (and the four

common species), we ran SCP analyses considering a typical conservation target

[23,24] to explore the spatial distribution of irreplaceable sites in a riverscape, and to

test for congruence and surrogacy in irreplaceable sites among species, and more

particularly between rare and common species. We further tested whether irreplaceable

sites identified using SCP analyses are more efficient for preserving intraspecific

genetic diversity than sites randomly selected or than sites selected on the basis of

traditional taxonomic criteria (e.g. sites displaying high species richness). We finally

tested whether irreplaceable sites were correctly predicted by classical indices of genetic

diversity such as allelic richness. We demonstrate that preserving the intraspecific

genetic diversity of a species assemblage using systematic conservation planning is an

efficient and relevant, yet complex, approach necessitating appropriate analyses to assist

the decision-making process.

2. Methods

<u>Data collection</u>

Biological models. We focused on an assemblage of six cyprinid freshwater fish

species. Three of them are widespread in Europe (i.e. *Squalius cephalus*, *Phoxinus*

phoxinus and *Barbatula barbatula*) whereas three of them (*Gobio occitaniae*, *Leuciscus*

burdigalensis and *Parachondrostoma toxostoma*) are endemic to Southern France [25].

This set of species covers a large functional trait space that is representative of many

freshwater fish communities. For instance, *S. cephalus* is a large-bodied fish species

with a long lifespan (i.e. it can be 60 cm long and live up to 15 years [25]) whereas at

the other extreme *Ph. phoxinus* is a small-bodied species with a shorter lifespan (i.e. it is

less than 12 cm long and usually lives up to 4-5 years [25]). From an ecological

perspective, *G. occitaniae, Pa. toxostoma* and *B. barbatula* are bottom feeders, whereas

S. cephalus and *Ph. phoxinus* are water column feeders and *L. burdigalensis* is more

opportunistic. Further, *B. barbatula* is mainly active during the night, while the other

species are particularly active during the day. Four of these species are relatively

abundant (i.e. *S. cephalus, G. occitaniae, Ph. phoxinus* and *B. barbatula*), although they

vary greatly in their ecological niche and hence their spatial occupancy in the river

network (see Figure S1 for maps representing the spatial distribution of sampling sites

for each species, which roughly corresponds to their spatial distribution in the Garonne-

Dordogne river basin). We will hereafter refer to this set of species as the "common"

species. The two other species are rare (*L. burdigalensis*) to very rare (*Pa. toxostoma;*

see Figure S1) in the Garonne-Dordogne river basin, and are of particular interest for

conservation. *Leuciscus burdigalensis* is a recently described species that is locally

experiencing both demographic and genetic bottlenecks in many populations [26,27].

Parachondrostoma toxostoma is a vulnerable species [28] listed in the IUCN red list, in

the Annex II of the European Union Habitats Directive and in Appendix III of the Bern

Convention [28].

Sampling design. During Spring/Summer 2010-2011, we used electrofishing to sample

92 sites distributed across 35 rivers from a whole European river basin, the Garonne-

Dordogne River basin (>100,000 km^2, South-Western France; Figure S1; Table S1).

Sampling sites were chosen to cover the entire distribution of each species at the

riverscape scale, and to allow characterising spatial patterns of genetic diversity for all

these species. Up to 25 individuals *per* species *per* site were sampled when possible.

Not all species were present at all sampling sites (Figure S1; Table S1), and not all sites

provided 25 individuals *per* species due to low densities. In these cases, we captured as

many individuals as possible. We anesthetized each individual and then we collected

and stored in 90% ethanol a fragment of their pelvic fin. All individuals were released

alive at their sampling location.

Genotyping. Genomic DNA was extracted using a salt-extraction protocol [29]. We

used multiplexed Polymerase Chain Reactions (PCRs) to co-amplify 8 to 15

microsatellite loci depending on the species (8 for *G. occitaniae*, 9 for *B. barbatula*, 10

for *S. cephalus* and *Ph. phoxinus*, 14 for *L. burdigalensis* and 15 for *Pa. toxostoma*). We

used 5-20 ng of genomic DNA and QIAGEN® Multiplex PCR Kits (Qiagen, Valencia,

CA, USA) to perform PCR amplifications. Details on loci, primer concentrations, PCR

conditions and multiplex sets can be found in Table S2. The genotyping was conducted

on an ABI PRISM™ 3730 Automated Capillary Sequencer (Applied Biosystems, Foster

City, CA, USA). The scoring of allele sizes was done using GENEMAPPER® v.4.0

(Applied Biosystems).

Genetic diversity assessment. Given the good spatial resolution of the sampling obtained

for the four common species (i.e. *S. cephalus*, *G. occitaniae*, *Ph. phoxinus* and *B.*

barbatula), descriptive genetic analyses were conducted for sampling sites displaying a

minimum sample size of N=10 individuals for these species, so as to maximize

consistency on subsequent allelic frequency-based genetic analyses (see Figure S1 and

Table S1 for details on sample sizes). For the two rare species, for which the sampling

was more restricted (i.e. *L. burdigalensis* and *Pa. toxostoma*), genetic analyses were

conducted for sampling sites displaying a minimum sample size of N=6 individuals, so

as to maximize the number of sampling sites included in the SCP procedures. This

allowed us to considerably increment the number of sites retained for genetic analyses

by 38.1% (21 to 29 sites) and 75% (from 8 to 14 sites) for *L. burdigalensis* and *Pa.*

toxostoma respectively and hence to improve the representation of the spatial

distribution of these two species. We then determined for each of the six species the

occurrence of null alleles and potential scoring errors with the program

MICROCHECKER 2.3 [30]. We tested for departures from Hardy-Weinberg (HW)

equilibrium with the 'adegenet' R package v1.6-2 [31]. The program GENEPOP v4.0

[32] was used to assess linkage disequilibrium among loci within sites. Levels of

significance for these multiple tests were adjusted using Bonferroni procedures. We

found significant deviations from HW for a few locus/population pairs for each of the

six species (see Appendix A1 and Supplementary Data File 1 for details and raw tables),

and significant linkage disequilibrium and homozygote excesses only for the four

common species (Appendix A1; Supplementary Data File 1). However, no clear

patterns were observed for any species across loci and populations for these deviations.

Given the small extent of these deviations and the large spatial extent of the databases,

we assumed that they weakly affected our main findings (Appendix A1; Supplementary

Data File 1).

To assess within-site intraspecific genetic diversity (i.e. α-IGD), we applied

rarefaction procedures implemented in ADZE v1.0 [33] to calculate both allelic richness

(AR [34]) and private allelic richness (PA [35]) at the sampling site level (based on a

minimum of N=10 individuals for common species or N=6 individuals for rare species).

To assess the among-site component of intraspecific genetic diversity (i.e. β-IGD), we

used the R package 'mmod' [36] to calculate –for each species– a pairwise genetic

differentiation index (i.e. D_{est} [37]). For each site (and species), we then derived the

averaged value of all pairwise D_{est} values estimated between one given site and all the

remaining sites, so as to obtain a single value *per* site. For rare species, the impact of

considering sites with N≤6 individuals instead of N≤10 on AR, PA and D_{est} was

relatively low (i.e. no clear impact neither on PA nor on D_{est} values, and a mean

decrease in AR values of 0.295 alleles *per* site for *Pa. toxostoma* and of 1.183 alleles *per*

site for *L. burdigalensis*). This impact was however overcompensated by the increase in

the number of sites retained for these two species when considering sites with N≤6

individuals.

<u>Testing the suitability of SCP analyses for intraspecific genetic diversity</u>

In the first step, we tested the influence of conservation targets and analytical

strategies on final conservation solutions. We focused specifically on data from the four

common species, as their large coverage of the sampling area is more suited for the

demonstrative exercise done in this step.

Spatial patterns of genetic diversity. We first used geostatistical modelling tools to

explore spatial patterns of α and β genetic diversity for the four common species at the

riverscape scale by predicting the distributions of *AR*, *PA* and D_{est} from the observed

empirical values using Generalized Linear Models for Spatial Stream Networks

(GLMSSN [38,39]). This was done using the 'STARS' toolset of ARCGIS v10.2 and

the R package 'SSN' [38,39]. We conducted a model selection procedure based on a

comparison of Akaike Information Criteria (AIC) estimated for several competing

GLMSSNs. These models were built by (i) assuming three geographic descriptors (i.e.

topological distance from the outlet, longitude and latitude) as explanatory variables,

and (ii) choosing a tail-down covariance structure model among the following ones:

exponential, Mariah, spherical, linear-with-sill and Epanechnikov. The model with the

lowest AIC score (see Supplementary Data File 1 for raw results) was considered as the

best model and was used to estimate the slope and the significance of the relationships

between explanatory variables and each genetic index, which allowed testing whether or

not spatial patterns of intraspecific genetic diversity can be detected. We finally used

predictions from the best models to produce krigged maps for each common species and

each genetic index, so as to visually represent the spatial distribution of intraspecific

genetic diversity across the whole river drainage, and to visually highlight hot- and

cold-spots of intraspecific genetic diversity. We also calculated Pearson's correlation

coefficients between values calculated at the site level for each genetic index (i.e. *AR*,

PA and D_{est}) for each pair of common species, so as to test for spatial congruence in

patterns of genetic diversity among common species.

Identification of irreplaceable sites. We then tested whether conservation targets (i.e.

the percentage of total number of conservation units to be present in the final

conservation solution) and analytical approaches (i.e. species-specific or species-pooled

analyses) influence the identification of irreplaceable sites. SCP methods traditionally

use species presence/absence data as input data to identify irreplaceable sites for the

conservation of taxonomic diversity at the community level (i.e. sites that cannot be

excluded from an optimal selection of sites for conservation [40]). Here, we replaced

species presence/absence data with allele presence-absence data as in [11] to identify

irreplaceable sites for the conservation of intraspecific genetic diversity of each species

in the river drainage. We used the program Marxan v2.1 [40] and genotypic data from

the common species to identify, for each common species independently, an optimal set

of sites that best represent at least 50, 75, 90 or 100% of the total number of alleles

present in the whole riverscape at a minimum "cost", which corresponds to four

conservation targets. Given the lack of ground estimates for conservation cost, we used

a constant cost *per* site, so our objective translated into identifying the minimum

number of sites that represent a given proportion of intraspecific genetic diversity (i.e.

50, 75, 90 or 100% of the total pool of alleles represented at least once [41]). We

arbitrarily choose the 50, 75 and 100% conservation targets to explore how the

proportion of alleles to protect affects the selection of irreplaceable sites. We

additionally tested the 90% conservation target, as it corresponds to a threshold target

being typically assumed in *ex-situ* conservation plans [23,24]. In order to estimate the

relevance of each site to preserve a given proportion of the allelic diversity in the river

basin, we used two different methods. In case of 100% of allelic diversity, we used the

traditional irreplaceability measure reported by Marxan that ranges between 0% (not

irreplaceable) and 100% (highly irreplaceable). This measure is estimated by running

the optimization algorithm a number of times (N=100 runs in our case) and then

computing the frequency of selection of each site within the solutions obtained. Sites

with unique allelic composition will be selected across all runs and reported as highly

irreplaceable, whereas sites with more common alleles, replaceable by other sites with

the same alleles, will appear poorly irreplaceable. For the other conservation targets, we

selected a random pull of 50, 75 and 90% of the total number of alleles existing at the

basin level and for each species. We then ran the optimization algorithm to identify the

minimum set of sites representing this particular selection of alleles. Given that the

analyses were run only on a subset of alleles, we replicated this process 100 times to

minimize the effect of the arbitrary selection of alleles. For each subset of alleles, we

ran the Marxan procedure as explained above (e.g. constant cost and 100 runs for each)

and retained the best solution for subsequent analyses (solution with the lowest value

for the objective function). We then calculated the irreplaceability as the frequency of

selection of each site within the 100 random pull of alleles. For a given species and a

given conservation target, we considered a site as irreplaceable for genetic diversity

conservation when it displayed an irreplaceability value of 100%. We selected such a

high threshold so as to be conservative. We tested and compared visually how the

proportions of irreplaceable sites vary among species and conservation targets.

To test how pooling data from several species affect the identification of

irreplaceable sites, we further performed a "pooled" analysis, in which all alleles found

for each common species at a site were pooled together in a single input dataset. We

then selected a random pull of 30, 50, 75, 90 and 100% of the total number of alleles

existing at the basin level (all common species confounded), and performed 100 Marxan

runs *per* conservation targets to identify the minimum set of sites representing these

particular selections of alleles. Allele counts for all species are provided in the

Supplementary Data File 1.

A real conservation-oriented study using SCP approaches

In this second step, we (i) explored the spatial distribution of irreplaceable sites,

(ii) tested whether conservation solutions found for each species are congruent among

species, (iii) tested whether solutions found for one species can be used as a surrogate

for other species, (iv) assessed the overall efficiency of the different conservation

solutions we identified for preserving the overall genetic diversity of all species (v)

compared the efficiency of conservation solutions identified through SCP to the

efficiency of alternative conservation solutions based on traditional taxonomic criteria

(e.g. in sites with high species richness), and finally (vi) tested whether –or not–

irreplaceable sites are correctly predicted by classical indices of genetic diversity. In this

step, we included the two rare species, since congruence and surrogacy are particularly

important to measure for rare species for which data are more difficult to collect. We

therefore focus more specifically on the comparisons comparing common *vs*. rare

species.

Identification of irreplaceable sites for genetic conservation. We focused only on

irreplaceable sites identified for the 90% target and used the program Marxan as

described above to identify these sites for *L. burdigalensis* and *Pa. toxostoma*

independently, in addition of the four common species.

We mapped these irreplaceable sites (for the six species pooled or independently)

on the river network, so as to test (i) whether or not specific areas harboured more

irreplaceable sites (e.g. upstream areas that are generally thought to be of high

conservation priority [42]) and (ii) if irreplaceable sites are spatially congruent among

species and, most notably, among common and rare species. In addition, we ran GLMs

(assuming a binomial error terms distribution) including whether or not a site has been

designated as an irreplaceable site at the 90% target as a binomial dependent variable,

and distance to the outlet of sampling sites and betweenness centrality values [43,44] for

each sites explanatory variables. Betweenness centrality is an index quantifying the

positional importance of each sampling site within the river basin [45,46]. The

significance of each term was tested at the $\alpha=0.05$ threshold.

Surrogacy in irreplaceable sites among species. We then estimated the levels of

surrogacy among species by calculating the percentage of the total number of alleles

observed for a given species that is secured by irreplaceable sites identified for another

species. Although surrogacy was calculated for all species pairs, we specifically focused

on rare species by calculating (i) the percentage of the total number of alleles observed

for rare species secured by all the irreplaceable sites identified for all the common

species, and (ii) the percentage of the total number of alleles observed for common

species secured by all the irreplaceable sites identified for the rare species.

Assessing the efficiency of SPC. The goal of this analysis was (i) to assess the efficiency

of SCP for preserving the overall levels of intraspecific genetic diversity observed at the

riverscape scale for the six species, and to (ii) compare the efficiency of SCP-based

conservation solutions to the efficiency of alternative conservation solutions selected on

the basis of traditional taxonomic criteria. We first computed an average accumulation

curve of "protected genetic diversity" by estimating the average cumulative percentage

of the total number of alleles (over all six species) secured by a cumulative number of

randomly selected sampling sites. Specifically, we increased the number of sites being

sampled from 1 to 92 with a continuous increment of one unit. At each incremental step,

we randomly selected sampling sites (without replacement) and we calculated the

percentage of the total number of alleles (over all species) secured by the random set of

sites. We repeated this random selection 1000 times for each incremental step and

averaged the percentages over these 1000 permutations to obtain the averaged

accumulation curve of protected genetic diversity. We then tested to which extent, for

an identical number of selected sites, the different conservation solutions we identified

through SCP (i.e. irreplaceable sites) secured higher percentages of genetic diversity

than those secured by a random selection of sites. Notably, we explored whether certain

conservation solutions (i.e. by considering irreplaceable sites identified for single

species or for the sets of common species, rare species and all the six species) were

more efficient than others to sustain a large percentage of the total number of alleles.

As in most conservation studies species diversity is the main conservation target [47],

we finally checked the efficiency of conservation solutions based on taxonomic criteria

for preserving the overall levels of genetic diversity of the six species of interest.

Specifically, we calculated the percentages of total number of alleles (over all species)

secured by sites (i) in which a number equal or greater than 10, 15 or 20 freshwater fish

species co-occur (the total number of fish species observed in the riverscape being 34

and the mean number of species observed at the site level being 12.77 +/- 0.55), and (ii)

in which the six fish species considered in our study co-occur. We finally compared the

efficiency of taxonomic-data-based conservation solutions to that of SCP-based

conservation solutions to assess if solutions based on taxonomic criteria are good

surrogates of the overall intraspecific genetic diversity of our species assemblage. Data

on fish species occurrence at the site level has been derived from [46] for 81 out of our

92 sampling sites.

Relationships between irreplaceable sites and indices of genetic diversity. To test the

ability of classical genetic indices to predict the propensity for a site to be irreplaceable

from a conservation viewpoint, we ran for each species Generalized Linear Models

(GLMs, with a binomial error terms distribution) including whether or not a site has

been designated as an irreplaceable site at the 90% target as a binomial dependent

variable, and AR, PA and D_{est} as explanatory variables. We tested the significance of

each term at the $\alpha=0.05$ threshold. Explanatory variables were centred and scaled, in

order to compare the relative strength of the predictors among species.

3. Results

Descriptive statistics. Overall, *Pa. toxostoma* (one of the two rare species) showed the

lowest within-site intraspecific genetic diversity estimates. Mean AR ranged from 2.114

for *Pa. toxostoma* to 5.821 for *Ph. phoxinus*, and mean PA ranged from 0.036 for *Pa.*

toxostoma to 0.162 for *L. burdigalensis* (Table S3). *Parachondrostoma toxostoma* also

showed the lowest mean D_{est} value (0.069), while the highest mean value was found for

B. barbatula (0.383; Table S3).

<u>Testing the suitability of SCP analyses for intraspecific genetic diversity</u>

Spatial patterns of genetic diversity. Spatial patterns of genetic diversity largely varied

and were actually poorly congruent among the four common fish species (Figure 1). As

extreme examples, hotspots of AR for *S. cephalus* where mainly found in the Western

part of the riverscape and on the core streams, whereas these same areas were identified

as coldspots of AR for *B. barbatula* (Figure 1A1-1A3). Similarly, hotspots of PA were

inversely related between *G. occitaniae* and *Ph. phoxinus* (Figure 1B2-1B4). Similar

conclusions were reached for D_{est} (Figure 1C). For instance, hotspots of D_{est} were

observed in opposite areas of the riverscape for the species pair *B. barbatula/Ph.*

phoxinus (Figure 1C3-1C4). As a consequence, the sign, slope and significance of

GLMSSNs explanatory variables strongly varied among species (Figure 1). This

incongruence in spatial patterns of genetic diversity among species was also reflected by

the low to moderate correlation coefficients measured among all possible pairs of

species and for each index of genetic diversity (i.e. *AR*, *PA* and D_{est}, Table S4). Indeed,

Pearson's correlation coefficients were lower than 0.6 for all comparisons but two (i.e.

between *B. barbatula*/*G. occitaniae* and between *Ph. phoxinus*/*B. barbatula* for *AR*, see

Table S4).

The influence of conservation targets and analytical strategies to identify irreplaceable

sites for genetic conservation. When species were analysed independently, we found

that the number of irreplaceable sites increased as the conservation target (i.e. the

percentage of alleles secured by irreplaceable sites) increased, with a steep increase for

conservation targets higher than 75% of the total number of alleles (Figure 2). However,

the percentage of irreplaceable sites strongly varied among species. For instance, for a

90% conservation target, the proportion of irreplaceable sites ranged from 3.61% of the

total number of sampled sites for *G. occitaniae* to 28.57% for *Ph. phoxinus* (Table S5;

Figure 2). For extreme conservation targets (100% of alleles to be secured), the

proportion of irreplaceable sites varied from 25.30% for *G. occitaniae* to 68.26% for

Ph. phoxinus (Table S5; Figure 2).

When alleles from the four common species were analysed in a single pooled

analysis, we similarly found that the proportion of irreplaceable sites increase as the

conservation target increases (Figure 2). Interestingly, we did not identify irreplaceable

sites for the 30% conservation goal, and only 3 irreplaceable sites were found for the

50% conservation goal (Figure 2). The proportion of irreplaceable sites increased

moderately to 17.39% for the 75% target, and then steeply increased for higher

conservation targets to reach 55.70% for the 90% target and 76.08% for the 100% target

(Figure 2). This later result suggests that almost all the riverscape should be protected to

reach high conservation targets when adopting a pooled strategy.

<u>A real conservation-oriented study using SCP approaches</u>

Identification of irreplaceable sites for genetic conservation. We first visually explored

the spatial distribution of irreplaceable sites. Overall, the localization of irreplaceable

sites in the riverscape strongly varied among species (Figure 3). We failed to identify

areas (e.g. upstream or downstream locations) clustering irreplaceable sites for any

species (Figure 3A-F). This apparent lack of clustering was statistically confirmed by

our generalized linear models (Table S6). Indeed, neither the distance from the outlet of

sampling sites nor the betweenness centrality of sampling sites were significant

predictors of the irreplaceability of sites for all species (the only exception being a

significant effect of distance from the outlet for *Ph. phoxinus*, Table S6).

Second, we tested whether conservation solutions found for each species were

spatially congruent among species. Over the six fish species, we identified forty-two

sites (45.65%) as irreplaceable at the 90% conservation target for at least one species

(Figure 4). Thirty-two of these forty-two sites were irreplaceable for at least one of the

four common species (Figure 4), and fourteen of the forty-two sites were irreplaceable

for at least one of the two rare species (Figure 4). Among the six species, only eight out

of these forty-two sites were irreplaceable for at least two species, and only one of these

sites was irreplaceable for three species (Figure 4).

Surrogacy in irreplaceable sites among species. Levels of surrogacy were generally low

to moderate, and strongly varied among species pairs (Table 1). For instance,

irreplaceable sites identified for *G. occitaniae* failed to secure the genetic diversity of

the two rare species, and covered only 18 to 43% of the total number of alleles of the

other common species. This indicates that irreplaceable sites identified for *G. occitaniae*

(i.e. the most widespread species) are poor surrogates for preserving the intraspecific

genetic diversity of other species (18.38% of surrogacy; Table 1). Conversely,

irreplaceable sites found for *Ph. phoxinus* and *L. burdigalensis* are better surrogates for

G. occitaniae, as 79.58% and 70.27% of the total number of alleles of *G. occitaniae* was

secured by irreplaceable sites identified for *Ph. phoxinus* and *L. burdigalensis*

respectively (Table 1). Overall, irreplaceable sites best sustaining genetic diversity of

other species were those identified for *B. barbatula*, which secured in average 68.28%

of the total number of alleles of other species (Table 1).

Interestingly, the thirty-two irreplaceable sites identified for the four common

species secured 81.79% and 90% of the total number of alleles of *L. burdigalensis* and

Pa. toxostoma respectively, suggesting that irreplaceable sites identified for a set of

common species can be good surrogates for intraspecific genetic diversity of rare

species. The fourteen irreplaceable sites identified for the two rare species secured a

total number of alleles ranging from 65.18% for *S. cephalus* to 79.40% for *B. barbatula*,

which is also not negligible given that conservation priorities are generally directed

towards rare species.

Assessing the efficiency of SPC. Overall, irreplaceable sites identified for single species

using SCP did not significantly secured higher percentages of genetic diversity than

randomly selected sites, excepting irreplaceable sites identified for *L. burdigalensis* and

B. barbatula (Figure 5), which were more efficient for preserving the overall genetic

diversity of all species than randomly selected sites. However, we found significantly

higher percentages of secured alleles than randomly selected sites on irreplaceable sites

identified through SCP for (i) the set of common species (ii) the set of rare species, and

(iii) all the six species (Figure 5).

Conservation solutions based on the co-occurrence of high numbers of species (i.e. sites

with numbers of co-occurring species equal or greater than 10, 15 or 20) did not secured

more alleles than those found for randomly selected sites, suggesting that raw

interspecific diversity (i.e. species richness) may not be a good surrogate of intraspecific

genetic diversity in our study (Figure 5). Nevertheless, we found that a higher

percentage of alleles would be secured by selecting sites on which the six species co-

occur than by randomly selecting sites in the riverscape (Figure 5).

Relationships between genetic indices and irreplaceable sites. For all species but *G.*

occitaniae and *Pa. toxostoma* (for which none of the variables were significant

predictors), *PA* was identified as the only variable that significantly predicted the

probability for a site to be identified as an irreplaceable site (Table S7). This probability

increased with the number of *PA* in a site.

4. Discussion

Intraspecific genetic diversity reflects the evolutionary and demographic history

of populations, and hence sustain the capacity of species and populations to cope with

environmental changes [14–17]. This biodiversity facet is the first that should respond

to global change [16], and is hence surprising that this biodiversity facet has yet rarely

integrated in dedicated optimization planning tools. We here fill this gap by

demonstrating that systematic conservation planning for intraspecific genetic diversity

is an efficient, yet complex, approach necessitating careful considerations.

From idiosyncratic distributions of genetic diversity...

525   Our results strongly suggest that the genetic diversity of the targeted species did

not follow a common spatial pattern, but rather species-specific (idiosyncratic) spatial

distributions. This conclusion holds true for all genetic diversity indices, and it

corroborates the few previous studies investigating simultaneously the spatial

distribution of genetic diversity at large spatial scales and for sympatric species (e.g.

[48,49]). This was however unexpected given that recent meta-analyses on freshwater

organisms demonstrated that α-IGD is generally higher in downstream than in upstream

areas [50], and that β-IGD tends to be higher in upstream than in downstream sections

[42]. Overall, these very general spatial patterns of intraspecific genetic diversity were

verified in our datasets. For instance, a negative relationship between allelic richness

and distance from the outlet is expected in freshwater organisms [50], and was actually

observed for three out of the six species considered in this study (Figure S2). However,

when using a precise and novel approach to map genetic diversity across the network,

we demonstrated that the distribution of cold- and hot-spots of α- and β-IGD was subtler

and idiosyncratic. This probably reflects interactions between recent colonization

histories, life-history-traits of species and the network structure, which are expected to

drive patterns of neutral genetic diversity in rivers [50–52].

...to systematic conservation planning for intraspecific genetic diversity.

The spatial mismatch in intraspecific genetic diversity among species probably

explains why we found that the level of congruence and surrogacy of irreplaceable sites

identified for each species was extremely low. For instance, there was an extremely low

proportion of irreplaceable sites that were common to two or three species (and never

more than three species; Figure 4). In the same way, we detected no clear patterns in the

spatial distribution of irreplaceable sites. Indeed, irreplaceable sites (for any of the six

species) were not particularly situated in specific locations such as upstream areas

and/or in areas of high connectivity (i.e. with high centrality values [45,46]), hence

suggesting that priority areas for the conservation of intraspecific genetic diversity of

species assemblages should not be restricted to specific areas in the riverscape. It is

however noteworthy that the greatly variable position of irreplaceable sites in the

riverscape found for the species we studied may reflect their own ecological niche and

spatial occupancy in the riverscape. Finally, the level of surrogacy among irreplaceable

sites was low to moderate, and never attained the 90% threshold we assumed when we

considered all species independently. Combined with our finding that the number of

irreplaceable sites can be relatively high for reasonable conservation targets (up to 46%

sites are identified as irreplaceable sites for at least one species at the conservation

target of 90%), we conclude that our ability to identify priority areas for intraspecific

genetic diversity may be highly species-specific and may depend on the capacity to

tackle the trade-off between the amount of genetic diversity to protect, and the extent of

priority areas we can realistically protect.

However, when surrogacy between all irreplaceable sites identified for the entire

set of common species and those identified for the rare species was tested, we reached

reasonable proportions of the total number of alleles to protect (~80-90%) for the two

rare species (i.e. *Pa. toxostoma* and *L. burdigalensis*). This result suggests that, in some

cases, genetic data obtained for a set of widely-distributed, "easier-to-sample" common species displaying varying life-history traits can be used for identifying protection areas for the intraspecific genetic diversity of other rare species that can be more problematic to sample. This would require, however, a good overlap between the spatial distributions of the common species used as surrogates and the spatial distributions of the targeted rare species, as well as a wide sampling design allowing representing the whole spatial occupancy of species in the riverscape. Failing to fulfil these two prerequisites may have a negative impact on conservation costs, as the number of irreplaceable sites necessary to preserve a predefined amount of genetic diversity of rare species would be higher than expected. In our demonstrative study for instance, thirty-four of the forty-two (81%) irreplaceable sites were sites in which at least four study species co-occur, and nineteen of these sites (54.3%) were sites in which the six species co-occur. Another important finding we report here is that selecting conservation areas based on traditional taxonomic criteria (i.e. by selecting sites with high species richness) is not an efficient strategy for preserving the overall genetic diversity of a set of species. This highlights the need for finding efficient ways for considering intraspecific genetic diversity in systematic conservation planning, as we and others [e.g. 9–13] proposed.

Overall, our results suggest that two different analytical strategies can be employed in real-case SCP studies aiming at preserving intraspecific genetic diversity of a species assemblage (i) identification of conservation areas for a set of rare species or (ii) identification of conservation areas for a set of representative common species. These two strategies have their own advantages and inconveniences. The first strategy may optimally preserve genetic diversity of rare species at competitive costs (e.g. only 14 irreplaceable sites to protect in our demonstrative study), but this at the expense of the genetic diversity of other sympatric species. Conversely, the second strategy may

optimally preserve genetic diversity of a set of common species while maintaining high

levels of genetic diversity for rare species, but this at a higher cost (e.g. 32 irreplaceable

sites to protect in our study). Whether to choose one of these two strategies will

therefore depend on many factors such as how difficult is the sampling of rare species

compared to common species, or the extent of the resources available for setting new

protected areas. We recommend adopting the second strategy when possible, since it

allows to simultaneously maintain genetic diversity from rare and common species.

Indeed, genetic diversity of common species is vital for ensuring ecosystem stability, as

it ultimately influences species interactions, population dynamics and ecosystem

functioning [53], and we argue that it should be considered in conservation plans. It is

noteworthy that further studies must be conducted to test if the two analytical strategies

we propose can be applicable across taxa and landscapes.

Although it was not a main goal of the study, we identified an alternative

strategy that may also be efficient for maintaining high levels of genetic diversity of a

species assemblage without requiring SCP analyses. Indeed, selecting sites where all the

studied species co-occur as conservation solutions allowed maintaining high levels of

genetic diversity for both common and rare species in our study, although this strategy

was less efficient than selecting irreplaceable sites found for common species through

SCP. The efficiency of this strategy also needs to be thoroughly tested across different

taxa and landscapes, as it can be a useful surrogate strategy in some cases (e.g. when

molecular data cannot be produced due to funding or technical limitations). It is

however noteworthy that such strategy requires highly reliable taxonomic data gathered

over the whole studied landscape.

Finally, we demonstrate that microsatellite-based genotypic data can be easily

used as direct input data in SCP to identify efficient conservation areas for intraspecific

genetic diversity of a species assemblage at the landscape scale. Relatively small

numbers of microsatellite markers such as those considered in our study (8 to 15) have

been proven to be sufficient for capturing the confounding genetic effects of recent past

demographic history, genetic drift and connectivity issues between populations

[27,50,51], and their efficiency for informing recent evolutionary processes can be equal

or higher than low to moderate numbers of SNP markers for instance (e.g. <300 SNPs;

[19]). However, microsatellite markers have the disadvantage of not being suitable for

drawing conclusions on the adaptive potential of populations, contrarily to other

markers derived at the genome scale. Given that our capacity to compile genetic

datasets at large spatial, temporal and taxonomic scales has greatly increased in recent

decades [54], a next methodological step may be the use of high-throughput techniques

to develop high numbers of markers informing both neutral and non-neutral processes

to assess whether SCP solutions obtained using different categories of markers allow

efficiently preserving both the whole genomic diversity of populations, and hence their

whole evolutionary history.

5. Conclusions

Our study provides novel, insightful and promising knowledge on the setting of

priority conservation areas for intraspecific diversity. It shows that systematic

conservation planning methods are useful and efficient objective tools for conservation

geneticists whose conservation solutions will strikingly depend on the species to be

conserved and the quantity of genetic information that managers aim at preserving in a

landscape. Given our results, we suggest that two strategies could be employed in real-

case conservation programs: (i) identification of priority conservation areas for a set of

rare species when resources allocated to conservation are scarce or (ii) identification of

priority conservation areas on the basis of the analysis of a set of representative

common species that may serve as "umbrellas" for rare sympatric species when

resources allocated to conservation are higher.

Our study also raises many additional questions that should be considered in the

near future. Among others, we believe that the next steps will be to formally identify

sound conservation targets for intraspecific diversity, to test whether neutral

intraspecific diversity appropriately mirrors quantitative and adaptive diversity [17], and

to quantify the influence of intraspecific diversity on ecosystem functioning and

services, so as to better evaluate the added value of preserving such a facet of

biodiversity [17,55].

Competing Interests statement

Authors declare not having any competing interest.

Author Contributions

The study was designed by SB, GL and IPV. IPV and SB wrote the manuscript with the

help of VH, GG, NP and GL. IPV, SB, GL and CV collected the samples. GL and CV

produced the genetic data. IPV, VH and SB conducted the population genetic and

statistical analyses.

Acknowledgements

We are very grateful to all the ONEMA staff that contributed to the field sampling, in

particular Laurence Blanc. Leslie Faggiano, Charlotte Evangelista, Christine Lauzeral,

Loïc Tudesque, Roselyne Étienne, Sébastien Villéger, Julien Cucherousset, Vincent

Dubut, Alain Blanchet and South-Western France ONEMA staff are thanked for their

help in the field. Roselyne Étienne and Vincent Dubut are also thanked for their help

with molecular analyses. Lisa Fourtune is acknowledged for having calculated the

betweenness centrality values of our river network. We also thank Marie-Hélène Lizée

for her help with GIS tools and Camille Pagès and Jérôme Prunier for their comments.

Further, the authors thank the "Génopole Toulouse" for help with genotyping and Keoni

Saint-Pée for correcting the English. This work has been done in two research units

(EDB & SETE) that are part of the "Laboratoire d'Excellence" (LABEX) entitled

TULIP (ANR-10-LABX-41).

Funding

IPV was financially supported by a MESR ("Ministère de l'Enseignement Supérieur et

de la Recherche") PhD scholarship during this study. VH was funded by the National

Environmental Research Program Northern Australia Hub and a "Ramon y Cajal"

contract (RYC-2013-13979) funded by the Spanish government. This study is part of

the European project "IMPACT". This project has been carried out with financial

support from the Commission of the European Communities, specific RTD programme

"IWRMNET".

References

1.   Watson JEM, Dudley N, Segan DB, Hockings M. 2014 The performance and
potential of protected areas. *Nature* **515**, 67–73.
691
692   2.   Filipe AF, Marques TA, Tiago P, Ribeiro F, Da Costa LM, Cowx IG, Collares-
693   Pereira MJ. 2004 Selection of priority areas for fish conservation in Guadiana river
694   basin, Iberian Peninsula. *Conserv. Biol.* **18**, 189–200.
695
696   3.   Kirkpatrick JB. 1983 An iterative method for establishing priorities for the
697   selection of nature reserves: An example from Tasmania. *Biol. Conserv.* **25**, 127–134.
698

699   4.   Margules CR, Pressey RL. 2000 Systematic conservation planning. *Nature* **405**,
700   243–253.

701
702   5.   Vane-Wright RI, Humphries CJ, Williams PH. 1991 What to protect?—
703   Systematics and the agony of choice. *Biol. Conserv.* **55**, 235–254.

704
705   6.   Faith DP. 1992 Conservation evaluation and phylogenetic diversity. *Biol.*
706   *Conserv.* **61**, 1–10.

707
708   7.   Asmyhr MG, Linke S, Hose G, Nipperess DA. 2014 Systematic conservation
709   planning for groundwater ecosystems using phylogenetic diversity. *PLoS ONE* **9**,
710   e115132.

711
712   8.   Buerki S, Callmander MW, Bachman S, Moat J, Labat J-N, Forest F. 2015
713   Incorporating evolutionary history into conservation planning in biodiversity hotspots.
714   *Philos. Trans. R. Soc. B Biol. Sci.* **370**, 20140014–20140014.

715
716   9.   Carvalho SB, Velo-Antón G, Tarroso P, Portela AP, Barata M, Carranza S,
717   Moritz C, Possingham HP. 2017 Spatial conservation prioritization of biodiversity
718   spanning the evolutionary continuum. *Nat. Ecol. Evol.* **1**, 0151.

719
720   10.   Thomassen HA *et al.* 2011 Mapping evolutionary process: a multi-taxa approach
721   to conservation prioritization. *Evol. Appl.* **4**, 397–413.

722
723   11.   Diniz-Filho JAF *et al.* 2012 Planning for optimal conservation of geographical
724   genetic variability within species. *Conserv. Genet.* **13**, 1085–1093.

725
726   12.   Hanson JO, Rhodes JR, Riginos C, Fuller RA. 2017 Environmental and
727   geographic variables are effective surrogates for genetic variation in conservation
728   planning. *Proc. Natl. Acad. Sci.*, 201711009.

729
730   13.   Hermoso V, Kennard MJ, Schmidt DJ, Bond N, Huey JA, Mondol RK,
731   Jamandre BW, Hughes JM. 2016 Species distributions represent intraspecific genetic
732   diversity of freshwater fish in conservation assessments. *Freshw. Biol.* **61**, 1707–1719.

733
734   14.   Hoffmann AA, Sgrò CM. 2011 Climate change and evolutionary adaptation.
735   *Nature* **470**, 479–485.

736
737   15.   Carroll SP, Jorgensen PS, Kinnison MT, Bergstrom CT, Denison RF, Gluckman
738   P, Smith TB, Strauss SY, Tabashnik BE. 2014 Applying evolutionary biology to
739   address global challenges. *Science* **346**, 1245993–1245993.

740
741   16.   Rey O, Danchin E, Mirouze M, Loot C, Blanchet S. 2016 Adaptation to global
742   change: A transposable element–epigenetics perspective. *Trends Ecol. Evol.* **31**, 514–
743   526.

744
745   17.   Mittell EA, Nakagawa S, Hadfield JD. 2015 Are molecular markers useful
746   predictors of adaptive potential? *Ecol. Lett.* **18**, 772–778.

748   18.   Pritchard JK, Stephens M, Donnelly P. 2000 Inference of population structure
749   using multilocus genotype data. *Genetics* **155**, 945–959.
750
751   19.   Putman AI, Carbone I. 2014 Challenges in analysis and interpretation of
752   microsatellite data for population genetic studies. *Ecol. Evol.* **4**, 4399-4428.
753
754   20.   Blanchet S, Prunier JG, De Kort H. 2017 Time to go bigger: Emerging patterns
755   in macrogenetics. *Trends Genet.* **33**, 579–580.
756
757   21.   Moritz C. 1994 Defining 'Evolutionarily Significant Units' for conservation.
758   *Trends Ecol. Evol.* **9**, 373–375.
759
760   22.   Waples RS *et al.* 2001 Characterizing diversity in salmon from the Pacific
761   Northwest. *J. Fish Biol.* **59**, 1–41.
762
763   23.   Frankham R, Briscoe DA, Ballou JD. 2002 *Introduction to conservation
764   genetics*. Cambridge, UK; New York: Cambridge University Press.
765
766   24.   Neel MC, Cummings MP. 2003 Effectiveness of conservation targets in
767   capturing genetic diversity. *Conserv. Biol.* **17**, 219–229.
768
769   25.   Kottelat M, Freyhof J. 2007 *Handbook of European freshwater fishes*. Cornol,
770   Switzerland: Publications Kottelat.
771
772   26.   Poulet N, Beaulaton L, Dembski S. 2011 Time trends in fish populations in
773   metropolitan France: insights from national monitoring data. *J. Fish Biol.* **79**, 1436–
774   1452.
775
776   27.   Paz-Vinas I, Quéméré E, Chikhi L, Loot G, Blanchet S. 2013 The demographic
777   history of populations experiencing asymmetric gene flow: combining simulated and
778   empirical data. *Mol. Ecol.* **22**, 3279–3291.
779
780   28.   Paz-Vinas I, Comte L, Chevalier M, Dubut V, Veyssiere C, Grenouillet G, Loot
781   G, Blanchet S. 2013 Combining genetic and demographic data for prioritizing
782   conservation actions: insights from a threatened fish species. *Ecol. Evol.* **3**, 2696–2710.
783
784   29.   Aljanabi SM, Martinez I. 1997 Universal and rapid salt-extraction of high
785   quality genomic DNA for PCR-based techniques. *Nucleic Acids Res.* **25**, 4692–4693.
786
787   30.   Van Oosterhout C, Hutchinson WF, Wills DPM, Shipley P. 2004 *micro-checker*:
788   software for identifying and correcting genotyping errors in microsatellite data. *Mol.
789   Ecol. Notes* **4**, 535–538.
790
791   31.   Jombart T. 2008 *adegenet*: a R package for the multivariate analysis of genetic
792   markers. *Bioinformatics* **24**, 1403–1405.
793
794   32.   Rousset F. 2008 *genepop'007*: a complete re-implementation of the genepop
795   software for Windows and Linux. *Mol. Ecol. Resour.* **8**, 103–106.
796

797   33.   Szpiech ZA, Jakobsson M, Rosenberg NA. 2008 *ADZE*: a rarefaction approach
798   for counting alleles private to combinations of populations. *Bioinforma. Oxf. Engl.* **24**,
799   2498–2504.
800
801   34.   Petit RJ, El Mousadik A, Pons O. 1998 Identifying populations for conservation
802   on the basis of genetic markers. *Conserv. Biol.* **12**, 844–855.
803
804   35.   Kalinowski ST. 2004 Counting alleles with rarefaction: Private alleles and
805   hierarchical sampling designs. *Conserv. Genet.* **5**, 539–543.
806
807   36.   Winter DJ. 2012 *mmod*: an R library for the calculation of population
808   differentiation statistics. *Mol. Ecol. Resour.* **12**, 1158–1160.
809
810   37.   Jost L. 2008 G(st) and its relatives do not measure differentiation. *Mol. Ecol.* **17**,
811   4015–4026.
812
813   38.   Peterson E, Ver Hoef JMV. 2014 *STARS*: An ArcGIS toolset used to calculate
814   the spatial information needed to fit spatial statistical models to stream network data. *J.*
815   *Stat. Softw.* **56**, 1–17.
816
817   39.   Ver Hoef J, Peterson E, Clifford D, Shah R. 2014 *SSN*: An R Package for Spatial
818   Statistical Modeling on Stream Networks. *J. Stat. Softw.* **56**, 1–43.
819
820   40.   Ball IR, Possingham HP, Watts M. 2009 Marxan and relatives: Software for
821   spatial conservation prioritisation. In *Spatial conservation prioritisation: Quantitative*
822   *methods and computational tools* (eds A Moilanen, KA Wilson, HP Possingham), pp.
823   185–195. Oxford Iniversity Press, Oxford, UK.
824
825   41.   Hermoso V, Kennard MJ. 2012 Uncertainty in coarse conservation assessments
826   hinders the efficient achievement of conservation goals. *Biol. Conserv.* **147**, 52–59.
827
828   42.   Finn DS, Bonada N, Múrria C, Hughes JM. 2011 Small but mighty: headwaters
829   are vital to stream network biodiversity at two levels of organization. *J. North Am.*
830   *Benthol. Soc.* **30**, 963–980.
831
832   43.   Freeman LC. 1977 A Set of measures of centrality bsed on betweenness.
833   *Sociometry* **40**, 35.
834
835   44.   Estrada E, Bodin Ö. 2008 Using network centrality measures to manage
836   landscape connectivity. *Ecol. Appl.* **18**, 1810–1825.
837
838   45.   Altermatt F. 2013 Diversity in riverine metacommunities: a network perspective.
839   *Aquat. Ecol.* **47**, 365–377.
840
841   46.   Fourtune L, Paz-Vinas I, Loot G, Prunier JG, Blanchet S. 2016 Lessons from the
842   fish: a multi-species analysis reveals common processes underlying similar species-
843   genetic diversity correlations. *Freshw. Biol.* **61**, 1830–1845.
844
845   47.   Myers N, Mittermeier RA, Mittermeier CG, da Fonseca GAB, Kent J. 2000
846   Biodiversity hotspots for conservation priorities. *Nature* **403**, 853–858.

848   48.   Fortuna MA, Albaladejo RG, Fernandez L, Aparicio A, Bascompte J. 2009
849   Networks of spatial genetic variation across species. *Proc. Natl. Acad. Sci.* **106**, 19044–
850   19049.
851

852   49.   Taberlet P *et al.* 2012 Genetic diversity in widespread species is not congruent
853   with species richness in alpine plant communities. *Ecol. Lett.* **15**, 1439–1448.
854

855   50.   Paz-Vinas I, Loot G, Stevens VM, Blanchet S. 2015 Evolutionary processes
856   driving spatial patterns of intraspecific genetic diversity in river ecosystems. *Mol. Ecol.*
857   **24**, 4586–4604.
858

859   51.   Paz-Vinas I, Blanchet S. 2015 Dendritic connectivity shapes spatial patterns of
860   genetic diversity: a simulation-based study. *J. Evol. Biol.* **28**, 986–994.
861

862   52.   Thomaz AT, Christie MR, Knowles LL. 2016 The architecture of river networks
863   can drive the evolutionary dynamics of aquatic populations. *Evolution* **70**, 731–739.
864

865   53.   Mimura M *et al.* 2017 Understanding and monitoring the consequences of
866   human impacts on intraspecific variation. *Evol. Appl.* **10**, 121–139.
867

868   54.   Pauls SU *et al.* 2014 Integrating molecular tools into freshwater ecology:
869   developments and opportunities. *Freshw. Biol.* **59**, 1559–1576.
870

871   55.   Siefert A *et al.* 2015 A global meta-analysis of the relative extent of intraspecific
872   trait variation in plant communities. *Ecol. Lett.* **18**, 1406-1419.
873

 Figure captions

Figure 1

Spatial distribution of observed (coloured circles) and interpolated (coloured lines)

values of AR, PA and D_{est} (A, B, C respectively in the figure) for *Squalius cephalus*,

Gobio occitaniae, *Barbatula barbatula,* and *Phoxinus phoxinus* (1, 2, 3, 4 respectively

in the figure) obtained with GLMSSNs. The width of coloured lines is inversely related

to the prediction standard error. The cursor on the vertical coloured scale indicates the

mean value of AR, PA and D_{est}. The slope (β) of each explanatory variable (i.e.

topological distance to the outlet, longitude and latitude) and its significance is also

reported. N.I. indicates that the explanatory variable has not been included in the model;

* p-value <0.05; ** p-value <0.01; *** p-value <0.001.

Figure 2

Percentage of irreplaceable sites identified by Marxan for conservation targets of 50, 75,

90 and 100% of the total number of alleles present in the river basin for each common

species and for a pooled analysis in which all alleles from all common species were

pooled together. N_{SITES} represents the number of sites included in the Marxan analyses.

Figure 3

Irreplaceable sites that have been identified for each species by Marxan for preserving

90% of the total number of alleles present in the river basin (red-filled circles).

Figure 4

Irreplaceable sites that have been identified by Marxan for at least one (unicoloured-

filled points), two (black dotted circles surrounding bicoloured points) or three (black

bolded circle surrounding a tricoloured point) species, assuming a conservation target of

90% of the total number of alleles present in the river basin when considering the six

species.

Figure 5

Mean percentage of the total number of alleles observed at the riverscape scale for all

the six species that are secured by (i) a cumulative number of randomly selected

sampling sites (blue line; average accumulation curve obtained under 1000

permutations of the sampling sites sites; grey vertical lines: standard error), (ii)

irreplaceable sites identified using SCP for each of the six species independently (red

dots), (iii) irreplaceable sites identified using SCP for different sets of species (red

squares), and (iv) sites identified on the basis of freshwater fish species co-occurrence

data (black dots).

TABLE 1: Table reporting for each species the percentages of their total number of alleles that are secured by the sampling sites having

been identified as irreplaceable sites for the other species (considering the 90% conservation target).

Proportion of alleles from (below) that fell within irreplaceable sites from (right):	*Squalius cephalus* *n=7*	*Gobio occitaniae* *n=3*	*Barbatula barbatula* *n=9*	*Phoxinus phoxinus* *n=18*	*Leuciscus burdigalensis* *n=11*	*Parachondrostoma toxostoma* *n=3*	Common species *n=32*	Rare species *n=14*
Squalius cephalus	*90.00*	17.66	61.52	56.52	61.60	39.73	—	65.18
Gobio occitaniae	65.60	*90.00*	66.76	79.58	70.27	50.97	—	73.40
Barbatula barbatula	49.79	31.65	*90.00*	79.57	76.78	38.95	—	79.40
Phoxinus phoxinus	48.25	42.60	68.43	*90.00*	63.49	44.61	—	68.88
Leuciscus burdigalensis	44.73	0	66.13	67.09	*90.00*	36.42	81.79	—
Parachondrostoma toxostoma	78.55	0	78.55	50.00	67.14	*90.00*	90.00	—
Average over all other species	*55.58*	*18.38*	*68.28*	*66.55*	*67.86*	*42.14*		

A *S. cephalus*

Legend
○ Non-irreplaceable site
● Irreplaceable site

0 40 80 km

N

B *G. occitaniae*

C *B. barbatula*

D *Ph. phoxinus*

E *L. burdigalensis*

F *Pa. toxostoma*

Non-irreplaceable site
S. cephalus (7)
G. occitaniae (3)
B. barbatula (9)
Ph. phoxinus (18)
Pa. toxostoma (3)
L. burdigalensis (11)

Irreplaceable site
(for at least one species)

Irreplaceable site (for at least two species)

Irreplaceable site (for at least three species)

N

0 40 80 km