Iroki: automatic customization for phylogenetic trees

Ryan M. Moore[1*], Amelia O. Harrison[2], Sean M. McAllister[4], Rachel Marine[3], Clara Chan[4],

and K. Eric Wommack[1]

[1]Center for Bioinformatics & Computational Biology, University of Delaware, Newark, DE,

USA

[2]Department of Entomology and Wildlife Ecology, University of Delaware, Newark, DE,

USA

[3]Centers for Disease Control, Atlanta, Georgia, USA

[4]School of Marine Science and Policy, University of Delaware, Newark, DE, USA

Corresponding author's information

*To whom correspondence should be addressed

Address: Delaware Biotechnology Institute, 15 Innovation Way, Newark, Delaware

19711

(Tel): (302) 831-4362

(Fax): (302) 831-3447

(E-mail): moorer@udel.edu

Abstract

Background

Phylogenetic trees are an important analytical tool for examining species and community

diversity and the evolutionary history of species. In the case of microorganisms,

decreasing sequencing costs have enabled researchers to generate ever-larger sequence

datasets, which in turn have begun to fill gaps in the evolutionary history of microbial

groups. However, phylogenetic analyses of large sequence datasets present challenges to

extracting meaningful trends from complex trees. Scientific inferences made by visual

inspection of phylogenetic trees can be simplified and enhanced by customizing various

parts of the tree, including label color, branch color, and other features. Yet, manual

customization is time-consuming and error prone, and programs designed to assist in

batch tree customization often require programming experience. To address these

limitations, we developed Iroki, a program for fast, automatic customization of

phylogenetic trees. Iroki allows the user to incorporate information on a broad range of

metadata for each experimental unit represented in the tree.

Results

Iroki was applied to four existing microbial sequence datasets to demonstrate its utility in

data exploration and presentation. Iroki was used to highlight connections between viral

phylogeny and host taxonomy, to explore the abundance of microbial groups across

samples of cattle hide, to examine short-term temporal dynamics of virioplankton

communities, and to search for trends in the biogeography of Zetaproteobacteria.

Conclusions

Iroki is an easy-to-use web app and command line application for fast, automatic

customization of phylogenetic trees based on user-provided categorical or continuous

metadata. Iroki allows for rapid hypothesis testing through visualizing custom

phylogenetic trees, streamlining the process of phylogenetic data exploration and

presentation.

Availability

Iroki can be accessed through a web app or via installation through RubyGems, from

source, or through the Iroki Docker image. All source code and documentation is

available under the GPLv3 license at https://github.com/mooreryan/iroki. The Iroki web-

app is accessible at www.iroki.net or through the Virome portal

(http://virome.dbi.udel.edu), and its source code is released under GPLv3 license at

https://github.com/mooreryan/iroki_web. The Docker image can be found here:

https://hub.docker.com/r/mooreryan/iroki.

Keywords

Phylogeny, visualization, sequence analysis, bioinformatics, metagenomics

Iroki: automatic customization for phylogenetic trees

Background

Studies in microbial ecology often use phylogenetic trees as a means for assessing the

diversity and evolutionary history of microorganisms. As the cost of sequencing has

declined, researchers have been able to gather ever-larger sequence datasets. While large

sequence datasets have begun to fill in the gaps in the evolutionary history of microbial

groups [1–5]; they have also posed new analytical challenges as extracting meaningful

trends within such highly dimensional datasets can be cumbersome. In particular,

scientific inferences made by visual inspection of phylogenetic trees can be simplified and

enhanced by customizing various parts of the tree including label and branch color,

branch width, and other features. Though many tree visualization packages allow for

manual modifications [6–9], the process can be time consuming and error prone

especially when the tree contains many nodes. While a handful of existing programs

address the issue of tree visualization, most are not capable of batch customization and

those that do often require programming experience [10–13].

Iroki, a program for fast, automatic customization of phylogenetic trees, was developed to

address these limitations and enable users to incorporate information on a broad range of

metadata for each experimental unit represented in the tree. Iroki is available for use

through a web interface at www.iroki.net, through the Virome portal

(http://virome.dbi.udel.edu), and through a UNIX command line tool. Results are saved in

the widely used Nexus format with color metadata tailored for use with FigTree [8] (a

freely available and efficient tree viewer).

Implementation

Iroki enhances visualization of phylogenetic trees by coloring node labels and branches

according to categorical metadata criteria or numerical data such as abundance

information. Iroki can also rename nodes in a batch process according to user

specifications so that node names are more descriptive. A tree file in Newick format

containing a phylogenetic tree is always required. Additional required input files depend

on the operation(s) desired. Coloring functions require a color map or a biom [14] file.

Node renaming functions require a name map. The color map, name map, and biom files

are created by the user and, along with the Newick file, form the inputs for Iroki.

Explicit tree coloring

Iroki's principle functionality involves coloring node labels and/or branches based on

information provided by the user in the color map. The color map text file contains either

two or three tab-delimited columns depending on how branches and labels are to be

colored. Two columns, pattern and color, are used when labels and branches are to have

the same color. Three columns, pattern, label color, and branch color, are used when

branches and labels are to have different colors. Patterns are searched against node labels

either as regular expressions or exact string matches.

Entries in the color column can be any of the 657 named colors in the R programming

language [15] (e.g., skyblue, tomato, goldenrod2, lightgray, black) or any valid

hexadecimal color code (e.g., #FF78F6). In addition, Iroki provides a 19 color palette

with complementary colors based on Kelly's color scheme for maximum contrast [16].

Nodes in the tree that are not in the color map will remain black.

Depending on user-specified options, a pattern match to node label(s) will trigger coloring

of the label and/or the branch directly connected to that label. Inner branches will be

colored to match their descendent branches if all descendants are the same color,

allowing quick identification of common ancestors and clades that share common

metadata.

Tree coloring based on numerical data

Iroki provides the ability to generate color gradients based on numerical data, such as

absolute or relative abundance, from a tab-delimited biom format file. Single-color

gradients use color saturation to illustrate numerical differences, with nodes at a higher

level being more saturated than those at a lower level. For example, highly abundant

nodes will be represented by more highly saturated colors. Two-color gradients show

numerical differences through both color mixing and luminosity. Additionally, the biom

file may specify numerical information for one group (e.g., abundance in a particular

sample) or for two groups (e.g., abundance in the treatment group vs. abundance in the

control group). For biom files with one group, single- or two-color gradients may be used.

However, biom files specifying two-group metadata may only use the two-color gradient.

Renaming nodes

Some packages for generating phylogenetic trees (RAxML [17], PHYLIP [18], etc.) require

node names to be ten characters or less. Name restrictions present challenges to scientific

interpretation of phylogenetic trees. Iroki's renaming function uses a two-column, tab-

delimited name map to associate current node names, exactly matching those in the tree

file, with new names. The new name column has no restrictions on name length or

character type. Iroki ensures name uniqueness by appending integers to the ends of

names, if necessary.

Combining the color map, name map, and biom files

Iroki can be used to make complex combinations of customizations by combining the

color map, name map, and biom files. For example, a biom file can be used to apply a

color gradient based on numerical data to the labels of a tree, a color map can be used to

separately color the branches based on user-specified conditions, and a name map can be

used to rename nodes in a single command or web request. Iroki follows a specific order

of precedence when applying multiple customizations. First, the color gradient inferred

from the biom file is applied. Next, the color map is applied to specified labels or

branches, overriding the gradient applied in the previous step if necessary. Finally, the

name map is used to map current names to the new names (Fig. 1).

Output

Iroki outputs the modified tree in the Nexus format. When building the phylogenetic tree,

FigTree uses the Nexus format file and interprets the color metadata output of Iroki.

Results & Discussion

Global diversity of bacteriophage

Viruses are the most abundant biological entity on Earth, providing an enormous reservoir

of genetic diversity, driving evolution of their hosts, influencing composition of microbial

communities, and affecting global biogeochemical cycles [19,20]. The current viral

taxonomic system is based on a suite of physical characteristics of the virion rather than

on genome sequences. The phage proteomic tree, created to provide a genome-based

taxonomic system for bacteriophage classification [21], was recently updated to include

hundreds of new phage genomes from the Phage SEED reference database [22], as well as

long assembled contigs from viral shotgun metagenomes (viromes) collected from the

Chesapeake Bay (SERC) [23] and the Mediterranean Sea [24].

Taxonomy and host information metadata was collected for the viral genome sequences, a

color map was created to assign colors based on viral family and host phyla, and Iroki was

used to add color metadata to branches and labels of the phage proteomic tree. Since a

large number of colors were required on the tree, Iroki's Kelly color palette was used to

provide clear color contrasts. The tree was rendered with FigTree (Fig. 2).

Adding color to the phage proteomic tree with Iroki shows trends in the data that would

be difficult to discern without color. Uncultured phage contigs from the SERC and

Mediterranean viromes make up a large portion of all phage sequences shown on the tree,

and are widely distributed among known phage. In general, viruses in the same family

claded together, e.g., branch coloring highlights large groups of closely related

Siphoviridae and Myoviridae. This label-coloring scheme also shows that viruses infecting

hosts within same phylum are, in general, phylogenetically similar. For example, viruses

within one of the multiple large groups of Siphoviridae across the tree infect almost

exclusively host species within the same phylum, e.g., Siphoviridae infecting

Actinobacteria clade away from Siphoviridae infecting Firmicutes or Proteobacteria.

Bacterial community diversity and prevalence of E. coli in beef cattle

Shiga toxin-producing *Escherichia coli* (STEC) are dangerous human pathogens that

colonize the lower gastrointestinal tracts of cattle and other ruminants. STEC-

contaminated beef and STEC shed in the feces of these animals are major sources of

foodborne illness. To identify possible interactions between STEC populations and the

commensal cattle microbiome, a recent study examined diversity of the bacterial

community associated with beef cattle hide [25]. Fecal and hide samples were collected

over twelve weeks and SSU rRNA amplicon libraries were constructed and analyzed by

Illumina sequencing [26]. The study indicated that the community structure of hide

bacterial communities was altered when the hides were positive for STEC contamination.

Iroki was used to visualize changes in the relative abundance of each cattle hide bacterial

OTU according to the presence or absence of STEC. A Mann-Whitney U test comparing

OTU abundance between STEC positive and STEC negative samples was performed, and

those bacterial OTUs showing a significant change in relative abundance ($p < 0.5$) were

placed on a phylogenetic tree according to the 16S rRNA sequence. Branches of the tree

were colored based on whether there was a significant change in relative abundance with

STEC contamination (red: $p <= 0.05$, blue: $p > 0.05$). Node labels were colored along a

blue-green color gradient representing the abundance ratio of OTUs between samples

with STEC (blue) and without (green). Additionally, label luminosity was determined

based on overall abundance (lighter: less abundant, darker: more abundant) (Fig. 3). Iroki

makes it clear that most OTUs on the tree showed a significant difference in abundance

(branch coloring) between STEC positive and STEC negative samples (node coloring).

Furthermore, we can see that most OTUs are at low abundance with only a few highly

abundant OTUs (label luminosity). The color gradient added by Iroki allows us to see that

the abundant OTUs were evolutionarily distant from one another and thus spread out

across many phylogenetic groups.

Iroki can be used to quickly test hypotheses without investing a large amount of time

annotating trees manually. A UPGMA tree was created based on unweighted UniFrac

distance [27] between 356 bacterial community profiles based on SSU rRNA amplicon

sequences from cattle hide and feces samples (Fig. 4). Iroki was used to evaluate

similarities in sample bacterial communities according to the sampling location. Iroki

colored branches based on whether the sample originated from feces (blue) or from hide

(red). The coloring added by Iroki shows a clear partitioning of bacterial communities on

the tree based on their sampling location (hide or feces). However, four fecal samples

claded with hide samples, and two hide samples claded with fecal samples, making them

be good candidates for more in-depth examination. Additionally, Iroki was used to

illustrate a correlation between one of the most abundant bacterial families,

Ruminococcaceae, and sampling location. Iroki colored node labels with a color gradient

based on Ruminococcaceae family abundance, utilizing both a single color gradient (Fig.

4A) and a two color gradient (Fig. 4B). Custom trees were visualized using FigTree.

Iroki's automatic color gradient and ability to label branches and nodes based on different

criteria clearly show that Ruminococcaceae is more abundant in fecal samples than in

hide samples.

Short-term dynamics of virioplankton

The gene encoding Ribonuclotide reductase (RNR) is common within viral genomes and

thus can be used as a marker gene for studying viral diversity. Moreover, RNR

polymorphism is predictive of some of the biological and ecological features of viral

populations [28]. A mesocosm experiment examined the short-term dynamics of phage

populations using RNR amplicon sequences, specifically, sequences of class II RNRs of

bacteriophages infecting cyanobacterial hosts. A phylogenetic tree was created from the

Cyano II RNR amplicon sequences and Iroki was used to color nodes and branches based

on the time point (0 h, 6 h, 12 h) at which each amplicon sequence was observed. The

customized tree was then visualized using FigTree (Fig. 5). Iroki's coloring shows that no

phylogenetic clade was dominated by OTUs observed in any particular time point; rather,

time points were spread relatively evenly across clades. This analysis demonstrates Iroki's

utility for exploring sequence datasets, allowing the researcher to quickly and easily test

hypotheses.

Phylogeny of Zetaproteobacteria within a biogeographic context

Biogeographical studies assess the distribution of an organism's biodiversity across space

and time. The extent to which microorganisms exhibit biogeography is an open question

in microbial ecology. The isolated nature of the microbial communities associated with

deep-ocean hydrothermal vents, provides an ideal system for studying the biogeography of

microbes. In particular, iron-oxidizing bacteria have been shown to thrive in vent fluids,

sediments, and iron-rich microbial mats associated with the vents. Globally, iron-

oxidizing bacteria make significant contributions to the iron and carbon cycles. A recent

study analyzed multiple SSU rRNA clone libraries to investigate the biogeography of

Zetaproteobacteria, a phyla containing many iron-oxidizing bacterial species, between

three sampling regions of the Pacific Ocean (central Pacific—Loihi seamount, western

Pacific—Southern Mariana Trough, and southern Pacific (Vailulu'u Seamount/Tonga

Arc/East Lau Spreading Center/Kermadec Arc) [29]. Sequences were aligned and a

phylogeny was inferred as described in [29]. Iroki was used to examine the relationship

between sampling location and phylotype by adding branch and label color based on

geographic location and renaming original node labels with OTU and location metadata.

The custom tree was visualized using FigTree (Fig. 6). In some cases, OTUs contained

sequences from only one sampling location (e.g., OTUs 12, 15, and 16), whereas other

OTUs are distributed among more than one sampling location (e.g., OTUs 1, 2, and 4).

Often, sequences samples from the same geographic location are in the same phylotype

despite being members of different OTUs (e.g., OTUs 10 and 19).

Availability and requirements

A web-based version of Iroki can be accessed online at www.iroki.net or through the

Virome portal (http://virome.dbi.udel.edu/). For users who wish to run Iroki locally, a

command line version of the program is installable via RubyGems, from GitHub

(https://github.com/mooreryan/iroki). A Docker image is available for users who desire the

flexibility of the command line tool, but do not want to install Iroki or manage its

dependencies (https://hub.docker.com/r/mooreryan/iroki). Docker is a convenient method

for packaging an application with all of its dependencies that is guaranteed to run the

same regardless of the user's environment [30,31]. The README provided with the source

code provides detailed instructions for setting up and running Iroki. Further

documentation and tutorials can be found at the Iroki Wiki

(https://github.com/mooreryan/iroki/wiki).

License

Iroki and its associated programs are released under the GNU General Public License

version 3 [32].

Conclusions

Iroki is a command line program and web app for fast, automatic customization of large

phylogenetic trees based on user specified configuration files describing categorical or

continuous metadata information. The output files include Nexus tree files with color

metadata tailored specifically for use with FigTree. Various example datasets from

microbial ecology studies were analyzed to demonstrate Iroki's utility. In each case, Iroki

simplified the processes of data exploration, data presentation, and hypothesis testing.

Iroki provides a simple and convenient way to rapidly customize phylogenetic trees,

especially in cases where the tree in question is too large to annotate manually or in

studies with many trees to annotate.

List of Abbreviations

OTU*:* operational taxonomic

RNR*:* Ribonucleotide reductase

STEC: Shiga-toxengenic *Escherichia coli*

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and materials

Data and code used to generate figures are available on GitHub at

https://github.com/mooreryan/iroki_manuscript_data

Funding

This project was supported by the USDA National Institute of Food and Agriculture award

number 2012-68003-30155 and the National Science Foundation Advances in

Bioinformatics program (award number DBI_1356374).

Competing Interests

The authors declare that they have no competing interests.

Authors' contributions

RMM and SMM conceived the project. RMM wrote the manuscript and implemented

Iroki. AOH and RM processed and analyzed Cyano II amplicons. All authors read,

edited, and approved the final manuscript.

Acknowledgements

We would like to acknowledge Daniel J. Nasko and Jessica M. Chopyk for their work on

the phage proteomic tree, and Barbra D. Ferrell for editing the manuscript.

References

1. Lan Y, Rosen G, Hershberg R. Marker genes that are less conserved in their sequences

are useful for predicting genome-wide similarity levels between closely related prokaryotic

strains. Microbiome. 2016;4:18.

2. Larkin A a, Blinebry SK, Howes C, Lin Y, Loftus SE, Schmaus CA, et al. Niche

partitioning and biogeography of high light adapted Prochlorococcus across taxonomic

ranks in the North Pacific. ISME J. 2016;1–13.

3. Simister RL, Deines P, Botté ES, Webster NS, Taylor MW. Sponge-specific clusters

revisited: A comprehensive phylogeny of sponge-associated microorganisms. Environ.

Microbiol. 2012;14:517–24.

4. Wu Z, Yang L, Ren X, He G, Zhang J, Yang J, et al. Deciphering the bat virome catalog

to better understand the ecological diversity of bat viruses and the bat origin of emerging

infectious diseases. ISME J. 2016;10:609–20.

5. Müller AL, Kjeldsen KU, Rattei T, Pester M, Loy A. Phylogenetic and environmental

diversity of DsrAB-type dissimilatory (bi)sulfite reductases. ISME J. 2015;9:1152–65.

6. University W. Phylogeny Programs.

http://evolution.genetics.washington.edu/phylip/software.html#Plotting. Accessed 2016 Jul

21.

7. Zhang H, Gao S, Lercher MJ, Hu S, Chen WH. EvolView, an online tool for visualizing,

annotating and managing phylogenetic trees. Nucleic Acids Res. 2012;40.

8. Rambaut A. FigTree. http://tree.bio.ed.ac.uk/software/figtree/. Accessed 2016 Jul 21.

9. Zmasek CM. Archaeopteryx.

https://sites.google.com/site/cmzmasek/home/software/archaeopteryx. Accessed 2016 Jul

21.

10. Paradis E, Claude J, Strimmer K. APE: Analyses of phylogenetics and evolution in R

language. Bioinformatics. 2004;20:289–90.

11. Revell LJ. phytools: An R package for phylogenetic comparative biology (and other

things). Methods Ecol. Evol. 2012;3:217–23.

12. Huerta-Cepas J, Dopazo J, Gabaldón T. ETE: a python Environment for Tree

Exploration. BMC Bioinformatics. 2010;11:24.

13. Chen W-H, Lercher MJ, Ganfornina M, Gutierrez G, Bastiani M, Sanchez D, et al.

ColorTree: a batch customization tool for phylogenic trees. BMC Res. Notes. BioMed

Central; 2009;2:155.

14. McDonald D, Clemente JC, Kuczynski J, Rideout J, Stombaugh J, Wendel D, et al. The

Biological Observation Matrix (BIOM) format or: how I learned to stop worrying and love

the ome-ome. Gigascience. 2012;1:7.

15. Ripley BD. The R project for statistical computing. 2001. p. 1–3.

16. Kelly KL. Twenty-two colors of maximum contrast. Color Eng. 1965. p. 26–7.

17. Stamatakis A. RAxML version 8: A tool for phylogenetic analysis and post-analysis of

large phylogenies. Bioinformatics. 2014;30:1312–3.

18. Felsenstein J. PHYLIP. http://evolution.gs.washington.edu/phylip.html. Accessed 2016

Jul 21.

19. Suttle CA. Marine viruses — major players in the global ecosystem. Nat. Rev.

Microbiol. Nature Publishing Group; 2007;5:801–12.

20. Rohwer F, Thurber RV. Viruses manipulate the marine environment. Nature. Nature

Publishing Group; 2009;459:207–12.

21. Rohwer F, Edwards R. The Phage Proteomic Tree: a genome-based taxonomy for

phage. J. Bacteriol. 2002;184:4529–35.

22. Phage SEED. http://www.phantome.org/PhageSeed/Phage.cgi. Accessed 2016 Jul 21.

23. Wommack KE, Nasko DJ, Chopyk J, Sakowski EG. Counts and sequences,

observations that continue to change our understanding of viruses in nature. J. Microbiol.

2015;53:181–92.

24. Mizuno CM, Rodriguez-Valera F, Kimes NE, Ghai R. Expanding the marine virosphere

using metagenomics. PLoS Genet. Public Library of Science; 2013;9:e1003987.

25. Chopyk J, Moore RM, DiSpirito Z, Stromberg ZR, Lewis GL, Renter DG, et al. Presence

of pathogenic Escherichia coli is correlated with bacterial community diversity and

composition on pre-harvest cattle hides. Microbiome. BioMed Central; 2016;4:9.

26. Fadrosh DW, Ma B, Gajer P, Sengamalay N, Ott S, Brotman RM, et al. An improved

dual-indexing approach for multiplexed 16S rRNA gene sequencing on the Illumina

MiSeq platform. Microbiome. 2014;2:6.

27. Lozupone C, Knight R. UniFrac: a New Phylogenetic Method for Comparing Microbial

Communities. Appl. Environ. Microbiol. American Society for Microbiology;

2005;71:8228–35.

28. Sakowski EG, Munsell E V., Hyatt M, Kress W, Williamson SJ, Nasko DJ, et al.

Ribonucleotide reductases reveal novel viral diversity and predict biological and

ecological features of unknown marine viruses. Proc. Natl. Acad. Sci. National Academy

of Sciences; 2014;111:15786–91.

29. McAllister SM, Davis RE, McBeth JM, Tebo BM, Emerson D, Moyer CL. Biodiversity

and emerging biogeography of the neutrophilic iron-oxidizing Zetaproteobacteria. Appl.

Environ. Microbiol. American Society for Microbiology (ASM); 2011;77:5445–57.

30. Biodocker. http://biodocker.org/. Accessed 2016 Jul 21.

31. Merkel D. Docker: lightweight Linux containers for consistent development and

deployment. Linux J. Belltown Media; 2014;2014:2.

32. GNU Operating System. http://www.gnu.org/licenses/. Accessed 2016 Jul 21.

Fig. 1: Precedence of Iroki's customization pipeline

Flowchart illustrating the precedence of steps when performing multiple customizations

with Iroki. Input/output files are purple, command line options are in blue, and processes

are orange. The choice of exact or regular expression matching guides each subsequent

step of the process. Iroki gives higher precedence to processes towards the bottom of the

diagram. For example, given that a user selects the options for coloring both labels and

branches, and provides both a biom file and color map with the color map specifying

colors for the labels only, then the branches will be colored according to the color

gradient inferred from the biom file, whereas the labels will be colored according to the

rules specified in the color map.

0.9

Fig. 2: Comparing phage and their host phyla

All phage genomes from Phage SEED with assembled virome contigs from the Chesapeake

Bay and Mediterranean Sea. Iroki highlights phylogenetic trends after coloring branches

according to viral family or sampling location in the case of virome contigs (marked with

an asterisk in the legend), and coloring node labels according to host phylum of the

phage.

Fig. 3: Changes in OTU abundance in two sample groups

Approximate-maximum likelihood tree of OTUs that showed significant differences in

relative abundance between STEC positive and STEC negative cattle hide samples.

Branches show significance based on coloring by the p-value of a Mann-Whitney U test

examining changes in abundance between samples positive for STEC (p < 0.05 – red) and

samples negative for STEC, (p >= 0.05 – blue). Label color on a blue-green color gradient

highlights OTU occurrence based on the abundance ratio between STEC positive samples

(blue) and STEC negative samples (green). For example, labels that are darker green had a

higher abundance in STEC negative samples. Node luminosity represents overall

abundance with lighter nodes being less abundant than darker nodes.

Fig. 4: Comparing cattle fecal and hide samples and the abundance of Ruminococcaceae

Phylogeny based on UPGMA tree of pairwise unweighted UniFrac distance between 356

bacterial community profiles based on SSU rRNA amplicon sequences from cattle hide

and feces. Branches are colored by feces (blue) and hide (red). Rapid testing of the

hypothesis that the abundance of one of the most abundant families, Ruminococcaceae,

and sample origin are correlated is enabled through node label coloring by (A) a green

single-color gradient (color saturation increases with increasing abundance of

Ruminococcaceae OTUs) and (B) a light green (low abundance of Ruminococcaceae

OTUs) to dark blue (high abundance of Ruminococcaceae OTUs) color gradient.

Fig. 5: Temporal dynamics of virioplankton populations according to Cyano II RNR

amplicon phylogeny

An approximately-maximum-likelihood phylogenetic tree of 200 randomly selected class

II Cyano RNR representative sequences from 98% percent clusters. Iroki was used to color

branches by time point: zero hours – yellow, six hours – orange, and twelve hours –

purple.

Fig. 6: Zetaproteobacteria show biogeographic partitioning

Phylogenetic tree showing placement of full-length Zetaproteobacteria SSU rRNA

sequences with outgroups. Iroki was used to color labels and branches by geographic

location of the sampling site (Loihi – green, Mariana – gold, South Pacific – purple, and

Other – gray), as well as to rename the nodes with OTU and sampling site metadata.