Accumulation and retention of radioactive elements in biofilm communities surrounding the

accident site of the Fukushima Daiichi Nuclear Power Plant

Shigeharu Moriya[1, 3, *], Hideaki Otsu[2], Kumiko Kihara[1, 4], Yukari Kato[1], Misao Itouga[1], Kenji

Sakata[1], Jun Kikuchi[1, 3], Hiroshi Yamakawa[5, 6] and Jiro Suga[6]

1) RIKEN Center for Sustainable Resource Science, 1-7-22 Suehiro-cho, Tsurumi-ku,

Yokohama city, Kanagawa 230-0045, JAPAN

Shigeharu Moriya, Kumiko Kihara, Yukari Kato, Misao Itouga, Kenji Sakata, and Jun

Kikuchi

2) RIKEN Nishina Center for Accelerator-Based Science, 2-1 Hirosamwa, Wako city,

Saitama 351-0198, JAPAN

Hideaki Otsu

3) Graduate School of Medical Life Science, Yokohama-city University, 1-7-29 Suehiro-cho,

Tsurumi-ku, Yokohama city, Kanagawa 230-0045, JAPAN

Shigeharu Moriya and Jun Kikuchi

4) Department of Biological and Chemical Systems Engineering, National Institute of

Technology, Kumamoto College, 2627 Hirayama-Shinmachi, Yatsushiro-shi, Kumamoto

866-8501, JAPAN

Kumiko Kihara

5) Tokyo University of Marine Science and Technology, 4-5-7 Konan, Minato-ku, Tokyo

108-8477, JAPAN

Hiroshi Yamakawa

6) Japan Academy of Underwater Sciences, 3-9-1-301 Botan, Koto-ku, Tokyo 135-0046,

JAPAN

Hiroshi Yamakawa and Jiro Suga

*Corresponding author: Shigeharu Moriya

RIKEN Center for Sustainable Resource Science, 1-7-22 Suehiro-cho, Tsurumi-ku,

Yokohama city, Kanagawa 230-0045, JAPAN

Tel +81-45-508-7221, Fax +81-45-508-7363

Email: smoriya@riken.jp

Abstract

After the Fukushima Daiichi Nuclear Power Plant accident, various surveys have been

performed to measure the extent of radioactive contamination in marine sediments, surface

waters, plankton, and fish. However, the radioactive contamination of one of the most

important ecological niches, biofilms, has not been investigated. Therefore, in this study, we

sampled biofilms from sea floor stones around Hisanohama Port, which is less than 30 km

south of the accident site, and then analyzed the microbial community structure and

element profiles, including those of radioactive elements, of these biofilms in order to

determine the accumulation and retention of radioactive elements in them. Our results

showed that the biofilm samples contained relatively high levels of radioactive cesium even

when the sampling was performed 8–11 months after the accident. Our results also

suggested that the structure of the biofilm organismal community is related to the element

profile of radioactive cesium. Thus, our study suggests that biofilms are a possible

radioactive compound accumulator in the natural environment and that they can retain

radioactive material for at least 8–11 months.

Introduction

Following the Great East Japan Earthquake and catastrophic tsunami in March of 2011,

the accident at the Fukushima Daiichi Nuclear Power Plant (FDNPP) led to the release of

large amounts of radioactive materials into the atmosphere and marine environment. This

release of radioactive elements from the FDNPP introduced a variety of radioisotopes,

especially radiocesium (^{134}Cs and ^{137}Cs), into marine waters. Public monitoring showed that

during several months after the first explosion at the FDNPP, a rapid decrease in

radioisotopes was observed in the nearby seawater. A joint official survey team from the

Tokyo Power Electric Company (TEPCO), Fukushima prefectural government, and the

Japanese Ministry of Education, Culture, Sports, Science and Technology (MEXT) reported

that the ^{137}Cs contamination level in the seawater quickly decreased from over 10 k Bq/L to

less than 100 Bq/L within a 2-month period. However, ^{137}Cs in the ocean sediments was

continuously observed over this period at levels between 100 Bq/dry kg and 1,000 Bq/dry kg

at least until September 2012 (Kanda, 2012). This disproportional distribution of

radioisotopes was also observed in marine fish by Wada et al., who reported that the

retention time of radiocesium appears to be remarkably longer in the larvae of demersal fish

such as *Sebastes cheni* than in surface fish such as *Eugraulis japonica* (Wada et al., 2013).

These disproportional distributions of radioactivity suggest that the sea floor eco-system,

including sedimental microbes and biofilms, may play role in the accumulation of

radioactive elements.

Therefore, in this study, we analyzed the biofilm community on the surface of sea floor rocks

with a focus on the element profiles, including those of radioactive elements. Our results

showed that all the biofilm community contained and retained relatively high levels of

radioactive cesium even when the sampling was performed 8–11 months after the accident.

Our findings indicate that biofilm organism community structures are likely related to their

element profiles, including that of radiocesium.

Materials and Methods

Sampling

Sampling was performed from November 11, 2011, to February 4, 2012. The sampling site

was the ocean along the coastline of Hisanohama, Iwaki City, Fukushima, Japan. We

collected samples 10 times within the established time period on November 11 (week 1),

November 20 (week 2), November 26 (week 3), December 3 (week 4), December 10 (week 5),

December 17 (week 6), January 14 (week 7), January 21 (week 8), January 29 (week 9), and

February 4 (week 10). Sampling was performed at six different locations near Hisanohama

Port: Area 1 (37.153762N 141.006167E, weeks 1 and 5), Area 2 (approximately 37.149863N

141.001017E, weeks 2, 8, and 9), Area 3 (approximately 37.146579N 141.002390E, week 4),

Area 3′ (approximately 37.147503N 141.008570E, week 10), Area 4 (approximately

37.133887N 141.002390E, weeks 3 and 7), and Area 4′ (approximately 37.142953N

141.003592E, week 6); all of these sampling sites are located 30 km south of the Fukushima

Nuclear Power Plant #1. Sampling was performed by divers using a self-contained

underwater breathing apparatus (SCUBA) at a depth of 5–10 m. The divers sampled the

bottom water and surface water (5 L) and collected bottom stones to obtain the biofilms. The

samplings were performed with permission from Fukushima Prefecture and the fishery

association of Iwaki City.

Sample preparation

The seawater samples were immediately filtered using two 24-mm-wide 0.22-μm Durapore

filters (Millipore, MA, USA) until they clogged. All the samples were then frozen at -20°C in

a portable freezer prior to transportation back to the laboratory.

In the laboratory, the water samples were dried by boiling. The remaining salts (seawater

salt) were allowed to settle into a U8 container, and the net weight (weight of container with

sample – weight of empty container) and height of the samples were measured in the

container.

In the case of biofilms, the bottom stones were rinsed with purified water and then brushed

down with a clean brush using purified water. The obtained biofilm suspension was then

divided into two aliquots. One aliquot was filtered through 24-mm-wide 0.22-µm Durapore

filters (Millipore, MA, USA) and stored at -20°C for ribosomal RNA gene analysis, and the

other one was dried at 105°C in an oven and then allowed to settle into a U8 container; the

weight and height were then measured in the container for measurement of radioactivity.

Measurement of radioactivity

The gamma-ray activity of the samples in the containers was measured using a Germanium

(Ge) detector. The detector setup was identical to that described previously (HABA et al.,

2012). The energy resolution of 2 keV full width at half maximum (FWHM) was achieved at

1332.5 keV gamma rays. The sample location and detector were surrounded with 15-cm lead

blocks to reduce the background radiation during gamma ray spectroscopy. The detection

efficiency of the gamma rays was calibrated with an accuracy of 2%–10% using a multiple

gamma ray standard source, which ranged from 88–1836 keV. A calibrated ^{134g}Cs source

was also used to correct for the coincidence summing for radioactivity determinations of

^{134g}Cs.

Every sample was contained in a U8 plastic container. The U8 container is a standard

container in Japan that is used for measurements of absolute radioactivity with high

accuracy using a Ge detector. The shape of the container is cylindrical, with a diameter of 47

mm and height of 60 mm. The volume is approximately 100 mL. The efficiency calibration

source is available within the container with a uniform density.

Every sample was measured using the Ge detector for at least 8 h to obtain sufficient

statistics for each gamma ray peak. The relative geometry between the sample and the Ge

detector was carefully reproduced by a guide apparatus at the sample location.

Gamma rays from ^{134}Cs and ^{137}Cs were observed for the biofilm samples. The radioactivity

levels for every isotope were determined from the yield of the gamma rays corrected using

the detection efficiency of the Ge detector and a geometrical acceptance between the sample

and the detector. The geometric difference in the height of the sample from the calibration

source was simulated using several electromagnetic simulation codes, including Geant4

(Agostinelli et al., 2003; Allison et al., 2006) and EGS5 (Hirayama et al., 2005). The

radioactivity of the samples were normalized by weight.

Element analysis

To quantitatively analyze the content of 37 elements (^{7}Li, ^{9}Be, ^{23}Na, ^{24}Mg, ^{27}Al, ^{39}K, ^{43}Ca,

^{45}Sc, ^{47}Ti, ^{51}V, ^{52}Cr, ^{55}Mn, ^{57}Fe, ^{59}Co, ^{60}Ni, ^{63}Cu, ^{66}Zn, ^{85}Rb, ^{88}Sr, ^{98}Mo, ^{111}Cd, ^{133}Cs, ^{139}La, ^{140}Ce,

^{141}Pr, ^{146}Nd, ^{152}Sm, ^{153}Eu, ^{158}Gd, ^{159}Tb, ^{164}Dy, ^{165}Ho, ^{166}Er, ^{169}Tm, ^{172}Yb, ^{175}Lu, and ^{202}Hg) in

the biofilm samples for measurement of radioactivity, the samples were predigested with 5

mL of concentrated HNO_3 for 1 h at room temperature. Next, the organic components were

completely decomposed by wet-ashing using a microwave sample preparation system

(Multi-Wave-3000, Perkin Elmer, MA USA) (Itouga et al., 2014). The digested samples were

then brought up to a volume of 50 mL using MilliQ water (MQW) and filtered through 5B

filter paper (Advantec, Tokyo, Japan). The concentrations of the mineral elements were

determined by Inductively Coupled Plasma Mass Spectrometry (ICP-MS, NexION300,

Perkin Elmer). For the ICP-MS analysis, a portion of the filtrated samples was diluted

appropriately with 0.01 mol/L HCl (Itouga et al., 2014). The mineral nutrient concentration

in the samples was calculated as a unit (mg/g in dry weight).

The obtained ICP-MS data were then normalized using the root-sum-of-squares (RSS) levels

for each element. Count data of radioactive cesium (^{134}Cs and ^{137}Cs) were treated with the

same calculation for normalization and then merged with the normalized ICP-MS dataset

(element table).

ssrDNA analysis

Genomic DNA was prepared from the stored biofilm samples by capturing on a 0.22-µm

Durapore filter, followed by bead-beating with phenol:chloroform:isoamyl alcohol (PCI)

extraction. Bead-beating was performed at 3000 rpm for 5 min (Micro Smash MS-100R,

TOMY, Tokyo, Japan), and nucleic acids were then recovered by ethanol precipitation.

PCR amplification was performed with the extracted genomic DNA using two primer sets

for 16S ssrDNA, 515F/806R (Caporaso et al., 2012; Caporaso et al., 2011) and 18S ssrDNA,

TAReuk454FWD1/TAReukREV3 (Stoeck et al., 2010). The primers were modified for

Illumina sequencing with a multi-index kit for the Nextera XT sequencing library

construction kit (Illumina, CA, USA). We also added a "read1" sequence (5′- TCG GCA GCG

TCA GAT GTG TAT AAG AGA CAG -3′) to the forward primers and a "read2" sequence (5′-

GTC TCG TGG GCT CGG AGA TGT GTA TAA GAG ACA G -3′) to the reverse primers. PCR

was performed with a standard reaction of ExTaq DNA polymerase according to the

manufacturer's instructions (Takara Bio, Kyoto Japan) with 25 pmol of each primer/50 μL

reaction mixture. The reactions were performed in a two-step PCR (10 cycles of 94°C for 30 s,

55°C for 45 s, and 72°C for 60 s, followed by 20 cycles of 94°C for 30 s and 72°C for 60 s) for

16S and a 3-step PCR (30 cycles of 94°C for 30 s, 55°C for 45 s and 72°C for 60 s) for 18S. The

amplified products were then purified via 1% agarose gel electrophoresis. The obtained

amplified products were applied to the NexteraXT (Illumina CA USA) sequencing library

construction process without the "tagmentation" process.

A multi-index sequencing library was applied to the MiSeq sequencing kit v1 and read using

Illumina MiSeq according to the instructions in the paired-end sequencing mode. Data were

treated with MiSeq reporter software, and the reads were obtained from individual samples.

After the extraction of the reads, the reads from weeks 4, 5, and 8 for the 18S ssrDNA were

found to be insufficient for analysis. Therefore, subsequent analyses were performed using

the reads from weeks 1, 2, 3, 6, 7, 9, and 10.

The obtained read data were then mapped onto the ribosomal RNA sequence library SILVA

release 108 with the program package QIIME (Caporaso et al., 2010) using "Prefix-suffix

operational taxonomy unit (OTU) picking," and the absolute abundance OTU table at the

family level was constructed. Count zero data were manually removed from the table, and

the relative abundance values were calculated. The 16S and 18S data were treated

independently and then merged into a single OTU table with the relative abundance values.

Sequence data have been deposited in the DDBJ sequence read archive under the accession

number DRA004367.

Statistical analysis

For the statistical analysis, we used the program package R (R Core Team, 2014). First, we

drew a heatmap chart with hierarchical cluster analysis (HCA). The heatmap.3 function in

the GMD package, which is a package for non-parametric distance measurements between

two discrete frequency distributions (Zhao et al., 2011), was used to draw the heatmap with

HCA. Independent heatmaps were drawn from each element table and the OTU table with a

two-axis HCA. A scaling option was used for the OTU table treatment according to the row

direction but not for the elements table treatment.

Principal components analysis (PCA) was performed with the merged dataset, including

both the OTU table and elements table, using the prcomp function in R. Score plots were

drawn with principal component 1 (PC1; x-axis) and PC2 (y-axis). The loading plots for

quadrants II and III were drawn manually.

Results and Discussion

The gamma-ray energy spectra revealed significant peaks corresponding to natural

radioisotope signals, such as that of ^{40}K (data not shown), for all samples. Furthermore, we

found significant peaks corresponding to ^{134}Cs and ^{137}Cs from the biofilm samples. However,

we did not detect signals for ^{134}Cs and ^{137}Cs from the sea water salts and particles captured

on filter (Table 1).

Table 1. Peak detection from samples

Material	Peak detection	
	134Cs	137Cs
Surface sea water salt	−	−
Bottom sea water salt	−	−
Captured particles in surface water	−	−
Captured particles in bottom water	−	−
Biofilms on bottom rocks	+	+

+: significant peak detected from samples

−: significant peak not detected from all samples

Captured particles were filtered from water by 0.22um filter and measure radioactivity

It is possible that the ^{134}Cs (half-life: 2.0652 years) and ^{137}Cs (half-life: 30.1 years) released

owing to prior nuclear weapons tests in the Pacific Ocean and the Chernobyl disaster have

remained in the seabed around Japan. Therefore, we also analyzed samples collected from

near Nishi-kawana in the Tokyo Bay (which is approximately 280 kilometers from the

FDNPP) as controls, but found no significant signals for the radioisotopes ^{134}Cs and ^{137}Cs

from these control samples. Therefore, we concluded that the radioisotopes detected in the

samples around Hisanohama port in this study originated from the FDNPP disaster rather

than from the environmental background or from past nuclear weapons tests. These results

indicated that the biofilms on the sea floor around Hisanohama retained radioactive

material for at least 8–11 months after the FDNPP accident.

However, the observed radioactivity level was different among the different biofilm samples

(Table 2). Therefore, we analyzed the relationship between the microbial consortia within

the biofilms and the elemental composition of the biofilms. To perform this analysis,

quantified element data were combined with radioactivity data for radiocesium (^{134}Cs and

^{137}Cs) and utilized for statistical analysis. Ribosomal RNA genes were sequenced and used

to construct a dataset for the statistical analysis.

Table 2. Radioactivity of biofilm samples

		Cs134 Bq/kg dry	Cs137 Bq/kg dry
Week1	Area1	1141.78	1655.17
Week2	Area2	671.01	932.18
Week3	Area4	306.90	429.21
Week4	Area3	429.21	2353.58
Week5	Area1	926.40	1436.73
Week6	Area4'	538.56	829.08
Week7	Area4	844.49	1543.05
Week8	Area2	818.66	1607.65
Week9	Area2	359.77	591.46
Week10	Area3'	190.29	294.20

A HCA heat map chart from the dataset of elements and radiocesium (Fig. 1) showed

relatively higher content of radiocesium, Cr, Ni, Sc, Rb, Li, and cesium (Cs cluster) for the

week 1 sample than for other samples. Another remarkable profile from the elemental

analysis was observed in the week 3 sample, where high relative amounts of lanthanide

elements were detected (Lanth cluster).

A PCA score plot (Fig. 2A) of the merged dataset between the elements/radiocesium and the

microbial consortia showed that the week 1 sample was placed in the PC1 negative-PC2

positive direction, and the week 3 sample was placed in the PC1 negative-PC2 negative

direction. As mentioned above, the week 1 sample contained high relative amounts of Cs

cluster signals, and the week 3 sample contained high relative amounts of the Lanth cluster

signals. Consistent with these findings, the PCA loading plot showed that the Cs cluster was

placed in the PC1 negative-PC2 positive direction (Fig. 2B upper panel, Table S1), and the

lanthanide cluster was placed in the PC1 negative-PC2 negative direction (Fig. 2B lower

panel, Table S2). The results of the HCA heat map chart and PCA suggested that a

"cesium-philic" biofilm occurred at the sampling point in the week 1 sample, and a

"lanthanide-philic" biofilm occurred at the sampling point in the week 3 sample.

The PCA results from the datasets of both the elements and the microbes showed candidate

organisms in the "cesium-philic" and "lanthanide-philic" biofilms on the loading plots,

because those plots were placed in same loading direction, e.g., PC1 negative-PC2 positive

(Cs cluster included) or PC1 negative-PC2 negative (Lanth cluster included, Fig. 2C, Table

S1 and S2). Indeed, the HCA heat map of the organismal consortia showed that the PC1

negative-PC2 negative–directed organisms mainly appeared in the week 1 column (Fig. 3

blue dots). The same tendency was observed with the Lanth cluster organisms (Fig. 3 green

dots). These results suggested that these two OTU sets are positively correlated with the

Lanth cluster (week 3 sample) and Cs cluster (week 1 sample) elements.

Table 3. Cs cluster taxon in Week1 sample

OTU ID	Taxon information	Relative intensity
Primer18 S_Euk_2	Eukaryota;Alveolata;Dinophyceae;Gonyaulacales;Gonyaulacaceae	0.02974
Primer18 S_Euk_4	Eukaryota;Metazoa;Annelida;Clitellata;Oligochaeta	0.10316
Primer18	Eukaryota;Metazoa;Annelida;Polychaeta;Scolecida	0.02144

ID	Primer	Taxonomy	Value
S_Euk_7	Primer18		
S_Euk_8	Primer18	Eukaryota;Metazoa;Arthropoda;Chelicerata;Arachnida	0.00847
S_Euk_11	Primer18	Eukaryota;Metazoa;Arthropoda;Crustacea;Maxillopoda	0.06880
S_Euk_12	Primer18	Eukaryota;Metazoa;Brachiopoda;Linguliformea;Lingulata	0.03599
S_Euk_15	Primer18	Eukaryota;Metazoa;Cnidaria;Anthozoa;Hexacorallia	0.14012
S_Euk_23	Primer18	Eukaryota;Metazoa;environmental samples;uncultured metazoan;Other	0.02215
S_Euk_25	Primer18	Eukaryota;Rhodophyta;Florideophyceae;Ceramiales;Ceramiaceae	0.05550
S_Euk_27	Primer18	Eukaryota;Rhodophyta;Florideophyceae;Ceramiales;Delesseriaceae	0.01796
S_Euk_34	Primer18	Eukaryota;environmental samples;uncultured eukaryote;Other;Other	0.03092
S_Euk_35		Eukaryota;environmental samples;uncultured freshwater eukaryote;Other;Other	0.00399
S_Euk_39	Primer18	Eukaryota;stramenopiles;Chrysophyceae;Hibberdiales;Hibberdiaceae	0.00977
S_Arch_1	Primer16	Archaea;Crenarchaeota;Marine Group I;uncultured archaeon;Other	0.06712
S_Bac_1	Primer16	Bacteria;Acidobacteria;RB25;uncultured bacterium;Other	0.00588
S_Bac_2		Bacteria;Actinobacteria;Acidimicrobiia;Acidimicrobiales;Sva0996 marine group	0.02444
S_Bac_3	Primer16	Bacteria;Actinobacteria;Acidimicrobiia;Acidimicrobiales;uncultured	0.02674
S_Bac_4	Primer16	Bacteria;Bacteroidetes;Cytophagia;Cytophagales;Flammeovirgaceae	0.07348
		Bacteria;Bacteroidetes;Flavobacteria;Flavobacteriales;Cryomorphaceae	0.00429

ID	Taxonomy	Value
Primer16 S_Bac_5		
Primer16 S_Bac_7	Bacteria;Bacteroidetes;Sphingobacteria;Sphingobacteriales;Saprospiraceae	0.01855
Primer16 S_Bac_9	Bacteria;Cyanobacteria;Chloroplast;uncultured bacterium;Other	0.00919
Primer16 S_Bac_16	Bacteria;Proteobacteria;Alphaproteobacteria;Rhizobiales;Hyphomicrobiaceae	0.00994
Primer16 S_Bac_18	Bacteria;Proteobacteria;Alphaproteobacteria;Rhizobiales;Phyllobacteriaceae	0.00703
Primer16 S_Bac_20	Bacteria;Proteobacteria;Alphaproteobacteria;Rhodobacterales;Rhodobacteraceae	0.15535
Primer16 S_Bac_21	Bacteria;Proteobacteria;Alphaproteobacteria;Sphingomonadales;Erythrobacteraceae	0.01248
Primer16 S_Bac_28	Bacteria;Proteobacteria;Deltaproteobacteria;Desulfobacterales;Desulfobacteraceae	0.01218
Primer16 S_Bac_29	Bacteria;Proteobacteria;Deltaproteobacteria;Desulfobacterales;Desulfobulbaceae	0.00744
Primer16 S_Bac_31	Bacteria;Proteobacteria;Deltaproteobacteria;Sh765B-TzT-29;uncultured delta proteobacterium	0.01246
Primer16 S_Bac_32	Bacteria;Proteobacteria;Epsilonproteobacteria;Campylobacterales;Helicobacteraceae	0.00372
Primer16 S_Bac_35	Bacteria;Proteobacteria;Gammaproteobacteria;BD7-8 marine group;uncultured bacterium	0.02819
Primer16 S_Bac_37	Bacteria;Proteobacteria;Gammaproteobacteria;E01-9C-26 marine group;uncultured bacterium HERMI11	0.00922
Primer16 S_Bac_38	Bacteria;Proteobacteria;Gammaproteobacteria;KI89A clade;uncultured gamma proteobacterium	0.11585
Primer16 S_Bac_39	Bacteria;Proteobacteria;Gammaproteobacteria;Oceanospirillales;OM182 clade	0.01344
Primer16 S_Bac_40	Bacteria;Proteobacteria;Gammaproteobacteria;Oceanospirillales;Oceanospirillaceae	0.04369
Primer16	Bacteria;Proteobacteria;Gammaproteobacteria;Oceanospirillales;SS1-B-0	0.00422

S_Bac_41	6−26	
Primer16	Bacteria;Proteobacteria;Gammaproteobacteria;Order Incertae	
S_Bac_42	Sedis;Family Incertae Sedis	0.04639
Primer16	Bacteria;Proteobacteria;Gammaproteobacteria;Xanthomonadales;Sinobact	
S_Bac_46	eraceae	0.02341
Primer16		
S_Bac_48	Bacteria;Verrucomicrobia;Verrucomicrobiae;Verrucomicrobiales;DEV007	0.00691

Table 3 lists the OTU set contents that correlated with Cs cluster elements, including

radiocesium, and the relative intensity of each OTU in the biofilm community. The most

abundant taxa in this OTU set were *Hexacorallia* and *Oligochaete* among the eukaryotes.

For bacteria, high relative intensities of *Bacteroidetes*, *Rhodobacteraceae*, and

Gammaproteobacteria were observed. Although a primer set that did not target Archaea

was used, archeal sequences were obtained in this work by sequencing amplicons generated

using the 16S ribosomal RNA primer set. An uncultured *Crenarcheota* in marine group I

was observed to be one of the dominant OTUs with a correlation with Cs cluster elements.

Previously, *Alpha-* and *Gamma-proteobacteria* have been obtained from biofilms on the

surface of nuclear fuel pools (Sarró et al., 2005). According to Sarro et al. (Sarró et al., 2005),

biofilms obtained from nuclear fuel pools accumulate ^{60}Co. In our study, cobalt accumulation

in the week 1 sample was suggested by the PCA data (Fig. 2B upper panel and Table S1)

despite a lack of ^{60}Co signal during gamma-ray spectroscopy. Microbial community retrieved

from nuclear fuel pools have been described previously as capable of accumulating both ^{60}Co

and ^{137}Cs (Tišáková et al., 2012). Our results suggest that the biofilm sampled in week 1 in

our study presents similar characteristics to the microbes described by Sarro et al. and

Tisakova et al. in the above-mentioned previous studies.

Conclusions

In this study, we found that biofilms retain a relatively high amount of radioactive cesium in

their microbial communities even after more than half a year from the release of radioactive

elements into oceans due to nuclear power plant accidents. Our statistical analysis also

indicated that different community structures of biofilms show different affinities for

radiocesium. Our results suggested biofilms as a point of entry for radioactive elements into

the eco-system owing to long-term retention of radiocesium in the biofilm communities.

In the future, further research on the accumulation and retention of radioactive elements by

marine biofilm communities needs to be undertaken to obtain information that can be used

for developing ways to eliminate radioactive contamination and for monitoring of

environmental pollution by radiocesium.

Acknowledgements

This work was supported by grants for the RIKEN Incentive Research Project (FY2011)

"Emergency monitoring for possible cycling of radio active compounds via oceanic food web"

awarded to SM and HO and a MEXT Grant-in-Aid for Scientific Research on Innovative

Areas (No. 23117003) awarded to SM and KK. We also wish to thank the Iwaki City Fishery

Association for permitting the sampling for our study and for providing the fishing vessel

Shyo-ei-maru and supporting crew belonging to the abalone section of the Iwaki Fishery

Association for our study. We are also grateful to Toshiyuki Suzuki for overseeing and

ensuring the diving safety of SM. The authors also wish to acknowledge the profound

academic comments and proofreading provided by R. Craig Everroad and Diogo M. O.

Ogawa.

References

Agostinelli, S., Allison, J., Amako, K., Apostolakis, J., Araujo, H., Arce, P., Asai, M., Axen, D.,

Banerjee, S., Barrand, G., et al. (2003). Geant4—a simulation toolkit. *Nuclear*

Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers,

Detectors and Associated Equipment 506, 250–303.

Allison, J., Amako, K. and Apostolakis, J. (2006). IEEE Xplore Abstract - Geant4

developments and applications. *Nuclear Science* 53, 270–278.

Caporaso, J. G., Kuczynski, J., Stombaugh, J., Bittinger, K., Bushman, F. D., Costello, E. K.,

Fierer, N., Peña, A. G., Goodrich, J. K., Gordon, J. I., et al. (2010). QIIME allows

analysis of high-throughput community sequencing data. *Nat. Methods* 7, 335–336.

Caporaso, J. G., Lauber, C. L., Walters, W. A., Berg-Lyons, D., Huntley, J., Fierer, N.,

Owens, S. M., Betley, J., Fraser, L., Bauer, M., et al. (2012). Ultra-high-throughput

microbial community analysis on the Illumina HiSeq and MiSeq platforms. *ISME J* 6,

1621–1624.

Caporaso, J. G., Lauber, C. L., Walters, W. A., Berg-Lyons, D., Lozupone, C. A., Turnbaugh,

P. J., Fierer, N. and Knight, R. (2011). Global patterns of 16S rRNA diversity at a depth

of millions of sequences per sample. *Proceedings of the National Academy of Sciences*

108 Suppl 1, 4516–4522.

HABA, H., KANAYA, J., MUKAI, H., KAMBARA, T. and KASE, M. (2012). One-year

monitoring of airborne radionuclides in Wako, Japan, after the Fukushima Dai-ichi

nuclear power plant accident in 2011. *Geochem. J.* 46, 271–278.

Hirayama, H., Namito, Y., Nelson, W. R. and Bielajew, A. F. (2005). The EGS5 code system.

Itouga M., Kato Y., Sakakibara H. (2014). Phenotypic plasticity and mineral nutrient

uptake of the moss Polytrichum commune Hedw. (Polytrichaceae, Bryophyta) during

acclimation to a change in light intensity. *Hikobia* 16, 459–466.

Kanda, J. (2012). Long-term Sources:To what extent are marine sediments, coastal

groundwater, and rivers a source of ongoing contaminaton? pp. 1–19. Tokyo.

Sarró, M. I., García, A. M. and Moreno, D. A. (2005). Biofilm formation in spent nuclear fuel

pools and bioremediation of radioactive water. *Int. Microbiol.* 8, 223–230.

Stoeck, T., Bass, D., Nebel, M., Christen, R., Jones, M. D. M., Breiner, H.-W. and Richards, T.

A. (2010). Multiple marker parallel tag environmental DNA sequencing reveals a highly

complex eukaryotic community in marine anoxic water. *Mol Ecol* 19 Suppl 1, 21–31.

Team, R. C. (2014). R: A language and environment for statistical computing.

www.R-project.org.

Tišáková, L., Pipíška, M., Godány, A., Horník, M., Vidová, B. and Augustín, J. (2012).

Bioaccumulation of 137Cs and 60Co by bacteria isolated from spent nuclear fuel pools. *J*

Radioanal Nucl Chem 295, 737–748.

Wada, T., Nemoto, Y., Shimamura, S., Fujita, T., Mizuno, T., Sohtome, T., Kamiyama, K.,

Morita, T. and Igarashi, S. (2013). Effects of the nuclear disaster on marine products in

Fukushima. *Journal of Environmental Radioactivity* 124, 246–254.

Zhao, X., Valen, E., Parker, B. J. and Sandelin, A. (2011). Systematic clustering of

transcription start site landscapes. *PLoS ONE* 6, e23409.

Figure legends

Figure 1

Heatmap chart of the elements and radioactivity data from the biofilm samples based on

hierarchical clustering analysis (HCA). The heatmap chart was sorted by HCA for both axes

(elements axis and sample axis). Boxes indicate clusters of radiocesium (Cs cluster) or

lanthanides (Lanth cluster).

Figure 2

Principal component analysis (PCA) results for the sampling sites with microbial consortia

data and element data with radiocesium. Panel A is the PCA score plot among sampling

sites. Each dot indicates one sampling site. Panel B indicates the loading plot of the

elements data. The upper panel shows the plot of elements that contribute to the

PC1-negative and PC2-positive directions (supplementary table 1), indicating that they

contribute to separate week 1 samples in the PC1-negative and PC2-positive directions. The

lower panel shows the plot of elements that contribute to the PC1-negative and

PC2-negative directions (supplementary table 2), indicating that they contribute to separate

week 3 samples in the PC1-negative and PC2-negative directions in the PCA score plot.

Panel C indicates the loading plot of the microbial consortia data. The upper panel displays

the plot of operational taxonomic units (OTUs) that contribute to the PC1-negative and

PC2-positive direction (supplementary table 1), indicating that they contribute to separate

week 1 samples in the PCA score plot. The lower panel shows the plot of the OTUs that

contribute to the PC1-negative and PC2-negative directions (supplementary table 2),

indicating that they contribute to separate week 3 samples in the PCA score plot.

Figure 3

Heatmap chart of microbial consortia data for biofilms with HCA.

The heatmap chart has been sorted by HCA for both axes (OTU axis and sample axis).

Green and blue dots indicate contributing OTUs for the PC1-negative and PC2-positive

directions and the PC1/PC2-negative direction, respectively. The relative intensity value

has been scaled and centered in the row direction.

Supplementary tables.xlsx

Supplementary table 1

Supplementary table 2

Fig.1

Fig.2

Fig.3

Row direction scaled and centered
relative abundance

-2   3