Title: Single neurons may encode simultaneous stimuli by switching between

activity patterns

Authors: Valeria C. Caruso[1,2,3,4]†, Jeff T. Mohl[1,2,3,4], Christopher Glynn[7,8], Jungah Lee[1,2,3,4]‡,

Shawn M. Willett[1,2,3,4], Azeem Zaman[6], Akinori F. Ebihara[10], Rolando Estrada[5,9], Winrich A.

Freiwald[10], Surya T. Tokdar[1,6]†, Jennifer M. Groh[1,2,3,4]†

Affiliations:

[1]Duke Institute for Brain Sciences, Duke University, Durham, NC 27708

[2]Center for Cognitive Neuroscience, Duke University

[3]Department of Psychology and Neuroscience, Duke University

[4]Department of Neurobiology, Duke University

[5]Department of Computer Science, Duke University

[6]Department of Statistical Science, Duke University

[7]Department of Decision Sciences, University of New Hampshire

[8]Department of Statistics, Harvard University

[9]Department of Computer Science, Georgia State University, Atlanta, GA

[10]The Rockefeller University, New York, New York

† Co-corresponding authors. Correspondence to VCC at v.caruso@duke.edu, or JMG at

jmgroh@duke.edu or ST at surya.tokdar@duke.edu

‡ Current address: Department of Psychology, University of Canterbury, New Zealand

Short title: Fluctuations as part of the neural code

Keywords: neural coding, time division multiplexing, stimulus multiplexing across time,

statistical methods, perception

ABSTRACT

How the brain preserves information about multiple simultaneous items is poorly understood. We report that single neurons can represent multiple different stimuli by interleaving different signals across time. We record single units in an auditory region, the inferior colliculus, while monkeys localize 1 or 2 simultaneous sounds. During dual-sound trials, we find that some neurons fluctuate between firing rates observed for each single sound, either on a whole-trial or on a sub-trial timescale. These fluctuations are correlated in pairs of neurons, can be predicted by the state of local field potentials prior to sound onset, and, in one monkey, can predict which sound will be reported first. We find corroborating evidence of fluctuating activity patterns in a separate data set involving responses of inferotemporal cortex neurons to multiple visual stimuli. Alternation between activity patterns corresponding to each of multiple items may therefore be a general strategy to enhance the brain processing capacity, potentially linking such disparate phenomena as variable neural firing, neural oscillations, and limits in attentional/memory capacity.

In the natural world many stimuli or events occur at the same time. Sensory neurons in the brain are broadly tuned and potentially responsive to more than one such stimulus, raising the question of how information about multiple simultaneous items can be preserved. We investigated whether the brain solves this problem at the neuronal level via activity patterns that fluctuate between those evoked by each stimulus alone (Figure 1A). If present, such fluctuations could allow each individual stimulus to be represented across time in a common neuronal ensemble. Such switching would contrast with possibilities such as signal summation or normalization/averaging, either of which results in information about each item being lost. Fluctuation is more related to winner-take-all but suggests that the winning stimulus might change across time for an individual neuron. [1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11]

Specifically, we tested how neurons in the monkey inferior colliculus (IC) respond when two sounds are presented simultaneously from different locations and both must be behaviorally reported. The IC is an early[12, 13] and nearly obligatory station along the auditory pathway [14], and thought to be essential for accurate sound localization behavior [15, 16, 17]. Importantly, the IC is thought to encode sound location via a monotonic firing rate (meter) code in which individual neurons respond broadly across a range of spatial positions but the level of activity is correlated

with the location of the sound [18, 19, 20, 21]. We developed a novel statistical approach to assess whether

signal fluctuations in the IC serve to interleave information about both sounds at a variety of time

scales.

We found that a subpopulation of IC neurons exhibited fluctuating activity consistent with

switching between individual sound responses at different time scales. Such activity fluctuations

were masked in conventional analysis of spiking activity (trial-and-time pooled spike counts),

which indicated that average dual-sound activity levels were intermediate between the average

single-sound responses. The state of the network prior to stimulus onset (assessed by the mean

voltage of the local field potentials) predicted the slower whole-trial spiking fluctuations, and in

turn, these fluctuations could predict which sound location the monkey reported first. We

replicated the key observations regarding fluctuating activity in a separate data set involving

inferotemporal cortex: neurons confronted with multiple object stimuli exhibited activity

fluctuations consistent with switching between individual object responses. These observations

support fluctuating activity as a viable and likely a general strategy for encoding simultaneously

presented stimuli. We conclude by discussing how such a code can be read out, especially with

regard to such factors as time scale and coordination across ensembles of neurons. Finally, we

consider several broad implications of activity fluctuations for interpreting variability and other

aspects of neural encoding.

Figure 1. Experimental rationale, task and visualization of individual trial activity.

(A) In telecommunications, multiple signals can be conveyed along a single channel by interleaving samples of each, thus increasing the amount of information transmitted by a single physical resource. Here we investigated whether the brain might employ a similar strategy: do neurons encode multiple items (A and B) using spike trains that alternate between the firing rates

corresponding to each item, at some unknown time scale? Such a strategy would preserve

information about both items, in contrast to alternatives such as winner-take-all, summation, or

*averaging, which involve varying degrees of information loss. (**B**) Sound localization task. Two*

monkeys were successfully trained to report one or two simultaneous (bandlimited noise) sounds

*by saccading at them. See Supplementary Figure 1 for accuracy. (**C**) Eye traces of the saccades*

*towards one (left) or two targets (right) during a sample session. (**D**) Time-and-trial aggregated*

dual-sound responses resemble the averaging more than the summation of single-sound responses.

For 81% of the triplets tested, the absolute values of the Z-score of each dual-sound response

*relative to the average were smaller than those relative to the sum. (**E-F**) Visualization of*

individual trials of two IC neurons in which dual-sound responses alternate between firing rates

corresponding to single-sounds, across trials for the neuron in E, or within trials for the neuron

in F. In each panel, the red and blue shaded areas indicate the median and central 50% of the

data on the single-sound trials. The black traces are the individual trials, for single-sound and

dual-sound trials as indicated above the panel. For the neuron in E, individual traces on dual-

sound trials were classified based on whether they matched the responses to single-sounds A and

B (A vs. B assignment score, see Methods) and are plotted in two separate panels accordingly.

For the neuron in F, the fluctuations occurred faster, within trials, and are plotted in the same

panel. See Supplementary Figure 4 for peristimulus time histograms and frequency sensitivity of

these two example neurons.

RESULTS

1. Time-and-trial pooled activity does not account for behavior

We first tested whether monkeys can perceptually preserve information about multiple

sounds presented simultaneously. Two monkeys performed a localization task (Figure 1B) in

which they made eye movements to each of the sounds they heard: one saccade on single-sound

trials and two saccades in sequence on dual-sound trials (Figure 1C and Supplementary Figure 1).

The sounds were separated horizontally by 30 degrees and consisted of band-limited noise with

different central frequencies. The sounds were thus physically distinguishable in principle, and

humans are able to do so [22, 23, 24]. The monkeys learned the task successfully, and, like humans,

typically performed better when the frequency separation between the two sounds was larger

(Figure 1C, Supplementary Figure 1, ~72 vs. ~77% correct for frequency differences of 3.4 vs. 6.8

semitones).

If the monkeys can report the locations of two sounds presented simultaneously, it follows

that their brains, and the IC in particular, must preserve information about both sound items. To

investigate the neural basis of this perceptual ability, we recorded 166 single units from the left

and right IC of two monkeys (see Methods), and evaluated the activity patterns evoked on dual-

sound trials (involving a given pair of sound locations and frequencies, AB) in comparison to the

activity patterns evoked on the matching single-sound trials (A and B). We refer to such related

sets of stimuli as triplets.

Conventional analysis of spike data typically involves two simplifications: spikes are

counted within a fairly long window of time, such as a few hundred milliseconds, and activity is

pooled across trials for statistical analysis. If IC neurons interleave signals related to each of the

two sounds, then they might appear to show averaging responses on dual (or AB) trials when

activity is pooled across time and across trials. But they should not appear to show summation

responses, i.e. in which the responses on dual-sound trials resemble the sum of the responses

exhibited on single-sound trials involving the component sounds. Such summation has been

observed in some neural populations in areas such as primary visual cortex [25, 26], the hippocampus

[27], and the superior colliculus [28] when multiple stimuli are presented.

Indeed, IC dual-sound responses do not generally appear to sum the responses to the

component single sounds. Using an analysis similar to that of [28], dual-sound responses were

converted to Z-scores relative to either the sum or the average of the corresponding single-sound

responses (see Methods). For 81% of the tested triplets, the Z-score values relative to the average

were smaller than those relative to the sum, indicating that the responses more closely resembled

averaging (Figure 1D, see also Supplementary Figure 2 and 3)

However, such apparent averaging response patterns appear inconsistent with the

behavioral results: if the neurons truly responded at an average firing rate, then presumably the

monkeys should respond to dual-sounds as if there were only a single sound at the midpoint of the

two sources (Supplementary Figure 2A). Since monkeys can indicate the locations of *both* sounds

(Figure 1C), fluctuating activity patterns that could preserve both items across time in a neuronal

ensemble might provide a better explanation for so-called averaging response patterns.

2. Evaluating activity fluctuations at various time scales

To determine whether neural activity fluctuates within and/or between trials, creating an overall averaging response but retaining information about each sound at distinct moments, we developed a series of statistical analyses that test for the presence of various forms of alternation in firing rates. Several unknown parameters must be taken into consideration when testing for activity fluctuations, specifically, the time scale, repeatability, and potential correlations across the neural population. We made minimal assumptions about the time scale at which neurons might alternate between encoding each stimulus, and considered that switching behaviors might vary from trial to trial and/or across time within a trial, as suggested by the visual inspection of single unit responses.

Figure 1E-F shows the activity of two example neurons on dual-sound trials compared to their matched single-sound trials. The colored backgrounds illustrate the median and 25-75% quantiles of the activity on single-sound trials, in 50ms time bins. Superimposed on these backgrounds is the activity on individual trials. Individual single-sound (A alone, B alone) trials align well with their corresponding 25-75% quantiles, by definition (Figure 1E-F). But on dual-sound (AB) trials, for any given trial or time bin, some individual traces correspond well to the 25-75% quantiles of one of the component sounds, and on other trials or time bins they correspond well to the 25-75% quantiles of the other component sound. For the neuron in Figure1E, there are whole trials in which the activity matches that evoked by sound A alone and others in which it better corresponds to that evoked by sound B alone. For the neuron in Figure 1F, the firing pattern on dual-sound trials appears to switch back and forth between the levels observed for sounds A and B as the trial unfolds. In short, for these two examples, the activity on dual-sound AB trials does not appear to occur at a consistent value intermediate between those evoked on single-sound A and B trials, but can fluctuate between those levels at a range of time scales. The following analyses will quantitatively assess these fluctuations across trials and/or time.

3. Evidence for fluctuations in whole-trial spike counts

If neurons alternate firing rates at the time scale of trials, as appears to be the case for the neuron in Figure 1E, then the spike counts from dual-sound responses should resemble a mix of

spike counts from each of the component single-sound responses. We statistically tested this hypothesis against other reasonable competing possibilities using the subset of triplets whose spike counts on single-sound A and B trials could be well modeled by Poisson distributions with statistically different mean rates λ^A and λ^B (N=363 triplets from N=108 neurons, see methods for details).

The competing scenarios to describe the corresponding dual-sound trials were:

(a) **Mixture**: The spike counts observed on individual trials are best described as draws from a mixture of $Poi(\lambda^A)$ and $Poi(\lambda^B)$ (Figure 2A). This is consistent with fluctuating activity across trials.

(b) **Intermediate**: A single Poisson distribution best describes the spike counts, and this Poisson has a rate λ^{AB} between λ^A and λ^B (Figure 2B). This is consistent with either fluctuations at faster, sub-trial time scales or with true averaging/normalization.

(c) **Outside**: A single Poisson distribution best describes the spike counts, but the rate λ^{AB} is outside the range of λ^A and λ^B (i.e it is greater than both or less than both; Figure 2C). Summation-type responses would be captured under this heading, as would inhibitory interactions.

(d) **Single**: A single Poisson describes the dual-sound trial spike counts, but the rate λ^{AB} is equal to one of the single-sound rates λ^A or λ^B (Figure 2D). A winner- (or loser-)-take-all pattern would fit this category.

In summary, these four models capture the spectrum of possibilities at the whole-trial time scale. For each triplet, we computed the posterior probabilities of the four competing models as a (Bayesian) measure of goodness of fit to the data [29] (see Methods). In other words, for each triplet the model with the highest posterior probability can be interpreted as the best fit, and the posterior probability itself is a measure of the confidence in the model. We consider three levels of confidence from high to low (posterior probability >0.95, >0.50, and no threshold on the confidence level, that is, posterior probability > the minimum, 0.33, figure 2I).

Figure 2. Whole-trial analysis. Panels A to D show the four models that could describe the distribution of spike counts on individual dual-sound trials (0-600 or 0-1000 ms after sound onset, see Methods). (A) Mixture of the Poisson distributions of spike counts for the component single-sound trials, (B) Intermediate Poisson distribution, with rate between the rates of single-sounds responses, (C) Outside, Poisson distribution with rate larger or lower than the rates of single-sounds responses, (D) Single, Poisson distribution with rate equal to one of the two single-sound rates. (E,F,G,H) Four examples of spike count distributions for triplets classified as Mixtures or

*Intermediates. Red and blue shades indicate distributions of spike counts for single-sounds; black outlines indicate distributions for dual sounds. The triplets in **E,F** were classified as Mixture with winning probability >0.95 (panel E shows the same triplet as Figure 1E; panel F shows the same triplet as Figure 3B). Triplets in G,H were classified as Intermediate with winning probability >0.95 (panel G shows the same triplet as Figure 1F and figure 3C ; panel H shows the same triplet as Figure 3D). (**I**) Population results of the whole-trial analysis. Shading indicates the confidence level of the assignment of individual triplets to winning models.*

The Mixture model had the strongest support of any of the models tested. For such triplets, the spike counts on dual-sound trials were better fit by a mixture of the single-sound Poisson distributions than by any single Poisson distribution (Figure 2I, bar labeled Mixture). These response patterns indicate the presence of fluctuating activity at the level of individual trials; the neurons illustrated in Figures 2E (same as Figure 1E) and 2F (same as Figure 3B) met these criteria. Of the 72 triplets (from N=46 neurons) in which one model had a winning probability >0.95, 50 triplets (69%) were categorized as Mixtures.

For the next largest category, the best fitting model involved a unique λ^{AB} between λ^A and λ^B (Figure 2I, bar labeled Intermediate). These triplets are ambiguous: they could exhibit a true intermediate firing rate on the dual-sound trials, or they could simply show alternation at a time scale more rapid than individual trials. The neurons illustrated in Figures 2G (same as Figure 1F and 3C) and 2H (same as figure 3D) were classified as Intermediate. Of the 72 triplets in which one model had a winning probability >0.95, 18 triplets (25%) were categorized this way.

The remaining triplets were categorized as Single, or $\lambda^{AB} = \lambda^A$ or λ^B (a narrowly defined category that consequently did not produce any winning model probabilities >0.95) or Outside, λ^{AB} greater or less than both λ^A and λ^B. Single can be thought of as a winner-take-all response pattern. Outside may be consistent with a modest degree of summation in the neural population, particularly as λ^{AB} was generally greater than both λ^A and λ^B in this subgroup. The small prevalence of the Single and Outside groups compared to the Mixture and Intermediates is in agreement with the aggregated analysis of spiking data (Supplementary Figures 2-3). The overall pattern of classification in the whole-trial analysis was similar in the population of triplets that exhibited responses to both sounds individually vs. those that responded to only one of the two sounds

(Supplementary Figure 5), with the exception that the bulk of the Outside triplets were indeed responsive to both A and B sounds individually.

4 Evidence for within-trial fluctuations vs. stable averaging

We next evaluated whether firing patterns fluctuated or remained stable across time within a trial. In particular, might triplets categorized as Intermediate in the whole trial analysis show evidence of fluctuating activity on a faster time scale?

We developed a novel statistical approach to study temporal patterns of the spike trains, the **Dynamic Admixture of Poisson Process (DAPP)** model. The computation is schematized in figure 3A (see also Methods). We focused on the same 363 triplets selected above. For each triplet, spike trains from individual single-sound trials were assumed to be independent realizations of a nonhomogeneous Poisson process with unknown time-dependent firing rates $\lambda^A(t)$ and $\lambda^B(t)$ for sounds A and B. Dual-sound trials were modeled as a weighted combination: $\lambda^{AB}(t) = \alpha(t)\lambda^A(t) + (1 - \alpha(t))\lambda^B(t)$ (Figure 3A). The weight function $\alpha(t)$ was unique to each dual-sound trial and quantified the relative contribution of sound A on that trial at time t, while $1 - \alpha(t)$ quantified the complementary contribution of sound B. Thus, the dynamics of the $\alpha(t)$ function characterize the dynamics of each dual-sound trial (see examples in figure 3B,C,D).

For each selected triplet, we used a Bayesian analysis to estimate the function valued model parameters (see Methods) and predict the $\alpha(t)$ curves that the corresponding cell was likely to produce on future dual-sound AB trials (figure 3A). Each predicted curve was summarized by two features: its time average over the response period of a given trial and its maximum swing size, that is, the difference between its highest peak and lowest trough on that trial. The triplet was then subjected to a two-way classification based on the distribution of these two features over the predicted curves (DAPP tags, Figure 3A). The triplet was categorized as Wavy vs. Flat depending on whether the distribution of the maximum swing size peaked at high or low values, and as Central vs. Extreme according to whether the distribution of the time average $\alpha(t)$ had a peak close to 0.5 or had one to two peaks at the extreme values of 0 and 1. In addition to this main classification scheme, triplets were subcategorized as exhibiting Symmetric or Skewed response patterns, indicating whether the $\alpha(t)$ curves reflected roughly equal contributions from the stimulus A and

stimulus B response patterns or whether one or the other tended to dominate (Supplementary

Figure 6).

Figure 3. Dynamic Admixture Point Process (DAPP) model: rationale and results (A) The

DAPP model fits smoothly time-varying weights (α and (1- α)) capturing the relative contribution

of A- and B-like response distributions to each AB dual-sound trial (point1). The dynamic
tendencies of the α curves were then used to generate projected new α curves for hypothetical
future draws from this distribution. The waviness and central tendencies were quantified by
computing the max swing size and trial-wise mean for an individual trial draw from the
distribution (point 2). Low max swing sizes indicate flat curves and higher values indicate wavy
ones (point 3, right panel). Similarly, the distribution of trial-wise means could be bimodal
(Extreme) or unimodal (Central) (point 3, left panel). (B,C,D) Fit alphas for three example triplets
(triplets in panels B,C,D are the same as in Figure 2 F,G,H, respectively) and the distributions of
trial-wise means and max swing sizes for future draws from the alpha curve generator. (E) The
pattern of DAPP results extended the whole-trial analysis results. Triplets categorized as Mixtures
with a win probability > 0.95 tended to be tagged as Flat-Extreme (as example in B). Triplets
categorized as Intermediates fell in two different main groups, Wavy-Central (as example in C)
and Flat-Central (as example in D). Information about the Skewed vs. Symmetric tag is not shown.
See Supplementary Table 1 and Supplementary Figures 6 and 7 for a complete listing of all the
tag combinations and additional analyses.

The DAPP tags confirmed and extended the results of the whole-trial analysis. Figure 3E
shows the different distributions of DAPP tags for the triplets categorized as Intermediate and
Mixture in the whole-trial analysis with winning probability >0.95 (see also Supplementary
Figures 6-7).   Mixture triplets tended to be classified as Flat-Extreme (70%), that is, the dynamics
of these dual-sound responses were flat in time and matching either the responses to A ($\alpha \approx 1$) or
B ($\alpha \approx 0$) on different trials. In addition, the distribution of the average α values tended to be either
Symmetric or unlabeled with regard to symmetry, a sanity check that excludes winner-take-all
responses (these would be characterized by Flat-Extreme-Skewed responses, Supplementary Table
1, Supplementary Figure 6).
In contrast, Intermediate triplets showed a combination of two types of labeling patterns
relevant to our hypothesis. Some (22%) were classified as Flat-Central (and Symmetric),
indicating stable α between 0 and 1, that is, stable firing at roughly the average of the responses
evoked by each sound separately (see example cell in figure 3D). One triplet was classified as Flat-
Extreme suggesting a stable α very close to either 0 or 1. Together these two firing patterns are
consistent with some form of normalization occurring in this subpopulation. Other triplets (28%)

were classified as Wavy-Central (and Symmetric) indicating responses that fluctuated
symmetrically around a central value (see example cell in figure 3C).

In short, the DAPP analysis shows that the dynamics of the Mixture responses were
consistent with fluctuations at the level of whole trials like the cell in figure 3B, whereas the
dynamics of the Intermediate responses classified as Wavy-Central are suggestive of a neural code
that could switch relatively rapidly between responses consistent with two single stimuli.

314
315

5 Fluctuations appear coordinated and predict behavior

We next considered the question of whether and how activity fluctuations are coordinated
across the neural population, in two ways: (1) by evaluating activity correlations across time within
trials between pairs of simultaneously recorded neurons, and (2) by evaluating whether the state
of the local field potential prior to sound onset predicts between-trial fluctuations in activity e.g. [30,
31]. Finally, we determined whether the observed fluctuations are of functional relevance for the
execution of the task by testing the relation between the trial-by-trial fluctuation and which target
the monkey looked at first on that trial.

First, we evaluated correlations in *within-trial* switching based on how A-like vs. how B-
like the responses were across time on individual trials. A total of 91 pairs of triplet conditions
from 34 pairs of simultaneously recorded neurons (from among the 363 triplets used for the
previous analyses) were assessed. For each 50 ms bin of a dual-sound trial in a given triplet, we
assigned a probability score between 0 and 1 that the spike count in the bin was drawn from the
Poisson distribution with rate equaling the bin's sound A rate. The complementary probability
indicated the likelihood that the count was drawn from a Poisson distribution with a rate equaling
the bin's sound B rate (Figure 4A; see Methods: A vs. B assignment scores). We normalized these
probabilities by converting them to Z-scores within a given time bin but across trials, to minimize
the contribution of shared correlations due to stimulus responsiveness or changes in motivational
state across time [32]. We then calculated the neuron-to-neuron correlation coefficients between the
normalized assignment scores across time bins within each trial (i.e. one correlation coefficient
value estimated per trial). This analysis is conceptually similar to conventional cross-correlation

analysis of spike trains in neural pairs, but does not focus on precise timing of spikes or the relative

latency between them [33, 34]. The 50 ms time scale is consistent with the frequency range in which

spike-to-spike coherence has been observed in visual attention paradigms [35].

The observed correlations were generally positive, indicating that the activity was

coordinated within the neural population. Figure 4 illustrates analysis of the dual-sound trials for

a particular triplet in an example pair of neurons (Figure 4A), and the distribution of the mean

neuron-to-neuron correlations in the population for all the dual-sound conditions (Figure 4B). The

distribution of mean correlation coefficients was skewed positive (t-test, $p = 6.8 \times 10^{-6}$). Similar

results were obtained when the raw spike counts were analyzed rather than the assignment scores

(Supplementary Figure 8). This was the case even though we included triplets that were not

categorized as showing wavy behavior in the DAPP analysis. It may be that coordinated activity

fluctuations occur in more neurons than those that met our statistical criteria.

Figure 4. Pairs of neurons tend to show positive within-trial correlations (A) Schematics of the

analysis. Raster plots of two neurons recorded simultaneously; trials shown are for a particular

set of dual-sound conditions. We evaluated the spike count in a given 50 ms time bin, trial, and

member of the neuron pair for a given set of dual-sound conditions to determine if it was more

similar to the spikes evoked during that bin on the corresponding sound A alone or B alone trials

(blue box). We then converted these A vs. B assignment probabilities to a Z-score based on the

mean and standard deviation of the assignment probabilities in that time bin on the other trials

that involved the same stimulus conditions (red box). Finally, we computed the correlation

coefficient between the set of Z score values for a given trial between the pair of simultaneously

*recorded neurons (green box). **(B)** Across the population of pairs of triplets recorded*

simultaneously, the distribution of mean correlation coefficients tended to be positive (t-test

comparing the mean correlation coefficients to zero; p = 6.8 x 10^{-6}). See Supplementary Figure 8

for the same analysis based on spike counts and broken down according to whether the neurons

in the pair shared the same or had different preferences for sound A vs. sound B.

Next, we determined whether the state of the local field potential prior to sound onset predicts between-trial fluctuations in activity. We analyzed the LFP data recorded simultaneously with single unit spiking data. We combined data across triplets, creating two groups of trials based on whether the whole-trial spike count on a given dual-sound trial more closely resembled the responses evoked by sound A alone (where A is the contralateral sound) or sound B alone (see Methods: A vs. B assignment scores). Figure 5A shows the average LFP for the two groups of dual-sound trials. We quantified differences between these two groups with a t-test in the 600ms windows *before* and *after* sound onset (each trial contributed one mean LFP value in each time window). As expected, the LFP signals statistically differed after sound onset in these two trial groupings (red vs. blue traces, time period 0-600 ms, p = 1.0474 X 10^{-05}). But the average LFP voltage also differed *prior* to sound onset (p = 0.0064), suggesting that the state of activity in the local network surrounding an individual neuron at the time of sound onset is predictive of whether the neuron encodes the contralateral or the ipsilateral sound on that particular trial. What exactly that network state consists of is unknown; but it could be an altered balance in the levels of activity in the ICs contralateral and ipsilateral to the first-saccade target.

If fluctuations in neural activity are coordinated across the population, it follows that there should be a relationship between variability in neural activity and behavior. Accordingly, we investigated whether the activity on individual trials predicted whether the monkey would look first to sound A or sound B on that trial. We focused here on the whole-trial time scale as the most robust measure of activity fluctuations in our data. As noted in Methods, we trained the monkeys on sequential sounds first and this training strategy tended to promote performing the task in a stereotyped sequence. Partway through neural data collection, we provided monkey Y with additional training on the non-sequential task, after which that monkey began displaying less

stereotypical behavior and sometimes saccaded first to A and sometimes first to B for a given AB

dual-sound combination (see Figure 5B for example). We then analyzed recording sessions after

this training (N=73 triplets) and we found that at both the whole trial and sub-trial time scales, the

activity of individual neurons was predictive of what saccade sequence the monkey would choose

on that particular trial. Specifically, the average dual-sound AB assignment score for a given triplet

was computed separately for trials in which the first saccade was toward A vs. toward B. The

average scores statistically differed between the two groups of dual-sound trials (t-test, p = 5 x

10^-9, Figure 5C) and in the expected direction, with more A-like scores occurring on trials in

which the monkey looked at A first. This relationship was also present when looking at finer, 50

ms bin time scales (Figure 5D).

Figure 5: Fluctuations can be predicted from pre-stimulus LFP and in turn predict behavior

(A) To assess the relation between the LFP prior *to sound onset and the spiking response* after

sound onset, we assigned each dual-sound trial to one of two groups, A-like or B-like, based on

*whether the spike count matched the response to single-sound A or B (see Methods). We then compared the average LFP voltage (without filtering for any particular frequency band) of the two groups. The average LFP (mean ± SE) is shown in blue for 1917 trials classified as A-like (A=contralateral sound) and in red for 1565 trials classified as B-like. The traces are significantly different in the 600ms before sound onset and in the 600ms after sound onset (two-tailed t-test, p < 0.01). (B,C,D) The target of the first saccade on dual-sound trials can be predicted by the spike count during sound presentation. (**B**) Eye trajectories during dual-sound trials to the same pair of single-sounds (one triplet). The traces are color-coded based on which of the two sounds the monkey looked at first in the response sequence. For clarity, all traces are aligned on a common starting position despite some variation in fixation accuracy. (**C**) The average assignment score of trials in which the monkey looked at sound A first is more A-like than that of trials in which the monkey looked at sound B first. Bars indicate SEM; p value is for a two-tailed t-test involving a total of n=1171 trials (**D**) The relationship between assignment score and first saccade target was also evident at the scale of 50 ms bins (green = positive correlation; * indicate p<0.05 for t-test of assignment score on A-first vs. B-first trials).*

6. Multiplexing across time may be a general brain phenomenon

The problem of encoding multiple simultaneous stimuli is neither limited to the auditory system nor to the localization of sounds, but ubiquitous in the brain and sensory information processing. To gain insight into whether activity fluctuations may contribute to preserving information about multiple stimuli in other systems, we turned to a different sensory system, vision, and a different neural substrate, cortex. Cells in an fMRI-identified face area, the middle fundus (MF) face patch [36] are highly face-selective and strongly tuned to head orientation. It is thus possible to find a stimulus eliciting a strong response in a given MF neuron, and another one that elicits a weak one (e.g. a non-face stimulus or a face at a non-preferred orientation). We found such stimuli for 105 MF neurons and recorded responses during single stimulus presentation or combinations of preferred and non-preferred stimuli within the receptive field (see Methods and [37]). Monkeys maintained fixation throughout stimulus presentation (400 ms).

Results of whole trial analysis are shown in figure 6A: the two most common response patterns at this time scale were Mixtures and Intermediates. The within-trial DAPP analysis (Figure 6B) confirmed that most Mixtures could be classified as showing Flat-Extreme response

patterns. For the Intermediates, in this data set we found little evidence for Flat-Central; rather the largest category showed a Wavy-Central activity pattern, consistent with fluctuations at the sub trial time scale. In short, the evidence for item-preserving activity fluctuations was at least as strong in face patch MF as in the IC data set.

***Figure 6: Evidence of fluctuations in face patch MF**. (**A**) Population results of whole-trial analysis. As in figure 2I, shading indicates the confidence level of the assignment of individual triplets to winning models. (**B**) Population results of DAPP analysis by winning model from whole-trial analysis (win probability > 0.95). Results resemble those found in the IC, except that there was no evidence of Flat-Central among the Intermediates; Wavy-Central was the predominant label for this group among triplets that could be categorized.*

DISCUSSION

Our results show that the activity patterns of IC neurons fluctuate, and that these fluctuations may be consistent with encoding of multiple items in the same processing channels (i.e. the set of neural spike trains occurring in the IC). The time scale of these fluctuations ranges from the level of individual trials down to at least 50 ms bins within a trial. The fluctuations are positively correlated across pairs of neurons (at least, those recorded within the IC on a given side of the brain), are reflective of the state of local field potentials at the time of sound onset, and are predictive of the behavioral response to follow. These fluctuations may be a necessary element of

the neural code, permitting the representation of more than one stimulus across time within a neural

ensemble, and they may serve as a critical element of the coding of natural scenes replete with

many simultaneous stimuli. The fact that similar activity patterns occurred in a completely

different sensory system and brain region, face patch MF, supports this general interpretation.

The notion that aspects of signal fluctuations can be used to encode multiple items in a

limited capacity channel has a history in telecommunications and engineering. Similar strategies

have been postulated to occur in some form in the brain [1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11], but empirical evidence

and statistical methods of assessment have been lacking.

A strength of our statistical approach is that we do not simply assess

unimodality/bimodality (e.g. [10]) but anchor our consideration of the dual stimulus response

distributions to the observed distributions for single stimulus trials. However, there are several

limitations to the present statistical approach. First, the analyses could only be conducted on a

subset of the data, requiring a good fit of a Poisson distribution to the single-sound trials and

adequate separation of the responses on those trials. For the moment, it is unknown whether any

of the excluded data exhibit meaningful response fluctuations. In principle, the modeling approach

can be extended to other types of response distributions which should reduce the amount of data

that is excluded. Second, the full range of time scales at which fluctuations occur is still

undetermined. We focused on the whole-trial timescale and on 50ms bins -- biologically plausible

for both the task and the stimuli. However, any faster fluctuations would likely have been

(erroneously) categorized as Flat-Central in the DAPP model. Indeed, the preponderance of Flat-

Extreme / Mixtures may not reflect the true state of the underlying population but rather the greater

sensitivity of the analysis method for detecting slower fluctuations rather than faster ones.   Third,

our statistical approach based on the DAPP model involves a categorization step that summarizes

the dominant features of a triplet. If a neuron sometimes behaves as a Flat-Extreme type and

sometimes as a Wavy-Central type for a given triplet of conditions, it would likely be categorized

as ambiguous. In other words, even though the DAPP model can pick up composite response

patterns, the results we present ignore the existence of any such patterns.

The observed fluctuations have broad implications because they provide a novel account

linking a number of other well-known aspects of brain function under a common explanation.

First, it is widely recognized that neural firing patterns are highly variable. This variability is often

thought to reflect some fundamental inability of neurons to code information accurately. Here, we

suggest that some of this variability may actually reflect interleaved periods of potentially quite

accurate coding of different items. What else individual neurons may commonly be coding for in

experiments involving presentation of only one stimulus at a time is not known, but possibilities

include stimuli not deliberately presented by the experimenter, memories of previous stimuli, or

mental imagery as suggested by the theory of embodied cognition [38]. Indeed, variability has been

found to be higher when no deliberate stimulus is present at all [39]. In the present study, we were

able to demonstrate signal in these fluctuations by virtue of statistical tests comparing each of the

trial types in A-B-AB triplets, but it may be the case that fluctuations were occurring in the single

stimulus trials as well. We could not test this because our analysis required having as benchmarks

the response distributions corresponding to the potentially encoded items.

Second, as a concept, stimulus multiplexing via activity fluctuations across time provides

insight into why limitations in certain types of cognition exist. Working memory capacity is

limited; attention filters stimuli to allow in-depth processing of a selected set of items. These

limitations may stem from using the same population of neurons for each attended or remembered

item, a situation that may arise whenever neurons are broadly tuned. If this is the case, then the

puzzle becomes why these limits are often greater than one. Stimulus multiplexing across time

suggests that cycling between different items allows evading what might otherwise be a one-item

limit [2]. Here, we investigated only two time scales, 50 ms and whole trials. Additional work is

needed to more fully explore the time scales on which this occurs and to tie the resulting

information on duty cycle to perceptual capacity.

Third, brain oscillations are ubiquitous, have been linked specifically to attentional and

memory processes [31, e.g. 40, see also 41], and have been suggested as having a connection to the coding

of multiple stimuli [2, 3, 4, 5, 6, 7, 8, 42, 43]. Field potential oscillations indicate that neural activity fluctuates,

although they capture only the portion of such fluctuation that is coordinated across the underlying

neural population and is regular in time. It remains to be determined to what degree field potential

oscillations reflect cause vs effect. In other words, field potential oscillations could stem from the

activity of neural circuits involved in controlling multiplexing, or they could reflect the activity of

the neural circuits subject to the effects of such control. In a highly interconnected system such as

the brain, both are likely to occur.

Our findings are particularly relevant to the domain of attention at both the behavioral and

neural mechanism levels. Multiplexing implies that multiple stimuli are handled by a common

circuit in a serial fashion. Serial processing of multiple stimuli has been suggested using several different behavioral paradigms such as searching for or tracking targets among distractors e.g. refs. [44, 45]. There is also evidence for sampling of the visual scene across time: the detectability of visual stimuli can fluctuate in an oscillatory fashion [46], and this is correlated with the phase of EEG signals in the 4-12 Hz range [47, 48]. A similar relationship between auditory perception and EEG phase has also recently been demonstrated [49]. On the whole, this line of investigation suggests that the brain samples stimuli in a periodic or sequential manner.

The scope of such sampling processes may be more extensive than is possible to demonstrate with behavioral assays alone. Our neural analyses do not require that activity fluctuations be either coordinated or periodic at the level of individual neurons. Instead, neural ensembles may fluctuate asynchronously with other ensembles and the fluctuations may not always be periodic. Perceptual studies such as those described above can likely only reveal effects when the underlying neural activity is highly coordinated across the neural population. This would account for why the perceptual impact of a sampling mechanism of the brain is not evident in daily life, but requires experiments designed to elicit or align with maximum synchrony across component neural ensembles to reveal it.

Single-neuron recording studies concerning the impact of attention on (chiefly visual) representations have debated two main possibilities: that attention biases the competition between stimuli for processing in neural populations vs. that it operates as either a filter or spotlight that adjusts the gain of neural responses to unattended vs. attended stimuli (for review, see ref. [50]). The biased competition view bears the closer relationship to multiplexing. Stated strongly, under this theory neurons should respond as if only the attended stimulus is present, ignoring distractors, and the response rate should be the same as if the attended stimulus was the only one presented. If attention were to shift between items across time, as it might have in our study, neural activity would be expected to fluctuate accordingly.

Most research to assess this theory has not investigated fluctuations but has instead pooled the activity across time and trials (see ref. [10] for an exception). The results have generally produced responses to dual stimuli that are intermediate between the responses evoked by each stimulus alone but with a bias in favor of the attended stimulus (e.g. refs. [51, 52]). It will be of interest to ascertain whether such intermediate responses and incomplete bias in favor of the attended stimulus reflects underlying activity fluctuations that serve to preserve information about both

stimuli while enhancing the salience of the attended one. Such a pattern would help account for

the otherwise curious observation that intermediate responses also apparently occur when *neither*

stimulus is attended (e.g. ref. [52]). In short, stimulus multiplexing across time may operate both in

concert with attention and independently of it. We note that we did not attempt to assess whether

attention was shifting between the two stimuli that were presented, so future work will be needed

to tease apart these possibilities.

The need for multiplexing extends to any domain where neural tuning is broad. The

meter/firing rate code for sound location in the IC is one example. The IC's sensitivity to sound

frequency can only partially ameliorate this problem, because here too the tuning is very broad: a

pure tone in the frequency range employed in this study has been shown to evoke activity in 40-

80% of IC neurons[53]. Indeed, about 2/3 of the triplets in the present sample responded to both of

the two sounds individually (Supplementary Figure 5). Most natural sounds are spectrally rich and

will activate hills of neural activity with even greater overlap. That said, it is possible that the

resolution of the frequency map might sharpen via lateral inhibition when more than one sound is

present; such a mechanism might work in concert with stimulus multiplexing across time.

In the case of our particular experimental paradigm, several additional questions arise.

How do signals related to different items come to be multiplexed? Are they later de-multiplexed?

If so, how?

To some degree, sounds are multiplexed in the world. That is, the sound waves from

multiple sources sum in the world and are never purely distinct from one another. The air pressure

waves arriving at each ear reflect the combined contribution of all sound sources. Where and how

signals may be de-multiplexed critically depend on the nature of the representation to which a de-

multiplexed output could be written. In barn owls, which have maps of auditory space, the coding

bottleneck intrinsic to meter/rate coding does not occur, and two sounds produce two separate

active populations [54, 55, 56, 57]. Such distinct peaks suggest that the multiplexed-in-the-air signals have

been de-multiplexed and segregated into two hills of activity.

In primates and several other mammals, neural representations of space employ meters

(rate codes) rather than maps throughout the pathway from sound input to eye movement output,

as far as is currently known [18, 19, 20, 21, 58, 59, 60, 61, 62]. This is the case even at the level of the superior

colliculus [63], an oculomotor structure which has a well-deserved reputation for mapping when

activity is evoked by non-auditory stimuli [64, 65].

Given that different types of codes exist in different species, and given that coding format

is not known in all the circumstances in which multiplexing might apply (e.g. attention, working

memory), we developed two different models to illustrate a range of different de-multiplexing

possibilities (Figure 7) based on the nature of the recipient representation. In the first (Figure 7A),

a multiplexed signal in a meter is converted into two hills of activity in a map, using a basic

architecture involving graded thresholds and inhibitory interneurons suggested previously[66].

Adding an integration mechanism such as local positive feedback loops would then serve to latch

activity on at the appropriate locations in the map, producing a more sustained firing pattern. No

clock signal is necessary for this model.

In the second model (Figure 7B), there are multiple output channels, each capable of

encoding one item. An oscillating circuit that knows about the timing of the input gates signals to

each output channel at the appropriate moments. As in the first model, a local positive feedback

mechanism acts to sustain the activity during the gaps in the input. This model thus retains the

efficient coding format of a meter but requires a controlling signal with knowledge of when to

latch input flow through to each output channel. In our data, fluctuations were at least somewhat

coordinated across pairs of simultaneously recorded neurons, in agreement with this model. It is

possible that within-trial fluctuating units lie at the input stage of such a circuit, and that between-

trial fluctuating units actually lie at the output stage. A given unit might be allocated to either the

A or the B pools based on state of the network (as detected by the LFP measurements) on different

trials.

Although the stimulus multiplexing across time that we observed here has parallels in

engineering (particularly with time division multiplexing in telecommunications) these two

models also help illustrate that multiplexing is unlikely to occur exactly the same way in biological

systems as it does in technological ones. For example, the time course may be more fluid in

biological systems, and different neural ensembles may operate asynchronously and/or

aperiodically as opposed to operating under a regular, shared, clock cycle of some kind. The

strongest shared element of the analogy with telecommunications is the transmission of more than

one item or type of information via signal fluctuations at some time scale.

Figure 7. Two possible mechanisms for de-multiplexing a fluctuating signal. *A clock signal that*
*knows about coding transitions is not necessarily needed if signals are read out into a map (**a**),*
*but is required if signals are retained in a meter or rate-coded format (**b**). Both models have an*
*input signal that employs a meter code for sound location (purple, **c** top panel), and this signal is*
*assumed to fluctuate between two response levels when two sounds are present (**c** bottom panel).*
In the meter-to-map model (a), the second stage consists of group of excitatory neurons (open
circles) with varying thresholds, paired with inhibitory neurons (filled circles) that have slightly
higher thresholds. These neurons all receive input from the input meter signal. An individual
excitatory neuron is activated when the input signal exceeds its own threshold and is lower than
the threshold of the paired inhibitory interneuron, producing tuning curves like those shown in
*panel **d**. The net drive across time to two examples A and B is shown schematically as the thin gray*
*line in the two inset graphs (**e**). These two neurons would turn on and off out of phase with each*
other based on their external inputs alone. Adding a positive autofeedback loop to each excitatory
neuron in the map (green) provides integration of the activity of each neuron and permits the
activity to be sustained across periods of time when there is no external drive (dark line,). This
model is derived from a portion of the Vector Subtraction model of [66]. In the meter-to-meter model
*(**b**), the input is forked to an A meter channel neuron and a B meter channel neuron. A bistable*
oscillator coupled to the same unknown clock (shared timing signal) that controls the input

*fluctuations would be needed to appropriately route the output to these two units. The resulting output, in the absence of positive autofeedback, would also fluctuate but between an off state and a level that corresponds to signaling the presence of sound A or B respectively (**f** gray lines). Adding positive autofeedback would again allow bridging across the off states (dark lines).*

Similarity of results in the IC with auditory stimuli and MF with visual stimuli suggest that multiplexing may be a general mechanism that is commonly at play to enhance the total processing power of the brain. The statistical tools developed here can be applied to any triplet data. Additional studies with both single stimulus conditions, to define the distributions of signals, and dual stimulus conditions, to evaluate fluctuations between membership in those distributions, will be important for delineating the extent of this phenomenon. Digging under the hood of the time-and-trial pooled activity to look at activity patterns on a moment by moment basis will be essential to advancing our understanding of how the brain operates dynamically to maximize its processing power.

REFERENCES

1. Meister M. Multineuronal codes in retinal signaling. *Proc Natl Acad Sci U S A* **93**, 609-614 (1996).

2. Lisman JE, Idiart MA. Storage of 7 +/- 2 short-term memories in oscillatory subcycles. *Science* **267**, 1512-1515 (1995).

3. Hoppensteadt FC, Izhikevich EM. Thalamo-cortical interactions modeled by weakly connected oscillators: could the brain use FM radio principles? *Bio Systems* **48**, 85-94 (1998).

4. Akam T, Kullmann DM. Oscillatory multiplexing of population codes for selective communication in the mammalian brain. *Nat Rev Neurosci* **15**, 111-122 (2014).

5. Tallon-Baudry C. The roles of gamma-band oscillatory synchrony in human visual cognition. *Front Biosci (Landmark Ed)* **14**, 321-332 (2009).

6. Lisman JE, Jensen O. The theta-gamma neural code. *Neuron* **77**, 1002-1016 (2013).

7. Siegel M, Warden MR, Miller EK. Phase-dependent neuronal coding of objects in short-term memory. *Proc Natl Acad Sci U S A* **106**, 21341-21346 (2009).

8. McLelland D, VanRullen R. Theta-Gamma Coding Meets Communication-through-Coherence: Neuronal Oscillatory Multiplexing Theories Reconciled. *PLoS computational biology* **12**, e1005162 (2016).

9. Jezek K, Henriksen EJ, Treves A, Moser EI, Moser MB. Theta-paced flickering between place-cell maps in the hippocampus. *Nature* **478**, 246-249 (2011).

678   10.   Li K, Kozyrev V, Kyllingsbaek S, Treue S, Ditlevsen S, Bundesen C. Neurons in Primate Visual Cortex
679   Alternate between Responses to Multiple Stimuli in Their Receptive Field. *Front Comput Neurosci* **10**, 141
680   (2016).
681
682   11.   Panzeri S, Brunel N, Logothetis NK, Kayser C. Sensory neural codes using multiplexed temporal scales.
683   *Trends Neurosci* **33**, 111-120 (2010).
684
685   12.   Adams JC. Ascending projections to the inferior colliculus. *J Comp Neurol* **183**, 519-538 (1979).
686
687   13.   Moore RY, Goldberg JM. Ascending projections of the inferior colliculus in the cat. *J Comp Neurol* **121**,
688   109-135. (1963).
689
690   14.   Aitkin LM, Phillips SC. Is the inferior colliculus an obligatory relay in the cat auditory system?
691   *Neuroscience Letters* **44**, 259-264 (1984).
692
693   15.   Jenkins WM, Masterton RB. Sound localization: effects of unilateral lesions in central auditory system. *J*
694   *Neurophysiol* **47**, 987-1016. (1982).
695
696   16.   Kelly JB, Kavanagh GL. Sound localization after unilateral lesions of inferior colliculus in the ferret (
697   Mustela putorius). *J Neurophysiol* **71**, 1078-1087 (1994).
698
699   17.   Litovsky RY, Fligor BJ, Tramo MJ. Functional role of the human inferior colliculus in binaural hearing.
700   *Hear Res* **165**, 177-188 (2002).
701
702   18.   Groh JM, Kelly KA, Underhill AM. A monotonic code for sound azimuth in primate inferior colliculus.
703   *Journal of Cognitive Neuroscience* **15**, 1217-1231 (2003).
704
705   19.   McAlpine D, Grothe B. Sound localization and delay lines--do mammals fit the model? *Trends Neurosci*
706   **26**, 347-350 (2003).
707
708   20.   Grothe B, Pecka M, McAlpine D. Mechanisms of sound localization in mammals. *Physiol Rev* **90**, 983-
709   1012 (2010).
710
711   21.   Groh JM. Moving with maps and meters. In: *Making space: How the brain knows where things are*
712   (ed^(eds). Harvard University Press (2014).
713
714   22.   Perrott DR. Discrimination of the spatial distribution of concurrently active sound sources: some
715   experiments with stereophonic arrays. *J Acoust Soc Am* **76**, 1704-1712 (1984).
716
717   23.   Perrott DR. Concurrent minimum audible angle: a re-examination of the concept of auditory spatial acuity.
718   *J Acoust Soc Am* **75**, 1201-1206 (1984).
719
720   24.   Best V, van Schaik A, Carlile S. Separation of concurrent broadband sound sources by human listeners. *J*
721   *Acoust Soc Am* **115**, 324-336 (2004).
722
723   25.   Carandini M, Heeger DJ, Movshon JA. Linearity and normalization in simple cells of the macaque primary
724   visual cortex. *J Neurosci* **17**, 8621-8644 (1997).
725
726   26.   Movshon JA, Thompson ID, Tolhurst DJ. Spatial summation in the receptive fields of simple cells in the
727   cat's striate cortex. *J Physiol* **283**, 53-77 (1978).
728
729   27.   Cash S, Yuste R. Linear summation of excitatory inputs by CA1 pyramidal neurons. *Neuron* **22**, 383-394
730   (1999).
731
28.   Alvarado JC, Vaughan JW, Stanford TR, Stein BE. Multisensory versus unisensory integration: contrasting
modes in the superior colliculus. *J Neurophysiol* **97**, 3193-3205 (2007).

29. Berger JO, Pericchi LR. The Intrinsic Bayes Factor for Model Selection and Prediction. *J Am Stat Assoc*
91, 109-122 (1996).

30. Haslinger R, Ulbert I, Moore CI, Brown EN, Devor A. Analysis of LFP phase predicts sensory response of
barrel cortex. *J Neurophysiol* **96**, 1658-1663 (2006).

31. Vanrullen R, Busch NA, Drewes J, Dubois J. Ongoing EEG Phase as a Trial-by-Trial Predictor of
Perceptual and Attentional Variability. *Front Psychol* **2**, 60 (2011).

32. Metzger RR, Greene NT, Porter KK, Groh JM. Effects of reward and behavioral context on neural activity
in the primate inferior colliculus. *J Neurosci* **26**, 7468-7476 (2006).

33. Gerstein GL, Perkel DH, Subramian KN. Identification of functionally related neural assemblies. *Brain Res*
140, 43-62 (1978).

34. Alonso JM, Usrey WM, Reid RC. Precisely correlated firing in cells of the lateral geniculate nucleus.
Nature **383**, 815-819 (1996).

35. Mitchell JF, Sundberg KA, Reynolds JH. Spatial attention decorrelates intrinsic activity fluctuations in
macaque area V4. *Neuron* **63**, 879-888 (2009).

36. Freiwald WA, Tsao DY. Functional compartmentalization and viewpoint generalization within the
macaque face-processing system. *Science* **330**, 845-851 (2010).

37. Ebihara AF. Normalization among heterogeneous population confers stimulus discriminability on the
macaque face patch neurons. . (ed^(eds). Rockefeller University (2015).

38. Barsalou LW. Grounded cognition. *Annu Rev Psychol* **59**, 617-645 (2008).

39. Churchland MM*, et al.* Stimulus onset quenches neural variability: a widespread cortical phenomenon. *Nat*
Neurosci **13**, 369-378 (2010).

40. Raghavachari S*, et al.* Gating of human theta oscillations by a working memory task. *J Neurosci* **21**, 3175-
3183 (2001).

41. Engel TA, Steinmetz NA, Gieselmann MA, Thiele A, Moore T, Boahen K. Selective modulation of cortical
state during spatial attention. *Science* **354**, 1140-1144 (2016).

42. Kienitz R, Schmiedt JT, Shapcott KA, Kouroupaki K, Saunders R, Schmid MC. Rhythmic neural spiking
and attentional sampling arising from cortical receptive field interactions. *bioRxiv*, (2018).

43. Rollenhagen JE, Olson CR. Low-frequency oscillations arising from competitive interactions between
visual stimuli in macaque inferotemporal cortex. *J Neurophysiol* **94**, 3368-3387 (2005).

44. Triesman A, Gelade G. A feature integration theory of attention. *Cognit Psychol* **12**, 97-136 (1980).

45. Holcombe AO, Chen WY. Splitting attention reduces temporal resolution from 7 Hz for tracking one object
to <3 Hz when tracking three. *J Vis* **13**, 12 (2013).

46. Landau AN, Fries P. Attention samples stimuli rhythmically. *Curr Biol* **22**, 1000-1004 (2012).

47. Busch NA, Dubois J, VanRullen R. The phase of ongoing EEG oscillations predicts visual perception. *J*
Neurosci **29**, 7869-7876 (2009).

48.   Busch NA, VanRullen R. Spontaneous EEG oscillations reveal periodic sampling of visual attention. *Proc*
790   *Natl Acad Sci U S A* **107**, 16048-16053 (2010).
791

792   49.   Ng BS, Schroeder T, Kayser C. A precluding but not ensuring role of entrained low-frequency oscillations
793   for auditory perception. *J Neurosci* **32**, 12268-12276 (2012).
794

795   50.   Desimone R, Duncan J. Neural correlates of selective visual attention. *Ann Rev Neurosci* **18**, 193-222
796   (1995).
797

798   51.   Moran J, Desimone R. Selective attention gates visual processing in the extrastriate cortex. *Science* **229**,
799   782-784 (1985).
800

801   52.   Reynolds JH, Chelazzi L, Desimone R. Competitive mechanisms subserve attention in macaque areas V2
802   and V4. *J Neurosci* **19**, 1736-1753 (1999).
803

804   53.   Bulkin DA, Groh JM. Systematic mapping of the monkey inferior colliculus reveals enhanced low
805   frequency sound representation. *J Neurophysiol* **105**, 1785-1797 (2011).
806

807   54.   Baxter CS, Nelson BS, Takahashi TT. The role of envelope shape in the localization of multiple sound
808   sources and echoes in the barn owl. *J Neurophysiol* **109**, 924-931 (2013).
809

810   55.   Takahashi TT, Keller CH, Nelson BS, Spitzer MW, Bala AD, Whitchurch EA. Object localization in
811   cluttered acoustical environments. *Biol Cybern* **98**, 579-586 (2008).
812

813   56.   Keller CH, Takahashi TT. Localization and identification of concurrent sounds in the owl's auditory space
814   map. *J Neurosci* **25**, 10446-10461 (2005).
815

816   57.   Takahashi TT, Keller CH. Representation of multiple sound sources in the owl's auditory space map.
817   *Journal of Neuroscience* **14**, 4780-4793 (1994).
818

819   58.   Salminen NH, May PJ, Alku P, Tiitinen H. A population rate code of auditory space in the human cortex.
820   *PLoS One* **4**, e7600 (2009).
821

822   59.   Salminen NH, Tiitinen H, Yrttiaho S, May PJ. The neural code for interaural time difference in human
823   auditory cortex. *J Acoust Soc Am* **127**, EL60-65 (2010).
824

825   60.   Werner-Reiss U, Groh JM. A rate code for sound azimuth in monkey auditory cortex: implications for
826   human neuroimaging studies. *J Neurosci* **28**, 3747-3758 (2008).
827

828   61.   Woods TM, Lopez SE, Long JH, Rahman JE, Recanzone GH. Effects of stimulus azimuth and intensity on
829   the single-neuron activity in the auditory cortex of the alert macaque monkey. *J Neurophysiol* **96**, 3323-
830   3337 (2006).
831

832   62.   Ashida G, Carr CE. Sound localization: Jeffress and beyond. *Curr Opin Neurobiol* **21**, 745-751 (2011).
833

834   63.   Lee J, Groh JM. Different stimuli, different spatial codes: a visual map and an auditory rate code for
835   oculomotor space in the primate superior colliculus. *PLoS One* **9**, e85017 (2014).
836

837   64.   Robinson DA. Eye movements evoked by collicular stimulation in the alert monkey. *Vision Res* **12**, 1795-
838   1807 (1972).
839

840   65.   Mohler CW, Wurtz RH. Organization of monkey superior colliculus: intermediate layer cells discharging
841   before eye movements. *J Neurophysiol* **39**, 722-744 (1976).
842

843   66.   Groh JM, Sparks DL. Two models for transforming auditory signals from head-centered to eye- centered
844   coordinates. *Biological Cybernetics* **67**, 291-302 (1992).

845
846   67.   Barton EJ, Sparks DL. Saccades to remembered targets exhibit enhanced orbital position effects in
847   monkeys. *Vision Res* **41**, 2393-2406. (2001).
848
849   68.   Metzger RR, Mullette-Gillman OA, Underhill AM, Cohen YE, Groh JM. Auditory saccades from different
850   eye positions in the monkey: implications for coordinate transformations. *J Neurophysiol* **92**, 2622-2627
851   (2004).
852
853   69.   Berger J. The case for objective Bayesian analysis. *Bayesian Anal* **1.3**, 385-402 (2006).
854
855   70.   Berger JO, Pericchi LR. The intrinsic Bayes factor for model selection and prediction. *J Am Stat Assoc*
856   **91.433**, 109-122 (1996).
857
858   71.   Escobar MD, West M. Bayesian Density-Estimation and Inference Using Mixtures. *J Am Stat Assoc* **90**,
859   577-588 (1995).
860
861
862
863

METHODS

IC data

General procedures

All procedures conformed to the guidelines of the National Institutes of Health (NIH Pub.

No. 86-23, Revised 1985) and were approved by the Institutional Animal Care and Use Committee

of Duke University. Two adult rhesus monkeys (Macaca mulatta) participated (monkey P, and

monkey Y, both female). Under general anesthesia and in sterile surgery we first implanted a head

post holder to restrain the head and a scleral search coil to track eye movements (Robinson 1963;

Judge et al. 1980). After recovery with suitable analgesics and veterinary care, we trained the

monkeys in the experimental task. In a second surgery, we implanted a recording cylinder (2 cm

diameter) over the right (monkey Y) or left (monkey Y, P) IC respectively. We determined the

location of the cylinder with stereotactic coordinates and verified it with MRI scans e.g. [53].

Sound localization task

The monkeys performed a single- or dual-sound localization task (Figure 1B) by making

saccades toward one or two simultaneously-presented auditory targets with one or two saccades

as appropriate. All sound targets were located in front of the monkey at eye level; the horizontal

location, frequency and intensity were varied pseudorandomly as described below (Recording

Procedures). Each trial began with 600-700ms of fixation of a visual stimulus (light emitting-

diode, LED, located straight ahead and 10-14° below the speakers). During fixation we presented

one sound (single-sound trials) or two simultaneous sounds (dual-sound trials). After a fixation

time of either 600-800 (Data Set I, some of Data Set II) or 1000-1100 (remainder of Data Set II),

the fixation light was extinguished and the monkey was required to make a single saccade on

single-sound trials or a sequence of two saccades (in either order) on dual-sound trials. Trials were

considered correct if each saccade was directed within 10-17.5 degrees horizontally and 20-40

degrees vertically of a target due to vertical inaccuracies in localizing non-visual targets in primates, [67] and if

the gaze was maintained on the final target for 100 – 200 ms. On correct trials monkeys were

rewarded with juice drops.

Training

Training was accomplished in three stages. We initially trained the monkeys to report the

location of single visual targets by saccading to them. We then introduced single auditory targets.

As these were novel and unexpected in the silent experimental booth, monkeys readily saccaded

to them [68]. To help the monkeys calibrate their auditory saccades, a visual feedback was added on

trials where the auditory saccade was not initiated correctly within 700 ms. The feedback was

presented only at the most peripheral target locations (+/- 24 degree) and only for a few initial days

of training. Finally, we trained monkey to localize dual-sound targets. Initially we presented the

two sounds sequentially in a specific order, then we gradually reduced the temporal gap between

them until the sounds were simultaneous.

In the final version of the task, monkeys were allowed to look at the targets in either order,

as noted above. However, due to the initial training with sequential sounds, they retained

stereotyped patterns of saccades in which they tended to look first to whichever sound location

had been presented first during the sequential and partial overlap stages of training. Monkey P was

trained with more central target locations (e.g. -6 or 6 degree targets) initially occurring first and

more peripheral targets (e.g. -24 or 24 degree targets) occurring second, and monkey Y was trained

with sounds initially occurring in the opposite sequence. Midway through neural data collection,

we provided additional training to monkey Y to encourage free choice of which sound to look at

first. This allowed us to investigate the relationship between each behavioral response and the

neural representation at that moment.

Recording procedure and strategy

Recordings were made with one or two tungsten electrodes (FHC, impedance between 1
and 3 MΩ at 1 kHz). Each electrode was lodged in a stainless-steel guide tube (manually advanced
through the dura) and controlled independently with an oil hydraulic pulse micropositioner
(Narishige International USA, Inc. and NAN INSTRUMENTS LTD, Israel). First, we localized
the IC (and isolated single neurons) while the monkey listened passively to sounds of different
frequencies. We then collected single unit spiking activity and local field potential while the
monkey performed the single- and dual-sound localization tasks. We used a Multichannel
Acquisition Processor (MAP system, Plexon Inc., Dallas, TX) and Sort Client software. The single
unit spiking activity was filtered between 150 Hz and 8 kHz and sampled at 20 kHz, while the LFP
signal was filtered between 0.7 and 300 Hz and sampled at either 20kHz or 1kHz (see Local Field
Potential). Data were collected as long as the neurons were well isolated and the monkey
performed the tasks.

Neural signals were recorded primarily from two functionally-defined subregions of the
IC, the low frequency area and the tonotopic area [53]. Neurons in the low frequency tuned area
generally respond best to low frequencies and there is little heterogeneity in tuning, whereas
neurons recorded in the tonotopic area had best frequencies that could be either low or high
depending on the position of the recording electrode.

Data Sets and Sound Stimuli: Locations, Frequencies, Levels

The spiking activity of 166 single neurons was recorded in two datasets involving the same
task but differing in which sound levels and frequencies were included. A total of 68 of these
neurons were recorded as pairs from separate electrodes positioned in the IC on the same side of
the brain at a minimum spatial separation of 2 mm. Local field potentials (LFP) were also recorded
from 87 of these recording sites.

In both datasets, the sounds consisted of bandpass noise with a bandwidth of +/- 200 Hz.
On dual-sound trials, the sounds were delivered from pairs of locations (24 degrees and -6
degrees), and (-24 and +6 degrees) i.e. 30 degrees apart. The two sounds differed in frequency,
with one of the two sounds having a 742 Hz center frequency and the other differing by at least
0.285 octaves or multiples of this distance. Single-sound trials involved the same set of locations
and frequencies as on dual-sound trials, but with only a single-sound presented at a time. All

sounds were <u>frozen</u> within an individual session; that is, all trials with a given set of auditory
parameters involved the same time series signal delivered to the relevant speaker.

In data set I (N=98 neurons), the sounds presented on dual-sound trials were 742 Hz and a
sound from the set (500, 609, 903, 1100 Hz); these frequencies were ±0.285 octave or ± 0.57
octaves above or below 742 Hz, or ±3.4 and 6.8 semitones. Combining two sounds will produce a
combination that is louder than either component. Sound levels were therefore calibrated to
provide two sets of conditions: dual sounds for which the component sounds involve the *same*
signals to the audio speakers as on single-sound trials, producing a *louder* dual sound, and dual
sounds for which the level of the component sounds was *reduced* so that the overall *loudness* was
the *same* on dual as on single trials. The levels used for the components were 51 and 55 dB,
producing sound levels of minimum 55 or maximum 60 dB on dual-sound trials. The same-signal
comparison involved using the 55 dB component levels, singly and on dual-sound trials. The
same-loudness comparison involved using the 55 dB levels on single-sound trials and the 51 dB
levels for the components of dual-sound trials. Calibrations were performed using a microphone
(Bruel and Kjaer 2237 sound level meter) placed at the position normally occupied by the animal's
head.

Because results did not differ substantively when comparisons were made between same-
signal and same-loudness conditions (Supplementary Figure 2 vs. Supplementary Figure 3), we
pooled across sound levels for subsequent analyses, and we dispensed with the multiple sound
levels for data set II (monkey Y only, N=68 neurons), using either 50 or 55 dB levels for all
components. We also incorporated additional sound frequencies, [1340 1632 1988 Hz], to improve
the odds that responses to each of the component sounds differed significantly. Again, one of the
two sounds on dual-sound trials was 742 Hz; the other sound frequency was either from the original
list of [500 609 903 1100] or from the new frequencies. Most of the neurons in this data set were
tested with [500 742 1632].

Cell Inclusion/Exclusion criteria and trial counts

The N=166 neurons (N=98 from Data Set I and N=68 from Data Set II) included for
analysis were drawn from a larger set of 325 neurons. Neurons were excluded from analysis if the
neuron proved unresponsive to sound (Student's t-test, spike counts during the 600 ms after sound
onset compared to the same period immediately prior to sound onset, one-tailed, p>0.05), or if

there were too few correct trials (minimum of five correct trials for each of the components [A, B,
and AB trials] that formed a given triplet of conditions or if there were technical problems during
data collection (e.g. problems with random interleaving of conditions or with computer crashes).
The average number of correct trials for a given set of stimulus conditions in the included dataset
(N=166) was 10.5 trials. The total number of included triplets was 1484.
981

Data Analysis
All analyses concerned correctly performed trials.
Summation vs. Averaging in time-and-trial pooled activity
To evaluate IC activity using conventional analysis methods that pool across time and/or
across trials, we counted action potentials during two standard time periods. The baseline period
(*Base*) was the 600ms period before target onset, and the sensory-related target period (*Resp*) was
the 600ms period after target onset (i.e. ending before, or at the time of, the offset of the fixation
light, Figure 1B).
Summation/Averaging Indices: We quantified the activity on dual-sound trials in
comparison to the sum and the average of the activity on single-sound trials, expressed in units of
standard deviation (Z-scores), similar to a method used by [28]. Specifically, we calculated,
$PredictedSum_{A,B} = mean(Resp_A) + mean(Resp_B) - mean(Base_{A,B})$ (1)
and
$PredictedAvg_{A,B} = (mean(Resp_A) + mean(Resp_B))/2$ (2)
where *Resp_A* and *Resp_B* were the number of spikes of a given neuron for a given set of single-sound
conditions *A* and *B* (location, frequency, and intensity) that matched the component sounds of the
dual-sound trials being evaluated. As the response may actually include a contribution from
spontaneous baseline activity, we subtracted the mean of the baseline activity for the single-sounds
(*Base_{A,B}*). Without this subtraction, the predicted sum would be artificially high because two copies
of baseline activity are included under the guise of the response activity.
The Z scores for the dual-sound trials were computed by subtracting these predicted values
from the mean of the dual-sound trials (mean(Resp_{AB})) and dividing by the mean of the standard
deviations of the responses on single-sound trials:
$$Zsum_{AB} = \frac{mean(Resp_{AB}) - PredictedSum_{A,B}}{mean(std(Resp_A), std(Resp_B))}$$ (3)
and

$$1007 \quad ZAvg_{AB} \; = \frac{mean\left(Resp_{AB}\right) - PredictedAvg_{A,B}}{mean(std\left(Resp_A\right), std\left(Resp_B\right))} \tag{4}$$

1008 If the dual response was within +/- 1.96 standard deviations of the predicted sum or predicted

1009 average, we could say the actual dual response was within the 95% confidence intervals for

1010 addition or averaging of two single responses, respectively.

1011

1012 **Analysis of whole-trial fluctuations and inclusion criteria**

1013

Our statistical tests for fluctuations in neural firing were conducted on triplets, or related

sets of single and dual-sound trials (A, B, AB trials). To evaluate whether neural activity fluctuates

across trials in a fashion consistent with switching between firing patterns representing the

component sounds, we evaluated the Poisson characteristics of the spike trains on matching dual

and single-sound trials (triplets: AB, A and B). Spike train data from each trial was summarized

by the total spike count between 0-600ms or 0-1000 ms from sound onset (i.e. whatever the

minimum duration of the overlap between fixation and sound presentation was for that recorded

neuron, see section Sound localization task). We modeled the distribution of spike counts in

response to single-sounds A and B as Poisson distributions with unknown rates λ^A, denoted

$Poi(\lambda^A)$, and λ^B, denoted $Poi(\lambda^B)$. Four hypotheses were considered for the distribution of sound

AB spike counts:

1. a mixture distribution $\alpha \cdot Poi(\lambda^A) + (1 - \alpha) \cdot Poi(\lambda^B)$ with an unknown mixing weight α

(Mixture)

2. a single $Poi(\lambda^{AB})$ with some λ^{AB} in between λ^A and λ^B (Intermediate)

3. a single $Poi(\lambda^{AB})$ where λ^{AB} is either larger or smaller than both λ^A and λ^B (Outside)

4. a single $Poi(\lambda^{AB})$ where λ^{AB} exactly equals one of λ^A and λ^B (Single)

Relative plausibility of these competing hypotheses was assessed by computing

their posterior probabilities with equal prior weights (1/4) assigned to the models, and with

default Jeffreys' prior [69] on model specific Poisson rate parameters, and a uniform prior on

the mixing weight parameter α. The Jeffreys' prior was truncated to appropriate ranges for

the Intermediate and Outside models. Posterior model probabilities were calculated by

computation of relevant intrinsic Bayes factors [70].

Triplets were excluded if either of the following applied: 1) the Poisson assumption on A and B trial counts was not supported by data; or 2) λ^A and λ^B were not well separated. To test the Poisson assumption on single-sound trials A and B of a given triplet, we used an approximate chi-square goodness of fit test with Monte Carlo p-value calculation. For each sound type, we estimated the Poisson rate by averaging counts across trials. Equal probability bins were constructed from the quantiles of this estimated Poisson distribution, with number of bins determined by expected count of 5 trials in each bin or at least 3 bins -- whichever resulted in more bins. A lack-of-fit statistic was calculated by summing across all bins the ratio of the square of the difference between observed and expected bin counts to the expected bin count. Ten thousand Monte Carlo samples of Poisson counts, with sample size given by the observed number of trials, were generated from the estimated Poisson distribution and the lack-of-fit statistic was calculated from each one of these samples. P-value was calculated as the proportion of these Monte Carlo samples with lack-of-fit statistic larger than the statistic value from the observed data. Poisson assumption was considered invalid if the resulting Monte Carlo p-value < 0.1.

For triplets with valid Poisson assumption on sound A and B spike counts, we tested for substantial separation between λ^A and λ^B, by calculating the intrinsic Bayes factor of the model $\lambda^A \neq \lambda^B$ against $\lambda^A = \lambda^B$ with the non-informative Jeffreys' prior on the λ parameters: λ^A, λ^B or their common value. The triplet was considered well separated in its single sounds if the logarithm of the intrinsic Bayes factor equaled 3 or more, which is the same as saying the posterior probability of $\lambda^B \neq \lambda^A$ exceeded 95% when a-priori the two models were given 50-50 chance.

It should be noted that the sensitivity/specificity of detection were not equal across the four competing hypotheses. Because the Single hypothesis is a limiting case of each of the other three hypotheses, the method's sensitivity to this response pattern is lower than for the other three possible outcomes. This was verified on simulated data for which the truth could be known; simulated Mixtures, Intermediates, and Outsides were commonly correctly categorized with >95% confidence, whereas Singles were correctly categorized but at a lower level of confidence.

Dynamic Admixture Point Process Model

To evaluate whether neural activity fluctuates *within* trials, we developed a novel analysis

method we call a Dynamic Admixture Point Process model (DAPP) which characterized the

dynamics of spike trains on dual-sound trials as an admixture of those occurring on single-sound

trials. The analysis was carried out by binning time into moderately small time intervals. Given a

predetermined bin-width $w = T/C$ for some integer C, we divided the response period into

contiguous time intervals $I_1 = [0;w); I_2 = [w; 2w)\ldots I_C = [(C-1)w, Cw)$ and reduced each trial to a

C-dimensional vector of bin counts $(X^e_{j1},\ldots,X^e_{jC})$ for $e \in \{A;B;AB\}$ and $j = 1,\ldots, n_e$. Mathematically,

$X^e_{jC} = N^e_j(Ic)$. The results reported here were based on $w = 50$ (with time measured in ms and $T =$

600 or 1000), but we also repeated the analyses with $w = 25$ and noticed little difference.

Our model for the bin counts was the following. Below we denote by t^*_c the mid-point $(c -$

$1/2)w$ of sub-interval I_c.

1. $X^e_{jc} \sim Poi(w \cdot \lambda^e(t^*_c))$, $e \in \{A, B\}$, $c \in \{1, \ldots, C\}$, $j \in \{1, \ldots, n_e\}$. We assume both $\lambda^A(t)$ and

$\lambda^B(t)$ are smooth functions over $t \in [0, T]$.

2. $X^{AB}_{jc} \sim Poi(w \cdot \lambda_j(t^*_c))$, where $\lambda_j(t) = \lambda_j(t) = \alpha_j(t)\lambda^A(t) + \{1 - \alpha_j(t)\}\lambda^B(t)$ with $\alpha_j \colon [O, T] \to$

$(0,1)$ being unknown smooth functions.

We modeled $\alpha_j(t) = S(\eta_j(t))$, where $S(t) = 1/(1 + e^{-t})$, and, each $\eta_j(t)$ was taken to

be a (smooth) Gaussian process with $E\{\eta_j(t)\} \equiv \phi_j, Var\{\eta_j(t)\} \equiv \psi_j$, and, $Cor\{\eta_j(t), \eta_j(t')\} =$

$\exp\{-0.5(t - t')^2/\ell_j^2\}$. The three parameters (ϕ_j, ψ_j, ℓ_j) respectively encoded the *long-term*

average value, the *total swing magnitude* and the *waviness* of the $\alpha_j(t)$ curve.—Intuitively, these

parameters can be thought of as related to the means and variances of the distribution of weight

values α regardless of time within a trial, as well as the correlation between the weight observed

at one point in time and the weight observed at another on any given trial. While the temporal

imprint carried by each α_j was allowed to be distinct, we enforced the dual trials to share dynamic

patterns by assuming $(\phi_j, \psi_j, \ell_j), j = 1, \ldots, n_{AB}$, were drawn from a common, unknown

probability distribution P, which we called a dynamic pattern generator and viewed as a

characteristic of the triplet to be estimated from the data.

To facilitate estimation of P, we assumed it decomposed as $P = P_{\phi\psi} \times P_\ell$, where $P_{\phi\psi}$ was

an unknown distribution on $(-\infty, \infty) \times (0, \infty)$ generating (ϕ_j, ψ_j), and, P_ℓ was an unknown

distribution on $(0, \infty)$ generating ℓ_j. To simplify computation, we restricted ℓ_j to take only a

finitely many positive values, representative of the waviness range we are interested in (in our

analyses, we took these representative values to be {75, 125, 200, 300, 500}, all in ms). This

restricted P_ℓ to be a finite dimensional probability vector.

We performed an approximate Bayesian estimation of model parameters. Note

that only $\lambda^A(t)$ and $\lambda^B(t)$ were informed by the single-sound trial data. All other model parameters

were informed only by the dual-sound trial data conditionally on the knowledge of $\lambda^A(t)$ and

$\lambda^B(t)$. To take advantage of this partial factorization of information sources, we first smoothed

each set of single-sound trial data to construct a conditional gamma prior for the corresponding

$\lambda^e(t_c^*), e \in \{A, B\}, c = 1, \dots, C$, where the gamma distribution's mean and standard deviation were

matched with the estimate and standard error of $\lambda^e(t_c^*)$. A formal Bayesian estimation was then

carried out on all model parameters jointly by (a) using only the dual-sound trial data, (b) utilizing

the conditional gamma priors on $\lambda^A(t)$ and $\lambda^B(t)$, and, (c) assuming a Dirichlet process prior [71]

on $P_{\phi\psi}$ and an ordinary Dirichlet prior on P_ℓ. This final step involved a Markov chain Monte Carlo

computation whose details will be reported in a separate paper.

Next, the estimate of the generator P was utilized to repeatedly simulate $\alpha(t)$ functions for

hypothetical, new dual trials for the triplet. For each simulated $\alpha(t)$ curve, we computed its

maximum swing size $|\alpha| = max_t\, \alpha(t) - min_t \alpha(t)$, and, time aggregated average value $\bar\alpha =$

$\int_0^T \alpha(t)dt / T$. The waviness index of the triplet was computed as the odds of seeing an $\alpha(t)$

function exhibiting a swing of at least 50% between its peak and trough:

1115
$$r_w = \frac{P(|\alpha|>0.5)}{P(|\alpha|<0.5)}$$

1116 where P denotes the sampling proportion of the simulated α draws. The triplet's extremeness index

1117 was computed as the odds of seeing an $\alpha(t)$ function with its long-term average $\bar\alpha$ being closer to

1118 the mid-way mark of 50% than the extremes:

1119
$$r_c = \frac{P(\bar\alpha \in (0.25, 0.75))}{P(\bar\alpha \notin (0.25, 0.75))}.$$

The two indices were then thresholded to generate a 2-way classification of all triplets. On

waviness, a triplet was categorized as Wavy, Flat or Ambiguous according to whether $r_w > 1.3$,

$r_w < 0.77$, or, $0.77 \leq r_w \leq 1.3$, respectively On extremeness, the categories were Central,

Extreme, or, Ambiguous according to whether $r_c > 3.24$, $r_c < 1.68$, or, $1.68 \leq r_c \leq 3.24$,

respectively.

In addition to the Flat/Wavy and Extreme/Central classification, a third parameter was

evaluated for each triplet: the degree of skewness in the distribution of $\overline{\alpha_*}$:

$$r_S = max\left(\left\{\frac{P(\overline{\alpha}_* < 0.5)}{P(\overline{\alpha}_* > 0.5)}, \frac{P(\overline{\alpha}_* > 0.5)}{P(\overline{\alpha}_* < 0.5)}\right\}\right)$$

1127

which ranges in $(1, \infty)$. Each triplet's Flat/Wavy/Central/Extreme tag could then be subcategorized

as either Skewed or Symmetric depending on whether $r_S > 4$ or $r_S < 2$ (with no label in the middle).

This subcategorization step was useful for distinguishing the dynamic admixtures associated with

the whole-trial categorizations of Single and Outside from Intermediate and Mixture, with Single

and Outside tending to be classified as Skewed. Supplementary Table 1 and Supplementary

Figures 6 and 7 give the full results of the main 2-way classification together with the

symmetry/skewness subclassification, cross tabulated with the classification done under the whole

trial spike count analysis.

1136

A vs. B assignment scores

1137

A vs. B assignment scores were computed for several analyses (the example shown in

Figure 1E-F; pairs of recorded neurons; the relationship between spiking activity and local field

potential; and the relationship between saccade sequences and spiking activity). For each triplet,

every dual-sound trial received an A-like score and a B-like score, either for the entire response

window (600-1000 ms after sound onset) or for 50 ms time bins. The scores were computed as the

posterior probability that the spike count in each dual-sound trial was drawn from the Poisson

distribution of single-sound spike counts.

For the pairs analysis, the A vs. B assignment scores were computed within each 50 ms

time bin independently for each pair of neurons recorded simultaneously. The scores were

normalized across trials by subtracting the mean score and dividing by the standard deviation of

scores for that bin (a Z-score in units of standard deviation). Only conditions for which both

recorded neurons exhibited reasonably different responses to the A vs. the B sound and for which

there were at least 5 correct trials for A, B, and AB trials were included (t-test, p < 0.05). A total

of 206 conditions were included in this analysis.

1152

Local field potential analysis

1153

1154 We analyzed the local field potential from 87 sites in both monkeys (30 sites from monkey

1155 P's left IC, 31 sites from monkey Y's right IC and 26 sites from monkey Y's left IC). The LFP

1156 acquisition was either recorded in discrete temporal epochs encompassing behavioral trials

1157 (roughly 1.2 to 2 seconds long) and at a sampling rate of 20 kHz (Dataset I, part of Dataset II), or

as a continuous LFP signal during each session, at a sampling rate of 20 kHz or 1kHz (rest of Dataset II). We standardized the LFP signals by trimming the continuous LFP into single trial intervals and down-sampling all signals to 1 kHz. The MAP system filters LFP signals between 0.7 and 300 Hz; no additional filtering was applied. For each site we subtracted the overall mean LFP value calculated over the entire session, to remove any DC shifts, and we excluded trials that exceeded 500mV.

For the voltage-and-time domain analysis presented in Figure 5A, for each triplet, we assigned individual dual-sound trials to two groups based on the total spike count in a 600 ms response window (see Methods: A vs. B assignment scores). The average LFP was then compared across the two groups in two 600 ms windows before and after sound onset (baseline and response periods). The results reported here refer to these mean-normalized LFP signals. We obtained similar results when the amplitude of each trial's LFP was scaled as a proportion of the maximum response within the session.

Face patch MF recording procedures

The full experimental procedure is described in [37]. We give a summary here. All procedures conformed to the US National Institutes of Health Guide for Care and Use of Laboratory Animals, and were approved by The Rockefeller University Institutional Animal Care and Use Committee (IACUC).

The localization of face patch MF was guided via fMRI as described in [37]. A total of 105 single neurons were recorded from MF in two male adult macaques (monkey Q, Macaca mulatta, and monkey J, Macaca fascicularis). The monkeys were head-restrained and performed a fixation task while viewing visual stimuli on a CRT screen placed 57 cm in front of them. Eye position was monitored with an infrared pupil tracking system (ETL-200, ISCAN Inc.,Burlington, MA). The monkeys were rewarded with juice for maintaining the eyes within a ≤2X2 degree square window around the fixation point.

All stimuli were controlled by custom software written in C (Visiko) running on a Windows PC. For each neuron, three visual stimuli (400 ms, 4X4 degrees in size) were selected from among a pool of face and object stimuli: a face that elicited a strong response, dubbed the preferred face; a face that elicited a weak or no response, dubbed the non-preferred face; and an object that also elicited a poor response, dubbed the non-preferred object. They were presented

either alone or in pairs consisting of the preferred face and one or the other of the non-preferred stimuli. Thus, there were two triplets per cell suitable for analysis. The stimuli were randomly interleaved with each other and with other conditions not analyzed here see [37].

Stimulus positions on the screen were such that the preferred face was always at the center of the neuron's receptive field whereas the non-preferred stimulus could occupy one of eight equidistant locations adjacent to the preferred face. The exact location of the non-preferred face/object was ignored in the present analysis, but excessive heterogeneity in the responses due to variation in location would have caused the triplet to be excluded on the grounds of not exhibiting a sufficiently Poisson-like spike count distribution on the relevant single-stimulus trials.

The data were otherwise analyzed as described in the preceding Analysis section.

Data Availability

The data and computer code that support the findings of this study are available from the corresponding authors upon reasonable request.

ACKNOWLEDGMENTS

We are grateful for expert technical assistance from Jessi Cruger, Karen Waterstradt, Christie Holmes, Stephanie Schlebusch, Tom Heil, and Eddie Ryklin. We have benefitted from thoughtful discussions with Michael Lindon, Liz Romanski, Marlene Cohen, Nicolas Brunel, Aaron Batista, Stephen Lisberger, Marty Woldorff, Daniel Pages, David Bulkin, Kurtis Gruters, Bryce Gessell, Na Young Jun, Cindy King, Luke Farrell and David Murphy. We thank Bao Tran-Phu, Will Hyung, Stephen Spear, Francesca Tomasi, and Ashley Wilson for assistance with animal training and/or recordings. Financial support for the research was provided by the National Science Foundation (0924750) to JMG and the National Institutes of Health (5R01DC013906-02) to STT and JMG.

CONTRIBUTIONS

V.C.C. and J.M.G. coordinated the project; V.C.C and J.L. recorded the IC data; A.F.E. and W.A.F. designed and executed the visual face patch recordings; S.T. and C.G. developed new statistical methods; S.T., C.G., A.Z., V.C.C., J.L., J.T.M, S.M.W., J.M.G. analyzed the data;

1219   V.C.C., J.M.G., S.T.T. wrote the manuscript; V.C.C., J.T.M., S.M.W., J.M.G., S.T., W.A.F.,

1220   R.E. reviewed the manuscript

1221

1222   **Competing interests**

1223   The authors declare no competing interests.

1224

1225   **SUPPLEMENTARY FIGURES AND TABLES**

1226

1227   ***Supplementary Figure 1: Monkeys can localize two sounds.***

1228   *Average accuracy for the dual-sound targets was overall over 70% correct. Performance was*

1229   *better for larger frequency separations. Data combined across all sessions in Data Set I.*

1230

1231

Supplementary Figure 2: Results of conventional analyses pooling across time and trials are

consistent with averaging but not summation.

*(**A**) Schematic activity patterns of IC neurons in response to single-sounds A and B, and predicted*

response for dual sounds if the neuron sums inputs corresponding to individual sounds (A+B) or

*averages them ((A+B)/2). (**B**) Observed Z-scores of activity on dual-sound trials differ from the*

*sum. **(C)** Observed Z-scores of activity on dual-sound trials correspond well to the average. The*

shaded areas indicate Z score values of +/- 1.96 standard deviations (95% confidence intervals)

of the sum (B) and of the average (C). This analysis was conducted on triplets of single- and dual-

sound trials with a given set of locations and frequencies, pooling across intensities. Triplets were

included if the single-sound responses differed (light bars, two-tailed t-test, p<0.05, n=761);

results were similar when single-sound responses were different and both excitatory (dark bars,

one-tailed t-test, p<0.05, n=486).

1244

1245
1246
Supplementary Figure 3. Dual-sound responses do not depend on sound intensities

Same analysis as Supplementary Figure 2, but comparing dual-sound trials to single-sound trials

that used the same signal sent to the speakers (left column), or for which the signal was amplified

on single-sound trials to match the dual-sounds in loudness (middle and right columns). The

results are essentially identical to each other.

Supplementary Figure 4. *Frequency responses of the example cells in the main text. For each*

cell, the average firing rate is computed across single-sound trials with the same central frequency

but pooling across speaker locations (20ms bins with no smoothing). (A) Same cell as Figure 1E,

2E (B) Same cell as Figure 1F, 2G, 3C.

.

1258

1259

1260

1261

Supplementary Figure 5. Whole-trial analysis results broken down by whether neurons were

responsive to both of the sounds in the triplet when presented individually (a) or only one of the

two sounds (b). *Results were largely similar except that Outsides were more common when the*

neuron was responsive to both sounds. Responsiveness was determined by comparing the firing

rate in the 600 ms period after target onset to the 500 ms period prior to sound onset (t-test,

p<0.05).

1268

1269

1270

	Intermediate	Mixture	Outside	Single	Total
Wavy-Central+Symmetric	13 (5)	2			15 (5)
Wavy-Central	1				1
Wavy+Symmetric	4	2			6

Flat-Central+Symmetric	7 (3)				7 (3)
Flat-Central	2 (1)	1			3 (1)
Flat-Extreme+Skewed	8 (1)	12 (3)	19 (4)	46	85 (8)
Flat-Extreme+Symmetric	2	22 (19)		4	28 (19)
Flat-Extreme	3	31 (13)	1	16	51 (13)
Flat+Skewed	4 (2)				4 (2)
Flat+Symmetric	6	11 (4)		4	21 (4)
Flat	13 (2)	1		2	16 (2)
Extreme+Symmetric	1	4 (4)		1	6 (4)
Extreme		3 (2)		1	4 (2)
Central+Symmetric	9 (2)	1		2	12 (2)
Symmetric	36 (1)	42 (5)		14	92 (6)
Skewed	1 (1)				1 (1)
Ambiguous	5	2		3	10
Total	115 (18)	134 (50)	20 (4)	93	362 (72)

1271

1272 ***Supplementary Table 1. Complete listing of tags applied in the DAPP analysis.*** *Categories not*

1273 *shown (e.g. Wavy-Central+Skewed) did not receive any members. The numbers in parentheses are*

1274 *for triplets that produced a winning model in the whole trial analysis with a probability > 95%.*

1275 *The total is 362, since for one Single triplet, the DAPP analysis failed to compute the necessary*

1276 *metrics.*

1277

a Mixtures

b Intermediates

c Singles

d Outsides

1278

Supplementary Figure 6. Hierarchical depiction of the DAPP tags, considering first Flat vs. Wavy, then Central vs. Extreme, then Symmetric vs. Skewed, as a function of whole trial model classification. As in Supplementary Table 1, numbers in parentheses are for triplets that produced a winning model in the whole trial analysis with a probability > 95% and the total included here is 362. For clarity, labels "0(0)" are not shown.

1284

DAPP tags vs. Whole-trial classification for different winning probablities

a   Whole trial model win prob > 0.95

b   Whole trial model win prob > 0.50

c   Whole trial model win prob > 0.33

Percent of triplets

1285

1286

1287   ***Supplementary Figure 7.*** *Relationship between the three relevant DAPP tag combinations (Flat-*
1288   *Extreme, Wavy-Central, and Flat-Central) and whole-trial Poisson classification, for different*
1289   *levels of winning probability for the whole-trial analysis. Panel A is identical to Figure 3E in the*
1290   *main text.*

1291

1292

Supplementary Figure 8. A. Same analysis as in Figure 4B, but shaded according to whether both neurons in the pair preferred the same sound (e.g.. $A_{resp}> B_{resp}$ or vice versa for both neurons in the pair; light gray) vs. different sounds (e.g. $A_{resp}> B_{resp}$ for one neuron and $A_{resp}< B_{resp}$. for the other; dark gray). B. The same analysis as panel A, but using spike counts in each bin instead of A vs. B assignment scores. Asterisks indicate the median; circles indicate the mean.

1298