Epidemic network analysis for mitigation of invasive pathogens in seed systems: Potato in Ecuador

J. F. Hernandez Nopsa[*]**, C. E. Buddenhagen**[*]**, K. F. Andersen, J. Andrade-Piedra, G. A. Forbes, P. Kromann, S. Thomas-Sharma, P. Useche, K. A. Garrett.**

First, second, third, and ninth authors: Plant Pathology Department, University of Florida, Gainesville, FL 32611, United States

First, second, third, and ninth authors: Institute for Sustainable Food Systems, University of Florida, Gainesville, FL 32611, United States

First, second, third, and ninth authors: Emerging Pathogens Institute, University of Florida, Gainesville, FL 32611, United States

Fourth author: International Potato Center, Lima, Peru

Fifth author: International Potato Center, Kunming, China

Sixth author: International Potato Center, Quito, Ecuador

Seventh author: Plant Pathology Department, Kansas State University, Manhattan, KS 66506, United States; current address: Department of Plant Pathology, University of Wisconsin-Madison, Madison, WI 53704

Eighth author: Food and Resource Economics Department, University of Florida, Gainesville, FL 32611, United States

*First and second authors made equivalent contributions

ABSTRACT

Seed system structure defines pathways for the spread of pathogens involved in seed degeneration and influences their ability to supply high quality seed to farmers. We evaluated seed system networks defined by a regional potato farmer consortium (CONPAPA) in Tungurahua, Ecuador. The structure of networks of farmer seed and potato transactions, and the linked network of information about pest and disease management, were estimated based on surveys. We performed a scenario analysis of disease spread in this multilayer network to identify key nodes for sampling and mitigation. The centrality of CONPAPA's leadership group in the network means that disease management interventions, such as training, monitoring and variety dissemination, should target CONPAPA staff and facilities. A market in the largest nearby town, Ambato, was the next most important node. Farmers reported receiving advice about disease and pest management through trusted CONPAPA technical staff. Advice from agrochemical stores was common but viewed as significantly less reliable. Farmer access to information (number and quality of sources) was similar for both genders. Women had a smaller amount of the market share, however. Understanding seed system networks provides a window into options for system improvement that include environmental and societal concerns.

Additional keywords: complex systems, multilayer networks, pests and diseases, seed degeneration, seed networks

Networks of crop seed distribution are an important factor in determining the ecology and

spatial distribution of crop plant genotypes (cultivars or landraces), and disease resistance genes,

as well as determining the potential for the spread of seedborne disease. Seed systems are human

mediated seed distribution networks that encompass biophysical elements as well as all the

stakeholders and activities that support the system, including interacting scientific (e.g.,

breeding, extension), management (e.g., agricultural practices, integrated pest management) and

regulatory aspects (e.g., legally certified seed standards; Almekinders et al. 2007; Devaux et al.

2014; Jaffee et al. 1992; Kromann et al. 2017; Thiele 1999; Thiele et al. 2011). Thus, seed

systems are best understood as a network of interacting biophysical and socioeconomic elements

(Leeuwis and Aarts 2011). Only the state regulated aspects of a system would be considered a

"formal" system (Sperling et al. 2013). Seed systems share many traits with other managed

ecological systems in which there are larger-scale human institutions driving some system

components and individual land managers who make choices about smaller units in the

landscape (e.g., farmers or conservation managers). Institutional interventions –or lack thereof–

can affect systems indirectly and directly via policy, training, funding or direct management. The

relationship between institutional interventions and on the ground management may be more or

less predictable, depending on adoption by land managers.

Ideally seed systems provide disease free, disease resistant, high quality seed to farmers,

through improved seed processing, multiplication, storage and distribution. Scientists also

contribute to seed systems by developing more disease resistant varieties with other positive

traits for dissemination through the system. Understanding seed systems can help scientists make

meaningful links among epidemiological patterns and socioeconomic factors across a range of

scales. While seed transaction networks have sometimes been studied and characterized, there is

great potential for developing new approaches to predict the spread of seedborne diseases and

help target disease detection efforts, training, treatments and other interventions (Andersen et al.

2017; Hernandez Nopsa et al. 2015; Pautasso et al. 2013; Tadesse et al. 2016). Here we use

epidemiological network analysis (Shaw and Pautasso 2014) of a seed potato network to

understand and predict disease risk, developing a new type of scenario analysis for interpreting

epidemic risk in seed systems.

Efforts to create new seed systems may be only partially successful, especially in

developing countries (Devaux et al. 2014; Devaux et al. 2010; Hirpa et al. 2010; Jaffee et al.

1992; Kromann et al. 2017; Panchi et al. 2012; Thiele et al. 2011; Thomas-Sharma et al.

2016). Understanding the structure and function of formal, informal, and mixed seed systems can

support the development of more sustainable seed systems. Aspects that determine the degree of

seed system utility, sustainability, and resilience include access to and availability of seed, seed

quality, cultivar quality (e.g., adapted, disease resistance, and matching user preferences),

affordability, and profitability (Sperling et al. 2013). There are tradeoffs in connectivity for

farmers, where high connectivity is good for getting access to new varieties and training, but can

increase the risk of being exposed to disease. Managing connectivity can help to increase system

resilience (Biggs et al. 2012).

Seed system resilience is tested when there are significant stressors or crises, be they

environmental (Violon et al. 2016), biotic (e.g. pathogen or pest outbreaks) or socioeconomic

(McGuire and Sperling 2013). Though broad categories of threats are predictable, some events

may be viewed as crises because they are spatially varied, temporally unpredictable, and may

have multiple distinct drivers (e.g. pathogen, drought, conflict and economic crises). Formal seed

systems can be "static and bureaucratic" (Lybbert and Sumner 2012) while farmers and markets

may be quite adaptable. Single optimal solutions are unlikely. Many development agencies orient

their interventions toward the development of demand-driven systems that support a for-profit

model of seed supply, believing them more sustainable and resilient (McGuire and Sperling

2013; Sperling et al. 2013), but governments and aid agencies continue to play important roles

in subsidized seed systems. A common belief is that diversity improves resilience, in terms of

crops and cultivars, or supply channels (McGuire and Sperling 2013). A frontier for plant

epidemiology is to better incorporate human decision making about disease management

(McRoberts et al. 2011). Seed system development efforts often attempt to foster equitable

access by stakeholders to services (Ricciardi 2015), although more needs to be understood about

gender effects on access. Epidemiological network analyses can help to identify systemic

vulnerabilities related to gender access to quality planting materials, integrated pest management

information, and the market for products (Tadesse et al. 2016). Clearly short- and long-term

planning could be required to meet challenges, and stakeholders need to be flexible to strike a

good balance between sustaining and transforming systems. Trade-offs are likely, with

interventions under one scenario or set of stressors potentially being counter indicative in another

scenario, or for some stakeholders.

There is always a risk that pathogens can move through a seed system network. Detection of

pathogens in the network in a timely manner could allow for mitigation measures to be

implemented. Important hubs in the network are obvious points of risk for disease spread, but

peripheral nodes could be the entry point for an invasion (Xing et al. 2017). Strategies for

dissemination of resistant varieties may need to change depending on network properties. In

addition, the spread of endemic pathogens like *Rhizoctonia* spp., or the potential arrival of

emerging diseases from distant locations (e.g., *Dickeya* spp.; Czajkowski et al. 2015;

Czajkowski et al. 2011; van der Wolf et al. 2014) can be modeled and mitigation strategies

tested using a multilayer network analysis (Garrett 2012, 2017). Our objectives in the analysis

presented here are as follows: (a) characterize cultivar dispersal through a potato seed system in

Ecuador defined by the potato consortium CONPAPA; (b) determine whether gender is

associated with different types of network transactions or access to information; (c) develop a

risk assessment for disease vulnerability of individual network nodes and examine their utility as

control points for pathogen mitigation measures; and (d) explore a scenario where the

CONPAPA leadership group no longer plays an organizing role in the seed system, and existing

seed multipliers must compensate for its absence.

119   **MATERIALS AND METHODS**

120   **Study system: the CONPAPA potato seed system in Tungurahua, Ecuador.** There

are approximately 50,000 ha of potato production in Ecuador, with 97% of this area located in

the Andes, and 87.5% of farms being less than 10 ha in size (Devaux et al. 2010). It is possible to

produce tubers all year, which has created a market that expects fresh potatoes for consumption

year round (Devaux et al. 2010). Seed is typically reused, i.e., tubers from the previous season

are planted in the next. This makes potato subject to seed degeneration and associated yield loss

(Thomas-Sharma et al. 2016). Seed degeneration refers to the reduction in yield or quality caused

by an accumulation of pathogens and pests in planting material over successive cycles of

vegetative propagation. The national agricultural research institute, INIAP (*Instituto Nacional de*

Investigaciones Agropecuarias) is the only agency in Ecuador registered to produce formal basic

seed potato. However, according to a 2012 estimate, less than 3% of the seed potato used in

Ecuador is from the formal system (Thomas-Sharma et al. 2016). Two preferred cultivars for

farmers in the Ecuadorian Andes are INIAP-*Fripapa* and *Superchola*. However, farmers also

grow many other cultivars, such as INIAP-*Gabriela*, INIAP-*Catalina*, and *Diacol-Capiro*. Seed

is produced by INIAP from pre-basic seed, which are mini-tubers produced from *in-vitro* plants.

Basic seed, the next generation, is multiplied in the field by INIAP or associated farmers. The

next three generations of seed include the following three seed categories; registered seed

(*semilla calidad I*), certified seed (*semilla calidad II*), and selected seed (*semilla calidad III*), and

are produced in the field by seed producers. Trained seed producers form a part of the

Consortium of Potato Producers (CONPAPA) and produce seed for member farmers (Fig. 1).

The yield increase associated with each of these three categories has been reported to be 17%,

11% and 6%, respectively, compared to the seed produced by the farmers in the informal system

(Devaux et al. 2010), although these estimates are low compared to the potential (30%) yield

increases reported globally from the use of quality seed potato (Thomas-Sharma et al. 2016).

Viruses such as *Potato virus Y* (PVY), *Potato virus X* (PVX) and *Potato leafroll virus*

(PLRV), are major causes of seed degeneration in many parts of the world (Frost et al. 2013;

Salazar 1996). Additionally, depending on the geographic region, fungi, bacteria, nematodes,

phytoplasmas, and insects can also play important roles in potato seed degeneration (Thomas-

Sharma et al. 2016). In high-elevation potato production regions of Ecuador, *Rhizoctonia solani*

is a major cause of seed degeneration (Fankhauser 2000), while in many other tropical and

subtropical countries *Ralstonia solanacearum* is a major concern (Mwangi et al. 2008). Adding

to this complex etiology, the rate of degeneration is also highly variable across geographical

regions. Factors such as host physiology, vector dynamics, environmental variability, and the

choice and success of management strategies can affect the rate of degeneration (Thomas-

Sharma et al. 2017; Thomas-Sharma et al. 2016). In high elevation regions, for example, lower

temperatures can limit vector activity and pathogen multiplication while also influencing host

physiology that limits pathogen transmission into daughter tubers (Bertschinger 1992; Navarrete

et al. 2017). In at least one case the presence of *Potato yellow vein virus* (PYVV) was associated

with small yield *improvements*, possibly via some sort of competitive interaction with other

viruses (Navarrete et al. 2017). In the Andes, evidence suggests virus transmission to daughter

tubers is usually incomplete with between 30 and 75% of tubers being infected (Bertschinger et

al. 2017). Similarly, the application of management strategies such as resistant cultivars, certified

seed material and other on-farm management strategies, individually and/or collectively, can

affect the spread of disease epidemics in a region (Thomas-Sharma et al. 2017). A better

understanding of these inter-related factors could contribute to the design of an integrated seed

health strategy for a geographic region (Thomas-Sharma et al. 2016).

Established in 2006, CONPAPA has a membership of ca. 300 farmers in central Ecuador

(principally in Tungurahua, Chimborazo and Bolivar Provinces). This organization is the current

realization of various aid and governmental efforts to improve livelihoods for small-scale potato

farmers (Kromann et al. 2017). It aims to support small-scale farmer associations that produce

seed potato and potato for consumption, through training, provision of quality assessed seed, and

by processing and marketing produce. It cleans and processes produce (e.g., for chips and fresh

potato) in regional processing facilities. It also sells potato on behalf of members. Annual mean,

production yield of table potato in CONPAPA (Tungurahua) ranges between 15 and 20 metric

tons per hectare, with production levels being influenced by management, variety, time of year,

and the number of generations since the seed was sourced from basic seed. Average production

reported by CONPAPA is higher than the 9.5 metric tons per hectare that has been reported for

Ecuador as a whole (Devaux et al. 2010). CONPAPA in Tungurahua reported

(www.conpapa.org) that it supplies more than 25 tons of potato for consumption per week to

meet market demand. Importantly, CONPAPA has trained seed multipliers who provide seed for

redistribution to member farmers.

Survey methods. This study focuses on 48 farmers who are members of CONPAPA in

the Tungurahua province. This is 66% of the 72 heads of households registered as members in

this region (Montesdeoca, pers. comm.). However, the 48 farmers in this study represent a

census of all the active farmers at the time of this study. Farmer network sizes and farmer

activity can change as farmers opportunistically pursue a variety of alternative livelihoods from

year to year, e.g., construction or service jobs, in response to changing conditions (in good and

bad years; Violon et al. 2016). A questionnaire was completed by scientists via on-farm

voluntary interviews of 48 farmers in the CONPAPA district of Tungurahua over three weeks in

November and December, 2015. In addition to demographic information, the questionnaire

documented the seed sources, cultivars planted, volume bought, and price paid for the last three

planting periods, as reported by farmers. Information was also collected about the sale or use of

potato for food, including destination, cultivar, volume, and price received. Information was

recorded about the principal pests and diseases that the farmers reported. Farmers were also

asked to describe their sources of advice regarding integrated pest management, and the

confidence they had in that advice. In some cases, there was missing data related to volume or

price information.

Data analysis and modeling. Networks of seed and potato transactions between the

farmers and other stakeholders were analyzed using igraph (Csárdi and Nepusz 2006) in the R

programming environment (R Core Team 2016). For cases where farmers reported a transaction

but did not give volume information, links were depicted in the network as dotted lines and given

a minimum visible width. Missing price and volume data were not treated as zeroes, but were

omitted from the calculation of means and percentages. Missing volume and price data are

reported in the results. An adjacency matrix based on reported sales was constructed, as well as

an adjacency matrix based on reported information flow. Transaction counts, volumes and prices

were compared with respect to potato cultivar and farmer gender, based on percentages, means,

and one-sided Wilcox tests (using the wilcox.text function in R). The frequency with which

common pests and diseases were reported by farmers, including diseases responsible for seed

degeneration, is reported overall and by gender (where gender differences were tested using chi-

squared tests).

210   **Rating the importance of nodes for sampling efforts.** An important question for

optimizing management of potential invasive pathogens in a seed system, is where the most

important geographic nodes are for sampling to detect disease (both in the field and in the

harvested tubers). Sampling some nodes will result in rapid detection of the pathogen, while

sampling other nodes will only detect the pathogen when it has already spread widely in the

network. In a scenario analysis, disease spread was simulated across the seed and table potato

distribution network, where the network was based on reports aggregated across the last three

plantings (and actual or anticipated harvest dates ranged from May 2014 to May 2016). In the

simplest version of the analysis, each node was considered equally likely to be the point of initial

introduction of a pathogen into the seed system network. Another version of the analysis drew

on the structure of the communication network. In this case, the probability that a pathogen

would be introduced into the network by a given farmer was weighted by a farmer's access to

information about pest and disease management (IPM), as a proxy for farmer ability to respond

effectively. During the survey interviews, farmers described their sources of information for pest

and disease management, and the trust that they placed in these. The probability that infection

would be introduced into the network by a given farmer was weighted by a function of the

number and quality of information sources about IPM. The idea is that a well-informed farmer

(with high node in-degree in communication networks, or with highly trusted sources) will be

less likely to be a point of disease establishment (with the probability of disease entering the

network at a node set to 0.8 to the power of the number of sources for that node). "In-degree" is

the number of directed links that point toward a node. In this case, it indicates the number of

sources of information that a farmer reports. This simulation generates an estimate of the number

of nodes infected before the disease will be detected at each potential sampling node, given that

each potential starting node has a weighted probability of being the initial source. The output

allows us to estimate relative risk in terms of the number of nodes that would be infected if only

the node in question were monitored.

Scenario analysis where the CONPAPA leadership group does not supply seed. The

CONPAPA leadership group is clearly central to this seed system, a key "cutpoint", or node

whose removal creates multiple disconnected components in the network. We explore how

resilient the seed system might be if the CONPAPA leadership group were removed. How

would other nodes need to compensate for its absence? We compared the scenario where the

CONPAPA leadership group provides seed to farmers and multipliers with a scenario where it

does not have a role in seed provision. For this alternative scenario we evaluated the reported

volumes for seed transactions over three plantings. Then where the CONPAPA leadership group

provided basic seed to multipliers we replaced these transactions with INIAP, the government

agency that provides basic seed to CONPAPA (GovtAgency1 in the Figures). Finally, where

farmers sourced their seed from the CONPAPA leadership group, they instead sourced their seed

from the geographically nearest multiplier (Farmers 7, 27, 34 and 46). The alternative scenario

thus maintains the same transaction volumes that were reported but removes the CONPAPA

leadership group as the go-between replacing these with the most plausible alternative. We

evaluate the structure of this new network.

RESULTS

Seed system: overview. The seed system in this study is centered around the CONPAPA

leadership group in Tungurahua, which provides and receives seed and table potato from

member farmers (Fig. 2). A total of 1157 quintals (100 lb bags), or 52 t (metric tons), of seed

was reported as used by farmers in the most recent planting, where CONPAPA provided 47%,

and 36% was self-supplied or reused seed, while the remaining 16% came from other sources.

CONPAPA was reported as receiving only 7 t of seed from trained (male) seed multipliers. Only

two women (F7 and F46) reported providing seed (Puca, Fripapa and Superchola) to CONPAPA

during this interval, although farmers 7, 8, 10, 19, 36, 40, 46, and 47 are women trained to be

seed multipliers. Of the 48 farmers that reported buying or selling potato or seed, 16 (33%) were

women. Farmers reported a total of 503.9 t potato being sold, with CONPAPA buying 414.7 t

(82%) of potato (where 28% of this was from women). Farmers reported selling 85.3 t directly to

local markets, and one farmer reported selling 3.2 t directly to a restaurant. It is important to note

that 262 transactions were reported in the most recent season but interviewees did not provide

volume for 71 transactions or price information for 58 transactions (including self-supplied seed

transactions). On a per transaction basis there was a difference between the volume of potato

product sold by women (mean=97 quintals) and men (mean=165 quintals; Wilcox test (one sided

alternative=less), W=1159 p-value = 0.03). There was also evidence for a difference in per

transaction volume for seed between women and men, with means of 5.6 and 16.2 quintals

respectively (Wilcox test, one sided alternative=less, W = 127.5, p-value = 0.003). There was not

evidence for a difference in per transaction prices for table potato for women and men, with

means of $13.5 and $12.5 USD, respectively (Wilcox test (one sided alternative=greater),

W=1456 p-value = 0.16). Prices were infrequently reported. These analyses are based on the

most recent season. Unreported here is the movement of pre-basic seed to CONPAPA from

INIAP. CONPAPA in Tungurahua may also receive seed from CONPAPA multipliers outside of

the region. Farmers reported replacing seed every 3-4 seasons. The evidence here is that

improved or healthy seed is bought but grown alongside seed saved from previous plantings.

Seed system: analysis by variety. Overall, while farmers planted on average two

cultivars, the median use was just one. In other words, about half of the farmers planted a single

cultivar, while the other half planted 2 to 5 different cultivars. Ranking the use of cultivars by the

numbers of farmers using them matches almost exactly the ranking by number of transactions

per cultivar (Table 1), which suggests that the high number of transactions for the main cultivars

is driven by their overall popularity. The 3 most commonly planted cultivars, according to these

criteria, are Superchola (33% of farmers planted it, its product transactions represent 36% of all

transactions, and its seed transactions 32%), Fripapa (17%, 20%, 22%) and Puca (13%, 10%,

10%), in respective order of ranking.

A second comparison of the total volume of transactions by cultivar, shows that the three

most frequently exchanged cultivars are also the ones with most transacted volume (Table 2).

Indeed, Superchola's transacted volume represents 40% of all volume transacted in terms of

product and 35% in terms of seed. Fripapa's seed volume transacted is higher than the product

volume transacted 26% vs. 21%. Finally, Puca variety volume represents 9% in terms of product

and 7% in seed. Interestingly, two varieties that are not used by a majority of farmers —Carrizo

and Victoria—represent 8 and 7% in terms of volume transacted, almost as much as Puca. This

related to a few farmers providing large volumes of product to a few non-CONPAPA buyers.

Finally, the percentage of volume transacted of Unica's seed is larger than Puca's seed volume

(9%) and Natividad is as large as Puca's (7%).

IPM information. Farmers largely reported obtaining information about integrated pest

and disease management (IPM) from the CONPAPA leadership group (mean in-degree for

information received by farmers was 3.5 overall; Fig. 3). There was not evidence for a difference

(t-test, p-value=0.39) between male (3.7) and female (3.2) in-degree with respect to number of

information sources reported. Importantly, farmers frequently reported receiving information

from agrichemical stores (green squares in Fig. 3). Family members also provided important

sources of information about IPM (Fig. 3). A quarter of the women reported their husband as a

source of information for IPM, but no men reported that their wife was a source of IPM

information. Farmer assessed trust levels could range between zero and five. There was some

evidence for a difference in mean trust levels reported by men (3.4) and women (3.8) (t-test, p-

value=0.08). The main sources of information were CONPAPA and agrochemical stores, where

the mean trust level farmers reported for all stores was 3.01 compared to 4.4 for CONPAPA (t-

test, p-value=1.873e-07). Only one farmer reported the internet as an important source of

information about management.

The most frequently reported diseases and pests were potato late blight, Andean potato

weevil, and potato black leg. Despite prompting, viruses were reported by only one percent of

farmers (Table 3). Slugs and leaf miners were more frequently reported as a problem by women

than men, though rates were low (Table 3).

Disease risk in the system. Under the scenarios we evaluated (Fig. 4A-C), the

CONPAPA leadership group is obviously the most effective place to monitor in order to detect a

disease before it has spread far. This reflects its central role in the network. Similarly, several

stakeholders and farmers at the periphery of the seed and potato network tend to be poor

locations for detecting potential disease in every simulation. This is because they only provided

seed rather than receiving seed or product (yellow) in this network, or had low in-degree (orange

or light orange; Fig. 4A-C). Weighting risk of establishment based on the quality of the

information sources about IPM causes some nodes at the periphery to become more important

for monitoring (colors are yellow in the equal weight scenario (Fig. 4A) versus darker orange

(Fig. 4B-C). In the scenario where farmer ability to prevent establishment was weighted by the

number of sources of information (Fig. 4C), we find that sampling "Market1" will lead to

relatively rapid detection of an incipient disease. This is the market in Ambato, the largest town

in the region, which has the highest reported in-degree of any of the five markets.

Scenario analysis where the CONPAPA leadership group does not supply seed. We

compared the scenario where the CONPAPA leadership group provides seed to farmers and

multipliers, and multipliers sell their seed to CONPAPA (Fig. 5A), with a scenario where the

CONPAPA leadership group does not have a role in seed provision (Fig. 5B). In this analysis,

based on a role of geographic proximity to multipliers, we see that multipliers do not have equal

access to all the seed buying farmers in the market (Fig. 5B). CONPAPA's role as distributor

and organizer of seed distribution (Fig. 5A) may result in all farmers having access to seed from

any of the multipliers.

DISCUSSION

In this analysis, we demonstrate an approach for identifying priorities for monitoring plant health

in seed systems. In this relatively small and centralized seed system, disease monitoring at

CONPAPA processing facilities is obviously a high priority for detection of incipient disease,

because it receives high quantities of table potato (it has high in-degree), and is the source of

most of the improved seed (it has high out-degree). Monitoring at the market in Ambato could

also be relatively effective. Similarly, mitigation measures during a disease outbreak – such as

dissemination of new resistant cultivars, training, or treatment of fields – would best focus on

these nodes in the network. Secondarily, the analysis identifies other nodes in the network that

can play a role in sampling and mitigation, offering a method to prioritize among these nodes for

sampling in the field and postharvest. Network models provide a window into the epidemiology

of plant diseases and strategies for efficient sampling for plant epidemic surveillance (Chadès et

al. 2011; Harwood et al. 2009; Hernandez Nopsa et al. 2015; Sanatkar et al. 2015; Sutrave et

al. 2012). We find that the CONPAPA leadership group and the Ambato market (Market1 in the

Figs. 2, 4 and 5) would be effective points for monitoring. By this we mean that if an invasive

disease entered the network from any node, it would tend to spread less through the network

before it was detected if sampling was at these key nodes (Fig. 4). Secondary nodes identified as

having some sampling value could also be ranked and prioritized to supplement sampling of the

two key nodes. An undetected disease at the CONPAPA leadership group or the Ambato market

would spread relatively quickly through the network.

Information about the dispersal of particular cultivars through the seed network can

provide insights into the likelihood of disease transmission, if cultivars have resistance to a

particular disease or if seed of a new cultivar is inadvertently a source of an introduced pathogen.

Good information is available about cultivar susceptibility to *Phytophthora infestans* (e.g.,

Forbes 2012; Kromann et al. 2009), but studies of viral infection rates for cultivars used in

Ecuador rarely consider more than a few varieties. Seed born viral incidence, especially PYVV,

PVS, and PVX were reported in one study for some of the cultivars used by CONPAPA farmers,

(from lowest to highest incidence: Fripapa, Gabriela, Yana, Unica, Dolores and Chaucha), but

per plant yield effects were negligible (Navarrete et al. 2017). High levels of PVY infection have

been reported occasionally in Ecuador for Superchola and Fripapa, but viral incidence seems to

depend on complex interactions between ecological conditions, on-farm management practices,

vector biology, seed sources and cultivar (Navarrete et al. 2017). Yana was reported as extremely

resistant to PLRV and PVY, while Unica was resistant to PVY but susceptible to PLRV

(Acquisition and Distribution Unit 2009).

In Ecuador seed degeneration, mostly attributable to viruses, can have important effects

on yield (7-17% loss, or even *gains* in the case of PYVV), but virus incidence is often low at

high altitude, even if levels vary widely from site to site (Peter Kromann unpublished; Devaux et

al. 2010; Navarrete et al. 2017; Panchi et al. 2012). It appears that the problem is still

understudied, under-appreciated or rarely recognized. For example, only one farmer reported

viruses as concern in this study. Yield losses of ±30% are common elsewhere in the world

(Thomas-Sharma et al. 2017). A large share of farmers report that they draw on advice from

agrochemical stores. Importantly, and perhaps with good reason, farmers do not report trusting

them highly as a source of information compared to technical staff working for CONPAPA.

Clearly training these store owners about disease and pest management has the potential to be an

effective measure to improve management outcomes for farmers inside and outside of the

consortium. However, it is unclear if training store owners would result in improved advice and

the sale of appropriate pesticides, or if potential economic conflicts of interest would influence

the quality of their advice. We modelled disease spread as a function of farmer information

quality and sources with respect to IPM. This usefully joins the network for the spread of

information about management with the biophysical network (seed network and disease

epidemiological models).

Women made up a third of the farmers and reported selling smaller volumes of potato

product on average. Clearly, they are making less money from potato farming than their male

counterparts. There were limited differences in gender access in terms of the number of

information sources, or the trust they placed in their information sources. It would be interesting

to determine if this is typical, or if less formal seed system networks in the region reveal larger

gender effects.

Modeling disease spread in seed and potato transaction networks can indicate the

structural effects of seed degeneration. In the case of viruses, most are transmitted to daughter

tubers and will be hitchhikers for each transaction of seed or potato. It is clear that some spread

can always occur via the seed system. Network dynamics change from year to year, so scenarios

should consider temporal dynamics (e.g., the different effects of wet and dry years; Violon et al.

2016). A more nuanced approach would also take into account different suites of viruses, and the

way their transmission rates from infected mother plants to daughter tubers vary depending on

varietal and environmental conditions (Bertschinger et al. 2017). Thus node (farmer)

vulnerability to infection could also be modeled in terms of specific diseases and scenarios, and

could account for varietal differences in resistance.

A key point to consider for potato seed systems is transmission mechanisms. As a case in

point, *Potato virus X* (PVX) and *Andean potato mottle comovirus* (APMoV) are transmitted by

contact while others such as *Potato virus Y* (PVY) and *Potato leafroll virus* (PLRV), *Potato*

yellow vein virus (PYVV) are vectored by aphids (Fankhauser 2000). Networks could include

both spread through seed transactions, and spread based on the spatial proximity of farm pairs (as

a proxy for the probability of vector movement between a pair). In this study, farms were widely

dispersed with both potato and other crops being grown in the intervening areas. Inoculum

sources could come from non-CONPAPA potato farmers, or non-potato host species. To

realistically model seed infection by vectors would require detailed disease specific data sets that

support accurate estimation of dispersal kernels, including the effects of infected volunteers and

tuber waste from potato harvesting.

Implementation of fully formal seed systems in many developing countries is beyond the

available resources of the agencies and farmers involved. Reaching the quality levels enshrined

in statutes may not be feasible. This means that most potato farmers in developing countries

operate wholly within informal seed systems. The CONPAPA seed system has been described as

a mixed formal and informal system (Kromann et al. 2017). CONPAPA defines seed quality

explicitly in three levels with real quality control measures in place. This means farmers can buy

improved seed of known quality with achievable quality levels for the stakeholders involved.

The adoption of this alternative seed quality assessment scheme has been incorporated into

formal Ecuadorian seed regulation (Kromann et al. 2017), thus formalizing the standards

CONPAPA developed. This has been described as "providing flexibility" (FAO 2006) and is

recommended as a means of achieving greater confidence by stakeholder and adoption of

improved seed. Therefore, the CONPAPA seed system could be characterized as predominantly

formal with the quality declared seed sources accounting for 47% of the seed in this study. In

practice, the mean time between seed replenishment was reported to be approximately 3-4

seasons, though we also found that improved seed are often planted together with reused seed in

any given year. This is a much higher rate of improved seed use than the 2-3% formal seed

sources reported for Ecuador and Bolivia (Almekinders et al. 2007; Devaux et al. 2010).

CONPAPA's cooperative model, combined with the seed quality assessment system could help

to overcome issues of access and household economic insecurity that determined participation in

formal seed systems elsewhere (Okello et al. 2016). This could have important consequences

since potato is becoming increasingly important as a staple crop in areas where informal seed

systems prevail (Devaux et al. 2014).

We evaluated the CONPAPA structure as a first step to support improved sampling, IPM,

risk assessment for pathogen and pest movement, and farmer decision-making. Identification of

key control points that influence the success of seed systems (e.g., farmers, farms, information

sources) supports enhancement of the system (e.g., maximizing the distribution of new seed

varieties using fewer distribution channels, managing disease outbreaks, and targeting

improvement of communication and infrastructure). Resources can be invested in particular

nodes to improve practices to control pest and disease outbreaks, leading to improvements in the

seed system. We present results for the CONPAPA system as part of an ongoing project to

develop general recommendations for improving seed system structure. While we illustrate here

how a seed system could potentially be resilient to removal of a key node (Fig. 5), the temporal

and structural dynamics of seed systems such as CONPAPA need to be better understood to

anticipate how they will react to important stressors, and to develop strategies for reducing

disease risk while increasing availability of improved varieties.

Seed system and network analyses provide one window into global change phenomena

encompassing environmental and societal concerns. The adoption of formal seed systems is

inherently a risk avoidance measure that aims to increase productivity and improve economic

outcomes for farmers, but the implications are wide reaching. Global change in land use, land

cover and biodiversity is often mediated through agricultural practices, development and trade.

Links are easily made between seed systems and land use change, agrochemical use,

biodiversity, climate change, and invasive species more broadly, in addition to disease impacts.

Local seed systems such as the one in this study are linked internationally via plant breeding

networks, through which resistance genes may be distributed, with the associated need to

manage connectivity for movement of pathogens (Garrett et al. 2017). Network analysis and a

systems approach can be used to expand our understanding of interacting biophysical,

socioeconomic and informational elements, and to put management interventions into their

proper context at local and regional scales.

462   **ACKNOWLEDGEMENTS**

This work received institutional review board approval through the University of Florida

IRB-02 #IRB201700024. We confirm farmer participation was voluntary and personal and

demographic information was protected. This research was undertaken as part of, and funded by,

the CGIAR Research Program on Roots, Tubers and Bananas (RTB) and supported by CGIAR

Fund Donors http://www.cgiar.org/about-us/governing-2010-june-2016/cgiar-fund/fund-donors-

2/ and the University of Florida. JHN designed the survey, collected data, wrote R scripts,

performed preliminary analyses, and wrote the paper; CEB wrote R scripts, performed analyses,

produced the figures, and wrote the paper; KFA contributed to R scripts and analyses; PK, GAF,

and JAP designed the study, and helped collect data; STS contributed to survey design and wrote

the paper; PU contributed to survey design and provided an economic analysis; and KAG

designed the study, wrote R scripts, mentored authors, and wrote the paper. We appreciate the

help of J. L. Brisbane and S. L. Lei in data management, and the help of I. Navarrete in testing

preliminary survey drafts. The authors confirm they have no conflicts of interest.

LITERATURE CITED

Acquisition and Distribution Unit. 2009. Catolog of potato varieties. Internation Potato Center (CIP), Lima, Peru.

Almekinders, C., Thiele, G., and Danial, D. L. 2007. Can cultivars from participatory plant breeding improve seed provision to small-scale farmers? Euphytica 153:363-372.

Andersen, K., Buddenhagen, C., Rachkara, P., Gibson, R., Kalule, S., Phillips, D., and Garrett, K. 2017. Analyzing key nodes and epidemic risk in seed networks: sweetpotato in northern Uganda. bioRxiv.

Bertschinger, L. 1992. Modelling of potato virus pathosystems by means of quantitative epidemiology: An exemplary case based on virus. Swiss Federal Institute Of Technology Zurich.

Bertschinger, L., Bühler, L., Dupuis, B., Duffy, B., Gessler, C., Forbes, G. A., Keller, E. R., Scheidegger, U. C., and Struik, P. C. 2017. Incomplete infection of secondarily infected potato plants – an environment dependent underestimated mechanism in plant virology. Frontiers in Plant Science 8:1-13.

Biggs, R., Schlüter, M., Biggs, D., Bohensky, E. L., BurnSilver, S., Cundill, G., Dakos, V., Daw, T. M., Evans, L. S., Kotschy, K., Leitch, A. M., Meek, C., Quinlan, A., Raudsepp-Hearne, C., Robards, M. D., Schoon, M. L., Schultz, L., and West, P. C. 2012. Toward principles for enhancing the resilience of ecosystem services. Annual Review of Environment and Resources 37:421-448.

Chadès, I., Martin, T. G., Nicol, S., Burgman, M. A., Possingham, H. P., and Buckley, Y. M. 2011. General rules for managing and surveying networks of pests, diseases, and endangered species. Proceedings of the National Academy of Sciences of the United States of America 108:8323-8328.

Csárdi, G., and Nepusz, T. 2006. The igraph software package for complex network research. InterJournal, Complex Systems 1695:http://igraph.org.

Czajkowski, R., Pérombelon, M. C. M., van Veen, J. A., and van der Wolf, J. M. 2011. Control of blackleg and tuber soft rot of potato caused by *Pectobacterium* and *Dickeya* species: a review. Plant Pathology 60:999-1013.

Czajkowski, R., Perombelon, M., Jafra, S., Lojkowska, E., Potrykus, M., van der Wolf, J., and Sledz, W. 2015. Detection, identification and differentiation of *Pectobacterium* and *Dickeya* species causing potato blackleg and tuber soft rot: a review. Ann Appl Biol 166:18-38.

Devaux, A., Kromann, P., and Ortiz, O. 2014. Potatoes for sustainable global food security. Potato Research 57:185-199.

Devaux, A., Ordinola, M., Hibon, A., and Flores, R. 2010. El Sector Papa en la Región Andina: Diagnóstico y Elementos para Una Visión Estratégica (Bolivia, Ecuador y Perú). Centro Internacional de la Papa, Lima, Peru.

Fankhauser, C. 2000. Seed-transmitted diseases as constraints for potato production in the tropical highlands of Ecuador. Swiss Federal Institute of Technology, Zürich.

FAO. 2006. Quality declared seed system. Page 243. FAO, Rome, Italy.

Forbes, G. A. 2012. Using host resistance to manage potato late blight with particular reference to developing countries. Potato Research 55:205-216.

520   Frost, K. E., Groves, R. L., and Charkowski, A. O. 2013. Integrated control of potato pathogens
521   through seed potato certification and provision of clean seed potatoes. Plant Disease
522   97:1268-1280.
523   Garrett, K., Andersen, K., Bowden, R., Forbes, G., Kulakow, P., and Zhou, B. 2017. Resistance
524   genes in global crop breeding networks. bioRxiv.
525   Garrett, K. A. 2012. Information networks for plant disease: Commonalities in human
526   management networks and within-plant signaling networks. Eur. J. Plant Pathol. 133:75-
527   88.
528   Garrett, K. A. 2017. Impact Network Analysis: a framework for evaluating the effects of
529   information and other technologies through linked socioeconomic and biophysical
530   networks. bioRxiv.
531   Harwood, T. D., Xu, X., Pautasso, M., Jeger, M. J., and Shaw, M. W. 2009. Epidemiological risk
532   assessment using linked network and grid based modelling: *Phytophthora ramorum* and
533   *Phytophthora kernoviae* in the UK. Ecol. Model. 220:3353-3361.
534   Hernandez Nopsa, J. F., Daglish, G. J., Hagstrum, D. W., Leslie, J. F., Phillips, T. W., Scoglio,
535   C., Thomas-Sharma, S., Walter, G. H., and Garrett, K. A. 2015. Ecological networks in
536   stored grain: Key postharvest nodes for emerging pests, pathogens, and mycotoxins.
537   BioScience 65:985-1002.
538   Hirpa, A., Meuwissen, M. P. M., Tesfaye, A., Lommen, W. J. M., Oude Lansink, A., Tsegaye,
539   A., and Struik, P. C. 2010. Analysis of seed potato systems in Ethiopia. American Journal
540   of Potato Research 87:537-552.
541   Jaffee, S., Srivastava, J., and Mundial, B. 1992. Seed system development: the appropriate roles
542   of the private and public sectors. World Bank Washington, DC.
543   Kromann, P., Montesdeoca, F., and Andrade-Piedra, J. 2017. Integrating formal and informal
544   potato seed systems in Ecuador. in: Case Studies of Roots, Tubers and Bananas Seed
545   Systems. J. L. Andrade-Piedra, J. W. Bentley, C. Almekinders, K. Jacobsen, S. Walsh
546   and G. Thiele, eds. CGIAR Research Program on Roots, Tubers and Bananas (RTB),
547   Lima, Peru.
548   Kromann, P., Taipe, A., Perez, W. G., and Forbes, G. A. 2009. Rainfall thresholds as support for
549   timing fungicide applications in the control of potato late blight in Ecuador and Peru.
550   Plant Disease 93:142-148.
551   Leeuwis, C., and Aarts, N. 2011. Rethinking communication in innovation processes: creating
552   space for change in complex systems. The Journal of Agricultural Education and
553   Extension 17:21-36.
554   Lybbert, T. J., and Sumner, D. A. 2012. Agricultural technologies for climate change in
555   developing countries: Policy options for innovation and technology diffusion. Food
556   Policy 37:114-123.
557   McGuire, S., and Sperling, L. 2013. Making seed systems more resilient to stress. Global
558   Environmental Change 23:644-653.
559   McRoberts, N., Hall, C., Madden, L. V., and Hughes, G. 2011. Perceptions of disease risk: From
560   social construction of subjective judgments to rational decision making. Phytopathology
561   101:654-665.
562   Mwangi, J. K., Nyende, A. B., Demo, P., and Matiru, V. N. 2008. Detection of latent infection
563   by *Ralstonia solanacearum* in potato (*Solanum tuberosum*) using stems instead of tubers.
564   African Journal of Biotechnology 7:1644-1649.

Navarrete, I., Panchi, N., Kromann, P., Forbes, G., and Andrade-Piedra, J. 2017. Health quality
of seed potato and yield losses in Ecuador. bioRxiv:108712.
Okello, J., Zhou, Y., Kwikiriza, N., Ogutu, S., Barker, I., Schulte-Geldermann, E., Atieno, E.,
and Ahmed, J. 2016. Determinants of the use of certified seed potato among smallholder
farmers: the case of potato growers in central and eastern Kenya. Agriculture 6:55.
Panchi, N., Navarrete, I., Taipe, A., Orellana, H., Pallo, E., Yumisaca, F., Montesdeoca, F.,
Kromann, P., and Andrade-Piedra, J. L. 2012. Incidencia, severidad y pérdidas causadas
por plagas de la semilla se papa en Ecuador. Pages 17-20 in: Congreso de la Asociación
Latinoamericana de la Papa—ALAP., Uberlandia, Brazil.
Pautasso, M., Aistara, G., Barnaud, A., Caillon, S., Clouvel, P., Coomes, O. T., Deletre, M.,
Demeulenaere, E., De Santis, P., Doring, T., Eloy, L., Emperaire, L., Garine, E.,
Goldringer, I., Jarvis, D., Joly, H. I., Leclerc, C., Louafi, S., Martin, P., Massol, F.,
McGuire, S., McKey, D., Padoch, C., Soler, C., Thomas, M., and Tramontini, S. 2013.
Seed exchange networks for agrobiodiversity conservation. A review. Agron. Sustain.
Dev. 33:151-175.
R Core Team. 2016. R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria.
Ricciardi, V. 2015. Social seed networks: Identifying central farmers for equitable seed access.
Agricultural Systems 139:110-121.
Salazar, L. F. 1996. Potato Viruses and their Control. International Potato Center, Lima, Peru.
Sanatkar, M. R., Scoglio, C., Natarajan, B., Isard, S., and Garrett, K. A. 2015. History, epidemic
evolution, and model burn-in for a network of annual invasion: Soybean rust.
Phytopathology 105:947-955.
Shaw, M. W., and Pautasso, M. 2014. Networks and plant disease management: concepts and
applications. Annual Review of Phytopathology 52:477-493.
Sperling, L., Ortiz, O., and Thiele, G. 2013. Seed systems. Conceptual frameworks for guiding
practical interventions. CGIAR.
Sutrave, S., Scoglio, C., Isard, S. A., Hutchinson, J. M. S., and Garrett, K. A. 2012. Identifying
highly connected counties compensates for resource limitations when evaluating national
spread of an invasive pathogen. PLoS ONE 7:e37793.
Tadesse, Y., Almekinders, C. J. M., Schulte, R. P. O., and Struik, P. C. 2016. Tracing the seed:
seed diffusion of improved potato varieties through farmers' networks in Chencha,
Ethiopia. Experimental Agriculture:1-16.
Thiele, G. 1999. Informal potato seed systems in the Andes: Why are they important and what
should we do with them? World Development 27:83-99.
Thiele, G., Devaux, A., Reinoso, I., Pico, H., Montesdeoca, F., Pumisacho, M., Andrade-Piedra,
J., Velasco, C., Flores, P., Esprella, R., Thomann, A., Manrique, K., and Horton, D. 2011.
Multi-stakeholder platforms for linking small farmers to value chains: evidence from the
Andes. International Journal of Agricultural Sustainability 9:423-433.
Thomas-Sharma, S., Andrade-Piedra, J., Carvajal Yepes, M., Hernandez Nopsa, J., Jeger, M.,
Jones, R., Kromann, P., Legg, J., Yuen, J., Forbes, G., and Garrett, K. A. 2017. A risk
assessment framework for seed degeneration: Informing an integrated seed health
strategy for vegetatively-propagated crops. bioRxiv: https://doi.org/10.1101/105361.
Thomas-Sharma, S., Abdurahman, A., Ali, S., Andrade-Piedra, J., Bao, S., Charkowski, A.,
Crook, D., Kadian, M., Kromann, P., Struik, P., Torrance, L., Garrett, K. A., and Forbes,

G. A. 2016. Seed degeneration in potato: the need for an integrated seed health strategy to
mitigate the problem in developing countries. Plant Pathology 65:3-16.
van der Wolf, J. M., Nijhuis, E. H., Kowalewska, M. J., Saddler, G. S., Parkinson, N.,
Elphinstone, J. G., Pritchard, L., Toth, I. K., Lojkowska, E., Potrykus, M., Waleron, M.,
de Vos, P., Cleenwerck, I., Pirhonen, M., Garlant, L., Helias, V., Pothier, J. F., Pfluger,
V., Duffy, B., Tsror, L., and Manulis, S. 2014. *Dickeya solani* sp. nov., a pectinolytic
plant-pathogenic bacterium isolated from potato (*Solanum tuberosum*). Int J Syst Evol
Microbiol 64:768-774.
Violon, C., Thomas, M., and Garine, E. 2016. Good year, bad year: changing strategies,
changing networks? A two-year study on seed acquisition in northern Cameroon.
Ecology and Society 21.
Xing, Y., Hernandez Nopsa, J., Andrade-Piedra, J., Beed, F., Blomme, G., Carvajal Yepes, M.,
Coyne, D., Forbes, G., Kreuze, J., Kroschel, J., Kumar, L., Legg, J., Parker, M., Schulte-
Geldermann, E., and Garrett, K. A. 2017. Global cropland connectivity: A risk factor for
invasion and saturation by emerging pathogens and pests.
bioRxiv:https://doi.org/10.1101/106542.
626

Tables

TABLE 1. The number of transactions and number of farmers using each cultivar for the current season, as reported in November 2015 (with percentages).

Cultivar	Total transactions	Seed potato		Table potato		No. Farmers using the cultivar	
Superchola	90	40	32%	50	36%	31	33%
Fripapa	56	28	22%	28	20%	16	17%
Puca-shungo	27	13	10%	14	10%	12	13%
Yana-shungo	17	9	7%	8	6%	8	8%
Unica	16	7	6%	9	7%	6	6%
Carolina	13	6	5%	7	5%	4	4%
Victoria	10	4	3%	6	4%	4	4%
Gabriela	8	3	2%	5	4%	3	3%
Chaucha	7	4	3%	3	2%	3	3%
Carrizo	6	3	2%	3	2%	2	2%
Suprema	4	2	2%	2	1%	2	2%
Americana	2	1	1%	1	1%	1	1%
Natividad	2	2	2%	0	0%	1	1%
Norteña	2	2	2%	0	0%	1	1%
Tulca	2	1	1%	1	1%	1	1%

634 TABLE 2. Volume of seed (quintals) and product exchanged (with percentages).

Variety	Total volume				Volume per transaction	
	product		Seed		Product	Seed
Superchola	4580	40%	425	35%	143	11
Fripapa	2405	21%	323	26%	172	15
Puca-shungo	999	9%	80	7%	111	7
Carrizo	960	8%	66	5%	480	22
Victoria	760	7%	48	4%	190	12
Unica	600	5%	116	9%	200	19
Carolina	470	4%	79	6%	118	13
Yana-shungo	350	3%	43	3%	58	5
Gabriela	90	1%	6	0%	45	3
Chaucha	80	1%	16	1%	27	5
Suprema	15	0%	21	2%	15	11
Americana	0	0%	1	0%	.	1
Tulca	0	0%	0	0%	.	.
Natividad		0%	90	7%		45
Norteña		0%	0	0%		.

637   TABLE 3. Pests and diseases reported by farmers in Tungurahua, Ecuador, in order by the frequency of reports.
638   Pests and diseases known to cause seed degeneration are indicated.
639
640

Pathogen/disease or pest	Causing degeneration	Women reporting	Men reporting	% farmers
Phytophthora infestans (Late blight)	Yes	15	30	94
Premnotrypes spp. (Andean potato weevil)	Yes	10	26	75
Rhizoctonia solani (Potato black leg)	Yes	7	16	48
Puccinia pittieriana (Common rust)	No	6	12	38
Epitrix spp. (Potato flea beetles)	Yes	3	9	25
Phtorimaea operculella, Symmestrichema tangolias, Tecia solanivora (Potato moths)	Yes	4	4	17
Fusarium spp. (Fusarium rot)	Yes	1	6	15
*Liriomyza spp.** (Leaf miner)	Yes	5	2	15
Slugs*	No	4	0	8
Frankliniella tuberosi (Thrips)	Yes	2	1	6
Nematodes	Yes	1	1	4
Spongospora subterranean (Powdery scab)	Yes	1	1	4
Septoria lycopersici (Annular leaf spot)	No	0	1	2
Viruses	Yes	1	0	2
White fly	Yes	1	0	2

641   *Gender differences (*) are significant in a Chi square test (a=0.05, df=1)*
642

Figures

Fig. 1. Potato production by farmers in the CONPAPA seed system in Tungurahua Province,

Ecuador (photos: J. F. Hernandez Nopsa).

Fig. 2. A seed system transaction network in which nodes represent 48 farmers associated with

CONPAPA in Tungurahua, Ecuador, along with other institutions and individuals linked with

them. Links indicate potato movement, and are weighted by the volume (proportional to line

thickness) of seed potato and table potato bought, sold, used or traded by farmers. Data are from

the most recent season reported in November 2015. Black lines indicate seed, and gray lines

represent potato for food consumption. Self-loops represent seed produced on-farm. Dotted lines

represent transactions where volumes were not reported.

Fig. 3. A network depicting farmer-reported information sources for integrated pest and disease

management (IPM). Link thickness is proportional to the reported level of trust that the farmer

has in that source of information.

Fig. 4. Disease invasion is simulated with initial infection starting at a random node and

proceeding through the network defined by farmer transactions for seed (black) lines and table

potato (grey). Link widths are scaled to volume of transaction. This network represents the last

two seasons as well as the current season reported in November 2015. The risk at each node is

evaluated in terms of the number of nodes that would become infected before the disease was

detected at that node, if it were the node being used for monitoring. Monitoring a low risk node

(blue) would mean that only a small number of nodes become infected before disease is detected

at that node. Three scenarios were evaluated, where the probability/risk of the disease starting at

a given farmer node is weighted differently for **A,** all farmers are equally likely to be an initial

source of spread; for **B,** risk of being an initial source decreases as the maximum quality of

information increases (per the IPM information network in Fig. 3); and for **C,** risk of being an

initial source decreases based on the number of information sources (node in-degree, not

including self-loops) as depicted in the IPM information network in Fig. 3.

Fig. 5. A scenario analysis evaluating potential compensation in the system if the CONPAPA

leadership group no longer played its central role. The figure compares the current scenario,

where it provides the majority of seed (**A),** versus a hypothetical scenario where farmers get their

seed from the nearest seed multiplier (**B),** and CONPAPA no longer plays a role. **A,** Seed

transactions weighted by the volume based on reports from the last three plantings, including

CONPAPA. **B,** Seed transactions weighted by volume in a scenario where seed normally going

from CONPAPA to multipliers was replaced with seed from the government agency (INIAP).

Seed that went from the CONPAPA leadership group to farmers is now provided by the nearest

multiplier. Active multipliers are farmers 7, 27, 34 and 46.

Fig. 1.

 Fig. 3.

 Fig 4. (A)

Farmer risk of invasion weighted equally

Hernandez Nopsa, Buddenhagen, et al. – 34

Farmer risk weighted by quality of information

Farmer risk weighted by number of information sources

690   **Fig. 5. (A)**

 Fig. 5. (B)