Stratified Linkage Disequilibrium Score Regression reveals enrichment of eQTL effects

on complex traits is not tissue specific

Authors

Hill F. Ip,[1]* Rick Jansen,[2] Abdel Abdellaoui,[1] Meike Bartels,[1,3,4] , UK Brain Expression Consortium,

Dorret I. Boomsma,[1,3,4] & Michel G. Nivard[1]**

Affiliations

[1] Department of Biological Psychology, Vrije Universiteit Amsterdam, the Netherlands

[2] Department of Psychiatry, VU University Medical Center, Amsterdam, the Netherlands

[3] Amsterdam Public Health Research Institute, Amsterdam, the Netherlands

[4] Neuroscience Amsterdam, the Netherlands

Corresponding authors

* h.f.ip@vu.nl

** m.g.nivard@vu.nl

Abstract

Both gene expression levels and eQTLs (expression quantitative trait loci) are partially tissue-specific, complicating the detection of eQTLs in tissues with limited sample availability, such as the brain. However, eQTL overlap between tissues might be non-trivial, allowing for inference of eQTL functioning in the brain via eQTLs measured in readily accessible tissues, e.g. whole blood. Using Stratified Linkage Disequilibrium Score Regression (SLDSR), we quantify the enrichment in GWAS signal of blood and brain eQTLs in genome-wide association study (GWAS) on 11 complex traits (schizophrenia, BMI, educational attainment, Crohn's disease, rheumatoid arthritis, ulcerative colitis, age at menarche, coronary artery disease, height, LDL levels, and smoking behavior). Our analyses established significant enrichment of blood and brain eQTLs in their effects across all traits. As we do not know the true number of causal eQTLs, it is difficult to determine the precise magnitude of enrichment. We found no evidence for tissue-specific enrichment in GWAS signal for either eQTLs uniquely seen in the brain or whole blood. To follow up on our findings, we tested tissue-specific enrichment of eQTLs discovered in 44 tissues by the Genotype-Tissue Expression (GTEx) consortium, and, again, found no tissue-specific eQTL effects. We further integrated the GTEx eQTLs with SNPs associated with tissue-specific histone modifiers, and interrogate its effect on rheumatoid arthritis and schizophrenia. We observed substantially enriched effects on schizophrenia, though again not tissue-specific. Finally, we extracted eQTLs in tissue-specific differentially expressed genes, and determined their effects on rheumatoid arthritis and schizophrenia. We conclude that, while eQTLs are strongly enriched in GWAS signal, the enrichment is not specific to the tissue used in eQTL discovery. Therefore, working with relatively accessible tissues, such as whole blood, as proxy for eQTL discovery is sensible; and restricting lookups for GWAS hits to a specific tissue might not be advisable.

Key words

49   eQTL; gene expression; tissue-specificity; complex traits; enrichment; genome-wide; brain; whole

50   blood; stratified linkage disequilibrium score regression; SLDSR; eQTL discovery

Introduction

The aim of genome-wide association studies (GWASs) is to detect statistically significant associations

between genetic variants, such as single nucleotide polymorphisms (SNPs), and a trait of interest

(Hirschhorn and Daly 2005). GWASs have identified many genetic variants and thereby provided

insights into the genetic architecture of complex traits (Hirschhorn and Daly 2005; Visscher et al.

2012). However, as a large number of variants identified through GWASs are located outside of

coding regions and specific knowledge of regulatory elements is limited, uncovering a relationship

between GWAS hits and biological function has proven to be complicated (Lowe and Reddy 2015).

Expression quantitative trait loci (eQTLs) contain SNPs that influence gene expression, and are not

necessarily located in coding regions. eQTLs may aid functional annotation of SNPs that have been

identified in a GWAS and are located outside of coding regions (Morley et al. 2004; Lowe and Reddy

2015). Previous work has found substantial enrichment of eQTLs among GWAS hits (Manolio et al.

2009; Nicolae et al. 2010; Torres et al. 2014) and an enrichment in their genome-wide effect on

complex traits (Davis et al. 2013). Therefore, eQTLs are viewed as an important tool in moving from

genome-wide association to biological interpretation.

67   As a result of difference in gene expression between cells originating from different tissues,

eQTLs are potentially tissue-specific (Hernandez et al. 2012; GTEx Consortium 2015). Tissue-

specificity poses no problem if the tissue of interest is readily available for research, such as whole

blood. However, discovery of eQTLs gets complicated when measurement of expression levels in a

tissue is limited by ethical and practical considerations, for example in brain tissue. Several studies

have shown that the overlap between eQTLs from different tissues might actually be larger than

initially assumed (Ding et al. 2010; Nica et al. 2011). The Genotype-Tissue Expression (GTEx)

consortium identified eQTLs in a wide range of human tissues and showed that 54-90% of the eQTLs

identified in one tissue are also designated as an eQTL in at least one other tissue (GTEx Consortium

2015; Aguet et al. 2016). In another study, Liu *et al* (2016) found a high average pairwise genetic

correlation (r_g=0.738) of local gene expression between tissues. Nevertheless, small differences in

terms of eQTL effect may be of considerable importance in terms of the effect an eQTL might have

on complex traits related to specific tissues. It is, therefore, worthwhile to investigate the specific

utility of tissue-specific eQTLs in their effect on complex traits, as studied in GWAS, as the discovery

of eQTLs for tissues such as the brain might be advanced by eQTLs discovered in more accessible

tissues, such as whole blood. The use of accessible tissues, though, depends on a substantial degree

of similarity of eQTL effect across tissue, and to what extend eQTL differences between tissues are

important in complex trait etiology.

Stratified Linkage Disequilibrium Score Regression (SLDSR) is a technique that estimates the

SNP-heritability (h^2_{SNP}) of a trait based on GWAS summary statistics (Bulik-Sullivan et al. 2015;

Finucane et al. 2015). By simultaneously analyzing multiple categories of SNPs (annotations), SLDSR

can partition h^2_{SNP} by annotation (h^2_{annot}) and thereby provides a way to jointly quantify the

enrichment in GWAS signal of several annotations. Here, we extend SLDSR by including annotations

containing *cis*-eQTLs, i.e. eQTLs located closely to the gene with which they associate (Brem et al.

2002; Ramasamy et al. 2014), discovered in multiple tissues. To this end, we perform analyses based

on representative eQTL resources, and consider a variety of traits as outcomes.

Firstly, we selected all eQTLs per gene discovered in large samples of RNA expression levels

assessed in whole blood (N=4896)(Wright et al. 2014; Jansen et al. 2017) and in brain tissues (N=134)

(Ramasamy et al. 2014), and quantified the contribution of these blood and brain eQTLs to the

genetic variance in complex traits captured in GWAS. We then estimated tissue-specific eQTL effects

on complex traits by quantifying the enrichments of eQTLs uniquely found in whole blood or

uniquely found in brain, conditional on the enrichment of the complete blood eQTL annotation or

complete brain eQTL annotation, respectively. We considered the effect of eQTLs on 11 complex

traits: schizophrenia, BMI, educational attainment, Crohn's disease, rheumatoid arthritis, ulcerative

colitis, age at menarche, coronary artery disease, height, LDL levels, and smoking behavior.

Secondly, we retrieved all eQTLs identified in any of the 44 tissues from the GTEx consortium

(N=70-361, median=126.5)(GTEx Consortium 2015; Aguet et al. 2016). We considered the

enrichment in GWAS signal of the union of all GTEx eQTLs, and, additionally, the enrichment of

tissue-specific eQTL effects on top of the union of all GTEx eQTLs. We expected to observe tissue-

specific enrichment of eQTLs in their effects on complex traits related to the tissue in question, e.g.

eQTLs discovered in immune-related tissues are expected to show higher enrichments in their effect

on immune-related traits compared to eQTLs found in skin tissue. We considered tissue-specific

enrichment of *cis*-eQTLs in their effect on schizophrenia (a disorder where there is strong prior

evidence for the involvement of processes in the brain) and rheumatoid arthritis (a disease with

strong prior evidence for the involvement of processes in immune tissue) as GWAS for these traits

are well powered for extended LD-score-based analyses. We further considered the enrichment of

the intersection of *cis*-eQTLs discovered in any tissue, and histone modification in a specific tissue

(i.e. tissue-specific epigenetically changed chromatin states in regulatory regions). Finally, we

explored the enrichment in GWAS signal of eQTLs for tissue-specific differentially expressed genes.

Our analyses were designed to elucidate the relation between eQTLs and complex traits, and

to quantify the extent to which this relation is dependent on the tissue used in eQTL discovery. Our

analysis further considered the enrichment of genomic regions related to gene expression and

epigenetically modified in specific tissues.

Material and Methods

SLDSR method

A measure of linkage disequilibrium (LD) for each SNP, called an "LD score", can be computed by

taking the sum of correlations between that SNP and all neighboring SNPs (Bulik-Sullivan et al. 2015;

Finucane et al. 2015). Under a polygenic model, LD scores are expected to show a linear relationship

with GWAS test statistics of corresponding SNPs, where the slope is proportional to h^2_{SNP}. For SLDSR,

LD scores are based on only (functional) parts of the genome and used as predictors in a multiple

linear regression (Finucane et al. 2015). In this manner, SLDSR is able to partition h^2_{SNP} into parts that

are explained by these parts of the genome (i.e. h^2_{annot}), while accounting for influences of the

remaining annotations in the model. The enrichment of an annotation is then obtained by taking the

ratio of h^2_{annot} over the proportion of SNPs that fall within that annotation. For eQTLs, the

denominator, i.e. the number of SNPs in the annotation, is a complicated quantity: not all significant

eQTLs are likely causal; whereas including only lead, or putative causal, eQTLs may result in very

small annotations located near genes and other regulatory elements, which presents a risk of

inflated estimates of the enrichment in GWAS signal. What constitutes an eQTL is sufficiently vague

and open for interpretation for us to consider the effect of multiple inclusion rules for inclusion of a

SNP into the eQTL annotation. Since eQTLs are essentially discovered in what amounts to a local

GWAS, we expect the average LD score of eQTLs to be higher than that of an average SNP, which

may influence the results of downstream SLDSR analysis. In order to break the relation between LD

score and probability of inclusion, we consider eQTL annotations which are based on a sample of

from all significant eQTLs for a given probe. First, we included the most strongly associated SNP, a

SNP with a high expected LD score. Second, we included one SNP per probe with a median p-value.

Third, we included one SNP per probe with a mean p-value. Fourth, we included the 10 most

strongly associated SNPs per probe. Finally, we included all SNPs significantly associated with gene

expression after FDR correction at $\alpha=0.05$. We add each annotation separately to the baseline

categories in an SLDSR model, and determined how the various p-value thresholds influence the

SLDSR coefficient and its test statistic. For each annotation, we looked up the SNPs in the baseline

category, and extracted their baseline LD scores and minor allele frequencies (MAF). We then

compared the mean LD score, median LD score and mean MAF between the annotations and the

entire baseline category. Based on the results (S1 Figure – S2 Figure, S3 Table), we consider all

significant *cis*-eQTLs as an annotation, and retain additional gene-centric and regulatory annotations

ins the model.

Target traits

As outcome for SLDSR, we used summary statistics of GWASs on Crohn's disease (Jostins et al. 2012),

rheumatoid arthritis (Okada et al. 2014), ulcerative colitis (Jostins et al. 2012), BMI (EK et al. 2010),

educational attainment (Rietveld et al. 2013), schizophrenia (Ripke et al. 2014), age at menarche

(Perry et al. 2014), coronary artery disease (Schunkert et al. 2011), height (Allen et al. 2010), LDL

levels (Teslovich et al. 2010), and smoking behavior (Furberg et al. 2010). The first three traits were

chosen because they are related to the immune system and are therefore expected to reveal

considerable enrichment of blood eQTL signal (Jostins et al. 2012; Okada et al. 2014). Similarly, brain

eQTLs are expected to show substantial enriched effects due to previous reports on the involvement

of the central nervous system (CNS) in schizophrenia (Ripke et al. 2014), educational attainment

(Rietveld et al. 2013), and BMI (Vimaleswaran et al. 2012). Of course, these traits do not perfectly

align with either tissue, e.g. the immune system has been implicated in the etiology of schizophrenia

(Andreassen et al. 2015) and BMI (Karalis et al. 2009), and might therefore also be enriched in their

effects for the other eQTL set. Enrichment of blood and brain eQTL effects on the remaining traits

was calculated to contrast the results with traits for which we do not have a strong *a priori*

expectation of the relationship between trait and tissue.

The discovery sample for detection of blood eQTLs (Wright et al. 2014; Jansen et al. 2017)

included participants from the Netherlands Twin Register (NTR)(Boomsma et al. 2008) and

participants from the Netherlands Study of Depression and Anxiety (NESDA)(Penninx et al. 2008).

These two cohorts did not participate in the GWAS for schizophrenia, Crohn's disease, rheumatoid

arthritis, ulcerative colitis, or coronary artery disease. However, participants from these two cohorts,

not necessarily the same ones, did participate in the GWAS for height, BMI, LDL levels, smoking

behavior, educational attainment, and age at menarche. For educational attainment and smoking

behavior, we were able to obtain summary statistics omitting subjects from NTR/NESDA. For both

these traits, we looked at trait-specific enrichment of blood and brain eQTL effect in GWAS signal,

comparing results from using publicly available datasets with using summary statistics based on the

same sample without subjects from the NTR or NESDA. The results did not reveal appreciable

differences between the respective datasets for educational attainment, but did show substantial

differences for smoking behavior (S4 Figure). This latter finding could conceivably be a function of

relatively strong effects of smoking behavior on gene-expression levels (Vink et al. 2015). Therefore,

the remaining analyses for smoking behavior were performed using the summary statistics omitting

the NTR and NESDA, whereas analyses for the remaining traits (height, BMI, LDL levels, and

educational attainment) were run using publicly available summary statistics. This caveat only

applies to eQTL annotations based on NTR/NESDA data (i.e. whole blood). We note that the issue of

overlap also applies to other techniques where the error covariance is assumed to be zero (e.g.

TWAS, mendelian randomization analysis, SMR, etc.).

Blood and brain eQTL enrichment

A catalog of whole blood *cis*-eQTLs was obtained from Jansen *et al* (2017; Wright et al. 2014), where

all eQTLs significantly associated with gene expression in whole blood for each probe set were

selected for inclusion in our whole blood eQTL annotation. A list of brain eQTLs was obtained from

the UK brain expression consortium (UKBEC), for which the analyses are described in Ramasamy *et al*

(2014). We based the brain eQTL annotation on SNPs that were significantly associated with the

average gene expression across 12 brain regions. SLDSR annotations were constructed as per the

instructions in Bulik-Sullivan et al. and Finucane et al. (2015). To guard against upward bias in the

eQTL enrichment signal, two extra annotations containing SNPs within a 500 base pair (bp) and

100bp window around any eQTL were constructed for each eQTL set (Finucane et al. 2015). To

ensure that the enrichment of eQTL effects in GWAS signal was not in fact caused by their proximity

to the genes they influence, an additional gene centric annotation was computed, which contained

all genes for which eQTLs were included. Finally, we performed an inverse-variance weighted meta-

analysis across the traits to determine the average effect of blood and brain eQTLs on complex traits

in general.

Tissue-specific eQTL enrichment

To distinguish between the effects of blood- and brain-specific eQTLs, we split each annotation into

two sets based on the overlap in genes that were tagged by eQTLs from both tissue. That is, the

brain eQTL annotation was split into an annotation of brain eQTLs which regulate genes for which

also at least one blood eQTL was found, and a second annotation of eQTLs that tagged genes for

which only brain eQTLs were found. Likewise, the blood eQTL annotation was split into an

annotation containing only eQTLs that tagged genes for which eQTLs from both tissue was found,

and an annotation consisting of blood eQTLs that tagged genes for which only eQTLs have been

found in blood. In doing so, we are saying that the effect of an eQTL is mediated via the gene it is

associated with. Then, if two different SNPs are associated with the expression of the same gene,

but in different tissues, this gene is likely the mechanism by which the SNP influences a trait.

Contrary, when the same SNP affects different genes in different tissues, this SNP can be seen to

have a tissue-specific mediation of its effect. We thus define eQTLs shared across tissue as eQTLs

that influence a common gene in separate tissues.

Enrichment of eQTLs from 44 tissues

There are several limitations to above mentioned analyses of tissue-specific enrichments of eQTL

effects in GWAS signal. The eQTLs are obtained from two different projects, which vary in terms of

sample size and their definition of an eQTL. To mitigate the heterogeneity between studies, and to

extend to additional tissues. We performed additional analyses using eQTLs obtained by a common

pipeline from 44 tissues (see S5 Table) and based on a broader eQTL locus definition (GTEx

Consortium 2015; Aguet et al. 2016). For each of the 44 tissues, we created annotations for analysis

229 in SLDSR following the previously described procedure. Analogous to the procedure of Finucane *et al*

230 (2015) for cell-type-specific analysis using SLDSR, we did not specify windows for the individual GTEx

231 annotations, but included an additional annotation that contained the union of all GTEx eQTLs, i.e.

232 all SNPs that are designated as part of at least one of the 44 individual GTEx annotations, and added

233 a 100bp and 500bp window around this union of GTEx eQTLs. SLDSR is essentially a multiple

234 regression and, due to the high number of models and high number of predictors (i.e. annotations)

235 in each model, has a high multiple testing burden. As such, it requires well-powered GWAS in order

236 to accurately partition the heritability the various annotations (Bulik-Sullivan et al. 2015; Finucane et

237 al. 2015). Based on the Z-score of the SNP-heritability and previous reports of substantial influence

238 of either tissue in the etiology of the traits (Okada et al. 2014; Ripke et al. 2014; Finucane et al. 2015;

239 Finucane et al. 2017), we obtained two well-powered traits, one for which we assume there to be

240 significant enrichment in signal for blood eQTLs (rheumatoid arthritis) and one for brain eQTLs

241 (schizophrenia). For each of these two traits, we ran one SLDSR model containing only the baseline

242 categories and the union of GTEx eQTLs, and 44 additional models with the two previous

243 annotations and one of the individual GTEx annotations at a time.

244  GTEx has relative small sample sizes for the brain eQTL discovery (mean=89 sample size,

245 range=72-103) compared to other tissues (mean=160 sample size, range=70-361) (GTEx Consortium

246 2015; Aguet et al. 2016). To investigate the effect of differences in sample size on estimates of

247 enrichments in GWAS signal, we collapsed the union of individual brain eQTL annotations into a

248 shared brain eQTL annotation (i.e. an eQTL found in at least one of the GTEx brain annotations was

249 included in the shared brain eQTL annotation). This annotation was then analyzed as an additional

250 GTEx eQTL annotation. We further tested the relationship between tissue sample size and tissue

251 eQTL enrichment.

253 **Enrichment of the intersection between eQTLs and histone marks**

The availability of annotations based on tissue-specific histone marks made it possible to create an

annotation that represents the intersection between eQTLs and this type of epigenetic modification

related to enhancers and promoters of actively transcribed genes. We obtained LD score

annotations of SNPs in regions that bear histone marks in cells from the CNS or immune system from

Finucane *et al* (2015). Out of the 220 cell-type-specific histone mark that were available, 101 were

found in the CNS or immune tissues. For each of the 101 annotations of SNPs in cell-type-specific

histone marks, we extracted its intersection with the union of GTEx eQTLs and made a new

annotation of eQTLs which intersected with histone marks (i.e. SNPs found in both annotations). We

then analyzed each of the intersection annotations individually in a model together with the baseline

categories, the union of GTEx eQTLs, and the corresponding cell-type-specific histone marks.

Enrichments in GWAS signal of the intersection should be interpreted as enrichment of genome-

wide SNP effects on a complex trait beyond the additive effects which work on all SNPs that are a

cis-eQTL and histone mark in question. In fact, we test whether the interaction between tissue-

specific chromatin state and eQTLs are enriched in their genome-wide effect on complex traits.

Intersection of GTEx eQTLs and tissue-specific differentially expressed genes

Finucane *et al* (2017) looked at tissue-specific gene expression and determined that the top 10% of

these differentially expressed genes are substantially enriched in their effects in GWAS signals for a

wide range of traits. Here, we build on these findings by taking the top 10% most highly differentially

expressed genes in the 44 GTEx tissues; obtaining the eQTLs for these specific genes, regardless of

the discovery tissue; and added these separately as an annotation to an SLDSR model together with

the baseline categories and union of all GTEx eQTLs. A significant increase in enrichment in GWAS

signal in the eQTLs compared to the genes themselves, would indicate that eQTLs explain part of the

enrichment seen by Finucane *et al*.

Results

Effect of eQTL selection on regression parameter and test statistics

We compare five annotations that included various SNPs based on the *p*-value of their associations with gene-expression levels (lead eQTL, median eQTL, mean eQTL, top 10 lead eQTLs, and all eQTLs). Supplementary table S3 shows various metrics of these annotations. Surprisingly, lead eQTLs had the lowest mean and median LD score amongst the annotations, indicating that the annotation contains less signal (S3 Table). However, it was still higher compared to the mean or median of all SNPs in the baseline annotation. Including all significant eQTLs in the annotation resulted in the highest mean and median LD score. All annotations had a mean MAF 0.27-0.28, whereas the mean MAF of the entire baseline category was 0.24. Smaller annotations had a higher enrichment in GWAS signal (S1 Figure). However, the enrichment in GWAS signal did not differ between taking the lead eQTL, eQTLs with a mean *p*-value, or eQTLs with a median *p*-value. Finally, coefficient test statistics did not differ much between the annotations (S2 Figure). Based on the explorative analysis, we selected the annotation based on all significant eQTLs for further analysis.

Blood and brain eQTL enrichment

We fitted an SLDSR model containing the baseline categories; the complete annotation for both brain and blood eQTL tissues, their 100 and 500bp windows, and gene-centric annotations to all traits (Crohn's disease, rheumatoid arthritis, ulcerative colitis, BMI, educational attainment, schizophrenia, age at menarche, coronary artery disease, height, LDL levels, and smoking behavior). We found significant effects of brain eQTLs on educational attainment, rheumatoid arthritis, smoking behavior, and schizophrenia, and significant effect of blood eQTLs on height and smoking behavior (see S6 Table). We then meta-analyzed the results for all annotations, both in the baseline model, and those associated with eQTLs across the 11 traits. Our analyses revealed significant effect of both blood ($p < 0.001$) and brain ($p < 0.001$) eQTL effects on all traits (Figure 1, S7 Table), exceeding, in terms of significance, all the baseline categories considered by Finucane *et al* (2015)

but conserved genomic regions. The gene-centric annotation for both blood and brain eQTLs showed

no effect on any trait.

We then separated the list of blood eQTLs into a list of unique and shared blood eQTLs

based on the overlap in target genes with brain eQTLs, and modelled the baseline categories

together with all blood eQTLs and the unique blood eQTLs. We did the same for brain eQTLs. We

observed no evidence for depletion of blood-specific eQTLs (relative to all blood eQTLs) on brain-

related traits, nor do we find significant depletion of effect on immune-related traits of eQTLs

associated with genes for which eQTLs were solely identified in brain tissue (Table I and Table II).

Enrichment of eQTLs from 44 tissues in GTEx

We interrogated the enrichment of the union of GTEx eQTLs and 44 individual GTEx annotations in

their effect on schizophrenia and rheumatoid arthritis. Figure 2 shows the coefficient of the 45 GTEx

annotations, sorted on their Z-scores for rheumatoid arthritis. In both cases, the union of GTEx

eQTLs contributed significantly to explaining the polygenic signal(S8 Table), indicating that eQTLs

were significantly enriched in their effects on complex traits. The individual annotations, however,

performed notably worse and in some cases even suggested depletion of genome-wide effects of

tissue-specific eQTLs on schizophrenia and rheumatoid arthritis. For rheumatoid arthritis, the

coefficient Z-scores of the whole blood annotation reached nominal significance (Z=2.251), but failed

correction for multiple testing. None of the other annotations reached nominal significance. The

union of all GTEx brain annotations did not contribute significantly to explaining h^2_{SNP} (Z=0.147,

p=0.441). Sample size in the eQTL discovery phase appears to be a strong determinant of tissue-

specific enrichment in GWAS signal. The correlation coefficients between the coefficient Z-scores

and sample sizes were 0.6453 (p=2.253*10^{-6}) and 0.4247 (p=0.004) for schizophrenia and

rheumatoid arthritis, respectively.

Enrichment of the intersection between eQTLs and histone marks

We interrogated the intersection of eQTLs and histone marks found in specific CNS and immune cells,

and estimated the enrichment of the intersection in its effect on rheumatoid arthritis and

schizophrenia. We found significant enrichment in GWAS signal for eQTLs that intersect with

histones that bear modification H3K4me1, a modification thought to be present in the enhancer of

actively transcribed genes (Zhou et al. 2011; Allis and Jenuwein 2016), in CNS cells for schizophrenia

(see Figure 3). There was some evidence for significant enrichment of eQTLs that intersected with

genomic regions in immune cells baring the H3K4me1 mark in their effect on schizophrenia, but not

on rheumatoid arthritis. Specifically, none of the intersecting annotations showed evidence of

enrichment for rheumatoid arthritis. For the separate annotations, we found significant enrichment

in GWAS signal across all histone marks found in CNS cells and three significant immune cell-types

that bear the H3K4me3 modification, a modification associated with transcriptional start sites and

promoters of actively transcribed genes (Zhou et al. 2011; Allis and Jenuwein 2016), for

schizophrenia (S9 Figure). The opposite picture was seen for rheumatoid arthritis: a wide variety of

immune-cell specific histone marks showed significant enrichments in GWAS signal, while all marks

found in CNS cells were below zero. The union of GTEx eQTLs reached statistical significance for all

models.

Intersection of GTEx eQTLs and tissue-specific differentially expressed genes

We extracted all eQTLs within the top 10% of tissue-specific differentially expressed genes in all 44

GTEx tissues. We then compare the enrichment in GWAS signal for these eQTLs with the genes

themselves. The correlation between the coefficients was 0.58 and 0.24 for schizophrenia and

rheumatoid arthritis, respectively. For schizophrenia, we see that the eQTL annotation most brain

tissues have the highest regression coefficient and Z-score, although none reached the significance

threshold (S10 Table). Furthermore, the eQTLs showed larger coefficients compared to the whole

genes, although the large standard errors made the difference non-significant. Interestingly, the top

10% differentially expressed genes within the nucleus accumbens showed a significant coefficient

comparable to the other brain regions, although the eQTLs for these genes showed a non-significant

depletion. For rheumatoid arthritis, whole blood showed the most significant coefficient, however,

again failed correction for multiple testing (S11 Figure). Furthermore both the whole genes as the

eQTLs for these genes showed a similar regression coefficient.

Discussion

Stratified Linkage Disequilibrium Score Regression provides a way to partition h^2_{SNP} into parts

explained by (functional) parts of the genome (Finucane et al. 2015). A "full baseline model"

containing 24 non-cell-type-specific annotations of SNPs, such as SNPs located in promoters or

coding regions, was developed previously for analysis using SLDSR. Here, we added annotations

containing eQTLs derived from whole blood and brain tissue into the model, and showed that eQTLs

were substantially stronger enriched in their effect on complex traits compared to all categories

considered by Finucane *et al* (2015). The complete brain eQTL annotation was significantly enriched

in GWAS signal for educational attainment, rheumatoid arthritis, smoking behavior, and

schizophrenia. This finding is consistent with previous estimates of eQTL effect enrichment (Davis et

al. 2013). Considerable enrichment for eQTLs, even for traits not apparently linked to the brain or

immune system (e.g. smoking behavior), suggested that non-trivial eQTL overlap across tissues

might be present.

Inclusion of both brain and blood eQTLs into the SLDSR model did not separate the signal

into tissue-specific effects. In general, we are not able to clearly identify tissue-specific eQTL signals

using these datasets and SLDSR. Our second analysis of eQTL enrichment based on 44 tissue-specific

cis-eQTL sets, obtained from the GTEx consortium (2015; Aguet et al. 2016), confirms the lack of

tissue-specific eQTL enrichment. While an annotation containing all eQTLs identified in GTEx is

significantly enriched in its effect on schizophrenia and rheumatoid arthritis (Z=5.501 and Z=3.802,

respectively, both *p*<0.001), none of the analyzed brain tissues are enriched beyond all eQTLs in

their effect on schizophrenia. Similarly, whole blood eQTLs are not significantly enriched beyond all

383 GTEx eQTLs taken together in their effect on rheumatoid arthritis. Again, these findings are not

384 consistent with the hypothesis of abundant tissue-specific *cis*-eQTLs with effects on complex traits

385 related to the specific tissue in question. Our findings are consistent with a lack of power to detect

386 any tissue-specific eQTL effects. Especially, when contrasted with tissue-specific gene expression

387 levels and tissue-specific histone modifications (Liu et al. 2016; Finucane et al. 2017), tissue-specific

388 eQTLs are of limited value in relating complex traits to a tissue. In fact, considering eQTLs associated

389 with genes expressed in a specific tissue improves our detection of tissue-specific effects. But, while

390 the regression parameter subsets eQTLs for specifically expressed genes have larger effects than the

391 rest of these genes, the significance of the enrichment is weak compared to the significance of the

392 tissue-specific enrichment of the whole gene body plus a 100kb window.

393   One of the limitations of the study presented here involves the substantial differences in

394 discovery sample size between the tissues, which influences the power to detect eQTLs (Lonsdale et

395 al. 2013). Even within the GTEx tissues, where differences in sample sizes are relatively small

396 compared to eQTLs obtained from Jansen *et al* and Ramasamy *et al*, we still see a significant

397 correlation between the discovery sample size and enrichment of eQTLs in GWAS signal. Several

398 methods have been developed to capitalize on cross-tissue overlap in eQTLs to improve power to

399 detect SNP effects on gene expression within tissue (Flutre et al. 2013; Li et al. 2013). The aim of this

400 paper was to explore the possibilities of assessing the effects of eQTLs expressed in whole blood on

401 presumably brain-related traits, and vice versa. In the analyses with eQTLs for differentially

402 expressed genes, we show that enrichment in GWAS signal is stronger in these eQTLs compared to

403 taking the all SNPs in the same genes. This indicates that eQTLs, irrespective of the tissue in which

404 they were discovered, play an important role in the etiology of complex traits, and do so via the

405 gene they are associated with. This, however, does not take away the need to increase sample sizes

406 when performing tissue-specific discovery of (*cis*-)eQTLs. Tissue specificity, in the end, is a relative

407 judgement best reached based on weighing multiple lines of evidence, among which are differential

408 expression, epigenetic regulation, and eQTLs. For eQTLs to play a large role in determining the

tissue-specific effects on complex traits, a continued investment in resources like GTEx is required in

order to increase sample sizes for detection, especially in rare tissues.

Our conclusions are limited to *cis*-eQTLs and it is not unlikely that trans-eQTLs behave

differently in terms of tissue-specificity. We do find evidence for possible enrichment for eQTLs that

intersect with tissue-specific H3K4me1 histone marks in the brain, but also immune cells, in their

effect on schizophrenia but not rheumatoid arthritis. This means that eQTLs in H3k4me1 marks are

enriched in their effect on schizophrenia above the expected enrichment based on the fact that

these SNPs are both eQTLs and located in H3K4me1 histone marks. What is of substantial interest is

that the e

nrichment in GWAS signal appears specific to H3K4me1 marks, and no other Histone marks,

suggesting that these marks specifically can aid in prioritizing genomic regions in which tissue-

specific eQTLs may reside. Though, again, the totality of evidence is inconclusive on the relevance of

tissue-specific eQTLs to variation in complex traits.

Our results are consistent with, and complimentary to, a study investigating the genetic

correlation between gene expression levels across 15 tissues (Liu et al. 2016). This study revealed

substantial correlations between *cis*-genetic effects on gene expression across 15 tissues (Liu et al.

2016). Our analyses confirmed the value of using whole blood as discovery tissue for detection of

cis-eQTLs and further demonstrated the usefulness of techniques that use *cis*-eQTLs discovered in

whole blood to study the etiology of complex traits related to different tissues (Gamazon et al. 2015;

Gusev et al. 2016). The results presented here highlight the overlap of *cis*-eQTL effects across tissues

on a genome-wide level. However, the effect of a *cis*-eQTL might vary substantially across tissues for

individual genes (Grundberg et al. 2012). Our conclusions are based on genome-wide enrichments

and therefore should not be interpreted as limited evidence for tissue-specific eQTL effects for

individual genes. Therefore, eQTL discovery in the tissue most relevant to a specific trait or disorder

remains important to further our understanding of the genetic regulation of tissue-specific gene

expression. What is also clear is that to discover those tissue-specific eQTLs that are of relevance to

the interpretation of GWASs of complex traits, tissue-specific eQTL discovery needs to be refined.

The practice of, as a post-hoc analysis to GWAS, performing eQTL lookup in a specific tissue linked to

a trait, when larger dataset for other accessible tissues are available, may be suboptimal. In fact, one

may prefer to perform a lookup in the overlap between histone modifications in a relevant tissue

and eQTLs regardless of tissue. One can further consider utilizing eQTLs to link GWAS findings to a

gene, and subsequently consider the differential expression of a gene to identify the tissue in which

the gene is most likely to act in effecting the trait. Tissue-specific differential gene expression vastly

outperforms eQTLs in tagging regions of the genome enriched in their effect on complex traits

(Finucane et al. 2017).

It is also evident that a limited dichotomous definition of eQTL/no-eQTL may be insufficient

to identify tissue-specific eQTL effects. An evident improvement would be to compute the *difference*

in eQTL effect on expression of the gene between tissues, and perform inference based on this

difference in effect. eQTLs are strongly enriched SNPs, with clear biological function and utility for

the translation of GWAS findings, though tissue-specific eQTL mechanisms remain elusive. The

discovery of tissue-specific eQTL effects, which can aid in linking complex trait to tissue, may require

novel research strategies.

Supplemental Data

Supplemental Data includes 4 figures and 7 tables

Compliance with Ethical Standards

Funding

MGN is supported by the Royal Netherlands Academy of Science Professor Award (PAH/6635) to DIB.

HFI is supported by the "Aggression in Children: Unraveling gene-environment interplay to inform

Treatment and InterventiON strategies" (ACTION) project. ACTION receives funding from the

European Union Seventh Framework Program (FP7/2007-2013) under grant agreement no 602768.

MB is supported by a University Research Chair of the Vrije Universiteit. The discovery of blood eQTL

was funded by the US National Institute of Mental Health (RC2 MH089951, principal Investigator PFS)

as part of the American Recovery and Reinvestment Act of 2009. We thank T. Lehner (National

Institute of Mental Health) for his support. We acknowledge Hillary Finucane, Raymond Walters and

Benjamin Neale for critical comments on our methods, design and manuscript.

Conflict of Interest

468   The authors declare that they have no conflict of interest

Ethical approval

471   This article does not contain any studies with human participants or animals performed by any of the

472   authors.

Web Resources

Age at menarche summary statistics, www.reprogen.org/data_download.html

Blood eQTLs, https://eqtl.onderzoek.io/

Brain eQTLs, http://www.braineac.org/

Coronary artery disease summary statistics, www.cardiogramplusc4d.org/data-downloads/

Crohn's disease and ulcerative colitis summary statistics, www.ibdgenetics.org/downloads.html

Educational attainment summary statistics, http://www.thessgac.org/data

Full baseline model LD scores, http://data.broadinstitute.org/alkesgroup/LDSCORE/

GTEx dataset, http://www.gtexportal.org/home/datasets

Height and BMI summary statistics,

www.broadinstitute.org/collaboration/giant/index.php/GIANT_consortium_data_files

LDL levels summary statistics, www.broadinstitute.org/mpg/pubs/lipids2010/

Rheumatoid arthritis summary statistics, http://plaza.umin.ac.jp/yokada/datasource/software.htm

Schizophrenia and smoking behavior summary statistics, www.med.unc.edu/pgc/results-and-

downloads

SLDSR software, https://github.com/bulik/ldsc/

References

Aguet F, Brown AA, Castel S, et al (2016) Local genetic effects on gene expression across 44 human

tissues. bioRxiv. doi: https://doi.org/10.1101/074450

Allen HL, Estrada K, Lettre G, et al (2010) Hundreds of variants clustered in genomic loci and

biological pathways affect human height. Nature 467:832–838.

Allis CD, Jenuwein T (2016) The molecular hallmarks of epigenetic control. Nat Rev Genet 17:487–

500.

Andreassen OA, Harbo HF, Wang Y, et al (2015) Genetic pleiotropy between multiple sclerosis and

schizophrenia but not bipolar disorder: differential involvement of immune-related gene loci.

Mol Psychiatry 20:207–14. doi: 10.1038/mp.2013.195

Boomsma DI, Willemsen G, Sullivan PF, et al (2008) Genome-wide association of major depression:

description of samples for the GAIN Major Depressive Disorder Study: NTR and NESDA biobank

projects. Eur J Hum Genet 16:335–342. doi: 10.1038/sj.ejhg.5201979

Brem RB, Yvert G, Clinton R, Kruglyak L (2002) Genetic dissection of transcriptional regulation in

budding yeast. Science 296:752–5. doi: 10.1126/science.1069516

Bulik-Sullivan BK, Finucane HK (2015) LD Score Estimation Tutorial.

https://github.com/bulik/ldsc/wiki/LD-Score-Estimation-Tutorial. Accessed 21 Nov 2015

Bulik-Sullivan BK, Loh P-R, Finucane HK, et al (2015) LD Score regression distinguishes confounding

from polygenicity in genome-wide association studies. Nat Genet 47:291–295. doi:

10.1038/ng.3211

Davis LK, Yu D, Keenan CL, et al (2013) Partitioning the heritability of Tourette syndrome and

obsessive compulsive disorder reveals differences in genetic architecture. PLoS Genet

9:e1003864. doi: 10.1371/journal.pgen.1003864

Ding J, Gudjonsson JE, Liang L, et al (2010) Gene expression in skin and lymphoblastoid cells: Refined

statistical method reveals extensive overlap in cis-eQTL signals. Am J Hum Genet 87:779–89.

doi: 10.1016/j.ajhg.2010.10.024

EK S, CJ W, SI B, et al (2010) Association analyses of 249,796 individuals reveal 18 new loci associated

with body mass index. Nat Genet 42:937.

Finucane H, Reshef Y, Anttila V, et al (2017) Heritability enrichment of specifically expressed genes

identifies disease-relevant tissues and cell types.

Finucane HK, Bulik-Sullivan B, Gusev A, et al (2015) Partitioning heritability by functional annotation

using genome-wide association summary statistics. Nat Genet 47:1228–1235. doi:

10.1038/ng.3404

Flutre T, Wen X, Pritchard J, Stephens M (2013) A Statistical Framework for Joint eQTL Analysis in

Multiple Tissues. PLoS Genet 9:e1003486. doi: 10.1371/journal.pgen.1003486

Furberg H, Kim Y, Dackor J, et al (2010) Genome-wide meta-analyses identify multiple loci associated

with smoking behavior. Nat Genet 42:441–7. doi: 10.1038/ng.571

Gamazon ER, Wheeler HE, Shah KP, et al (2015) A gene-based association method for mapping traits

using reference transcriptome data. Nat Genet 47:1091–1098. doi: 10.1038/ng.3367

Grundberg E, Small KS, Hedman ÅK, et al (2012) Mapping cis- and trans-regulatory effects across

multiple tissues in twins. Nat Genet 44:1084–9. doi: 10.1038/ng.2394

GTEx Consortium (2015) The Genotype-Tissue Expression (GTEx) pilot analysis: Multitissue gene

regulation in humans. Science (80-) 348:648–660.

Gusev A, Ko A, Shi H, et al (2016) Integrative approaches for large-scale transcriptome-wide

association studies. Nat Genet 48:245–252. doi: 10.1038/ng.3506

Hernandez DG, Nalls MA, Moore M, et al (2012) Integration of GWAS SNPs and tissue specific

expression profiling reveal discrete eQTLs for human traits in blood and brain. Neurobiol Dis

47:20–8. doi: 10.1016/j.nbd.2012.03.020

Hirschhorn JN, Daly MJ (2005) Genome-wide association studies for common diseases and complex

traits. Nat Rev Genet 6:95–108. doi: 10.1038/nrg1521

Jansen R, Hottenga J-J, Nivard MG, et al (2017) Conditional eQTL Analysis Reveals Allelic

Heterogeneity of Gene Expression. Hum Mol Genet. doi: 10.1093/hmg/ddx043

Jostins L, Ripke S, Weersma RK, et al (2012) Host-microbe interactions have shaped the genetic

architecture of inflammatory bowel disease. Nature 491:119–124.

Karalis KP, Giannogonas P, Kodela E, et al (2009) Mechanisms of obesity and related pathology:

linking immune responses to metabolic stress. FEBS J 276:5747–5754.

Li G, Shabalin AA, Rusyn I, et al (2013) An Empirical Bayes Approach for Multiple Tissue eQTL

Analysis.

Liu X, Finucane HK, Gusev A, et al (2016) Functional partitioning of local and distal gene expression

regulation in multiple human tissues. bioRxiv 46383. doi: 10.1101/046383

Lonsdale J, Thomas J, Salvatore M, et al (2013) The Genotype-Tissue Expression (GTEx) project. Nat

Genet 45:580–585. doi: 10.1038/ng.2653

Lowe WL, Reddy TE (2015) Genomic approaches for understanding the genetics of complex disease.

Genome Res 25:1432–1441. doi: 10.1101/gr.190603.115

Manolio TA, Collins FS, Cox NJ, et al (2009) Finding the missing heritability of complex diseases.

Nature 461:747–53. doi: 10.1038/nature08494

Morley M, Molony CM, Weber TM, et al (2004) Genetic analysis of genome-wide variation in human

gene expression. Nature 430:743–7. doi: 10.1038/nature02797

Nica AC, Parts L, Glass D, et al (2011) The architecture of gene regulatory variation across multiple

human tissues: the MuTHER study. PLoS Genet 7:e1002003. doi:

10.1371/journal.pgen.1002003

Nicolae DL, Gamazon E, Zhang W, et al (2010) Trait-associated SNPs are more likely to be eQTLs:

annotation to enhance discovery from GWAS. PLoS Genet 6:e1000888. doi:

10.1371/journal.pgen.1000888

Okada Y, Wu D, Trynka G, et al (2014) Genetics of rheumatoid arthritis contributes to biology and

drug discovery. Nature 506:376–381.

Penninx BWJH, Beekman ATF, Smit JH, et al (2008) The Netherlands Study of Depression and Anxiety

(NESDA): rationale, objectives and methods. Int J Methods Psychiatr Res 17:121–140.

Perry JRB, Day F, Elks CE, et al (2014) Parent-of-origin-specific allelic associations among 106 genomic

loci for age at menarche. Nature 514:92–97.

Ramasamy A, Trabzuni D, Guelfi S, et al (2014) Genetic variability in the regulation of gene expression

in ten regions of the human brain. Nat Neurosci 17:1418–28. doi: 10.1038/nn.3801

Rietveld CA, Medland SE, Derringer J, et al (2013) GWAS of 126,559 individuals identifies genetic

variants associated with educational attainment. Science (80-) 340:1467–1471.

Ripke S, Neale BM, Corvin A, et al (2014) Biological insights from 108 schizophrenia-associated

genetic loci. Nature 511:421–427. doi: 10.1038/nature13595

Schunkert H, König IR, Kathiresan S, et al (2011) Large-scale association analysis identifies 13 new

susceptibility loci for coronary artery disease. Nat Genet 43:333–338.

Teslovich TM, Musunuru K, Smith A V, et al (2010) Biological, clinical and population relevance of 95

loci for blood lipids. Nature 466:707–713.

Torres JM, Gamazon ER, Parra EJ, et al (2014) Cross-tissue and tissue-specific eQTLs: partitioning the

heritability of a complex trait. Am J Hum Genet 95:521–34. doi: 10.1016/j.ajhg.2014.10.001

Vimaleswaran KS, Tachmazidou I, Zhao JH, et al (2012) Candidate genes for obesity-susceptibility

show enriched association within a large genome-wide association study for BMI. Hum Mol

Genet 21:4537–42. doi: 10.1093/hmg/dds283

Vink JM, Jansen R, Brooks A, et al (2015) Differential gene expression patterns between smokers and

non-smokers: cause or consequence?

Visscher PM, Brown MA, McCarthy MI, Yang J (2012) Five Years of GWAS Discovery. Am J Hum Genet

90:7–24. doi: 10.1016/j.ajhg.2011.11.029

Wright FA, Sullivan PF, Brooks AI, et al (2014) Heritability and genomics of gene expression in

peripheral blood. Nat Genet 46:430–437. doi: 10.1038/ng.2951

Zhou VW, Goren A, Bernstein BE (2011) Charting histone modifications and the functional

organization of mammalian genomes. Nat Rev Genet 12:7–18.

Figure Titles and Legends

Figure 1. Average enrichment in GWAS signal of the 24 baseline annotations, 4 brain eQTL annotations and 4 blood eQTL annotations.

Bar plot of the average enrichment in GWAS signal across all traits for the 24 main baseline annotations and 8 main eQTL annotations. Grey beans represent the baseline categories. Blue beans represent eQTLs. Black bars indicate average enrichment. Boxes show upper- and lower-bounds of the 95% confidence interval of the mean. Red dots show enrichments for immune-related traits. Horizontal red line indicates enrichment of 1, i.e. no enrichment.

Figure 2. Coefficient Z-scores of the 45 GTEx annotations

Barplot of coefficient z-scores for all GTEx annotations for schizophrenia (grey) and rheumatoid arthritis (red). Bars are sorted from highest to lowest based on the results from schizophrenia. Horizontal dotted line indicates Bonferroni threshold for 45 tests. Two asterisks indicate bars passing Bonferroni correction for multiple testing.

Figure 3. Coefficient Z-score of intersection between union of GTEx eQTLs and cell-type-specific histone marks

Top two graphs show coefficient Z-scores for schizophrenia. Bottom two graphs show the same for rheumatoid arthritis. Grey bars indicate histone marks found in cells from the central nervous system. Red bars represent histone marks found in cells from the immune system. From dark to light, shades of the bars indicate histone marks H3K27ac, H3K4me1, H3K4me3, and H3K9ac. Vertical dotted lines indicate separation between histone marks. One asterisk above the bars indicate annotations passing FDR correction for multiple testing. Two asterisks indicate bars passing Bonferroni correction for multiple testing. Horizontal dotted line indicates Bonferroni threshold for 101 tests.

Tables

Table I. (see attachments)

Table II. (see attachments)