Title: Field-based species identification in eukaryotes using real-time nanopore

sequencing.

Authors: Joe Parker[1], Andrew J. Helmstetter[1], Dion Devey[1] & Alexander S.T.

Papadopulos[1]

[1]Jodrell Laboratory, Royal Botanic Gardens, Kew, Richmond, Surrey UK. TW9 3AB

Correspondence to: a.papadopulos@kew.org and joe.parker@kew.org

Keywords: Nanopore, MinION, onsite DNA sequencing, phylogenomics

Advances in DNA sequencing and informatics have revolutionised biology over the

past four decades, but technological limitations have left many applications

unexplored[1,2]. Recently, portable, real-time, nanopore sequencing (RTnS) has

become available. This offers opportunities to rapidly collect and analyse genomic

data anywhere[3–5]. However, the generation of datasets from large, complex

genomes has been constrained to laboratories[6,7]. The portability and long DNA

sequences of RTnS offer great potential for field-based species identification, but

the feasibility and accuracy of these technologies for this purpose have not been

assessed. Here, we show that a field-based RTnS analysis of closely-related plant

species (*Arabidopsis spp.*)[8] has many advantages over laboratory-based high-

throughput sequencing (HTS) methods for species level identification-by-

sequencing and *de novo* phylogenomics. Samples were collected and sequenced in

a single day by RTnS using a portable, "*al fresco*" laboratory. Our analyses

demonstrate that correctly identifying unknown reads from matches to a reference database with RTnS reads enables rapid and confident species identification. Individually annotated RTnS reads can be used to infer the evolutionary relationships of *A. thaliana*. Furthermore, hybrid genome assembly with RTnS and HTS reads substantially improved upon a genome assembled from HTS reads alone. Field-based RTnS makes real-time, rapid specimen identification and genome wide analyses possible. These technological advances are set to revolutionise research in the biological sciences[9] and have broad implications for conservation, taxonomy, border agencies and citizen science.

DNA sequencing used to be a slow undertaking, but the past decade has seen an explosion in HTS methods[2,10]. DNA barcoding (i.e., the use of a few, short DNA sequences to identify organisms) has benefited from this sequencing revolution[11,12], but has never become fully portable. Samples must be returned to a laboratory for testing and the discrimination of closely related species using few genes can be problematic due to evolutionary phenomena (e.g. lineage sorting, shared polymorphism and hybridisation)[10]. While typical barcoding approaches have been effective for generic level identification, accuracy is much more limited at the species level[11,13] and concerns remain[14]. Species delimitation using limited sequencing information has also been problematic and is thought to heavily underestimate species diversity[11,15]. Consequently, increasingly elaborate analytical methods have been spawned to mitigate the inherent limitations of short sequences[13,16]. The Oxford Nanopore Technologies® MinION® is one of a new generation of RTnS DNA sequencers that is small enough to be portable for fieldwork and produces data within minutes[17,18]. These properties suggest species identification could be conducted using genome scale data generated at the point of sample collection. Furthermore, the large number of long reads generated[17] may provide more accurate species-level identification than current approaches. This application offers great potential

for conservation, environmental biology, evolutionary biology and combating wildlife crime,

however, this potentially exciting combination of methods has not yet been tested in the

field for eukaryotes.

Our experiment was designed to determine whether DNA reads produced entirely in

the field could accurately identify and distinguish samples from closely-related species (*A.*

thaliana (L.) Heynh. and *A. lyrata* (L.) O'Kane & Al-Shehbaz). Recent analyses have

shown that gene flow has been common and shared polymorphisms are abundant

between the morphologically distinct species in *Arabidopsis.* Indeed, the two study species

share >20,000 synonymous SNPs[8], making this a good stress test of genome scale RTnS

sequencing for species discrimination.

The first goal was to extract and sequence shotgun genomic data from higher plant

species in the field using RTnS technology in sufficient quantity for downstream analyses

within hours of the collection of plant tissue (Extended Figure 1). On consecutive days,

tissue was collected from three specimens each of *A. thaliana* and *A. lyrata subsp. petraea*

(Figs. 1b,c) in Snowdonia National Park, and prepared, sequenced and analysed

outdoors in the Croesor Valley (Fig. 1a). Only basic laboratory equipment was used for

DNA extraction and MinION sequencing-library preparation; we did not use a PCR

machine (Fig. 1d; Extended Data Table 1). One specimen of each species was sequenced

with both R7.3 and R9 MinION chemistries. For *A. thaliana,* the RTnS experiment

generated 97k reads with a total yield of 204.6Mbp over fewer than 16h of sequencing

(see Extended Data Table 2)*.* Data generation was slower for *A. lyrata*, over ~90h

sequencing (including three days of sequencing at RBG Kew following a 16h drive), 26k

reads were generated with a total yield of 62.2Mbp. At the time, a limited implementation

of local basecalling was available for the R7.3 data only. Of 1,813 locally basecalled reads,

281 had successful BLAST matches to the reference databases with a correct to incorrect

species ID ratio of 223:30. The same samples were subsequently sequenced using HTS

short read technology (Illumina MiSeq™, paired-end, 300bp; Supplementary note 1).

Mapping reads to available reference genomes for the *A. thaliana* (TAIR10 release[19]) and

two *A. lyrata* assemblies[20,21] indicates approximate RTnS coverage of 2.0x, 0.3x, and 0.3x

for *A. thaliana*, *A. lyrata*, and *A. lyrata ssp. petraea*, respectively; and 19.5x, 11.9x and

12.0x respectively for HTS reads (Extended Data Tables 2 and 3, Supplementary note 2).

These results demonstrate that the entire process (from sample collection thorough to

genome scale sequencing) is now feasible for eukaryotic species within a few hours in

field conditions. As the technology develops, run yields are expected to improve and

implementation of sample indexing will allow many samples to be run on a single flow cell.

88   As expected given the developmental stages of the technologies, the quality and

yield of field sequenced RTnS data was lower than the HTS data (Extended Data Tables 2

and 4). *Arabidopsis thaliana* RTnS reads could be aligned to approx. 50% of the reference

genome (53Mbp) with an average error rate of 20.9%. Indels and mismatches were

present in similar proportions. The *A. lyrata* RTnS data were more problematic with

significantly poorer mapping to the two *A. lyrata* assemblies, whereas, the HTS data

performed relatively well. For the limited number of alignable RTnS reads, error rates were

slightly higher than for *A. thaliana* (22.5% and 23.5%). The poorer RTnS results for *A.*

lyrata may be a consequence of temperature-related reagent degradation in the field or

due to unknown contaminants in the DNA extraction that inhibited library preparation

and/or RTnS sequencing. Despite the smaller yield and lower accuracy of the RTnS

compared to HTS data, the RTnS reads were up to four orders of magnitude longer than

the HTS reads and we predicted they would be useful for species identification, hybrid

genome assembly and phylogenomics.

To explore the utility of these data for species identification, the statistical

performance of field-sequenced (RTnS) and lab-sequenced (HTS) read data was

assessed. Datasets for each species were compared to two databases via BLASTN,

retaining single best-hits: one database contained the *A. thaliana* reference genome and

the second was composed of the two draft *A. lyrata* genomes combined. Reads which

matched a single database were counted as positive matches for that species. The

majority of matching reads hit both databases, which is expected given the close

evolutionary relationships of the species. In these cases, positive identifications were

determined based on four metrics; a) the longest alignment length, b) the highest %

sequence identities and c) the largest number of sequence identities d) the lowest *E*-value

(Extended Data Table 4). Test statistics for each of these metrics were calculated as the

difference of scores (length, % identities, or *E*-value) between 'correct' and 'incorrect'

database matches. The performance of these difference statistics for binary classification

was assessed by investigating the true and false positive rates (by reference to the known

sample species) across a range of threshold difference values (Fig. 2a-d and Extended

Data Figs. 2-4). For both short- and long-read data at thresholds greater than 100bp, the

differences in total alignment lengths (ΔL_T) or number of identities (ΔL_I) are superior to e-

value or % identity biases (Figs. 2a-d). Furthermore, at larger thresholds (i.e., more

conservative tests), RTnS reads retained more accuracy in true- and false-positive

discrimination than HTS data. This proves that whole genome shotgun RTnS is a powerful

method for species identification. We posit that the extremely long length of the observed

'true positive' alignments compared with an inherent length ceiling on false-positive

alignments in a typical BLASTn search is largely responsible for this property.

To evaluate the speed with which species identification can be carried out, we

performed *post hoc* analyses by subsampling the RTnS *A. thaliana* dataset. This simulated

the rate of improvement in species assignment confidence over a short RTnS run. We

classified hits among the subsampled reads based on (i) ΔL_I over a range of threshold

values (ii) mean ΔL_I and (iii) aggregate ΔL_I (Fig. 3). This demonstrates that a high degree

of confidence can be assigned to species identifications over the timescales needed to

generate this much data (i.e., < one hour) and that variation in the accuracy of

identifications quickly stabilises above 1000 reads. Aggregate ΔL_l values rapidly exclude

zero (no signal) or negative (incorrect assignment) values, making this simple and rapidly-

calculated statistic particularly useful for species identification. In a multispecies context,

the slopes of several such log-accumulation curves could be readily compared, for

example (see Supplementary Discussion).

Assembly of large and complicated eukaryotic genomes with RTnS data alone

would require a greater volume of data than available here[7,22–24]. Field extracted samples

are unlikely to be of similar purity to those obtained with more sophisticated laboratory-

based methods, leading to lower yields. As expected, *de novo* assembly of our RTnS data

performed poorly, likely due to insufficient coverage. However, these data do have

potential for hybrid genome assembly approaches. We assembled the HTS data *de novo*

using ABYSS[25] and produced a hybrid assembly with both RTnS and HTS datasets using

HybridSPAdes[26]. The hybrid assembly was an improvement over the HTS-only assembly

(see Extended Data Table 5) with fewer contigs, a total assembly length closer to the

reference (119.0Mbp), N50 and longest contig statistics both increasing substantially and

estimated completeness (CEGMA[27]) of coding loci increased to ~99%. These results

suggest that relatively small quantities of long and short reads can produce useful genome

assemblies when analysed together, an important secondary benefit of field–sequenced

data. The length of typical RTnS reads is similar to that of genomic coding sequences (1-

10kb)[17]. This raises the possibility of extracting useful phylogenetic signal from such data,

despite the relatively high error rates of individual reads. We annotated individual raw *A.*

thaliana reads directly, without genome assembly, which recovered over 2,000 coding loci

from the data sequenced in the first three hours (Fig. 2e). These predicted gene

sequences were combined with a published dataset spanning 852 orthologous, single-

copy genes[28], downsampled to 6 representative taxa. Of our gene models, 207 were

present in the Wickett *et al.*[28] dataset and the best 56 matches were used for

phylogenomic analysis (see Supplementary Methods for details). The resulting

phylogenetic trees (Fig. 2f and Extended Data Fig. 5) are consistent with the established

intergeneric relationships[28]. Although the taxonomic scale used here for phylogenomics is

coarse it highlights an additional benefit to rapid, in-the-field sequencing for evolutionary

research.

This experiment is the first to demonstrate field-based sequencing of higher plant

species. When directly compared to lab-based HTS, our experiment highlights key

discriminatory metrics for highly accurate species identifications using portable RTnS

sequencing. Few approaches can boast this level of discriminatory power and none of

these have the same degree of portability[10,11]. The data produced for identification is also

useful for genome assembly. Entire coding sequences can be recovered from single reads

and incorporated into evolutionary analyses. Clearly, data generated with the goal of

accurate species identification has much broader usefulness for genomic and evolutionary

research. Few technical barriers remain to prevent the adoption of portable RTnS by non-

specialists, or even keen amateurs and schoolchildren. As these tools mature, and the

number of users expands, portable RTnS sequencing can revolutionise the way in which

researchers and practitioners can approach ecological, evolutionary and conservation

questions.

Methods summary: Genomic DNA was extracted from two plant specimens and

sequenced on Oxford Nanopore MinION devices according to manufacturers'

recommendations in a portable outdoor laboratory. Offline basecalling software and local

BLAST (v2.2.31) were used to identify individual reads on-site. Short reads were

sequenced in the laboratory from the same extracted DNA using an Illumina MiSeq. Local

BLAST was used to identify reads from all four datasets (2 field x 2 species) by

comparison to available published reference genomes. Gene models were predicted

directly from individual DNA reads using SNAP (v2006-07-28), matched to existing

phylogenomic datasets and used to infer plant phylogenies using MUSCLE (v3.8.31) and

RAxML (v7.2.8). *de novo* genome assemblies were performed using Abyss (v1.5.2) and

Hybrid-SPAdes (v3.5.1) with completeness assessed with QUAST (v4.0) and CEGMA. R

(v3.1.3) was used to perform statistical analyses. Additional details are given in the

Supplementary Methods.

References:

1.   Hebert, P. D. N., Hollingsworth, P. M., Hajibabaei, M. & Hebert, P. D. N. From

       writing to reading the encyclopedia of life. *Philos. Trans. R. Soc. London B Biol. Sci.*

       371, 1–9 (2016).

2.   Hajibabaei, M., Baird, D. J., Fahner, N. A., Beiko, R. & Golding, G. B. A new way to

       contemplate Darwin's tangled bank: how DNA barcodes are reconnecting

       biodiversity science and biomonitoring. *Philos. Trans. R. Soc. London B Biol. Sci.*

       371, 20150330 (2016).

3.   Quick, J. *et al.* Real-time, portable genome sequencing for Ebola surveillance.

       Nature **530,** 228–32 (2016).

4.   Faria, N. R. *et al.* Mobile real-time surveillance of Zika virus in Brazil. *Genome Med.*

       8, 97 (2016).

5.   Edwards, A., Debbonaire, A. R., Sattler, B., Mur, L. A. & Hodson, A. J. Extreme

       metagenomics using nanopore DNA sequencing: a field report from Svalbard, 78 N.

       bioRxiv 73965 (2016). doi:10.1101/073965

6.   Schmidt, K. *et al.* Identification of bacterial pathogens and antimicrobial resistance

       directly from clinical urines by nanopore-based metagenomic sequencing. *J.*

       Antimicrob. Chemother. dkw397 (2016). doi:10.1093/jac/dkw397

7.   Datema, E. *et al.* The megabase-sized fungal genome of Rhizoctonia solani

assembled from nanopore reads only. *bioRxiv* (2016). doi:10.1101/084772

8.   Novikova, P. Y. *et al.* Sequencing of the genus Arabidopsis identifies a complex

history of nonbifurcating speciation and abundant trans-specific polymorphism. *Nat.*

Genet. **48,** 1077–1082 (2016).

9.   Erlich, Y. A vision for ubiquitous sequencing. *Genome Res.* **25,** 1411–1416 (2015).

10. Mallo, D. & Posada, D. Multilocus inference of species trees and DNA barcoding.

Philos. Trans. R. Soc. London B **371,** 20150335 (2016).

11. CBOL Plant Working Group *et al.* A DNA barcode for land plants. *Proc. Natl. Acad.*

Sci. U. S. A. **106,** 12794–7 (2009).

12. Hollingsworth, P. M., Li, D.-Z., van der Bank, M. & Twyford, A. D. Telling plant

species apart with DNA: from barcodes to genomes. *Philos. Trans. R. Soc. B Biol.*

Sci. **371,** 20150338 (2016).

13. Little, D. P. DNA barcode sequence identification incorporating taxonomic hierarchy

and within taxon variability. *PLoS One* **6,** (2011).

14. Collins, R. A. & Cruickshank, R. H. The seven deadly sins of DNA barcoding. *Mol.*

Ecol. Resour. **13,** 969–975 (2013).

15. Tang, C. Q. *et al.* The widely used small subunit 18S rDNA molecule greatly

underestimates true diversity in biodiversity surveys of the meiofauna. *Proc. Natl.*

Acad. Sci. **109,** 16208–16212 (2012).

16. Zhang, A. B. *et al.* A fuzzy-set-theory-based approach to analyse species

membership in DNA barcoding. *Mol. Ecol.* **21,** 1848–1863 (2012).

17. Laver, T. *et al.* Assessing the performance of the Oxford Nanopore Technologies

MinION. *Biomol. Detect. Quantif.* **3,** 1–8 (2015).

18. Kasianowicz, J. J., Brandin, E., Branton, D. & Deamer, D. W. Characterization of

individual polynucleotide molecules using a membrane channel. *Proc. Natl. Acad.*

Sci. U. S. A. **93,** 13770–3 (1996).

19.   Lamesch, P. *et al.* The Arabidopsis Information Resource (TAIR): improved gene

annotation and new tools. *Nucleic Acids Res.* **40,** D1202–D1210 (2012).

20.   Hu, T. T. *et al.* The Arabidopsis lyrata genome sequence and the basis of rapid

genome size change. *Nat. Genet.* **43,** 476–81 (2011).

21.   Akama, S., Shimizu-Inatsugi, R., Shimizu, K. K. & Sese, J. Genome-wide

quantification of homeolog expression ratio revealed nonstochastic gene regulation

in synthetic allopolyploid Arabidopsis. *Nucleic Acids Res.* **42,** (2014).

22.   Goodwin, S., Gurtowski, J., Ethe-sayers, S., Deshpande, P. & Michael, C. Oxford

Nanopore Sequencing and de novo Assembly of a Eukaryotic Genome. (2015).

23.   Mikheyev, A. S. & Tin, M. M. Y. A first look at the Oxford Nanopore MinION

sequencer. *Mol. Ecol. Resour.* **14,** 1097–1102 (2014).

24.   Jansen, H. J. *et al.* Rapid de novo assembly of the European eel genome from

nanopore sequencing reads. *bioRxiv* (2017). doi:10.1101/101907

25.   Simpson, J. T., Wong, K., Jackman, S. D., Schein, J. E. & Jones, S. J. M. ABySS : A

parallel assembler for short read sequence data ABySS : A parallel assembler for

short read sequence data. 1117–1123 (2009). doi:10.1101/gr.089532.108

26.   Antipov, D., Korobeynikov, A., McLean, J. S. & Pevzner, P. A. HybridSPAdes: An

algorithm for hybrid assembly of short and long reads. *Bioinformatics* **32,** 1009–1015

(2016).

27.   Parra, G., Bradnam, K. & Korf, I. CEGMA: A pipeline to accurately annotate core

genes in eukaryotic genomes. *Bioinformatics* **23,** 1061–1067 (2007).

28.   Wickett, N. J. *et al.* Phylotranscriptomic analysis of the origin and early diversification

of land plants. *Proc. Natl. Acad. Sci.* **111,** E4859–E4868 (2014).

Supplementary Information is linked to the online version of the paper.

Acknowledgements This work was funded by a Pilot Study Grant to JDP and a Howard

Lloyd Davies legacy grant to ASTP. JDP was also supported by funding from the Calleva

Foundation Phylogenomic Research Programme and the Sackler Trust. The authors also

thank The Botanical Society of Britain & Ireland. Natural Resources Wales, Tim Wilkinson,

Robyn Cowan, and Patricia and David Brandwood for assistance.

Author contributions ASTP and JDP conceived the study and obtained funding. ASTP,

DD and JDP designed and conducted fieldwork. ASTP designed and conducted field-

based labwork with input from JDP, AH and DD. AH conducted lab-based sequencing.

JDP conducted bioinformatics and phylogenomic analyses with contributions from AH.

ASTP and JDP prepared the manuscript with contributions from DD and AH.

Author information Basecalled read data for Illumina and Oxford Nanopore sequencing

runs are available via the EBI ENA at XXXXX. The authors declare competing financial

interests: Oxford Nanopore Technologies contributed MinION sequencing reagents and

flowcells for this research. JDP and ASTP received travel remuneration and free tickets to

present an early version of this work at a conference (London Calling 2016).

Correspondence and requests for materials should be addressed to ASTP

(a.papadopulos@kew.org) and JDP (joe.parker@kew.org).

© Crown Copyright and Database Rights 2017 Ordnance Survey

Figure 1 | Logistics and scope of field-based sequencing. a, Location of sample collection and extraction, sequencing and analyses in the Snowdonia National Park, Wales. **b,** *Arabidopsis thaliana*. **c,** *A. lyrata ssp. petraea*. **d,** The portable field laboratory used for the research. Ambient temperatures varied between 7-16ºC with peak humidity >80%. A portable generator was used to supply electrical power.

Figure 2 | Sample identification and phylogenomics using field-sequenced RTnS data. a-d Orthogonal species identification using BLASTN difference statistics: HTS data (red) and RTnS (black) matched to reference databases via BLASTN. **a, c** Receiver operating characteristic (ROC; estimated false-positive rate vs. estimated true positive rate) and **b, d** estimated true- (solid lines) and false-positive (dashed lines) rates. **a, b** ΔL_T statistic; **c, d,** ΔL_I statistic. **e,** Accumulation curves for *ab initio* gene models predicted directly from individual *A. thaliana* reads over time. Count of unique TAIR10 genes (solid

296   line) and total number of gene models (dashed line). Shaded boxes represent periods

297   where the MinION devices were halted while the laboratory was dismantled and moved. **f,**

298   phylogenetic tree inferred under the multispecies coalescent from RTnS reads.

Figure 3 | Simulated accumulation curves for rapid species identification by DNA sequencing. 34k pairwise BLASTN hits of *A. thaliana* RTnS reads were subsampled without replacement to simulate an incremental accumulation of data (10^4 reads; 10^3 replicates). For each read the total identities bias (ΔL_I) is the number of identities with the *A. thaliana* reference minus the number of identities with the *A. lyrata* reference. **a)** the proportion of *A. thaliana* reads correctly identified on a per-read basis, classified as *A. thaliana* where $\Delta L_I >$ threshold cutoff (0, 1, 10 or 100). **b)** Mean ΔL_I in the simulated dataset rapidly stabilises on the population mean (+754bp, e.g. an average matching read alignment to *A. thaliana* is 754bp longer than to *A. lyrata*). **c)** Cumulative aggregate Δ_{LI}; negative or zero ΔL_I can rapidly be excluded. Typical data throughput rates exceed 10^4 reads per hour of sequencing.

Extended Methods

Study site and sample collection. On consecutive days, tissue was collected from three specimens each of *A. thaliana* and *A. lyrata subsp. petraea* in Snowdonia National Park and sequenced and analysed in a tent. *A. lyrata* was collected from the summit of Moelwyn Mawr (52.985168° N, 4.003754° W; OL17 65554500; SH6558244971) and *Arabidopsis thaliana* was collected at Plâs Tan-y-Bwlch (52.945976° N 4.002730° W; OL18 65604060; SH6552940610). Representative voucher specimens of each species are deposited at RBG, Kew. DNA extractions, library preparation and DNA sequencing with the MinION technology were all conducted using portable laboratory equipment in the Croesor valley on the lower slopes of Moelwyn Mawr immediately following sample collection (52.987463°N 4.028517° W; OL17 63904530; SH6392745273). Laboratory reagents were stored in passively-cooled polystyrene boxes with internal temperatures monitored using an Arduino Uno. Only basic laboratory equipment was used (including two MinION sequencers and three laptops; see Extended Data Table 1).

DNA extraction. The standard Qiagen DNeasy plant mini prep kit was used to extract genomic DNA from *Arabidopsis spp.* with the exception that the two batches were pooled at the DNeasy mini spin column step to maximise the DNA yield.

DNA library preparation and sequencing. An R7.3 and R9 1D MinION library preparation were performed for each species according to the manufacturer's instructions using a developer access programme version of the commercially available Nanopore RAD-001 library kit (Oxford Nanopore Technologies). No PCR machine was used. Lambda phage DNA was added to *A. thaliana* R9 library for quality control. For *A. thaliana,* the MinION experiment generated 96,845 1D reads with a total yield of 204.6Mbp over fewer than 16h of sequencing. Data generation was slower for *A. lyrata*, possibly due to

temperature-related reagent degradation or unknown contaminants in the DNA extraction.

Over ~90h sequencing, 25,839 1D reads were generated with a total yield of 62.2Mbp; this

included three days of sequencing at RBG Kew following a 16h drive, during which

reagents and flowcell were stored sub-optimally (near room-temperature). BLASTN 2.4.0[29]

was used to remove 5,130 reads with identity to phage lambda. Data are given in

Extended Data Table 2. The following week in a laboratory, NEBNext Ultra II sequencing

libraries were prepared for four field-extracted samples (two individuals from each species)

and sequenced on an Illumina MiSeq (300bp, paired end). In total, 11.3Gbp and 37.8M

reads were generated (each ~ 8M reads and 2Gbp; see Supplementary Note 1).

Field offline basecalling and bioinformatics in real-time. Offline basecalling using

nanocall 0.6.13[30] was applied to the R7.3 data as no offline R9 basecaller was available at

the time. Basecalled reads were compared to the reference genomes of *A. thaliana*

(TAIR10 release) and *A. lyrata subsp petraea* (1.0 release). In total, 119 reads were

processed in real-time with six reads making significant hits by BLASTN that scored

correctly : incorrectly for species ID in a 2:1 ratio. After the sequencer had been halted a

larger dataset of 1,813 reads gave 281 hits, with correct : incorrect : tied identifications in a

223:30:28 ratio.

Accuracy and mapping rates of short- and long-read data. Both lab-sequenced NGS

reads (trimmed with Trimmomatic[31]) and untrimmed, field-sequenced RTnS reads were

aligned to the appropriate reference genomes using the BWAv0.7.12-r1039[32] and

LASTv581[33] , to estimate depth of coverage and nominal error rate in mapped regions

(see Supplementary Note 2). For all *A. thaliana* datasets (short and long-read), average

mapped read depths were approximately equal to the gross coverage. MinION reads could

be aligned to 53Mbp of the reference genome with LAST (approx. 50% of the total genome

length). The nominal average error rate in these alignments was 20.9%). For both MinION

and MiSeq datasets, mapping and alignment to the *A. lyrata* and *A. lyrata ssp. petraea*

assemblies was more problematic. For alignable MinION reads, error rates were slightly

higher than for *A. thaliana* at 22.5% and 23.5%, estimated against *A. lyrata* and *A. lyrata*

ssp. petraea assemblies, respectively. We note that these assemblies are poorer quality

than the *A. thaliana* TAIR10 release; total genome lengths differ (206Mbp and 202Mbp,)

and contiguity is relatively poor in both (695 and 281,536 scaffolds).

Determination of true- and false-positive detection rates, sensitivity, and specificity.

Each of the four datasets (HTS and RTnS, for each species) was matched against two

custom databases (the *A. thaliana* reference genome and the two draft *A. lyrata* genomes

combined) separately with BLASTN, retaining only the best hit for each query. Queries

matching only a single database were counted as positive matches for that species

(Extended Data Table 4). Non-matching reads were treated as negative results

(Supplementary Methods). Queries matching both databases were defined as positives

based on: a) longest alignment length (L_T); b) highest % sequence identities, c) longest

alignment length counting only identities (L_I), or c) lowest E-value. Test statistics for each

of these metrics were simply calculated as the difference of scores (length (ΔL_T), %

identities, identities (ΔL_I), or E-value) between 'true' and 'false' hits. The statistical

performance of these statistics (true- and false-positive rates, and accuracy) in putative

analyses under varying threshold values were calculated and visualized using the ROCR

package in R[34]. The high proportion of reads with significant hits to both species is

expected given the close evolutionary relationships of the species. Analyses to determine

the best statistics to discriminate between species using reads which aligned to both

databases strongly indicated that difference in alignment lengths between the best

discriminator, shown in Figure 2a-d and Extended Data Figures 2, 3 & 4. Overall these

show that the difference in alignment length is a powerful indicator for both short- and

long-read data at any threshold ≥ ~100bp. Furthermore, and surprisingly, at this and more

conservative (greater difference) threshold, long-read field-sequenced reads had

substantially more accuracy in true- and false-positive discrimination than short-read data.

This suggests that this method provides a powerful means of species identification and we

posit that the extremely long length of 'true positive' alignments compared with the natural

length ceiling on false-positive alignments is largely responsible for this property.

Accumulation curves for simulated identification. 33,806 pairwise BLASTN hits

obtained above in identification against *A. thaliana* and *A.lyrata* genomic reference

databases were subsampled without replacement to simulate incremental accumulation of

BLASTN hit data during progress of a hypothetical sequencing experiment producing

10,000 reads produced in total. 1,000 replicates were used to calculate means and

variances for data accumulation in 0.1 log-increments from r=1 read to 10^4 reads total. For

each read, ΔL_I, 'number of identities bias', was calculated as the difference (number of

identities in *A. thaliana* alignment – number of identities in *A. lyrata* alignment). Each read

was assigned to *A. thaliana* or not if it ΔL_I exceeded a given threshold, repeated at four

possible values, $L_{threshold}$ ={0, 1, 10, 100}. Mean and aggregate (total) ΔL_I values were also

calculated for each replicate over the progress of the simulated data collection. Results are

shown in Figure 3.

***De novo* genome assembly.** Short-read HTS data was assembled *de novo* using ABYSS

v1.9.0[35]. A hybrid assembly with both HTS and RTnS datasets was performed with

HybridSPAdes v3.5.0[36]. Assemblies were completed for *A. thaliana* (sample AT2a) and *A.*

lyrata (sample AL1a). Assembly statistics were calculated in Quast v4.3[37]. Completeness

of the final hybrid assemblies was assessed using CEGMA v2.5[38]. Results of *de novo*

genome assemblies are given in Extended Data Table 5. Analyses of genome contiguity

and correctness and conserved coding loci completeness indicated that assembly of HTS

data performed as expected (20x coverage produced ~25,000 contigs covering

approximately 82% of the reference genome at an N50 of 7,853bp). By contrast, the hybrid

assembly of *A. thaliana* illumina MiSeq and Oxford Nanopore MinION data significantly

improved on the MiSeq-only assembly: 24,999 contigs reduced to 10,644; total assembly

length increased to close to the length of the reference genome (119.0Mbp) with nearly

89% mappable; N50 and longest contig statistics both improved (N50 7,853 → 48,730bp)

indicating better contiguity from the addition of long reads. Completeness of coding loci as

estimated by CEGMA (Extended Data Table 5) greatly increased to ~99%. Long reads did

not compromise the accuracy of high-coverage short-read data; basewise error rates were

not significantly worse.

Direct gene annotation of single unprocessed field-sequenced reads. The length of

typical individual RTnS reads is of similar magnitude to genomic coding sequences.

Consequently, useful phylogenomic information could potentially be obtained by

annotating reads directly, without a computationally expensive genome assembly step.

Raw, unprocessed *A. thaliana* reads were individually annotated directly without assembly

via SNAP[39]. To verify which gene predictions were genuine, the DNA sequences (and 1kb

flanking regions, where available) were matched to available *A. thaliana* (TAIR10) genes

with default parameters. BLAST hits were further pruned based on quality (based on 1st-

quartile quality scores: alignments length bias $\Delta L_T \geq$ +570bp / % identities bias $\geq$ +78.68 /

E-value bias $\geq$ 0), reducing the number of hits from 18,098 to 10,615. Sample read

alignments and details of SNAP output BLAST score summary statistics are given in

Supplementary Table 1 and encounter curves-through-time are shown in Figure 2e.

Phylogenomics of raw-read-annotated *A. thaliana* genes. Predicted *A. thaliana* gene

sequences were combined with a published phylogenomic dataset spanning 852

orthologous, single-copy genes in plants and algae[28], downsampled to 6 representative

taxa for speed: *Equisetum diffusum, Juniperus scopulorum, Oryza sativa, Zea mays, Vitis*

vinifera and *A. thaliana*. Our putative gene models were assigned identity based on

reciprocal best-hit BLASTN matching with the *A. thaliana* sequences in these alignments,

yielding 207 matches, of which the top 56 were used for phylogenomic analysis

(Supplementary Table 1), only 18 having no missing taxa in the Wickett et al.[28] dataset.

Alignments were refined using MUSCLE v3.8.31[40] and trimmed with a 50% missing-data

filter (using trimAL v1.4rev15[41]) then used to infer species trees in two ways: (i) single

gene phylogenies inferred separately (using RAxML v7.2.8[42]) under the GTRCAT

substitution model with 10 discrete starting trees then combined into a summary tree using

TreeAnnotator v.1.7.4[43]; (ii) a species tree inferred directly from the data under the

multispecies coalescent[44], implemented in *BEAST v2.4.4[45] (with adequate MCMC

performance confirmed using Tracer v1.5). A maximum clade credibility (MCC) tree was

produced using TreeAnnotator v.1.7.4. Phylogenies inferred by orthodox (RAxML) and

multispecies coalescent (*BEAST) methods are shown in Extended Data Figure 5 and

agreed with each other and the established phylogeny presented in Wickett et al.[28]

29. C., Coulouris G., Avagyan V., Ma N., Papadopoulos J., Bealer K., & Madden T.L.

(2008) BLAST+: architecture and applications. *BMC Bioinformatics* **10**:421.

30. David, M., Dursi, L.J., Yao, D., Boutros, P.B. & Simpson, J.T. (2017) Nanocall: An

Open Source Basecaller for Oxford Nanopore Sequencing Data. Bioinformatics

33(1):49-55. doi:10.1093/bioinformatics/btw569

31. Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: A flexible trimmer for

Illumina Sequence Data. Bioinformatics 30(15):2114-2120.

32. Li, H. & Durbin, R. (2009) Fast and accurate short read alignment with Burrows–

Wheeler transform. Bioinformatics 25(14):1754–1760.

33. Kiełbasa SM, Wan R, Sato K, Horton P, Frith MC. (2011) Adaptive seeds tame

genomic sequence comparison. Genome Res. 21(3):487-93.

34. Sing, T., Sander, O., Beerenwinkel, N. & Lengauer, T. (2005) ROCR: Visualizing

classifier performance in R. Bioinformatics 21(20):3940-3941.

35.   Simpson, J. T., Wong, K., Jackman, S.D., Schein, J.E., Jones, S.J.M. & Birol, I.

(2009) ABySS: a parallel assembler for short read sequence data. *Genome*

research 19(6):1117-1123.

36.   Antipov D, Korobeynikov A, McLean JS, Pevzner PA. (2016) hybridSPAdes: an

algorithm for hybrid assembly of short and long reads. Bioinformatics 32(7):1009-15.

doi: 10.1093/bioinformatics/btv688.

37.   Gureyvich, A., Saveliev, V., Vyahhi, N. & Tesler, G. (2013) QUAST: quality

assessment tool for genome assemblies. Bioinformatics 29(8):1072-1075.

38.   Parra, G., Bradnam, K. & Korf, I. (2007) CEGMA: a pipeline to accurately annotate

core genes in eukaryotic genomes. Bioinformatics 23(9):1061-1067.

39.   Korf I. (2004) Gene finding in novel Genomes. BMC Bioinformatics 5:59.

40.   Edgar, R.C. (2004) MUSCLE: multiple sequence alignment with high accuracy and

high throughput. Nucleic Acids Res 32(5):1792-97.

41.   Salvador Capella-Gutierrez; Jose M. Silla-Martinez; Toni Gabaldon. (2009) trimAl: a

tool for automated alignment trimming in large-scale phylogenetic analyses. Salvador

Capella-Gutierrez; Jose M. Silla-Martinez; Toni Gabaldon. Bioinformatics 25:1972-

1973.

42.   Stamatakis, A. (2014) RAxML version 8: a tool for phylogenetic analysis and post-

analysis of large phylogenies. *Bioinformatics* **30**(9):1312-1313.

43.   Drummond, A.J., Suchard, M.A., Xie, D. & Rambaut, A. (2012) Bayesian

phylogenetics with BEAUti and the BEAST 1.7 Molecular Biology And

Evolution 29:1969-1973.

44.   Heled, J. and Drummond, A.J. (2010) Bayesian inference of species trees from

multilocus data. Mol. Biol. Evol. 27 (3):570-580.

45.   Bouckaert, R., Heled, J., Kühnert, D., Vaughan, T., Wu, C-H., Xie, D., Suchard, MA.,

Rambaut, A., & Drummond, A. J. (2014). BEAST 2: A software platform for bayesian

evolutionary analysis. PLoS Computational Biology, 10(4):e1003537.

Extended Data

a

b

c

Extended Data Figure 1 | Schematic of experimental workflows. a, Sampling-to-sequencing

workflow. **b,** Sample identification workflow via BLASTN. **c,** Outline for direct annotation of raw

RTnS reads followed by phylogenomic inference. See Methods for details.

Extended Data Figure 2 | Distribution of difference statistics in BLASTN comparisons for

species ID. Empirical distribution of test statistics for each-way congeneric sample ID (binary

classification) using BLASTN evaluated for RTnS MinION (*left column*) and NGS MiSeq (*right*

column) platforms. Difference (test) statistics were calculated for each alignment as (true positive

(TP) score – false positive (FP) score) for each of: alignment length; % identities; and number of

identities. Reads sequenced from *A. thaliana* samples (comprising nominal true positives and

false-positives (contaminants) are shown in red; reads from *A. lyrata* samples (nominal true

negatives) shown in blue. 'True' and 'false' distributions' overlap is small, while alignment length

and number of identities' distributions are both unimodal, showing a simple cutoff-based classifier

should perform well to discriminate between 'true' and 'false' cases.

Extended Data Figure 3 | Performance of difference statistics in BLASTN comparisons for
species ID. Performance of test statistics for each-way congeneric sample ID (binary
classification) using BLASTN evaluated for MinION (*black*) and MiSeq (*red*) platforms. Difference
(test) statistics were calculated amongst reads matching both databases for each alignment as
(true positive (TP) score – false positive (FP) score) for each of e-value, length, % identities and
number of identities. Reads matching only one or neither database were additionally included with
either 'all-false' encoding (dashed green line) or mixed false and true encoding depending on
sample origin (blue line; see Methods and Supplementary Information for details). RTnS reads'
true- and false-positive rates are comparable to, and in some cases better than, NGS reads'
performance; while the longer length of RTnS reads permits the use of high thresholds where
greater confidence is desired (perhaps in the case of very closely related specimens). *Top row:*
true-positive (TP) vs false-positive (FP) rate; classical receiver operating curve. *Second row:* FP
rate with varying test statistic threshold. *Third row:* TP rate with varying test statistic threshold.
Bottom row: Accuracy with varying test statistic threshold. Accuracy estimated as (TP+TN / (P +
N)). *Columns (L-R):* Difference statistics for *e*-value, total alignment length, % identities, and
number of identities, respectively.

Extended Data Figure 4 | Modelling potential effect of incorrectly estimated TN/FN proportions. Red and black lines show empirically estimated statistical performance for species ID via BLASTN comparison of HTS and RTnS reads respectively (as for Extended Data Figure 3; see Methods for details). Reads that produced no hits to either database might represent false negatives (sequencing error, or genomic regions not represented in the reference genome BLAST databases) or true negatives (sequencing contaminants and sequencer noise). These nonmatching reads were to reflect 'true negative':'false negative' ('TN:FN') mixtures in 10%

increments shown from light to dark green shading. Plots (**a-d**) show results for bias statistics in *E*-

value; total alignment length; % identities; and total alignment identities, respectively. Extreme

TN:FN mixtures still display adequate true-positive vs. false-positive rates; empirical data is

approximated by the 30% TN:FN mixture, approximately reflecting the proportions of *A. lyrata* to *A.*

thaliana nonmatching reads in the dataset.

a

b

Extended Data Figure 5 | Phylogeny of species spanning major plant groups. Putative coding

sequences recovered from single, unassembled raw RTnS reads using SNAP *ab initio* gene

prediction could be aligned to existing phylogenomic data from other taxa and used to infer a

phylogeny consistent with accepted plant relationships. **a,** Multispecies coalescent species tree

inferred from 18 gene trees (genes predicted directly from raw nanopore reads). Inferred using

multispecies coalescent implemented in BEAST 2.4.4; **b,** consensus species tree inferred by

majority-rule from 18 gene trees, inferred with RAxML 7.2.8.

Extended Data Table 1 | List of field-sequencing equipment.

Item / model	Quantity	Supplier
Laptops, Portege R830-1DZ	2	Toshiba
Laptop, MacBook Pro	1	Apple
Portable firewire HDD, 1Tb	2	LaCie
Portable AC generator, IMPAX IM800I 700W	1	ScrewFix
Thermal control textiles (socks) for MinION	Pair	ScrewFix
AC extension plugs, 4-way, 3-20M	3	Argos
Portable folding tables	2	Argos
Weather meter, Kestrel 5500	1	KestrelMeters
Fluorometer, Quantus	1	Promega
Water bath, GD100	1	Grant
Microcentrifuge, 5415C	1	Eppendorf
Glass thermometers, 300mm, range 263.3-383.3°K	2	
Arduino Uno	1	Maplins
Themal transducers, LM335AZ	4	RS Components
Polystyrene thermal control boxes, various sizes	3	
Freezer coolpacks	14	
Pipettes, Gilson:	-	
P2	1	ThermoFischer
P20	1	ThermoFischer
P200	1	ThermoFischer
P1000	1	ThermoFischer
Waste containers	2	
Laboratory consumables:	-	
PCR tubes, thin-walled	Box	ThermoFischer
Eppendorf tubes, 1.5ml DNA lo-bind	Box	ThermoFischer
Eppendorf tubes, 2.0ml DNA lo-bind	Box	ThermoFischer
Plastic pestles	20	
Sterilized sand	5g	
Reagent tubes, screw-top, 50ml	10	ThermoFischer

Extended Data Table 2 | Performance of MinION sequencing runs

Species	*A. thaliana*	*A. thaliana*	*A. lyrata*	*A. lyrata*
Reaction chemistry	R7.3, 1D	R9, 1D	R7.3, 1D	R9, 1D
Run IDs	2507 2126 3637	5913 2144 0509	4901 1842 1201	1222 5458 1958 0824 2912 5201
Start time	18/05/2016 21:36	18/05/2016 18:21	19/05/2016 16:10	19/05/2016 16:39:20
Latest read*	19/05/2016 15:17:00	19/05/2016 15:18:39	24/05/2016 01:14	22/05/2016 07:08:56
Disc space (raw)	2.9Gb	106.8Gb	3.0Gb	35.1Gb
Metrichor ID	115360	115459	115375	115432
# reads	4,152	92,693	2,387	23,452
Yield (bp):				
total	7,351,585	233,244,147	16,092,487	46,118,754
mean	1,771	2,516	6,742	1,967
median	586	1,396	120	305
maximum	434,377	170,598	1,114,970	177,310
N25	19,244	9,651	574,112	24,254
N50	4,771	4,410	309,034	7,926
N75	2,041	2,121	62,360	3,374

Data produced with field-extracted and field-sequenced in conditions ranging from 6-14°C and up

to 100% humidity, producing genome-scale sequence data despite several pauses in sequencing

to dismantle, relocate, and reassemble equipment. The two latest sequencing runs (*A. lyrata*

samples) performed markedly worse than the two earliest runs (*A. thaliana)*, possibly due to the

impact of storage temperature (passively controlled, and steadily rising over the week) on reagent

performance. Yield summary statistics refer to untrimmed raw reads, including phage-lambda

experimental control in the case of *A. thaliana* R9 data (filtered from subsequent steps). See

Methods for details. Note: *Final *A. lyrata* sequencing phase performed in laboratory owing to time

constraints on-site.

Extended Data Table 3 | Statistics of reference genomes used

Species	Arabidopsis lyrata ssp. petraea	A. lyrata	A. thaliana
Version	1.0	1.0	TAIR10
Date accessed	25/05/2016	17/05/2016	17/05/2016
Accession / ID	NCBI: assembly= GCA_000524985.1	NCBI: genome=493 genome_assembly=29434	NCBI: genome=4 genome_assembly=22492
Total assembly length	202,972,003	206,667,935	119,667,750
Total gap length	20,456,163	22,960,134	185,644
Number of scaffolds	281,536	695	7
Scaffold N50	7,848	24,464,547	23,459,830
Scaffold L50	6,426	4	3
Number of contigs	369,168	3,645	102
Contig N50	2,321	227,391	11,194,537
Contig L50	16,831	247	5
Total chromosomes & plasmids	0	0	7

585   *A. thaliana* TAIR10 release is considerably more complete than either of the draft *A. lyrata*

586   assemblies.

590   **Extended Data Table 4 | Sample identification via BLASTN**

Sample	A. thaliana	A.lyrata	A. thaliana	A.lyrata
"TRUE" database	A. thaliana	A.lyrata combined*	A. thaliana	A.lyrata combined*
"FALSE" database	A.lyrata combined*	A. thaliana	A.lyrata combined*	A. thaliana
Source data	ONT 1D	ONT 1D	MiSeq	MiSeq
# Reads, total	91,715	25,839	9,476,598	9,659,489
# Reads with BLASTN hits:				
1-way TRUE[†]	10,322	76	2,140,403	2,907,921
1-way FALSE[‡]	378	2	53,056	24,329
2-way BOTH[§]	22,386	101	7,098,032	6,256,969
0-way ZERO[¶]	58,629	25,660	185,107	470,270
Proportion of reads:				
1-way TRUE	0.113	0.003	0.226	0.301
1-way FALSE	0.004	0.000	0.006	0.003
2-way BOTH	0.244	0.004	0.749	0.648
0-way ZERO	0.639	0.993	0.020	0.049
Biases [#]:				
Mean length	1,323.87	698.25	83.61	108.99
Mean identities	1,115	575	96	117
Mean % identities	37.97	61.41	30.16	42.38
Mean E-values	4.80E-07	1.05E-04	1.07E-08	3.52E-09

Individual RTnS and NGS reads were aligned to *A. thaliana* and *A. lyrata* databases (designated

TRUE or FALSE depending on sample origin) with BLASTN, keeping the single best-hit alignment

for each database. More '1-way' (only one database matched) hits to TRUE than FALSE

databases accumulated in all sample / technology combinations. Amongst '2-way' hits, positive

differences in the metrics were consistent with correct sample identification. For RTnS reads

differences between TRUE and FALSE hits were considerably larger than amongst HTS reads (by

an order of magnitude for length or number of identities), showing that confident identification could

be made with fewer RTnS reads. Notes: *A. lyrata* and *A. lyrata ssp. petraea* databases combined,

see Methods; †Total number of reads matching only conspecific database ('true-positives'); ‡Total

number of reads matching only pairwise-compared database ('false-positives' in the case of a

mixed /multiplexed sample, or 'false-negatives' in the case of a single sample); §Total number of

reads matching both databases; ¶Total number of reads with no hits in either comparison, e.g.

'false-negatives'; #Difference statistics for each query read calculated as (score conspecific

comparison – score congener comparison), for BLASTN alignment length, alignment identities,

alignment % identities and *E*-value; ☆Mean bias across all reads.

Extended Data Table 5 | Performance of *de novo* genome assembly

Species	*Arabidopsis thaliana*		*A. lyrata ssp. petraea*	
Data	MiSeq	MiSeq + MinION	MiSeq	MiSeq + MinION
Assembler *	Abyss	hybridSPAdes	Abyss	hybridSPAdes
Illumina MiSeq NGS reads, 300bp paired-end	8,033,488	8,033,488	8,143,010	8,143,010
NGS total yield	2,418,079,888	2,418,079,888	2,451,046,010	2,451,046,010
Oxford Nanopore MinION RTnS reads, R7.3 + R9, N50 ~ 4,410bp	n/a	96,845	n/a	25,839
RTnS reads total yield	n/a	240,597,532	n/a	62,211,241
# contigs	24,999	10,644	37,568	85,599
Largest contig	89,717	413,462	101,114	38,313
Total length	106,455,313	119,031,857	151,562,895	117,256,694
Reference length	*119,667,750* [†]	*119,667,750*	*183,707,801* [‡]	*183,707,801*
GC content (%)	35.97	36.20	36.16	36.55
N50 [§]	7,853	48,730	9,605	1,686
Unaligned length	7,121,882	6,737,059	36,669,847	35,287,390
Genome fraction (%)	82.0	88.7	53.4	43.7
Duplication ratio	1.01	1.058	1.17	1.02
# N's per 100 kbp	1.72	5.41	0.22	7.09
# mismatches / 100 kbp	518	588	1,297	1,097
# indels / 100 kbp	120	130	334	271
Largest alignment	76,935	264,039	44,515	17,201
Total aligned length	98,382,255	108,086,256	100,502,092	80,814,492
Coding loci completeness[¶]:				
# genes, 'complete'	**219**	**245**	n/a	n/a
% genes, 'complete'	88.31%	98.79%		
# genes 'partial'	**238**	**246**	n/a	n/a
% genes, 'partial'	95.97%	99.19%		

Field-extracted DNA material was of sufficient quality to enable a *de novo* assembly with lab-

sequenced NGS data. Furthermore, field-sequenced RTnS reads considerably augmented the

NGS data in hybrid assembly, greatly improving contiguity and estimated coding loci coverage

substantially without substantially raising basewise error rates. Notes: **de novo* genome

assemblies used either lab-sequenced short-read HTS data only (Abyss) or both HTS and field-

sequenced RTnS datasets (Hybrid-SPAdes). †TAIR10 release. ‡INSDC: *A. lyrata*:

ADBK00000000.1 (Hu *et al.,* 2011); *A. lyrata ssp. petraea*: BASP00000000.1 (Akama *et al.*, 2014).

§Assembly statistics calculated using QUAST 4.0. ¶Approximate completeness of coding loci

assessed via CEGMA. See Methods for details.

621