1   **Title:** Pattern separation of spiketrains in hippocampal neurons

2   **Short Title**: Temporal pattern separation in the hippocampus

3   **Authors:** Antoine D. Madar*[1], Laura A. Ewell*[2], Mathew V. Jones[1]

4   * These authors contributed equally to the manuscript.

Authors affiliations

6   [1]Department of Neuroscience, University of Wisconsin-Madison, Madison, WI 53705, USA

7   [2]Institute of Experimental Epileptology and Cognition Research, University of Bonn, Bonn,

8   Germany

Corresponding authors

10 Antoine D. Madar (madar@wisc.edu)

Author contributions

12 Conceptualization: MVJ, LAE. Data curation: LAE, ADM. Formal analysis: MVJ, ADM.

13 Funding acquisition: MVJ, ADM. Investigation/data collection: LAE, ADM. Methodology:

14 MVJ, LAE, ADM. Project administration: MVJ, LAE, ADM. Resources: N/A. Software: MVJ,

15 ADM. Supervision: MVJ. Validation: MVJ, LAE, ADM. Visualization: MVJ, LAE, ADM.

16 Writing – original draft: ADM. Writing – review & editing: MVJ, LAE, ADM.

Funding

18 This work was supported by the University of Wisconsin Institute for Clinical and Translational

19 Research (M.V.J.; NIH/NCATS UL1TR000427) and Lily's Fund for Epilepsy Research

20 (A.D.M.; 2015 fellow).

22 The authors declare no competing interests in relation to the present work.

Abstract

Pattern separation is a process that minimizes overlap between patterns of neuronal activity representing similar experiences. Theoretical work suggests that the dentate gyrus (DG) performs this role for memory processing but a direct demonstration is lacking. One limitation is the difficulty to measure DG inputs and outputs simultaneously. To rigorously assess pattern separation by DG circuitry, we used mouse brain slices to stimulate DG afferents and simultaneously record DG granule cells (GCs) and interneurons. Output spiketrains of GCs are more dissimilar than their input spiketrains, demonstrating for the first time temporal pattern separation at the level of single neurons in the DG. Pattern separation is larger in GCs than in fast-spiking interneurons and hilar mossy cells, and is amplified in CA3 pyramidal cells. Analysis of the neural noise and computational modelling suggest that this form of pattern separation is not explained by simple randomness and arises from specific presynaptic dynamics.

Introduction

How does the brain allow us to discriminate between similar events in our past? This question is a central challenge in the neurobiology of memory and remains elusive. To prevent confusion between memories that share similar features, the brain needs to store distinct activity patterns to represent distinct memories. In the influential Hebb-Marr framework of episodic memory (McNaughton & Morris, 1987; Treves et al., 2008), representations are stored in area CA3 of hippocampus, an auto-associative network where plastic recurrent excitatory connections facilitate recall of stored patterns in response to partial cues (McNaughton & Morris, 1987;

Rolls, 2010). However, strong recurrent excitation severely limits the number of patterns that can be stored without overlap (O'Reilly & McClelland, 1994; Rolls, 2010). Such overlap would lead, when a partial cue common to several patterns is presented, to the reactivation of many patterns and thus to confusion or confabulation. To avoid these interferences, the Hebb-Marr framework proposes that redundancy between input patterns is reduced before they are stored. This process of transforming similar input patterns into less similar output patterns is termed *pattern separation* (O'Reilly & McClelland, 1994; Santoro, 2013).

Theoretical models suggest that the dentate gyrus (DG) performs pattern separation of cortical inputs before sending its differentiated outputs to CA3 (Rolls, 2010; Treves et al., 2008). Indeed, DG is ideally located to do this, receiving signals via the major projection from entorhinal cortex (EC), the perforant path (PP), and sending signals to CA3 via the axons of granule cells (GCs) (Amaral et al., 2007). In addition, behavioral studies have shown that DG lesions impair mnemonic discrimination (Baker et al., 2016; Dillon et al., 2017; Kesner et al., 2016; Kesner & Rolls, 2015; McHugh et al., 2007). However, although experimental reports have concluded that pattern separation is performed by the DG (Deng et al., 2013; Leutgeb et al., 2007; Marrone et al., 2011; Nakazawa, 2017; Neunuebel & Knierim, 2014), they only directly show that similar environments or events are represented differently in the DG. The separation could be done by upstream structures and simply be reported by the DG. Hence, it is still unknown whether DG itself performs pattern separation. A rigorous demonstration would require simultaneous knowledge of the inputs arriving at the DG and the processed outputs from DG to CA3 (Santoro, 2013).

Another difficulty in studying pattern separation is in defining the nature of "activity patterns". Previous studies have focused on spatial patterns of "active neurons", with little

reference to the dynamics of neural activity. For example, computational models predict that DG

separates overlapping populations of active EC neurons into less overlapping populations of

active GCs (Chavlis et al., 2017; Myers & Scharfman, 2009; Neher et al., 2015; O'Reilly &

McClelland, 1994; Yim et al., 2014). Immediate-early genes (IEG) expression studies have

confirmed that distinct events drive plasticity in different populations of GCs (Deng et al., 2013;

Schmidt et al., 2012) and that overlap in these representations causes mnemonic interference

(Ramirez et al., 2013). In contrast, *in vivo* single-unit recordings in the DG found that similar

contexts are represented by the same population of active neurons, but differences are encoded

by different spatially tuned firing patterns (Leutgeb et al., 2007; Nakazawa, 2017; Neunuebel &

Knierim, 2014).

These conflicting results show that pattern separation could correspond to different

computations depending on the type of patterns investigated, and that multiple forms of pattern

separation could in theory be implemented by the DG (Santoro, 2013). For example, because *in

vivo* recordings suggest that the same neurons can be used to code different environments

(Leutgeb et al., 2007; Nakazawa, 2017; Neunuebel & Knierim, 2014), it is possible that pattern

separation is performed at the level of single GCs, each disambiguating the activity patterns that

it receives. Such disambiguation could be done by changing firing rates, or alternatively, by

changing spike timing. Previous experimental investigations of pattern separation in DG

examined population vectors of place fields averaged over minutes (Leutgeb et al., 2007;

Neunuebel & Knierim, 2014; Senzai & Buzsáki, 2017), but place cells also carry information at

shorter timescales (Buzsaki, 2010; Eichenbaum, 2014; Kelemen & Fenton, 2016). So far, pattern

separation has not been well characterized on the scale of milliseconds, and never where patterns

are explicitly afferent and efferent trains of action potentials.

Thus, whether the DG network per se reduces the overlap between similar inputs, and

how it performs this computation remains a mystery, especially at short timescales. Here, we set

out to test the hypothesis that the DG performs pattern separation of cortical spiketrains at the

level of single GCs. We designed a novel pattern separation assay in acute brain slices to take

advantage of the experimental control afforded to slice electrophysiology. Complex input

spiketrains of varying similarities were fed into the DG via its afferents, and the output of a

single GC was simultaneously recorded, allowing the first direct measure of pattern separation

(by comparing input similarity versus output similarity), on timescales relevant to neuronal

encoding and synaptic plasticity. Finally, we explored whether other cell types in the DG and

CA3 exhibited this form of pattern separation and investigated the role of neuronal noise and

synaptic dynamics in supporting this computation.

Results

Temporal pattern separation by individual dentate granule cells

A direct test of pattern separation in single GCs requires knowledge of the similarity between

input patterns arriving via the PP and comparison with the similarity between GC output

patterns. Here, we define input and output patterns as rasters of spiketrains. The similarity

between two spiketrains was assessed by computing their pairwise Pearson's correlation

coefficient (R) using a binning window τ_w of 10 ms (unless otherwise specified). We generated

sets of Poisson input spiketrains (simulating trains of incoming cortical action potentials), with

each set having an average correlation R_{input} (**Figure 1A** and **Methods – Pattern separation**

experiments). We then recorded the spiking responses of GCs to these sets of input trains

delivered to lateral PP fibers (**Figure 1B-C**) (102 recording sets from 28 GCs in adolescent

mice), allowing us to compute the average output correlation (R_{output}) (**Figure 2A-B**).

For every recording set, R_{output} was lower than the R_{input} of the associated input set,

indicating a decorrelation of the output spiketrains compared to their inputs (**Figure 2C**). This

was also the case in GCs recorded in slices from adult mice (35 recording sets from 14 neurons)

(**Figure 9D**). These results are the first direct experimental evidence that single GCs, the output

neurons of DG, exhibit pattern separation. The effective decorrelation, defined as the difference

between R_{input} and R_{output}, was statistically significant for every input set (**Figure 2D left**), but

was larger when input spiketrains were highly correlated (**Figure 2D**). This is consistent with the

role of DG in discriminating between similar memories more than already dissimilar ones

(Kesner & Rolls, 2015). Note, however, that the decorrelation normalized to R_{input} is invariant:

whatever the input set, the output trains were always decorrelated to ~70% of R_{input} (**Figure 2E**).

Such invariance suggests that the same decorrelating mechanism is used on all input sets.

These results constitute the first demonstration that input spiketrains are decorrelated in

the DG at the level of single GCs.

Temporal pattern separation in other cell types of the DG network

Any channel processing inputs and returning outputs, and thus any brain network, performs

either pattern separation or pattern convergence to some degree (Santoro, 2013). Thus, GCs are

unlikely to be the only neurons to exhibit temporal pattern separation of spiketrains. However,

we would expect pattern separation to be at its greatest in GCs, at least among DG cells, because

they are the output neurons of the DG and thus should provide the most separated patterns to

CA3 before they are stored. To test this hypothesis, we performed the same pattern separation

assay while recording from fast-spiking interneurons (FS, 20 recording sets) (**Figure 3A**) or hilar

mossy cells (HMC, 18 recording sets) (**Figure 3B**).

At the 10 ms timescale, the distributions of average (R_{input}, R_{output}) were significantly

different between FSs and GCs, with the R_{output} of simultaneously recorded GCs always lower

than their corresponding FS (**Figure 3A4**). This indicates that FSs perform lower levels of

decorrelation than GCs (**Figure 3A5**). For HMCs, although their current-clamp responses and

spiking behavior looked different from GCs (**Figure 3B2**), the distribution of average (R_{input},

R_{output}) appeared only slightly different from GCs with this analysis (**Figure 3B3**, but see below).

Temporal pattern separation in CA3 pyramidal cells

Our FS recordings have shown that not all neurons necessarily exhibit as high levels of temporal

pattern separation as GCs. We thus asked whether this form of pattern separation is specific to

the DG output, or whether neurons in other hippocampal regions can perform temporal pattern

separation. To answer this question, we adapted our pattern separation assay to CA3 pyramidal

cells (PCs) (**Methods – Pattern separation experiments**) and recorded their output spiketrains

in response to direct stimulation of the GCL, i.e. in response to its input from the DG (**Figure

4A**). Due to strong feedforward inhibition, CA3 PCs do not generally spike in response to

external stimulation of afferents in slices, unless inhibition is totally blocked (Rebola et al., 2017;

Sachidhanandam et al., 2009; Sun et al., 2017). By using 30 Hz Poisson input sets (**Figure 4B**)

and adding 100 nM of gabazine to the bath, which only slightly decreases the amplitude of

GABA-A-mediated IPSCs (see **Methods**), we managed to record for the first time the spiking

output of CA3 PCs in response to complex input spiketrains while preserving some inhibition in

the network (**Figure 4C**). PCs fired during periods of high input frequency, probably due to

depression of local inhibitory transmission (Mori et al., 2004; Rebola et al., 2017) combined with

facilitation at the mossy fiber-CA3 PCs synapses (Vyleta et al., 2016), and as a result action

potentials often appeared more clustered than in control GC recordings (**Figure 4C**). Despite this

spiking pattern, we found that, at the 10 ms timescale, CA3 PCs average (R_{input}, R_{output})

distribution was slightly but significantly lower than for GCs recorded in similar

pharmacological conditions (**Figure 4D**). This indicates that CA3 PCs are different from GCs in

respect to temporal pattern separation, but that, surprisingly, their output spiketrains are even

more decorrelated. Temporal pattern separation is thus not specific to the DG.

Temporal pattern separation at different timescales across celltypes

Because the timescales meaningful for the brain remain uncertain, it is important to assess the separation of spiketrains at different timescales. We therefore binned spiketrains using a range of τ_w from 5 ms to 250 ms and performed a finer grained analysis using pairwise R_{input} and associated pairwise R_{output} instead of the average across the ten pairs of input trains (**Figure 5A**). We discovered that pattern separation levels can dramatically change as a function of τ_w. In GCs, the larger the timescale the less they exhibit decorrelation of their input spiketrains, especially at high R_{input} (**Figure 5B**). Nonetheless, GCs still exhibit relatively high levels of pattern separation of highly similar input spiketrains, even at long timescales (**Figure 5B**).

This analysis confirmed that, at short timescales, FSs exhibit less pattern separation than GCs (**Figure 5C**) and revealed significant differences between HMCs and GCs, especially for pairs of input spiketrains with low R_{input} (**Figure 5D**), which were not as obvious in our previous coarse analysis (**Figure 3B3**). At longer timescales, the variability across neurons has a tendency to increase for all celltypes, but the average levels of decorrelation of both FSs and HMCs stayed highly significantly different from those of GCs (**Figure 5C-D**). Interestingly, the difference between FSs and GCs was larger at short timescales, whereas the difference between HMCs and GCs increased with larger timescales. Indeed, at τ_w above 100 ms, HMCs can often exhibit pattern convergence instead of pattern separation, especially for pairs of already dissimilar input spiketrains (low or negative pairwise R_{input}), whereas FSs just show weak or no pattern separation.

Concerning CA3 PCs, they still exhibit high levels of temporal pattern separation at long timescales and the difference with GCs increases (**Figure 5E**). Interestingly, in contrast to all

other tested celltypes and conditions (**Figure 5B-D**), PCs even show a dramatic increase of their

average levels of decorrelation at the 250 ms timescale (**Figure 5E**).

Overall, these results show that among the tested DG celltypes, GCs exhibit the highest

levels of temporal pattern separation of cortical spiketrains across all timescales. Our findings

also suggest that the high level of separation by the DG is amplified in CA3.

Mechanism of temporal pattern separation: neural noise

To determine what mechanisms might support temporal pattern separation in GCs, it is necessary

to understand its dynamics first. Limiting our analysis to the first presentation of an input set

revealed that outputs were already significantly decorrelated (**Figure 6A-B**). This shows that the

separation mechanism is fast, consistent with the fact that the brain generally does not have the

opportunity to average repeated signals and that separation must happen immediately during

encoding. In addition, analysis of the last presentation revealed only modestly more separation

than for the first one, and only for high input correlations (**Figure 6C**), suggesting that learning

to recognize the input pattern is not critical.

Because the mechanism for temporal pattern separation is fast and does not require

learning, we asked first whether intrinsic properties of GCs could play a role. Linear regression

analysis revealed that the membrane capacitance, resistance, time constant as well as the resting

membrane potential are not predictors of decorrelation in GCs (see low R^2 in **Table 1**). Another

hypothesis is that randomness in neuronal responses drives the decorrelation. Indeed, when the

same input spiketrain is repeated (e.g. $R_{input} = 1$) the output spiketrains are not well correlated

(as shown by the mean spiketrain reliability R_w) (**Figure 6A**), consistent with well-known trial-

to-trial variability in single neuron responses (Dobrunz & Stevens, 1999; Faisal et al., 2008;

Rieke et al., 1999). Theoretical investigation of pattern separation often relies on some sort of random process such as probabilistic neuronal activation (O'Reilly & McClelland, 1994) or stochastic firing (Aimone et al., 2009), which suggests that "neural noise" is a likely contributor to any form of pattern separation. However, because "neural noise" can cover multiple different definitions and phenomena (Faisal et al., 2008), determining its role in a complex computation is not trivial.

The noisiness in neural communication is often understood as the unreliability of spiking after a single input spike, and the jitter of the delay between an input spike and an output spike (Mongiat et al., 2009; Savanthrapadian et al., 2014). We thus assessed such spike-wise noise directly from recording sets from our pattern separation experiments (**Figure 7B**, **S1**, and **Materials and methods – Analysis of output spiketrains**). Although the spike probability (SP), delay and jitter of GCs were slightly higher in our experiments than in a previous report (but recording and analysis methods were different), the variability between GCs was consistent (Mongiat et al., 2009). We then asked whether these spike-wise noise parameters could predict the degree of decorrelation by GCs. First, linear regression analysis shows no good relationship with any parameter, the SP being a mediocre predictor at best (**Figure S2A** and **Table 2**). Moreover, the average firing rate of a GC output set (a measure dependent on SP) is not well correlated with the degree of decorrelation either (**Table 3** and **Figure S7A**). Temporal pattern separation in GCs seems to not be achieved merely because their output spiketrains are a sparser and jittered version of their inputs.

To more carefully test the hypothesis that random spiking failures and random delays support fast temporal pattern separation, we produced a shuffled dataset and a simulated dataset only governed by spike-wise noise statistics comparable to the original data (**Figure S1, S2B,**

Table 2 and **Methods – Computational models**). The distribution of (R_{input}, R_{output}) was

significantly higher in the original data (**Figure 7** and **S4**), showing that simple random

processes yield greater levels of separation than real GCs, especially for highly similar inputs

(**Figure 7E** and **S4A**).

In addition to the spike-wise noise, we considered neural noise at the level of spiketrains

by computing the average correlation coefficient R_w between "children" output spiketrains from

the same "parent" input train (**Figure 7A**). R_w characterizes the more complex notion of

spiketrain reliability, that is the ability of a neuron to reproduce the same output spiketrain in

response to repetitions of the same input spiketrain. R_w is not dependent on intrinsic cellular

properties (**Table 1**) and only moderately determined by spike-wise noise parameters (**Table 2**),

suggesting that the rather low R_w of GCs is the expression of more complex noisy biophysical

processes. Consistently, R_w was significantly lower for shuffled and simulated data than in real

GCs (**Figure 7D-E** and **S4B**). This indicates that the output spiketrains of GCs are more reliable

than if their output was entirely determined by simple random processes. Overall, the lower

R_{output} and R_w distributions of random datasets compared to GCs (**Figure 7**) clearly show that

simple noise cannot fully underlie the operations performed by GCs on input spiketrains.

The analysis above also suggests that there might be an unavoidable trade-off between

achieving pattern separation and reliable information transmission about input spiketrains. To

further investigate this, we looked at the relationship between the spiketrain reliability R_w and the

decorrelation levels in GC recordings and found a strong anticorrelation (**Figure 8A** and **Table**

3). This linear relationship is clear evidence that a biological process leading to sweep-to-sweep

variability is a powerful mechanism for temporal pattern separation in DG.

However, spiketrain unreliability is not the only reason GCs output spiketrains are

decorrelated. Indeed, three lines of evidence supports the idea that even if GCs were perfectly

reliable from sweep to sweep, their output spiketrains would still be less similar than their inputs.

First, when averaging out the variability between spiketrains associated to the same input (i.e. by

computing the PSTH), a significant level of decorrelation is still detected (**Figure 8B-D**).

Second, the linear model describing the relationship between R_w and decorrelation levels suggest

that even at $R_w = 1$, ~10% of decorrelation would still be achieved (**Figure 8A**). Finally, high

levels of pattern separation are performed upon the first presentation of an input set (**Figure 6**).

We can thus conclude that temporal pattern separation is supported by mechanisms in addition to

spiketrain unreliability.

Taken together, these results suggest that complex but noisy biophysical mechanisms

allow GCs to balance temporal pattern separation and reliable signaling about their inputs.

Mechanism of temporal pattern separation: short-term synaptic dynamics

The fact that intrinsic properties of GCs do not predict their decorrelation levels (**Table**

1) suggests that temporal pattern separation comes from the DG network in which each recorded

GC is embedded. We thus hypothesize that the specific short-term dynamics of synapses in the

DG network implement temporal pattern separation in GCs. This is a likely mechanism because:

1) Synaptic transmission is a stochastic process and generally considered the main source of

noisiness in the spiking output of neurons (Faisal et al., 2008; Mainen & Sejnowski, 1995),

which would fit our conclusion that temporal pattern separation is dependent on neural noise (see

paragraph above), 2) Short-term plasticity makes the probability of synaptic release dependent

on the timing of the preceding input spikes, which could explain why temporal pattern separation

and spiketrain reliability are not purely governed by noise. To test the idea that short-term

synaptic dynamics is a potential mechanism controlling the pattern separation/spiketrain

reliability ratio in GCs, we developed a computational model of a spiking GC with dynamic and

probabilistic PP-GC synapses (see **Methods – Computational models**). The parameters of the

model controlling the facilitation, depression and variability of presynaptic release, and thus the

amplitude of EPSCs , were constrained to match the whole-cell voltage-clamp recording of a GC

in response to a pattern separation protocol (10 Hz input trains, $R_{input} = 0.76$) (**Figure 9A-C** and

see **Methods – Computational models**). Simulations of pattern separation experiments with this

model resulted in high levels of output spiketrains decorrelation, akin to real GCs, but also in

relatively high levels of spiketrain reliability (**Figure 9D-E**). This is a proof of principle that the

probabilistic and dynamic nature of presynaptic release at the PP-GC synapse is a potential

mechanism balancing temporal pattern separation and information transmission in the DG.

Sources of pattern separation differences between celltypes

To further understand the mechanisms behind temporal pattern separation in single hippocampal

cells, we investigated the sources of the difference in decorrelation levels between the tested

celltypes. We considered differences in intrinsic cellular properties (**Table 1**), firing rate and

probability of bursting (**Figure S6-7**), spike-wise noise (**Figure S2-3**) and spiketrain reliability

(**Figure 8A** and **S5**).

Both FS and HMC recordings displayed bursts of spikes (defined as more than one output

spike between two input spikes), which was very rarely seen in our GC and CA3 recordings

(**Figure 3-4, S6C-E,** and **S7C, E**). As a result, both FSs and HMCs had significantly higher

firing rates than GCs (**Figure S6A-B**) (although the effect size was smaller for HMCs, because

they burst less often than FSs and they have generally less spikes per bursts. **Figure S6D-E**).

Then, are the firing rate or the probability of bursting predictive of differences in pattern

separation? The relationships are unclear, but it seems that both could partially explain the lower

decorrelations observed in some HMCs and FSs (**Figure S7A-B**). To more directly test the role

of bursting, we processed all FS and HMC recordings by removing all but the first spike in each

detected burst (**Figure S7C, E** and see **Methods – Analysis of the output spiketrains**). These

resulting "non-bursty" datasets (nbFS and nbHMC) still exhibited significantly less pattern

separation than GCs (and even pattern convergence of dissimilar input trains, in the case of

HMCs) (**Figure S7D, F**). Therefore, bursting and high firing rates, although a source of

differences in temporal pattern separation, are not a sufficient explanation.

Although FSs show less pattern separation than GCs, it is interesting that they do exhibit

some amount of separation, as opposed to pattern convergence (Santoro, 2013) which one could

have expected from their reputation of having a much more reliable and precise spiking behavior

than principal neurons (Hu et al., 2014; Savanthrapadian et al., 2014). The high fidelity in

relaying input spikes (Savanthrapadian et al., 2014) might still explain the difference in pattern

separation ability between FSs and GCs, although, to our knowledge, they have never been

formally compared. We thus first confirmed the idea that FS show much less spike-wise noise

than GCs (**Figure S3**). Then, linear regressions revealed that spiking probability (SP) is a good

predictor of both spiketrain unreliability $(1 - R_w)$ and decorrelation performance in FSs (**Figure**

S2B and **Table 2**). Surprisingly, the membrane resistance of FSs was also a good predictor

(**Table 1**). Thus, contrarily to GCs, FS pattern separation behavior is strongly and linearly

determined by some intrinsic and spike-wise properties, even though it is in principle hazardous

to anticipate complex neuronal operations from such low-level characteristics, as our previous

analysis on GCs illustrated. Indeed, spike-wise noise parameters of HMCs were very close to

those of GCs (**Figure S3**) and they showed striking pattern separation differences nonetheless.

Across all celltypes, it is clear that higher SP correlates with lower decorrelation levels

(**Figure S2D**). This suggests that sparseness can be a mechanism partially supporting temporal

pattern separation. Only partially, because the correlation is mediocre ($R^2 = 57\%$), and although

it is a better predictor in FSs, FSs decorrelation levels are consistently lower than expected by the

linear model fitting all celltypes (**Figure S2D**).

In the end, the best (but partial) predictor of decorrelation levels across all celltypes and

conditions is R_w, the spiketrain reliability (**Figure 8A** and **S5**). This result emphasizes the

unexpected idea that the biophysics of neurotransmission in hippocampal networks imposes a

trade-off between temporal pattern separation and reliable information transmission.

Overall, these results show that the differences in temporal pattern separation between

different hippocampal celltypes result from a combination of various sources, each celltype with

a unique combination.

Discussion

We report that similar cortical input spiketrains are transformed in the DG network, leading to

less similar output spiketrains in GCs. Our findings provide the first experimental demonstration

that a form of pattern separation is performed within the DG itself and exhibited at the level of

single neurons at different timescales. This computation arises from noisy but specific

biophysical processes (e.g. synaptic dynamics) in the DG network, where interneurons do not

exhibit as much temporal pattern separation as the final DG output. In turn, the CA3 network

seems to amplify this separation even more at the level of single PCs, suggesting that, at least in

the hippocampus, it is not a computation specific to the DG.

A novel way to test pattern separation

In contrast to *in vivo* experiments that have difficulty identifying recorded units with certainty

(Danielson et al., 2017; GoodSmith et al., 2017; Neunuebel & Knierim, 2012; Senzai & Buzsáki,

2017) and simultaneously recording the direct inputs of these units (Deng et al., 2013; Leutgeb et

al., 2007; Marrone et al., 2011; Nakazawa, 2017; Neunuebel & Knierim, 2014), *in vitro* brain

slices that preserve the lamellar connections of the hippocampus offer a more accessible

platform. For example, a similar experimental setup to ours was used to show that spatially

segregated inputs are represented by distinct spatiotemporal patterns in populations of DG

neurons (Hyde & Strowbridge, 2012; Zylberberg et al., 2016). However, our study is the first to

perform an experimental analysis of pattern separation within DG by manipulating the similarity

of the inputs and comparing it to the similarity of the outputs. Such a systematic approach had so

far only been done in computational studies (Chavlis & Poirazi, 2017). Although a rigorous

comparison is impossible because the activity patterns considered were defined differently, the

general pattern separation behavior of those models is confirmed by our experimental results: the

DG itself performs a form of pattern separation, especially for input patterns that are highly

similar (**Figure 2C-D**).

Pattern separation in the time dimension

Until now, most studies of pattern separation in the DG assumed that neural activity patterns

were ensembles of ON/OFF neurons (Deng et al., 2013; Marrone et al., 2011; Myers &

Scharfman, 2009; O'Reilly & McClelland, 1994; Rolls, 2010), sometimes considering a rate code

averaged over minutes in addition to this population code (Aimone et al., 2009; Leutgeb et al.,

2007; Neher et al., 2015; Neunuebel & Knierim, 2014). Because neurons carry information at

timescales shorter than minutes (Buzsaki, 2010; Eichenbaum, 2014; Kelemen & Fenton, 2016;

Rieke et al., 1999; VanRullen et al., 2005) and because the sparse firing of active GCs during a

brief event (Diamantaki et al., 2016; Nakazawa, 2017; Pernía-Andrade & Jonas, 2014) precludes

an efficient rate code (Rieke et al., 1999), we studied pattern separation at sub-second timescales.

Relevant scales are given by the time constant over which neurons can integrate synaptic

inputs (Buzsaki, 2010): 10-50 ms for GCs and ~100 ms for the "reader" CA3 pyramidal cells.

Windows of ~10 ms and ~100 ms, corresponding respectively to gamma and theta rhythms, have

been shown to organize CA neuronal assemblies (Buzsaki, 2010; Kelemen & Fenton, 2016;

Lisman & Jensen, 2013; Wikenheiser & Redish, 2015). Due to specific network properties

allowing persistent activity, the DG might also integrate information over several hundreds of

milliseconds (Hyde & Strowbridge, 2012; Larimer & Strowbridge, 2010; Zylberberg et al.,

2016). The point is that multiple timescales can be relevant simultaneously, and because it is still

uncertain which ones are the most important to episodic memory and hippocampal coding, we

investigated a range from 5 ms to 250 ms (**Figure 5**).

Most of our results are reported at a 10 ms resolution, which corresponds approximately

to the spike jitter in GCs (**Figure S1**) as well as their membrane time constant and the gamma

rhythm. This choice of temporal resolution is similar to a recent computational study of pattern

separation within a DG model, which used a 20 ms resolution on short spiketrains (30 ms inputs,

200 ms outputs) (Yim et al., 2014). Yet, our study was the first to investigate pattern separation

at a range of timescales. We found that temporal pattern separation in the DG output was best at

short timescales. This relationship was generally conserved across celltypes, with HMCs even

achieving the opposite of pattern separation, pattern convergence, at timescales above 100 ms.

Note, however, that temporal pattern separation is not necessarily a monotonically decreasing

function of the time resolution, as CA3 PCs exhibited a surprising sharp increase of temporal

pattern separation for low input similarity at 250 ms (**Figure 5**).

Our study of pattern separation is the first to focus on temporal patterns, as opposed to

spatial ones; but neural activity patterns are spatiotemporal. More work is clearly needed to test

whether the DG is a pattern separator at the spatiotemporal, population level, but the discovery

of temporal pattern separation in single hippocampal neurons has some implications for

population dynamics. The decorrelation of spiketrains at small timescales suggests that spikes

are constantly rearranged in different time windows, thus enforcing very small neuronal

assemblies (Buzsaki, 2010). In other words, it insures that a minimal number of output neurons

are active at the same time, and such sparsity in active neuronal population is known from

computational studies to be critical for efficient population pattern separation (Chavlis et al.,

2017; O'Reilly & McClelland, 1994; Severa et al., 2016).

Mechanisms of temporal pattern separation

The mechanisms supporting pattern separation within DG had so far never been experimentally

investigated. The decorrelation of sequentially presented input patterns can in theory be

explained by: 1) adaptive mechanisms, involving learning and recognition of input patterns,

comparison with previously stored ones and the pruning out of common features, 2) non-

adaptive (intrinsic) mechanisms, 3) or both (Friedrich & Wiechert, 2014). First, concerning

adaptive mechanisms, it has been suggested that Hebbian learning could enhance population

pattern separation in the DG (Fontanari et al., 1995), but computational models testing different forms of long-term synaptic learning found that it would actually impair this type of pattern separation (Neher et al., 2015; O'Reilly & McClelland, 1994). As for temporal pattern separation, our data show that it hardly benefits from the repetition of input patterns (**Figure 6**). We also offer indirect evidence that non-adaptive decorrelation processes support temporal pattern separation because output patterns are always decorrelated to the same proportion (**Figure 2E**), a feat that a simple random process can achieve (**Figure S4C**), suggesting that input patterns do not need to be recognized. Third, adaptive and non-adaptive mechanisms are not mutually exclusive: previous learning over days, during the neuron maturation process, could tune single GCs only to specific input patterns, allowing rapid pattern separation (Aimone et al., 2011). Indeed, a computational study suggested that adaptive networks can mature to perform a fast, non-adaptive orthogonalization of the population activity by the decorrelation between individual information channels (Wick et al., 2010).

Adaptive or not, what is the biological source of the temporal decorrelation we observed? We first determined that intrinsic membrane properties do not predict decorrelation levels (**Table 1**), and that celltypes ability to fire bursts only moderately affects pattern separation (**Figure S7**). Simple randomness was not sufficient to reproduce our results, even though the spiking probability, a form of spike-wise neural noise, plays a partial role (**Figure 8** and **S2**). In the end, it seems that the history-dependent noise of the specific presynaptic dynamics targeting GCs is the strongest candidate to support temporal pattern separation (**Figure 9**). Indeed, GCs are embedded in a network, notably receiving feedforward excitation from the PP, feedforward and feedback inhibition from FSs (Ewell & Jones, 2010; Hu et al., 2014) and both excitation and disynaptic inhibition from HMCs (Scharfman, 2016). Thus, although both FSs and HMCs

showed some temporal pattern separation (mostly at short timescales), our results suggests that

the final DG output is inherited from synaptic interactions between all DG celltypes, resulting in

maximal separation at the level of GCs.

More work is needed to clarify the exact role of FSs and HMCs in DG computations.

Computational and experimental studies have suggested that HMCs are involved in some forms

of pattern separation (Danielson et al., 2017; Myers & Scharfman, 2009; Nakazawa, 2017). Our

finding of pattern convergence in HMCs for dissimilar inputs appears in contradiction with the

recent reports that HMCs remap their firing rate in dissimilar environments (Nakazawa, 2017),

but this discrepancy likely comes from differences in the time resolution (sub-second vs

minutes). Overall, we show that HMCs can exhibit both separation or convergence, depending

on the time resolution and the amount of similarity between input patterns (**Figure 5**). The

impact of such a behavior on pattern separation at the level of GCs remains to be studied.

Concerning FSs, they exhibit a poor ability to separate spiketrains (**Figure 3, 5**), but the somatic

inhibition they provide could be the source of low spiking probability in GCs and thus improve

pattern separation by making GCs responses sparse (O'Reilly & McClelland, 1994; Temprana et

al., 2015). On the other hand, their ability to relay information reliably (Hu et al., 2014) (**Figure

S3, 5**) and to precisely control spike timing in target neurons (Hu et al., 2014) might actually

provide a mechanism that counteracts noisiness in GCs, helping them balance effective

separation with fidelity of information transmission to CA3.

The role of sweep-to-sweep variability

Because the brain needs to be able to recognize when situations are exactly the same, our finding

that pattern separation occurs even when the same input pattern is repeated (**Figure 7A**) might

seem counter-intuitive at first. However, in theory, the separation and the recognition functions

do not have to be supported by the same network. The Hebb-Marr framework actually

hypothesizes that the CA3 recurrent, auto-associative network is able to recall the original

pattern from a noisy input from DG. Even though most computational models that tested the

effect of repetition were consistent with the intuitive view (O'Reilly & McClelland, 1994; Yim et

al., 2014), this was likely because they used deterministic neurons. A model considering

variability across GCs and a probabilistic spiking behavior resulted, as in our experiments, in

separation of repetitions of the same pattern (Aimone et al., 2009).

In the cortex, the well-known variability of single neuron activity between trials is often

supposed to be "averaged out" at the population level so that the output of the population is

reliable (Faisal et al., 2008). It is thus conceivable that considering an ensemble of GCs would

increase the signal-to-noise ratio. In fact, when we average out the sweep-to-sweep variability,

GCs exhibit pattern separation for highly similar patterns but almost no separation for identical

ones (**Figure 8D**).

However, this variability, or "noise", is not necessarily meaningless (Faisal et al., 2008).

Our results suggest it might be a mechanism amplifying pattern separation (**Figure 8**). The

variability might even be just apparent, if we consider that when the same input is repeated it is

at different points in time: each repetition could be considered as a different event that needs to

be encoded slightly differently. The DG would thus meaningfully add some noise to transform

input spiketrains so that cortical information about an event is stored in the hippocampus with a

unique random time-stamp, consistent with the index theory of episodic memory (Teyler &

Rudy, 2007).

Pattern separation and pattern completion in CA3

In the Hebb-Marr framework, CA3 is thought to be a recurrent, autoassociative network that can perform pattern completion, and thus can support the recall of a full memory from a partial cue of the original event(Rolls, 2010). Although CA3 recurrent connections are sparser than previously thought (Guzman et al., 2016), computational models generally confirmed that CA3 could perform pattern completion of population patterns (Guzman et al., 2016; Neher et al., 2015; Rolls, 2010). Direct experimental evidence are scarce and unclear, both in vitro (Jackson, 2013), and in vivo (Denny et al., 2014; Marrone et al., 2014; Sun et al., 2017). In addition, the fact that neuronal representations of similar environments are more correlated in CA3 than in the DG was thought to constitute indirect evidence (Leutgeb et al., 2007; Neunuebel & Knierim, 2014), but recent reports suggesting that the DG representations previously measured were not coming solely from DG output cells have clouded these initial conclusions (GoodSmith et al., 2017; Senzai & Buzsáki, 2017).

Assuming that pattern completion is a process realized by CA3, this implies that, when presented with different but similar partial cues of the same initial memory, the final output of CA3 should converge towards the same representation. Our finding that CA3 PCs exhibit high levels of temporal pattern separation might then come as a surprise (**Figure 4-5**). Several lines of reasoning could explain this result. First, we focused on temporal patterns in single cells, and it is possible that a network can perform population pattern completion in addition to temporal pattern separation. Actually, different environments can be represented in CA3 by a different population of PCs, or by the same PCs with remapped firing rates (GoodSmith et al., 2017; Senzai & Buzsáki, 2017). Those reports are consistent with our finding that PCs are better than GCs at temporal pattern separation, as they show that single GCs remap much less than PCs.

Second, we tested CA3 under partial block of inhibition in order to allow PCs to fire. Given that

the number of active CA3 PCs is generally higher than DG GCs in vivo (GoodSmith et al., 2017;

O'Reilly & McClelland, 1994; Schmidt et al., 2012; Senzai & Buzsáki, 2017), it suggests that our

experiment may not have modelled physiological conditions well. Third, CA3 is known to be

physiologically and functionally heterogeneous along its proximodistal axis (Sun et al., 2017),

with studies suggesting that the PCs closest to DG perform population pattern separation

(Marrone et al., 2014). We recorded from PCs at the CA3b/c border, and it is possible that more

distal PCs would exhibit less temporal pattern separation. Last but not least, it is important to

note that *pattern completion* and *pattern separation* are not opposite, mutually exclusive

computations. Pattern separation or convergence describes the similarity of multiple patterns

from an input network (e.g. EC) compared to the similarity of the corresponding patterns of

another network (e.g. DG) (Santoro, 2013), whereas pattern completion is the process, happening

over time, of retrieval of a previously learned full pattern in a network (e.g. CA3) from a partial

seed pattern in the same network (O'Reilly & McClelland, 1994). Our experiments did not test

for pattern completion, as the different trains of an input set were not degraded versions of a

previously learned pattern. Moreover, pattern separation and completion are complementary in

the sense that pattern completion would benefit from initial input patterns being as separated as

possible (O'Reilly & McClelland, 1994; Rolls, 2010). It thus makes sense that CA3 would start

by amplifying the separation of input patterns, as is the case in our data, either before encoding

or before proceeding to completion of the seed pattern. In fact, our results suggest that CA3

might complement the separation effectuated in the DG, as CA3 is able to perform high

decorrelation levels at long timescales (> 100 ms) when the DG doesn't (**Figure 5**). CA3 could

thus ensure that seed input patterns are well separated at all timescales.

Materials and methods

Animals and dissection

All experiments were performed in accordance with the National Institute of Health guidelines

outlined in the National Research Council Guide for the Care and Use of Laboratory Animal

(2011) and regularly monitored and approved by the University of Wisconsin Institutional

Animal Care and Use Committee.

Horizontal slices of the ventral and intermediate hippocampus (400 μm) were prepared

from the brains of young (p15-25) or adult (p121 ± 15 days) C57BL/6 male mice

(Harlan/Envigo). Adult animals were only used for **Figure 9**. Mice were anesthetized with

isoflurane, decapitated, and the brain was removed quickly and placed in ice-cold cutting

sucrose-based solution (two different compositions were used. For HMC, CA3 pyramidal cells,

their GC controls and GCs from adult mice, we used version #2 (Yi et al., 2015)). The cutting

solutions contained (version #1 / version #2, in mM): 83 / 80 NaCl, 26 / 24 $NaHCO_3$, 2.5 / 2.5

KCl, 1 / 1.25 NaH2PO4, 0.5 / 0.5 $CaCl_2$, 3.3 / 4 MgCl2, 22 / 25 D-Glucose, 72 / 75 Sucrose, 0 / 1

Na-L-Ascorbate, 0 / 3 Na-Pyruvate, bubbled with 95% O_2 and 5% CO_2. During the dissection,

brain hemispheres were prepared following the "magic cut" procedure (Bischofberger et al.,

2006) with α-angle around 10 to 15° and β-angle around 5 to 10° for GC and FS recordings, and

with α close to 0° and β between 0 and -5° for HMC and CA3 pyramidal cells recordings

(Guzman et al., 2016). Slices were cut using a vibratome (Leica VT1000S) then placed in an

incubation chamber in standard artificial cerebrospinal fluid (aCSF) containing (in mM) 125

NaCl, 25 NaHCO3, 2.5 KCl, 1.25 NaH2PO4, 2 CaCl2, 1 MgCl2, and 25 D-Glucose (or in a

50/50 mix of standard aCSF and cutting solution. 100% cutting solution for HMC, CA3

recordings and their GC controls, and GCs from adult mice) at 35°C, for 15-30 minutes after

dissection. Slices were stored in the incubation chamber at room temperature for at least 30

minutes before being used for recordings.

Electrophysiology

All recordings were done in standard aCSF adjusted to 325 mOsm, at physiological temperature

(33-35 °C). Whole cell patch-clamp recordings were made using an upright microscope

(Axioskop FS2, Zeiss, Oberkochen, Germany) with infra-red differential interference contrast

optics. Patch pipettes pulled from thin-walled borosilicate glass (World Precision Instruments,

Sarasota, FL) had a resistance of 2-5 MΩ when filled with intracellular solution containing (in

mM) 140 K-gluconate, 10 EGTA, 10 HEPES, 20 phosphocreatine, 2 Mg_2ATP, 0.3 NaGTP. For

the dataset from adult mice, we used a slightly different recipe: 135 K-gluconate, 5 KCl, 0.1

EGTA, 10 HEPES, 20 Na-Phosphocreatine, 2 Mg_2-ATP, 0.3 Na-GTP, 0.25 $CaCl_2$. Intracellular

solutions were adjusted to pH 7.3 and 310 mOsm with KOH and H_2O. Both recipes yielded

similar electrophysiological behaviors.

Recordings were done using one or two Axopatch 200B amplifiers (Axon Instruments,

Foster City, CA), filtered at 5 kHz using a 4-pole Bessel filter and digitized at 10 kHz using a

Digidata 1320A analog-digital interface (Axon Instruments). Data were acquired to a Macintosh

G4 (Apple Computer, Cupertino, CA) using Axograph X v1.0.7 (AxographX.com). Stimulation

pipettes were pulled from double barrel borosillicate theta-glass (~10 μm tip diameter, Harvard

Apparatus, Edenbridge, U.K.) and filled with ACSF or a 1M NaCl solution and connected to a

constant current stimulus isolator used to generate 0.1-10 mA pulses, 100 microseconds in

duration.

Series resistance and cellular intrinsic properties were assessed online in Axograph from

the fit of the electrical responses to repetitions of 5-10 mV, 25 ms steps, holding the potential at -

65 mV. Neurons used for analysis were stable across a whole recording session as judged by

monitoring of series resistance and resting potential.

Dentate granule cells (GC) were visually identified as small cells in the granule cell layer

(GCL). GCs from young mice had the following intrinsic properties (mean ± sem) : resting

potential (V_{rest}) -69.3 ± 1.3 mV; input resistance (R_i) 171 ± 16MΩ and capacitance (C_m) 23 ± 2

pF. GCs from adult mice had the following intrinsic properties: V_{rest} -78.8 ± 1.8 mV; R_i = 137 ±

14 MΩ and C_m = 18 ± 1 pF.

Fast-spiking interneurons (FS) were identified as neurons with large somata at the hilus-

GCL border and a high firing rate response during large depolarizing current steps, and a large

after-hyperpolarization (AHP) (Ewell & Jones, 2010; Harney & Jones, 2002) (**Figure 3**). They

had the following intrinsic properties: V_{rest}= -66.7 ± 3.5 mV; R_i = 59 ± 10 MΩ and C_m = 19 ± 3

pF.

Hilar Mossy Cells (HMC) were identified as large neurons in the deep hilus (> 60 μm

away from the GCL) with a regular firing response to current steps and a small AHP, as well as

with a high frequency of spontaneous EPSPs (Larimer & Strowbridge, 2008; Scharfman, 2016)

(**Figure 3**). Their intrinsic properties were: V_{rest} = -69.7 ± 2.3 mV; R_i = 198 ± 12 MΩ and C_m =

33 ± 3 pF.

For CA3 pyramidal cells, because recent studies suggest a proximodistal gradient of

physiological and computational properties in the CA3 network (Marrone et al., 2014; Sun et al.,

2017), we avoided the extremity of the CA3c region and targeted large neurons around the

CA3b/c border in the pyramidal layer. Their intrinsic properties under partial inhibitory block

(aCSF + 100 nM gabazine) were: V_{rest} = -72.7 ± 2.2 mV; R_i = 186 ± 12 MΩ and C_m = 36 ± 2 pF.

The intrinsic properties of GCs recorded under the same pharmacological conditions were: V_{rest}

= -77.3 ± 1.9 mV; R_i = 244 ± 13 MΩ and C_m = 25 ± 2 pF.

Pattern separation experiments

We designed multiple sets of input patterns, each with a prespecified average Pearson's

correlation coefficient (R_{input}) computed with a binning window (τ_w) of 10 ms, using two

different algorithms (one developed in-house based on iterative modifications, and a more

efficient one from (Macke et al., 2009) based on mathematical principles and a preset covariance

matrix). An input set consisted of five different input spiketrains, each 2 s trains of impulses

simulating cortical spiketrains, with interspike intervals following a Poisson distribution. Each

pair of input trains had a correlation coefficient close to the R_{input} of its set (at τ_w = 10 ms, the

average relative standard error is 4% across all input sets). For all experiments except in **Figure**

4 (CA3 and GC control), the mean frequency of input trains was set close to 10 Hz (11.9 ± 0.7

Hz). This input firing rate was chosen to be consistent with the frequency of EPSCs recorded in

GCs of behaving mice (Pernía-Andrade & Jonas, 2014), and is known to promote a high

probability of spiking in GCs in slices (Andersen et al., 1966; Ewell & Jones, 2010).

The responses of one or two DG neurons were recorded in whole-cell mode while stimuli

were delivered to the outer molecular layer (OML). Stimulus current intensity and location were

set so that the recorded neuron spiked occasionally in response to electrical impulses (see range

of spike probability in **Figure S1-3**) and the stimulation electrode was at least 100 μm away

from the expected location of the dendrites of the recorded neuron. Once stimulation parameters

were set, a pattern separation protocol was run: the five trains of a given input set were delivered

one after the other, separated by 5 s of relaxation, and this was repeated ten times. The ten

repetitions of the sequence of five patterns were implemented to take into account any potential

variability in the output, and the non-random sequential scheme was used to avoid repeating the

same input spiketrain close in time. Each protocol yielded a recording set of fifty output

spiketrains, each associated with one of the five input trains of an input set (**Figure 1C**). A given

cell was recorded in response to up to five input sets with different R_{input} (i.e. a recorded cell

produced between one and five data points on **Figure 2C**).

The membrane potential baseline was maintained around -70mV during both current-

clamp and voltage-clamp recordings, consistent with the V_{rest} of mature GCs recorded in

behaving mice (Pernía-Andrade & Jonas, 2014). For comparison, FS and HMC current-clamp

recordings were also held at -70 mV. The output spiking frequency of GCs was variable (6.3 ±

0.3Hz, see **Figure S6-7**) but consistent with sparse activity generally observed in GCs *in vivo*

(Danielson et al., 2016; Diamantaki et al., 2016; Pernía-Andrade & Jonas, 2014; Senzai &

Buzsáki, 2017) and in slices under conditions of drive comparable to ours (Dieni et al., 2016;

Mongiat et al., 2009; Pardi et al., 2015).The output firing rates of FS and HMC were higher than

their GC controls (**Figure S6-7**), as expected from recent research (Danielson et al., 2017; Hu et

al., 2014; Senzai & Buzsáki, 2017).

To perform pattern separation experiments in CA3 pyramidal cells (PCs), we had to

change several parameters in order to make CA3 PCs spike. Indeed, PCs firing is controlled by

strong feedforward inhibition (Rebola et al., 2017; Sun et al., 2017), and all tested stimulation

sites led to net IPSPs or, rarely, to weak EPSPs. To make PCs fire in response to external

electrical stimulations: 1) the stimulating electrode was placed in the inferior blade of the GCL to

make the DG output cells fire, 2) we targeted CA3b PCs, which have the highest E/I ratio across

CA3 in slices and still receive strong connections from mossy fibers (Sun et al., 2017), 3) we

used input sets with 30 Hz input trains to promote the depression of inhibitory transmission

(Rebola et al., 2017), 4) we maintained the membrane potential baseline between -60 and -70

mV, and 5) we had to add 100 nM gabazine to the bath of standard aCSF in order to slightly

decrease IPSCs amplitude (recordings started at least 15 min after the slice was placed in the

bath, to insure equilibrium was reached). 100 nM of gabazine (SR-95531, Sigma-Aldrich) has

been shown to correspond to a 70% availability of GABA-A receptors (Jones et al., 2001) and, in

our conditions, consistently decreased spontaneous IPSCs mean amplitude by 30% in voltage-

clamp recordings of GCs and CA3 PCs held at -40 mV. These conditions allowed us to record

for the first time the spiking output of CA3 PCs in response to complex input spiketrains while

preserving some inhibition in the network (Rebola et al., 2017). PCs output firing rate were on

average below 10Hz (**Figure S6-7**), but close to mean rates observed in vivo (Leutgeb et al.,

2007; Neunuebel & Knierim, 2014; Senzai & Buzsáki, 2017),

Analysis of the output spiketrains

For each recording set, the similarity between pairs of spiketrains was computed as the Pearson's

correlation coefficient between the spiketrains rasters binned at a τ_w timescale. Sweeps without

spikes were excluded from further analysis.

We did not use separate protocols to assess the *firing rates*, *probability of bursting* and

spike-wise noise parameters (spike probability, delay, jitter), but computed them directly for each

recording set of spiketrains from a pattern separation experiment. The mean firing rate was

computed as the average firing rate across all fifty output spiketrains. A burst was defined as the

occurrence of more than one output spike in the interval of two input spikes (see **Figure S7**).

We define the spike-wise neural noise as the *probability of spiking* at least once after an

input spike, the *delay* of an output spike after an input spike and its average *jitter*. To assess these

parameters, we computed the cross-occurrence between input spikes and output spikes in a [-15

ms, 50ms] interval with 1 ms bins. The resulting histogram of counts of output spikes occurring

in the vicinity of an input spike was fitted with a Gaussian distribution $N(\mu,\sigma, baseline)$, where μ

is the mean delay of an output spike and σ is the jitter of this delay. The baseline corresponds to

the background firing, occurring by chance or caused by neighboring inputs. After subtracting

the baseline and extracting the probability of spiking by dividing the counts of output spikes by

the total number of input spikes, we defined the *spike probability* (SP) as the sum of probabilities

of an output spike in the predefined time interval around an input spike (**Figure 7B** and **Figure**

S1A). For FS and HMC, spike-wise noise parameters were computed on a non-bursty dataset

(nbFS and nbHMC) where all but the first spike in a burst were excluded. Spike-wise noise

parameters were not computed for CA3 PCs and their GC control because the input frequency

was too high to allow a good fit of the distribution.

Computational models

To assess the role of spike-wise neural noise in pattern separation, we generated two data sets.

First, we simulated output spiketrains in response to our input sets (for each input set, we

simulated ten output sets of fifty synthetic spiketrains). This simulation was entirely based on the

average spike-wise noise parameters computed from the original GC recordings (see above): the

matrix of input spike times was replicated ten times, and for each of the fifty resulting sweeps,

spikes were deleted randomly following a binomial distribution B(Nspk, F), where Nspk is the

number of input spikes in a sweep, and F the probability of not spiking (F = 1-mean SP = 1-

0.42). A random delay, sampled from a Gaussian distribution $N(\mu,\sigma)$, was added to each

resulting spike times, with μ and σ being respectively the mean delay and mean jitter in the

original recordings. The noise statistics of the resulting simulated data set is shown in **Figure**

S1C. Second, we created a surrogate data set by randomly shuffling the output spikes of the

original GC recordings: the delay of each spike was conserved but it was relocated to follow a

randomly selected input spike in the same input train (from a uniform distribution). This strategy

yielded a data set with noise statistics closer to the original data (**Figure S1D)**.

To test whether probabilistic synaptic dynamics is a potential mechanism of temporal

pattern separation, we used an Izhikevich model of a regular spiking neuron (Izhikevich, 2003)

with an adapted Tsodyks-Markram (TM) synapse model (Fuhrmann et al., 2002) designed to

capture short-term plasticity of stochastic neurotransmission at the LPP-GC synapse. The TM

synapse model consists of a system of two ordinary differential equations describing the

dynamics of X and U. X is the probability of a presynaptic vesicle being available for release, the

decrease of which leads to synaptic depression, and U is the probability of release for an

available vesicule, the increase of which models facilitation:

$$\frac{dX}{dt} = U(t).X(t).\delta\big(t - t_{ap}\big) + \frac{1 - X(t)}{\tau_{rec}} \quad (\mathbf{1})$$

$$\frac{dU(t)}{dt} = U_{SE}.\big(1 - U(t)\big).\delta(t - t_{AP}) - \frac{U(t)}{\tau_{facil}} \quad (\mathbf{2})$$

With X(0) = 1 and U(0) = 0, where τ_{rec} is the time constant of vesicule recovery after release,

τ_{facil} is the decay of U controlling synaptic facilitation, U_{SE} a factor determining the probability of

release Pr at the time of the first spike, δ the Dirac delta function, and t_{AP} the time of arrival of an

input spike. The system of ODEs was solved using the ode23 MATLAB solver.

Based on the quantal theory of neurotransmission, the model assumes that, at each time

point t, *k* vesicles are released, drawn from a binomial distribution *B*(N, Pr), where N is the

number of release sites and Pr is the probability of release defined as the product of X and U.

The current *I* is then the product of *k* with the quantal size *q*. Based on previous data from our

lab, we considered *q* = 20 pA (Ewell & Jones, 2010). The version of the TM model we

implemented only focuses on the dynamics of the presynaptic vesicles, ignoring postsynaptic

dynamics. To match the decay of real EPSCs recorded at the soma, *I* was convolved with a

template EPSC shape:

$$-A \cdot \left(e^{-T/\tau_{rise}} - e^{-T/\tau_{decay}} \right) \quad (3)$$

Where T = 50 ms, τ_{rise} = 0.5 ms, τ_{decay} = 5 ms and A is a factor normalizing the area of the

resulting shape to 10. In addition, a synaptic delay of 4 ms was introduced. The resulting current

(**Figure 9B-C**) is then passed to a non-dimensional regular spiking Izhikevich neuron, an

acceptable model for GCs (Hananeia & Benuskova, 2016). We used the following parameters: a

= 0.02, b = 0.2, c = -65 mV, d = 6, time resolution = 0.1 ms, max voltage = 30 mV, initial values

V(0) = -70 mV and x(0) = b.V(0).

To optimize the non-fixed parameters of the TM-Izhikevich model, we simulated output

spiketrains in response to ten repetitions of a single input set (10 Hz, R_{input} = 0.76) and compared

to the current dynamics and spiking behavior of a representative GC that was recorded both in

voltage-clamp and current-clamp in response to the same input set. For the TM model, the

following parameters were adjusted to match the dynamics and the peak of the current averaged

over ten repetitions, as well as the variability in the peak amplitude of the first EPSC: τ_{rec} = 500

ms, τ_{facil} = 9 ms, U_{SE} = 13, N = 40 release sites. Finally, because the Izhikevich model is non-

dimensional, the input current had to be scaled using the dividing constant K = 62, such that the

average firing rate of the simulation matched the average firing rate of the representative GC to

the same input set (3.5 Hz). In the end, the standard deviation of the firing rate was ~0.6 Hz for

simulations, compared to 0.8 Hz for the original recording. The only source of variability was

from presynaptic dynamics of vesicle release modelled as a binomial process.

Software and statistics

Data analysis was performed using custom-written routines in MATLAB (Mathworks, Natick,

MA, USA), including functions from toolboxes cited above. Sample sizes were chosen based on

the literature and estimations of the variance and effect size from preliminary data. All values are

reported as mean ± S.E.M. unless otherwise noted. The one-sample Kolmogorov-Smirnov test

was used to verify the normality of data distributions. Parametric or non-parametric statistical

tests were appropriately used to assess significance (p-value < 0.05). Assumptions on equal

variances between groups were avoided when necessary. All T and U tests were two-tailed. To

determine whether two distributions of data points are significantly different (e.g. R_{output} as a

function of R_{input}, for GC compared to FS, see **Figure 3-7 and Figure S4 and S7**), we performed

an analysis of the covariance (ANCOVA) using a linear or parabolic regression on the two data

sets as well as on the combined data set, and assessed significance via an F-test comparing the

goodness of fits (Motulsky & Ransnas, 1987). Because R_{input} can also be considered as a

categorical variable, we performed a two-way ANOVA before using post-hoc tests correcting for

multiple comparisons in order to determine at which R_{input} groups two conditions were

significantly different. When comparing FSs, HMCs or CA3 with GCs, different GCs were used

as control and were recorded under different protocols, which is why we did not need to control

for multiple comparisons across celltypes.

Figure 1. Pattern separation assay in acute brain slices at the single cell level.

(A) Examples of input sets. *Top*: each input set comprises five different trains of electrical pulses. *Bottom*: correlation coefficient matrix for each input set, each square representing the correlation coefficient between two input trains measured with a binning window (τ_w) of 10 ms. R_{input} is the average of coefficients, diagonal excluded.

(B) Histology of the DG in a horizontal slice (Cresyl violet/Nissl staining; scale bar: 250 μm), overlaid with a schematic of the experimental setup: a theta pipette in the ML (input) is used to focally stimulate the PP while a responding GC (output) is recorded via whole-cell patch-clamp.

(GCL: granule cell layer, H: hilus, ML: molecular layer, FS: fast-spiking interneuron. Solid lines

represent dendrites and dashed lines axons)

(C) Current-clamp recordings of the membrane potential of two different GCs in response to

different input sets (*Top*: $R_{input} = 1$; *Bottom*: $R_{input} = 0.76$). Each input set (five input traces) is

repeated ten times (only three repetitions are shown, with spikes truncated at 0 mV). In the

bottom graph, input trains and their respective children output spiketrains have matching colors.

Figure 2. Input spiketrains are decorrelated at the level of individual granule cells.

(A) Example of a recording set (input set + output set): the raster plot shows one set of input

spiketrains and the children output spiketrains recorded from one GC, reordered to display output

subsets (i.e., the ten children coming from one parent input spiketrain) together and with the

same color.

(B) Corresponding 55x55 correlation coefficient matrix using a binning window (τ_w) of 10 ms.

Each small square represents the correlation coefficient between two spiketrains. R_{output} is

defined as the mean of correlations between individual output spiketrains driven by different

input spiketrains, as outlined by the bold blue border, which excludes comparisons between

outputs generated from the same parent input.

(C) Data points, corresponding to 102 recording sets (28 GCs), are all below the identity line

(dashed line). This means that R_{output} was lower than R_{input} for all recordings, thus demonstrating

pattern separation.

(D) *Left*: Effective decorrelation averaged over all recording sets as a function of R_{input}. Although

there is a significant decorrelation for all tested input sets (one-sample T-tests: the blue shade

indicates the 95% confidence interval that average decorrelation is significantly above 0), they

are effectively decorrelated to different magnitudes (one-way ANOVA, $p < 0.0001$). *Right:*

Matrix of p-values from post-hoc Tukey-Kramer tests comparing effective decorrelation levels

across all pairs of parent input sets. The asterisk and star correspond to the comparisons

displayed in the left panel. This analysis shows that the decorrelation is significantly different

(higher) for highly similar input spiketrains than for already dissimilar inputs.

(E) When the effective decorrelation is normalized to the correlation of the input set, there is no

significant difference between input sets (ANOVA, $p = 0.18$).

In all graphs, $\tau_w = 10$ ms. Means and SEM in black.

Figure 3. Different levels of temporal pattern separation in different DG cell types

(A1) Picture of a recorded FS filled with biocytin (black). In the case of simultaneous recordings, the recorded GCs were close to the FS, as depicted by the schematic in green. In different experiments, we recorded from putative hilar mossy cells (HMC, orange). Full lines represent dendrites, dashed lines axons.

(A2-3) Example of a simultaneous whole-cell recording of a GC and a neighboring FS.

(A2) Simultaneous membrane potential recordings (baseline at -60mV) of a FS and a GC to the same set of current steps (-25, 100, 500 and 1000 pA).

(A3) Simultaneous current-clamp recordings of the same FS and GC as in A2 in response to the

five input traces of an input set with $R_{input} = 0.65$. Simultaneous input and output trains have the

same color.

(A4) R_{output} versus R_{input} for FSs and GCs. Data points correspond to recording sets: 20 for FS

(red, 4 cells, 4 per input set), and 61 for GC (green, with a darker shade open circle when

simultaneously recorded with a FS, 13 cells, 11-13 per input set). All GC recordings done at the

same input correlations as FS recordings were used for an unpaired comparison showing that FS

exhibit less spiketrain decorrelation than GCs: ANCOVA: $p < 0.0001$, 95% confidence bounds

around the linear fits shown as shaded areas; two-way ANOVA: input sets: $p = 0.0016$, cell-

types: $p < 0.0001$, interaction: $p = 0.72$. Post-hoc T-tests with Bonferroni corrections for 5

comparison groups: all $p < 0.05$ (for decreasing R_{input}, $p = 0.0307, 0.0181, 0.0007, 0.0001$,

0.0122)

(A5) Effective decorrelation ($R_{input} - R_{output}$) for FS and GC. Shaded areas represent the 95%

confidence interval of a bootstrap test comparing the amount of decorrelation between both

celltypes: GCs exhibit significantly more pattern separation than FS for all input sets.

(**A4-5**) Note that when comparing only the simultaneous GC and FS recordings, we found a

similarly significant difference between celltypes.

(**B1**) Membrane potential of a hilar neuron in response to current steps (-100, 0, 100, 400 pA;

baseline at -70mV), showing a spontaneous barrage of EPSPs, regular spiking, and a lack of

large after-hyperpolarization, all typical features of HMCs (Larimer & Strowbridge, 2008).

(**B2**) Current-clamp recordings of same HMC in response to a set of five input trains ($R_{input} =$

0.49, $\tau_w = 10ms$). HMCs fire occasional bursts of spikes (marked by asterisks) in response to a

single input, which was not seen in GCs.

(**B3**) R_{output} versus R_{input} for HMCs and GCs. Data points correspond to recording sets: 18 for

HMC (orange, 11 cells, 5-7 per input set), and 22 for GC (green, 11 cells, 4-10 per input set). An

unpaired comparison suggests that HMCs and GCs show only slight differences in pattern

separation measured at τ_w = 10 ms: ANCOVA: p = 0.15, 95% confidence bounds around the

linear fits shown as shaded areas; two-way ANOVA: input sets: p = 0.0004, cell-types: p =

0.074, interaction: p = 0.57. Post-hoc T-tests with Bonferroni corrections for three comparison

groups (for decreasing R_{input}): p = 1, 0.05, 0.21.

Figure 4. CA3 pyramidal cells exhibit more temporal pattern separation than GCs

(**A**) Schematic of the experimental setup: a theta pipette in the granule cell layer (GCL) is used to

stimulate GCs and their mossy fibers (dashed line) to evoke action potentials in CA3 pyramidal

cells recorded via whole-cell patch-clamp. To limit feedforward inhibition and allow pyramidal

cells to spike, experiments were performed under partial block of inhibition (100 nM of

gabazine). Control experiments were performed in GCs under the same pharmacological

conditions but with OML stimulations.

(**B**) Two input sets of 30 Hz Poisson trains were used. *Top*: rasters of the five spiketrains of each

set. *Bottom*: correlation coefficient matrix for each input set (τ_w = 10 ms).

(**C**) Example of current-clamp recordings from a pyramidal cell and a GC from the same animal.

Left: Membrane potential responses to current steps (-100pA, 100pA, 350pA). Scale: 100 ms

(horizontal) and 50 mV (vertical). *Right:* example of 5 output responses (sweeps 11-15) to an

input set ($R_{input} = 0.76$).

(D) R_{output} versus R_{input} for CA3 pyramidal cells and GCs. Data points correspond to recording

sets: 15 for CA3 (black, 14 cells, 6-9 recordings per input set), and 22 for GC (green, 13 cells, 11

per input set). The CA3 distribution is lower and significantly different from the GC distribution:

ANCOVA: $p < 0.01$, 95% confidence bounds around the linear fits shown as shaded areas; two-

way ANOVA: input sets: $p < 0.0001$, cell-types: $p = 0.0036$, interaction: $p = 0.24$. Post-hoc T-

tests with Bonferroni corrections for two comparison groups: $p = 0.032$ for $R_{input} = 0.76$, $p = 0.1$

for $R_{input} = 0.21$.

Figure 5. Differences in temporal pattern separation between hippocampal celltypes depend on the timescale.

(**A**) Pairwise analysis. Instead of averaging across all five input trains of a set (here $R_{input} = 0.76$ at $\tau_w = 10$ ms, same matrix as in Fig. 2B), we average children output coefficients corresponding to a single pair of input trains (identified by color-coded squares.). This finer analysis is necessary at τ_w higher than 20 ms because the *pairwise input correlation coefficients* are not well constrained around their mean anymore (see right panel). *Middle* and *Right*: mean across cells, following the same color-code as displayed in the matrix (*Left*).

(**B**) Pairwise R_{output} vs pairwise R_{input} for GCs, measured with different binning windows τ_w.

Only the means across cells are displayed, but the distribution of data points for single recording

sets is fitted with a linear model. The larger τ_w, the less GCs exhibit pattern separation.

(**C**) *Top:* Pairwise analysis on FS recordings (same color code as in B). *Bottom*: At $\tau_w = 100$ ms

and 250 ms, FS and GC distributions are still different, especially at high input similarity:

ANCOVA: $p < 0.0001$; two-way ANOVA: input sets: $p < 0.0001$, cell-types: $p < 0.0001$,

interaction: $p = 0.98$. Note that the larger τ_w, the lower the difference between FSs and GCs.

(**D**) *Top:* Pairwise analysis on HMC recordings (same color code as in B). *Bottom*: At $\tau_w = 100$

ms and 250 ms, HMC and GC distributions are different, especially at lower input similarity:

ANCOVA: $p < 0.0001$; two-way ANOVA: input sets: $p < 0.0001$, cell-types: $p < 0.0001$,

interaction: $p < 0.0001$. Note that under this analysis, HMCs and GCs are also significantly

different at $\tau_w = 10$ ms (ANCOVA: $p < 0.0001$; two-way ANOVA: input sets: $p < 0.0001$, cell-

types: $p < 0.0001$, interaction: $p = 0.3$). But, the larger τ_w, the higher the difference is.

(**E1**) Pairwise analysis on CA3 pyramidal cells and GCs, both under partial inhibition block and

under 30 Hz inputs (same color code as in B).

(**E2**) At $\tau_w = 100$ ms and 250 ms, CA3 and GC distributions are still different: ANCOVA: $p <$

0.0001; two-way ANOVA: input sets: $p < 0.0001$, cell-types: $p < 0.0001$, interaction: $p < 0.0001$.

Bars are SEM, asterisks denote $p < 0.05$.

Figure 6. Input spiketrains are efficiently separated upon their first presentations.

(A) Two of five inputs are shown with corresponding output spiketrains. The first output sweep is marked with a pink bar (right) and last sweep is marked with a blue bar.

(B) R_{output}, computed from the first sweep of five output trains only (pink), as a function of R_{input}, fitted with a parabola. All data points are below the identity line indicating that outputs are effectively decorrelated compared to their inputs even when input patterns have only been presented once each. The average decorrelation (R_{input}-R_{output}) is significant for all input sets (one-sample T-tests, $p < 0.01$) except for $R_{input} = 0.11$ ($p = 0.1$).

(C) *Left*: Average output correlations between spiketrains of the first sweep (pink) and the last sweep (blue). There is no significant difference (F-test comparing the two distributions using parabolic regressions, $p = 0.47$. Error bars are SEM). *Right:* When taking into account that the two distributions are paired, we detect that a few output correlations are significantly lower for

912 the last sweep than for the first one (one-sample T-test on the difference between R_{output} of the

913 first and last sweep of each recording set, asterisks signify $p < 0.05$). This is evidence, though

914 weak, that repetition of input spiketrains might improve pattern separation for highly similar

915 inputs.

Figure 7. Pattern separation in single GCs is not explained by simple neural noise.

(A) The variability of output spiketrains in response to the same input train sets the upper bound

for R_{output}. *Left:* Correlations between pairs of output spiketrains associated with different input

trains (enclosed by red, R_{output}) and pairs of different output spiketrains associated with the same

input train (enclosed by green, R_w: spiketrain reliability, the reproducibility of the output given

the same input). *Right:* Empirical probability distribution of R_w for all recordings (dark green

line is the mean: $\langle R_w \rangle = 0.3$), overlaid on the distribution of R_{output} as a function of R_{input} (102

recording sets) fitted with a parabola. Note that means $\langle R_w \rangle$ and $\langle R_{output} \rangle$ for $R_{input} = 1$ are close

because they both assess the reproducibility of the output when the input is the same.

(B) Characterization of neural noise. *Top:* example of input and output spiketrains illustrating

variable delay of the response spike after an input spike (d1, d2) or failure to spike after an input

spike (red cross). *Bottom*: Example from one GC recording. The spike-wise noise in output

spiketrains is characterized by the average spike delay, the standard deviation of this delay

(jitter) and the probability of spiking after an input spike (spiking probability, SP).

(C-D) Effect of random shuffling on R_{output} and R_w.

(C) Patterns are *less* separated for GC than for random shuffling (F-test using parabolic

regressions: $p < 0.0001$).

(D) GC output is *more* reliable than for random shuffling (unpaired T-test, $p < 0.0001$; mean

$<Rw>\{shuffle\} = 0.19$).

(E) Paired statistical tests show that shuffling leads to smaller R_{output} and R_w than original

recordings. *Top:* paired T-test on all recording sets, $p < 0.0001$. *Bottom*: one-sample T-test on

difference between shuffled and original data R_{output} performed on each R_{input} group. Black

symbols correspond to the means, bars to SEM. The purple shade indicates the 95% confidence

interval. Asterisks signify $p < 0.001$.

Figure 8. Unreliability in spiketrain transmission is a major but not unique source of temporal pattern separation.

(**A**) Spiketrain reliability (R_W) is an excellent predictor of normalized decorrelation (defined in Figure 2E) for all celltypes and conditions. Notice that, despite the strong anticorrelation, the intercept of the linear model at $R_W = 1$ predicts that even a perfect reliability could still allow 10% of decorrelation. See Table 3 for linear regressions on single celltypes.

(**B**) To assess the amount of decorrelation not due to spiketrain unreliability, the ten children output spiketrains of each of the five trains of an input set can be averaged to give the five output peristimulus histograms (PSTH). The 10 ms binned PSTHs of the output rasters in Figure 2A are shown.

(**C**) Correlation coefficients between all pairs of the five output PSTHs. The mean correlation (PSTH R_{output}) is the average of coefficients inside the red border, and excludes self-comparisons.

956   **(D)** *Left:* PSTH R_{output} as a function of R_{input} (102 recording sets, in red), fitted with a parabola

957   (black). All points are below the identity line indicating decorrelation of outputs compared to

958   inputs. *Right:* Average effective decorrelation (R_{input} – PSTH R_{output}) as a function of R_{input} (bars

959   are SEM) reveals a significant decorrelation for all input sets except for the most dissimilar (one-

960   sample T-tests; shaded area is the 95% confidence interval for significant decorrelation).

Figure 9. Probabilistic presynaptic release is a potential mechanism balancing temporal

pattern separation and spiketrain reliability

(A) Non simultaneous current-clamp ($V_{rest} \approx$ -70mV) and voltage-clamp recordings (V_{hold} = -

70mV) in response to a set of input patterns were successively performed in GCs from adult

mice. The example shows the first sweep of responses from a single cell to the five trains of an

input set (R_{input} = 0.76, τ_w = 10 ms).

(B-E) The recording set of the example cell in A was used to fit a computational model of a

spiking neuron with dynamic and probabilistic synapses.

(B) *Top*: four first input spikes of input train #1. *Middle*: Dynamics of the variables of a

probabilistic Tsodyks-Markram model of a synapse in response to the four input spikes. X

represents the probability of vesicle availability at the presynaptic site, U represents the

probability of release of available vesicles, and Pr is the probability of release for all vesicles.

The number of released vesicles was simulated at every time point from a binomial process.

Bottom: Parameters of the model were adjusted so that the resulting current (averaged over the

10 repetitions, in dark orange) would reasonably match the peak of the corresponding average

EPSCs in the original voltage-clamp recordings (baselined and inverted display with partially

blanked stimulation artifact, in black).

(C) Parameters of the model were also adjusted to match the variability in the amplitude of the

first EPSC in the original recording. *Top*: average EPSC. *Middle*: individual EPSCs evoked from

the 10 repetitions of the first input spike in input train #1. The range of peak amplitudes is

similar between model and data. *Bottom*: The coefficient of variation of the current in the model

is close to the data.

(D-E) Temporal pattern separation and spiketrain reliability for GCs from adult mice and a

single model regular spiking Izhikevich neuron with probabilistic synapses as described above.

Data points correspond to recording/simulation sets: 35 for GCs (green circles, 14 cells, 6-16 per

input set), and 5 simulations at all input sets (dark orange triangles), with the model fitted to a

single recording set ($R_{input} = 0.76$; R_{output} and R_w shown in dark green circles for all input sets).

Figure S1 related to Figure 7. Spike delay, jitter and reliability distributions for real data,

simulations and shuffled data

(**A**) Cross-occurrence method to measure spike delay, jitter and spiking reliability of a neuron

during a given recording session. *Top*: Example histogram of output spikes occurring after input

spikes, fitted (red curve) with a Gaussian distribution N(μ,σ, baseline), where μ is the mean

delay and σ is the jitter of this delay. Lag 0 ms corresponds to the input spike time. In this

example, output spikes are generated on average 16 ms after a stimulation impulse (delay) with a

jitter (σ) of 8.7 ms. *Bottom***:** the baseline is subtracted and the histogram divided by the number

of input spikes during the recording session. This gives the distribution of the probability of

spiking after an input spike, the sum of probabilities defining the spiking probability (SP) of the

cell during the recording session. Here the neuron fires 39 % of the time after an input spike.

(B-D) Delay, jitter and spiking probability (SP) distributions as a function of input sets, for (B)

the original recordings, (C) the simulations and (D) the shuffled data set. Dashed horizontal red

lines are means.

1006

Figure S2 related to Figure 7. Spike-wise neural noise characteristics are not good predictors of spiketrain decorrelation by single GCs.

Plots of the normalized decorrelation, i.e. $(R_{input} - R_{output}) / R_{input}$, of each recording set ($\tau_w = 10$ ms) : 102 for GC original and shuffled recordings **(A-B)**, 20 for FS **(C)**, and for GC, FS and HMC pooled together (**D**) as a function of spike-wise noise characteristics (spike delay, jitter and probability). Solid green lines are the best linear fit, with R^2 and p-values noted in each panel. These plots illustrate **Table 2**. Note that decorrelation is poorly explained (low R^2) by either the spike delay or its jitter in all cell-types. In contrast, the spiking probability (SP) is a good predictor of decorrelation in shuffled GC recordings (i.e. recordings entirely dominated by spike-wise noise) and even more so in FS recordings (for FS, SP was computed from nbFS data. See Figure S7). This suggests that a low SP can be a potent mechanism for decorrelation, and that FS show different levels of decorrelation than GCs partly because they are more reliable. However, the regression line for FS is lower than for GCs (even FS with low SP show less decorrelation than GCs with similar SP), and SP is only a mediocre predictor of decorrelation for GCs, thus confirming that temporal pattern separation in single GCs cannot be the result of simple neural noise.

1025

Figure S3 related to Figure 7. Spike-wise noise characteristics of granule cells are different than those of DG interneurons.

Spike delay, jitter and spiking probability for FS interneurons compared to GC recordings associated with the same input sets than FS (*top*) (20 FS recordings, 61 GC recordings, see Figure 3A) and HMC interneurons compared to a different set of GC recordings associated with the same input sets than HMC (*bottom*) (18 HMC recordings, 22 GC recordings, see Figure 3B). A U-test was applied to each pair of comparison, showing that FS are much faster and less noisy than GCs, whereas HMC are similar to GCs (slightly larger jitter, but slightly higher SP).

1034

1035

Figure S4 related to Figure 7. R_{output} and spiketrain reliability are lower for a random spike

generator than for GCs.

Simulations of output spiketrains to the different input sets with random spiking following a

Gaussian distribution defined by the mean spike delay, mean jitter and mean reliability of the

original recordings.

(A) R_{output} distribution at $\tau_W = 10$ ms, for simulated data and original data. Distributions are fitted

with parabolae and significantly different (F-test using parabolic regressions: $p < 0.0001$).

(B) R_w distributions are significantly different (unpaired T-test, $p < 0.0001$; $<Rw>\{simul\}$ =

0.14).

(C) Like in the original data (**Fig 2E**), the average normalized decorrelation ((R_{input}-R_{output})/R_{input})

seems invariant. Bars are SEM.

(A-B) Asterisks signify statistical significance $p < 0.05$.

1048

1049

Figure S5 related to Figure 8A. Spiketrain reliability in different hippocampal celltypes

FS have a higher R_w than GCs (n = 20 vs 61 recording sets, unpaired T-test: $p < 0.0001$). Note

that when comparing only the simultaneous FS and GC recordings (dark green), we found a

similarly significant difference. HMCs R_w is only slightly higher than GCs (n = 18 vs 22

recording sets, unpaired T-test: $p = 0.0963$). Under partial block of inhibition, CA3 pyramidal

cells have a lower R_w than GCs (n = 15 vs 22 recording sets, unpaired T-test: $p = 0.0052$).

Asterisks denote $p < 0.05$

1057

1058

Figure S6 related to Figure 3. DG interneurons fire bursts and have a higher firing rate than GCs

(A-B) Comparison of the firing rate in all celltypes and conditions. **(A)** For each comparison the GC dataset is different and matched to the non-GC dataset, as in Figure 3-4. FS and HMC have a higher firing rate than their GC control (U-test: $p < 0.0001$ and $p = 0.0005$ respectively) whereas CA3 and GC recordings under gabazine and 30 Hz input trains do not differ (U-test: $p = 0.17$).

(B) Cumulative distributions of the firing rate for all recordings of all celltypes (n = 102 GC, 20 FS, 18 HMC, 15 CA3+gzn, 22 GC+gzn).

(C-D) Same as A-B for the probability of bursting in a recording. Bursting was defined as the

occurrence of at least two action potentials between two input pulses. **(C)** FS and HMC have a

higher propensity to fire bursts than their GC control (U-test: $p < 0.0001$) whereas CA3

pyramidal cells and their GC control do not differ (U-test: $p = 0.74$).

(E) The distribution of the number of spikes between two input pulses, for all celltypes, show

that FS are the only neurons that consistently fire large bursts. Note that GCs almost never fire

more than twice, and when they do it generally corresponds to two preceding input pulses

occurring close to each other (the first spike being in reality associated to a different input than

the next spike).

1076

1077

Figure S7 related to Figure 3 and 5. Differences in pattern separation between GC and DG

interneurons are not solely due to different firing rates and bursting behaviors.

(**A**) The relationship between the firing rate and decorrelation levels in all recordings of all

celltypes. In contrast to GCs, for FS neurons there is a strong correlation between the firing rate

of a recording set and the associated normalized decorrelation. See **Table 3** for more details.

(**B**) The relationship between the probability of bursting (see Figure S6) and the decorrelation

levels in all recordings of all celltypes is less clear, but cells bursting more than 30% of the time

have the lowest decorrelation levels.

(**C**) Example of bursts in a FS recording (*bottom*) in response to input spikes (*top*). To assess the

effect of bursting on R_{output}, we truncated each recorded spiketrain from FS neurons to keep only

the first output spike between two input spikes, thus removing any burst without altering the SP

of the cell. The blue shaded areas highlight the spikes that were removed. The resulting truncated

dataset was termed "nbFS" for "non-burst FS".

(**D**) R_{output} versus R_{input} for nbFS and GC at $\tau_w = 10$ms (to compare to Figure 3A4). Both

distributions are still significantly different, suggesting the bursting behavior of FS is not

sufficient to explain the difference in pattern separation (Unbalanced two-way ANOVA.

Stimulation groups: $p = 0.002$, cell-types: $p < 0.0001$, interaction: $p = 0.84$. Post-hoc Tukey-

Kramer tests: nbFS and GC R_{output} are significantly different ($p < 0.05$) for stimulation groups

with the three highest R_{input})

(**E**) Bursts in HMC recordings were truncated to produce an "nbHMC" dataset.

(**F**) Pairwise analysis on nbHMC and GC recordings at $\tau_w = 250$ ms show that the distributions

are still different between the two celltypes (to compare to Figure 5D). ANCOVA: $p < 0.0001$;

two-way ANOVA: input sets: $p < 0.0001$, cell-types: $p < 0.0001$, interaction: $p < 0.0001$. Note

that under this analysis, nbHMCs and GCs are also significantly different at lower timescales (τ_w

= 100 and 50 ms) but with a lower effect size as τ_w decreases.

1103

Table 1-3. **Linear regressions goodness-of-fit, p-value and slope.** The predictor variables (x-axis)
correspond to columns, and the variables to be explained (y-axis) correspond to rows. Red highlights
significant regressions that explain more than 50% of the variance ($R^2 > 50\%$). Blue highlights
regressions that are significant ($p < 0.01$) but that explain less than 50% of the variance. The values used
for Normalized Decorrelation, i.e. (R_{input} - R_{output})/ R_{input}, and for Spiketrain Reliability (Rw) were
computed with a binning window of 10 ms, unless specified. Abbreviations: GC yo = from young mice,
GC ad = from adult mice, ALL = dataset pooling all celltypes and conditions recorded in young mice
1111

Table 1. Intrinsic electrophysiological cell properties

y-axis ↓ / x-axis →		Membrane Capacitance (Cm)	Membrane Resistance (Rm)	Membrane Time Constant = Rm.Cm	Resting Membrane Potential (Vrest)
Normalized Decorrelation	**GC yo**	$R^2 = 4\%$ p = 0.08 slope = -0.2	$R^2 = 5\%$ p = 0.06 slope = -0.03	$R^2 = 8\%$ p = 0.013 slope = -1.2	$R^2 = 3\%$ p = 0.17 slope = -0.2
	FS	$R^2 = 47\%$ p = 0.0008 slope = -1.5	$R^2 = 77\%$ p <0.0001 slope = 0.6	$R^2 = 5\%$ p = 0.4 slope = 13.6	$R^2 = 46\%$ p = 0.0009 slope = 1.4
	HMC	$R^2 = 33\%$ p = 0.013 slope = -0.4	$R^2 = 3\%$ p = 0.48 slope = 0.06	$R^2 = 7\%$ p = 0.3 slope = -1.8	$R^2 = 0.5\%$ p = 0.8 slope = 0.1
	CA3 +gzn	$R^2 = 0.1\%$ p = 0.9 slope = -0.05	$R^2 = 2\%$ p = 0.59 slope = 0.03	$R^2 = 0\%$ p = 1 slope = 1.2	$R^2 = 29\%$ p = 0.04 slope = 0.7
	GC +gzn	$R^2 = 30\%$ p = 0.008 slope = 1.6	$R^2 = 0.3\%$ p = 0.8 slope = 0.01	$R^2 = 2\%$ p = 0.5 slope = 3.5	$R^2 = 13\%$ p = 0.09 slope = -0.7
	ALL	$R^2 = 0.6\%$ p = 0.35 slope = -0.09	$R^2 = 3\%$ p = 0.04 slope = 0.03	$R^2 = 1\%$ p = 0.2 slope = 0.55	$R^2 = 0.2\%$ p = 0.55 slope = -0.08
	GC ad	$R^2 = 4\%$ p = 0.24 slope = -0.4	$R^2 = 10\%$ p = 0.06 slope = -0.07	$R^2 = 0.6\%$ p = 0.7 slope = -4.3	$R^2 = 5\%$ p = 0.2 slope = -0.3
Spiketrain Reliability (R_W)	**GC**	$R^2 = 2\%$ p =0.2 slope = 0.001	$R^2 = 5\%$ p = 0.03 slope = 4e-3	$R^2 = 7\%$ p = 0.02 slope = 0.01	$R^2 = 3\%$ p = 0.17 slope = 0.002
	FS	$R^2 = 48\%$ p =0.0006, slope = 0.01	$R^2 = 70\%$ p < 0.0001 slope = -0.007	$R^2 = 6\%$ p = 0.29 slope = -0.12	$R^2 = 39\%$ p = 0.003 slope = -0.014
	HMC	$R^2 = 29\%$ p = 0.19 slope = 0.005	$R^2 = 24\%$ p = 0.06 slope = -0.001	$R^2 = 9\%$ p = 0.2 slope = 0.006	$R^2 = 3\%$ p = 0.5 slope = -0.002
	CA3 +gzn	$R^2 = 4\%$ p = 0.5 slope = 0.003	$R^2 = 9\%$ p = 0.29 slope = 6e-4	$R^2 = 4\%$ p = 0.45 slope = -0.01	$R^2 = 10\%$ p = 0.24 slope = -0.004
	GC +gzn	$R^2 = 2\%$ p = 0.5 slope = -0.003	$R^2 = 22\%$ p = 0.03 slope = 8e-4	$R^2 = 17\%$ p = 0.05 slope = -0.02	$R^2 = 29\%$ p = 0.01 slope = 0.008
	ALL	$R^2 = 0.4\%$ p = 0.5 slope = 7e-4	$R^2 = 7\%$ p = 0.0013 slope = -4e-4	$R^2 = 3\%$ p = 0.045 slope = -0.008	$R^2 = 1\%$ p = 0.24 slope = 0.001
	GC ad	$R^2 = 13\%$ p =0.03 slope = 0.006	$R^2 = 6\%$ p = 0.17 slope = 4e-4	$R^2 = 17 \%$ p = 0.01 slope = 0.04	$R^2 = 8.5\%$ p = 0.09 slope = 0.004

Table 2. Spike-wise neural noise

y-axis ↓	x-axis →	Delay	Jitter	Spike Probability (SP)
Normalized Decorrelation	**GC**	$R^2 = 39.5\%$ $p < 0.0001$ slope = 1.8	$R^2 = 43\%$ $p < 0.0001$ slope = 1.3	$R^2 = 45\%$ $p < 0.0001$ slope = -47.3
	FS	$R^2 = 13\%$ $p = 0.1$ slope = 2.2	$R^2 = 22\%$ $p = 0.04$ slope = 9.2	$R^2 = 88\%$ $p < 0.0001$ slope = -42.4
	HMC	$R^2 = 9\%$ $p = 0.21$ slope = 2.1	$R^2 = 18\%$ $p = 0.08$ slope = 3.4	$R^2 = 34\%$ $p = 0.011$ slope = -42.0
	ALL	$R^2 = 35\%$ $p < 0.0001$ slope = 1.9	$R^2 = 23\%$ $p < 0.0001$ slope = 1.4	$R^2 = 57\%$ $p < 0.0001$ slope = -50.6
	GC ad	$R^2 = 2\%$ $p = 0.38$ slope = 0.9	$R^2 = 4\%$ $p = 0.23$ slope = -3.5	$R^2 = 42\%$ $p < 0.0001$ slope = -45.7
	Shuffle (GC yo)	$R^2 = 11\%$ $p = 0.02$ slope = 0.9	$R^2 = 24\%$ $p = 0.0003$ slope = 0.8	$R^2 = 65\%$ $p < 0.0001$ slope = -41.1
Spiketrain Reliability (R_W)	**GC**	$R^2 = 33\%$ $p < 0.0001$ slope = -0.016	$R^2 = 43\%$ $p < 0.0001$ slope = -0.013	$R^2 = 46\%$ $p < 0.0001$ slope = 0.5
	FS	$R^2 = 15\%$ $p = 0.09$ slope = -0.024	$R^2 = 15\%$ $p = 0.09$ slope = -0.078	$R^2 = 85\%$ $p < 0.0001$ slope = 0.4
	HMC	$R^2 = 35\%$ $p = 0.009$ slope = -0.030	$R^2 = 53\%$ $p = 0.0007$ slope = -0.043	$R^2 = 31\%$ $p = 0.016$ slope = 0.3
	ALL	$R^2 = 40\%$ $p < 0.0001$ slope = -0.020	$R^2 = 28\%$ $p < 0.0001$ slope = -0.016	$R^2 = 61\%$ $p < 0.0001$ slope = 0.5
	GC ad	$R^2 = 7\%$ $p = 0.14$ slope = -0.01	$R^2 = 0.03\%$ $p = 0.9$ slope = 0.003	$R^2 = 16\%$ $p = 0.017$ slope = 0.2
	Shuffle (GC yo)	$R^2 = 9\%$ $p = 0.04$ slope = -0.008	$R^2 = 23\%$ $p = 0.0004$ slope = -0.008	$R^2 = 65\%$ $p < 0.0001$ slope = 0.4

Table 3. Spiketrain-wise properties

		Overall Firing Rate	Spiketrain Reliability (R_W)
	GC	$R^2 = 15\%$ $p < 0.0001$ slope = - 1.3	$R^2 = 81\%$ $p < 0.0001$ slope = -87
	FS	$R^2 = 65\%$ $p < 0.0001$ slope = - 0.5	$R^2 = 90\%$ $p < 0.0001$ slope = -85
	HMC	$R^2 = 35\%$ $p = 0.0010$ slope = - 2.8	$R^2 = 61\%$ $p = 0.0001$ slope = -105
Normalized Decorrelation	**CA3 +gzn**	$R^2 = 69\%$ $p = 0.0001$ slope = - 1.7	$R^2 = 55\%$ $p = 0.0016$ slope = -76
	GC +gzn	$R^2 = 0.6\%$ $p = 0.73$ slope = - 0.2	$R^2 = 28\%$ $p = 0.0111$ slope = -75
	ALL	$R^2 = 37\%$ $p < 0.0001$ slope = - 0.8	$R^2 = 79\%$ $p < 0.0001$ slope = -91
	GC ad	$R^2 = 30\%$ $p = 0.0006$ slope = - 4.1	$R^2 = 52\%$ $p < 0.0001$ slope = -90

y-axis ↓   x-axis ➔

1128 References
1129

Aimone, J. B., Deng, W., & Gage, F. H. (2011). Resolving new memories: a critical look at the dentate
gyrus, adult neurogenesis, and pattern separation. *Neuron, 70*(4), 589-596. doi:
10.1016/j.neuron.2011.05.010
Aimone, J. B., Wiles, J., & Gage, F. H. (2009). Computational influence of adult neurogenesis on memory
encoding. *Neuron, 61*(2), 187-202. doi: 10.1016/j.neuron.2008.11.026
Amaral, D. G., Scharfman, H. E., & Lavenex, P. (2007). The dentate gyrus: fundamental neuroanatomical
organization (dentate gyrus for dummies). *163*, 3-790. doi: 10.1016/s0079-6123(07)63001-5
Andersen, P., Holmqvist, B., & Voorhoeve, P. E. (1966). Excitatory synapses on hippocampal apical
dendrites activated by entorhinal stimulation. *Acta Physiol Scand, 66*(4), 461-472. doi:
10.1111/j.1748-1716.1966.tb03224.x
Baker, S., Vieweg, P., Gao, F., Gilboa, A., Wolbers, T., Black, S. E., & Rosenbaum, R. S. (2016). The Human
Dentate Gyrus Plays a Necessary Role in Discriminating New Memories. *Curr Biol*. doi:
10.1016/j.cub.2016.07.081
Bischofberger, J., Engel, D., Li, L., Geiger, J. R., & Jonas, P. (2006). Patch-clamp recording from mossy
fiber terminals in hippocampal slices. *Nat Protoc, 1*(4), 2075-2081. doi: 10.1038/nprot.2006.312
Buzsaki, G. (2010). Neural syntax: cell assemblies, synapsembles, and readers. *Neuron, 68*(3), 362-385.
doi: 10.1016/j.neuron.2010.09.023
Chavlis, S., Petrantonakis, P. C., & Poirazi, P. (2017). Dendrites of dentate gyrus granule cells contribute
to pattern separation by controlling sparsity. *Hippocampus*.
Chavlis, S., & Poirazi, P. (2017). Pattern separation in the hippocampus through the eyes of
computational modeling. *Synapse*. doi: 10.1002/syn.21972
Danielson, N. B., Kaifosh, P., Zaremba, J. D., Lovett-Barron, M., Tsai, J., Denny, C. A., . . . Hen, R. (2016).
Distinct contribution of adult-born hippocampal granule cells to context encoding. *Neuron,
90*(1), 101-112.
Danielson, N. B., Turi, G. F., Ladow, M., Chavlis, S., Petrantonakis, P. C., Poirazi, P., & Losonczy, A. (2017).
In Vivo Imaging of Dentate Gyrus Mossy Cells in Behaving Mice. *Neuron, 93*(3), 552-559. e554.
Deng, W., Mayford, M., & Gage, F. H. (2013). Selection of distinct populations of dentate granule cells in
response to inputs as a mechanism for pattern separation in mice. *Elife, 2*, e00312. doi:
10.7554/eLife.00312
Denny, C. A., Kheirbek, M. A., Alba, E. L., Tanaka, K. F., Brachman, R. A., Laughman, K. B., . . . Hen, R.
(2014). Hippocampal memory traces are differentially modulated by experience, time, and adult
neurogenesis. *Neuron, 83*(1), 189-201. doi: 10.1016/j.neuron.2014.05.018
Diamantaki, M., Frey, M., Berens, P., Preston-Ferrer, P., & Burgalossi, A. (2016). Sparse activity of
identified dentate granule cells during spatial exploration. *Elife, 5*. doi: 10.7554/eLife.20252
Dieni, C. V., Panichi, R., Aimone, J. B., Kuo, C. T., Wadiche, J. I., & Overstreet-Wadiche, L. (2016). Low
excitatory innervation balances high intrinsic excitability of immature dentate neurons. *Nat
Commun, 7*, 11313. doi: 10.1038/ncomms11313
Dillon, S. E., Tsivos, D., Knight, M., McCann, B., Pennington, C., Shiel, A. I., . . . Coulthard, E. J. (2017). The
impact of ageing reveals distinct roles for human dentate gyrus and CA3 in pattern separation
and object recognition memory. *Sci Rep, 7*(1), 14069. doi: 10.1038/s41598-017-13853-8
Dobrunz, L. E., & Stevens, C. F. (1999). Response of hippocampal synapses to natural stimulation
patterns. *Neuron, 22*(1), 157-166.
Eichenbaum, H. (2014). Time cells in the hippocampus: a new dimension for mapping memories. *Nat
Rev Neurosci, 15*(11), 732-744. doi: 10.1038/nrn3827
Ewell, L. A., & Jones, M. V. (2010). Frequency-tuned distribution of inhibition in the dentate gyrus. *J
Neurosci, 30*(38), 12597-12607. doi: 10.1523/jneurosci.1854-10.2010

Faisal, A. A., Selen, L. P., & Wolpert, D. M. (2008). Noise in the nervous system. *Nat Rev Neurosci, 9*(4), 292-303. doi: 10.1038/nrn2258

Fontanari, J., Rolls, E., & Costa, D. F. (1995). A model of the operation of the hippocampus and entorhinal cortex in memory. *International Journal of Neural Systems, 6*(1), 51-70.

Friedrich, R. W., & Wiechert, M. T. (2014). Neuronal circuits and computations: pattern decorrelation in the olfactory bulb. *FEBS Lett, 588*(15), 2504-2513. doi: 10.1016/j.febslet.2014.05.055

Fuhrmann, G., Segev, I., Markram, H., & Tsodyks, M. (2002). Coding of temporal information by activity-dependent synapses. *J Neurophysiol, 87*(1), 140-148.

GoodSmith, D., Chen, X., Wang, C., Kim, S. H., Song, H., Burgalossi, A., . . . Knierim, J. J. (2017). Spatial representations of granule cells and mossy cells of the dentate gyrus. *Neuron, 93*(3), 677-690. e675.

Guzman, S. J., Schlögl, A., Frotscher, M., & Jonas, P. (2016). Synaptic mechanisms of pattern completion in the hippocampal CA3 network. *Science, 353*(6304), 1117-1123.

Hananeia, N., & Benuskova, L. (2016). Computational simulation of dentate gyrus granule cell—The role of metaplasticity. *Neurocomputing, 175*, 300-309.

Harney, S. C., & Jones, M. V. (2002). Pre- and postsynaptic properties of somatic and dendritic inhibition in dentate gyrus. *Neuropharmacology, 43*(4), 584-594.

Hu, H., Gan, J., & Jonas, P. (2014). Interneurons. Fast-spiking, parvalbumin(+) GABAergic interneurons: from cellular design to microcircuit function. *Science, 345*(6196), 1255263. doi: 10.1126/science.1255263

Hyde, R. A., & Strowbridge, B. W. (2012). Mnemonic representations of transient stimuli and temporal sequences in the rodent hippocampus in vitro. *Nat Neurosci, 15*(10), 1430-1438. doi: 10.1038/nn.3208

Izhikevich, E. M. (2003). Simple model of spiking neurons. *IEEE Trans Neural Netw, 14*(6), 1569-1572. doi: 10.1109/tnn.2003.820440

Jackson, M. B. (2013). Recall of spatial patterns stored in a hippocampal slice by long-term potentiation. *J Neurophysiol, 110*(11), 2511-2519. doi: 10.1152/jn.00533.2013

Jones, M. V., Jonas, P., Sahara, Y., & Westbrook, G. L. (2001). Microscopic kinetics and energetics distinguish GABA A receptor agonists from antagonists. *Biophysical journal, 81*(5), 2660-2670.

Kelemen, E., & Fenton, A. A. (2016). Coordinating different representations in the hippocampus. *Neurobiol Learn Mem, 129*, 50-59. doi: 10.1016/j.nlm.2015.12.011

Kesner, R. P., Kirk, R. A., Yu, Z., Polansky, C., & Musso, N. D. (2016). Dentate gyrus supports slope recognition memory, shades of grey-context pattern separation and recognition memory, and CA3 supports pattern completion for object memory. *Neurobiol Learn Mem, 129*, 29-37. doi: 10.1016/j.nlm.2015.08.010

Kesner, R. P., & Rolls, E. T. (2015). A computational theory of hippocampal function, and tests of the theory: new developments. *Neurosci Biobehav Rev, 48*, 92-147. doi: 10.1016/j.neubiorev.2014.11.009

Larimer, P., & Strowbridge, B. W. (2008). Nonrandom local circuits in the dentate gyrus. *J Neurosci, 28*(47), 12212-12223. doi: 10.1523/jneurosci.3612-08.2008

Larimer, P., & Strowbridge, B. W. (2010). Representing information in cell assemblies: persistent activity mediated by semilunar granule cells. *Nat Neurosci, 13*(2), 213-222. doi: 10.1038/nn.2458

Leutgeb, J. K., Leutgeb, S., Moser, M. B., & Moser, E. I. (2007). Pattern separation in the dentate gyrus and CA3 of the hippocampus. *Science, 315*(5814), 961-966. doi: 10.1126/science.1135801

Lisman, J. E., & Jensen, O. (2013). The theta-gamma neural code. *Neuron, 77*(6), 1002-1016. doi: 10.1016/j.neuron.2013.03.007

Macke, J. H., Berens, P., Ecker, A. S., Tolias, A. S., & Bethge, M. (2009). Generating spike trains with
specified correlation coefficients. *Neural Comput, 21*(2), 397-423. doi: 10.1162/neco.2008.02-
08-713

Mainen, Z. F., & Sejnowski, T. J. (1995). Reliability of spike timing in neocortical neurons. *Science,*
268(5216), 1503-1506.

Marrone, D. F., Adams, A. A., & Satvat, E. (2011). Increased pattern separation in the aged fascia
dentata. *Neurobiol Aging, 32*(12), 2317 e2323-2332. doi: 10.1016/j.neurobiolaging.2010.03.021

Marrone, D. F., Satvat, E., Odintsova, I. V., & Gheidi, A. (2014). Dissociation of spatial representations
within hippocampal region CA3. *Hippocampus*, n/a-n/a. doi: 10.1002/hipo.22367

McHugh, T. J., Jones, M. W., Quinn, J. J., Balthasar, N., Coppari, R., Elmquist, J. K., . . . Tonegawa, S.
(2007). Dentate gyrus NMDA receptors mediate rapid pattern separation in the hippocampal
network. *Science, 317*(5834), 94-99. doi: 10.1126/science.1140263

McNaughton, B. L., & Morris, R. G. M. (1987). Hippocampal synaptic enhancement and information
storage within a distributed memory system. *Trends in Neuroscience, 10*(10), 408-415.

Mongiat, L. A., Esposito, M. S., Lombardi, G., & Schinder, A. F. (2009). Reliable activation of immature
neurons in the adult hippocampus. *PLoS One, 4*(4), e5320. doi: 10.1371/journal.pone.0005320

Mori, M., Abegg, M. H., Gahwiler, B. H., & Gerber, U. (2004). A frequency-dependent switch from
inhibition to excitation in a hippocampal unitary circuit. *Nature, 431*(7007), 453-456. doi:
10.1038/nature02854

Motulsky, H. J., & Ransnas, L. A. (1987). Fitting curves to data using nonlinear regression: a practical and
nonmathematical review. *FASEB J, 1*(5), 365-374.

Myers, C. E., & Scharfman, H. E. (2009). A role for hilar cells in pattern separation in the dentate gyrus: a
computational approach. *Hippocampus, 19*(4), 321-337. doi: 10.1002/hipo.20516

Nakazawa, K. (2017). Dentate Mossy Cell and Pattern Separation. *Neuron, 93*(3), 465-467.

Neher, T., Cheng, S., & Wiskott, L. (2015). Memory storage fidelity in the hippocampal circuit: the role of
subregions and input statistics. *PLoS Comput Biol, 11*(5), e1004250. doi:
10.1371/journal.pcbi.1004250

Neunuebel, J. P., & Knierim, J. J. (2012). Spatial firing correlates of physiologically distinct cell types of
the rat dentate gyrus. *J Neurosci, 32*(11), 3848-3858. doi: 10.1523/JNEUROSCI.6038-11.2012

Neunuebel, J. P., & Knierim, J. J. (2014). CA3 retrieves coherent representations from degraded input:
direct evidence for CA3 pattern completion and dentate gyrus pattern separation. *Neuron,*
81(2), 416-427. doi: 10.1016/j.neuron.2013.11.017

O'Reilly, R. C., & McClelland, J. L. (1994). Hippocampal conjunctive encoding, storage, and recall:
avoiding a trade-off. *Hippocampus, 4*(6), 661-682. doi: 10.1002/hipo.450040605

Pardi, M. B., Ogando, M. B., Schinder, A. F., & Marin-Burgin, A. (2015). Differential inhibition onto
developing and mature granule cells generates high-frequency filters with variable gain. *Elife, 4*,
e08764. doi: 10.7554/eLife.08764

Pernía-Andrade, Alejandro J., & Jonas, P. (2014). Theta-Gamma-Modulated Synaptic Currents in
Hippocampal Granule Cells In Vivo Define a Mechanism for Network Oscillations. *Neuron, 81*(1),
140-152. doi: 10.1016/j.neuron.2013.09.046

Ramirez, S., Liu, X., Lin, P. A., Suh, J., Pignatelli, M., Redondo, R. L., . . . Tonegawa, S. (2013). Creating a
false memory in the hippocampus. *Science, 341*(6144), 387-391. doi: 10.1126/science.1239073

Rebola, N., Carta, M., & Mulle, C. (2017). Operation and plasticity of hippocampal CA3 circuits:
implications for memory encoding. *Nat Rev Neurosci, 18*(4), 208-220. doi: 10.1038/nrn.2017.10

Rieke, F., Warland, W., de Ruyter van Steveninck, R., & Bialek, W. (1999). *Spikes: Exploring the Neural*
Code: A Bradford Book.

Rolls, E. T. (2010). A computational theory of episodic memory formation in the hippocampus. *Behav*
Brain Res, 215(2), 180-196. doi: 10.1016/j.bbr.2010.03.027

Sachidhanandam, S., Blanchet, C., Jeantet, Y., Cho, Y. H., & Mulle, C. (2009). Kainate receptors act as
conditional amplifiers of spike transmission at hippocampal mossy fiber synapses. *J Neurosci,*
29(15), 5000-5008. doi: 10.1523/jneurosci.5807-08.2009
Santoro, A. (2013). Reassessing pattern separation in the dentate gyrus. *Front Behav Neurosci, 7*, 96. doi:
10.3389/fnbeh.2013.00096
Savanthrapadian, S., Meyer, T., Elgueta, C., Booker, S. A., Vida, I., & Bartos, M. (2014). Synaptic
Properties of SOM- and CCK-Expressing Cells in Dentate Gyrus Interneuron Networks. *J Neurosci,*
34(24), 8197-8209. doi: 10.1523/JNEUROSCI.5433-13.2014
Scharfman, H. E. (2016). The enigmatic mossy cell of the dentate gyrus. *Nat Rev Neurosci, 17*(9), 562-
575. doi: 10.1038/nrn.2016.87
Schmidt, B., Marrone, D. F., & Markus, E. J. (2012). Disambiguating the similar: the dentate gyrus and
pattern separation. *Behav Brain Res, 226*(1), 56-65. doi: 10.1016/j.bbr.2011.08.039
Senzai, Y., & Buzsáki, G. (2017). Physiological properties and behavioral correlates of hippocampal
granule cells and mossy cells. *Neuron, 93*(3), 691-704. e695.
Severa, W., Parekh, O., James, C. D., & Aimone, J. B. (2016). A Combinatorial Model for Dentate Gyrus
Sparse Coding. *Neural Comput*, 1-24. doi: 10.1162/NECO_a_00905
Sun, Q., Sotayo, A., Cazzulino, A. S., Snyder, A. M., Denny, C. A., & Siegelbaum, S. A. (2017).
Proximodistal Heterogeneity of Hippocampal CA3 Pyramidal Neuron Intrinsic Properties,
Connectivity, and Reactivation during Memory Recall. *Neuron, 95*(3), 656-672 e653. doi:
10.1016/j.neuron.2017.07.012
Temprana, S. G., Mongiat, L. A., Yang, S. M., Trinchero, M. F., Alvarez, D. D., Kropff, E., . . . Schinder, A. F.
(2015). Delayed coupling to feedback inhibition during a critical period for the integration of
adult-born granule cells. *Neuron, 85*(1), 116-130. doi: 10.1016/j.neuron.2014.11.023
Teyler, T. J., & Rudy, J. W. (2007). The hippocampal indexing theory and episodic memory: updating the
index. *Hippocampus, 17*(12), 1158-1169. doi: 10.1002/hipo.20350
Treves, A., Tashiro, A., Witter, M. P., & Moser, E. I. (2008). What is the mammalian dentate gyrus good
for? *Neuroscience, 154*(4), 1155-1172. doi: 10.1016/j.neuroscience.2008.04.073
VanRullen, R., Guyonneau, R., & Thorpe, S. J. (2005). Spike times make sense. *Trends Neurosci, 28*(1), 1-
4. doi: 10.1016/j.tins.2004.10.010
Vyleta, N. P., Borges-Merjane, C., & Jonas, P. (2016). Plasticity-dependent, full detonation at
hippocampal mossy fiber–CA3 pyramidal neuron synapses. *Elife, 5*.
Wick, S. D., Wiechert, M. T., Friedrich, R. W., & Riecke, H. (2010). Pattern orthogonalization via channel
decorrelation by adaptive networks. *J Comput Neurosci, 28*(1), 29-45. doi: 10.1007/s10827-009-
0183-1
Wikenheiser, A. M., & Redish, A. D. (2015). Decoding the cognitive map: ensemble hippocampal
sequences and decision making. *Curr Opin Neurobiol, 32*, 8-15.
Yi, F., Catudio-Garrett, E., Gabriel, R., Wilhelm, M., Erdelyi, F., Szabo, G., . . . Lawrence, J. (2015).
Hippocampal "cholinergic interneurons" visualized with the choline acetyltransferase promoter:
anatomical distribution, intrinsic membrane properties, neurochemical characteristics, and
capacity for cholinergic modulation. *Front Synaptic Neurosci, 7*, 4. doi:
10.3389/fnsyn.2015.00004
Yim, M. Y., Hanuschkin, A., & Wolfart, J. (2014). Intrinsic rescaling of granule cells restores pattern
separation ability of a dentate gyrus network model during epileptic hyperexcitability.
Hippocampus. doi: 10.1002/hipo.22373
Zylberberg, J., Hyde, R. A., & Strowbridge, B. W. (2016). Dynamics of robust pattern separability in the
hippocampal dentate gyrus. *Hippocampus, 26*(5), 623-632. doi: 10.1002/hipo.22546
1316
1317