1   **Title:** Pattern separation of spiketrains in hippocampal neurons
2
3   **Short Title**: Temporal pattern separation in the hippocampus
4

5   **Authors:** Antoine D. Madar*[#1,2], Laura A. Ewell[#1,3], Mathew V. Jones[1]
6   * Corresponding author: madar@wisc.edu, madar@uchicago.edu
7   # contributed equally to the manuscript.
8
9   **Authors affiliations**
10 [1]Department of Neuroscience, University of Wisconsin-Madison, WI 53705, USA
11 [2]Department of Neurobiology, Neuroscience Institute, University of Chicago, IL 60637, USA
12 [3]Institute of Experimental Epileptology and Cognition Research, University of Bonn, Germany
13

Author contributions
Conceptualization: MVJ, LAE. Data curation: LAE, ADM. Formal analysis: MVJ, ADM.
Funding acquisition: MVJ, ADM. Investigation/data collection: LAE, ADM. Methodology:
MVJ, LAE, ADM. Project administration: MVJ, LAE, ADM. Resources: N/A. Software: MVJ,
ADM. Supervision: MVJ. Validation: MVJ, LAE, ADM. Visualization: MVJ, LAE, ADM.
Writing – original draft: ADM. Writing – review & editing: MVJ, LAE, ADM.
20

21 **Acknowledgements**
22 We would like to thank Jesse Pfammatter and Linda Overstreet-Wadiche for many helpful
23 discussions and feedback on early versions of the manuscript.
24

25 **Funding**
26 This work was supported by the University of Wisconsin Institute for Clinical and Translational
27 Research (M.V.J.; NIH/NCATS UL1TR000427) and Lily's Fund for Epilepsy Research
28 (A.D.M.; 2015 fellow).
29

30 **Competing interest statement**
31 The authors declare no financial or non-financial competing interests in relation to the present
32 work.
33

Abstract

Pattern separation is a process that minimizes overlap between patterns of neuronal activity representing similar experiences. Theoretical work suggests that the dentate gyrus (DG) performs this role for memory processing but a direct demonstration is lacking. One limitation is the difficulty to measure DG inputs and outputs simultaneously. To rigorously assess pattern separation by DG circuitry, we used mouse brain slices to stimulate DG afferents and simultaneously record DG granule cells (GCs) and interneurons. Output spiketrains of GCs are more dissimilar than their input spiketrains, demonstrating for the first time temporal pattern separation at the level of single neurons in the DG. Pattern separation is larger in GCs than in fast-spiking interneurons and hilar mossy cells, and is amplified in CA3 pyramidal cells. Analysis of the neural noise and computational modelling suggest that this form of pattern separation is not explained by simple randomness and arises from specific presynaptic dynamics. Overall, by reframing the concept of pattern separation in dynamic terms and by connecting it to the physiology of different types of neurons, our study offers a new window of understanding in how hippocampal networks might support episodic memory.

Introduction

How does the brain allow us to discriminate between similar events in our past? This question is a central challenge in the neurobiology of memory and remains elusive. To prevent confusion between memories that share similar features, the brain needs to store distinct activity patterns to represent distinct memories. In the influential Hebb-Marr framework of episodic memory[1,2], representations are stored in area CA3 of hippocampus, an auto-associative network where plastic recurrent excitatory connections facilitate recall of stored patterns in response to partial

cues[3,4]. However, strong recurrent excitation severely limits the number of patterns that can be

stored without overlap[4,5]. Such overlap would lead, when a partial cue common to several

patterns is presented, to the reactivation of many patterns and thus to confusion or confabulation.

To avoid these interferences, the Hebb-Marr framework proposes that redundancy between input

patterns is reduced before they are stored. This process of transforming similar input patterns

into less similar output patterns is termed *pattern separation*[5,6].

Theoretical models suggest that the dentate gyrus (DG) performs pattern separation of

cortical inputs before sending its differentiated outputs to CA3[1,2,4]. Indeed, DG is ideally located

to do this, receiving signals via the major projection from entorhinal cortex (EC), the perforant

path (PP), and sending signals to CA3 via the axons of granule cells (GCs)[7]. In addition,

behavioral studies have shown that DG lesions impair mnemonic discrimination[8-12] and several

experimental reports have shown that similar environments or events are represented differently

in the DG[13-17]. However, this separation of DG representations could be inherited from upstream

structures (e.g. EC) and simply reported by DG. Therefore, a rigorous demonstration that pattern

separation is performed by DG requires simultaneous knowledge of its inputs and outputs[6]. Some

electrophysiological studies suggest that EC spatial representations are on average more

correlated than in DG[13,14,18,19], but the recorded EC neurons were unlikely to contact the recorded

DG neurons, and were not recorded at the same time: a direct test of whether DG itself performs

pattern separation on EC inputs is thus still lacking.

Another difficulty in studying pattern separation is in defining the nature of "activity

patterns". Previous studies have focused on spatial patterns of "active neurons", with little

reference to the dynamics of neural activity. For example, computational models predict that DG

separates overlapping populations of active EC neurons into less overlapping populations of

active GCs[5,20-23]. Immediate-early genes (IEG) expression studies have confirmed that distinct

events drive plasticity in different populations of GCs[15,24] and that overlap in these

representations causes mnemonic interference[25]. In contrast, *in vivo* single-unit recordings in the

DG found that similar contexts are represented by the same population of active neurons, but

differences are encoded by different spatially tuned firing patterns[13,14,17].

These conflicting results show that pattern separation could correspond to different

computations depending on the type of patterns investigated, and that multiple forms of pattern

separation could in theory be implemented by the DG[6]. For example, because *in vivo* recordings

suggest that the same neurons can be used to code different environments[13,14,17], it is possible that

pattern separation is performed at the level of single GCs, each disambiguating the activity

patterns that it receives. Such disambiguation could be done by changing firing rates, or

alternatively, by changing spike timing. Previous experimental investigations of pattern

separation in DG examined population vectors of place fields averaged over minutes[13,14,26], but

place cells also carry information at shorter timescales[27-30]. So far, pattern separation has not

been well characterized on the scale of milliseconds, and never where patterns are explicitly

afferent and efferent trains of action potentials.

Thus, whether the DG network per se reduces the overlap between similar inputs, and

how it performs this computation remains a mystery, especially at short timescales. Here, we set

out to test the hypothesis that the DG performs pattern separation of cortical spiketrains at the

level of single GCs. We designed a novel pattern separation assay in acute brain slices to take

advantage of the experimental control afforded to slice electrophysiology. Complex input

spiketrains of varying similarities were fed into the DG via its afferents, and the output of a

single GC was simultaneously recorded, allowing the first direct measure of pattern separation

103 (by comparing input similarity versus output similarity), on timescales relevant to neuronal

104 encoding and synaptic plasticity. Finally, we explored whether other cell types in the DG and

105 CA3 exhibited this form of pattern separation and investigated the role of neuronal noise and

106 synaptic dynamics in supporting this computation.

Results

Temporal pattern separation by individual dentate granule cells

A direct test of pattern separation in single GCs requires knowledge of the similarity between

input patterns arriving via the PP and comparison with the similarity between GC output

patterns. Here, we define input and output patterns as rasters of spiketrains. The similarity

between two spiketrains was assessed by computing their pairwise Pearson's correlation

coefficient (R) using a binning window τ_w of 10 ms (unless otherwise specified). We generated

sets of Poisson input spiketrains (simulating trains of incoming cortical action potentials), with

each set having an average correlation R_{input} (**Figure 1a** and **Methods – Pattern separation**

experiments). We then recorded the spiking responses of GCs to these sets of input trains

delivered to lateral PP fibers (**Figure 1b-c**) (102 recording sets from 28 GCs in adolescent mice),

allowing us to compute the average output correlation (R_{output}) (**Figure 2a-b**).

For every recording set, R_{output} was lower than the R_{input} of the associated input set,

indicating a decorrelation of the output spiketrains compared to their inputs (**Figure 2c**). This

was also the case in GCs recorded in slices from adult mice (35 recording sets from 14 neurons)

(**Figure 9d**). The effective decorrelation, defined as the difference between R_{input} and R_{output}, was

statistically significant for every input set (**Figure 2d left**), but was larger when input spiketrains

were highly correlated (**Figure 2d**). This is consistent with the role of DG in discriminating

between similar memories more than already dissimilar ones [9]. Note, however, that the

decorrelation normalized to R_{input} is invariant: whatever the input set, the output trains were

always decorrelated to ~70% of R_{input} (**Figure 2e**). Such invariance suggests that the same

decorrelating mechanism is used on all input sets.

These results constitute the first direct experimental demonstration that single GCs, the

output neurons of DG, exhibit temporal pattern separation. A comparison of the spiking patterns

recorded in multiple GCs also shows that different GCs tend to process the same input spiketrain

in different ways (**Figure S1**). Such diversity could support pattern separation at the population

level.

Temporal pattern separation in other cell types of the DG network

Any channel processing inputs and returning outputs, and thus any brain network, performs

either pattern separation or pattern convergence to some degree[6]. Thus, GCs are unlikely to be

the only neurons to exhibit temporal pattern separation of spiketrains. However, we would

expect pattern separation to be at its greatest in GCs, at least among DG cells, because they are

the output neurons of the DG and thus should provide the most separated patterns to CA3 before

they are stored. To test this hypothesis, we performed the same pattern separation assay while

recording from fast-spiking interneurons (FS, 20 recording sets) (**Figure 3a**) or hilar mossy cells

(HMC, 18 recording sets) (**Figure 3b**).

At the 10 ms timescale, the distributions of average (R_{input}, R_{output}) were significantly

different between FSs and GCs, with the R_{output} of simultaneously recorded GCs always lower

than their corresponding FS (**Figure 3a4**):FSs exhibit lower levels of decorrelation than GCs

(**Figure 3a5**). For HMCs, although their current-clamp responses and spiking behavior looked

different from GCs (**Figure 3b2**), the distribution of average (R_{input}, R_{output}) appeared only

slightly different from GCs with this analysis (**Figure 3b3**, but see below, in **Figure 5d**).

Temporal pattern separation in CA3 pyramidal cells

Our FS recordings have shown that not all neurons necessarily exhibit as high levels of temporal

pattern separation as GCs. We thus asked whether this form of pattern separation is specific to

the DG output, or whether neurons in other hippocampal regions can perform temporal pattern

separation. To answer this question, we adapted our pattern separation assay to CA3 pyramidal

cells (PCs) (**Methods – Pattern separation experiments**) and recorded their output spiketrains

in response to direct stimulation of the GCL, i.e. in response to its input from the DG (**Figure**

4a). Due to strong feedforward inhibition, CA3 PCs do not generally spike in response to

external stimulation of afferents in slices, unless inhibition is totally blocked[31-33]. By using 30 Hz

Poisson input sets (**Figure 4b**) and adding 100 nM of gabazine to the bath, which only slightly

decreases the amplitude of GABA-A-mediated IPSCs (see **Methods**), we managed to record for

the first time the spiking output of CA3 PCs in response to complex input spiketrains while

preserving some inhibition in the network (**Figure 4c**). PCs fired during periods of high input

frequency, probably due to depression of local inhibitory transmission[31,34] combined with

facilitation at the mossy fiber-CA3 PCs synapses[35], and as a result action potentials often

appeared more clustered than in control GC recordings (**Figure 4c**). Despite this spiking pattern,

we found that, at the 10 ms timescale, CA3 PCs average (R_{input}, R_{output}) distribution was slightly

but significantly lower than for GCs recorded in similar pharmacological conditions (**Figure 4d**).

This indicates that CA3 PCs are different from GCs in respect to temporal pattern separation, but

that, surprisingly, their output spiketrains are even more decorrelated. Temporal pattern

separation is thus not specific to the DG.

Temporal pattern separation at different timescales across celltypes

Because the timescales meaningful for the brain remain uncertain, it is important to assess the

separation of spiketrains at different timescales. We therefore binned spiketrains using a range of

τ_w from 5 ms to 250 ms and performed a finer grained analysis using pairwise R_{input} and

associated pairwise R_{output} instead of the average across the ten pairs of input trains (**Figure 5a**).

We discovered that pattern separation levels can dramatically change as a function of τ_w. In GCs,

the larger the timescale the less they exhibit decorrelation of their input spiketrains, especially at

high R_{input} (**Figure 5b**). Nonetheless, GCs still exhibit relatively high levels of pattern separation

of highly similar input spiketrains, even at long timescales (**Figure 5b**).

This analysis confirmed that, at short timescales, FSs exhibit less pattern separation than

GCs (**Figure 5c**) and revealed significant differences between HMCs and GCs, especially for

pairs of input spiketrains with low R_{input} (**Figure 5d**), which were not as obvious in our previous

coarse analysis (**Figure 3b3**). At longer timescales, the variability across neurons has a tendency

to increase for all celltypes, but the average levels of decorrelation of both FSs and HMCs stayed

highly significantly different from those of GCs (**Figure 5c-d**). Interestingly, the difference

between FSs and GCs was larger at short timescales, whereas the difference between HMCs and

GCs increased with larger timescales. Indeed, at τ_w above 100 ms, HMCs can often exhibit

pattern convergence instead of pattern separation, especially for pairs of already dissimilar input

spiketrains (low or negative pairwise R_{input}), whereas FSs just show weak or no pattern

separation.

Concerning CA3 PCs, they still exhibit high levels of temporal pattern separation at long

timescales and the difference with GCs increases (**Figure 5e**). Interestingly, in contrast to all

other tested celltypes and conditions (**Figure 5b-d**), PCs even show a dramatic increase of their

average levels of decorrelation at the 250 ms timescale (**Figure 5e**).

Overall, these results show that among the tested DG celltypes, GCs exhibit the highest

levels of temporal pattern separation of cortical spiketrains across all timescales. Our findings

also suggest that the high level of separation by the DG is amplified in CA3.

Mechanism of temporal pattern separation: neural noise

To determine what mechanisms might support temporal pattern separation in GCs, it is necessary

to understand its dynamics first. Limiting our analysis to the first presentation of an input set

revealed that outputs were already significantly decorrelated (**Figure 6a-b**). This shows that the

separation mechanism is fast, consistent with the fact that the brain generally does not have the

opportunity to average repeated signals and that separation must happen immediately during

encoding. In addition, analysis of the last presentation revealed only modestly more separation

than for the first one, and only for high input correlations (**Figure 6c**), suggesting that learning to

recognize the input pattern is not critical.

Because the mechanism for temporal pattern separation is fast and does not require

learning, we asked first whether intrinsic properties of GCs could play a role. Linear regression

analysis revealed that the membrane capacitance, resistance, time constant as well as the resting

membrane potential are not predictors of decorrelation in GCs (see low R^2 in **Table S1**). Another

hypothesis is that randomness in neuronal responses drives the decorrelation. Indeed, when the

same input spiketrain is repeated (e.g. $R_{input} = 1$) the output spiketrains are not well correlated

(as shown by the mean spiketrain reliability R_w) (**Figure 7a**), consistent with well-known trial-

to-trial variability in single neuron responses[36-38]. Theoretical investigation of pattern separation

often relies on some sort of random process such as probabilistic neuronal activation[5] or

stochastic firing[39], which suggests that "neural noise" is a likely contributor to any form of

pattern separation. However, because "neural noise" can cover multiple different definitions and

phenomena[36], determining its role in a complex computation is not trivial.

The noisiness in neural communication is often understood as the unreliability of spiking

after a single input spike, and the jitter of the delay between an input spike and an output

spike[40,41]. We thus assessed such spike-wise noise directly from recording sets from our pattern

separation experiments (**Figure 7b**, **S2**, and **Materials and methods – Analysis of output**

spiketrains). Although the spike probability (SP), delay and jitter of GCs were slightly higher in

our experiments than in a previous report (but recording and analysis methods were different),

the variability between GCs was consistent[41]. We then asked whether these spike-wise noise

parameters could predict the degree of decorrelation by GCs. First, linear regression analysis

shows no good relationship with any parameter, the SP being an average predictor at best

(**Figure S3a** and **Table S2**). Moreover, the average firing rate of a GC output set (a measure

dependent on SP) is not well correlated with the degree of decorrelation either (**Table S3** and

Figure S8a). Temporal pattern separation in GCs seems to not be achieved merely because their

output spiketrains are a sparser and jittered version of their inputs.

To more carefully test the hypothesis that random spiking failures and random delays

support fast temporal pattern separation, we simulated recording sets based on computational

models only governed by spike-wise noise statistics comparable to the original data (**Figure S2,**

S3b, Table S2 and M**ethods – Computational models**). The distribution of (R_{input}, R_{output}) was

significantly higher in the original data (**Figure 7** and **S5**), showing that simple random

processes yield greater levels of separation than real GCs, especially for highly similar inputs

(**Figure 7e** and **S5a**).

In addition to the spike-wise noise, we considered neural noise at the level of spiketrains

by computing the average correlation coefficient R_w between "children" output spiketrains from

the same "parent" input train (**Figure 7a**). R_w characterizes the more complex notion of

spiketrain reliability, that is the ability of a neuron to reproduce the same output spiketrain in

response to repetitions of the same input spiketrain. R_w is not dependent on intrinsic cellular

properties (**Table S1**) and only moderately determined by spike-wise noise parameters (**Table**

S2), suggesting that the rather low R_w of GCs is the expression of more complex noisy

biophysical processes. Consistently, R_w was significantly lower for shuffled and simulated data

than in real GCs (**Figure 7d-e** and **S5b**). This indicates that the output spiketrains of GCs are

more reliable than if their output was entirely determined by simple random processes. Overall,

GCs have both a higher R_{output} and R_w distributions than datasets based on random spiking

(**Figure 7**), which clearly shows that simple noise cannot fully underlie the operations performed

by GCs on input spiketrains.

The fact that random spiking yields better temporal pattern separation but less reliable

information transmission than GCs also suggests that there might be an unavoidable trade-off

between achieving pattern separation and reliable information transmission about input

spiketrains. To further investigate this, we looked at the relationship between the spiketrain

reliability R_w and the decorrelation levels in GC recordings and found a strong anticorrelation

(**Figure 8a** and **Table S3**). This linear relationship is clear evidence that biological processes

leading to sweep-to-sweep variability is a powerful mechanism for temporal pattern separation in

DG.

However, spiketrain unreliability is not the only reason GCs output spiketrains are

decorrelated. Indeed, three lines of evidence support the idea that even if GCs were perfectly

reliable (i.e. a given input train always leads to the same output train), the set of output

spiketrains would still be less similar than the set of inputs. First, the linear model describing the

relationship between Rw and decorrelation levels suggest that even at Rw = 1, ~10% of

decorrelation would still be achieved (**Figure 8a**). Moreover, high levels of pattern separation

are performed upon the first presentation of an input set (**Figure 6**): this means that even if the

full output set was composed of exact repetitions of the set of output trains recorded during the

first sweep (e.g. Input trains 1 and 2 always lead to the same Output trains 1 and 2, respectively)

pattern separation would still occur, at the levels measured for the first sweep (difference

between Outputs 1 and 2 > difference between Inputs 1 and 2). In other words, perfect reliability

does not equate the absence of pattern separation. Finally, when averaging out the variability

between spiketrains associated to the same input (i.e. by computing the PSTH), a significant

level of decorrelation is still detected (**Figure 8b-d**). We can thus conclude that temporal pattern

separation is supported by some mechanisms in addition to those that produce spiketrain

unreliability.

Taken together, these results suggest that complex but noisy biophysical mechanisms

allow GCs to balance temporal pattern separation and reliable signaling about their inputs.

Mechanism of temporal pattern separation: short-term synaptic dynamics

The fact that intrinsic properties of GCs do not predict their decorrelation levels (**Table S1**)

suggests that temporal pattern separation comes from the DG network in which each recorded

GC is embedded. We thus hypothesize that the specific short-term dynamics of synapses in the

DG network implement temporal pattern separation in GCs. This is a likely mechanism because:

1) Synaptic transmission is a stochastic process and generally considered the main source of

noisiness in the spiking output of neurons[36,42], which would fit our conclusion that temporal

pattern separation is dependent on neural noise (see paragraph above), 2) Short-term plasticity

makes the probability of synaptic release dependent on the timing of the preceding input spikes,

which could explain why temporal pattern separation and spiketrain reliability are not purely

governed by noise. To test the idea that short-term synaptic dynamics is a potential mechanism

controlling the pattern separation/spiketrain reliability ratio in GCs, we developed a

computational model of a spiking GC with dynamic and probabilistic PP-GC synapses (see

Methods – Computational models). The parameters of the model controlling the facilitation,

depression and variability of presynaptic release, and thus the amplitude of EPSCs , were

constrained to match the whole-cell voltage-clamp recording of a GC in response to a pattern

separation protocol (10 Hz input trains, R_{input} = 0.76) (**Figure 9a-c** and see **Methods –**

Computational models). Simulations of pattern separation experiments with this model resulted

in high levels of output spiketrains decorrelation, akin to real GCs, but also in relatively high

levels of spiketrain reliability (**Figure 9d-e**). This is a proof of principle that the probabilistic and

dynamic nature of presynaptic release at the PP-GC synapse is a potential mechanism balancing

temporal pattern separation and information transmission in the DG.

Sources of pattern separation differences between celltypes

To further understand the mechanisms behind temporal pattern separation in single hippocampal

cells, we investigated the sources of the difference in decorrelation levels between the tested

celltypes. We considered differences in intrinsic cellular properties (**Table S1**), firing rate and

probability of bursting (**Figure S7-8**), spike-wise noise (**Figure S3-4**), spiketrain reliability

(**Figure 8a** and **S6**) and synaptic transmission dynamics (**Figure S9**).

Both FS and HMC recordings displayed bursts of spikes (defined as more than one output

spike between two input spikes), which was very rarely seen in our GC and CA3 recordings

(**Figure 3-4, S7c-e,** and **S8c, e**). As a result, both FSs and HMCs had significantly higher firing

rates than GCs (**Figure S7a-b**) (although the effect size was smaller for HMCs, because they

burst less often than FSs and they have generally less spikes per bursts. **Figure S7d-e**). Then, are

the firing rate or the probability of bursting predictive of differences in pattern separation? The

relationships are unclear, but it seems that both could partially explain the lower decorrelations

observed in some HMCs and FSs (**Figure S8a-b**). To more directly test the role of bursting, we

processed all FS and HMC recordings by removing all but the first spike in each detected burst

(**Figure S8c, e** and see **Methods – Analysis of the output spiketrains**). These resulting "non-

bursty" datasets (nbFS and nbHMC) still exhibited significantly less pattern separation than GCs

(and even pattern convergence of dissimilar input trains, in the case of HMCs) (**Figure S8d, f**).

Therefore, bursting and high firing rates, although a source of differences in temporal pattern

separation, are not a sufficient explanation.

Although FSs show less pattern separation than GCs, it is interesting that they do exhibit

some amount of separation, as opposed to pattern convergence[6] which one could have expected

from their reputation of having a much more reliable and precise spiking behavior than principal

neurons[40,43]. The high fidelity in relaying input spike[40] might still explain the difference in

pattern separation ability between FSs and GCs, although, to our knowledge, they have never

been formally compared. We thus first confirmed the idea that FS show much less spike-wise

noise than GCs (**Figure S4**). Linear regressions then revealed that spiking probability (SP) is a

good predictor of both spiketrain unreliability $(1 - R_w)$ and decorrelation performance in FSs

(**Figure S3b** and **Table S2**). Surprisingly, the membrane resistance of FSs was also a good

predictor (**Table S1**). Thus, contrarily to GCs, FS pattern separation behavior is strongly and

linearly determined by some intrinsic and spike-wise properties, even though it is in principle

hazardous to anticipate complex neuronal operations from such low-level characteristics, as our

previous analysis on GCs illustrated. Indeed, spike-wise noise parameters of HMCs were very

close to those of GCs (**Figure S4**) and they nonetheless showed striking pattern separation

differences.

Across all celltypes, it is clear that higher SP correlates with lower decorrelation levels

(**Figure S3d**). This suggests that sparseness can be a mechanism partially supporting temporal

pattern separation. Only partially, because: 1) the correlation is not perfect ($R^2 = 57\%$), 2)

although the correlation is better in FSs, FSs decorrelation levels are consistently lower than

expected by the linear model fitting all celltypes (**Figure S3d**), 3) There is no difference in terms

of sparseness between CA3 PCs and their GC controls (**Figure S7**) even though there are clear

differences in terms of pattern separation (**Figure 4-5**).

In the end, the best predictor of decorrelation levels for all celltypes and conditions is R_w,

the spiketrain reliability (**Figure 8a** and **S6**). This result emphasizes the unexpected idea that the

biophysics of neurotransmission in hippocampal networks imposes a trade-off between temporal

pattern separation and reliable information transmission. Figure 9 suggests that the balance

between separation and reliability can depend on synaptic dynamics. Because different celltypes

have synapses with unique properties, we hypothesize that those differences produce varying

degrees of pattern separation behaviors not explained by spiking sparseness (SP, FR, pBurst).

For example, in contrast to PP-GCs synapses that mostly exhibit depression, the monosynaptic

connection from GCs to CA3 PCs is made through giant mossy fiber buttons with low initial

probability of release and short-term facilitation under high input frequency[31,44] (**Figure S9a**).

By modelling a spiking neuron with synapses inspired by GC-CA3 connections and comparing it

to the model from Figure 9, under conditions yielding similar FR, we confirm that differences in

short-term synaptic dynamics alone can lead to obvious pattern separation differences (**Figure**

S9b-c).

Overall, these results show that the differences in temporal pattern separation between

different hippocampal celltypes result from a combination of various sources, each celltype with

a unique combination.

Discussion

We report that similar cortical input spiketrains are transformed in the DG network, leading to

less similar output spiketrains in GCs. Our findings provide the first experimental demonstration

that a form of pattern separation is performed within the DG itself and exhibited at the level of

single neurons at different timescales. This computation arises from noisy but specific

biophysical processes (e.g. synaptic dynamics) in the DG network, where interneurons do not

exhibit as much temporal pattern separation as the final DG output. In turn, the CA3 network

seems to amplify this separation even more at the level of single PCs, suggesting that, at least in

the hippocampus, it is not a computation specific to the DG.

A novel way to test pattern separation

In contrast to *in vivo* experiments that have difficulty identifying the cell-type of recorded units with certainty[26,45-47] and simultaneously recording the direct inputs of these units[13-17], *in vitro* brain slices that preserve the lamellar connections of the hippocampus offer a more accessible platform. For example, a similar experimental setup to ours was used to show that spatially segregated axonal inputs are represented by distinct spatiotemporal patterns in populations of DG neurons[48,49]. However, our study is the first to perform an experimental analysis of pattern separation within DG by directly manipulating the similarity of the inputs and comparing it to the similarity of simultaneously recorded outputs. Such a systematic approach had so far only been done in computational studies[50]. Although a rigorous comparison is impossible because the activity patterns considered were defined differently, the general pattern separation behavior of those models is confirmed by our experimental results: the DG itself performs a form of pattern separation, especially for input patterns that are highly similar (**Figure 2c-d**).

Pattern separation in the time dimension

Until now, most studies of pattern separation in the DG assumed that neural activity patterns were ensembles of ON/OFF neurons[4,5,15,16,20], sometimes considering a rate code averaged over minutes in addition to this population code[13,14,23,39]. Because neurons carry information at timescales shorter than minutes[27-29,38,51] and because the sparse firing of active GCs during a brief event[17,52,53] precludes an efficient rate code[38], we studied pattern separation at sub-second timescales.

Relevant scales are given by the time constant over which neurons can integrate synaptic

inputs[28]: 10-50 ms for GCs and ~100 ms for the "reader" CA3 pyramidal cells. Windows of ~10

ms and ~100 ms, corresponding respectively to gamma and theta rhythms, have been shown to

organize CA neuronal assemblies[27,28,54,55]. Due to specific network properties allowing persistent

activity, the DG might also integrate information over several hundreds of milliseconds[48,49,56].

The point is that multiple timescales can be relevant simultaneously, and because it is still

uncertain which ones are the most important to episodic memory and hippocampal coding, we

investigated a range from 5 ms to 250 ms (**Figure 5**).

Most of our results are reported at a 10 ms resolution, which corresponds approximately

to the spike jitter in GCs (**Figure S2**) as well as their membrane time constant and the gamma

rhythm. This choice of temporal resolution is similar to a recent computational study of pattern

separation within a DG model, which used a 20 ms resolution on short spiketrains (30 ms inputs,

200 ms outputs)[21]. Yet, our study was the first to investigate pattern separation at a range of

timescales. We found that temporal pattern separation in the DG output was best at short

timescales. This relationship was generally conserved across celltypes, with HMCs even

achieving the opposite of pattern separation, pattern convergence, at timescales above 100 ms.

Note, however, that temporal pattern separation is not necessarily a monotonically decreasing

function of the time resolution, as CA3 PCs exhibited a surprising sharp increase of temporal

pattern separation for low input similarity at 250 ms (**Figure 5**).

Our study of pattern separation is the first to focus on temporal patterns, as opposed to

spatial ones; but neural activity patterns are spatiotemporal. More work is clearly needed to test

whether the DG is a pattern separator at the spatiotemporal, population level, but the discovery

of temporal pattern separation in single hippocampal neurons has some implications for

population dynamics. The decorrelation of spiketrains at small timescales, in addition to the fact

that different GCs respond differently to the same inputs (**Figure S1**), suggests that spikes are

constantly rearranged in different time windows, thus enforcing very small neuronal

assemblies[28]. In other words, it ensures that a minimal number of output neurons are active at the

same time, and such sparsity in active neuronal population is known from computational studies

to be critical for efficient population pattern separation[5,22,57].

Mechanisms of temporal pattern separation

The mechanisms supporting pattern separation within DG had so far never been experimentally

investigated. The decorrelation of sequentially presented input patterns can in theory be

explained by: 1) adaptive mechanisms, involving learning and recognition of input patterns,

comparison with previously stored ones and the pruning out of common features, 2) non-

adaptive (intrinsic) mechanisms, 3) or both [58]. First, concerning adaptive mechanisms, it has

been suggested that Hebbian learning could enhance population pattern separation in the DG[59],

but computational models testing different forms of long-term synaptic learning found that it

would actually impair this type of pattern separation[5,23]. As for temporal pattern separation, our

data show that it hardly benefits from the repetition of input patterns (**Figure 6**). We also offer

indirect evidence that non-adaptive decorrelation processes support temporal pattern separation

because output patterns are always decorrelated to the same proportion (**Figure 2e**), a feat that a

simple random process can achieve (**Figure S5c**), suggesting that input patterns do not need to

be recognized. Third, adaptive and non-adaptive mechanisms are not mutually exclusive:

previous learning over days, during the neuron maturation process, could tune single GCs only to

specific input patterns, allowing rapid pattern separation[60]. Indeed, a computational study

suggested that adaptive networks can mature to perform a fast, non-adaptive orthogonalization of

the population activity by the decorrelation between individual information channels[61].

Adaptive or not, what is the biological source of the temporal decorrelation we observed?

We first determined that intrinsic membrane properties do not predict decorrelation levels (**Table**

S1), and that celltypes ability to fire bursts only moderately affects pattern separation (**Figure**

S8). Simple randomness was not sufficient to reproduce our results, even though the spiking

probability, a form of spike-wise neural noise, plays a partial role (**Figure 8** and **S3**). In the end,

temporal pattern separation seems most likely supported by specific short-term synaptic

dynamics, thanks to the synergy between the probabilistic nature of neurotransmission and its

dependency on the spike-timing history imposed by depression or facilitation (**Figure 9, S9**).

Various levels of pattern separation/convergence might thus be achieved in different celltypes

due to a unique combination of synaptic properties.

Indeed, GCs are embedded in a network of synapses, notably receiving feedforward

excitation from the PP, feedforward and feedback inhibition from FSs[43,62] and both excitation

and disynaptic inhibition from HMCs[63]. Thus, although both FSs and HMCs showed some

temporal pattern separation (mostly at short timescales), our results suggest that the final DG

output is inherited from synaptic interactions between all DG celltypes, resulting in maximal

separation at the level of GCs.

More work is needed to clarify the exact role of FSs and HMCs in DG computations.

Computational and experimental studies have suggested that HMCs are involved in some forms

of pattern separation[17, 20, 47]. Our finding of pattern convergence in HMCs for dissimilar inputs

may appear in contradiction with the recent reports that HMCs spatial representations remap in

dissimilar environments[17,26,46,47], but 1) activity patterns were defined differently and averaged

over longer timescales (minutes) and 2) as argued above, remapping is not a direct measure of

pattern separation without knowledge of the input patterns. Overall, we show that HMCs can

exhibit both separation or convergence, depending on the time resolution and the amount of

similarity between input patterns (**Figure 5**). The impact of such a behavior on pattern separation

at the level of GCs remains to be studied. Concerning FSs, they exhibit a poor ability to separate

spiketrains (**Figure 3, 5**), but the somatic inhibition they provide could be the source of low

spiking probability in GCs and thus improve pattern separation by making GCs responses

sparse[5,64]. On the other hand, their ability to relay information reliably[43] (**Figure S4, S6**) and to

precisely control spike timing in target neurons[43] might actually provide a mechanism that

counteracts noisiness in GCs, helping them balance effective separation with fidelity of

information transmission to CA3.

The role of sweep-to-sweep variability

Because the brain needs to be able to recognize when situations are exactly the same, our finding

that pattern separation occurs even when the same input pattern is repeated (**Figure 7a**) might

seem counter-intuitive at first. However, in theory, the separation and the recognition functions

do not have to be supported by the same network. The Hebb-Marr framework actually

hypothesizes that the CA3 recurrent, auto-associative network is able to recall the original

pattern from a noisy input from DG. Even though most computational models that tested the

effect of repetition were consistent with the intuitive view[5,21], this was likely because they used

deterministic neurons. A model considering variability across GCs and a probabilistic spiking

behavior resulted, as in our experiments, in separation of repetitions of the same pattern[39].

In the cortex, the well-known variability of single neuron activity between trials is often

supposed to be "averaged out" at the population level so that the output of the population is

reliable[36]. It is thus conceivable that considering an ensemble of GCs would increase the signal-

to-noise ratio. In fact, when we average out the sweep-to-sweep variability, GCs exhibit pattern

separation for highly similar patterns but almost no separation for identical ones (**Figure 8d**).

However, this variability, or "noise", is not necessarily meaningless[36]. Our results

suggest it might be a mechanism amplifying pattern separation (**Figure 8**). The variability might

even be just apparent, if we consider that when the same input is repeated it is at different points

in time: each repetition could be considered as a different event that needs to be encoded slightly

differently. The DG would thus meaningfully add some noise to transform input spiketrains so

that cortical information about an event is stored in the hippocampus with a unique random time-

stamp, consistent with the index theory of episodic memory[65].

Pattern separation and pattern completion in CA3

In the Hebb-Marr framework, CA3 is thought to be a recurrent, autoassociative network that can

perform pattern completion, and thus can support the recall of a full memory from a partial cue

of the original event[4]. Although CA3 recurrent connections are sparser than previously thought[66],

computational models generally confirmed that CA3 could perform pattern completion of

population patterns[4,23,66]. Direct experimental evidence are scarce and unclear, both in vitro[67],

and in vivo[32,68,69]. In addition, the fact that neuronal representations of similar environments are

more correlated in CA3 than in the DG was thought to constitute indirect evidence[13,14], but recent

reports suggesting that the DG representations previously measured were not coming solely from

DG output cells have clouded these initial conclusions[26,46].

Assuming that pattern completion is a process realized by CA3, this implies that, when

presented with different but similar partial cues of the same initial memory, the final output of

CA3 should converge towards the same representation. Our finding that CA3 PCs exhibit high

levels of temporal pattern separation might then come as a surprise (**Figure 4-5**). Several lines of

reasoning could explain this result. First, we focused on temporal patterns in single cells, and it is

possible that a network can perform population pattern completion in addition to temporal

pattern separation. Actually, different environments can be represented in CA3 by different

populations of PCs, or by the same PCs with remapped firing rates[26,46], which has led some to

conclude that CA3 could perform pattern separation[13]. A recent report even showed that single

GCs remap much less than CA3 PCs[26]: assuming GCs and PCs receive inputs with similar

overlap, this is consistent with our finding that PCs are better than GCs at temporal pattern

separation. Second, we tested CA3 under partial block of inhibition in order to allow PCs to fire.

Given that the number of active CA3 PCs is generally higher than DG GCs in vivo[5,24,26,46], it

suggests that our experiment may not have modelled physiological conditions well. Third, CA3

is known to be physiologically and functionally heterogeneous along its proximodistal axis[32],

with studies suggesting that the PCs closest to DG perform population pattern separation[68,70,71].

We recorded from PCs at the CA3b/c border, and it is possible that more distal PCs would

exhibit less temporal pattern separation. Last but not least, it is important to note that *pattern*

completion and *pattern separation* are not opposite, mutually exclusive computations. Pattern

separation and its actual opposite, pattern convergence, describe the similarity of multiple

patterns from an input network (e.g. EC) compared to the similarity of the corresponding patterns

of another network (e.g. DG)[6]. On the other hand, pattern completion is the process, happening

over time, of retrieval of a previously learned full pattern in a network (e.g. CA3) from a partial

seed pattern in the same network[5]. Our experiments did not test for pattern completion, as the

different trains of an input set were not degraded versions of a previously learned pattern.

Moreover, pattern separation and completion are complementary in the sense that pattern

completion would benefit from initial input patterns being as separated as possible[4,5]. It thus

makes sense that CA3 would start by amplifying the separation of input patterns, as is the case in

our data, either before encoding or before proceeding to completion of the seed pattern. In fact,

our results suggest that CA3 might complement the separation effectuated in the DG, as CA3 is

able to perform high decorrelation levels at long timescales (> 100 ms) when the DG doesn't

(**Figure 5**). CA3 could thus ensure that seed input patterns are well separated at all timescales.

Materials and methods

Animals and dissection

All experiments were performed in accordance with the National Institute of Health guidelines

outlined in the National Research Council Guide for the Care and Use of Laboratory Animal

(2011) and regularly monitored and approved by the University of Wisconsin Institutional

Animal Care and Use Committee.

Horizontal slices of the ventral and intermediate hippocampus (400 μm) were prepared

from the brains of young (p15-25) or adult (p121 ± 15 days) C57BL/6 male mice

(Harlan/Envigo). Adult animals were only used for **Figure 9**. Mice were anesthetized with

isoflurane, decapitated, and the brain was removed quickly and placed in ice-cold cutting

sucrose-based solution (two different compositions were used. For HMC, CA3 pyramidal cells,

their GC controls and GCs from adult mice, we used version #2 [72]). The cutting solutions

contained (version #1 / version #2, in mM): 83 / 80 NaCl, 26 / 24 NaHCO$_3$, 2.5 / 2.5 KCl, 1 /

1.25 NaH2PO$_4$, 0.5 / 0.5 CaCl$_2$, 3.3 / 4 MgCl2, 22 / 25 D-Glucose, 72 / 75 Sucrose, 0 / 1 Na-L-

Ascorbate, 0 / 3 Na-Pyruvate, bubbled with 95% O$_2$ and 5% CO$_2$. During the dissection, brain

hemispheres were prepared following the "magic cut" procedure [73] with α-angle around 10 to 15°

and β-angle around 5 to 10° for GC and FS recordings, and with α close to 0° and β between 0

and -5° for HMC and CA3 pyramidal cells recordings [66]. Slices were cut using a vibratome

(Leica VT1000S) then placed in an incubation chamber in standard artificial cerebrospinal fluid

(aCSF) containing (in mM) 125 NaCl, 25 NaHCO3, 2.5 KCl, 1.25 NaH2PO4, 2 CaCl2, 1

MgCl2, and 25 D-Glucose (or in a 50/50 mix of standard aCSF and cutting solution. 100%

cutting solution for HMC, CA3 recordings and their GC controls, and GCs from adult mice) at

35°C, for 15-30 minutes after dissection. Slices were stored in the incubation chamber at room

temperature for at least 30 minutes before being used for recordings.

Electrophysiology

All recordings were done in standard aCSF adjusted to 325 mOsm, at physiological temperature

(33-35 °C). Whole cell patch-clamp recordings were made using an upright microscope

(Axioskop FS2, Zeiss, Oberkochen, Germany) with infra-red differential interference contrast

optics. Patch pipettes pulled from thin-walled borosilicate glass (World Precision Instruments,

Sarasota, FL) had a resistance of 2-5 MΩ when filled with intracellular solution containing (in

mM) 140 K-gluconate, 10 EGTA, 10 HEPES, 20 phosphocreatine, 2 Mg$_2$ATP, 0.3 NaGTP. For

the dataset from adult mice, we used a slightly different recipe: 135 K-gluconate, 5 KCl, 0.1

EGTA, 10 HEPES, 20 Na-Phosphocreatine, 2 Mg$_2$-ATP, 0.3 Na-GTP, 0.25 CaCl$_2$. Intracellular

solutions were adjusted to pH 7.3 and 310 mOsm with KOH and H_2O. Both recipes yielded

similar electrophysiological behaviors.

Recordings were done using one or two Axopatch 200B amplifiers (Axon Instruments,

Foster City, CA), filtered at 5 kHz using a 4-pole Bessel filter and digitized at 10 kHz using a

Digidata 1320A analog-digital interface (Axon Instruments). Data were acquired to a Macintosh

G4 (Apple Computer, Cupertino, CA) using Axograph X v1.0.7 (AxographX.com). Stimulation

pipettes were pulled from double barrel borosillicate theta-glass (~10 μm tip diameter, Harvard

Apparatus, Edenbridge, U.K.) and filled with ACSF or a 1M NaCl solution and connected to a

constant current stimulus isolator used to generate 0.1-10 mA pulses, 100 microseconds in

duration.

Series resistance and cellular intrinsic properties were assessed online in Axograph from

the fit of the electrical responses to repetitions of 5-10 mV, 25 ms steps, holding the potential at -

65 mV. Neurons used for analysis were stable across a whole recording session as judged by

monitoring of series resistance and resting potential.

Dentate granule cells (GC) were visually identified as small cells in the granule cell layer

(GCL). GCs from young mice had the following intrinsic properties (mean ± sem) : resting

potential (V_{rest}) -69.3 ± 1.3 mV; input resistance (R_i) 171 ± 16MΩ and capacitance (C_m) 23 ± 2

pF. GCs from adult mice had the following intrinsic properties: V_{rest} -78.8 ± 1.8 mV; R_i = 137 ±

14 MΩ and C_m = 18 ± 1 pF.

Fast-spiking interneurons (FS) were identified as neurons with large somata at the hilus-

GCL border and a high firing rate response during large depolarizing current steps, and a large

after-hyperpolarization (AHP) [62,74] (**Figure 3**). They had the following intrinsic properties:

V_{rest}= -66.7 ± 3.5 mV; R_i = 59 ± 10 MΩ and C_m = 19 ± 3 pF.

Hilar Mossy Cells (HMC) were identified as large neurons in the deep hilus (> 60 μm

away from the GCL) with a regular firing response to current steps and a small AHP, as well as

with a high frequency of spontaneous EPSPs, all characteristics that allow to distinguish them

from other neurons in the hilus [63,75] (**Figure 3**). Their intrinsic properties were: V_{rest} = -69.7 ± 2.3

mV; R_i = 198 ± 12 MΩ and C_m = 33 ± 3 pF.

For CA3 pyramidal cells, because recent studies suggest a proximodistal gradient of

physiological and computational properties in the CA3 network [32,68,70,71], we avoided the

extremity of the CA3c region and targeted large neurons around the CA3b/c border in the

pyramidal layer. Their intrinsic properties under partial inhibitory block (aCSF + 100 nM

gabazine) were: V_{rest} = -72.7 ± 2.2 mV; R_i = 186 ± 12 MΩ and C_m = 36 ± 2 pF. The intrinsic

properties of GCs recorded under the same pharmacological conditions were: V_{rest} = -77.3 ± 1.9

mV; R_i = 244 ± 13 MΩ and C_m = 25 ± 2 pF.

Pattern separation experiments

We designed multiple sets of input patterns, each with a prespecified average Pearson's

correlation coefficient (R_{input}) computed with a binning window (τ_w) of 10 ms, using two

different algorithms (one developed in-house based on iterative modifications, and a more

efficient one from [76] based on mathematical principles and a preset covariance matrix). An input

set consisted of five different input spiketrains, each 2 s trains of impulses simulating cortical

spiketrains, with interspike intervals following a Poisson distribution. Each pair of input trains

had a correlation coefficient close to the R_{input} of its set (at τ_w = 10 ms, the average relative

standard error is 4% across all input sets). For all experiments except in **Figure 4** (CA3 and GC

control), the mean frequency of input trains was set close to 10 Hz (11.9 ± 0.7 Hz). This input

firing rate was chosen to be consistent with the frequency of EPSCs recorded in GCs of behaving

mice [52], and is known to promote a high probability of spiking in GCs in slices [62,77].

The responses of one or two DG neurons were recorded in whole-cell mode while stimuli

were delivered to the outer molecular layer (OML). Stimulus current intensity and location were

set so that the recorded neuron spiked occasionally in response to electrical impulses (see range

of spike probability in **Figure S2-4**) and the stimulation electrode was at least 100 μm away

from the expected location of the dendrites of the recorded neuron. Once stimulation parameters

were set, a pattern separation protocol was run: the five trains of a given input set were delivered

one after the other, separated by 5 s of relaxation, and this was repeated ten times. The ten

repetitions of the sequence of five patterns were implemented to take into account any potential

variability in the output, and the non-random sequential scheme was used to avoid repeating the

same input spiketrain close in time. Each protocol yielded a recording set of fifty output

spiketrains, each associated with one of the five input trains of an input set (**Figure 1c**). A given

cell was recorded in response to up to five input sets with different R_{input} (i.e. a recorded cell

produced between one and five data points on **Figure 2c**).

The membrane potential baseline was maintained around -70mV during both current-

clamp and voltage-clamp recordings, consistent with the V_{rest} of mature GCs recorded in

behaving mice [52]. For comparison, FS and HMC current-clamp recordings were also held at -70

mV. The output spiking frequency of GCs was variable (6.3 ± 0.3Hz, see **Figure S7-8**) but

consistent with sparse activity generally observed in GCs *in vivo* during behavior [26,52,53,78] and in

slices under conditions of drive comparable to ours [41,79,80].The output firing rates of FS and HMC

were higher than their GC controls (**Figure S7-8**), as expected from recent research [26,43,47].

To perform pattern separation experiments in CA3 pyramidal cells (PCs), we had to

change several parameters in order to make CA3 PCs spike. Indeed, PCs firing is controlled by

strong feedforward inhibition [31,32], and all tested stimulation sites led to net IPSPs or, rarely, to

weak EPSPs. To make PCs fire in response to external electrical stimulations: 1) the stimulating

electrode was placed in the inferior blade of the GCL to make the DG output cells fire, 2) we

targeted CA3b PCs, which have the highest E/I ratio across CA3 in slices and still receive strong

connections from mossy fibers [32], 3) we used input sets with 30 Hz input trains to promote the

depression of inhibitory transmission [31], 4) we maintained the membrane potential baseline

between -60 and -70 mV, and 5) we had to add 100 nM gabazine to the bath of standard aCSF in

order to slightly decrease IPSCs amplitude (recordings started at least 15 min after the slice was

placed in the bath, to insure equilibrium was reached). 100 nM of gabazine (SR-95531, Sigma-

Aldrich) has been shown to correspond to a 70% availability of GABA-A receptors [81] and, in our

conditions, consistently decreased spontaneous IPSCs mean amplitude by 30% in voltage-clamp

recordings of GCs and CA3 PCs held at -40 mV. These conditions allowed us to record for the

first time the spiking output of CA3 PCs in response to complex input spiketrains while

preserving some inhibition in the network [31]. PCs output firing rate were on average below 10Hz

(**Figure S7-8**), but close to mean rates observed in vivo during behavior [13,14,26].

Analysis of the output spiketrains

For each recording set, the similarity between pairs of spiketrains was computed as the Pearson's

correlation coefficient between the spiketrains rasters binned at a τ_w timescale. Sweeps without

spikes were excluded from further analysis.

We did not use separate protocols to assess the *firing rates*, *probability of bursting* and

spike-wise noise parameters (spike probability, delay, jitter), but computed them directly for each

recording set of spiketrains from a pattern separation experiment. The mean firing rate was

computed as the average firing rate across all fifty output spiketrains. A burst was defined as the

occurrence of more than one output spike in the interval of two input spikes (see **Figure S8**).

We define the spike-wise neural noise as the *probability of spiking* at least once after an

input spike, the *delay* of an output spike after an input spike and its average *jitter*. To assess these

parameters, we computed the cross-occurrence between input spikes and output spikes in a [-15

ms, 50ms] interval with 1 ms bins. The resulting histogram of counts of output spikes occurring

in the vicinity of an input spike was fitted with a Gaussian distribution $N(\mu,\sigma, \text{baseline})$, where μ

is the mean delay of an output spike and σ is the jitter of this delay. The baseline corresponds to

the background firing, occurring by chance or caused by neighboring inputs. After subtracting

the baseline and extracting the probability of spiking by dividing the counts of output spikes by

the total number of input spikes, we defined the *spike probability* (SP) as the sum of probabilities

of an output spike in the predefined time interval around an input spike (**Figure 7b** and **Figure

S2a**). For FS and HMC, spike-wise noise parameters were computed on a non-bursty dataset

(nbFS and nbHMC) where all but the first spike in a burst were excluded. Spike-wise noise

parameters were not computed for CA3 PCs and their GC control because the input frequency

was too high to allow a good fit of the distribution.

Computational models

To assess the role of spike-wise neural noise in pattern separation, we generated two data sets.

First, we simulated output spiketrains in response to our input sets (for each input set, we

simulated ten output sets of fifty synthetic spiketrains). This simulation was entirely based on the

average spike-wise noise parameters computed from the original GC recordings (see above): the

matrix of input spike times was replicated ten times, and for each of the fifty resulting sweeps,

spikes were deleted randomly following a binomial distribution B(Nspk, F), where Nspk is the

number of input spikes in a sweep, and F the probability of not spiking (F = 1-mean SP = 1-

0.42). A random delay, sampled from a Gaussian distribution $N(\mu,\sigma)$, was added to each

resulting spike times, with μ and σ being respectively the mean delay and mean jitter in the

original recordings. The noise statistics of the resulting simulated data set is shown in **Figure**

S2c. Second, we created a surrogate data set by randomly shuffling the output spikes of the

original GC recordings: the delay of each spike was conserved but it was relocated to follow a

randomly selected input spike in the same input train (from a uniform distribution). This

manipulation yielded a dataset with noise statistics very close to the original data (**Figure 2d**).

Using this strategy, we performed spike shuffling in each GC recording set a hundred times,

yielding a dataset of 10,200 simulated recording sets, or, in other words, 100 datasets of 102

simulated recording sets directly paired to the original 102 GC recording sets.

To test whether probabilistic synaptic dynamics is a potential mechanism of temporal

pattern separation, we used an Izhikevich model of a regular spiking neuron [82] with an adapted

Tsodyks-Markram (TM) synapse model [83] designed to capture short-term plasticity of stochastic

neurotransmission at the LPP-GC synapse. The TM synapse model consists of a system of two

ordinary differential equations describing the dynamics of X and U. X is the probability of a

presynaptic vesicle being available for release, the decrease of which leads to synaptic

depression, and U is the probability of release for an available vesicule, the increase of which

models facilitation:

$$\frac{dX}{dt} = U(t).X(t).\delta(t - t_{ap}) + \frac{1 - X(t)}{\tau_{rec}} \quad (1)$$

$$\frac{dU(t)}{dt} = U_{SE}.(1 - U(t)).\delta(t - t_{AP}) - \frac{U(t)}{\tau_{facil}} \quad (2)$$

With $X(0) = 1$ and $U(0) = 0$, where τ_{rec} is the time constant of vesicule recovery after release,

τ_{facil} is the decay of U controlling synaptic facilitation, U_{SE} a factor determining the probability of

release Pr at the time of the first spike, δ the Dirac delta function, and t_{AP} the time of arrival of an

input spike. The system of ODEs was solved using the ode23 MATLAB solver.

Based on the quantal theory of neurotransmission, the model assumes that, at each time

point t, k vesicles are released, drawn from a binomial distribution $B(N, Pr)$, where N is the

number of release sites and Pr is the probability of release defined as the product of X and U.

The current I is then the product of k with the quantal size q. Based on previous data from our

lab, we considered $q = 20$ pA [62]. The version of the TM model we implemented only focuses on

the dynamics of the presynaptic vesicles, ignoring postsynaptic dynamics. To match the decay of

real EPSCs recorded at the soma, I was convolved with a template EPSC shape:

$$-A.\left(e^{-T/\tau_{rise}} - e^{-T/\tau_{decay}}\right) \quad (3)$$

Where $T = 50$ ms, $\tau_{rise} = 0.5$ ms, $\tau_{decay} = 5$ ms and A is a factor normalizing the area of the

resulting shape to 10. In addition, a synaptic delay of 4 ms was introduced. The resulting current

(**Figure 9b-c**) is then passed to a non-dimensional regular spiking Izhikevich neuron, an

acceptable model for GCs [84]. We used the following parameters: a = 0.02, b = 0.2, c = -65 mV, d

= 6, time resolution = 0.1 ms, max voltage = 30 mV, initial values $V(0)$ = -70 mV and $x(0)$ =

b.$V(0)$.

To optimize the non-fixed parameters of the TM-Izhikevich model, we simulated output

spiketrains in response to ten repetitions of a single input set (10 Hz, R_{input} = 0.76) and compared

to the current dynamics and spiking behavior of a representative GC that was recorded both in

voltage-clamp and current-clamp in response to the same input set. For the TM model, the

following parameters were adjusted to match the dynamics and the peak of the current averaged

over ten repetitions, as well as the variability in the peak amplitude of the first EPSC: τ_{rec} = 500

ms, τ_{facil} = 9 ms, U_{SE} = 13, N = 40 release sites. Finally, because the Izhikevich model integrates

non-dimensional currents, the input current had to be scaled using the dividing constant K = 62,

such that the average firing rate of the simulation matched the average firing rate of the

representative GC to the same input set (3.5 Hz). K can be considered as a constant modelling a

tonus of inhibition: the higher K is, the more difficult it is for an EPSC to make the Izhikevich

neuron fire. In the end, the standard deviation of the firing rate was ~0.6 Hz for simulations,

compared to 0.8 Hz for the original recording. The only source of variability was from

presynaptic dynamics of vesicle release modelled as a binomial process.

To determine the influence of synaptic transmission parameters on pattern separation, we

compared two TM-Izhikevich models with different parameters. Model 1 has exactly the same

parameters as described above except that, in order to model the presence of gabazine, the

inhibition constant K was set at 40: with this value, Model 1 responses to 30 Hz input trains have

a FR ~ 7 Hz, corresponding to the mean FR of real GCs recorded under partial inhibitory block

(**Figure S7**). For model 2, q = 29 pA, τ_{rec} = 100 ms, τ_{facil} = 500 ms, U_{SE} = 0.01, K = 57, and all

other parameters were the same as for model 1. U_{SE}, q and N (kept at 40) were chosen based on

estimates from the literature on mossy fiber buttons, the giant GC-CA3 PCs synapses[31,44]. τ_{rec},

τ_{facil} and K were adjusted to model short-term synaptic facilitation observed at mossy fiber

buttons and to match the average firing rate that we observed in our CA3 current-clamp

recordings made under gabazine (~7Hz, **Figure S7**). In order to specifically study the impact of

pre-synaptic dynamics, post-synaptic (i.e. EPSC shape) and Izhikevich parameters were kept the

same for all models. Note that none of our models are intended to closely match spiking

behaviors observed in real GCs or CA3 PCs.

Software and statistics

Data analysis was performed using custom-written routines in MATLAB (2017a), including

functions from toolboxes cited above. Sample sizes were chosen based on the literature and

estimations of the variance and effect size from preliminary data. All values are reported as mean

± S.E.M. unless otherwise noted. The one-sample Kolmogorov-Smirnov test was used to verify

the normality of data distributions. Parametric or non-parametric statistical tests were

appropriately used to assess significance (p-value < 0.05). Assumptions on equal variances

between groups were avoided when necessary. All T and U tests were two-tailed. To determine

whether two distributions of data points are significantly different (e.g. R_{output} as a function of

R_{input}, for GC compared to FS, see **Figure 3-6 and Figure S5 and S8**), we performed an analysis

of the covariance (ANCOVA) using linear regressions on the two data sets as well as on the

combined data set, and assessed significance via an F-test comparing the goodness of fits[85].

Because R_{input} can also be considered as a categorical variable, we performed a two-way

ANOVA before using post-hoc tests correcting for multiple comparisons in order to determine at

which R_{input} groups two conditions were significantly different. When comparing FSs, HMCs or

CA3 with GCs, different GCs were used as control and were recorded under different protocols,

which is why we did not need to control for multiple comparisons across celltypes. In order to

determine whether distributions were significantly different in the case of our spike shuffling

analysis (**Figure 7**), we designed Monte-Carlo exact hypothesis tests[86]. **Table S4** provides details

on all statistical tests conducted in this study.

Data availability

788   Data used in this manuscript are freely available at: https://www.ebi.ac.uk/biostudies/studies/S-

789   BSST219

790   References

1   Treves, A., Tashiro, A., Witter, M. P. & Moser, E. I. What is the mammalian dentate gyrus good for? *Neuroscience* **154**, 1155-1172, doi:10.1016/j.neuroscience.2008.04.073 (2008).

2   McNaughton, B. L. & Nadel, L. Hebb-Marr networks and the neurobiological representation of action in space. *Neuroscience and connectionist theory*, 1-63 (1990).

3   McNaughton, B. L. & Morris, R. G. M. Hippocampal synaptic enhancement and information storage within a distributed memory system. *Trends in Neuroscience* **10**, 408-415 (1987).

4   Rolls, E. T. A computational theory of episodic memory formation in the hippocampus. *Behavioural brain research* **215**, 180-196, doi:10.1016/j.bbr.2010.03.027 (2010).

5   O'Reilly, R. C. & McClelland, J. L. Hippocampal conjunctive encoding, storage, and recall: avoiding a trade-off. *Hippocampus* **4**, 661-682, doi:10.1002/hipo.450040605 (1994).

6   Santoro, A. Reassessing pattern separation in the dentate gyrus. *Frontiers in behavioral neuroscience* **7**, 96, doi:10.3389/fnbeh.2013.00096 (2013).

7   Amaral, D. G., Scharfman, H. E. & Lavenex, P. The dentate gyrus: fundamental neuroanatomical organization (dentate gyrus for dummies). **163**, 3-790, doi:10.1016/s0079-6123(07)63001-5 (2007).

8   McHugh, T. J. *et al.* Dentate gyrus NMDA receptors mediate rapid pattern separation in the hippocampal network. *Science* **317**, 94-99, doi:10.1126/science.1140263 (2007).

9   Kesner, R. P. & Rolls, E. T. A computational theory of hippocampal function, and tests of the theory: new developments. *Neuroscience and biobehavioral reviews* **48**, 92-147, doi:10.1016/j.neubiorev.2014.11.009 (2015).

10   Kesner, R. P., Kirk, R. A., Yu, Z., Polansky, C. & Musso, N. D. Dentate gyrus supports slope recognition memory, shades of grey-context pattern separation and recognition

memory, and CA3 supports pattern completion for object memory. *Neurobiology of*
learning and memory **129**, 29-37, doi:10.1016/j.nlm.2015.08.010 (2016).
11   Baker, S. *et al.* The Human Dentate Gyrus Plays a Necessary Role in Discriminating
New Memories. *Current biology : CB*, doi:10.1016/j.cub.2016.07.081 (2016).
12   Dillon, S. E. *et al.* The impact of ageing reveals distinct roles for human dentate gyrus
and CA3 in pattern separation and object recognition memory. *Scientific reports* **7**,
14069, doi:10.1038/s41598-017-13853-8 (2017).
13   Leutgeb, J. K., Leutgeb, S., Moser, M. B. & Moser, E. I. Pattern separation in the dentate
gyrus and CA3 of the hippocampus. *Science* **315**, 961-966, doi:10.1126/science.1135801
(2007).
14   Neunuebel, J. P. & Knierim, J. J. CA3 retrieves coherent representations from degraded
input: direct evidence for CA3 pattern completion and dentate gyrus pattern separation.
Neuron **81**, 416-427, doi:10.1016/j.neuron.2013.11.017 (2014).
15   Deng, W., Mayford, M. & Gage, F. H. Selection of distinct populations of dentate
granule cells in response to inputs as a mechanism for pattern separation in mice. *eLife* **2**,
e00312, doi:10.7554/eLife.00312 (2013).
16   Marrone, D. F., Adams, A. A. & Satvat, E. Increased pattern separation in the aged fascia
dentata. *Neurobiology of aging* **32**, 2317 e2323-2332,
doi:10.1016/j.neurobiolaging.2010.03.021 (2011).
17   Nakazawa, K. Dentate Mossy Cell and Pattern Separation. *Neuron* **93**, 465-467 (2017).
18   Neunuebel, J. P., Yoganarasimha, D., Rao, G. & Knierim, J. J. Conflicts between local
and global spatial frameworks dissociate neural representations of the lateral and medial
entorhinal cortex. *The Journal of neuroscience : the official journal of the Society for*
Neuroscience **33**, 9246-9258, doi:10.1523/JNEUROSCI.0946-13.2013 (2013).
19   Knierim, J. J. & Neunuebel, J. P. Tracking the flow of hippocampal computation: Pattern
separation, pattern completion, and attractor dynamics. *Neurobiology of learning and*
memory **129**, 38-49, doi:10.1016/j.nlm.2015.10.008 (2016).
20   Myers, C. E. & Scharfman, H. E. A role for hilar cells in pattern separation in the dentate
gyrus: a computational approach. *Hippocampus* **19**, 321-337, doi:10.1002/hipo.20516
(2009).
21   Yim, M. Y., Hanuschkin, A. & Wolfart, J. Intrinsic rescaling of granule cells restores
pattern separation ability of a dentate gyrus network model during epileptic
hyperexcitability. *Hippocampus*, doi:10.1002/hipo.22373 (2014).
22   Chavlis, S., Petrantonakis, P. C. & Poirazi, P. Dendrites of dentate gyrus granule cells
contribute to pattern separation by controlling sparsity. *Hippocampus* (2017).
23   Neher, T., Cheng, S. & Wiskott, L. Memory storage fidelity in the hippocampal circuit:
the role of subregions and input statistics. *PLoS computational biology* **11**, e1004250,
doi:10.1371/journal.pcbi.1004250 (2015).
24   Schmidt, B., Marrone, D. F. & Markus, E. J. Disambiguating the similar: the dentate
gyrus and pattern separation. *Behavioural brain research* **226**, 56-65,
doi:10.1016/j.bbr.2011.08.039 (2012).
25   Ramirez, S. *et al.* Creating a false memory in the hippocampus. *Science* **341**, 387-391,
doi:10.1126/science.1239073 (2013).
26   Senzai, Y. & Buzsáki, G. Physiological properties and behavioral correlates of
hippocampal granule cells and mossy cells. *Neuron* **93**, 691-704. e695 (2017).

27 Kelemen, E. & Fenton, A. A. Coordinating different representations in the hippocampus.
Neurobiology of learning and memory **129**, 50-59, doi:10.1016/j.nlm.2015.12.011
(2016).
28 Buzsaki, G. Neural syntax: cell assemblies, synapsembles, and readers. *Neuron* **68**, 362-
385, doi:10.1016/j.neuron.2010.09.023 (2010).
29 Eichenbaum, H. Time cells in the hippocampus: a new dimension for mapping memories.
Nature reviews. Neuroscience **15**, 732-744, doi:10.1038/nrn3827 (2014).
30 van Dijk, M. T. & Fenton, A. A. On How the Dentate Gyrus Contributes to Memory
Discrimination. *Neuron* **98**, 832-845 e835, doi:10.1016/j.neuron.2018.04.018 (2018).
31 Rebola, N., Carta, M. & Mulle, C. Operation and plasticity of hippocampal CA3 circuits:
implications for memory encoding. *Nature reviews. Neuroscience* **18**, 208-220,
doi:10.1038/nrn.2017.10 (2017).
32 Sun, Q. *et al.* Proximodistal Heterogeneity of Hippocampal CA3 Pyramidal Neuron
Intrinsic Properties, Connectivity, and Reactivation during Memory Recall. *Neuron* **95**,
656-672 e653, doi:10.1016/j.neuron.2017.07.012 (2017).
33 Sachidhanandam, S., Blanchet, C., Jeantet, Y., Cho, Y. H. & Mulle, C. Kainate receptors
act as conditional amplifiers of spike transmission at hippocampal mossy fiber synapses.
The Journal of neuroscience : the official journal of the Society for Neuroscience **29**,
5000-5008, doi:10.1523/jneurosci.5807-08.2009 (2009).
34 Mori, M., Abegg, M. H., Gahwiler, B. H. & Gerber, U. A frequency-dependent switch
from inhibition to excitation in a hippocampal unitary circuit. *Nature* **431**, 453-456,
doi:10.1038/nature02854 (2004).
35 Vyleta, N. P., Borges-Merjane, C. & Jonas, P. Plasticity-dependent, full detonation at
hippocampal mossy fiber–CA3 pyramidal neuron synapses. *eLife* **5** (2016).
36 Faisal, A. A., Selen, L. P. & Wolpert, D. M. Noise in the nervous system. *Nature reviews.*
Neuroscience **9**, 292-303, doi:10.1038/nrn2258 (2008).
37 Dobrunz, L. E. & Stevens, C. F. Response of hippocampal synapses to natural
stimulation patterns. *Neuron* **22**, 157-166 (1999).
38 Rieke, F., Warland, W., de Ruyter van Steveninck, R. & Bialek, W. *Spikes: Exploring the*
Neural Code. (THE MIT PRESS: A Bradford Book, 1999).
39 Aimone, J. B., Wiles, J. & Gage, F. H. Computational influence of adult neurogenesis on
memory encoding. *Neuron* **61**, 187-202, doi:10.1016/j.neuron.2008.11.026 (2009).
40 Savanthrapadian, S. *et al.* Synaptic Properties of SOM- and CCK-Expressing Cells in
Dentate Gyrus Interneuron Networks. *The Journal of neuroscience : the official journal*
of the Society for Neuroscience **34**, 8197-8209, doi:10.1523/JNEUROSCI.5433-13.2014
(2014).
41 Mongiat, L. A., Esposito, M. S., Lombardi, G. & Schinder, A. F. Reliable activation of
immature neurons in the adult hippocampus. *PloS one* **4**, e5320,
doi:10.1371/journal.pone.0005320 (2009).
42 Mainen, Z. F. & Sejnowski, T. J. Reliability of spike timing in neocortical neurons.
Science **268**, 1503-1506 (1995).
43 Hu, H., Gan, J. & Jonas, P. Interneurons. Fast-spiking, parvalbumin(+) GABAergic
interneurons: from cellular design to microcircuit function. *Science* **345**, 1255263,
doi:10.1126/science.1255263 (2014).

44 Evstratova, A. & Toth, K. Information processing and synaptic plasticity at hippocampal
mossy fiber terminals. *Frontiers in cellular neuroscience* **8**, 28,
doi:10.3389/fncel.2014.00028 (2014).
45 Neunuebel, J. P. & Knierim, J. J. Spatial firing correlates of physiologically distinct cell
types of the rat dentate gyrus. *The Journal of neuroscience : the official journal of the*
Society for Neuroscience **32**, 3848-3858, doi:10.1523/JNEUROSCI.6038-11.2012
(2012).
46 GoodSmith, D. *et al.* Spatial representations of granule cells and mossy cells of the
dentate gyrus. *Neuron* **93**, 677-690. e675 (2017).
47 Danielson, N. B. *et al.* In Vivo Imaging of Dentate Gyrus Mossy Cells in Behaving Mice.
Neuron **93**, 552-559. e554 (2017).
48 Hyde, R. A. & Strowbridge, B. W. Mnemonic representations of transient stimuli and
temporal sequences in the rodent hippocampus in vitro. *Nature neuroscience* **15**, 1430-
1438, doi:10.1038/nn.3208 (2012).
49 Zylberberg, J., Hyde, R. A. & Strowbridge, B. W. Dynamics of robust pattern
separability in the hippocampal dentate gyrus. *Hippocampus* **26**, 623-632,
doi:10.1002/hipo.22546 (2016).
50 Chavlis, S. & Poirazi, P. Pattern separation in the hippocampus through the eyes of
computational modeling. *Synapse (New York, N.Y.)*, doi:10.1002/syn.21972 (2017).
51 VanRullen, R., Guyonneau, R. & Thorpe, S. J. Spike times make sense. *Trends in*
neurosciences **28**, 1-4, doi:10.1016/j.tins.2004.10.010 (2005).
52 Pernía-Andrade, Alejandro J. & Jonas, P. Theta-Gamma-Modulated Synaptic Currents in
Hippocampal Granule Cells In Vivo Define a Mechanism for Network Oscillations.
Neuron **81**, 140-152, doi:10.1016/j.neuron.2013.09.046 (2014).
53 Diamantaki, M., Frey, M., Berens, P., Preston-Ferrer, P. & Burgalossi, A. Sparse activity
of identified dentate granule cells during spatial exploration. *eLife* **5**,
doi:10.7554/eLife.20252 (2016).
54 Lisman, J. E. & Jensen, O. The theta-gamma neural code. *Neuron* **77**, 1002-1016,
doi:10.1016/j.neuron.2013.03.007 (2013).
55 Wikenheiser, A. M. & Redish, A. D. Decoding the cognitive map: ensemble hippocampal
sequences and decision making. *Current opinion in neurobiology* **32**, 8-15 (2015).
56 Larimer, P. & Strowbridge, B. W. Representing information in cell assemblies: persistent
activity mediated by semilunar granule cells. *Nature neuroscience* **13**, 213-222,
doi:10.1038/nn.2458 (2010).
57 Severa, W., Parekh, O., James, C. D. & Aimone, J. B. A Combinatorial Model for
Dentate Gyrus Sparse Coding. *Neural computation*, 1-24, doi:10.1162/NECO_a_00905
(2016).
58 Friedrich, R. W. & Wiechert, M. T. Neuronal circuits and computations: pattern
decorrelation in the olfactory bulb. *FEBS letters* **588**, 2504-2513,
doi:10.1016/j.febslet.2014.05.055 (2014).
59 Fontanari, J., Rolls, E. & Costa, D. F. A model of the operation of the hippocampus and
entorhinal cortex in memory. *International Journal of Neural Systems* **6**, 51-70 (1995).
60 Aimone, J. B., Deng, W. & Gage, F. H. Resolving new memories: a critical look at the
dentate gyrus, adult neurogenesis, and pattern separation. *Neuron* **70**, 589-596,
doi:10.1016/j.neuron.2011.05.010 (2011).

61 Wick, S. D., Wiechert, M. T., Friedrich, R. W. & Riecke, H. Pattern orthogonalization
via channel decorrelation by adaptive networks. *Journal of computational neuroscience*
28, 29-45, doi:10.1007/s10827-009-0183-1 (2010).
62 Ewell, L. A. & Jones, M. V. Frequency-tuned distribution of inhibition in the dentate
gyrus. *The Journal of neuroscience : the official journal of the Society for Neuroscience*
30, 12597-12607, doi:10.1523/jneurosci.1854-10.2010 (2010).
63 Scharfman, H. E. The enigmatic mossy cell of the dentate gyrus. *Nature reviews.*
Neuroscience **17**, 562-575, doi:10.1038/nrn.2016.87 (2016).
64 Temprana, S. G. *et al.* Delayed coupling to feedback inhibition during a critical period for
the integration of adult-born granule cells. *Neuron* **85**, 116-130,
doi:10.1016/j.neuron.2014.11.023 (2015).
65 Teyler, T. J. & Rudy, J. W. The hippocampal indexing theory and episodic memory:
updating the index. *Hippocampus* **17**, 1158-1169, doi:10.1002/hipo.20350 (2007).
66 Guzman, S. J., Schlögl, A., Frotscher, M. & Jonas, P. Synaptic mechanisms of pattern
completion in the hippocampal CA3 network. *Science* **353**, 1117-1123 (2016).
67 Jackson, M. B. Recall of spatial patterns stored in a hippocampal slice by long-term
potentiation. *Journal of neurophysiology* **110**, 2511-2519, doi:10.1152/jn.00533.2013
(2013).
68 Marrone, D. F., Satvat, E., Odintsova, I. V. & Gheidi, A. Dissociation of spatial
representations within hippocampal region CA3. *Hippocampus*, n/a-n/a,
doi:10.1002/hipo.22367 (2014).
69 Denny, C. A. *et al.* Hippocampal memory traces are differentially modulated by
experience, time, and adult neurogenesis. *Neuron* **83**, 189-201,
doi:10.1016/j.neuron.2014.05.018 (2014).
70 Lee, H., Wang, C., Deshmukh, S. S. & Knierim, J. J. Neural population evidence of
functional heterogeneity along the CA3 transverse axis: pattern completion versus pattern
separation. *Neuron* **87**, 1093-1105 (2015).
71 Lu, L., Igarashi, K. M., Witter, M. P., Moser, E. I. & Moser, M.-B. Topography of place
maps along the CA3-to-CA2 axis of the hippocampus. *Neuron* **87**, 1078-1092 (2015).
72 Yi, F. *et al.* Hippocampal "cholinergic interneurons" visualized with the choline
acetyltransferase promoter: anatomical distribution, intrinsic membrane properties,
neurochemical characteristics, and capacity for cholinergic modulation. *Frontiers in*
synaptic neuroscience **7**, 4, doi:10.3389/fnsyn.2015.00004 (2015).
73 Bischofberger, J., Engel, D., Li, L., Geiger, J. R. & Jonas, P. Patch-clamp recording from
mossy fiber terminals in hippocampal slices. *Nature protocols* **1**, 2075-2081,
doi:10.1038/nprot.2006.312 (2006).
74 Harney, S. C. & Jones, M. V. Pre- and postsynaptic properties of somatic and dendritic
inhibition in dentate gyrus. *Neuropharmacology* **43**, 584-594 (2002).
75 Larimer, P. & Strowbridge, B. W. Nonrandom local circuits in the dentate gyrus. *The*
Journal of neuroscience : the official journal of the Society for Neuroscience **28**, 12212-
12223, doi:10.1523/jneurosci.3612-08.2008 (2008).
76 Macke, J. H., Berens, P., Ecker, A. S., Tolias, A. S. & Bethge, M. Generating spike trains
with specified correlation coefficients. *Neural computation* **21**, 397-423,
doi:10.1162/neco.2008.02-08-713 (2009).

77 Andersen, P., Holmqvist, B. & Voorhoeve, P. E. Excitatory synapses on hippocampal
apical dendrites activated by entorhinal stimulation. *Acta physiologica Scandinavica* **66**,
461-472, doi:10.1111/j.1748-1716.1966.tb03224.x (1966).
78 Danielson, N. B. *et al.* Distinct contribution of adult-born hippocampal granule cells to
context encoding. *Neuron* **90**, 101-112 (2016).
79 Dieni, C. V. *et al.* Low excitatory innervation balances high intrinsic excitability of
immature dentate neurons. *Nature communications* **7**, 11313, doi:10.1038/ncomms11313
(2016).
80 Pardi, M. B., Ogando, M. B., Schinder, A. F. & Marin-Burgin, A. Differential inhibition
onto developing and mature granule cells generates high-frequency filters with variable
gain. *eLife* **4**, e08764, doi:10.7554/eLife.08764 (2015).
81 Jones, M. V., Jonas, P., Sahara, Y. & Westbrook, G. L. Microscopic kinetics and
energetics distinguish GABA A receptor agonists from antagonists. *Biophysical journal*
81, 2660-2670 (2001).
82 Izhikevich, E. M. Simple model of spiking neurons. *IEEE transactions on neural*
networks **14**, 1569-1572, doi:10.1109/tnn.2003.820440 (2003).
83 Fuhrmann, G., Segev, I., Markram, H. & Tsodyks, M. Coding of temporal information by
activity-dependent synapses. *Journal of neurophysiology* **87**, 140-148 (2002).
84 Hananeia, N. & Benuskova, L. Computational simulation of dentate gyrus granule cell—
The role of metaplasticity. *Neurocomputing* **175**, 300-309 (2016).
85 Motulsky, H. J. & Ransnas, L. A. Fitting curves to data using nonlinear regression: a
practical and nonmathematical review. *FASEB journal : official publication of the*
Federation of American Societies for Experimental Biology **1**, 365-374 (1987).
86 Waller, L. A., Smith, D., Childs, J. E. & Real, L. A. Monte Carlo assessments of
goodness-of-fit for ecological simulation models. *Ecological Modelling* **164**, 49-63
(2003).
1021
1022

1025
Figure 1. Pattern separation assay in acute brain slices at the single cell level.

(a) Examples of input sets. *Top*: each input set comprises five different trains of electrical pulses.

Bottom: correlation coefficient matrix for each input set, each square representing the correlation

coefficient between two input trains measured with a binning window (τ_w) of 10 ms. R_{input} is the

average of coefficients, diagonal excluded.

(b) Histology of the DG in a horizontal slice (Cresyl violet/Nissl staining; scale bar: 250 µm),

overlaid with a schematic of the experimental setup: a theta pipette in the ML (input) is used to

focally stimulate the PP while a responding GC (output) is recorded via whole-cell patch-clamp.

(GCL: granule cell layer, H: hilus, ML: molecular layer, FS: fast-spiking interneuron. Solid lines

represent dendrites and dashed lines axons)

(c) Current-clamp recordings of the membrane potential of two different GCs in response to

different input sets (*Top*: $R_{input} = 1$; *Bottom*: $R_{input} = 0.76$). Each input set (five input traces) is

repeated ten times (only three repetitions are shown, with spikes truncated at 0 mV). In the

bottom graph, input trains and their respective children output spiketrains have matching colors.

1040

1041
Figure 2. Input spiketrains are decorrelated at the level of individual granule cells.

(a) Example of a recording set (input set + output set): the raster plot shows one set of input

spiketrains and the children output spiketrains recorded from one GC, reordered to display output

subsets (i.e., the ten children coming from one parent input spiketrain) together and with the

same color.

(b) Corresponding 55x55 correlation coefficient matrix using a binning window (τ_w) of 10 ms.

Each small square represents the correlation coefficient between two spiketrains. R_{output} is

defined as the mean of correlations between individual output spiketrains driven by different

input spiketrains, as outlined by the bold blue border, which excludes comparisons between

outputs generated from the same parent input.

(c) Data points, corresponding to 102 recording sets (28 GCs), are all below the identity line

(dashed line). This means that R_{output} was lower than R_{input} for all recordings, thus demonstrating

pattern separation.

(d) *Left*: Effective decorrelation averaged over all recording sets as a function of R_{input}. Although

there is a significant decorrelation for all tested input sets (one-sample T-tests: the blue shade

indicates the 95% confidence interval that average decorrelation is significantly above 0), they

are effectively decorrelated to different magnitudes (one-way ANOVA, p < 0.0001). *Right:*

Matrix of p-values from post-hoc Tukey-Kramer tests comparing effective decorrelation levels

across all pairs of parent input sets. The asterisk and star correspond to the comparisons

displayed in the left panel. This analysis shows that the decorrelation is significantly different

(higher) for highly similar input spiketrains than for already dissimilar inputs.

(e) When the effective decorrelation is normalized to the correlation of the input set, there is no

significant difference between input sets (ANOVA, p = 0.18).

In all graphs, τ_w = 10 ms. Means and SEM in black.

Figure 3. Different levels of temporal pattern separation in different DG cell types

(a1) Picture of a recorded FS filled with biocytin (black). In the case of simultaneous recordings, the recorded GCs were close to the FS, as depicted by the schematic in green. In different experiments, we recorded from putative hilar mossy cells (HMC, orange). Full lines represent dendrites, dashed lines axons.

(a2-3) Example of a simultaneous whole-cell recording of a GC and a neighboring FS.

(a2) Simultaneous membrane potential recordings (baseline at -60mV) of a FS and a GC to the same set of current steps (-25, 100, 500 and 1000 pA).

(a3) Simultaneous current-clamp recordings of the same FS and GC as in a2 in response to the five input traces of an input set with $R_{input} = 0.65$. Simultaneous input and output trains have the same color.

(a4) R_{output} versus R_{input} for FSs and GCs. Data points correspond to recording sets: 20 for FS (red, 4 cells, 4 per input set), and 61 for GC (green, with a darker shade open circle when simultaneously recorded with a FS, 13 cells, 11-13 per input set). All GC recordings done at the same input correlations as FS recordings were used for an unpaired comparison showing that FS exhibit less spiketrain decorrelation than GCs: ANCOVA: $p < 0.0001$, 95% confidence bounds around the linear fits shown as shaded areas; two-way ANOVA: input sets: $p = 0.0016$, cell-types: $p < 0.0001$, interaction: $p = 0.72$. Post-hoc T-tests with Bonferroni corrections for 5 comparison groups: all $p < 0.05$ (for decreasing R_{input}, $p = 0.0307, 0.0181, 0.0007, 0.0001, 0.0122$)

(a5) Effective decorrelation ($R_{input} - R_{output}$) for FS and GC. Shaded areas represent the 95% confidence interval of a bootstrap test comparing the mean decorrelation of both celltypes: GCs exhibit significantly more pattern separation than FS for all input sets.

(**a4-5**) Note that when comparing only the simultaneous GC and FS recordings, we found a similarly significant difference between celltypes.

(**b1**) Membrane potential of a hilar neuron in response to current steps (-100, 0, 100, 400 pA; baseline at -70mV), showing a spontaneous barrage of EPSPs, regular spiking, and a lack of large after-hyperpolarization, all typical features of HMCs (Larimer & Strowbridge, 2008).

(**b2**) Current-clamp recordings of same HMC in response to a set of five input trains ($R_{input} = 0.49$, $\tau_w = 10$ms). HMCs fire occasional bursts of spikes (marked by asterisks) in response to a single input, which was not seen in GCs.

(**b3**) R_{output} versus R_{input} for HMCs and GCs. Data points correspond to recording sets: 18 for

HMC (orange, 11 cells, 5-7 per input set), and 22 for GC (green, 11 cells, 4-10 per input set). An

unpaired comparison suggests that HMCs and GCs show only slight differences in pattern

separation measured at $\tau_w = 10$ ms: ANCOVA: $p = 0.15$, 95% confidence bounds around the

linear fits shown as shaded areas; two-way ANOVA: input sets: $p = 0.0004$, cell-types: $p =$

0.074, interaction: $p = 0.57$. Post-hoc T-tests with Bonferroni corrections for three comparison

groups (for decreasing R_{input}): $p = 1, 0.05, 0.21$.

1106

1107
Figure 4. CA3 pyramidal cells exhibit more temporal pattern separation than GCs

(a) Schematic of the experimental setup: a theta pipette in the granule cell layer (GCL) is used to

stimulate GCs and their mossy fibers (dashed line) to evoke action potentials in CA3 pyramidal

cells recorded via whole-cell patch-clamp. To limit feedforward inhibition and allow pyramidal

cells to spike, experiments were performed under partial block of inhibition (100 nM of

gabazine). Control experiments were performed in GCs under the same pharmacological

conditions but with OML stimulations.

(b) Two input sets of 30 Hz Poisson trains were used. *Top*: rasters of the five spiketrains of each

set. *Bottom*: correlation coefficient matrix for each input set ($\tau_w = 10$ ms).

(c) Example of current-clamp recordings from a pyramidal cell and a GC from the same animal.

Left: Membrane potential responses to current steps (-100pA, 100pA, 350pA). Scale: 100 ms

(horizontal) and 50 mV (vertical). *Right:* example of 5 output responses (sweeps 11-15) to an

input set ($R_{input} = 0.76$).

(d) R_{output} versus R_{input} for CA3 pyramidal cells and GCs. Data points correspond to recording

sets: 15 for CA3 (black, 14 cells, 6-9 recordings per input set), and 22 for GC (green, 13 cells, 11

per input set). The CA3 distribution is lower and significantly different from the GC distribution:

ANCOVA: $p < 0.01$, 95% confidence bounds around the linear fits shown as shaded areas; two-

way ANOVA: input sets: $p < 0.0001$, cell-types: $p = 0.0036$, interaction: $p = 0.24$. Post-hoc T-

tests with Bonferroni corrections for two comparison groups: $p = 0.032$ for $R_{input} = 0.76$, $p = 0.1$

for $R_{input} = 0.21$.

1128

Figure 5. Differences in temporal pattern separation between hippocampal celltypes depend on the timescale.

(**a**) Pairwise analysis. Instead of averaging across all five input trains of a set (here $R_{input} = 0.76$ at $\tau_w = 10$ ms, same matrix as in Fig. 2b), we average children output coefficients corresponding to a single pair of input trains (identified by color-coded squares.). This finer analysis is necessary at τ_w higher than 20 ms because the *pairwise input correlation coefficients* are not well constrained around their mean anymore (see right panel). *Middle* and *Right*: mean across cells, following the same color-code as displayed in the matrix (*Left*).

(**b**) Pairwise R_{output} vs pairwise R_{input} for GCs, measured with different binning windows τ_w. Only the means across cells are displayed, but the regression line was fitted on the full distribution of data points. The larger τ_w, the less GCs exhibit pattern separation.

(**c**) *Top:* Pairwise analysis on FS recordings (same as in b). *Bottom*: mean± SEM and regression lines for τ_w = 100 ms: FS and GC distributions are still different, especially at high input similarity (ANCOVA: $p < 0.0001$ for τ_w = 10 up to 250 ms). Lower right inset: effect size as a function of the timescale τ_w (up to 250 ms): Mean ± SEM (red) and maximum (black dots) of the absolute difference between FSs and GCs Routput mean values for all pairwise R_{input}. Note a decreasing effect of larger τ_w on the difference between FSs and GCs in terms of pattern separation.

(**d**) *Top:* Pairwise analysis on HMC recordings (same color code as in b). *Bottom*: At τ_w = 250 ms, HMC and GC distributions are different (ANCOVA: $p < 0.0001$ for τ_w = 10 up to 250 ms), especially at lower input similarity: notice the points above the identity line, showing pattern convergence for HMCs in contrast to GCs. Under this pairwise analysis (in contrast to Fig. 3b), HMCs and GCs are significantly different at τ_w = 10 ms (comparison graph not shown, see inset for effect size). However, the inset (see c for legends) shows an increasing effect of τ_w on the difference between HMCs and GCs in terms of pattern separation.

(**e1**) Pairwise analysis on CA3 pyramidal cells and GCs, both under partial inhibition block and under 30 Hz inputs (same color code as in b).

(**e2**) At τ_w = 100 ms and 250 ms, CA3 and GC distributions are still different (ANCOVA: $p < 0.0001$). The inset (see c for legends) shows an increasing effect of τ_w (plateauing at large timescales) on the difference between CA3 PCs and GCs.

1160

Figure 6. Input spiketrains are efficiently separated upon their first presentations.

(a) Two of five inputs are shown with corresponding output spiketrains. The first output sweep is marked with a pink bar (right) and last sweep is marked with a blue bar.

(b) R_{output}, computed from the first sweep of five output trains only (pink), as a function of R_{input}, fitted with a parabola. All data points are below the identity line indicating that outputs are effectively decorrelated compared to their inputs even when input patterns have only been presented once each. The average decorrelation (R_{input}-R_{output}) is significant for all input sets (one-sample T-tests, $p < 0.01$) except for $R_{input} = 0.11$ ($p = 0.1$).

(c) *Left*: Output correlations (Mean ± SEM) between spiketrains of the first sweep (pink) and the last sweep (blue). There is no significant difference (ANCOVA, $p = 0.33$). *Right:* When taking

1171   into account that the two distributions are paired, we detect that a few output correlations are

1172   significantly lower for the last sweep than for the first one (one-sample T-test on the difference

1173   between R_{output} of the first and last sweep of each recording set, asterisks signify $p < 0.05$). This

1174   is evidence, though weak, that repetition of input spiketrains might improve pattern separation

1175   for highly similar inputs.

1176

Figure 7. Pattern separation in single GCs is not explained by simple neural noise.

(a) The variability of output spiketrains in response to the same input train sets the upper bound for R_{output}. *Left:* Correlations between pairs of output spiketrains associated with different input trains (R_{output}) and pairs of different output spiketrains associated with the same input train (enclosed by green, R_w: spiketrain reliability, the reproducibility of the output given the same input). *Right:* Frequency distribution of R_w for all recordings (green; dark green line is the mean: $<R_w> = 0.3$), overlaid on the distribution of 102 (R_{output}, R_{input}) points and its regression line (black). Note that means $<R_w>$ and $<R_{output}>$ for $R_{input} = 1$ are close because they both assess the reproducibility of the output when the input is the same.

(b) Characterization of neural noise. *Top:* example of input and output spiketrains illustrating variable delay of the response spike after an input spike (d1, d2) or failure to spike after an input spike (red cross). *Bottom*: Example from one GC recording. The spike-wise noise in output spiketrains is characterized by the average spike delay, the standard deviation of this delay (jitter) and the probability of spiking after an input spike (spiking probability, SP).

(c-e) Effect of spike shuffling on R_{output} and R_w. For each output spiketrain, each original spike was reassigned to the time of a randomly chosen input spike. This shuffling was performed 100 times for each of the 102 original GC recording sets, producing 10,200 shuffled recording sets.

(c) The (R_{output}, R_{input}) GC distribution (green) is overlaid with the 90% sample interval of the distribution of the 10,200 shuffled recording sets (grey area: 5 to 95 percentiles). A linear regression was performed for each of the 100 shuffling distributions (102 simulated data points per shuffle): the 90% confidence interval (CI) around the mean regression line is represented by the two dashed lines (short dash) very close to each other, and the GC regression line (green) falls out of this range. As illustrated in the inset, 100% of the shuffling regression lines (grey squares) have a lower slope and higher intercept than the regression line for the original GC dataset (i.e. the green dot is outside of the grey cloud). This Monte-Carlo exact test shows that spiketrains are significantly *less* separated in GCs than would be expected from random spiking.

(d) Frequency distributions of R_w for 102 GC recording sets (green, same as in a, solid line is mean) and 10,200 shuffled recording sets (grey). Dashed lines represent the 90% CI of the shuffling mean (see inset). Inset: distribution of the mean R_w for the 100 shufflings. GCs mean R_w (0.3) is outside of this distribution, showing that GCs output is significantly *more* reliable than expected from random spiking.

(e) Paired statistical tests (based on difference between each GC recording and its 100 associated

shuffled recording sets) show that spike shuffling leads to smaller R_{output} and R_w than original GC

recordings. *Top:* Frequency distribution of the difference of R_w (10,200 data points). Monte-

Carlo exact tests: 99.25% of data points (i.e. $p = 0.0075$) and 100% of the shuffling means are

above 0. *Bottom*: Difference of R_{output} as a function of R_{input}. The grey area represents the sample

interval where 90% of the 10,200 data points fall (5-95 percentiles): for high R_{input} values, all

points are above 0 (see table S4 for details). Solid grey line: means; dashed lines: 90% CI.

Monte-Carlo exact tests based on proportion of shuffling means above 0: asterisks denote

significance (see table S4 for details).

1218

1219

Figure 8. Unreliability in spiketrain transmission is a major but not unique source of

temporal pattern separation.

(a) Spiketrain reliability (R_W) is an excellent predictor of normalized decorrelation (defined in

Figure 2E) for all celltypes and conditions. Notice that, despite the strong anticorrelation, the

intercept of the linear model at $R_w = 1$ predicts that even a perfect reliability could still allow

10% of decorrelation. See Table S3 for linear regressions on single celltypes.

(b) To assess the amount of decorrelation not due to spiketrain unreliability, the ten children

output spiketrains of each of the five trains of an input set can be averaged to give the five output

peristimulus histograms (PSTH). The 10 ms binned PSTHs of the output rasters in Figure 2a are

shown.

(c) Correlation coefficients between all pairs of the five output PSTHs. The mean correlation

(PSTH R_{output}) is the average of coefficients inside the red border, and excludes self-comparisons.

1232 **(d)** *Left:* PSTH R_{output} as a function of R_{input} (102 recording sets, in red), fitted with a parabola

1233 (black). All points are below the identity line indicating decorrelation of outputs compared to

1234 inputs. *Right:* Average effective decorrelation (R_{input} – PSTH R_{output}) as a function of R_{input} (bars

1235 are SEM) reveals a significant decorrelation for all input sets except for the most dissimilar (one-

1236 sample T-tests; shaded area is the 95% confidence interval for significant decorrelation).

1237

1238

Figure 9. Probabilistic presynaptic release is a potential mechanism balancing temporal

pattern separation and spiketrain reliability

(a) Non simultaneous current-clamp ($V_{rest} \approx -70mV$) and voltage-clamp recordings ($V_{hold} = -70mV$) in response to a set of input patterns were successively performed in GCs from adult

mice. The example shows the first sweep of responses from a single cell to the five trains of an

input set ($R_{input} = 0.76$, $\tau_w = 10$ ms).

(b-e) The recording set of the example cell in a was used to fit a computational model of a

spiking neuron with dynamic and probabilistic synapses.

(b) *Top*: four first input spikes of input train #1. *Middle*: Dynamics of the variables of a

probabilistic Tsodyks-Markram model of a synapse in response to the four input spikes. X

represents the probability of vesicle availability at the presynaptic site, U represents the

probability of release of available vesicles, and Pr is the probability of release for all vesicles.

The number of released vesicles was simulated at every time point from a binomial process.

Bottom: Parameters of the model were adjusted so that the resulting current (averaged over the

10 repetitions, in dark orange) would reasonably match the peak of the corresponding average

EPSCs in the original voltage-clamp recordings (baselined and inverted display with partially

blanked stimulation artifact, in black).

(c) Parameters of the model were also adjusted to match the variability in the amplitude of the

first EPSC in the original recording. *Top*: average EPSC. *Middle*: individual EPSCs evoked from

the 10 repetitions of the first input spike in input train #1. The range of peak amplitudes is

similar between model and data. *Bottom*: The coefficient of variation of the current in the model

is close to the data.

(d-e) Temporal pattern separation and spiketrain reliability ($\tau_w = 10$ ms) for GCs from adult mice

and a single model regular spiking Izhikevich neuron with probabilistic synapses as described

above. Data points correspond to recording/simulation sets: 35 for GCs (green circles, 14 cells,

6-16 per input set), and 5 simulations at all input sets (dark orange triangles), with the model

fitted to a single recording set ($R_{input} = 0.76$; R_{output} and R_w shown in dark green circles for all

input sets).

1267

1 **Supplements**
2
3 Tables and supplementary figures for "Pattern separation of spiketrains in hippocampal neurons"
4 Authors: A.D. Madar, L.A. Ewell, M.V. Jones.
5
6

7
Figure S1. Different granule cells process identical inputs differently.
(A) The similarity between pairs of spiketrains coming from two different output sets but associated to the
same input set and with the same sweep number is assessed with the Pearson's correlation coefficient (τ_w
= 10 ms). The fifty resulting coefficients are then averaged to give $R_{cell-to-cell}$, a single number measuring
the overall similarity of all output spiketrains between two output sets. **(B)** Probability distribution of $R_{cell-to-cell}$ (green histogram) across all GC recordings (all combinations of pairs of GC output sets from the
same input set were compared). The distribution of $R_{cell-to-cell}$ (black circles) is not dependent on R_{input}.
14

Figure S2 related to Figure 7. Spike delay, jitter and reliability distributions for real data,
simulations and shuffled data
(**A**) Cross-occurrence method to measure spike delay, jitter and spiking reliability of a neuron during a
given recording session. *Top*: Example histogram of output spikes occurring after input spikes, fitted (red
curve) with a Gaussian distribution N(μ,σ, baseline), where μ is the mean delay and σ is the jitter of this
delay. Lag 0 ms corresponds to the input spike time. In this example, output spikes are generated on
average 16 ms after a stimulation impulse (delay) with a jitter (σ) of 8.7 ms. *Bottom***:** the baseline is
subtracted and the histogram divided by the number of input spikes during the recording session. This
gives the distribution of the probability of spiking after an input spike, the sum of probabilities defining
the spiking probability (SP) of the cell during the recording session. Here the neuron fires 39 % of the
time after an input spike.
(**B-D**) Delay, jitter and spiking probability (SP) distributions as a function of input sets, for (**B**) the
original GC recordings, (**C**) the simulations of binomially random spiking with Gaussian delay (n = 110)
and (**D**) one spike shuffling dataset (n = 102). Dashed horizontal red lines are means.

Figure S3 related to Figure 7. Spike-wise neural noise characteristics are not good predictors of spiketrain decorrelation by single GCs.

Plots of the normalized decorrelation, i.e. $(R_{input} - R_{output}) / R_{input}$, of each recording set ($\tau_w = 10$ ms) : 102 for GC original and shuffled recordings (**A-B**), 20 for FS (**C**), and for GC, FS and HMC pooled together (**D**) as a function of spike-wise noise characteristics (spike delay, jitter and probability). Solid green lines are the best linear fit, with R^2 and p-values noted in each panel. These plots illustrate **Table S2**. Note that decorrelation is poorly explained (low R^2) by either the spike delay or its jitter in all cell-types. In contrast, the spiking probability (SP) is a good predictor of decorrelation in shuffled GC recordings (n = 102 recordings entirely dominated by spike-wise noise. See Figure S2) and even more so in FS recordings (for FS, SP was computed from nbFS data. See Figure S7). This suggests that a low SP can be a potent mechanism for decorrelation, and that FS show different levels of decorrelation than GCs partly because they are more reliable. However, the regression line for FS is lower than for GCs (even FS with low SP show less decorrelation than GCs with similar SP), and SP is only an average predictor of decorrelation for GCs, thus confirming that temporal pattern separation in single GCs cannot be the result of simple neural noise.

Figure S4 related to Figure 7. Spike-wise noise characteristics of granule cells are different than those of DG interneurons.

Mean +/- SEM (bars) and individual recordings (dots). Spike delay, jitter and spiking probability (SP) for FS interneurons compared to GC recordings associated with the same input sets than FS (*top*) (20 FS recordings, 61 GC recordings, see Figure 3A) and HMC interneurons compared to a different set of GC recordings associated with the same input sets than HMC (*bottom*) (18 HMC recordings, 22 GC recordings, see Figure 3B). A U-test was applied to each pair of comparison, showing that FS are much faster and less noisy than GCs, whereas HMC are similar to GCs (slightly larger jitter, but slightly higher SP).

58
Figure S5 related to Figure 7. R_{output} and spiketrain reliability are lower for a random spike
generator than for GCs.
A Binomial/Gaussian random spike generator with parameters based on the mean spike-wise neural noise
measured in GCs leads to more decorrelation but less spiketrain reliability than in GCs.
(A) R_{output} distribution at $\tau_w = 10$ ms, for simulated and GC recordings. (ANCOVA: $p < 0.0001$). Shaded
areas (green and grey) represent the 95% CI of the regressions.
(B) R_w distributions are significantly different (unpaired T-test correcting for unequal variances: $p <$
0.0001, $<Rw>simul = 0.14$).
(C) Like in the original data (**Fig 2E**), the average normalized decorrelation ((R_{input}-R_{output})/R_{input}) seems
invariant. Bars are SEM.
(A-B) Asterisks: $p < 0.05$.
70

71
Figure S6 related to Figure 8A. Spiketrain reliability in different hippocampal celltypes
FS have a higher R_w than GCs (n = 20 vs 61 recording sets, unpaired T-test: p < 0.0001). Note that when
comparing only the simultaneous FS and GC recordings (dark green), we found a similarly significant
difference. HMCs R_w are not significantly higher than in GCs (n = 18 vs 22 recording sets, unpaired T-
test: p = 0.0963). Under partial block of inhibition, CA3 pyramidal cells have a lower R_w than GCs (n =
15 vs 22 recording sets, unpaired T-test: p = 0.0052). Asterisks: p < 0.05
78

79
Figure S7 related to Figure 3. DG interneurons fire bursts and have a higher firing rate than GCs
(A-B) Comparison of the firing rate in all celltypes and conditions. **(A)** For each comparison the GC
dataset is different and matched to the non-GC dataset, as in Figure 3-4. FS and HMC have a higher firing
rate than their GC control (U-test: $p < 0.0001$ and $p = 0.0005$ respectively) whereas CA3 and GC
recordings under gabazine and 30 Hz input trains do not differ (U-test: $p = 0.18$).
(B) Cumulative distributions of the firing rate for all recordings of all celltypes (n = 102 GC, 20 FS, 18
HMC, 15 CA3+gzn, 22 GC+gzn).
(C-D) Same as A-B for the probability of bursting in a recording. Bursting was defined as the occurrence
of at least two action potentials between two input pulses. **(C)** FS and HMC have a higher propensity to
fire bursts than their GC control (U-test: $p < 0.0001$) whereas CA3 pyramidal cells and their GC control
do not differ (U-test: $p = 0.74$).
(E) The distribution of the number of spikes between two input pulses, for all celltypes, show that FS are
the only neurons that consistently fire large bursts. Note that GCs almost never fire more than twice, and
when they do it generally corresponds to two preceding input pulses occurring close to each other (the
first spike being in reality associated to a different input than the next spike).
95

Figure S8 related to Figure 3 and 5. Differences in pattern separation between GC and DG interneurons are not solely due to different firing rates and bursting behaviors.

(A) The relationship between the firing rate and decorrelation levels in all recordings of all celltypes. In contrast to GCs, for FS neurons there is a strong correlation between the firing rate of a recording set and the associated normalized decorrelation. See **Table S3** for more details.

(B) The relationship between the probability of bursting (see Figure S7) and the decorrelation levels in all recordings of all celltypes is less clear, but cells bursting more than 30% of the time have the lowest decorrelation levels.

(C) Example of bursts in a FS recording (*bottom*) in response to input spikes (*top*). To assess the effect of bursting on R_{output}, we truncated each recorded spiketrain from FS neurons to keep only the first output spike between two input spikes, thus removing any burst without altering the SP of the cell. The blue shaded areas highlight the spikes that were removed. The resulting truncated dataset was termed "nbFS" for "non-burst FS".

(D) R_{output} versus R_{input} for nbFS and GC at $\tau_w = 10$ms (to compare to Figure 3A4). Both distributions are still significantly different, suggesting the bursting behavior of FS is not sufficient to explain the difference in pattern separation: ANCOVA: $p < 0.0001$.

(E) Bursts in HMC recordings were truncated to produce an "nbHMC" dataset.

(F) Pairwise analysis on nbHMC and GC recordings at $\tau_w = 250$ ms show that the distributions are still different between the two celltypes (to compare to Figure 5D): ANCOVA: $p < 0.0001$. Under this analysis, nbHMCs and GCs are also significantly different at lower timescales ($\tau_w = 100, 50$ and 10 ms) but with a lower effect size as τ_w decreases (not shown).

119
Figure S9 related to Figure 5 and 9. Short-term synaptic dynamics differences can drive differences
in terms of temporal pattern separation.
(A) Examples of current-clamp recordings in one GC and one CA3 PC, under partial block of inhibition,
in response to a 30 Hz input train (input pulses are noticeable as downward artefacts). EPSPs visually
appear quite different between GCs and CA3 PCs, with clear facilitation in CA3 PCs, which leads them to
spike mostly during periods of high input frequencies.
(B-C) We designed two models of spiking neurons only differing by a few synaptic parameters. Model 1
is the same as the model presented in Figure 9, with depressing EPSC dynamics, except that the inhibition
constant was decreased to match FR observed in real GC+gzn recordings. Model 2 was inspired by GC-
to-CA3 mossy fiber synapses that exhibit low initial probability of release and short-term facilitation, and
the inhibition constant was set so to match the mean FR of real CA3+gzn recordings (~7Hz). Model 1 and
2 were thus designed to have different synaptic dynamics but yield similar FR. Spiking responses to the
pattern separation protocols used on real CA3 PCs and their GC controls (30 Hz input trains: $R_{input} = 0.21$
and 0.76, shown in Figure 4) were then simulated (n = 5 simulated output sets for each model and each
R_{input}).
(B) Example of current and voltage responses of model 2 to a 30 Hz input train. Asterisks correspond to
spike times of a single sweep from a CA3 PC recording (different than in A).
(C) Pairwise correlation analysis, as in Figure 5, shows that the two models yield visually obvious
differences in terms of temporal pattern separation (crosses and bars: mean +/- SEM; solid lines: linear
regression on data points). Although model 1 and 2 do not reproduce well the pattern separation levels
observed in real GCs and CA3 PCs (Figure 5E), they qualitatively go in the same direction: the GC-to-
CA3 inspired model 2 produces visually lower R_{output} than the PP-GC inspired model 1, at short
timescales (10 ms) and large ones (250 ms, for low R_{input} values). Overall, this analysis shows that
differences in terms of synaptic dynamics are sufficient to cause differences in terms of temporal pattern
separation.

Table S1-3. **Linear regressions goodness-of-fit, p-value and slope.** The predictor variables (x-axis)
correspond to columns, and the variables to be explained (y-axis) correspond to rows. Red highlights
significant regressions that explain more than 50% of the variance ($R^2 > 50\%$). Blue highlights
regressions that are significant ($p < 0.01$) but that explain less than 50% of the variance. The values used
for Normalized Decorrelation, i.e. (R_{input} - R_{output})/ R_{input}, and for Spiketrain Reliability (R_w) were
computed with a binning window of 10 ms, unless specified. Abbreviations: GC yo = from young mice,
GC ad = from adult mice, ALL = dataset pooling all celltypes and conditions recorded in young mice
152
153

Table S1. Intrinsic electrophysiological cell properties

y-axis ↓ / x-axis →		Membrane Capacitance (Cm)	Membrane Resistance (Rm)	Membrane Time Constant = Rm.Cm	Resting Membrane Potential (Vrest)
Normalized Decorrelation	**GC yo**	$R^2 = 4\%$ p = 0.08 slope = -0.2 F(1,100) = 3.2	$R^2 = 5\%$ p = 0.06 slope = -0.03 F(1,100) = 3.7	$R^2 = 8\%$ p = 0.013 slope = -1.2 F(1,100) = 6.5	$R^2 = 3\%$ p = 0.17 slope = -0.2 F(1,100) = 1.9
	FS	$R^2 = 47\%$ p = 0.0008 slope = -1.5 F(1,18) = 15.8	$R^2 = 77\%$ p <0.0001 slope = 0.6 F(1,18) = 58.9	$R^2 = 5\%$ p = 0.4 slope = 13.6 F(1,18) = 1.2	$R^2 = 46\%$ p = 0.0009 slope = 1.4 F(1,18) = 15.6
	HMC	$R^2 = 33\%$ p = 0.013 slope = -0.4 F(1,16) = 7.8	$R^2 = 3\%$ p = 0.48 slope = 0.06 F(1,16) = 0.5	$R^2 = 7\%$ p = 0.3 slope = -1.8 F(1,16) = 1.6	$R^2 = 0.5\%$ p = 0.8 slope = 0.1 F(1,16) = 0.08
	CA3 +gzn	$R^2 = 0.1\%$ p = 0.9 slope = -0.05 F(1,13) = 0.01	$R^2 = 2\%$ p = 0.59 slope = 0.03 F(1,13) = 0.3	$R^2 = 4\%$ p = 0.60 slope = 1.2 F(1,13) = 0.6	$R^2 = 29\%$ p = 0.04 slope = 0.7 F(1,13) = 5.4
	GC +gzn	$R^2 = 30\%$ p = 0.008 slope = 1.6 F(1,20) = 8.6	$R^2 = 0.3\%$ p = 0.8 slope = 0.01 F(1,20) = 0.06	$R^2 = 2\%$ p = 0.5 slope = 3.5 F(1,20) = 4.1	$R^2 = 13\%$ p = 0.09 slope = -0.7 F(1,20) = 3.1
	ALL	$R^2 = 0.6\%$ p = 0.35 slope = -0.09 F(1,175) = 0.9	$R^2 = 3\%$ p = 0.04 slope = 0.03 F(1,175) = 4.2	$R^2 = 1\%$ p = 0.2 slope = 0.55 F(1,175) = 1.5	$R^2 = 0.2\%$ p = 0.55 slope = -0.08 F(1,175) = 0.3
	GC ad	$R^2 = 4\%$ p = 0.24 slope = -0.4 F(1,33) = 1.4	$R^2 = 11\%$ p = 0.06 slope = -0.07 F(1,33) = 3.9	$R^2 = 17\%$ p = 0.015 slope = -4.3 F(1,33) = 6.6	$R^2 = 5\%$ p = 0.2 slope = -0.3 F(1,33) = 1.7
Spiketrain Reliability (R_W)	**GC yo**	$R^2 = 2\%$ p =0.2 slope = 0.001 F(1,100) = 1.8	$R^2 = 6\%$ p = 0.03 slope = 4e-3 F(1,100) = 4.7	$R^2 = 7\%$ p = 0.02 slope = 0.01 F(1,100) = 6.1	$R^2 = 3\%$ p = 0.17 slope = 0.002 F(1,100) = 1.9
	FS	$R^2 = 48\%$ p =0.0006, slope = 0.01 F(1,18) = 16.8	$R^2 = 70\%$ p < 0.0001 slope = -0.007 F(1,18) = 41.7	$R^2 = 6\%$ p = 0.29 slope = -0.12 F(1,18) = 0.7	$R^2 = 39\%$ p = 0.003 slope = -0.014 F(1,18) = 11.6
	HMC	$R^2 = 29\%$ p = 0.19 slope = 0.005 F(1,16) = 6.5	$R^2 = 24\%$ p = 0.06 slope = -0.001 F(1,16) = 5.0	$R^2 = 9\%$ p = 0.2 slope = 0.006 F(1,16) = 0.3	$R^2 = 3\%$ p = 0.5 slope = -0.002 F(1,16) = 0.5
	CA3 +gzn	$R^2 = 4\%$ p = 0.5 slope = 0.003 F(1,13) = 0.5	$R^2 = 9\%$ p = 0.29 slope = 6e-4 F(1,13) = 1.2	$R^2 = 4\%$ p = 0.45 slope = -0.01 F(1,13) = 0.45	$R^2 = 10\%$ p = 0.24 slope = -0.004 F(1,13) = 1.5
	GC +gzn	$R^2 = 2\%$ p = 0.5 slope = -0.003 F(1,20) = 0.4	$R^2 = 22\%$ p = 0.03 slope = 8e-4 F(1,20) = 5.7	$R^2 = 17\%$ p = 0.05 slope = -0.02 F(1,20) = 3.9	$R^2 = 29\%$ p = 0.01 slope = 0.008 F(1,20) = 8.0
	ALL	$R^2 = 0.4\%$ p = 0.5 slope = 7e-4 F(1,175) = 0.5	$R^2 = 7\%$ p = 0.0013 slope = -4e-4 F(1,175) = 10.7	$R^2 = 3\%$ p = 0.045 slope = -0.008 F(1,175) = 4.1	$R^2 = 1\%$ p = 0.24 slope = 0.001 F(1,175) = 1.4
	GC ad	$R^2 = 13\%$ p =0.03 slope = 0.006 F(1,33) = 4.9	$R^2 = 6\%$ p = 0.17 slope = 4e-4 F(1,33) = 2.0	$R^2 = 24\%$ p = 0.003 slope = 0.04 F(1,33) = 10.3	$R^2 = 8.5\%$ p = 0.09 slope = 0.004 F(1,33) = 3.05

Table S2. Spike-wise neural noise

y-axis ↓	x-axis →	Delay	Jitter	Spike Probability (SP)
Normalized Decorrelation	**GC yo**	$R^2 = 20\%$ $p < 0.0001$ slope = 1.1 $F(1,100) = 24.5$	$R^2 = 31\%$ $p < 0.0001$ slope = 1.05 $F(1,100) = 45.1$	$R^2 = 41\%$ $p < 0.0001$ slope = -37.6 $F(1,100) = 69.8$
	FS	$R^2 = 11\%$ $p = 0.15$ slope = 2.3 $F(1,18) = 2.2$	$R^2 = 18\%$ $p = 0.06$ slope = 9.6 $F(1,18) = 3.9$	$R^2 = 88\%$ $p < 0.0001$ slope = -49.1 $F(1,18) = 129.8$
	HMC	$R^2 = 9\%$ $p = 0.22$ slope = 2.1 $F(1,16) = 1.6$	$R^2 = 18\%$ $p = 0.08$ slope = 3.4 $F(1,16) = 3.5$	$R^2 = 34\%$ $p = 0.011$ slope = -42.0 $F(1,16) = 8.1$
	ALL	$R^2 = 35\%$ $p < 0.0001$ slope = 1.9 $F(1,175) = 74.3$	$R^2 = 23\%$ $p < 0.0001$ slope = 1.4 $F(1,175) = 41.0$	$R^2 = 57\%$ $p < 0.0001$ slope = -50.6 $F(1,175) = 183.8$
	GC ad	$R^2 = 2\%$ $p = 0.38$ slope = 0.9 $F(1,33) = 0.8$	$R^2 = 4\%$ $p = 0.23$ slope = -3.5 $F(1,33) = 1.4$	$R^2 = 42\%$ $p < 0.0001$ slope = -45.7 $F(1,33) = 23.8$
	Shuffle (GC yo)	$R^2 = 5\%$ $p = 0.02$ slope = 0.6 $F(1,100) = 5.8$	$R^2 = 20\%$ $p < 0.0001$ slope = 0.8 $F(1,100) = 25.6$	$R^2 = 61\%$ $p < 0.0001$ slope = -36.5 $F(1,100) = 160.0$
Spiketrain Reliability (R_W)	**GC yo**	$R^2 = 23\%$ $p < 0.0001$ slope = -0.012 $F(1,100) = 29.9$	$R^2 = 37\%$ $p < 0.0001$ slope = -0.012 $F(1,100) = 58.9$	$R^2 = 48\%$ $p < 0.0001$ slope = 0.42 94.1
	FS	$R^2 = 10\%$ $p = 0.16$ slope = -0.025 $F(1,18) = 2.1$	$R^2 = 11\%$ $p = 0.15$ slope = -0.084 $F(1,18) = 2.2$	$R^2 = 87\%$ $p < 0.0001$ slope = 0.55 $F(1,18) = 117.7$
	HMC	$R^2 = 36\%$ $p = 0.009$ slope = -0.030 $F(1,16) = 8.8$	$R^2 = 53\%$ $p = 0.0007$ slope = -0.043 $F(1,16) = 17.7$	$R^2 = 31\%$ $p = 0.016$ slope = 0.30 $F(1,16) = 7.3$
	ALL	$R^2 = 40\%$ $p < 0.0001$ slope = -0.020 $F(1,175) = 93.4$	$R^2 = 28\%$ $p < 0.0001$ slope = -0.016 $F(1,175) = 53.4$	$R^2 = 61\%$ $p < 0.0001$ slope = 0.5 $F(1,175) = 219.1$
	GC ad	$R^2 = 7\%$ $p = 0.14$ slope = -0.01 $F(1,33) = 2.3$	$R^2 = 0.03\%$ $p = 0.9$ slope = 0.003 $F(1,33) = 0.01$	$R^2 = 16\%$ $p = 0.017$ slope = 0.2 $F(1,33) = 6.3$
	Shuffle (GC yo)	$R^2 = 5\%$ $p = 0.03$ slope = -0.006 $F(1,100) = 5.0$	$R^2 = 21\%$ $p < 0.0001$ slope = -0.008 $F(1,100) = 26.2$	$R^2 = 62\%$ $p < 0.0001$ slope = 0.4 $F(1,100) = 162$

Table S3. Spiketrain-wise properties

		Overall Firing Rate	Spiketrain Reliability (R_W)
Normalized Decorrelation	**GC yo**	$R^2 = 15\%$ $p < 0.0001$ slope = - 1.3 $F(1,100) = 18.4$	$R^2 = 81\%$ $p < 0.0001$ slope = -87 $F(1,100) = 430.1$
	FS	$R^2 = 65\%$ $p < 0.0001$ slope = - 0.5 $F(1,18) = 33.5$	$R^2 = 90\%$ $p < 0.0001$ slope = -85 $F(1,18) = 169.0$
	HMC	$R^2 = 35\%$ $p = 0.0010$ slope = - 2.8 $F(1,16) = 8.6$	$R^2 = 61\%$ $p = 0.0001$ slope = -105 $F(1,16) = 25.5$
	CA3 +gzn	$R^2 = 69\%$ $p = 0.0001$ slope = - 1.7 $F(1,13) = 28.7$	$R^2 = 55\%$ $p = 0.0016$ slope = -76 $F(1,13) = 15.8$
	GC +gzn	$R^2 = 0.6\%$ $p = 0.73$ slope = - 0.2 $F(1,20) = 0.1$	$R^2 = 28\%$ $p = 0.0111$ slope = -75 $F(1,20) = 7.8$
	ALL	$R^2 = 37\%$ $p < 0.0001$ slope = - 0.8 $F(1,175) = 101.5$	$R^2 = 79\%$ $p < 0.0001$ slope = -91 $F(1,175) = 673.1$
	GC ad	$R^2 = 30\%$ $p = 0.0006$ slope = - 4.1 $F(1,33) = 14.5$	$R^2 = 52\%$ $p < 0.0001$ slope = -90 $F(1,33) = 36.2$

x-axis ➔

y-axis ⬇

Table S4. Statistics

Figure	Test	N		Descriptive stats	p-value	Degrees of freedom & F/t/z/R/etc value	other
2d	One-sample T-test	8 8 13 13 10 13 11 11 8 4 3	recorded neurons	mean +/- SEM.	< 0.0001 < 0.0001 < 0.0001 < 0.0001 < 0.0001 < 0.0001 < 0.0001 < 0.0001 < 0.0001 0.0002 0.0093	t(7) = 21.81 t(7) = 12.81 t(12) = 22.70 t(12) = 24.49 t(9) = 19.91 t(12) = 33.62 t(10) = 41.62 t(10) = 24.15 t(7) = 20.56 t(3) = 22.28 t(2) = 10.325	Rinput = 1 0.95 0.88 0.84 0.76 0.74 0.65 0.56 0.47 0.26 0.11
2d	Unbalanced one-way ANOVA	8, 8, 13, 13, 10, 13, 11, 11, 8, 4, 3	11 groups of 102 recording sets from 28 neurons	mean +/- SEM	< 0.0001	F(10,91) = 30.05	
2d	Post-hoc Tukey-Kramer Multiple-comparison	8, 8, 13, 13, 10, 13, 11, 11, 8, 4, 3	recorded neurons	mean +/- SEM	see Fig2d right		
2e	unbalanced	8, 8,	11 groups	mean +/- SEM	0.19	F(10,91) = 1.42	

	one-way ANOVA	13, 13, 10, 13, 11, 11, 8, 4, 3	of 102 recording sets (see above)				
3A4	ANCOVA	20, 61	recording sets (FS, GC)	Linear best fit, 95% CI	< 0.0001	F(2,77) = 38.70	
3A4	Unbalanced two-way ANOVA	20, 61	recording sets (FS, GC)		0.016 < 0.0001 0.72	F(4,71) = 4.86 F(1,71) = 69.65 F(4,71) = 0.52	Input groups Cell types Interaction
3A4	Post-hoc two-sample T-test with Bonferroni correction for 5 comparison groups	4, 13 4, 13 4, 13 4, 11 4, 11	Recorded neurons (FS, GC)	R_{output} Mean +/- SEM 0.26±0.05 | 0.12±0.02 0.34±0.04 | 0.17±0.02 0.40±0.04 | 0.18±0.02 0.42±0.04 | 0.21±0.02 0.45±0.04 | 0.23±0.02	0.0122 0.0001 0.0007 0.0181 0.0307	t(15) = -3.19 t(15) = -3.44 t(15) = -5.10 t(13) = -6.48 t(13) = -3.75	R_{input} = 0.88 0.84 0.74 0.65 0.56

3A4	one-sample T-test on difference between R_{output} of simultaneously recorded GC and FS	3 3 3 3 3	Pairs of recorded neurons	Mean +/- SEM -0.4038 ± 0.02 -0.3443± 0.03 -0.3266± 0.03 -0.1683± 0.002 -0.2397± 0.007	0.0024 0.0079 0.0061 0.0002 0.0008	t(2) = -20.23 t(2) = -11.18 t(2) = -12.74 t(2) = -78.87 t(2) = -35.94	R_{input} = 0.88 0.84 0.74 0.65 0.56
3B3	ANCOVA	18, 22	recording sets (HMC, GC)	Linear best fit, 95% CI	0.15	F(2,36) = 1.97	
3B3	Unbalanced two-way ANOVA	18, 22	recording sets (HMC, GC)		0.0004 0.074 0.57	F(2, 34) = 9.76 F(1, 34) = 3.39 F(2, 34) = 0.58	Input groups Cell types Interaction
3B3	Post-hoc two-sample T-test with Bonferroni correction for 3 comparison groups	10, 7 8, 5 4, 6	Recorded neurons (HMC, GC)		1 0.05 0.21	t(15) = -0.24 t(11) = -2.81 t(8) = -2.09	R_{input} = 0.76 0.26 0.11

4D	ANCOVA	15, 22	recording sets (CA3, GC)	Linear best fit, 95% CI	0.0083	F(2,33) = 5.49	
4D	Unbalanced two-way ANOVA	15, 22	recording sets (CA3, GC)		<0.0001 0.0036 0.24	F(1, 33) = 118.21 F(1, 33) = 9.82 F(1, 33) = 1.45	Input groups Cell types Interaction
4D	Post-hoc two-sample T-test with Bonferroni correction for 2 comparison groups	6, 11 9, 11	Recorded neurons (CA3, GC)		0.032 0.1	t(15) = 2.14 t(18) = 2.65	R_{input} = 0.76 0.11
5C	ANCOVA	200, 610	Recordings for all pairs of input spiketrains (FS, GC)	Linear best fit	<0.0001 <0.0001 <0.0001 <0.0001	F(2,806)=364.8 F(2,806)=59.32 F(2,806)=66.36 F(2,806) = 56.70	Timescale = 10ms 50ms 100ms 250ms

5D	ANCOVA	180, 220	Recordings for all pairs of input spiketrains (HMC, GC)	Linear best fit	<0.0001 <0.0001 <0.0001 <0.0001	F(2,396)=15.10 F(2,396)=21.34 F(2,396)=19.33 F(2,396) = 24.30	Timescale = 10ms 50ms 100ms 250ms
5E	ANCOVA	150, 220	Recordings for all pairs of input spiketrains (CA3+gzn, GC+gzn)	Linear best fit	<0.0001 <0.0001 <0.0001 <0.0001	F(2,366)=33.47 F(2,366)=17.52 F(2,366)=34.20 F(2,366) = 108.9	Timescale = 10ms 50ms 100ms 250ms
6B	One-sample T-test	8 8 13 13 10 13 11 11 8 4 3	recorded neurons	parabolic best fit	< 0.0001 0.0001 < 0.0001 < 0.0001 < 0.0001 < 0.0001 < 0.0001 < 0.0001 0.0001 0.0042 0.1125	t(7) = 11.23 t(7) = 7.43 t(12) = 12.63 t(12) = 16.615 t(9) = 8.57 t(12) = 17.59 t(10) = 13.65 t(10) = 17.72 t(7) = 8.15 t(3) = 7.91 t(2) = 2.72	Rinput = 1 0.95 0.88 0.84 0.76 0.74 0.65 0.56 0.47 0.26 0.11
6C left	ANCOVA	102, 102	Recordings	mean +/- SEM	0.33	F(2,200) = 1.09	

6C right	one-sample T-test	8 8 13 13 10 13 11 11 8 4 3	recording sets	mean +/- SEM	0.63 0.09 0.03 0.06 0.96 0.06 0.73 0.46 0.77 0.85 0.60	t(7) = -0.51 t(7) = -1.99 t(12) = -2.47 t(12) = -2.06 t(9) = -0.05 t(12) = -2.07 t(10) = 0.36 t(10) = -0.77 t(7) = -0.30 t(3) = -0.21 t(2) = -0.6	Rinput = 1 0.95 0.88 0.84 0.76 0.74 0.65 0.56 0.47 0.26 0.11
7E bottom	Monte-Carlo exact test (based on proportion of data points above 0)	800 800 1300 1300 1000 1300 1100 1100 800 400 300	GC - Shuffle recording sets	Mean +/- 90% CI	0 0 0 0 0.055 0 0.0009 0.0845 0.0088 0.26 0.5867		Rinput = 1 0.95 0.88 0.84 0.76 0.74 0.65 0.56 0.47 0.26 0.11

		N	Data	Statistic	p-value	Test statistic	Rinput
7E bottom	Monte-Carlo exact test (based on proportion of means above 0)	100 100 100 100 100 100 100 100 100 100 100 100	GC means - Shuffle means		0 0 0 0 0 0 0 0 0 0 0.01 0.85		Rinput = 1 0.95 0.88 0.84 0.76 0.74 0.65 0.56 0.47 0.26 0.11
8A	Linear regression F-test	102+20+18+15+22 = 177	Recording sets (GC,FS,HMC,CA3+gzn,GC+gzn)	$R^2 = 79.4\%$	< 0.0001	F(1, 175) = 673.1	
8D	One-sample T-test	8 8 13 13 10 13 11 11 8 4 3	Recording sets	PSTH Rout means +/- SEM	0.0013 0.0125 < 0.0001 < 0.0001 0.0006 < 0.0001 < 0.0001 < 0.0001 0.0002 0.0107 0.0905	t(7) = 5.17 t(7) = 3.335 t(12)= 6.85 t(12) = 7.65 t(9) = 5.12 t(12) = 9.05 t(10) = 8.08 t(10) = 8.32 t(7) = 7.05 t(3) = 5.69 t(2) = 3.095	Rinput = 1 0.95 0.88 0.84 0.76 0.74 0.65 0.56 0.47 0.26 0.11

S3A	Linear regression F-test	102	recording sets	R^2: 20% 31% 41%	<0.0001 <0.0001 <0.0001	$F(1,100) =$ 24.5 45.1 69.8	Delay Jitter SP
S3B	Linear regression F-test	102	recording sets	R^2: 5% 20% 61%	0.02 <0.0001 < 0.0001	$F(1,100) =$ 5.8 25.6 160.0	Delay Jitter SP
S3C	Linear regression F-test	20	recording sets	R^2: 11% 18% 88%	0.15 0.06 <0.0001	$F(1,18) =$ 2.2 3.9 129.8	Delay Jitter SP
S3D	Linear regression F-test	177	recording sets	R^2: 35% 23% 57%	<0.0001 <0.0001 <0.0001	$F(1,175) =$ 74.3 41.0 183.8	Delay Jitter SP
S4 top	Wilcoxon rank sum test	20, 61	Recording sets (FS, GC)	Mean +/- SEM	< 0.0001 <0.0001 0.0004	Z = 6.1716 Z= 5.6787 Z = -3.5212	Delay Jitter SP
S4 bottom	Wilcoxon rank sum test	18,22	Recording sets (HMC, GC)	Mean +/- SEM	0.5773 0.0267 0.0795	Z = 0.5573 Z= -2.2157 Z = -1.7535	Delay Jitter SP
S5A	ANCOVA	110, 102	recording sets (Simulation , GC)	Linear best fit	< 0.0001	F(2,208) = 92.7	
S5B	Two-sample T-test corrected for samples	110, 102	recording sets (Simulation , GC)		< 0.0001	T(104) = 15.18	

	with unequal variances					
S6(FS)	Two sample T-test	20, 61	recording sets (FS, GC)	Mean +/- SEM	< 0.0001	T(79) = -8.81
S6 (HMC)	Two sample T-test	18, 22	recording sets (HMC, GC)	Mean +/- SEM	0.0963	T(38) = -1.70
S6 (CA3)	Two sample T-test	15, 22	recording sets (FS, GC)	Mean +/- SEM	0.0052	T(35) = 2.98
S7A (FS)	Wilcoxon rank sum test	20, 61	recording sets (FS, GC)	Mean +/- SEM	< 0.0001	Z = -4.2550
S7C (FS)	One-sided Wilcoxon rank sum test	20, 61	recording sets (FS, GC)	Mean +/- SEM	< 0.0001	Z = -4.7815
S7A (HMC)	Wilcoxon rank sum test	18, 22	recording sets (HMC, GC)	Mean +/- SEM	0.0005	Z = -3.4663
S7C (HMC)	One-sided Wilcoxon rank sum test	18, 22	recording sets (HMC, GC)	Mean +/- SEM	< 0.0001	Z = -4.9732
S7A (CA3)	Wilcoxon rank sum test	15, 22	recording sets (HMC, GC)	Mean +/- SEM	0.18	Z = 1.3456

S7C (CA3)	One-sided Wilcoxon rank sum test	15, 22	recording sets (HMC, GC)	Mean +/- SEM	0.74	Z = 0.6499	
S8D	ANCOVA	20, 61	Recording sets (nbFS, GC)	Linear best fit	<0.0001	F(2,77)=30.1	
S8D	Unbalanced two-way ANOVA	20, 61	recording sets (nbFS, GC)		0.002 <0.0001 0.84	F(4, 71) = 4.7 F(1, 71) = 55.0 F(4, 71) = 0.35	Input groups Cell types Interaction
S8F	ANCOVA	180, 220	Recordings for all pairs of input spiketrains (nbHMC, GC)	Linear best fit	0.0001 <0.0001 <0.0001 <0.0001	F(2,396)=9.4 F(2,396)=18.2 F(2,396)=9.7 F(2,396) = 21.8	Timescale = 10ms 50ms 100ms 250ms