Variability in fitness effects and the limitations of fitness optimization

Christopher J Graves[1*] and Daniel M Weinreich[1*]

[1] Brown University, Department of Ecology and Evolutionary Biology and Center for

Computational and Molecular Biology. Providence, RI, USA

* Corresponding authors: christopher_graves@brown.edu and daniel_weinreich@brown.edu;

Box G-W, 80 Waterman Street

Brown University, Providence, RI 02912. Phone: 401-863-3937.

Running title: Limitations of fitness optimization

Keywords: Natural selection, Fitness optimization, Varying environments, Group selection,

Inclusive fitness, Mutation rate

Abstract

Evolutionary biologists commonly assess the evolutionary advantage of an allele based on its

effects on the lifetime survival and reproduction of individuals. However, alleles affecting traits

like sex, evolvability, and cooperation can cause fitness effects that depend heavily on

differences in the environmental, genetic, and social context of individuals carrying the allele.

This variability makes it difficult to summarize the evolutionary fate of an allele based solely on

its effects on any one individual. In this review we show how attempts to average over variability

in the fitness effects of an allele can sometimes cause misleading results. We then describe a

number of intriguing new evolutionary phenomena that have emerged in studies that explicitly

model the fate of alleles that influence long-term lineage dynamics. We conclude with prospects

for generalizations of population genetics theory and discuss how this theory might be applied to

the evolution of infectious diseases.

Contents

 I. **Introduction**

Evolution by natural selection is driven by heritable differences in the reproductive success of

individuals. However, the long-term outcome of natural selection depends not only on the effects

of an allele on individual bearers but also on its effects across its entire lineage of descendants-

defined here as the genealogy of an allele from its origination to its ultimate fixation or

extinction in the population (Sidebar 1). When fitness effects are invariant across a lineage, the

long-term fate of an allele can be deduced in a relatively straightforward manner from its

recursive effects on survival and reproduction across descendent members of the lineage. In

other cases, the evolutionary success of an allele is not an obvious consequence of its effects on

individuals. For example, variable environments can cause the same allele to have differing

effects on fitness depending on an individuals' environmental context. Similarly, fitness effects

may vary due to the presence of other alleles in the genome, which are themselves polymorphic

in the population. In such cases, it is often presumed that traits will tend to spread by natural

selection so long as they are beneficial to their carriers on average (Eshel 1973, Nunney 1999).

This implies that natural selection favors traits that are beneficial not strictly to individuals, but

to genetic lineages as a whole.

The concept that natural selection may optimize quantities related to the average success of an

allele across a lineage has arisen in a wide range of problems ranging from varying environments

to the evolution of sex and cooperation (Akçay and Van Cleve 2016, Eshel 1973, Kussell and

Leibler 2005, Lehmann, et al. 2016, Nunney 1999, Nunney 1999). In general, this idea arises

when the fitness effect of an allele varies between individual carriers, thereby limiting the ability

to infer the long-term success of an allele based on measures of individual fitness alone. A large

class of evolutionary problems fit this description and they can be classified by whether the

variability across a lineage arises due to environmental, genetic, or social factors. We outline

examples of each in Table 1 and describe them in more detail in the main text. Each source of

variation has largely been discussed within its own body of literature, where equivalent concepts

are used to describe a distinct set of adaptations, often with distinct terminology. Despite some

obvious similarities, there have been few attempts at synthesizing what is known in each of these

cases into a formal quantitative theory of the evolution of alleles with lineage-variable fitness

effects.

Averaging the fitness effects of an allele across a lineage shifts the target of adaptation from

individuals to lineages. However, one must acknowledge possible limitations in the ability of

natural selection to favor traits that confer a long-term benefit to a lineage. Specifically, natural

selection is myopic in nature- acting to increase the frequency of traits that confer an immediate

advantage to individuals without regard to their future utility to a lineage. This shortsightedness

can have dramatic consequences, particularly if it results in the permanent extinction of an allele

prior to it realizing any average benefit in the long-term. Indeed, the notion that natural selection

will act most strongly on alleles that confer a short-term advantage was championed by Maynard

Smith (1964) and Williams (1966) in their now famous critique of group selection, and is still in

use (Lynch 2007, Sniegowski and Murphy 2006). When does natural selection favor traits that

confer a long-term benefit averaged across a lineage and when does shortsighted selection limit

this ability?

After briefly summarizing results from classical, lineage-invariant theory that successfully

relates individual fitness to a lineage's eventual fate, we discuss a diversity of examples of

lineage-variable fitness, i.e., cases in which the fitness effects of an allele vary across its lineage

of descendants. We illustrate the shortcomings of averaging variability across the lineage in

finite populations, in which alleles that are beneficial in the long-term are nevertheless

vulnerable to extinction. Consequently, shortsighted selection in finite populations can limit the

ability of natural selection to optimize even these measures of fitness. Finally, we discuss other

counterintuitive results that emerge in examples where lineage-variable fitness is modeled

explicitly. These results show that the fate of an allele can be sensitive not only to its fitness

effects across a lineage, but to features unrelated to classical notions of fitness, such as the

population size. We conclude by highlighting implications for the evolution of infectious

diseases and directions for future work.

II.   Lineage-Invariant Fitness Effects

Evolutionary biologists are fundamentally concerned with understanding the outcome of natural

selection on traits that influence the survival and reproduction of their carriers. Before discussing

cases in which the fitness effects of an allele are variable across its lineage we first consider the

case where fitness effects are invariant across a lineage. Our approach throughout will be on the

field of population genetics, which has a rich tradition of analyzing dynamical models that

combine various evolutionary forces including natural selection, genetic drift, and mutation.

Such dynamical treatments of evolution provide a comprehensive analysis of a lineage – starting

from its origination in the population and ending with its ultimate fixation or extinction. We will

therefore be decidedly brief in our overview of other aspects of evolutionary theory, which

include techniques such as game theory and quantitative genetics.

Consider an allele that influences the expected number of surviving offspring produced over the

lifetime of its carriers. Formally, we allow the precise number of offspring produced by any

particular individual in this lineage to be a Poisson random variable drawn independently from

an identical distribution, with mean defined as the Wrightian fitness, w. This concept of fitness

articulates well with the Darwinian notion of fitness as lifetime reproductive success. The most

fundamental consideration regarding the fate of an allele by natural selection is whether the allele

influences this measure of fitness relative to the resident "wild type" in the population. In

population genetics, this fitness effect is most often denoted with the selection coefficient, s,

defined as the proportional change in expected number of offspring relative to the wild-type: $s \equiv$

$w_{\mathrm{mut}}/w_{\mathrm{wt}} - 1$.

Now consider a population with constant size, N. Since the number of surviving offspring born

to an individual is a random variable, we allow for random fluctuations in the number of

individuals carrying an allele as the basis for genetic drift. Assuming that generations are discrete

and non-overlapping and approximating the Poisson offspring distribution with a binomial so

that the population size remains fixed, we can describe the allele frequency dynamics using the

Wright-Fisher model. We emphasize that the Wright-Fisher process and related models capture

the interplay between natural selection and genetic drift in finite populations by incorporating

stochasticity in the number of surviving offspring born to each individual. However, by

definition all models of lineage-invariant selection assume that the distribution in that number

remains constant across the lineage (Figure 1A).

Given this framework, we can obtain solutions for a number of quantities pertaining to the fate of

a mutant allele based on its selection coefficient, s. Of particular interest given our concern with

the ultimate fate of a lineage is the probability that an allele eventually displaces all alternatives

in the population, known as the probability of fixation, P_{fix}. Kimura (Kimura 1962) found this

quantity for a mutation starting at frequency x_0, in a haploid, randomly mating population of size

N, using a continuous diffusion approximation of the Wright-Fisher process:

$$P_{\text{fix}}(N, s, x_0) = \frac{1 - e^{-Nsx_0}}{1 - e^{-Ns}}. \tag{1}$$

This result highlights many of the key features of classical population genetics theory. Solving

for the limit as s approaches zero leads to $P_{\text{fix}} = x_0$. This defines the neutral expectation that the

probability of fixation of an allele is simply equal to its starting frequency. Focusing on the case

where an allele starts from a single mutation in the population, we take $x_0 = 1/N$. Now consider a

beneficial mutation, $s > 0$. Here, $P_{\text{fix}} > 1/N$, but only asymptotically approaches s, even as

population size N tends to infinity (Haldane 1927). In other words, fixation of even a strongly

beneficial mutation is not assured, reflecting the fact genetic drift dominates allele frequency

dynamics until there are roughly $1/s$ copies in the population. This effect is worsened in small

populations since $1/s$ copies may be an appreciable fraction of the population. Thus as s or N get

small, $1/s$ approaches N and genetic drift comes to dominate selection. This result implies that

mutations are effectively neutral from the standpoint of natural selection, unless $s > 1/N$. Finally,

and somewhat less intuitively, Kimura's formula also shows that even deleterious mutations ($s <$

0), can have a nonzero fixation probability. Here again, genetic drift can overwhelm natural

selection in populations roughly no larger than $1/|s|$ individuals.

III.   Lineage-Variable Fitness Effects

Under the assumption that an allele exerts a constant, lineage-invariant effect on fitness,

Equation 1 demonstrates that a mutant's fitness effect is sufficient to predict the fate of its

lineage. We now turn to cases where variability in the fitness effects of an allele can cause this

result to fail. Examples of lineage-variable fitness effects emerge under many realistic biological

scenarios, where alleles do not act alone to influence fitness but interact with different

environmental, genetic, or social factors (Table 1, Figure 1). Consequently, the number of

offspring produced by individuals in a lineage may not be drawn from any fixed distribution,

violating the assumption of lineage invariance underlying Equation 1. We emphasize that such

variability in offspring number is beyond that captured in models like Wright-Fisher, which

typically require the distribution of offspring number to be fixed. Our goal in this section is to

highlight some of the relevant examples of variability in fitness of an allele represented by the

three classes in Table 1, and to build some intuition for how they have been handled in the

literature. We also seek to show that adaptations associated with each example depend uniquely

on the effects of an allele on the fate of a lineage rather than on individual success.

Environmental interactions

Natural environments are inherently variable and therefore present an obvious challenge to the

assumption that an allele will have the same effect on fitness for all members of a lineage.

Variation in the environment over time will cause contemporary members of a lineage to

experience the same distribution of offspring number, but this distribution now depends on time

(Figure 1B). Contrastingly, under spatial variation in the environment, contemporary members

will experience fitness effects that depend on the interaction between their shared allele and the

local environment they encounter. This again implies that no single distribution in offspring

number will be generally applicable. In either case, if environmental change is so rapid that

individuals encounter a succession of different environments in their lifetime, then fitness can be

described as a lifetime average of total survival and reproduction (Levins 1968). We will

therefore focus on the more interesting case where environments vary on a timescale greater than

the generation time of the organism; here averaging can often be misleading.

The greatest progress has been made in models of temporally varying environments, in which

case the selection coefficient s is no longer a constant, but a time-dependent quantity, $s(t)$.

Formal analysis typically requires specifying a particular form of $s(t)$ at the expense of

generality. It is commonly assumed that environments are randomly drawn from a fixed

distribution or that that the population size is infinite (Dempster 1955, Gillespie 1973, Kussell

and Leibler 2005, Lewontin and Cohen 1969). Under these assumptions, a diverse set of models

can be integrated based on how variation in fitness correlates within and between members of

two competing lineages (Frank and Slatkin 1990). We note, however, that such an approach is

limited to deriving the instantaneous change in allele frequency rather than explicitly modeling

lineage dynamics. Another consequence of assuming random environmental change and infinite

populations is that natural selection will favor alleles that increase the long-term growth rate of a

lineage, averaged over all environments (Dempster 1955, Gillespie 1973, Kussell and Leibler

2005, Lewontin and Cohen 1969, Stearns 2000). Formally, this corresponds to an increase in the

geometric mean fitness, or equivalently, the mean intrinsic growth rate (Sidebar 2), and is

generalizable to other forms of $s(t)$ (Cvijović, et al. 2015). Importantly, even arbitrarily large but

finite populations experience genetic drift, which can limit the ability to maximize the long-term

growth rate of lineages in certain environmental scenarios. We discuss these limitations in more

detail below.

Despite its limitations in finite populations, the principle that in variable environments natural

selection acts to increase geometric mean fitness is a key theoretical insight, and it is presumed

to underlie numerous adaptations. These include strategies like developmental and phenotypic

plasticity that allow adaptive phenotypic responses to environmental conditions that may not be

encountered by all individuals (Meyers and Bull 2002, Via, et al. 1995). Most notable is the

evolution of bet-hedging traits in which an allele causes the exaggeration of phenotypic noise

among members of a lineage, thereby allowing a single genotype to spread environmental risk

among different phenotypes that are suited to different environments (Fraser and Kaern 2009,

Gillespie 1974, Kussell and Leibler 2005, Philippi and Seger 1989). Such a strategy is inherently

dependent on selection favoring traits that confer a long-term benefit across a lineage, since

individuals will experience differing fitness values depending on their phenotype and the

environment they encounter. By spreading the risk of fitness losses under future environmental

uncertainty across members of a lineage, bet-hedging helps to ensure survival and reproduction

across the lineage as a whole, regardless of the environment. Examples of adaptive bet-hedging

strategies have been noted in plants (Childs, et al. 2010, Clauss and Venable 2000, Gremer and

Venable 2014), insects (Hopper 1999, Menu, et al. 2000), and microbes (Balaban, et al. 2004,

Jones and Lennon 2010, Levy, et al. 2012).

Genetic interactions

Alleles don't influence fitness alone but do so as part of an integrated genome. The genetic

background of an allele is therefore another important source of variability in fitness across a

lineage. Perhaps the most obvious example is that of epistasis (Phillips 1998), in which the

fitness effect of a mutation depends on its genetic context. Empirical evidence suggests that

epistasis among alleles is widespread (Costanzo, et al. 2016, Kryazhimskiy, et al. 2014, Wang, et

al. 2014, Weinreich, et al. 2013) and therefore provides an important source of variability in the

fitness effects of an allele, particularly in sexual populations. Similar variation in fitness can

occur in asexual populations due to secondary mutations that arise on the genetic background of

an allele as it spreads. This effect is most important in large populations or under high mutation

rates. Such conditions lead to clonal interference (Gerrish and Lenski 1998), in which multiple

asexual lineages carry competing beneficial mutations, thereby interfering with one another's

fixation. The fate of a lineage under clonal interference cannot be decided by the selection

coefficient of a single allele, but instead depends on the process of successive mutations

accumulating along a series of competing asexual lineages (Desai and Fisher 2007, Lang, et al.

2013). Indeed, this presents a major hurdle to evolving asexual populations, since the lack of

recombination implies strict genetic linkage among mutations that occur on the same

background. This lack of recombination can also lead to Muller's ratchet (Haigh 1978, Muller

1964), in which the serial fixation of deleterious mutations by genetic drift can cause fitness to

erode along an asexual lineage.

The constraints on asexual adaptation due to clonal interference and Muller's ratchet provide

strong arguments for why so many organismal life cycles include periods of recombination or

sexual reproduction. These arguments are invariably related to the idea that alleles influencing

sex can increase the long-term average evolutionary success of a lineage (Nunney 1999). This is

because sex is inherently costly to individuals, who must invest time and energy in mating and

further invest resources into the production of males, which are not capable of independent

reproduction (Maynard Smith 1978). These costs could, however, be balanced if sex increases

the long-term fitness of lineages (Nunney 1989, Nunney 1999). For example, under certain

conditions of epistasis, recombination can accelerate both the pace of adaptation (Eshel and

Feldman 1970) and the ability of populations to purge deleterious mutations and fend off

Muller's ratchet (Kondrashov 1988). Furthermore, sexual reproduction can increase rates of

adaptation by allowing beneficial mutations that arise on different backgrounds to be combined

into a single genotype, thereby limiting the constraints imposed by clonal interference (Cooper

2007, McDonald, et al. 2016). Finally, the red-queen hypothesis (Hamilton, et al. 1990, Van

Valen 1973), asserts that the constant creation of new genotypes under recombination can allow

organisms to more readily compete in a co-evolutionary arms race with parasites. Indeed, sex is

likely to have evolved for a combination of reasons and empirical observations support many of

the hypotheses that have been put forth (Colegrave 2002, Cooper 2007, Goddard, et al. 2005,

McDonald, et al. 2016, Morran, et al. 2011).

Sex and recombination are not the only processes that increase rates of adaptation. There has

been substantial recent attention on whether natural selection can act more generally on the

ability of populations to adapt, or its evolvability. Selection for evolvability is contentious, since

the ability to adapt to future contingencies is a feature of populations and would therefore appear

to require evolutionary foresight and group selection operating on biological populations (Lynch

2007, Pigliucci 2008, Sniegowski and Murphy 2006, but see Watson and Szathmáry 2016).

However, traits that increase evolvability could also arise by the process of natural selection

favoring traits that are beneficial on average, with lineages being more likely to persist over

longer evolutionary periods if they are able to adapt to future conditions (Eshel 1973). While

numerous traits could increase evolvability (Wagner and Altenberg 1996), there has been a great

deal of attention paid to the evolution of alleles which influence the mutation rate – known as

mutation rate modifiers (Denamur and Matic 2006, Sniegowski, et al. 2000). Mutation rate

modifiers have been observed in microbial populations both in the lab (Sniegowski, et al. 1997)

and in nature (LeClerc, et al. 1996, Matic, et al. 1997). The fate of such "mutator" alleles is

intriguing, since they often arise without a direct effect on fitness themselves (Chao and Cox

1983, Sniegowski, et al. 1997). In asexual populations, mutators are still physically linked to the

mutations they produce and can thereby influence the statistical properties and long-term fate of

lineages (Figure 1C). In such scenarios, evolvability arises as a by-product of indirect selection

and genetic hitchhiking of mutators (Sniegowski and Murphy 2006). However, there are also

notable exceptions in which histories of repeated environmental change could directly favor the

evolution of traits that increase evolvability. This appears to be the case in pathogens, where

elevated mutation rates in antigens to increase the capacity to adapt to a dynamic vertebrate

immune response (Graves, et al. 2013, Moxon, et al. 1994).

Social interactions

Fitness is influenced not only by environmental and genetic factors but also by interactions with

other conspecifics. These interactions can create a type of lineage-variable fitness known as

frequency dependent selection, where the fitness effects of an allele are dependent on the

frequency of the allele in the population. Frequency dependence is conveniently analyzed in the

context of evolutionary game theory (Sidebar 3), which allows one to consider the ability of an

initially rare allele to invade a population fixed for a wild-type allele (Maynard Smith 1982,

Maynard Smith and Price 1973). This approach provides a generalization of the concept of a

selection coefficient to instances where fitness cannot be wholly represented by a constant value.

A classic example of frequency-dependent selection arises when considering cooperative traits.

Here, cooperative acts incur a cost to individuals and are therefore susceptible to invasion by

selfish "cheater" strategies that avoid the cost of cooperating while still reaping the benefit.

Cheaters are typically beneficial when rare, since their fitness advantage requires interactions

with other cooperators. Despite the inherent susceptibility to cheaters, cooperation is common in

nature and is presumed to underlie major transitions in evolutionary history, such as the

evolution of multicellularity (Szathmary and Maynard Smith 1995). The mechanisms promoting

the evolution and maintenance of cooperation are therefore of long-standing interest to

biologists.

Significant theoretical progress on the evolution of cooperation arose with the formulation of

inclusive fitness theory. Hamilton (1964) showed that genes controlling cooperation may be

beneficial on average so long as the beneficiary of cooperative actions are kin, which are likely

to share the genes controlling cooperation by common descent. The key realization of this theory

is that cooperative acts need not directly increase the reproductive success of individual bearers,

but instead must increase the average effect of a gene across the lineage of cooperators (Akçay

and Van Cleve 2016). Cooperation can also be stable under cases of multi-level selection (Luo

2014, Simon, et al. 2013, Traulsen and Nowak 2006). The formation and dissolution of new

groups is itself a reproductive process and the long-term fate of a lineage is therefore sensitive to

the influence of an allele on group-level reproduction (Figure 1D). A well-known example is

infectious diseases, discussed below, in which individual cells or viral particles must replicate

within hosts and also spread among hosts to establish new infections.

There is ample empirical evidence for the stability of cooperative traits in nature if selection

favors traits that increase the long-term growth rate of a lineage at the expense of individual

fitness. For example, a large number of studies have shown how cooperation, which appears

costly to individuals, can prevail through the action of group selection and kin selection (Gore, et

al. 2009, Koschwanez, et al. 2013, Rainey and Rainey 2003, Turner and Chao 1999, Velicer, et

al. 2000). Perhaps more intriguing is the evolution of "policing" phenotypes that function to

reduce the short-term benefits of selfish cheater phenotypes and thereby stabilize cooperation

(Frank 1995, Nunney 1999, Travisano and Velicer 2004). For example, in social insects,

reproduction by the worker caste constitutes a selfish trait that can undermine colony

reproductive interests. To prevent selfish reproduction among workers, social insects have

evolved anti-cheater strategies, where colony members will systematically destroy eggs laid by

workers (Ratnieks and Visscher 1989). Tumor suppressor genes of multi-cellular organisms

perform a similar function by recognizing and destroying cells that violate normal growth

regulation and thereby preventing outgrowths of genetically selfish cancer cells (Nunney 1999).

Finally, group selection dynamics can even result in Simpson's paradox, in which the overall

frequency of cooperators increases despite their systematic tendency to decrease within groups

(Chuang, et al. 2009). The fact that a trait can spread even as it selects against in every individual

carrier shows the potential for selection on long-term growth rate to prevail over selection on

individuals.

IV.   Limitations of fitness averages

Limitations due to short-sighted selection

A central theme in many of the treatments of lineage-variable fitness effects is that fitness

differences can be averaged across a lineage using concepts like geometric mean fitness and

inclusive fitness. These extended fitness averages provide a convenient way to determine if an

allele has a positive or negative affect on a lineage- by instead determining whether it increases

long-term the rate of spread of a lineage as a whole. We also note the equivalency between these

concepts and several related averages that depict long-term growth rates. For example, pathogens

are widely assumed to maximize their long-term transmission success, R_0 (Alizon, et al. 2009,

Anderson and May 1982). Similarly, Lyapunov exponents are sometimes used to derive long-

term growth rates in variable environments (Kussell and Leibler 2005) and the concept of

invasion fitness in evolutionary game theory (Sidebar 3) indicates whether natural selection tends

to favor a trait under frequency dependence (Lehmann, et al. 2016). Similar averages have been

used to deal with variation in an allele's genetic background (Falconer 1994, Livnat and

Papadimitriou 2016). In general, averages across the variability in reproductive success are

meant to allow one to directly define an "effective" selection coefficient in order to identify

which allele increases fitness. An even more ambitious goal would be to salvage Equation 1, as

is the case under scenarios of rapid environmental change (Cvijović, et al. 2015).

Unfortunately, there are fundamental problems with the use of these averages that can preclude

natural selection from maximizing fitness averages. Specifically, shortsighted selection can drive

alleles to extinction, regardless of their long-term benefit to a lineage. This is most readily seen

in the case of a changing environment (Figure 2), where it has been noted in several contexts

(Cvijović, et al. 2015, Gerland and Hwa 2009, King and Masel 2007, Masel, et al. 2007).

Assume that a mutation arises in an environment in which it is beneficial and that the

environment is constant for τ generations. Provided it survives genetic drift, the allele will

increase in frequency following a logistic function and reach a frequency of one in

approximately $2 \cdot \ln(Ns)/s$ generations (Desai and Fisher 2007). Thus, if $\tau \gg 2 \cdot \ln(Ns)/s$, then

alleles will tend to arise and fix all in the same environment (Cvijović, et al. 2015). This provides

a straightforward threshold, beyond which natural selection is blind to the allele's long-term

benefit. Of course, this threshold is derived under the assumption of a well-mixed population of

constant size, and other factors such as demographic changes and population subdivision could

substantially extend this upper bound. Still, these considerations demonstrate an inherent time-

constraint imposed by evolution in finite populations, which only disappear as a mathematical

artifact in infinite populations (Figure 2C).

Similar limitations can be seen whenever the timescale of change in the fitness effects of an

allele are greater than the time needed for natural selection to fix alleles conferring a short-term

advantage. For example, models of multi-level selection become dominated by shortsighted

selection of selfish phenotypes whenever group-level reproductive events are rare (Luo 2014).

This breakdown in favor of shortsighted selection is analogous to that in variable environments

(compare Figures 2B and 2D) and can be understood by considering the relative effects of

individual and group selection on changes to allele frequency. Natural selection takes about s

generations to double the frequency of a selfish trait within groups, where s denotes the within-

group benefit of a selfish trait. On the other hand, increased rates of group reproduction in groups

of non-selfish individuals will double the frequency of a cooperative trait after approximately $w{\cdot}r$

generations, where r is the group-level selection coefficient and w is the number of individual

generations between group reproductive events. This heuristic reasoning implies that

shortsighted selection in favor of a selfish trait will dominate allele frequency changes and

preclude the evolution of cooperation whenever $s \gg w{\cdot}r$, which very closely matches results

derived by formal analysis (Luo 2014).

Beyond fitness averages

In addition to the role of extinction in tipping the outcome of selection toward shortsighted traits,

studies explicitly modeling variability across a lineage have yielded a number of additional

results that are not readily captured by Equation 1. Recently, Cvijović, et al. (2015) examined the

case of a periodic environment that alternates between two states. An allele that is favored in one

environment but disfavored in the next can follow unintuitive dynamics, particularly when large

changes in allele frequency occur within environmental epochs. In the classic, lineage-invariant

scenario discussed above, fixation of a neutral allele from a single starting copy requires

traversing from a starting frequency of $1/N$ to a frequency of 1 by the action of genetic drift

alone. In contrast, mutations in a fluctuating environment experience selective pressures

continually, albeit of varying signs and intensities. This means that alleles can be driven to very

high or very low frequencies by natural selection and then achieve fixation or loss due to genetic

drift with far greater probability than predicted by Equation 1. This effect can cause the fixation

probability of an allele to increase well beyond the neutral expectation of $1/N$, even when alleles

are neutral or deleterious on average. Furthermore, natural selection becomes less efficient at

recognizing long-term fitness effects- causing mutations to behave as though they were

effectively neutral, even when they are beneficial or deleterious on average. Finally, as

populations become smaller or swings in frequency more dramatic, fixation can become

independent of the average selection coefficient, creating conditions where the fixation

probability is not even a monotonically increasing function of long-term fitness.

Another intriguing result emerges when the mean reproductive success across a lineage is held

constant but its variance is altered. For example, Gillespie (1974) considered a model meant to

capture spatial variation in the environment by relaxing the assumption of a Poisson-distributed

number of offspring. Gillespie found that the natural way to quantify fitness is $w = \mu - \frac{1}{N}\sigma^2$

where μ is the mean number of offspring, σ^2 is its variance, and N is the population size. A

striking feature of this model is the appearance of population size in the definition of fitness,

which suggests that the same allele can be favored or disfavored depending solely on the

population size. This same sort of dependence on population size arises in a model of fluctuating

environments (Takahata, et al. 1975), as well as in mutators (André and Godelle 2006, Raynes, et

al. 2014, Wylie, et al. 2009). We emphasize that the population size dependence in the above

examples is distinct from that of Equation 1, where population size influences the efficiency of

natural selection but does affect its sign. Instead, variability in fitness across a lineage makes it

possible that a subset of individuals will experience strong selective pressures that are not

dominated by drift, even in small populations. This implies that genetic drift and natural

selection do not, in general, scale according the relationship in Equation 1.

Perhaps the most intriguing feature of lineage variability is the possibility that the fate of an

allele may not always be reducible to a selection coefficient at all. This is certainly the case for

the evolution of mutation rate modifiers, where the succession of *de novo* beneficial and

deleterious mutations results not only in variability in the distribution of offspring numbers

across a lineage, but also in temporal autocorrelation in this distribution among the resulting sub-

lineages (Figure 2C). Consequently, the offspring distribution is not only changing through time,

but is also inherently linked to the underlying lineage dynamics. This implies that one is unable

to define any selection coefficient for a mutator that predicts $P_{\text{fix.}}$, but must instead derive P_{fix}

directly under models that explicitly capture the dynamics of secondary mutations and clonal

interference (Good and Desai 2016). Although one could then use P_{fix} to retrospectively define

an effective coefficient for any given population size using Equation 1 (Wylie, et al. 2009), it

seems that one cannot generally define such a selection coefficient *a priori*. Moreover, even

given such an effective selection coefficient, true P_{fix} doesn't scale with N in the manner

predicted by Equation 1 (Good and Desai 2016). It remains to be seen whether a similar inability

to reduce lineage fate to any effective selection coefficient might emerge in the context of

variable environments and other examples of lineage-variable fitness effects.

V.   Implications for infectious disease evolution

One of the most promising applications of models considering lineage-variable fitness effects is

in predicting and controlling the evolution of infectious diseases. Medically important traits such

as pathogen virulence and drug resistance evolve rapidly and there has been considerable interest

in the development of evolution-proof vaccines and antibiotics (Allen, et al. 2014, Day and Read

2016, Huijben, et al. 2013, Read, et al. 2011). Pathogen lineages experience a variety of extrinsic

environmental changes including a dynamic immune response, a diverse set of tissues and hosts,

and varying exposure to drugs. Additionally, since reproduction occurs both within and between

hosts, multi-level selection can create conflicting selective pressures operating over different

timescales (Kawashima, et al. 2009, Levin and Bull 1994). Finally, the dynamic immune

response targeting antigenic epitopes has resulted in the selective pressures favoring mutator

genes capable of immune evasion and antigenic evolvability (Deitsch, et al. 2009, Graves, et al.

2013, Moxon, et al. 1994). Variability across lineages therefore appears to be the rule rather than

the exception in infectious disease evolution.

Predicting pathogen evolution and designing evolution-proof drugs will be greatly aided by

models that combine the various selective pressures operating at different levels and timescales

during the pathogen life-cycle. Traditional models have generally assumed that natural selection

will favor traits that increase the long-term epidemiological success. For example, virulence is

widely regarded as an adaptation to balance the increased rate of transmission by more

aggressive diseases with the reduced duration of infection caused by host mortality or immune

selection (Alizon, et al. 2009, Alizon and Michalakis 2015, Anderson and May 1982, Bull and

Lauring 2014). However, the assumption that natural selection will maximize transmission

success is analogous to selection maximizing other long-term measures of lineage success, like

geometric mean fitness, and is therefore sensitive to the limitations discussed above (Figure 2).

Specifically, shortsighted selection occurring within-hosts may act as a barrier for traits that

could increase long-term transmission success (Levin and Bull 1994; Sidebar 2). Indeed, models

that include mutation or competition between strains within-hosts or other ecological dynamics

have demonstrated the inability of selection to maximize transmission success (Alizon, et al.

2013, Bonhoeffer and Nowak 1994, Day 2003).

There is broad support for the prediction that shortsighted selection and selection acting to

increase traits that are beneficial on average can interact to shape infectious disease traits. For

example, empirical studies in HIV (Alizon and Fraser 2013) and enteric bacteria (Giraud, et al.

2001) show how short-sighted selection can dominate patterns of evolution and lead to

reductions in long-term transmission success. In *Salmonella enterica*, the need to maintain costly

virulence factors that are susceptible to short-sighted selection for cheaters appears to have

favored a strategy of cheater prevention that help to stabilize long-term infectivity (Diard, et al.

2013, Frank 2013, Mulder and Coombes 2013). Further theoretical progress on the role of

interaction between the differing timescales of selection in pathogens could come from models

that explicitly combine mechanistic within-host processes with long-term epidemiological

dynamics (Coombs, et al. 2007, Day and Gandon 2007, Gilchrist and Coombs 2006, Mideo, et

al. 2008). In addition, new experimental technologies such as lineage tracking of pathogens using

barcode deep-sequencing (Blundell and Levy 2014, Levy, et al. 2015) offer exciting

opportunities to measure selective pressures occurring within-hosts and integrate them with more

traditional epidemiological data.

VI.   Conclusions

Despite historical emphasis on individual fitness effects shaping the fate of an allele, such a

measure of fitness cannot always capture long-term evolutionary behavior when variability in

fitness effects arise due to environmental, genetic or social interactions (Table 1, Figure 1). In

some cases, averaging lineage-variable fitness across the various environmental, genetic, and

social contexts an allele encounters allows for the application of classical population genetics

results based on traditional notions of fitness related to individual survival and reproduction.

However, this approach can fail in finite populations where an allele's predicted fate can be

interrupted by fixation or extinction due to shortsighted selection (Figure 2B). Furthermore,

genetic drift and natural selection interact in unexpected ways when variability in fitness effects

occurs over a comparable timescale to allele frequency (Cvijović et al. 2015, Figure 2D). More

strikingly, examples from studies of mutation rate modifiers indicate that there may be no way to

summarize the direction of natural selection on an allele without simply modeling its long-term

lineage dynamics (Good and Desai 2016). Taken together, these findings may have particular

relevance for the study of infectious pathogens, where alleles are likely to experience variability

due to a combination of environmental, genetic, and social interactions.

Variability in the fitness effects of an allele challenge the conventional premise of population

genetics that individual offspring number can be drawn from a fixed distribution for all members

of a lineage (Figure 1). Cases where the typical assumption of a Poisson offspring distribution

have been relaxed (Gillespie 1974) have yielded intriguing new evolutionary properties such as

dependence on both the mean and variance in fitness effects and a critical effect of population

size in determining whether an allele is beneficial. Other examples allow properties of the

offspring distribution to vary in time, but still assume that the form of the distribution is fixed

(Cvijović, et al. 2015). In yet other cases, it appears that allele frequency dynamics cannot

always be reduced to one of independent draws from any offspring distribution, time-dependent

or not. This effect is most recognizable in mutators, where the offspring distribution changes in a

manner that is inseparable from the underlying lineage dynamics caused by secondary mutations

and selection on sub-lineages (Figure 1C). Thus, while theoretical progress has been made in

understanding processes where the offspring distribution takes on more general forms (Cannings

1974, Der, et al. 2011), we are still far from a population genetics theory with which to predict

the fate of an allele in general scenarios of lineage-variable fitness effects.

Lineage variability also highlights the need for caution when interpreting the adaptive

significance of biological traits in nature. Emphasis has often been placed on individual fitness

effects at the expense of neglecting the ability of selection to favor traits that have longer term

consequences on the fate of an allele (Williams 1966). Indeed, there are a plurality of definitions

of fitness (Orr 2009) with each generalizing the concept of fitness under a particular source of

lineage variability but none that appear sufficiently general to account for all examples. Caution

is warranted when considering traits in the context of their long-term effects on a lineage, since

such traits are inherently susceptible to shortsighted selection (Figure 2). Thus, while it is often

safe to assume that selection will favor traits on the basis of extended fitness metrics, it is also

important to consider the inherent limitations in the ability of natural selection to optimize any

measure of fitness.

Acknowledgements

The authors gratefully acknowledge Erol Akçay, Michael Desai, Steven A. Frank, Joanna Masel,

Paul Rainey, Richard Watson, Eugene Raynes and other members of the Weinreich lab for

numerous constructive comments on an earlier draft. C.J.G. is supported in part by NSF

Graduate Research Fellowship 1644760, NSF 1501355 and NSF DGE 0966060. D.M.W.

supported in part by NSF DEB-1556300 and NIH R01GM095728.

Literature cited

Akçay E, Van Cleve J. 2016. There is no fitness but fitness, and the lineage is its bearer. *Philos.*

Trans. R. Soc. Lond., B, Biol. Sci. 371: 20150085

Alizon S, de Roode JC, Michalakis Y. 2013. Multiple infections and the evolution of virulence.

Ecol. Lett. 16: 556-67

Alizon S, Fraser C. 2013. Within-host and between-host evolutionary rates across the HIV-1

genome. *Retrovirology* 10: 49

Alizon S, Hurford A, Mideo N, Van Baalen M. 2009. Virulence evolution and the trade-off

hypothesis: history, current state of affairs and the future. *J. Evol. Biol.* 22: 245-59

Alizon S, Michalakis Y. 2015. Adaptive virulence evolution: the good old fitness-based

approach. *Trends Ecol. Evol.* 30: 248-54

Allen RC, Popat R, Diggle SP, Brown SP. 2014. Targeting virulence: can we make evolution-

proof drugs? *Nature Rev. Microbiol.* 12: 300-08

Anderson RM, May RM. 1982. Coevolution of hosts and parasites. *Parasitology* 85 (Pt 2): 411-

26

583 André J-B, Godelle B. 2006. The Evolution of Mutation Rate in Finite Asexual Populations.

584   *Genetics* 172: 611-26

586 Balaban NQ, Merrin J, Chait R, Kowalik L, Leibler S. 2004. Bacterial persistence as a

587   phenotypic switch. *Science* 305: 1622-5

589 Blundell JR, Levy SF. 2014. Beyond genome sequencing: Lineage tracking with barcodes to

590   study the dynamics of evolution, infection, and cancer. *Genomics* 104: 417-30

592 Bonhoeffer S, Nowak MA. 1994. Mutation and the Evolution of Virulence. *Proc. R. Soc. B* 258:

593   133-40

595 Bull JJ, Lauring AS. 2014. Theory and empiricism in virulence evolution. *PLoS Pathog.* 10:

596   e1004387

598 Cannings C. 1974. The latent roots of certain Markov chains arising in genetics: a new approach,

599   I. Haploid models. *Adv. Appl. Probab.* 6: 260-90

601 Chao L, Cox EC. 1983. Competition between high and low mutating strains of *Escherichia coli*.

602   *Evolution* 37: 125-34

604 Childs DZ, Metcalf C, Rees M. 2010. Evolutionary bet-hedging in the real world: empirical

605   evidence and challenges revealed by plants. *Proc. R. Soc. B* 277: 3055-64

Chuang JS, Rivoire O, Leibler S. 2009. Simpson's Paradox in a Synthetic Microbial System.

Science 323: 272-75

Clauss M, Venable D. 2000. Seed germination in desert annuals: an empirical test of adaptive bet

hedging. *Am. Nat.* 155: 168-86

Colegrave N. 2002. Sex releases the speed limit on evolution. *Nature* 420: 664-66

Coombs D, Gilchrist MA, Ball CL. 2007. Evaluating the importance of within- and between-host

selection pressures on the evolution of chronic pathogens. *Theor. Popul. Biol.* 72: 576-91

Cooper TF. 2007. Recombination speeds adaptation by reducing competition between beneficial

mutations in populations of Escherichia coli. *PLoS biology* 5: e225

Costanzo M, VanderSluis B, Koch EN, Baryshnikova A, Pons C, et al. 2016. A global genetic

interaction network maps a wiring diagram of cellular function. *Science* 353: aaf1420

Cvijović I, Good BH, Jerison ER, Desai MM. 2015. Fate of a mutation in a fluctuating

environment. *Proc. Natl. Acad. Sci. USA* 112: E5021-E28

Day T. 2003. Virulence evolution and the timing of disease life-history events. *Trends Ecol.*

Evol. 18: 113-18

Day T, Gandon S. 2007. Applying population-genetic models in theoretical evolutionary

epidemiology. *Ecol. Lett.* 10: 876-88

Day T, Read AF. 2016. Does high-dose antimicrobial chemotherapy prevent the evolution of

resistance? *PLoS Comput. Biol.* 12: e1004689

Deitsch KW, Lukehart SA, Stringer JR. 2009. Common strategies for antigenic variation by

bacterial, fungal and protozoan pathogens. *Nature Rev. Microbiol.* 7: 493-503

Dempster ER. 1955. Maintenance of genetic heterogeneity. *Cold Spring Harb. Symp. Quant.*

Biol. 20: 25-31

Denamur E, Matic I. 2006. Evolution of mutation rates in bacteria. *Mol. Microbiol.* 60: 820-27

Der R, Epstein CL, Plotkin JB. 2011. Generalized population models and the nature of genetic

drift. *Theor. Popul. Biol.* 80: 80-99

Desai MM, Fisher DS. 2007. Beneficial mutation selection balance and the effect of linkage on

positive selection. *Genetics* 176: 1759-98

Diard M, Garcia V, Maier L, Remus-Emsermann MNP, Regoes RR, et al. 2013. Stabilization of

cooperative virulence by the expression of an avirulent phenotype. *Nature* 494: 353-56

Eshel I. 1973. Clone-Selection and Optimal Rates of Mutation. *J Appl. Probab.* 10: 728-38

Eshel I, Feldman MW. 1970. On the evolutionary effect of recombination. *Theor. Popul. Biol.* 1:

88-100

Eshel I, Feldman MW, Bergman A. 1998. Long-term Evolution, Short-term Evolution, and

Population Genetic Theory. *Journal of Theoretical Biology* 191: 391-96

Falconer DS. 1994. *Introduction to Quantitative Genetics*. Essex, England: Longman Scientific

and Technical. 438 pp.

Frank SA. 1995. Mutual policing and repression of competition in the evolution of cooperative

groups. *Nature* 377: 520-22

Frank SA. 2013. Microbial Evolution: Regulatory Design Prevents Cancer-like Overgrowths.

Curr. Biol. 23: R343-R46

Frank SA, Slatkin M. 1990. Evolution in a variable environment. *Am. Nat.* 136: 244-60

Fraser D, Kaern M. 2009. A chance at survival: gene expression noise and phenotypic

diversification strategies. *Mol Microbiol* 71: 1333-40

Gerland U, Hwa T. 2009. Evolutionary selection between alternative modes of gene regulation.

Proc. Natl. Acad. Sci. USA 106: 8841-46

Gerrish PJ, Lenski RE. 1998. The fate of competing beneficial mutations in an asexual

population. *Genetica* 102: 127

Gilchrist MA, Coombs D. 2006. Evolution of virulence: interdependence, constraints, and

selection using nested models. *Theor. Popul. Biol.* 69: 145-53

Gillespie JH. 1973. Natural selection with varying selection coefficients–a haploid model. *Genet.*

Res. 21: 115-20

Gillespie JH. 1974. Natural selection for within-generation variance in offspring number.

Genetics 76: 601-06

Giraud A, Matic I, Tenaillon O, Clara A, Radman M, et al. 2001. Costs and benefits of high

mutation rates: adaptive evolution of bacteria in the mouse gut. *Science* 291: 2606-8

Goddard MR, Godfray HCJ, Burt A. 2005. Sex increases the efficacy of natural selection in

experimental yeast populations. *Nature* 434: 636-40

Good BH, Desai MM. 2016. Evolution of Mutation Rates in Rapidly Adapting Asexual

Populations. *Genetics* 204: 1249-66

Gore J, Youk H, Van Oudenaarden A. 2009. Snowdrift game dynamics and facultative cheating

in yeast. *Nature* 459: 253-56

Graves CJ, Ros VI, Stevenson B, Sniegowski PD, Brisson D. 2013. Natural selection promotes

antigenic evolvability. *PLoS Pathog.* 9: e1003766

Gremer JR, Venable DL. 2014. Bet hedging in desert winter annual plants: optimal germination

strategies in a variable environment. *Ecol. Lett.* 17: 380-7

Haigh J. 1978. The accumulation of deleterious genes in a population—Muller's Ratchet. *Theor.*

Popul. Biol. 14: 251-67

Haldane JBS. 1927. A mathematical theory of natural and artificial selection. Part V: Selection

and mutation. *Proc. Camb. Phil. Soc* 23: 838-44

Hamilton WD. 1964. The genetical evolution of social behaviour. I. *Theor. Popul. Biol.* 7: 1-16

Hamilton WD, Axelrod R, Tanese R. 1990. Sexual reproduction as an adaptation to resist

parasites (a review). *Proc. Natl. Acad. Sci. USA* 87: 3566-73

Hopper KR. 1999. Risk-spreading and bet-hedging in insect population biology 1. *Annu. Rev.*

Entomol. 44: 535-60

Huijben S, Bell AS, Sim DG, Tomasello D, Mideo N, et al. 2013. Aggressive chemotherapy and

the selection of drug resistant pathogens. *PLoS Pathog* 9: e1003578

Jones SE, Lennon JT. 2010. Dormancy contributes to the maintenance of microbial diversity.

Proc Natl Acad Sci U S A 107: 5881-6

Kawashima Y, Pfafferott K, Frater J, Matthews P, Payne R, et al. 2009. Adaptation of HIV-1 to

human leukocyte antigen class I. *Nature* 458: 641-45

Kimura M. 1962. On the probability of fixation of mutant genes in a population. *Genetics* 47:

713-19

King OD, Masel J. 2007. The evolution of bet-hedging adaptations to rare scenarios. *Theor.*

Popul. Biol. 72: 560-75

Kondrashov AS. 1988. Deleterious mutations and the evolution of sex. *Nature* 336: 435-40

Koschwanez JH, Foster KR, Murray AW. 2013. Improved use of a public good selects for the

evolution of undifferentiated multicellularity. *eLife* 2: e00367

Kryazhimskiy S, Rice DP, Jerison ER, Desai MM. 2014. Global epistasis makes adaptation

predictable despite sequence-level stochasticity. *Science* 344: 1519-22

Kussell E, Leibler S. 2005. Phenotypic Diversity, Population Growth, and Information in

Fluctuating Environments. *Science* 309: 2075-78

Lang GI, Rice DP, Hickman MJ, Sodergren E, Weinstock GM, et al. 2013. Pervasive genetic

hitchhiking and clonal interference in forty evolving yeast populations. *Nature* 500: 571-

4

LeClerc JE, Li B, Payne WL, Cebula TA. 1996. High mutation frequencies among Escherichia

coli and Salmonella pathogens. *Science* 274: 1208

Lehmann L, Mullon C, Akçay E, Van Cleve J. 2016. Invasion fitness, inclusive fitness, and

reproductive numbers in heterogeneous populations. *Evolution* 70: 1689-702

Levin BR, Bull JJ. 1994. Short-sighted evolution and the virulence of pathogenic

microorganisms. *Trends in Microbiology* 2: 76-81

Levins R. 1968. *Evolution in changing environments: Some theoretical explorations*. Princeton,

NJ: Princeton University Press. 120 pp.

Levy SF, Blundell JR, Venkataram S, Petrov DA, Fisher DS, Sherlock G. 2015. Quantitative

evolutionary dynamics using high-resolution lineage tracking. *Nature* 519: 181-+

Levy SF, Ziv N, Siegal ML. 2012. Bet hedging in yeast by heterogeneous, age-correlated

expression of a stress protectant. *PLoS Biol.* 10: e1001325

Lewontin RC, Cohen D. 1969. On population growth in a randomly varying environment. *Proc.*

Natl. Acad. Sci. USA 62: 1056-60

Livnat A, Papadimitriou CH. 2016. Sex as an Algorithm: The Theory of Evolution Under the

Lens of Computation. *Commun. ACM* 59: 84-93

Luo S. 2014. A unifying framework reveals key properties of multilevel selection. *Theor. Popul.*

Biol. 341: 41-52

Lynch M. 2007. The frailty of adaptive hypotheses for the origins of organismal complexity.

Proc. Natl. Acad. Sci. USA 104: 8597-604

Masel J, King Oliver D, Maughan H. 2007. The Loss of Adaptive Plasticity during Long Periods

of Environmental Stasis. *Am. Nat.* 169: 38-46

Matic I, Radman M, Taddei F, Picard B, Doit C, et al. 1997. Highly variable mutation rates in

commensal and pathogenic Escherichia coli. *Science* 277: 1833-34

Maynard Smith J. 1964. Group Selection and Kin Selection. *Nature* 201: 1145-47

Maynard Smith J. 1978. *The evolution of sex*: Cambridge Univ Press

Maynard Smith J. 1982. *Evolution and the Theory of Games*: Cambridge Univ Press

Maynard Smith J, Price G. 1973. The logic of animal conflict. *Nature* 246: 15

McDonald MJ, Rice DP, Desai MM. 2016. Sex speeds adaptation by altering the dynamics of

molecular evolution. *Nature* 531: 233-36

Menu F, Roebuck J-P, Viala M. 2000. Bet-hedging diapause strategies in stochastic

environments. *Am. Nat.* 155: 724-34

Meyers LA, Bull JJ. 2002. Fighting change with change: adaptive variation in an uncertain

world. *Trends Ecol. Evol.* 17: 551-57

Mideo N, Alizon S, Day T. 2008. Linking within- and between-host dynamics in the

evolutionary epidemiology of infectious diseases. *Trends Ecol .Evol.* 23: 511-7

Morran LT, Schmidt OG, Gelarden IA, Parrish RC, Lively CM. 2011. Running with the Red

Queen: host-parasite coevolution selects for biparental sex. *Science* 333: 216-18

Moxon ER, Rainey PB, Nowak MA, Lenski RE. 1994. Adaptive evolution of highly mutable

loci in pathogenic bacteria. *Curr. Biol.* 4: 24-33

Mulder DT, Coombes BK. 2013. Infection biology: Cheats never prosper. *Nature* 494: 321-22

Muller HJ. 1964. The relation of recombination to mutational advance. *Mutat. Res.* 106: 2-9

Nunney L. 1989. The Maintenance of Sex by Group Selection. *Evolution* 43: 245-57

Nunney L. 1999. Lineage selection and the evolution of multistage carcinogenesis. *Proc. R. Soc.*
B 266: 493-98

Nunney L. 1999. Lineage selection: natural selection for long-term benefit. In *Levels of selection*
in evolution, ed. L Keller, pp. 238-52. Princeton, NJ: Princeton University Press

Orr HA. 2009. Fitness and its role in evolutionary genetics. *Nature Rev. Genet.* 10: 531-9

Philippi T, Seger J. 1989. Hedging one's evolutionary bets, revisited. *Trends Ecol. Evol.* 4: 41-4

Phillips PC. 1998. The language of gene interaction. *Genetics* 149: 1167-71

Pigliucci M. 2008. Is evolvability evolvable? *Nature Rev. Genet.* 9: 75-82

Rainey PB, Rainey K. 2003. Evolution of cooperation and conflict in experimental bacterial
populations. *Nature* 425: 72-74

Ratnieks FL, Visscher PK. 1989. Worker policing in the honeybee. *Nature* 342: 796-97

Raynes Y, Halstead AL, Sniegowski PD. 2014. The effect of population bottlenecks on mutation

rate evolution in asexual populations. *J. Evol. Biol.* 27: 161-69

Read AF, Day T, Huijben S. 2011. The evolution of drug resistance and the curious orthodoxy of

aggressive chemotherapy. *Proc. Natl. Acad. Sci. USA* 108: 10871-77

Simon B, Fletcher JA, Doebeli M. 2013. Towards a general theory of group selection. *Evolution*

67: 1561-72

Sniegowski PD, Gerrish PJ, Johnson T, Shaver A. 2000. The evolution of mutation rates:

separating causes from consequences. *Bioessays* 22: 1057-66

Sniegowski PD, Gerrish PJ, Lenski RE. 1997. Evolution of high mutation rates in experimental

populations of *E. coli*. *Nature* 387: 703-05

Sniegowski PD, Murphy HA. 2006. Evolvability. *Curr. Biol.* 16: R831-R34

Stearns SC. 2000. Daniel Bernoulli (1738): evolution and economics under risk. *J. Biosci.* 25:

221-8

Szathmary E, Maynard Smith J. 1995. The major evolutionary transitions. *Nature* 374: 227-32

Takahata N, Ishii K, Matsuda H. 1975. Effect of temporal fluctuation of selection coefficient on

gene frequency in a population. *Proc. Natl. Acad. Sci. USA* 72: 4541-45

Traulsen A, Hauert C. 2009. Stochastic evolutionary game dynamics. In *Reviews of nonlinear*

dynamics and complexity, ed. HG Schuster, pp. 25-61. Weinheim, Germany: Wiley-VCH

Verlag GmbH & Co.

Traulsen A, Nowak MA. 2006. Evolution of cooperation by multilevel selection. *Proc. Natl.*

Acad. Sci. USA 103: 10952-55

Travisano M, Velicer GJ. 2004. Strategies of microbial cheater control. *Trends Microbiol.* 12:

72-78

Turner PE, Chao L. 1999. Prisoner's dilemma in an RNA virus. *Nature* 398: 441-43

Van Valen L. 1973. A new evolutionary law. *Evolutionary theory* 1: 1-30

Velicer GJ, Kroos L, Lenski RE. 2000. Developmental cheating in the social bacterium

Myxococcus xanthus. *Nature* 404: 598-601

Via S, Gomulkiewicz R, De Jong G, Scheiner SM, Schlichting CD, Van Tienderen PH. 1995.

Adaptive phenotypic plasticity: consensus and controversy. *Trends Ecol. Evol.* 10: 212-7

Wagner GP, Altenberg L. 1996. Perspective: Complex adaptations and the evolution of

evolvability. *Evolution* 50: 967-76

Wang X, Fu AQ, McNerney ME, White KP. 2014. Widespread genetic epistasis among cancer

genes. *Nature communications* 5

Watson RA, Szathmáry E. 2016. How Can Evolution Learn? *Trends in Ecology & Evolution* 31:

147-57

Weinreich DM, Lan Y, Wylie CS, Heckendorn RB. 2013. Should Evolutionary Geneticists

Worry about High Order Epistasis? *Current Opinion in Development and Genetics* 23:

700-07

Williams GC. 1966. *Adaptation and natural selection: A critique of some current evolutionary*

thought: Princeton university press

Wylie CS, Ghim CM, Kessler DA, Levine H. 2009. The fixation probability of rare mutators in

finite asexual populations. *Genetics* 181: 1595-602

Sidebar 1 – What is a lineage? (Typeset near 'Introduction')

We define a lineage as the full genealogy of descendent copies of an allele starting from the

original copy and ending at its long-term fate: extinction, fixation, or maintenance as a stable

polymorphism in the population. A traditional approach in population genetics has been to

describe the long-term evolutionary fate of a mutant allele influencing some biological trait

under the combined influence of evolutionary processes like mutation, genetic drift, migration,

and natural selection. Using analytical approaches from stochastic process theory, this work

seeks to calculate the probability that such an allele ultimately reaches a frequency of one, or

achieves fixation, in the population. This approach places emphasis not solely on the individual

reproductive process but also on the long-term fate of a genetic lineage carrying the mutation. It

therefore captures a much larger class of phenomena where fitness may not be directly affected

among individual carriers of an allele but the allele instead influences the statistical properties of

a lineage (Wylie, et al. 2009). Our focus will be primarily on lineage of asexual haploid lineages,

which are easier to analyze and depict. However, the approach and definition of a lineage given

here extends naturally to sexual diploid organisms.

Sidebar 2 – Geometric mean fitness (Typeset near beginning of 'Lineage-Variable Fitness

Effects')

A widely appreciated result regarding adaptations to varying environments is the principle that

natural selection will favor traits based on their geometric mean fitness. When reproductive

success changes between generations, natural selection favors traits that increase the long-term

geometric mean fitness (GMF). Reflecting the multiplicative nature of reproduction, GMF is the

product of fitness in each generation, raised to the reciprocal of the number of generations.

Algebraically, GMF = $(\prod_{t=1}^{n} w_t)^{1/n}$, where w_t is the Wrightian fitness of a trait in generation *t*.

The same quantity can be expressed as a linear average over the natural log of this fitness value,

GMF = $\exp\left[\frac{1}{n}\sum_{t=1}^{n} \ln(w_t)\right]$. In practice, approximations are used such as GMF $\approx \mu - \sigma^2/\mu$,

where μ is the arithmetic mean fitness and σ^2 is the variance in fitness. This formula explicates

the fact that natural selection favors increases in mean fitness, but also decreases in the variance

of fitness. This implies that natural selection can be risk averse, favoring alleles with lower

variance in fitness even at the expense of decreasing fitness on average.

Sidebar 3 – Evolutionary game theory (Typeset near end of 'Lineage-Variable Fitness Effects' or

beginning of 'Limitations of Fitness averages')

Evolutionary game theory (Maynard Smith 1982) analyzes an interaction among a set of

competing alleles or "strategies" and summarizes their effect in a matrix representing the fitness

payoff of all pairwise competitions among competitors. Such a framework is most useful in the

context of frequency dependent selection, where the fitness effects of an allele are not easily

summarized by a constant selection coefficient. Such a framework provides a natural way to

determine whether a new allele starting from a single copy will tend to increase in frequency or

"invade" a population that is fixed for an alternative allele. This leads to the concept of an

evolutionarily stable strategy or ESS, which is defined as a strategy or allele that cannot be

invaded by any alternative strategy starting at an initially small frequency. The ability of an allele

to invade, or invasion fitness, is a generalization of the notion of a selection coefficient to the

case of frequency dependent selection and describes the long-term stability of a mutation against

other competing mutations (Eshel, et al. 1998, Lehmann, et al. 2016). While there are notable

exceptions (Traulsen and Hauert 2009, Traulsen and Nowak 2006), game theoretic models are

typically deterministic and describe the central tendency for allele frequency change but not the

statistical properties of lineages in finite populations.

Glossary of terms (*To appear adjacent to first use of each term or phrase*)

Lineage-variable fitness effects: Differing fitness effects of an allele between individuals due to

genetic, social, or environmental interactions.

Offspring distribution: A discrete probability distribution that captures the stochasticity in an

individual organism's reproductive success.

Cheater: An individual that benefits from cooperative interactions of other individuals without

itself contributing to the cost of cooperating.

Frequency dependent selection: A model in which the fitness of an allele depends on its

frequency in the population as a consequence of interactions between organisms.

Epistasis: The phenotypic effect of a mutation varies with genetic context.

Modifier loci: Loci responsible for genetic properties of a genome, such as mutation rate,

recombination rate and mutational robustness.

Indirect selection: Selection acting on a modifier locus mediated by genetic linkage with fitness

effects at other loci in the genome.

Clonal interference: Competition between mutational independent asexual lineages, each

carrying one or more beneficial mutations.

Genetic drift: Stochastic variation in allele frequency as a consequence of stochasticity in

reproduction inherent in finite populations.

Figure Captions

Figure 1. Variability in fitness across a lineage in diverse models. A large number of realistic

biological scenarios can result in the presence of variation in fitness across a lineage either

among contemporary individuals (vertical axis) or between individuals in time (horizontal axis).

Genealogies are shown for two competing allelic lineages indicated by circles. The focal lineage

is shaded yellow and the wild-type lineage is shaded black. Curves on the bottom of each panel

depict changes to the mean (blue) and variance (orange) of fitness across the focal lineage. **A.**

Lineage carrying a beneficial allele (yellow) rising to fixation under the classical scenario of

lineage invariance. **B.** Lineage carrying an allele that alternates from beneficial to deleterious in

a variable environment. Contemporary individuals share an identical fitness, and hence an

identical selection coefficient, but this quantity changes over time. **C.** Evolution of a mutator

lineage that experiences increased rates of both deleterious (red dots) and beneficial (grey

background) mutations. **D.** A cooperative lineage under a group selection model. Within-group

selective pressures cause the allele to be disfavored over short timescales. Groups with more

cooperative alleles tend to displace other groups over longer timescales (shown with solid grey

lines). In both **C** and **D**, fitness in the lineage varies both among contemporary individuals and

over time.

1000

Figure 2. Limitations of fitness averages in finite populations. Evolution in a periodic

environment results in four distinct regimes characterized by the relative timescale of natural

selection ($1/s$) and environmental change (τ). Simulation results of the model described by

Cvijović et al. (2015) are shown in blue and the expected change of an allele with the same

average fitness effect in a constant environment is shown in red. Unless otherwise noted,

simulations are conducted with an average selection coefficient of 0.05 and a population size of

100,000. Selection coefficients are held constant at $\pm\ 0.06$ within each environment while the

timescale of environmental change is varied (beneficial environmental epochs are shaded grey

while deleterious epochs are unshaded). **A.** When the environment changes fast relative to

changes in allele frequency (small $s\tau$), the average change in allele frequency is well

approximated by a fitness average like geometric mean fitness. **B.** When the environment

changes slower than the time of fixation of an allele (large $s\tau$), mutations tend to arise and fix all

in the same environment, regardless of their average fitness effect. **C.** In infinite populations,

averages like the geometric mean fitness are accurate regardless of the timescale of

environmental change. This is an artifact of the fact that, in the absence of genetic drift, allele

frequencies can become arbitrarily close to zero or one but never permanently achieve fixation or

extinction. **D.** Average fitness breaks down when large fluctuations in allele frequency occur on

a similar timescale to environmental change (intermediate $s\tau$). This is due to the amplification of

fluctuations by genetic drift whenever alleles reach very high or very low frequencies (Cvijović

et al. 2015). Note that genetic drift occurring as the frequency of the allele approaches 1 causes it

to respond only modestly to the second deleterious epoch. The allele subsequently achieved

fixation much sooner than would be expected on the basis of its average fitness effect.

1023

1024 **Table 1.** Sources of variability across a lineage and associated adaptations (*Typeset near*

1025 *'Introduction'*)

	Basis of variability in fitness		
	Environmental	**Genetic**	**Social**
Specific examples	Spatial variation	Sex	Kin selection
	Temporal variation	Mutation	Multi-level selection
		Clonal interference	
		Epistasis	
Adaptations	Bet-hedging	Sex/recombination	Cooperation
	Phenotypic plasticity	Mutation rate modifiers	Policing
		Evolvability	

1026

Figure 1

Figure 2