Variability in fitness effects and the limitations of lineage selection

Christopher J Graves[1*] and Daniel M Weinreich[1*]

[1] Brown University, Department of Ecology and Evolutionary Biology and Center for

Computational and Molecular Biology. Providence, RI, USA

* Corresponding authors: christopher_graves@brown.edu and daniel_weinreich@brown.edu;

Box G-W, 80 Waterman Street

Brown University, Providence, RI 02912. Phone: 401-863-3937.

Running title: Lineage selection and its limitations

Keywords: Natural selection, Lineage selection, Varying environments, Group selection,

Inclusive fitness, Mutation rate

Abstract

Natural selection is sensitive not only to the effect of a trait on total number of offspring produced but also to how a trait affects an individual's entire lineage of descendants. Here we show how a large number of seemingly disparate evolutionary problems, including sex, evolvability, and cooperation, all share the property that fitness varies among members of a lineage. This feature makes it difficult to summarize the evolutionary fate of an allele based solely on its effects on individual reproduction. We show that attempts to average over this variability are often justified, but can sometimes cause misleading results. We then describe a number of intriguing new evolutionary phenomena that have emerged in studies that explicitly model the fate of alleles that influence long-term lineage dynamics. We conclude with prospects for generalizations of population genetics theory and discuss how this theory might be applied to the evolution of infectious diseases.

Contents

I. INTRODUCTION

II. LINEAGE INVARIANT SELECTION

III. LINEAGE SELECTION

Environmental Interactions

Genetic Interactions

Social Interactions

IV. LIMITS TO LINEAGE SELECTION

Limits of fitness averages

Beyond fitness averages

V. IMPLICATIONS FOR INFECTIOUS DISEASE EVOLUTION

VI. CONCLUSIONS

I.   Introduction

Evolution by natural selection is driven by heritable differences in the reproductive success of individuals. However, the long-term outcome of natural selection depends not only on the effects of an allele on individual bearers but also on its effects across its entire lineage of descendants (Sidebar 1). When fitness effects are invariant across a lineage, the long-term fate of an allele can be deduced in a relatively straightforward manner from its recursive effects on survival and reproduction across descendent members of the lineage. In other cases, the evolutionary success of an allele is not an obvious consequence of its effects on individuals. For example, variable environments can cause the same allele to have differing effects on fitness depending on an individuals' environmental context. Similarly, fitness effects may vary due to the presence of other alleles in the genome, which are themselves polymorphic in a population. In such cases, it is often presumed that natural selection will favor traits providing a net increase in fitness averaged across a lineage via the process of lineage selection (Eshel, 1973, Nunney, 1999b). In other words, natural selection is predicted to favor traits that are beneficial not strictly to individuals, but to genetic lineages.

The notion of lineage selection has arisen independently in a variety of evolutionary problems (Kussell and Leibler, 2005, Nunney, 1999a, Akçay and Van Cleve, 2016, Lehmann et al., 2016, Eshel, 1973, Nunney, 1999b). In general, lineage selection applies to cases where the fitness effects of an allele are variable across a genetic lineage, thereby limiting the ability to infer the long-term success of an allele based on individual fitness. A large class of evolutionary problems fit this description and they can be classified based on whether the variability across a lineage

arises due to environmental, genetic, or social factors. We outline examples of each in Table 1

and describe them in more detail in the main text. Each source of variation has largely been

discussed within its own body of literature, where lineage selection or equivalent concepts are

used to describe a distinct set of adaptations, often with distinct terminology. Despite some

obvious similarities, there have been few attempts at synthesizing what is known in each of these

cases into a formal quantitative theory of lineage selection.

A common feature of lineage selection is the notion that variability in fitness across a lineage can

be averaged to define a quantity representing an "effective" fitness. We discuss the equivalence

of different approaches to averaging across a lineage below – which include familiar notions

such as geometric mean fitness and inclusive fitness. Generally, such averages shift the focus

from individuals to lineages and suggest that natural selection will favor traits that increase the

long-term average growth of a genetic lineage. However, natural selection is myopic in nature-

acting to increase the frequency of traits that confer an immediate advantage without regard to

their future utility. This shortsightedness can have dramatic consequences, particularly if it

results in the permanent extinction of an allele prior to it realizing any long-term benefit. Indeed,

the notion that natural selection will act most strongly on alleles that confer a short-term

advantage was championed by Maynard Smith (1964) and Williams (1966) in their now famous

critique of group selection, and is still in use (Sniegowski and Murphy, 2006, Lynch, 2007).

When does natural selection favor traits that confer a long-term benefit to a lineage and when

does shortsighted evolution limit this ability?

After briefly summarizing results from classical, lineage-invariant theory that successfully

relates individual fitness to a lineage's eventual fate, we discuss a diversity of examples of

lineage selection and emphasize the shared theme of variability in fitness across a lineage. We

illustrate the shortcomings of averaging variability across a lineage in the context of finite

populations, in which alleles that are beneficial in the long-term are nevertheless vulnerable to

extinction. Consequently, shortsighted evolution in finite populations can limit the ability of

natural selection to optimize even these measures of fitness. Finally, we discuss other

counterintuitive results that emerge in examples where lineage variability is modeled explicitly,

which provide more general insights into underappreciated features the evolutionary process. We

conclude by highlighting implications for the evolution of infectious diseases and directions for

future work.

II.   Lineage Invariant Selection

Evolutionary biologists are fundamentally concerned with understanding the outcome of natural

selection on traits that influence the fate of an individuals' descendent lineage. Before discussing

the realm of lineage selection – where fitness varies among members of a lineage – we will first

consider the case where fitness effects are invariant across a lineage. Our emphasis throughout

will be on the field of population genetics, which has a rich tradition of analyzing dynamical

models that combine various evolutionary forces including natural selection, genetic drift, and

mutation. This emphasis reflects not only our own expertise but also the fact that such dynamical

treatments of evolution provide a comprehensive analysis of a lineage – starting from its

origination in the population and ending with its ultimate fixation or extinction. We will

therefore be decidedly brief in our overview of other aspects of evolutionary theory, which

include techniques such as game theory and quantitative genetics that capture effects over shorter

intervals of the evolutionary process.

Consider an allele that influences the expected number of surviving offspring produced over the

lifetime of its carriers. Formally, we allow the precise number of offspring produced by any

particular individual in this lineage to be a Poisson random variable drawn independently from

an identical distribution, with mean defined as the Wrightian fitness, w. This concept of fitness

articulates well with the Darwinian notion of fitness as lifetime reproductive success. The most

fundamental consideration regarding the fate of an allele by natural selection is to consider

whether the allele influences this measure of fitness relative to the resident "wild type" in the

population. In population genetics, this fitness effect is most often denoted with the selection

coefficient, s, defined as the proportional change in expected number of offspring relative to the

wild-type: $s \equiv w_{\text{mut}}/w_{\text{wt}} - 1$.

Now consider a population with constant size, N. Since the number of surviving offspring born

to an individual is a random variable, we allow for random fluctuations in the number of

individuals carrying an allele as the basis for genetic drift. Assuming that generations are discrete

and non-overlapping we can approximate the allele frequency dynamics using the Wright-Fisher

model. We emphasize that the Wright-Fisher process and related models capture the interplay

between natural selection and genetic drift in finite populations by incorporating stochasticity in

the number of surviving offspring born to each individual. However, they assume that the

distribution in that number remains constant across a lineage, and we refer to this assumption as

lineage invariant selection (Figure 1A).

Given this framework, we can obtain solutions for a number of quantities pertaining to the fate of

a mutant allele based on its selection coefficient, *s*. Of particular interest given our concern with

the ultimate fate of a lineage is the ultimate probability that an allele displaces all alternatives in

the population, known as the probability of fixation, P_{fix}. Kimura (1962) found this quantity for a

mutation starting at frequency x_0, in a haploid, randomly mating population of size *N*, using a

continuous diffusion approximation of the Wright-Fisher process:

$$P_{fix}(N, s, x_0) = \frac{1 - e^{-Nsx_0}}{1 - e^{-Ns}}. \tag{1}$$

This result highlights many of the key features of classical population genetics theory. Focusing

on the case where an allele starts from a single mutation in the population, we will assume that x_0

= 1/*N*. Solving for the limit as s approaches zero leads to P_{fix} = 1/*N*. This defines the neutral

expectation that the probability of fixation of an allele is simply equal to its starting frequency.

Now consider a beneficial mutation, s > 0. Here, P_{fix} > 1/*N*, but only asymptotically approaches

1.0 as *s* grows, even as population size *N* tends to infinity (Haldane, 1927). In other words,

fixation of even a strongly beneficial is not assured due to the fact genetic drift dominates allele

frequency dynamics until there are roughly 1/*s* copies in the population. This effect is worsened

in small populations since 1/*s* copies may be an appreciable fraction of the population. Thus as *s*

or *N* get small, 1/*s* approaches *N* and genetic drift comes to dominate selection. This result –

known as the drift barrier – implies that mutations are effectively neutral from the standpoint of

natural selection, unless *s* > 1/*N*. Finally, and somewhat less intuitively, Kimura's formula also

shows that even deleterious mutations (*s* < 0), can have a nonzero fixation probability. Here

again, genetic drift can overwhelm natural selection in populations roughly no larger than $1/|s|$

individuals.

III.   Lineage Selection

Under the assumption that an allele exerts a constant, lineage invariant, effect on fitness,

Equation 1 demonstrates that a mutant's fitness effect is sufficient predict the fate of its lineage.

We now turn to cases of lineage selection, where variability in the fitness effects of an allele can

cause this result to fail. Lineage selection emerges under more realistic biological scenarios,

where alleles do not act alone to influence fitness but interact with different environmental,

genetic, or social factors (Table 1, Figure 1). Consequently, the number of offspring produced by

individuals in a lineage may not be drawn from any fixed distribution, violating the assumption

of lineage invariance underlying Equation 1. We emphasize that such variability in offspring

number is beyond that captured in models like Wright-Fisher, which require the distribution of

offspring number to be fixed. Our goal in this section is to highlight some of the relevant

examples of variability in fitness of an allele represented by the three classes in Table 1, and to

build some intuition for how they have been handled in the literature. We also seek to show that

adaptations associated with each example depend uniquely on the effects of an allele on the fate

of a lineage rather than on individual success.

Environmental interactions

Natural environments are inherently variable and therefore present an obvious challenge to the

assumption that an allele will have the same effect on fitness for all members of a lineage.

Variation in the environment over time will cause contemporary members of a lineage to

experience the same distribution of fitness values, but this distribution now depends on time

(Figure 1B). Contrastingly, under spatial variation in the environment contemporary members

will experience fitness effects that depend on the interaction between their shared allele and the

local environment they encounter. This implies that no single distribution in offspring number

will be generally applicable (e.g., Gillespie, 1974). In either case, if environmental change is so

rapid that individuals encounter a succession of different environments in their lifetime, then

fitness can be described as a lifetime average of total survival and reproduction (Levins, 1968).

We will therefore focus on the more interesting case where environments vary on a timescale

greater than the generation time of the organism.

The greatest progress has been made in models of temporally varying environments, in which

case the selection coefficient s is no longer a constant, but a time-dependent quantity, $s(t)$.

Formal analysis typically requires specifying a particular form of $s(t)$ at the expense of generality

across all types of variation. It is commonly assumed that environments are randomly drawn

from a fixed distribution (Lewontin and Cohen, 1969, Dempster, 1955, Gillespie, 1973, Kussell

and Leibler, 2005). Under these assumptions, a diverse set of models can be integrated based on

how variation in fitness correlates within and between members of two competing lineages

(Frank and Slatkin, 1990). We note, however, that such an approach is limited to deriving the

average direction of change in allele frequency rather than explicitly modeling lineage dynamics.

Another consequence of assuming random environmental change is that natural selection will

favor alleles that increase the long-term growth rate of a lineage, averaged over all environments

(Lewontin and Cohen, 1969, Gillespie, 1973, Stearns, 2000, Kussell and Leibler, 2005,

Dempster, 1955). Formally, this corresponds to an increase in the geometric mean fitness, or

equivalently, the mean intrinsic growth rate (Sidebar 2), and is generalizable to other forms of

$s(t)$ (Cvijović et al., 2015).

The principle that natural selection in variable environments acts to increase geometric mean

fitness is one of the key theoretical results on variable environments and it is presumed to

underlie numerous adaptations. These include strategies like developmental and phenotypic

plasticity that allow adaptive phenotypic responses to environmental conditions that may not be

encountered by all individuals (Meyers and Bull, 2002, Via et al., 1995). More notable is the

evolution of bet-hedging traits in which an allele causes the exaggeration of phenotypic noise

among members of a lineage, thereby allowing a single genotype to spread environmental risk

among different phenotypes that are suited to different environments (Philippi and Seger, 1989,

Fraser and Kaern, 2009, Gillespie, 1974, Kussell and Leibler, 2005). Such a strategy is inherently

dependent on lineage selection, since individuals will experience differing fitness values

depending on their phenotype and the environment they encounter. By spreading the risk of

fitness losses under future environmental uncertainty across members of a lineage, bet-hedging

helps to ensure survival and reproduction across the lineage as a whole, regardless of the

environment. Examples of adaptive bet-hedging strategies have been noted in plants (Gremer

and Venable, 2014, Clauss and Venable, 2000, Childs et al., 2010), insects (Hopper, 1999, Menu

et al., 2000), and microbes (Balaban et al., 2004, Jones and Lennon, 2010, Levy et al., 2012).

Genetic interactions

Alleles don't influence fitness alone but do so as part of an integrated genome. The genetic

background of an allele is therefore another important source of variability in fitness across a

lineage. Perhaps the most obvious example is that of epistasis (Phillips, 1998), in which the

fitness effect of a mutation depends on its genetic context. Empirical evidence suggests that

epistasis among alleles is widespread (Costanzo et al., 2016, Wang et al., 2014, Weinreich et al.,

2013, Kryazhimskiy et al., 2014, Mossman et al., 2016) and therefore provides an important

source of variability in the fitness effects of an allele, particularly in sexual populations. Similar

variation in fitness can occur in asexual populations due to secondary mutations that arise on the

genetic background of an allele as it spreads. This effect is most important in large populations

or under high mutation rates. Such conditions lead more generally to a condition of clonal

interference (Gerrish and Lenski, 1998), in which multiple asexual lineages carry competing

beneficial mutations, thereby interfering with one another's fixation. The fate of a lineage under

clonal interference cannot be decided by the selection coefficient of a single allele, but instead

depends on the process of successive mutations accumulating along a series of competing

asexual lineages (Lang et al., 2013, Desai and Fisher, 2007). Indeed, this presents a major hurdle

to adapting populations, since the lack of recombination implies strict genetic linkage among

mutations that occur on the same background. This lack of recombination can also lead to

Muller's ratchet (Haigh, 1978, Muller, 1964), in which the serial fixation of deleterious

mutations by genetic drift can cause fitness to erode along an asexual lineage.

The constraints on asexual adaptation due to clonal interference and Muller's ratchet provide

strong arguments for why so many organismal life cycles include periods of recombination or

sexual reproduction. Indeed, numerous theories for the evolution of sex have been put forth,

mostly relying on arguments for why sex is beneficial in terms of the long-term evolutionary

success of a lineage (Nunney, 1999b). This is because sex is inherently costly to individuals, who

must invest time and energy in mating and further invest resources into the production of males,

which are not capable of independent reproduction (Maynard Smith, 1978). These costs could,

however, be balanced by the fact that sex appears to increase the long-term adaptive potential of

lineages (Nunney, 1989, Nunney, 1999b). For example, under certain conditions of epistasis,

recombination can accelerate both the pace of adaptation (Eshel and Feldman, 1970) and the

ability of populations to purge deleterious mutations and fend off Muller's ratchet (Kondrashov,

1988). Furthermore, sexual reproduction can increase rates of adaptation by allowing beneficial

mutations that arise on different backgrounds to be combined into a single genotype, thereby

limiting the constraints imposed by clonal interference (McDonald et al., 2016, Cooper, 2007).

Finally, the red-queen hypothesis (Van Valen, 1973, Hamilton et al., 1990), asserts that the

constant creation of new genotypes under recombination can be a strategy allowing organisms to

more readily compete in a co-evolutionary arms race with parasites. Indeed, sex is likely to have

evolved for a combination of reasons and empirical observations support many of the hypotheses

that have been put forth (Colegrave, 2002, Goddard et al., 2005, Cooper, 2007, McDonald et al.,

2016, Morran et al., 2011).

Sex and recombination are not the only processes that increase rates of adaptation. There has

been substantial recent attention on whether natural selection can act more generally on the

ability of populations to adapt, or its evolvability. Selection for evolvability is contentious, since

the ability to evolve is a feature of populations and would therefore appear to require a form of

group selection operating on biological populations (Sniegowski and Murphy, 2006, Lynch,

2007, Pigliucci, 2008). However, traits that increase evolvability could also arise by the process

of lineage selection, with lineages being more likely to persist over longer evolutionary periods if

they are able to adapt to future conditions (Eshel, 1973). Since mutation is the raw material

required for adaptation, there has been a great deal of attention paid to the evolution of alleles

which influence the mutation rate – known as mutation rate modifiers (Sniegowski et al., 2000,

Denamur and Matic, 2006). Mutation rate modifiers have been observed in microbial populations

both in the lab (Sniegowski et al., 1997) and in nature (LeClerc et al., 1996, Matic et al., 1997).

The fate of such "mutator" alleles is intriguing, since they often arise without a direct effect on

fitness themselves (Chao and Cox, 1983, Sniegowski et al., 1997). In asexual populations,

mutators are still physically linked to the mutations they produce and can thereby influence the

statistical properties and long-term fate of lineages (Figure 1C). In such scenarios, evolvability

arises as a by-product of indirect selection and genetic hitchhiking of mutators (Sniegowski and

Murphy, 2006). However, there are notable exceptions in which selection on evolvability may be

more direct. This appears to be the case in pathogens, where lineage selection has favored

elevated mutation rates in antigens to increase the capacity to adapt to a dynamic vertebrate

immune response (Moxon et al., 1994, Graves et al., 2013).

Social interactions

Fitness can be influenced not only by environmental and genetic factors but also by interactions

with other conspecifics. These interactions can create a type of variability across a lineage

known as frequency dependent selection, where the fitness effects of an allele are dependent on

the frequency of the allele in the population. Frequency dependence is conveniently analyzed in

the context of evolutionary game theory (Sidebar 3), which allows one to consider the ability of

an initially rare allele to invade a population fixed for a wild-type allele (Maynard Smith, 1982,

Maynard Smith and Price, 1973). This approach provides a generalization of the concept of a

selection coefficient to instances where fitness cannot be wholly represented by a constant value.

A classic example of frequency-dependent selection arises when considering cooperative traits.

Here, cooperative acts incur a cost to individuals and are therefore susceptible to invasion by

selfish "cheater" strategies that avoid the cost of cooperating while still reaping the benefit.

Cheaters are typically beneficial when rare, since their fitness advantage requires interactions

with other cooperators. Despite the inherent susceptibility to cheaters, cooperation is common in

nature and is presumed to underlie underlying major transitions in evolutionary history, such as

the evolution of multicellularity (Szathmary and Maynard Smith, 1995). The mechanisms

promoting the evolution and maintenance of cooperation are therefore of long-standing interest

to biologists.

Significant theoretical progress on the evolution of cooperation arose with the formulation of

inclusive fitness theory. Hamilton (1964) showed that genes controlling cooperation may be

beneficial on average so long as the beneficiary of cooperative actions are kin, which are likely

to share the genes controlling cooperation by common descent. The key realization of this theory

is that cooperative acts need not directly increase the reproductive success of individual bearers,

but instead must increase the average effect of a gene across the lineage of cooperators (Akçay

and Van Cleve, 2016). Cooperation can also be stable under cases of multi-level selection

(Traulsen and Nowak, 2006, Luo, 2014, Simon et al., 2013). The formation and dissolution of

new groups is itself a reproductive process and the long-term fate of a lineage is therefore

sensitive to the influence of an allele on group-level reproduction (Figure 1D). A well-known

example is infectious diseases, discussed below, in which individual cells or viral particles

replicate within hosts but also spread among hosts to establish new infections.

There is ample empirical evidence for the stability of cooperative traits in nature due to lineage

selection. For example, a large number of studies have shown how cooperation can prevail

through the action of group selection and kin selection (Velicer et al., 2000, Koschwanez et al.,

2013, Gore et al., 2009, Turner and Chao, 1999, Rainey and Rainey, 2003). Perhaps a more

intriguing result of lineage selection is the evolution of "policing" phenotypes that function to

reduce the potential benefits to cheaters (Frank, 1995, Nunney, 1999b, Travisano and Velicer,

2004). For example, in social insects, reproduction by the worker caste constitutes a selfish trait

that can undermine colony reproductive interests. To prevent selfish reproduction among

workers, social insects have evolved anti-cheater strategies, where colony members will

systematically destroy eggs laid by workers (Ratnieks and Visscher, 1989). Tumor suppressor

genes of multi-cellular organisms perform a similar function by recognizing and destroying cells

that violate normal growth regulation and thereby preventing outgrowths of genetically selfish

cancer cells (Nunney, 1999a). Finally, group selection dynamics can even result in Simpson's

paradox (Blyth, 1972) in which the overall frequency of cooperators increases despite their

systematic tendency to decrease within groups (Chuang et al., 2009). The fact that a trait can

spread even as it selects against in every individual carrier shows the potential for lineage

selection to prevail over selection on individuals.

IV.   Limitations to lineage selection

Limitations of fitness averages

A central theme in many of the treatments of lineage selection described above is that fitness

differences can be averaged across a lineage using concepts like geometric mean fitness and

inclusive fitness. These extended fitness averages provide a convenient way to determine if an

allele has a positive or negative affect on a lineage. We also note the equivalency between these

concepts and several related averages. For example, pathogens are widely assumed to maximize

their long-term transmission success, R_0 (Anderson and May, 1982, Alizon et al., 2009).

Similarly, Lyapunov exponents are sometimes used to derive long-term growth rates in variable

environments (Kussell and Leibler, 2005) and the concept of invasion fitness in evolutionary

game theory (Sidebar 3) indicates whether natural selection tends to favor a trait under frequency

dependence (Lehmann et al., 2016). Similar averages have been used to deal with variation in an

allele's genetic background (Livnat and Papadimitriou, 2016, Falconer, 1994). In general,

averages across the variability in reproductive success are meant to allow one to directly define a

selection coefficient in order to identify which allele increases fitness. An even more ambitious

goal would be if Equation 1 could be salvaged altogether, as is the case under scenarios of rapid

environmental change (Cvijović et al., 2015).

Unfortunately, there are fundamental problems with the use of these averages that can preclude

natural selection from maximizing fitness averages. Specifically, shortsighted evolution can

drive alleles permanently extinct, regardless of their long-term benefit to a lineage. This is most

readily seen in the case of a changing environment (Figure 2), where it has been noted in several

contexts (Masel et al., 2007, King and Masel, 2007, Gerland and Hwa, 2009, Cvijović et al.,

2015). Assume that a mutation arises in an environment in which it is beneficial and that the

environment is constant for τ generations. Provided it survives genetic drift, the allele will

increase in frequency following a logistic function and reach a frequency of one in

approximately $2 \cdot \ln(Ns)/s$ generations (Desai and Fisher, 2007). Thus, if $\tau \gg 2 \cdot \ln(Ns)/s$, then

alleles will arise and fix all in the same environment (Cvijović et al., 2015). This provides a

straightforward threshold, beyond which natural selection is blind to the allele's long-term

benefit. Of course, this threshold is derived under the assumption of a well-mixed population of

constant size, and other factors such as demographic changes and population subdivision could

substantially extend this upper bound. Still, these considerations demonstrate an inherent time-

constraint imposed by evolution in finite populations, which only disappear as a mathematical

artifact in infinite populations (Figure 2C).

Similar limitations can be seen whenever the timescale of change in the fitness effects of an

allele are greater than the time needed for natural selection to fix alleles conferring a short-term

advantage. For example, models of multi-level selection become dominated by short-sighted

evolution of selfish phenotypes whenever group-level reproductive events are rare (Luo 2014,

Doebeli). This breakdown in favor of shortsighted evolution is analogous to that in variable

environments (Figure 2B) and can be understood by considering the relative effects of individual

and group selection on changes to allele frequency. Natural selection takes about s generations to

double the frequency of a selfish trait within groups, where s denotes the within-group benefit of

a selfish trait. On the other hand, increased rates of group reproduction in groups of non-selfish

individuals will double the frequency of a cooperative trait after approximately $w \cdot r$ generations,

where r is the group-level selection coefficient and w is the number of individual generations

between group reproductive events. This heuristic reasoning implies that short-sighted evolution

in favor of a selfish trait will dominate allele frequency changes and preclude the evolution of

cooperation whenever $s \gg w \cdot r$, which very closely matches results derived by formal analysis

(Luo, 2014).

Beyond fitness averages

In addition to the role of extinction in tipping the outcome of selection toward shortsighted traits,

studies explicitly modeling variability across a lineage have yielded a number of results that are

not readily captured by Equation 1. Recently, Cvijović et al. (2015) examined the case of a

periodic environment that alternates between two states. An allele that is favored in one

environment but disfavored in the next can follow unintuitive dynamics, particularly when large

changes in allele frequency occur within environmental epochs. In the classic, lineage invariant

scenario discussed above, fixation of a neutral allele from a single starting copy requires

traversing from a starting frequency of $1/N$ to a frequency of 1 by the action of genetic drift

alone. In contrast, mutations in a fluctuating environment experience selective pressures

continually, albeit of varying signs and intensities. This means that alleles can be driven to very

high or very low frequencies by natural selection and then achieve fixation or loss due to genetic

drift with far greater probability than predicted by Equation 1. This effect can cause the fixation

probability of an allele to increase well beyond the neutral expectation of $1/N$, even when alleles

are neutral or deleterious on average. Furthermore, the drift barrier is substantially greater

compared to that in a constant environment, which implies that natural selection also becomes

less efficient at distinguishing beneficial and deleterious mutations. Finally, as populations

become smaller or swings in frequency more dramatic, fixation becomes independent of the

average selection coefficient, creating conditions where the fixation probability is not even a

monotonically increasing function of long-term fitness.

Another intriguing result emerges when the mean reproductive success across a lineage is held

constant but its variance is altered. For example, Gillespie (1974) considered a model meant to

capture spatial variation in the environment by relaxing the assumption of a Poisson-distributed

number of offspring. Gillespie found that the natural way to quantify fitness is $w = \mu - \frac{1}{N}\sigma^2$

where μ is the mean number of offspring, σ^2 is its variance, and N is the population size. A

striking feature of this model is the appearance of population size in the definition of fitness,

which suggests that the same allele can be favored or disfavored depending solely on the

population size. This same sort of dependence on population size arises in a model of fluctuating

environments (Takahata et al., 1975), as well as in mutators (Wylie et al., 2009, André and

Godelle, 2006, Raynes et al., 2014). We emphasize that the population size dependence in the

above examples is distinct from that of Equation 1, where population size influences the

efficiency of natural selection but does affect its sign. Instead, variability in fitness across a

lineage makes it possible that a subset of individuals will experience strong selective pressures

that are not dominated by drift, even in small populations. This implies that genetic drift and

natural selection do not, in general, scale according the relationship in Equation 1.

Perhaps the most intriguing feature of lineage variability is the possibility that the fate of an

allele may not always be reducible to a selection coefficient at all. This is certainly the case for

the evolution of mutation rate modifiers, where the succession of *de novo* beneficial and

deleterious mutations results not only in variability in the distribution of offspring numbers

across a lineage, but also in temporal autocorrelation in this distribution among the resulting sub-

lineages (Figure 2C). Consequently, the offspring distribution is not only changing through time,

but is also inherently linked to the underlying lineage dynamics. This implies that one is unable

to define any selection coefficient for a mutator that predicts $P_{\text{fix.}}$, but must instead derive P_{fix}

directly under models that explicitly capture the dynamics of secondary mutations and clonal

interference (Good and Desai, 2016). Although one could then use P_{fix} to retrospectively define

an effective coefficient using Equation 1 (Wylie et al., 2009), it seems that one cannot generally

define such a selection coefficient *a priori*. It is conceivable that similar properties might emerge

in the context of variable environments and other examples of lineage selection, though such

results have not been described to our knowledge.

V.   Implications for infectious disease evolution

One of the most promising applications for theoretical developments related to variability in

fitness across a lineage is in predicting and controlling the evolution of infectious diseases.

Medically important traits such as pathogen virulence and drug resistance evolve rapidly and

there has been considerable interest in the development of evolution-proof vaccines and

antibiotics (Day and Read, 2016, Huijben et al., 2013, Read et al., 2011, Allen et al., 2014).

Pathogen lineages experience a variety of extrinsic environmental changes including a dynamic

immune response, a diverse set of tissues and hosts, and varying exposure to drugs. Additionally,

since reproduction occurs both within and between hosts, multi-level selection can create

conflicting selective pressures operating over different timescales (Levin and Bull, 1994,

Kawashima et al., 2009). Finally, the dynamic immune response targeting antigenic epitopes has

resulted in the selective pressures favoring mutator genes capable of immune evasion and

antigenic evolvability (Deitsch et al., 2009, Graves et al., 2013, Moxon et al., 1994). Variability

across lineages therefore appears to be the rule rather than the exception in infectious disease

evolution.

Predicting pathogen evolution and designing evolution-proof drugs will be greatly aided by

models that combine the various selective pressures operating at different levels and timescales

during the pathogen life-cycle. Traditional models have generally assumed that natural selection

will favor traits that increase the long-term epidemiological success. For example, virulence is

widely regarded as an adaptation to balance the increased rate of transmission by more

aggressive diseases with the reduced duration of infection caused by host mortality or immune

selection (Anderson and May, 1982, Alizon et al., 2009, Alizon and Michalakis, 2015, Bull and

Lauring, 2014). However, the assumption that natural selection will maximize transmission

success is analogous to selection maximizing other long-term measures of lineage success, like

geometric mean fitness, and is therefore sensitive to the limitations discussed above (Figure 2).

Specifically, shortsighted evolutionary processes occurring within-hosts may act as a barrier for

traits that could increase long-term transmission success (Levin and Bull, 1994; Sidebar 2).

Indeed, models that include mutation or competition between strains within-hosts or other

ecological dynamics have demonstrated the inability of selection to maximize transmission

success (Bonhoeffer and Nowak, 1994, Day, 2003, Alizon et al., 2013).

There is broad support for the prediction that shortsighted evolution and lineage selection can

influence the evolution of infectious diseases. For example, empirical studies in HIV (Alizon and

Fraser, 2013) and enteric bacteria (Giraud et al., 2001) show how short-sighted evolution can

dominate patterns of evolution and lead to reductions in long-term transmission success. In

Salmonella enterica, lineage selection appears to have favored a strategy to preclude shortsighted

evolution and stabilize long-term infectivity (Diard et al., 2013, Frank, 2013, Mulder and

Coombes, 2013). Further theoretical progress on the role of lineage selection in pathogens could

come from models that explicitly combine mechanistic within-host processes with long-term

epidemiological dynamics (Coombs et al., 2007, Gilchrist and Coombs, 2006, Day and Gandon,

2007, Mideo et al., 2008). In addition, new experimental technologies such as lineage tracking of

pathogens using barcode deep-sequencing (Blundell and Levy, 2014, Levy et al., 2015) offer

exciting opportunities to measure selective pressures occurring within-hosts and integrate them

with more traditional epidemiological data.

VI.   Conclusions

Evolutionary biologists have traditionally assumed that natural selection acts to favor traits that

increase individual survival and reproductive success. However, individual fitness cannot always

capture the long-term evolutionary fate of an allele when variability in fitness effects arise due to

environmental, genetic or social interactions (Table 1, Figure 1). Lineage selection seeks to

address this variability by averaging fitness across the various environmental, genetic, and social

contexts an allele encounters. However, this approach can fail in finite populations where an

allele's predicted fate can be interrupted by fixation or extinction due to shortsighted evolution

(Figure 2B). Furthermore, genetic drift and natural selection interact in unexpected ways when

variability in fitness effects occurs over a comparable timescale to allele frequency (Cvijović et

al. 2015, Figure 2D). More strikingly, examples from studies of mutation rate modifiers indicate

that there may be no way to summarize the direction of natural selection on an allele without

simply modeling its long-term lineage dynamics. Taken together, these findings may have

particular relevance for the study of infectious pathogens, where alleles are likely to experience

variability due to a combination of environmental, genetic, and social interactions.

Variability in the fitness effects of an allele challenge the conventional premise of population

genetics which assumes that individual offspring number can be drawn from a fixed distribution

for all members of a lineage (Figure 1). Cases where the typical assumption of a Poisson

offspring distribution have been relaxed (Gillespie, 1974) have yielded intriguing new

evolutionary properties such as dependence on both the mean and variance in fitness effects and

a critical effect of population size in determining whether an allele is beneficial. Other examples

allow properties of the offspring distribution to vary in time, but still assume that the form of the

distribution is fixed (Cvijović et al., 2015). In other cases, it appears that allele frequency

dynamics cannot always be reduced to one of independent draws from any offspring distribution,

time-dependent or not. This effect is most recognizable in mutators, where the offspring

distribution changes in a manner that is inseparable from the underlying lineage dynamics caused

by secondary mutations and selection on sub-lineages (Figure 1C). Thus, while theoretical

progress has been in understanding processes where the offspring distribution takes on more

general forms (Cannings, 1974, Der et al., 2012, Der et al., 2011), we are still far from a

population genetics theory with which to predict the fate of an allele in general scenarios of

lineage selection.

Lineage variability also highlights the need for caution when interpreting the adaptive

significance of biological traits in nature. Emphasis has often been placed on individual fitness

effects at the expense of neglecting the ability of selection to favor traits that have longer term

consequences on the fate of an allele (Williams, 1966). Indeed, there are a plurality of definitions

of fitness (Orr, 2009) with each generalizing the concept of fitness under a particular source of

lineage variability but none that appear sufficiently general to account for all cases of lineages

selection. Caution is warranted when considering traits in the context of their long-term effects

on a lineage, since such traits are inherently susceptible to shortsighted evolution (Figure 2).

Thus, while it is often safe to assume that selection will favor traits on the basis of extended

fitness metrics, it is also important to consider the inherent limitations in the ability of natural

selection to optimize any measure of fitness.

Literature cited

Akçay, E. & Van Cleve, J. 2016. There is no fitness but fitness, and the lineage is its bearer.

Philos. Trans. R. Soc. Lond., B, Biol. Sci., 371**:**20150085.

Alizon, S., De Roode, J. C. & Michalakis, Y. 2013. Multiple infections and the evolution of

virulence. *Ecol. Lett.,* 16**:**556-567.

Alizon, S. & Fraser, C. 2013. Within-host and between-host evolutionary rates across the HIV-1

genome. *Retrovirology,* 10**:**49.

Alizon, S., Hurford, A., Mideo, N. & Van Baalen, M. 2009. Virulence evolution and the trade-

off hypothesis: History, current state of affairs and the future. *J. Evol. Biol.,* 22**:**245-59.

Alizon, S. & Michalakis, Y. 2015. Adaptive virulence evolution: The good old fitness-based

approach. *Trends Ecol. Evol.,* 30**:**248-54.

Allen, R. C., Popat, R., Diggle, S. P. & Brown, S. P. 2014. Targeting virulence: Can we make

evolution-proof drugs? *Nature Rev. Microbiol.,* 12**:**300-308.

Anderson, R. M. & May, R. M. 1982. Coevolution of hosts and parasites. *Parasitology,* 85 (Pt

2)**:**411-26.

André, J.-B. & Godelle, B. 2006. The evolution of mutation rate in finite asexual populations.

Genetics, 172**:**611-626.

Balaban, N. Q., Merrin, J., Chait, R., Kowalik, L. & Leibler, S. 2004. Bacterial persistence as a

phenotypic switch. *Science,* 305**:**1622-5.

Blundell, J. R. & Levy, S. F. 2014. Beyond genome sequencing: Lineage tracking with barcodes

to study the dynamics of evolution, infection, and cancer. *Genomics,* 104**:**417-430.

Blyth, C. R. 1972. On Simpson's paradox and the sure-thing principle. *J. Amer. Statist. Assoc.,*

67**:**364-366.

Bonhoeffer, S. & Nowak, M. A. 1994. Mutation and the evolution of virulence. *Proc. R. Soc. B,*

258**:**133-140.

Bull, J. J. & Lauring, A. S. 2014. Theory and empiricism in virulence evolution. *PLoS Pathog.,*

10**:**e1004387.

Cannings, C. 1974. The latent roots of certain markov chains arising in genetics: A new

approach, i. Haploid models. *Adv. Appl. Probab.,* 6**:**260-290.

Chao, L. & Cox, E. C. 1983. Competition between high and low mutating strains of *Escherichia*

coli. *Evolution,* 37**:**125-134.

Childs, D. Z., Metcalf, C. & Rees, M. 2010. Evolutionary bet-hedging in the real world:

Empirical evidence and challenges revealed by plants. *Proc. R. Soc. B,* 277**:**3055-64.

Chuang, J. S., Rivoire, O. & Leibler, S. 2009. Simpson's paradox in a synthetic microbial system.

Science, 323**:**272-275.

Clauss, M. & Venable, D. 2000. Seed germination in desert annuals: An empirical test of

adaptive bet hedging. *Am. Nat.,* 155**:**168-186.

Colegrave, N. 2002. Sex releases the speed limit on evolution. *Nature,* 420**:**664-666.

Coombs, D., Gilchrist, M. A. & Ball, C. L. 2007. Evaluating the importance of within- and

between-host selection pressures on the evolution of chronic pathogens. *Theor. Popul.*

Biol., 72**:**576-91.

Cooper, T. F. 2007. Recombination speeds adaptation by reducing competition between

beneficial mutations in populations of *Escherichia coli*. *PLoS Biology,* 5**:**e225.

Costanzo, M., Vandersluis, B., Koch, E. N., Baryshnikova, A., Pons, C., Tan, G., Wang, W., et

al. 2016. A global genetic interaction network maps a wiring diagram of cellular function.

Science, 353**:**aaf1420.

Cvijović, I., Good, B. H., Jerison, E. R. & Desai, M. M. 2015. Fate of a mutation in a fluctuating

environment. *Proc. Natl. Acad. Sci. USA,* 112**:**E5021-E5028.

Day, T. 2003. Virulence evolution and the timing of disease life-history events. *Trends Ecol.*

Evol., 18**:**113-118.

Day, T. & Gandon, S. 2007. Applying population-genetic models in theoretical evolutionary

epidemiology. *Ecol. Lett.,* 10**:**876-888.

Day, T. & Read, A. F. 2016. Does high-dose antimicrobial chemotherapy prevent the evolution

of resistance? *PLoS Comput. Biol.,* 12**:**e1004689.

Deitsch, K. W., Lukehart, S. A. & Stringer, J. R. 2009. Common strategies for antigenic

variation by bacterial, fungal and protozoan pathogens. *Nature Rev. Microbiol.,* 7**:**493-

503.

Dempster, E. R. 1955. Maintenance of genetic heterogeneity. *Cold Spring Harb. Symp. Quant.*

Biol. , 20**:**25-31.

Denamur, E. & Matic, I. 2006. Evolution of mutation rates in bacteria. *Mol. Microbiol.,* 60**:**820-

827.

Der, R., Epstein, C. & Plotkin, J. B. 2012. Dynamics of neutral and selected alleles when the

offspring distribution is skewed. *Genetics,* 191**:**1331-1344.

Der, R., Epstein, C. L. & Plotkin, J. B. 2011. Generalized population models and the nature of

genetic drift. *Theor. Popul. Biol.,* 80**:**80-99.

Desai, M. M. & Fisher, D. S. 2007. Beneficial mutation selection balance and the effect of

linkage on positive selection. *Genetics,* 176**:**1759-98.

Diard, M., Garcia, V., Maier, L., Remus-Emsermann, M. N. P., Regoes, R. R., Ackermann, M. &

Hardt, W.-D. 2013. Stabilization of cooperative virulence by the expression of an

avirulent phenotype. *Nature,* 494**:**353-356.

Eshel, I. 1973. Clone-selection and optimal rates of mutation. *J Appl. Probab.,* 10**:**728-738.

Eshel, I. & Feldman, M. W. 1970. On the evolutionary effect of recombination. *Theor. Popul.*

Biol., 1**:**88-100.

Falconer, D. S. 1994. *Introduction to quantitative genetics,* Essex, England, Longman Scientific

and Technical.

Frank, S. A. 1995. Mutual policing and repression of competition in the evolution of cooperative

groups. *Nature,* 377**:**520-522.

Frank, Steven a. 2013. Microbial evolution: Regulatory design prevents cancer-like overgrowths.

Curr. Biol., 23**:**R343-R346.

Frank, S. A. & Slatkin, M. 1990. Evolution in a variable environment. *Am. Nat.,* 136**:**244-260.

Fraser, D. & Kaern, M. 2009. A chance at survival: Gene expression noise and phenotypic

diversification strategies. *Mol. Microbiol.,* 71**:**1333-40.

Gerland, U. & Hwa, T. 2009. Evolutionary selection between alternative modes of gene

regulation. *Proc. Natl. Acad. Sci. USA,* 106**:**8841-8846.

Gerrish, P. J. & Lenski, R. E. 1998. The fate of competing beneficial mutations in an asexual

population. *Genetica,* 102**:**127.

Gilchrist, M. A. & Coombs, D. 2006. Evolution of virulence: Interdependence, constraints, and

selection using nested models. *Theor. Popul. Biol.,* 69**:**145-53.

Gillespie, J. H. 1973. Natural selection with varying selection coefficients–a haploid model.

Genet. Res., 21**:**115-120.

Gillespie, J. H. 1974. Natural selection for within-generation variance in offspring number.

Genetics, 76**:**601-606.

Giraud, A., Matic, I., Tenaillon, O., Clara, A., Radman, M., Fons, M. & Taddei, F. 2001. Costs

and benefits of high mutation rates: Adaptive evolution of bacteria in the mouse gut.

Science, 291**:**2606-8.

Goddard, M. R., Godfray, H. C. J. & Burt, A. 2005. Sex increases the efficacy of natural

selection in experimental yeast populations. *Nature,* 434**:**636-640.

Good, B. H. & Desai, M. M. 2016. Evolution of mutation rates in rapidly adapting asexual

populations. *Genetics,* 204**:**1249-66.

Gore, J., Youk, H. & Van Oudenaarden, A. 2009. Snowdrift game dynamics and facultative

cheating in yeast. *Nature,* 459**:**253-6.

Graves, C. J., Ros, V. I., Stevenson, B., Sniegowski, P. D. & Brisson, D. 2013. Natural selection

promotes antigenic evolvability. *PLoS Pathog.,* 9**:**e1003766.

Gremer, J. R. & Venable, D. L. 2014. Bet hedging in desert winter annual plants: Optimal

germination strategies in a variable environment. *Ecol. Lett.,* 17**:**380-7.

Haigh, J. 1978. The accumulation of deleterious genes in a population—Muller's ratchet. *Theor.*

Popul. Biol., 14**:**251-267.

Haldane, J. B. S. 1927. A mathematical theory of natural and artificial selection. Part V:

Selection and mutation. *Proc. Camb. Phil. Soc,* 23**:**838-844.

Hamilton, W. D. 1964. The genetical evolution of social behaviour. I. *Theor. Popul. Biol.,* 7**:**1-

16.

Hamilton, W. D., Axelrod, R. & Tanese, R. 1990. Sexual reproduction as an adaptation to resist

parasites (a review). *Proc. Natl. Acad. Sci. USA,* 87**:**3566-3573.

Hopper, K. R. 1999. Risk-spreading and bet-hedging in insect population biology. *Annu. Rev.*

Entomol., 44**:**535-560.

Huijben, S., Bell, A. S., Sim, D. G., Tomasello, D., Mideo, N., Day, T. & Read, A. F. 2013.

Aggressive chemotherapy and the selection of drug resistant pathogens. *PLoS Pathog.,*

9**:**e1003578.

Jones, S. E. & Lennon, J. T. 2010. Dormancy contributes to the maintenance of microbial

diversity. *Proc. Natl. Acad. Sci. USA,* 107**:**5881-6.

Kawashima, Y., Pfafferott, K., Frater, J., Matthews, P., Payne, R., Addo, M., Gatanaga, H., et al.

2009. Adaptation of HIV-1 to human leukocyte antigen class I. *Nature,* 458**:**641-645.

Kimura, M. 1962. On the probability of fixation of mutant genes in a population. *Genetics,*

47**:**713-719.

King, O. D. & Masel, J. 2007. The evolution of bet-hedging adaptations to rare scenarios. *Theor.*

Popul. Biol., 72**:**560-575.

Kondrashov, A. S. 1988. Deleterious mutations and the evolution of sex. *Nature,* 336**:**435-440.

Koschwanez, J. H., Foster, K. R. & Murray, A. W. 2013. Improved use of a public good selects

for the evolution of undifferentiated multicellularity. *eLife,* 2**:**e00367.

Kryazhimskiy, S., Rice, D. P., Jerison, E. R. & Desai, M. M. 2014. Global epistasis makes

adaptation predictable despite sequence-level stochasticity. *Science,* 344**:**1519-1522.

Kussell, E. & Leibler, S. 2005. Phenotypic diversity, population growth, and information in

fluctuating environments. *Science,* 309**:**2075-2078.

Lang, G. I., Rice, D. P., Hickman, M. J., Sodergren, E., Weinstock, G. M., Botstein, D. & Desai,

M. M. 2013. Pervasive genetic hitchhiking and clonal interference in forty evolving yeast

populations. *Nature,* 500**:**571-4.

Leclerc, J. E., Li, B., Payne, W. L. & Cebula, T. A. 1996. High mutation frequencies among

Escherichia coli and *Salmonella* pathogens. *Science,* 274**:**1208-11.

Lehmann, L., Mullon, C., Akçay, E. & Van Cleve, J. 2016. Invasion fitness, inclusive fitness,

and reproductive numbers in heterogeneous populations. *Evolution,* 70**:**1689-1702.

Levin, B. R. & Bull, J. J. 1994. Short-sighted evolution and the virulence of pathogenic

microorganisms. *Trends Microbiol.,* 2**:**76-81.

Levins, R. 1968. *Evolution in changing environments: Some theoretical explorations,* Princeton,

NJ, Princeton University Press.

Levy, S. F., Blundell, J. R., Venkataram, S., Petrov, D. A., Fisher, D. S. & Sherlock, G. 2015.

Quantitative evolutionary dynamics using high-resolution lineage tracking. *Nature,*

519**:**181-6.

Levy, S. F., Ziv, N. & Siegal, M. L. 2012. Bet hedging in yeast by heterogeneous, age-correlated

expression of a stress protectant. *PLoS Biol.,* 10**:**e1001325.

Lewontin, R. C. & Cohen, D. 1969. On population growth in a randomly varying environment.

Proc. Natl. Acad. Sci. USA, 62**:**1056-1060.

Livnat, A. & Papadimitriou, C. H. 2016. Sex as an algorithm: The theory of evolution under the

lens of computation. *Commun. ACM,* 59**:**84-93.

Luo, S. 2014. A unifying framework reveals key properties of multilevel selection. *Theor. Popul.*

Biol., 341**:**41-52.

Lynch, M. 2007. The frailty of adaptive hypotheses for the origins of organismal complexity.

Proc. Natl. Acad. Sci. USA, 104**:**8597-8604.

Masel, J., King, Oliver d. & Maughan, H. 2007. The loss of adaptive plasticity during long

periods of environmental stasis. *Am. Nat.,* 169**:**38-46.

Matic, I., Radman, M., Taddei, F., Picard, B., Doit, C., Bingen, E., Denamur, E., et al. 1997.

Highly variable mutation rates in commensal and pathogenic *Escherichia coli*. *Science,*

277**:**1833-1834.

Maynard Smith, J. 1964. Group selection and kin selection. *Nature,* 201**:**1145-1147.

Maynard Smith, J. 1978. *The evolution of sex*, Cambridge Univ Press.

Maynard Smith, J. 1982. *Evolution and the theory of games*, Cambridge University Press.

Maynard Smith, J. & Price, G. 1973. The logic of animal conflict. *Nature,* 246**:**15.

McDonald, M. J., Rice, D. P. & Desai, M. M. 2016. Sex speeds adaptation by altering the

dynamics of molecular evolution. *Nature,* 531**:**233-236.

Menu, F., Roebuck, J.-P. & Viala, M. 2000. Bet-hedging diapause strategies in stochastic

environments. *Am. Nat.,* 155**:**724-734.

Meyers, L. A. & Bull, J. J. 2002. Fighting change with change: Adaptive variation in an

uncertain world. *Trends Ecol. Evol.,* 17**:**551-557.

Mideo, N., Alizon, S. & Day, T. 2008. Linking within- and between-host dynamics in the

evolutionary epidemiology of infectious diseases. *Trends Ecol .Evol.,* 23**:**511-7.

Morran, L. T., Schmidt, O. G., Gelarden, I. A., Parrish, R. C. & Lively, C. M. 2011. Running

with the red queen: Host-parasite coevolution selects for biparental sex. *Science,*

333**:**216-218.

Mossman, J. A., Biancani, L. M., Zhu, C.-T. & Rand, D. M. 2016. Mitonuclear epistasis for

development time and its modification by diet in *Drosophila*. *Genetics,* 203**:**463-84.

Moxon, E. R., Rainey, P. B., Nowak, M. A. & Lenski, R. E. 1994. Adaptive evolution of highly

mutable loci in pathogenic bacteria. *Curr. Biol.,* 4**:**24-33.

Mulder, D. T. & Coombes, B. K. 2013. Infection biology: Cheats never prosper. *Nature,*

494**:**321-322.

Muller, H. J. 1964. The relation of recombination to mutational advance. *Mutat. Res.,* 106**:**2-9.

Nunney, L. 1989. The maintenance of sex by group selection. *Evolution,* 43**:**245-257.

Nunney, L. 1999a. Lineage selection and the evolution of multistage carcinogenesis. *Proc. R.*

Soc. B, 266**:**493-498.

Nunney, L. 1999b. Lineage selection: Natural selection for long-term benefit. *In:* Keller, L. (ed.)

Levels of selection in evolution. Princeton, NJ: Princeton University Press.

Orr, H. A. 2009. Fitness and its role in evolutionary genetics. *Nature Rev. Genet.,* 10**:**531-9.

Philippi, T. & Seger, J. 1989. Hedging one's evolutionary bets, revisited. *Trends Ecol. Evol.,*

4**:**41-4.

Phillips, P. C. 1998. The language of gene interaction. *Genetics,* 149**:**1167-1171.

Pigliucci, M. 2008. Is evolvability evolvable? *Nature Rev. Genet.,* 9**:**75-82.

Rainey, P. B. & Rainey, K. 2003. Evolution of cooperation and conflict in experimental bacterial

populations. *Nature,* 425**:**72-74.

Ratnieks, F. L. & Visscher, P. K. 1989. Worker policing in the honeybee. *Nature,* 342**:**796-797.

Raynes, Y., Halstead, A. L. & Sniegowski, P. D. 2014. The effect of population bottlenecks on

mutation rate evolution in asexual populations. *J. Evol. Biol.,* 27**:**161-169.

Read, A. F., Day, T. & Huijben, S. 2011. The evolution of drug resistance and the curious

orthodoxy of aggressive chemotherapy. *Proc. Natl. Acad. Sci. USA,* 108**:**10871-10877.

Simon, B., Fletcher, J. A. & Doebeli, M. 2013. Towards a general theory of group selection.

Evolution, 67**:**1561-1572.

Sniegowski, P. D., Gerrish, P. J., Johnson, T. & Shaver, A. 2000. The evolution of mutation

rates: Separating causes from consequences. *Bioessays,* 22**:**1057-1066.

Sniegowski, P. D., Gerrish, P. J. & Lenski, R. E. 1997. Evolution of high mutation rates in

experimental populations of *E. coli*. *Nature,* 387**:**703-705.

Sniegowski, P. D. & Murphy, H. A. 2006. Evolvability. *Curr. Biol.,* 16**:**R831-R834.

Stearns, S. C. 2000. Daniel Bernoulli (1738): Evolution and economics under risk. *J. Biosci.,*

25**:**221-8.

Szathmary, E. & Maynard Smith, J. 1995. The major evolutionary transitions. *Nature,* 374**:**227-

232.

Takahata, N., Ishii, K. & Matsuda, H. 1975. Effect of temporal fluctuation of selection

coefficient on gene frequency in a population. *Proc. Natl. Acad. Sci. USA,* 72**:**4541-4545.

Traulsen, A. & Hauert, C. 2009. Stochastic evolutionary game dynamics. *In:* Schuster, H. G.

(ed.) *Reviews of nonlinear dynamics and complexity.* Weinheim, Germany: Wiley-VCH

Verlag GmbH & Co.

Traulsen, A. & Nowak, M. A. 2006. Evolution of cooperation by multilevel selection. *Proc.*

Natl. Acad. Sci. USA, 103**:**10952-10955.

Travisano, M. & Velicer, G. J. 2004. Strategies of microbial cheater control. *Trends Microbiol.,*

12**:**72-78.

Turner, P. E. & Chao, L. 1999. Prisoner's dilemma in an RNA virus. *Nature,* 398**:**441-443.

Van Valen, L. 1973. A new evolutionary law. *Evolutionary theory,* 1**:**1-30.

Velicer, G. J., Kroos, L. & Lenski, R. E. 2000. Developmental cheating in the social bacterium

Myxococcus xanthus. *Nature,* 404**:**598-601.

Via, S., Gomulkiewicz, R., De Jong, G., Scheiner, S. M., Schlichting, C. D. & Van Tienderen, P.

H. 1995. Adaptive phenotypic plasticity: Consensus and controversy. *Trends Ecol. Evol.,*

10**:**212-7.

Wang, X., Fu, A. Q., Mcnerney, M. E. & White, K. P. 2014. Widespread genetic epistasis among

cancer genes. *Nature communications,* 5.

Weinreich, D. M., Lan, Y., Wylie, C. S. & Heckendorn, R. B. 2013. Should evolutionary

geneticists worry about high order epistasis? *Current Opinion in Development and*

Genetics, 23**:**700-707.

Williams, G. C. 1966. *Adaptation and natural selection: A critique of some current evolutionary*

thought, Princeton university press.

Wylie, C. S., Ghim, C. M., Kessler, D. A. & Levine, H. 2009. The fixation probability of rare

mutators in finite asexual populations. *Genetics,* 181**:**1595-1602.

Sidebar 1 – What is a lineage? (Typeset near 'Introduction')

We define a lineage as the full genealogy of descendent copies of an allele starting from the

original copy and ending at its long-term fate: extinction, fixation, or maintenance as a stable

polymorphism in the population. A traditional approach in population genetics has been to

describe the long-term evolutionary fate of a mutant allele influencing some biological trait

under the combined influence of evolutionary processes like mutation, genetic drift, migration,

and natural selection. Using analytical approaches from stochastic process theory, this work

seeks to calculate the probability that such an allele ultimately reaches a frequency of one, or

achieves fixation, in the population. This approach places emphasis not solely on the individual

reproductive process but also on the long-term fate of a genetic lineage carrying the mutation. It

therefore captures a much larger class of phenomena where fitness may not be directly affected

among individual carriers of an allele but the allele instead influences the statistical properties of

a lineage (Wylie et al., 2009). Our focus will be primarily on lineage of asexual haploid lineages,

which are easier to analyze and depict. However, the approach and definition of a lineage given

here extends naturally to sexual diploid organisms.

Sidebar 2 – Geometric mean fitness (Typeset near beginning of 'Lineage selection')

A widely appreciated result regarding adaptations to varying environments is the principle that

natural selection will favor traits based on their geometric mean fitness. When reproductive

success changes between generations, natural selection favors traits that increase the long-term

geometric mean fitness (GMF). Reflecting the multiplicative nature of reproduction, GMF is the

product of fitness in each generation, raised to the reciprocal of the number of generations.

Algebraically, GMF = $(\prod_{t=1}^{n} w_t)^{1/n}$, where w_t is the Wrightian fitness of a trait in generation t.

The same quantity can be expressed as a linear average over the natural log of this fitness value,

GMF = $\exp(\frac{1}{n}\sum_{t=1}^{n} \log(w_t))$. In practice, approximations are used such as GMF ≈ $\mu - \sigma^2/\mu$,

where μ is the arithmetic mean fitness and σ^2 is the variance in fitness. This formula explicates

the fact that natural selection favors increases in mean fitness, but also decreases in the variance

of fitness. This implies that natural selection can be risk averse, favoring alleles with lower

variance in fitness even at the expense of decreasing fitness on average.

Sidebar 3 – Evolutionary game theory (Typeset near end of 'Lineage selection' or beginning of

'Limitations of Fitness averages')

Evolutionary game theory (Maynard Smith, 1982) analyzes an interaction among a set of

competing alleles or "strategies" and summarizes their effect in a matrix representing the fitness

payoff of all pairwise competitions among competitors. Such a framework is most useful in the

context of frequency dependent selection, where the fitness effects of an allele are not easily

summarized by a constant selection coefficient. Such a framework provides a natural way to

determine whether a new allele starting from a single copy will tend to increase in frequency or

"invade" a population that is fixed for an alternative allele. This leads to the concept of an

evolutionarily stable strategy or ESS, which is defined as a strategy or allele that cannot be

invaded by any alternative strategy starting at an initially small frequency. The ability of an allele

to invade, or invasion fitness, is a generalization of the notion of a selection coefficient to the

case of frequency dependent selection (Lehmann et al., 2016). While there are notable exceptions

(Traulsen and Nowak, 2006, Traulsen and Hauert, 2009), game theoretic models are typically

deterministic and describe the tendency for allele frequency change but not the statistical

properties of lineages in finite populations.

Glossary of terms (*To appear adjacent to first use of each term or phrase*)

Lineage selection: Competition between two or more lineages in population.

Offspring distribution: A discrete probability distribution that captures the stochasticity in an

individual organism's reproductive success

Cheater: An organism that produces little or no public good but utilizes those goods produced

by other organisms.

Frequency dependent selection: A model in which the fitness of an allele depends on its

frequency in the population as a consequence of interactions between organisms.

Epistasis: The phenotypic effect of a mutation varies with genetic context.

Modifier loci: Loci responsible for genetic properties of a genome, such as mutation rate,

recombination rate and mutational robustness.

Indirect selection: Selection acting on a modifier locus mediated by its effect on the fitness at

other loci in the genome. Indirect selection models commonly assume that the modifier is

intrinsically selectively neutral, and require that recombination rates are low.

Clonal interference: Competition between mutational independent lineages, each carrying one

or more beneficial mutations. Clonal interference is common in asexual populations in which the

beneficial mutation rate is larger than approximately the reciprocal of the population size.

Genetic drift: stochastic variation in allele frequency as a consequence of stochasticity in

reproduction inherent in finite populations.

Drift barrier: the limit on the efficiency of natural selection imposed by genetic drift.

    Figure Captions

Figure 1. Variability in fitness across a lineage in diverse models. The defining feature of

lineage selection is the presence of variability across a lineage either among contemporary

individuals (vertical axis) or between individuals in time (horizontal axis). Genealogies are

shown for two competing allelic lineages indicated by circles. The focal lineage is shaded yellow

and the wild-type lineage is shaded black. **A.** Lineage carrying a beneficial allele (yellow) rising

to fixation under the classical scenario of lineage invariance. **B.** Lineage carrying an allele that

alternates from beneficial to deleterious in a variable environment. Contemporary individuals

share an identical fitness, and hence an identical selection coefficient, but this quantity changes

between generations. **C.** Evolution of a mutator lineage that experiences increased rates of both

deleterious (red dots) and beneficial (grey background) mutations. Fitness in the lineage varies

both among contemporary individuals and between generations. **D.** A cooperative lineage under

a group selection model. Within-group selective pressures cause the allele to be disfavored over

short timescales. Groups with more cooperative alleles tend to displace other groups over longer

timescales (shown with solid grey lines).

Figure 2. Limitations of fitness averages in finite populations. Evolution in a periodic

environment results in four distinct regimes characterized by the relative timescale of natural

selection ($1/s$) and environmental change (τ). Simulation results of the model described by

Cvijović et al. (2015) are shown in blue and the expected change of an allele with the same

average fitness effect in a constant environment is shown in red. Unless otherwise noted,

simulations are conducted with an average selection coefficient of 0.05 and a population size of

100,000. Selection coefficients are held constant at ± 0.06 within each environment while the

timescale of environmental change is varied (beneficial environmental epochs are shaded grey

while deleterious epochs are unshaded). **A.** When the environment changes fast relative to

changes in allele frequency (small $s\tau$), the average change in allele frequency is well

approximated by a fitness average like geometric mean fitness. **B.** When the environment

changes slower than the time of fixation of an allele (large $s\tau$), mutations tend to arise and fix all

in the same environment, regardless of their average fitness effect. **C.** In infinite populations,

averages like the geometric mean fitness are accurate regardless of the timescale of

environmental change. This is an artifact of the fact that, in the absence of genetic drift, allele

frequencies become arbitrarily close to zero or one but never permanently achieve fixation or

extinction. **D.** Average fitness breaks down when large fluctuations in allele frequency occur on

a similar timescale to environmental change (intermediate $s\tau$). This is due to the amplification of

fluctuations by genetic drift whenever alleles reach very high or very low frequencies (Cvijović

et al. 2015). Note that genetic drift occurring as the frequency of the allele approaches 1 causes it

to respond only modestly to the second deleterious epoch. The allele subsequently achieved

fixation much sooner than would be expected on the basis of its average fitness effect.

1016

1017 **Table 1.** Sources of variability across a lineage and associated adaptations (*Typeset near*

1018 *'Introduction'*)

	Basis of variability in fitness		
	Environmental	**Genetic**	**Social**
Specific examples	Spatial variation	Sex	Kin selection
	Temporal variation	Mutation	Multi-level selection
		Clonal interference	
		Epistasis	
Adaptations	Bet-hedging	Sex/recombination	Cooperation
	Phenotypic plasticity	Mutation rate modifiers	Policing
		Evolvability	

1019

Figure 1

1023   **Figure 2**

1024