Response to nitrogen addition reveals metabolic and ecological strategies of soil

bacteria

Md Sainur Samad[1], Charlotte Johns[2], Karl G. Richards[3], Gary J. Lanigan[3], Cecile A.

M. de Klein[4], Timothy J. Clough[2*], Sergio E. Morales[1*]

[1]*Department of Microbiology and Immunology, Otago School of Medical Sciences,*

University of Otago, Dunedin, New Zealand

[2]*Department of Soil and Physical Sciences, Lincoln University, Lincoln, New Zealand*

[3]*Teagasc, Environmental Research Centre, Johnstown Castle, Wexford, Ireland*

[4]*AgResearch Invermay, Mosgiel, New Zealand*

Running head: Community successions in N treated soils

Key words: denitrification; nitrification; greenhouse gases; N_2O; r /K selection

theory; grasslands; soil

*Corresponding Authors

E-mails:

sergio.morales@otago.ac.nz (S. E. Morales)

Timothy.Clough@lincoln.ac.nz (T. J. Clough)

Summary

The nitrogen (N) cycle represents one of the most well studied systems yet the taxonomic diversity of the organisms that contribute to it is mostly unknown, or linked to poorly characterized microbial groups. While progress has allowed functional groups to be refined, they still rely on *a priori* knowledge of enzymes involved, and the assumption of functional conservation, with little connection to the role the transformation plays for specific organisms. Here, we use soil microcosms to test the impact of N deposition on prokaryotic communities. By combining chemical, genomic and transcriptomic analysis we are able to identify and link changes in community structure to specific organisms catalyzing given chemical reactions. Urea deposition led to a decrease in prokaryotic richness, and a shift in community composition. This was driven by replacement of stable native populations, which utilize energy from N-linked redox reactions for maintenance, with fast responding populations that use this energy for growth. This model can be used to predict response to N disturbances and allows us to identify putative life strategies of different functional, and taxonomic, groups thus providing insights into how they persist in ecosystems by niche differentiation.

Introduction

Modern microbiological techniques have given us unprecedented access to the microbial world (Spiro, 2012; Rinke *et al.*, 2013), yet soil communities remain poorly understood (Delmont *et al.*, 2015). While many studies have focused on the diversity or abundance of key populations (Taylor *et al.*, 2012; Gubry-Rangin *et al.*, 2015; Wei *et al.*, 2015), fewer have looked at the transcriptional profiles over time (Nicol *et al.*, 2008; Morales and Holben, 2013), and even less have done so for multiple groups at the same time (Liu *et al.*, 2010; 2014; Brenzinger *et al.*, 2015). This is particularly true of organisms involved in nitrogen (N) cycling in soils. The complexity of the underlying processes combined with the diversity of microbes contributing to each process provides a large challenge to identifying mechanisms active at any given time (Butterbach-Bahl *et al.*, 2013). Currently we lack enough information to understand basic ecological concepts linked to N cycling *in situ* such as: i) substrate competition at both inter and intra species level, ii) full diversity of both present and active N cycling populations, iii) and the life strategies of these populations which in turn control their responses (both as observed growth or transcriptional changes).

Where progress has been made, groundbreaking findings have followed. The initial discovery of ammonia oxidizing archaea (AOA) and its subsequent rise to major player in the N cycle (Leininger *et al.*, 2006; Hatzenpichler, 2012; Stahl and la Torre, 2012) highlighted the unexpected gaps in knowledge. Later studies have suggested different life strategies for AOA when compared to ammonia oxidizing bacteria (AOB) (Sterngren *et al.*, 2015), but this may be complicated by variance across strains (Bayer *et al.*, 2015). One major unknown is whether observations made in studies, or organisms, from one ecosystem translate to others.

It is well established that individual intermediates in the N cycle can be used

for specific reasons (i.e. ammonia oxidation provides electrons, while denitrification

intermediates accept reducing equivalents), but the purpose of the reactions for any

organism is another major unknown. That is, while some organisms carry out these

processes for electrogenic purposes that can result in growth, others do it in order to

maintain redox homeostasis (Simon and Klotz, 2013). Unfortunately examples where

an organism harbours multiple versions of the same enzyme for completely different

purposes (respiration vs. redox balance) exist (Hartsock and Shapleigh, 2011), and are

likely to limit generalizations.

Despite this, studies focusing on population changes in response to

manipulations have consistently recorded conserved patterns (e.g. growth of AOB but

not AOA (Jia and Conrad, 2009; Di *et al.*, 2009; Pratscher *et al.*, 2011)) suggesting

that responses by specific populations in a given location or ecosystem are

predictable. However, the debate continues on whether niche specialization and

differentiation can be determined based solely on correlations, without analyzing the

wider array of processes that contribute or influence any given N transformation

(Prosser and Nicol, 2012). This is relevant in ecosystems such as agricultural

grassland where an understanding of N cycling is crucial for management of both

productivity and greenhouse gases (Herrero *et al.*, 2016), of which nitrous oxide

(N_2O) is a key player (Reay *et al.*, 2012).

In grazed pastures (i.e. agricultural grasslands) N deposition through ruminant

urine drives the emissions of N_2O (Saggar *et al.*, 2013). In this system a full cascade

of transformations begin with urea and can result in accumulation of any intermediate

depending on conditions, but with a final end product of N_2 or N_2O. While the

chemical transformations have been explored (Hamonts *et al.*, 2013; Baral *et al.*,

2014; de Klein, Shepherd, *et al.*, 2014; de Klein, Luo, *et al.*, 2014), mechanistic

understanding of the populations catalyzing the reactions, and the purpose they serve

for the organisms is less clear. In this study we aimed to identify active biochemical

pathways involved in N loss (through gases) in response to urea (simulated ruminant

urine deposition event) and varying moisture content. Observed phenotypes (chemical

transformations) were linked to changes in genotype (functional potential through

DNA; a proxy for population changes), expression of genotype (RNA profiles), and

total community composition (specific taxonomically defined populations based on

16S). We hypothesized that sequential transformation of nitrogenous intermediates

would be coupled to changes in expression of functional genes catalyzing production

and consumption of intermediates. Alternatively, transformations not linked to

population, or expression changes, would be driven by other (non-biochemical)

pathways. We also hypothesized that changes in transcription, or population size,

could serve to determine life strategies of microbes utilizing each intermediate

(whether they are used for growth vs. maintenance). To test this we mimicked a

ruminant urine-N deposition event using repacked soil cores (1.1 mg m^{-3}) on tension

tables monitored for 63 days. Soils were treated with urea under two different

moisture contents: high (near saturation; -1.0 kPa) and low (field capacity; -10 kPa)

moisture. Simultaneous measurements of soil chemistry, gas kinetics, microbial

community composition (by 16S amplicon sequencing) and functional gene

abundance (for nitrification and denitrification) at DNA (gene) and RNA (transcript)

levels were performed to determine the active populations and pathways.

1.

Results:

Soil pH and N transformation dynamics in response to urea

Soil pH increased from acidic (pH = 5.5) to alkaline reaching a maximum (pH

= 8.5) at day 3 in urea treated soils. Return to baseline pH was modulated by soil

moisture with high moisture (HM; -1.0kPa) soil reaching baseline at day 35 and low

moisture soils (LM; -10kPa) doing so at day 53 (Fig. 1). This shift in pH was linked to

a successive N transformation process initiated with urea hydrolysis and leading to

nitrification and denitrification: urea $\rightarrow NH_4^+ \rightarrow NO_2^- \rightarrow NO_3^- \rightarrow N_2O \rightarrow N_2$ (Fig. 1).

Sequential peak activity was observed for each transformation with the response

modified by moisture. Maximum production (mean µg N g^{-1} soil) for each

transformation was observed at day 3, 21 and 35 respectively for NH_4^+ (HM+N =

1758; LM+N= 1730), NO_2^- (HM+N = 79.2; LM+N= 39.7) and NO_3^- (HM+N = 429.2;

LM+N= 335). Two distinct production peaks were observed for N_2O, with a short

pulse (0 to 5 days) reaching a maximum at day 2 for HM soils (11602.8 µg m^{-2} h^{-1})

and day 3 for LM soils (46.8 µg m^{-2} h^{-1}) (Fig. 1 and Supplementary Fig. S1). A

second, longer duration (10 to ~50 days), N_2O pulse reached a maximum at day 28 for

HM soils (6405.1 µg m^{-2} h^{-1}) and day 30 for LM soils (448.9 µg m^{-2} h^{-1}). The large

N_2O spike (first peak) between days 0 to 5 in the HM+N treatment was about 11.6%

of the total N_2O cumulative flux over 63 days, whereas in the LM+N treatment the 0

to 5 day periods accounted for 22.3% of the total N_2O cumulative flux over 63 days.

Population and transcription dynamics for nitrogen related functional groups

Significant changes (ANOVA, p<0.05) in relative activity (mRNA copy

numbers/16S rRNA gene copy numbers) were observed promptly between day 0 & 3

for all functional groups (except AOA and N-fixers in HM soil) in response to urea

(Fig. 1). However, maximum relative transcription did not match maximum

production peaks for corresponding substrates, or products, for each functional group.

Nitrifiers (ammonia oxidizers) displayed niche differentiation, with time, length and

strength of response differing between bacterial (AOB) and archaeal ammonia

oxidizers (AOA). Relative activity of AOA increased (4.6-fold for LM and 1.6-fold

for HM) under urea treatments at day 3 only, with a subsequent decrease (-19.3-fold

for LM and -7-fold for HM) resulting in lower expression than in untreated soils (Fig.

1). AOB relative activity also increased but was sustained for a much longer period

(3-63 days), with maximum activity (>11-fold change) seen at 21 and 35 days for

HM+N and LM+N respectively (Fig. 1). Denitrifiers (both nitrite and nitrous oxide

reducers) showed similar responses as AOA, with peak activity at day 3 and a rapid

return to baseline, in the case of nitrite reducers decreasing to levels below those

observed in non-urea treated soils (Fig. 1). To account for endogenous sources of N,

N_2 fixers were monitored through the activity of the nitrogenase gene (*nifH*). No

significant changes were observed except for day 3 (LM +N only), with a subsequent

decrease in activity below background. This decrease below background was

observed for all N treated samples.

Changes in the relative contribution to total community composition were

calculated by normalizing functional gene copy numbers to total 16S rRNA gene copy

numbers per sample for each functional group (Fig. 1). The maximum observed

relative abundance of each functional group differed for each group (HM and LM

respectively): AOB, 19 & 12 %; AOA, 8 & 13 %; *nirS*, 6.3 & 2.9 %; *nosZI*, 3.3 & 3.4

%; *nifH*, 4.7 & 4.32 %. Further, large population changes over time were mostly

limited to AOB. Generally, AOB comprised <1 % of the total community, but in

response to urea increased up to 29-fold to make up 19 % (day 21 for HM) and 20-

fold to make up 12 % (day 35 for LM) of the community in urea treated soils. In

contrast, AOA were found at consistently high levels (median=4.2 %) in untreated

soils, but numbers decreased >7-fold in response to urea (~1.3 % at least 63 day).

Similarly, other functional groups (*nosZI*, *nifH*) decreased or remained stable (*nirS*) in

response to urea. Similar patterns for both activity and population changes were

observed when absolute values were analyzed (Supplementary Fig. S2).

N deposition induces both a genotypic and a transcriptional response at the

community level that is modified by soil moisture content

Urea deposition imposed a general negative selective pressure leading to

decreases in OTU level prokaryotic diversity (Shannon, -1.2-fold change) and

richness (-1.5-fold change) at DNA level (Fig. 2a, Supplementary Fig. S3). The same

pattern was observed when active microbes (based on RNA) were analyzed with

decreases in OTU level prokaryotic diversity (Shannon, -1.3-fold change) and

richness (-1.9-fold change). Moisture was found to play a modifier role, with LM

samples consistently resulting in lower diversity and richness when compared to their

HM pairs. Richness and diversity losses were not recovered even after 63 days. In

contrast, samples where no urea was applied remained stable (i.e. constant diversity

and richness).

Urea deposition significantly altered community structure as shown in a non-

metric multidimensional scaling (NMDS) plot using a Bray-Curtis dissimilarity

matrix (Fig. 2b and Supplementary Fig. S4). At both DNA and RNA level community

changes along the first axis corresponded with changes in response to urea treatment,

with the second axis accounting for changes in moisture. A pvclust analysis

(hierarchical clustering with p-values calculated via multiscale bootstrap resampling,

Supplementary Fig. S5) confirmed two major clusters [100% AU (Approximately

Unbiased) and 100% BP (Bootstrap Probability)] formed by urea treated (HM+N and

LM+N samples, excluding day 0), vs. untreated soils (HM-N, LM-N, field samples,

and HM+N & LM+N at Day 0). Temporal variance within each cluster was

confirmed using a Mantel correlogram analysis (Fig. 2c). Urea treated samples had

significant changes in community composition immediately upon treatment (Day 0 to

7), with no return to baseline conditions by the end of the experiment. In contrast,

untreated samples did not change significantly over time (Supplemental Fig. S6)

Changes in community structure were associated with shifts in major

taxonomic lineages (Fig. 3). In general, phylum level changes in abundance and

transcription where correlated to each other (Supplementary Table S2 and Fig. S7,

S8). Urea deposition induced temporal changes in phylum level abundance with

observed maximum fold changes per group (HM & LM at DNA level) being:

Acidobacteria, -4.6 & -3.7; Actinobacteria, 2.4 & 5.3; Bacteroidetes, 4.6 & 2.2;

Candidate Division WS3, -10.5 & -7; Chloroflexi, -2.9 & -2.6; Firmicutes, 10.8 &

16.2; Gemmatimonadetes, 2 & 3.3; Nitrospirae, -3.2 & -2; Planctomycetes, -3.7 & -

2.5; Thaumarchaeota, -5.2 & -3.6; Verrucomicrobia, -2.5 & -2; Alphaproteobacteria,

1.4 & 1.7; Betaproteobacteria, 4 & 2; Deltaproteobacteria, -2.2 & -1.4;

Gammaproteobacteria, 1.5 & 2.6. Normalized transcriptional activity (reads of 16S

rRNA/reads of 16S rDNA) identified the Firmicutes and members within classes of

the Proteobacteria as the most transcriptionally active. While abundant phyla tended

to have high levels of normalized transcription, less abundant organisms like the

Thaumarchaeota, were observed to have high normalized transcriptional activity

especially under background conditions (Supplementary Fig. S7). In contrast, groups

traditionally considered slow growers (e.g. Nitrospirae and Gemmatimonadetes) had

low normalized transcription. It was also noted that while normalized transcription

levels remained stable without urea, N deposition induced changes. These changes in

normalized activity did not always match trends observed at individual DNA or RNA

level (e.g. Firmicutes).

Shifts in N and moisture status trigger OTU response linked to divergent life

strategies

Since Fig. 3 only represents a taxonomic summary of all OTUs (irrespective

of their response to treatments), it does not provide a clear indication of who is

changing and why. To account for this, urea responsive OTUs were identified

independently in RNA and DNA profiles (under each treatment) through a SIMPER

analysis. OTUs accounting for 50% of the variance were analyzed (Fig. 4). Response

patterns for detected OTUs were conserved between RNA and DNA profiles.

However, while some OTUs responded similarly to urea under varying moisture

conditions, marked differences were observed with no detectable pattern based on

taxonomy.

OTUs within the Proteobacteria identified in the SIMPER analysis did not

display a conserved response to urea, however when lower taxonomic levels were

examined patterns emerged. A consistent positive response was seen for OTUs within

the class Betaproteobacteria and the family Hyphomicrobiaceae, amongst others.

Conserved positive responses to urea where also observed at the phylum level for the

Firmicutes, Bacteroidetes, Actinobacteria, Gemmatimonadetes and Planctomycetes,

although the level of response varied across lower taxonomic levels. In contrast, with

only some exceptions, OTUs within the phyla Acidobacteria, Verrucomicrobia,

Nitrospirae, Candidate Division WS3 (also referred to as candidate phylum

Latescibacteria) and the Thaumarchaeota all were negatively impacted by urea

deposition.

To account for response patterns over time, we focused on OTUs that

accounted for 30% of the variance in the SIMPER analysis (36 total), with individual

OTU contributions ranging from 5 to 0.1 percent at the DNA level and 5 to 0.06

percent at the RNA level (Table S3). Temporal patterns were conserved between

DNA and RNA profiles (Supplementary Fig. S9, S10), despite differences in absolute

abundance. Once again, moisture acted as a modulator of response with the extent of

impact dependent on the OTU (Fig. 5 and 6). While most functional groups responded

immediately (at both DNA and RNA level), positively affected OTU responses were

observed along all time points creating a succession of positively selected organisms.

In contrast, negatively affected OTUs all responded within the first 2 time points

indicating an immediate negative selective pressure (Fig. 6). Large variances in

absolute changes were observed, even within similar organisms (e.g. Pedobacter),

with fold changes ranging from -10.5 to 410 across both positively and negatively

affected OTUs. Despite this, OTU response was noted to correspond to taxonomy,

with both the effect (positive or negative) and the extent of response (fold change or

total abundance) in line with predicted ecological growth strategies (r vs. k) predicted

for different taxa. To test this, we predicted rRNA operon copy numbers (rrn) for all

36 OTUs and compared them to the observed maximum abundance, max fold change

in population or observed growth rate per day. We consistently observed a non-linear

response with an asymptote reached at higher copy numbers (Fig. 7). These trends

were consistent independent of which moisture conditions were present at the time of

response. To account for preferential response due to moisture, we selected the

highest response for each organism and saw no clear difference in patterns. To

account for potential biases due to uneven representation, OTUs were grouped into

low (1-2 copies of rrn) or high (>2) copy number organisms (Supplementary Table

S4). While significant changes ($p < 0.05$, Supplementary Fig. S11) were observed in

most instances, exceptions were noted (e.g. growth rate under HM).

Discussion

Functional profiling (identification and quantification of specific functional

genes/transcripts) is normally utilized to link chemical transformations to specific

microbial populations capable of catalyzing reactions. However, functional groups are

comprised of taxonomically diverse species of microbes with different lifestyle

strategies that are unlikely to share a conserved response to an ecosystem disturbance

(Ho *et al.*, 2012). While functional profiling allows us to measure the net response of

a functional group, and could serve as a proxy for determining the importance of the

group in a sample, it does not identify how specific organisms benefit from a

catalyzed transformation. Here we used a controlled microcosm experiment to

measure the response of soil communities to a disturbance in the form of changes in

moisture and nitrogen (urea) deposition. Functional analysis (qPCR) demonstrated a

biological response to urea, but differing responses to moisture depending on group

(Fig. 1). Responses are potentially linked to different life strategies amongst these

groups. Ammonia oxidizers displayed contrasting population and expression profiles,

suggesting niche differentiation driven by time and/or substrate concentration. AOA

responded early, and declined, as new N was made available while AOB responded

later with population swings spanning from near detection limit to most dominant

group. These observations match prior reports showing AOA prefer low N conditions,

while AOB respond vigorously to N deposition (Di *et al.*, 2010; Sterngren *et al.*,

2015). This has been interpreted as evidence for differing lifestyles for AOB and

AOA, with AOA preferring nutrient poor conditions and AOB dominating in rich

ones (Sterngren *et al.*, 2015). However, prior assertions that AOB are solely important

for driving nitrification might be overstated given that transcriptional activity for both

groups is comparable if compared at peak time (Di *et al.*, 2009). This contrasting use

of energy between functionally redundant organisms might explain the low

correlations between processes and the abundance of their respective functional

populations (Rocca *et al.*, 2015). When we examine the response of other functional

groups benefiting from influxes of N, like denitrifiers, we see no significant change in

population sizes suggesting that either energy is being utilized for maintenance or

otherwise for redox balance (Hartsock and Shapleigh, 2011; Y. Li *et al.*, 2012;

Dietrich *et al.*, 2013). In contrast, organism which benefit from low levels of N, like N

fixers, decline in response to exogenous N demonstrating real time selective pressure

in a complex ecosystem. These responses also highlight the temporal nature of these

relationships and how by following niche differentiation high number of functionally

redundant organisms can be maintained (Stempfhuber *et al.*, 2016).

Observations also highlight how lifestyle preferences for organisms are

reflected in their dominance in the ecosystem. Prior work suggests that AOA

dominate in soils with low N inputs, but AOB numbers are higher at times of high N

loading or in ecosystems with consistent N deposition (Gong *et al.*, 2013; Sterngren *et*

al., 2015; Venterea *et al.*, 2015; C. Li *et al.*, 2016). This would suggest that a dynamic

ecosystem with varying nutrient levels would select for a higher diversity of

organisms that maintain ecosystem processes stable over time and space (Wang and

Loreau, 2014). Indeed, our data supports this with alpha diversity (calculated based on

16S amplicon analysis at both DNA and RNA) decreasing in response to urea. This is

inconsistent with plant responses to nutrient deposition in which multiple resources

need to be added to elicit a response (Harpole *et al.*, 2016), although contrasting

results have been observed (Suding *et al.*, 2005; Bai *et al.*, 2010; Song *et al.*, 2011;

2012). For microbes, high site to site variance is reported (De Schrijver *et al.*, 2011;

Leff *et al.*, 2015), but similar negative responses are suggested and could be linked to

increased competition in the absence of natural ecosystem variability. However, links

between microbial and plant response suggest interplay between the response of

macro and microbiota (Zeng *et al.*, 2016). While previous work suggests an important

role for moisture in controlling community composition (Waldrop and Firestone,

2006), we only observed a modifier role in our experiment.

Although broad observations align with ecological theory, precise

identification of responsive organisms is rarely carried out. Here we note that while at

phylum level clear responses (+/- fold change) are observed, variance is seen at the

OTU level suggesting the intra-taxonomic diversity we hypothesized reflected the life

history strategies of the organisms. Attempts to link specific transformations to

organisms failed, potentially due to the succession of functionally redundant

organisms that respond at different time with non-overlapping optima. That is, while

functional gene abundance provides the population size of organisms capable of

carrying out a process, the group may be composed of many OTUs with divergent life

strategies or metabolic potentials that affect when they can respond. This makes

functional gene measurement an average of all OTU subpopulations carrying that

gene. However, community response allows us to identify OTUs responsive to N

deposition, which when analyzed independently, provides insights into metabolic

preferences (i.e. aerobic vs. anaerobic, nitrifier vs. denitrifier) based on time and

response to treatments. Taxonomic groups regularly recognized as native to, or

abundant in, oligotrophic conditions declined in the presence of urea. Most of these

groups are still poorly understood, and included the Acidobacteria, Verrucomicrobia,

Nitrospirae, Candidate Division WS3 (also referred to as candidate phylum

Latescibacteria) and the Thaumarchaeota. These organisms are predicted to be slow

growers with the Thaumarchaeal response confirming the AOA patterns observed at

the functional level. In contrast, positively responding organisms are those generally

associated with groups considered eutrophic or capable of fast response. This

discrepancy based on life history strategies has been proposed and applied to

microbes previously, and suggests that an organisms' ability to grow, utilize carbon,

generate proteins and efficiently transform resources to biomass, amongst others, is

related to its rRNA operon copy number (Klappenbach *et al.*, 2000; Stevenson and

Schmidt, 2004; Dethlefsen and Schmidt, 2007; Roller *et al.*, 2016). When applied to

communities, it is associated with microbial successions in which decreases in copy

numbers are associated with later stages of succession including in soils (Nemergut *et*

al., 2015). For example, two OTUs matching the Verrucomicrobial OTU DA101

where found to be negatively affected by urea, and at least one was found to be highly

abundant under background conditions. DA101 seems to be a common soil (and

grassland) organism identified throughout the world (Felske and Akkermans, 1998;

O'Farrell and Janssen, 1999; Brewer *et al.*, 2016). Based on growth (Sangwan *et al.*,

2005) and genome reconstructions (Brewer *et al.*, 2016), these organisms are

predicted to be slow but efficient growers (k strategists). In contrast, most of the

positively affected organisms seemed to posses higher rrn copy numbers and included

members of the Proteobacteria and Bacteroidetes in line with prior predictions (Fierer

et al., 2007). Statistical analysis supported this interpretation with low copy numbers

(1-2) significantly associated to a negative response to N deposition, while high copy

numbers (>2) were linked to increased capacity for growth, growth rate and maximum

abundance. However, we found a non-linear relationship between increased rrn copy

numbers and growth capacity, best fitted by models reaching an asymptote (like

Michaelis-Menten and logistic curves). These are first order models that suggest that

while a benefit exists where increased copy numbers lead to increased growth rate,

after a certain threshold other variables might limit any benefit. Alternatively, a

decrease in growth rate might be observed with increasing copy numbers once a

tradeoff threshold is passed (Lipson, 2015). However, when rrn copy numbers are

log2 transformed, a significant linear fit was observed as seen in prior studies (Roller

et al., 2016). In our study these predictions are made complicated due to the observed

intra-taxonomic variance that can arise from the lack of accurate knowledge of copy

numbers for many organisms, or from metabolic plasticity at higher taxonomic levels.

In addition, our analysis focused on N responsive organisms only, and with only 38

identified it indicates that most organisms were neither positively nor negatively

affected. This could explain why certain organisms (e.g. Actinobacteria) expected to

be k strategist, based on their ability to produce secondary metabolites (Abdelmohsen

et al., 2015) and compete with other organisms (Barka *et al.*, 2015), showed a positive

response to N deposition. Alternatively, the low number of responsive organisms

could indicate that our false discovery rate corrections were too restrictive.

These findings help us get closer to understanding not just the metabolic

potential of organisms in soils, but the role specific pathways play for an organism. It

also allows us to understand the repercussion of disturbances and management of soils

on below ground biodiversity. The knowledge gained through these type of

observations, and integration of life history strategies into microbial ecology, will get

us one step closer to microbiome management as part of soil care.

Experimental procedure

Sample collection and experimental design

A detailed methodology can be found in (Clough *et. al*., In review). In brief,

soil (8^o15'W, 52^o9'N) was collected from a permanently grazed agricultural grassland

(dairy pasture) in March (early spring) at the Teagasc Moorepark Research Centre,

County Cork, Ireland. Soil was sampled after the turf was removed and a spade was

used to randomly sample the A-horizon (5-20 cm depth, excluding grass layer). To

avoid fresh N loading, fields had not been grazed for over a month. Field moist

samples were immediately shipped to Lincoln University, New Zealand and kept at

4^oC until processed. Prior to use, soil was sieved ($\leq$ 2 mm) to remove any stones,

plant roots or earthworms and packed into stainless steel rings (7.3 cm internal

diameter, 7.4 cm deep) to a depth of 4.1 cm at *in situ* soil bulk density (1.1 Mg m^{-3}

with a gravimetric water content (θg) of 0.24 g water g^{-1} soil). The resulting cores had

a total porosity of 0.58 cm^3 pores cm^{-3} soil and were arranged in a factorial

experiment replicated four times. Soil cores were maintained at two moisture

contents: high (near saturated; -1.0 kPa) and low (field capacity; -10 kPa) moisture

using tension tables (Romano et al., 2002). These moisture contents, -1 and -10 kPa

respectively, corresponded to 53% and 30% volumetric water content, or 91% and

52% water-filled pore space (WFPS). Nitrogen was applied as a urea solution at 2141

kg urea/ha dry soil (equivalent to a single urination event at the higher rate expected

under bovine urine deposition of 1000 kg N ha^{-1}). Four treatments in total were

carried out (replicated four times each for a total of 112 cores analyzed) representing

two levels of urea and two levels of moisture: urea + high moisture (HM +N; Urea _-

1.0kPa), urea + low moisture (LM +N; Urea _-10kPa), no urea + high moisture (HM

–N; No Urea _-1.0kPa) and no urea + low moisture (LM –N; No Urea_-10kPa). All

cores where held at 20^{o}C for a period of 63 days.

Soil pH, and inorganic-N measurements

Soil pH was monitored throughout the experiment using a flat surface pH

electrode (Broadley James Corp., Irvine, California). Inorganic N concentrations

(NH_4^+, NO_2^-, NO_3^-) were determined by destructively sampling batches of soil cores.

Each core was homogenized and a subsample was extracted (10 g dry soil: 100 ml 2M

KCl shaken for 1 hour), filtered (Whatman 42) and analyzed using flow injection

analysis (Blakemore et al., 1987). N_2O flux was determined by placing a soil core into

a 1-L stainless steel tin fitted with a gas-tight lid and rubber septa. The headspace was

sampled after 15 and 30 minutes and analyzed using an automated gas chromatograph

(8610; SRI Instruments, Torrance, CA), linked to an autosampler (Gilson 222XL;

Gilson, Middleton, WI) as previously described (Clough et al., 2006).

Nucleic acids extraction

Samples for RNA and DNA extraction were collected simultaneously with

samples for inorganic N analysis, but only samples at 0, 7, 14, 21, 35, 63 days were

processed for nucleic acids. Each biological replicate was extracted and analyzed

separately. For each extraction 2 g (wet weight) of soil were processed using the

PowerSoil Total RNA Isolation and DNA Elution Accessory Kits (MoBio, Carlsbad,

CA) as per manufacturer's instructions, with slight modifications. Bead beating was

done in a Geno/Grinder 2010 (SPEX SamplePrep, LLC, Metuchen, NJ) using two

rounds of beating (1750 strokes/min) for 15 s with a 1 min pause in between. The

total elution volume for RNA and DNA was 60 µl and 100 µl respectively. RNA was

treated with DNase I (RNase-Free) (New England Biolabs, USA) as per the

manufacturer's protocol. RNA quality was assessed by denaturing gel electrophoresis.

RNA and DNA concentration, purity and humic acid contamination were determined

using a Nanodrop Spectrophotometer, ND-1000 (Thermo Scientific). All extractions

were stored at -80 ^{o}C until downstream analyses.

Reverse transcription (RT)

Triplicate cDNA conversions (technical replicates) were performed for each

RNA extraction using the Maxima H Minus First Strand cDNA Synthesis Kit

(Thermo Scientific) according to manufacturer's protocol. Each 20 µl reaction

contained: 13 µl of RNA (208 ng Total RNA), 1 µl of random hexamers (100 pmol),

1µl of dNTP mix (0.5 mM final conc.) and 5 µl of master mix (4 µl of 5X RT buffer

and 1 µl Maxima H Minus reverse transcriptase). All technical replicates for a sample

were combined and stored at -80^{o}C until further analysis. All further analyses were

performed on the same cDNA pool for each sample.

16S rRNA gene amplicon sequencing

16S rRNA gene amplicon sequencing was performed using primers

515F/806R (V4 region of the 16S gene) and the Earth Microbiome Project conditions

(Version 4_13) (Caporaso *et al.*, 2012). All samples were run simultaneously on a

single Illumina MiSeq run. Sequences were first processed in Qiime (version 1.9.1)

using default parameters (Caporaso *et al.*, 2010). Sequences were clustered into

Operational Taxonomic Units (OTUs) at 97% sequence similarity using the SILVA

(version 119) reference library (Quast *et al.*, 2012) and UCLUST (Edgar, 2010)

following the open-reference Operational Taxonomic Unit (OTU) picking protocol.

Taxonomic identification was done using BLAST against the SILVA database (max-e

value = 0.001) (Altschul *et al.*, 1990). Subsampling and rarefactions (10 times) were

performed to equal read depths of 7,400 per sample, and samples below that threshold

were removed. After rarefaction, all 10 OTU tables were merged and exported for

further processing in R (R Development Core Team, 2008). The 16S amplicon

sequences are available in the NCBI SRA database (accession numbers SRP091980).

Quantification of gene and transcript abundance

Quantitative PCR (qPCR) was performed in 384-well plates using the ViiA7

real-time PCR system (Applied Biosystems, Carlsbad, CA). Absolute quantification

was done using a 10-fold serial dilution (10^8 to 10^1) of known copy numbers of

pGEM-T easy (Promega, Madison, Wisconsin, USA) cloned templates as standards.

For all targets qPCR runs included cloned standards, no template control and no

reverse transcription controls (RNA) run in triplicate. No inhibition or positive

amplification on negative controls was observed for any target. All DNA and cDNA

samples were run in quadruplicates to determine abundance of: prokaryotes (16S

rRNA gene), ammonia oxidizers (archaeal [AOA] & bacterial [AOB] ammonia

monooxygenase gene; *amoA*), denitrifiers (cytochrome cd1-type nitrite reductase

gene; *nirS,* and Clade I nitrous oxide reductase gene; *nosZI*) and nitrogen fixers

(nitrogenase gene; *nifH*).

All reactions were performed in 10 µl volumes containing: 1× Master Mix

(Fast SYBR Green Master Mix, ABI), 0.2-0.6 µM of each primer [0.2 µM for AOA

(Tourna *et al.*, 2008), 0.6 µM for AOB (Avrahami *et al.*, 2003); 0.5 µM for 16S

rRNA (Hartman *et al.*, 2009); *nirS* (Throbäck *et al.*, 2004; Yergeau *et al.*, 2007),

nosZI (Henry *et al.*, 2006) & *nifH* (Rösch and Bothe, 2005)], 2 µl of template [DNA

(1 ng total) or cDNA (80× diluted RT reaction, i.e. total 0.13 ng RNA)] and

autoclaved Milli-Q H_2O to a final volume of 10 µl. Primers and qPCR conditions are

summarized in Table S1. A melt curve analysis (95^{o}C for 15 s, 60^{o}C for 1 min then

increasing 0.05^{o}C/s (data acquisition) until 95^{o}C) was performed to test for specificity

and to confirm no amplification in the negative controls.

Statistical analyses

All statistical analyses were performed in R (R Development Core Team,

2008) using the phyloseq (McMurdie and Holmes, 2013), pvclust (Suzuki and

Shimodaira, 2006), vegan (Oksanen *et al.*) and mpmcorrelogram packages. Detailed

descriptions can be found in supplemental methods.

Growth rate estimation and prediction of rRNA operon (rrn) copy numbers

rrn copy numbers for identified OTUs were predicted using the ribosomal RNA

operon copy number database (rrnDB) (Stoddard *et al.*, 2015). For each OTU,

information from the closest strain available was selected. In instances where a

closely related organism was not available, the mean copy number for the closest

taxonomic group (i.e. genus, class, etc.) was used. Copy numbers where then

compared to the maximum observed abundance and the maximum observed fold

change (calculated based on lowest observed abundance for the same organism in a

preceding time point for OTUs showing growth or succeeding time points for those

decreasing in abundance). An estimated growth rate was calculated for OTUs

showing increases in population size in response to N using the following formula:

$$N_t = N_{0*}e^{rt}$$

where: N_t: The amount at time t; N_0: The amount at time 0; r: exponential growth rate;

514   t: Time passed

Fit model for rrn copy numbers

Both non-linear (Michaelis-Menten) and linear regressions were used to fit rrn

copy numbers and population changes (i.e. maximum abundance and fold-change),

and growth rate (per day). The fit model was performed in R using "drc" and

"ggplot2" packages.

Acknowledgements

This work was funded by the New Zealand Government through the New

Zealand Fund for Global Partnerships in Livestock Emissions Research to support the

objectives of the Livestock Research Group of the Global Research Alliance on

Agricultural Greenhouse Gases (Agreement number: 16084) awarded to SEM and the

University of Otago.

References

Abdelmohsen, U.R., Grkovic, T., Balasubramanian, S., Kamel, M.S., Quinn, R.J., and Hentschel, U. (2015) Elicitation of secondary metabolism in actinomycetes. *Biotechnology Advances* **33**: 798–811.

Altschul, S.F., Gish, W., Miller, W., Myers, E.W., and Lipman, D.J. (1990) Basic local alignment search tool. *J Mol Biol* **215**: 403–410.

Avrahami, S., Liesack, W., and Conrad, R. (2003) Effects of temperature and fertilizer on activity and community structure of soil ammonia oxidizers. *Environ Microbiol* **5**: 691–705.

Bai, Y., Wu, J., Clark, C.M., Naeem, S., and Pan, Q. (2010) Tradeoffs and thresholds in the effects of nitrogen addition on biodiversity and ecosystem functioning: evidence from inner Mongolia Grasslands. *Glob Change Biol* **16**: 358–372.

Baral, K.R., Thomsen, A.G., Olesen, J.E., and Petersen, S.O. (2014) Controls of nitrous oxide emission after simulated cattle urine deposition. *Agr Ecosyst Environ* **188**: 103–110.

Barka, E.A., Vatsa, P., Sanchez, L., Gaveau-Vaillant, N., Jacquard, C., Klenk, H.P., et al. (2015) Taxonomy, Physiology, and Natural Products of Actinobacteria. *Microbiol Mol Biol R* **80**: 1–43.

Bayer, B., Vojvoda, J., Offre, P., Alves, R.J.E., Elisabeth, N.H., Garcia, J.A., et al. (2015) Physiological and genomic characterization of two novel marine thaumarchaeal strains indicates niche differentiation. **10**: 1051–1063.

Brenzinger, K., Dörsch, P., and Braker, G. (2015) pH-driven shifts in overall and transcriptionally active denitrifiers control gaseous product stoichiometry in growth experiments with extracted bacteria from soil. *Front Microbiol* **6**: 1226.

Brewer, T., Handley, K., Carini, P., Gibert, J., and Fierer, N. (2016) Genome reduction in an abundant and ubiquitous soil bacterial lineage.

Butterbach-Bahl, K., Baggs, E.M., Dannenmann, M., Kiese, R., and Zechmeister-Boltenstern, S. (2013) Nitrous oxide emissions from soils: how well do we understand the processes and their controls? *Phil Trans R Soc B* **368**: 20130122–20130122.

Caporaso, J.G., Kuczynski, J., Stombaugh, J., Bittinger, K., Bushman, F.D., Costello, E.K., et al. (2010) QIIME allows analysis of high-throughput community sequencing data. *Nat Meth* **7**: 335–336.

Caporaso, J.G., Lauber, C.L., Walters, W.A., Berg-Lyons, D., Huntley, J., Fierer, N., et al. (2012) Ultra-high-throughput microbial community analysis on the Illumina HiSeq and MiSeq platforms. *ISME J* **6**: 1621–1624.

Clough, T.J., Lanigan, G.J., de Klein, C.A.M., Samad, M.S., Morales, S.E., Rax, D., et al. (n.d.). Influence of soil moisture on co-denitrification fluxes from a urea-affected pasture soil. In Review.

de Klein, C.A., Shepherd, M.A., and van der Weerden, T.J. (2014) Nitrous oxide emissions from grazed grasslands: interactions between the N cycle and climate change — a New Zealand case study. *Curr Opin Environ Sustain* **9-10**: 131–139.

de Klein, C.A.M., Luo, J., Woodward, K.B., Styles, T., Wise, B., Lindsey, S., and Cox, N. (2014) The effect of nitrogen concentration in synthetic cattle urine on nitrous oxide emissions. *Agr Ecosyst Environ* **188**: 85–92.

De Schrijver, A., De Frenne, P., Ampoorter, E., Van Nevel, L., Demey, A., Wuyts, K., and Verheyen, K. (2011) Cumulative nitrogen input drives species loss in terrestrial ecosystems. *Global Ecology and Biogeography* **20**: 803–816.

Delmont, T.O., Eren, A.M., Maccario, L., Prestat, E., Esen, Ö.C., Pelletier, E., et al.

576   (2015) Reconstructing rare soil microbial genomes using in situ enrichments and
577    metagenomics. *Front Microbiol* **6**: 1–15.
578 Dethlefsen, L. and Schmidt, T.M. (2007) Performance of the translational apparatus
579    varies with the ecological strategies of bacteria. *J Bacteriol* **189**: 3237–3245.
580 Di, H.J., Cameron, K.C., Shen, J.-P., Winefield, C.S., O'Callaghan, M., Bowatte, S.,
581    and He, J.-Z. (2010) Ammonia-oxidizing bacteria and archaea grow under
582    contrasting soil nitrogen conditions. *Fems Microbiol Ecol* **72**: 386–394.
583 Di, H.J., Cameron, K.C., Shen, J.P., Winefield, C.S., O'Callaghan, M., Bowatte, S.,
584    and He, J.Z. (2009) Nitrification driven by bacteria and not archaea in nitrogen-
585    rich grassland soils. *Nature Geoscience* **2**: 621–624.
586 Dietrich, L.E.P., Okegbe, C., Price-Whelan, A., Sakhtah, H., Hunter, R.C., and
587    Newman, D.K. (2013) Bacterial Community Morphogenesis Is Intimately Linked
588    to the Intracellular Redox State. *J Bacteriol* **195**: 1371–1380.
589 Edgar, R.C. (2010) Search and clustering orders of magnitude faster than BLAST.
590    *Bioinformatics* **26**: 2460–2461.
591 Felske and Akkermans (1998) Prominent occurrence of ribosomes from an uncultured
592    bacterium of the Verrucomicrobiales cluster in grassland soils. *Letters in Applied*
593    *Microbiology* **26**: 219–223.
594 Fierer, N., Bradford, M.A., and Jackson, R.B. (2007) Toward an ecological
595    classification of soil bacteria. *Ecology* **88**: 1354–1364.
596 Gong, P., Zhang, L., Wu, Z., Shang, Z., and Li, D. (2013) Does the nitri cation
597    inhibitor dicyandiamide affect the abundance of ammonia-oxidizing bacteria and
598    archaea in a Hap-Udic Luvisol? *J Soil Sci Plant Nutr* **13**: 35–42.
599 Gubry-Rangin, C., Kratsch, C., Williams, T.A., McHardy, A.C., Embley, T.M.,
600    Prosser, J.I., and Macqueen, D.J. (2015) Coupling of diversification and pH
601    adaptation during the evolution of terrestrial Thaumarchaeota. *Proc Nat Acad Sci*
602    **112**: 9370–9375.
603 Hamonts, K., Balaine, N., Moltchanova, E., Beare, M., Thomas, S., Wakelin, S.A., et
604    al. (2013) Influence of soil bulk density and matric potential on microbial
605    dynamics, inorganic N transformations, N_2O and N_2 fluxes following urea
606    deposition. *Soil Biol Biochem* **65**: 1–11.
607 Harpole, W.S., Sullivan, L.L., Lind, E.M., Firn, J., Adler, P.B., Borer, E.T., et al.
608    (2016) Addition of multiple limiting resources reduces grassland diversity. *Nature*
609    **537**: 93–96.
610 Hartman, A.L., Lough, D.M., Barupal, D.K., Fiehn, O., Fishbein, T., Zasloff, M., and
611    Eisen, J.A. (2009) Human gut microbiome adopts an alternative state following
612    small bowel transplantation. *Proc Nat Acad Sci* **106**: 17187–17192.
613 Hartsock, A. and Shapleigh, J.P. (2011) Physiological Roles for Two Periplasmic
614    Nitrate Reductases in Rhodobacter sphaeroides 2.4.3 (ATCC 17025). *J Bacteriol*
615    **193**: 6483–6489.
616 Hatzenpichler, R. (2012) Diversity, physiology, and niche differentiation of ammonia-
617    oxidizing archaea. *Appl Environ Microb* **78**: 7501–7510.
618 Henry, S., Bru, D., Stres, B., Hallet, S., and Philippot, L. (2006) Quantitative
619    detection of the *nosZ* gene, encoding nitrous oxide reductase, and comparison of
620    the abundances of 16S rRNA, *narG*, *nirK*, and *nosZ* genes in soils. *Appl Environ*
621    *Microb* **72**: 5181–5189.
622 Herrero, M., Henderson, B., Havlík, P., Thornton, P.K., Conant, R.T., Smith, P., et al.
623    (2016) Greenhouse gas mitigation potentials in the livestock sector. *Nature*
624    *Climate Change* **6**: 452–461.
625 Ho, A., Kerckhof, F.-M., Luke, C., Reim, A., Krause, S., Boon, N., and Bodelier,

P.L.E. (2012) Conceptualizing functional traits and ecological characteristics of
methane-oxidizing bacteria as life strategies. *Environmental Microbiology*
Reports **5**: 335–345.
Jia, Z. and Conrad, R. (2009) Bacteriarather than Archaea dominate microbial
ammonia oxidation in an agricultural soil. *Environ Microbiol* **11**: 1658–1671.
Klappenbach, J.A., Dunbar, J.M., and Schmidt, T.M. (2000) rRNA Operon Copy
Number Reflects Ecological Strategies of Bacteria. *Appl Environ Microb* **66**:
1328–1333.
Leff, J.W., Jones, S.E., Prober, S.M., Barberán, A., Borer, E.T., Firn, J.L., et al.
(2015) Consistent responses of soil microbial communities to elevated nutrient
inputs in grasslands across the globe. *Proc Nat Acad Sci* **112**: 10967–10972.
Leininger, S., Urich, T., Schloter, M., Schwark, L., Qi, J., Nicol, G.W., et al. (2006)
Archaea predominate among ammonia-oxidizing prokaryotes in soils. *Nature*
442: 806–809.
Li, C., Di, H.J., Cameron, K.C., Podolyan, A., and Zhu, B. (2016) Effect of different
land use and land use change on ammonia oxidiser abundance and N_2O
emissions. *Soil Biol Biochem* **96**: 169–175.
Li, Y., Katzmann, E., Borg, S., and Schuler, D. (2012) The Periplasmic Nitrate
Reductase Nap Is Required for Anaerobic Growth and Involved in Redox Control
of Magnetite Biomineralization in Magnetospirillum gryphiswaldense. *J Bacteriol*
194: 4847–4856.
Lipson, D.A. (2015) The complex relationship between microbial growth rate and
yield and its implications for ecosystem processes. *Front Microbiol* **6**: 1–5.
Liu, B., Liu, B., Frostegard, Å., Frostegård, Å., Bakken, L.R., and Bakken, L.R.
(2014) Impaired Reduction of N_2O to N_2 in Acid Soils Is Due to a
Posttranscriptional Interference with the Expression of nosZ. *mBio* **5**: e01383–14–
e01383–14.
Liu, B., Mørkved, P.T., Frostegård, Å., and Bakken, L.R. (2010) Denitrification gene
pools, transcription and kinetics of NO, N_2O and N_2 production as affected by soil
pH. *Fems Microbiol Ecol* **72**: 407–417.
McMurdie, P.J. and Holmes, S. (2013) phyloseq: an R package for reproducible
interactive analysis and graphics of microbiome census data. *PLoS ONE*.
Morales, S.E. and Holben, W.E. (2013) Functional response of a near-surface soil
microbial community to a simulated underground CO_2 storage leak. *PLoS ONE* **8**:
e81742–10.
Nemergut, D.R., Knelman, J.E., Ferrenberg, S., Bilinski, T., Melbourne, B., Jiang, L.,
et al. (2015) Decreases in average bacterial community rRNA operon copy
number during succession. **10**: 1147–1156.
Nicol, G.W., Leininger, S., Schleper, C., and Prosser, J.I. (2008) The influence of soil
pH on the diversity, abundance and transcriptional activity of ammonia oxidizing
archaea and bacteria. *Environ Microbiol* **10**: 2966–2978.
Oksanen, J., Blanchet, F.G., Kindt, R., and Legendre, P. Vegan: Community Ecology
Package (2013) R package version 2.0-7. https://cran.r-
project.org/package=vegan.
O'Farrell, K.A. and Janssen, P.H. (1999) Detection of Verrucomicrobia in a Pasture
Soil by PCR-Mediated Amplification of 16S rRNA Genes. *Appl Environ Microb*
65: 4280–4284.
Pratscher, J., Dumont, M.G., and Conrad, R. (2011) Ammonia oxidation coupled to
CO_2 fixation by archaea and bacteria in an agricultural soil. pp. 4170–4175.
Prosser, J.I. and Nicol, G.W. (2012) Archaeal and bacterial ammonia-oxidisers in soil:

the quest for niche specialisation and differentiation. *Trends in Microbiology* **20**:
523–531.
Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., et al. (2012)
The SILVA ribosomal RNA gene database project: improved data processing and
web-based tools. *Nucleic Acid Res* **41**: D590–D596.
R Development Core Team (2008) R: A language and environment for statistical
computing. *R Foundation for Statistical Computing, Vienna, Austria*.
Rinke, C., Schwientek, P., Sczyrba, A., Ivanova, N.N., Anderson, I.J., Cheng, J.F., et
al. (2013) Insights into the phylogeny and coding potential of microbial dark
matter. *Nature* **499**: 431–437.
Rocca, J.D., Hall, E.K., Lennon, J.T., Evans, S.E., Waldrop, M.P., Cotner, J.B., et al.
(2015) Relationships between protein-encoding gene abundance and
corresponding process are commonly assumed yet rarely observed. **9**: 1693–1699.
Roller, B.R.K., Stoddard, S.F., and Schmidt, T.M. (2016) Exploiting rRNA operon
copy number to investigate bacterial reproductive strategies. *Nature Microbiology*
1–7.
Rösch, C. and Bothe, H. (2005) Improved assessment of denitrifying, N_2-fixing, and
total-community bacteria by terminal restriction fragment length polymorphism
analysis using multiple restriction enzymes. *Appl Environ Microb* **71**: 2026–2035.
Saggar, S., Jha, N., Deslippe, J., Bolan, N.S., Luo, J., Giltrap, D.L., et al. (2013)
Denitrification and $N_2O:N_2$ production in temperate grasslands: Processes,
measurements, modelling and mitigating negative impacts. *Sci Total Environ* **465**:
173–195.
Sangwan, P., Kovac, S., Davis, K.E.R., Sait, M., and Janssen, P.H. (2005) Detection
and cultivation of soil verrucomicrobia. *Appl Environ Microb* **71**: 8402–8410.
Simon, J. and Klotz, M.G. (2013) Diversity and evolution of bioenergetic systems
involved in microbial nitrogen compound transformations. *Biochimica et
Biophysica Acta (BBA)-Bioenergetics* **1827**: 114–135.
Song, L., Bao, X., Liu, X., and Zhang, F. (2012) Impact of nitrogen addition on plant
community in a semi-arid temperate steppe in China. *Journal of Arid Land* **4**: 3–
10.
Song, L., Bao, X., Liu, X., Zhang, Y., Christie, P., Fangmeier, A., and Zhang, F.
(2011) Nitrogen enrichment enhances the dominance of grasses over forbs in a
temperate steppe ecosystem. *Biogeosciences* **8**: 2341–2350.
Spiro, S. (2012) Nitrous oxide production and consumption: regulation of gene
expression by gas-sensitive transcription factors. *Phil Trans R Soc B* **367**: 1213–
1225.
Stahl, D.A. and la Torre, de, J.R. (2012) Physiology and diversity of ammonia-
oxidizing archaea. *Annu. Rev. Microbiol.* **66**: 83–101.
Stempfhuber, B., Richter-Heitmann, T., Regan, K.M., Kölbl, A., Wüst, P.K., Marhan,
S., et al. (2016) Spatial Interaction of Archaeal Ammonia-Oxidizers and Nitrite-
Oxidizing Bacteria in an Unfertilized Grassland Soil. *Front Microbiol* **6**: 861–15.
Sterngren, A.E., Hallin, S., and Bengtson, P. (2015) Archaeal ammonia oxidizers
dominate in numbers, but bacteria drive gross nitrification in N-amended
grassland soil. *Front Microbiol* **6**: 1620–8.
Stevenson, B.S. and Schmidt, T.M. (2004) Life History Implications of rRNA Gene
Copy Number in Escherichia coli. *Appl Environ Microb* **70**: 6670–6677.
Stoddard, S.F., Smith, B.J., Hein, R., Roller, B.R.K., and Schmidt, T.M. (2015)
rrnDB: improved tools for interpreting rRNA gene abundance in bacteria and
archaea and a new foundation for future development. *Nucleic Acid Res* **43**:

D593–D598.
Suding, K.N., Collins, S.L., Gough, L., Clark, C., Cleland, E.E., Gross, K.L., et al.
(2005) Functional- and abundance-based mechanisms explain diversity loss due
to N fertilization. *Proc Nat Acad Sci* **102**: 4387–4392.
Suzuki, R. and Shimodaira, H. (2006) Pvclust: an R package for assessing the
uncertainty in hierarchical clustering. *Bioinformatics* **22**: 1540–1542.
Taylor, A.E., Zeglin, L.H., Wanzek, T.A., Myrold, D.D., and Bottomley, P.J. (2012)
Dynamics of ammonia-oxidizing archaea and bacteria populations and
contributions to soil nitrification potentials. **6**: 2024–2032.
Throbäck, I.N., Enwall, K., Jarvis, Å., and Hallin, S. (2004) Reassessing PCR primers
targeting *nirS*, *nirK* and *nosZ* genes for community surveys of denitrifying
bacteria with DGGE. *Fems Microbiol Ecol* **49**: 401–417.
Tourna, M., Freitag, T.E., Nicol, G.W., and Prosser, J.I. (2008) Growth, activity and
temperature responses of ammonia-oxidizing archaea and bacteria in soil
microcosms. *Environ Microbiol* **10**: 1357–1364.
Venterea, R.T., Clough, T.J., Coulter, J.A., Breuillin-Sessoms, F., Wang, P.,
Sadowsky, M.J., and A, R.P.L. (2015) Ammonium sorption and ammonia
inhibition of nitrite-oxidizing bacteria explain contrasting soil N_2O production.
Sci. Rep. **5**: 12153–15.
Waldrop, M.P. and Firestone, M.K. (2006) Response of microbial community
composition and function to soil climate change. *Microb Ecol* **52**: 716–724.
Wei, W., Isobe, K., Nishizawa, T., Zhu, L., Shiratori, Y., Ohte, N., et al. (2015)
Higher diversity and abundance of denitrifying microorganisms in environments
than considered previously. 1–12.
Yergeau, E., Kang, S., He, Z., Zhou, J., and Kowalchuk, G.A. (2007) Functional
microarray analysis of nitrogen and carbon cycling genes across an Antarctic
latitudinal transect. *ISME J* **1**: 163–179.
Zeng, J., Liu, X., Song, L., Lin, X., Zhang, H., Shen, C., and Chu, H. (2016) Nitrogen
fertilization directly affects soil bacterial diversity and indirectly affects bacterial
community composition. *Soil Biol Biochem* **92**: 41–49.
756

757
Figures

Fig. 1 Chemical transformations and biological (functional group) response in soils

treated with urea (+/- 1000 μg N/g dry soil) under two moisture conditions (LM = low

moisture [-10kPa]; HM = high moisture [-1.0kPa]). Error bars are the standard error

of the mean (n ≥ 3, except gene abundance data of day 7 [n=1; LM soil] and day 21

[n=1; LM soil]) for replicate mesocosms. Gene and transcript abundance were

measured by qPCR targeting: nitrifiers (AOA, ammonia oxidizing archaea; AOB,

ammonia oxidizing bacteria), denitrifiers (*nirS,* cytochrome cd $_1$-containing nitrite

reductase*; nosZI*, nitrous oxide reductase) and nitrogen fixers (*nifH*, nitrogenase

reductase). All qPCR results are normalized to 16S rRNA copy numbers and

presented as percent of the nucleic acid pool.

Fig. 2 Total microbial community response (based on 16S rRNA gene amplicon

profiling and clustering of sequences at OTU level (97% sequence similarity)) to urea

(+/-1000 μg N/g dry soil) under two moisture conditions (LM = low moisture [-

10kPa]; HM = high moisture [-1.0kPa]) at both DNA and RNA level. Error bars are

the standard error of the mean (n = 3, except day 7 [n=1; LM soil] and day 21 [n=1;

LM soil]) for replicate mesocosms. (a) Changes in microbial diversity (Shannon)

index over time in response to treatment. (b) Non-metric multidimensional scaling

(NMDS) ordination plots based on Bray-Curtis distances showing relationships

among samples based on OTU level changes in community composition. (c) Mantel

correlogram showing autocorrelation on community composition by performing

sequential Mantel tests between the Bray-Curtis dissimilarities and the grouping of

samples using a time period index (index 1 represents 0-7 days; 2 represents 7-14; 3

represents 14-21; 4 represents 21-35; 5 represents 35-63). Filled circles represent

 significant correlation (p < 0.05) in community composition at specific time periods,

788 with open circles indicating no significant correlation.

Fig. 3 Phylum and class level (for Proteobacteria only) changes in abundance (DNA)

representing relative contribution >1% of all detected phyla (based on OTUs clustered

at 97% sequence similarity). A total of 7,400 sequences were examined per sample.

Error bars are the standard error of the mean (n = 3, except day 7 [n=1; LM soil] and

day 21 [n=1; LM soil]) for replicate mesocosms. Treatments = +/- N [+/-1000 µg N/g

dry soil] under two moisture conditions (LM = low moisture [-10kPa]; HM = high

moisture [-1.0kPa]). Abbreviations: c: Class; p: Phylum. See supplemental Fig. S8 for

relative abundance.

Fig. 4 Taxonomic summary of OTUs responsive to urea treatment identified through similarity percentages (SIMPER) analysis (representing top 50% cumulative sum). The 4 outer rings represent fold changes in response to urea under high and low moisture content (MH & LM respectively) at either DNA or RNA level, with blank gaps indicating OTUs not identified in SIMPER analysis under the specified ring condition. Nodes on the tree (moving outwards from center) correspond to taxonomic level [Domain, Phylum, Class, Order, Family, Genus and Species/OTUs]. Nodes are colored based on dominant response (>50% conserved fold change response across OTUs within a node) with black notes indicating equal representation of positive and

 negatively responding OTUs. Shaded areas of branches delineate defined taxonomic

 groups. See Supplementary file (Table S3) for full classification.

Fig. 5 Population (16S rDNA) changes (absolute abundance based on 7400 reads per samples) for OTUs identified as positively responsive to urea treatment based on similarity percentages (SIMPER) analysis (representing top 30% cumulative sum).

818   Treatments = +/- N [+/-1000 μg N/g dry soil] under two moisture conditions (LM =

819   low moisture [-10kPa]; HM = high moisture [-1.0kPa]).

825   **Fig. 6** Population (16S rDNA) changes (absolute abundance based on 7,400 reads per

826   samples) for OTUs identified as negatively responsive to urea treatment based on

827   similarity percentages (SIMPER) analysis (representing top 30% cumulative sum).

828   Treatments = +/- N [+/-1000 μg N/g dry soil] under two moisture conditions (LM =

829   low moisture [-10kPa]; HM = high moisture [-1.0kPa]).

Fig. 7 Relationship between predicted ribosomal RNA operon (rrn) copy numbers and

growth rate (per day), maximum observed population change, or fold change in

response to N treatment under both high moisture (HM) content, low moisture (LM)

content and best growth either in HM or in LM (based on maximum observed

growth). Copy number was estimated using rrn database (Stoddard *et al.*, 2015). Copy

number values were obtained by finding the closest match (lowest taxonomic level

possible) to each OTU and retrieving the mean rRNA copy number for that group.

852