Parsing propagule pressure: Simulated and experimental disentanglement of introduction

size and number of introductions for colonizing individuals

Michael J Koontz [a,b,c,1], Meagan F Oldfather [d,2], Brett A Melbourne[e,3], Ruth A Hufbauer[b,c,4]

a. Graduate Group in Ecology, University of California, Davis, CA 95616

b. Graduate Degree Program in Ecology, Colorado State University, Fort Collins, CO 80523

c. Department of Bioagricultural Science and Pest Management, Colorado State University, Fort

Collins, CO 80523

d. Department of Integrative Biology, University of California, Berkeley, CA 94720

e. Department of Ecology and Evolutionary Biology, University of Colorado, Boulder, CO

80309

Running header: Parsing propagule pressure

[1] Email: mkoontz@ucdavis.edu
[2] Email: meagan_oldfather@berkeley.edu
[3] Email: brett.melbourne@colorado.edu
[4] Email: hufbauer@colostate.edu

ABSTRACT

Colonization of novel habitats is more likely with increasing propagule pressure— the total

number of individuals introduced to a site. Two key components of propagule pressure are the

number of introduction events and the size of those introduction events. It is unclear which of

these components is more important for colonization success, or under what environmental

conditions their relative importance might shift. Using demographic simulations paired with a

Tribolium flour beetle microcosm experiment, we introduced 20 total individuals into replicated

novel habitats of stable or fluctuating quality and manipulated the number and size of

introduction events used to distribute them through time. After seven generations, we evaluated

how different combinations of the number and size of introductions affected establishment

probability, size of established populations, and whether effects depended on the variability of

the recipient environment. We found no effect of biologically realistic environmental

stochasticity on establishment probability or size of established populations in the demographic

simulations. However, there was strong evidence that establishment probability was enhanced

with more, smaller introductions. In the microcosm, we similarly found no effect of

environmental stochasticity on establishment probability, but unlike the simulations found that

populations that established were larger in the stable environment, especially with more

introduction events. The microcosm experiment yielded greater overall establishment probability

and larger populations compared to the demographic simulations. Genetic mechanisms likely

underlie these differences in results and thus deserve more attention in efforts to parse propagule

pressure. Our results highlight the importance of preventing further introductions of undesirable

species to invaded sites, and suggest conservation efforts should focus on increasing the number

of introductions or re-introductions of desirable species rather than increasing the size of those

introduction events.

KEYWORDS: invasion, reintroduction, propagule pressure, biocontrol, conservation,

microcosm, population dynamics, stochasticity, simulation

INTRODUCTION

Colonization is the ecologically fundamental process of population establishment in an

unoccupied site, and it underlies the past, present, and future distributions of species.

Colonization occurs naturally, but is increasingly prevalent due to anthropogenic influences

(Sakai et al. 2001, Cassey et al. 2005, Ricciardi 2007). Incipient populations often face

environments that are entirely novel, which is especially likely in the case of anthropogenic

colonization (Cassey et al. 2005, Ricciardi 2007). Regardless of whether colonization events to

novel habitats are natural (e.g., range expansion) or human-mediated (e.g., biological invasions,

reintroductions of rare species, release of biological control agents), their successes or failures

have significant implications for natural resource managers and society (Mack et al. 2000).

Most introductions to novel habitats fail, and colonization success can be difficult to predict

(Lockwood et al. 2005, Zenni and Nuñez 2013). Incipient populations are commonly small, and

face threats from environmental, demographic, and genetic stochasticity (Lande 1988, 1993,

Fauvergue et al. 2012). Furthermore, the success of any given population can be idiosyncratic

with respect to taxonomy and geography (Lodge 1993, Lockwood et al. 2005). Thus, it is crucial

to understand more general features of the colonization process beyond the particular invading

organism or the particular invaded environment (Lockwood et al. 2005). Propagule pressure is

one such general feature that is the only known consistent predictor of colonization success in

novel habitats (Lockwood et al. 2005, Colautti et al. 2006, Simberloff 2009).

Propagule pressure is the total number of potentially reproductive individuals (e.g*.,* adults, eggs,

seeds, vegetative material) introduced to an area (Novak 2007). It is often described in this broad

sense, which belies its complexity. Two important components of propagule pressure are the

number of introduction events— sometimes termed propagule number, and the number of

individuals per introduction event— sometimes termed propagule size (*sensu* Fauvergue et al.

2012). Here, we use "introduction regime" to refer to different combinations of the number of

introduction events and the number of individuals introduced per event. The same total

propagule pressure, N, can be achieved by different introduction regimes depending on how

those N individuals are distributed in time or space (Haccou and Iwasa 1996). An introduction

regime of N individuals lies on a continuum bounded by maximizing the number of individuals

introduced per event (all N individuals introduced in one event to the same location) and

maximizing the number of introduction events (one individual introduced in each of N sequential

events through time or to each of N unique locations).

Colonization success increases with total propagule pressure, but it is unclear whether the

correlation is driven by the number of individuals introduced per event or the number of

introduction events (Lockwood et al. 2005, Colautti et al. 2006, Simberloff 2009). Historical data

suggest that multiple introductions can facilitate population establishment, but conclusions from

these studies are limited by the inability to control for the total number of individuals introduced

(Hopper and Roush 1993, Blackburn and Duncan 2001, Grevstad et al. 2011, Fauvergue et al.

2012). Models that hold the total propagule pressure constant agree that multiple, small

introductions distributed across space will lead to greater establishment probability compared to

a single, large introduction when Allee effects are weak (Haccou and Iwasa 1996, Grevstad 1999,

Schreiber and Lloyd-Smith 2009). However, both modeling and empirical approaches have

generated conflicting views on how an introduction regime affects colonization success when

introductions are distributed through time and total propagule pressure is constant.

There is evidence from both models and experiments that colonization success can increase with

more, smaller introduction events through time. Branching process models show that, in the long

run, several, small introductions will always be more likely to successfully establish a population

than a single, large introduction (Haccou and Iwasa 1996, Haccou and Vatutin 2003). This

finding was corroborated by simulations with no Allee effects (Drolet and Locke 2016). In a

Daphnia microcosm experiment, increasing introduction frequency (proportional to the number

of introductions) positively affected population growth, but the number of individuals per

introduction had no detectable effect. This pattern did not hold for establishment probability,

which was only affected by total propagule pressure (Drake et al. 2005). However, Drake et al.

(2005) did not continue scheduled introductions if a population went extinct, denying those

populations one of the main benefits of repeated introductions and creating variability in the total

propagule pressure. In a more recent experiment, several, small introductions led to a 65%

increase in abundance of successfully colonizing invasive Pacific Oyster (*Crassostrea gigas*)

compared to a single, large introduction (Hedge et al. 2012).

There is also evidence from both models and experiments that colonization success increases

with fewer, larger introduction events through time. In simulations of bird invasions, a single,

large introduction event always led to the greatest establishment probability (Cassey et al. 2014).

When Drolet and Locke (2016) included Allee effects in their simulations, their previously

observed pattern reversed and colonization success was instead more likely with fewer, larger

introductions than with more, smaller introductions. In a field experiment with the psyllid

biocontrol agent, *Arytainilla spartiophila*, the number of individuals per introduction event was a

better predictor of establishment success than the number of introduction events (Memmott et al.

2005). In this case however, the introduction regimes with the most individuals per event also

had the highest total propagule pressure (Memmott et al. 2005). In an experiment that controlled

total propagule pressure, a single, large introduction of the non-native mysid, *Hemimysis*

anomala, led to larger populations and greater survival probabilities compared to several, small

introductions through time (Sinclair and Arnott 2015).

Environmental stochasticity in the recipient environment may also affect which introduction

regime is optimal for colonization. Branching process models show that a more variable

environment can reduce the probability of population establishment for all introduction regimes

(Haccou and Iwasa 1996, Haccou and Vatutin 2003), and simulations of introductions distributed

in space suggest that greater environmental variability will magnify the benefit of multiple

introductions (Grevstad 1999). However, simulations of introductions through time did not

support either of these outcomes— a single, large introduction was most likely to establish a

population even with extreme levels of environmental stochasticity (Cassey et al. 2014).

Thus, modeling and empirical approaches have not resolved how different introduction regimes

with a fixed total propagule pressure will affect colonization success when introductions are

distributed through time. Further, there has been no experimental test of whether variability in

the recipient environment interacts with the introduction regime to affect colonization. We paired

demographic simulations with a laboratory microcosm experiment using the red flour beetle,

Tribolium castaneum, to reconcile conflicts in the literature and test how different introduction

regimes implemented through time affected colonization success in novel habitats. We explicitly

manipulated whether the novel habitat was stable or randomly fluctuating in quality to assess

how the success of different introduction regimes may depend upon variability in the recipient

environment. With the total number of individuals introduced held constant at 20, we varied the

size and number of introduction events used to distribute those individuals in four different

introduction regimes. We evaluated establishment probability and population size over 7 discrete

generations to ask: 1) does colonization success increase with more introduction events or with

more individuals per introduction event?, and 2) does the effect of the introduction regime on

colonization success depend on whether the recipient novel environment is stable or fluctuating

through time?

METHODS

General Framework

In simulations and a microcosm experiment, we evaluated the outcome of introducing 20 total

individuals to one of two environmental contexts (a stable or fluctuating novel environment),

varying the number of introduction events used to distribute those individuals through time. This

total was low enough to allow some population extinction within a reasonable timeframe, but

high enough to be representative of documented introductions in the literature (Simberloff 1989,

Simberloff 2009; Grevstad 1999; Berggren 2001; Taylor et al. 2005; Drake et al. 2005). The

introduction regimes were: 20 individuals introduced in the first generation, 10 individuals

introduced in each of the first 2 generations, 5 individuals introduced in each of the first 4

generations, and 4 individuals introduced in each of the first 5 generations. To create the

fluctuating environment, we imposed a magnitude of variability corresponding to environmental

stochasticity in nature, which leads to frequent, mild to moderate perturbations in population

growth rate due to external forces (Lande 1993). Populations were tracked for seven generations

following the initial introduction.

Establishment probability and the size of established populations were used as measures of

colonization success. Populations were deemed 'established' for any time step in which they

were extant and population size was noted for all extant populations in every time step. For

analysis, establishment probability and mean population size were assessed in generation 7.

Study System

Our simulations and microcosm experiment use *Tribolium castaneum* (red flour beetles) to

model the life history of organisms with discrete, non-overlapping generations (e.g., annual

insects and fishes) following Melbourne and Hastings (2008). Individual simulated colonists

were randomly sourced from a randomly mating, infinitely large population. Individual

experimental colonists came from a thoroughly mixed source population maintained at 800

individuals in each of the 4 temporal blocks over which the experiment was replicated. To obtain

colonists, beetles from the source population were mixed freely on a plate, selected from many

sections of the plate, and introduced to subsets of the experiment populations in a random order

(about 24 subsets per block). All migrant females were assumed to arrive mated from the source

population. The strong maternal effects exhibited by *Tribolium* flour beetles were reduced by

rearing individuals from the source population on novel growth medium for 1 generation prior to

using them as colonists (Van Allen and Rudolf 2013, Hufbauer et al. 2015).

Simulations

Population dynamics were simulated with a Negative Binomial-Binomial gamma (NBBg) Ricker

model (Melbourne and Hastings 2008) with an additional mating function of the form:

$$N_{t+1,i} \sim NegBinom\left(\mu_i, k_D F_{mated_{(t,i)}}\right)$$

$$\mu_i = \frac{F_{mated_{(t,i)}}}{p} R_{E_i} e^{-\alpha N_{t,i}}$$

$$R_{E_i} \sim Gamma\left(k_E, \frac{R_0}{k_E}\right)$$

$$N_{t,i} = N_{migrants_{(t,i)}} + N_{residents_{(t,i)}}$$

$$F_{mated_{(t,i)}} = \begin{cases} F_{migrants_{(t,i)}}, & F_{migrants_{(t,i)}} + F_{residents_{(t,i)}} = N_{t,i} \\ F_{migrants_{(t,i)}} + F_{residents_{(t,i)}}, & F_{migrants_{(t,i)}} + F_{residents_{(t,i)}} < N_{t,i} \end{cases}$$

$$F_{migrants_{(t,i)}} \sim Binom\left(N_{migrants_{(t,i)}}, p\right)$$

$$F_{residents_{(t,i)}} \sim Binom(N_{residents_{(t,i)}}, p)$$

$$p = 0.5$$

$$R_0 \sim Gamma(2.6, 1)$$

$$\alpha \sim Gamma(0.0037, 1)$$

$$k_E \sim Gamma(17.6, 1)$$

$$k_D \sim Gamma(1.07, 1)$$

(Eq. 1)

Where $N_{t+1,i}$ is the size of population i at time $t+1$, k_D represents stochasticity due to

demographic heterogeneity, μ_i is the expectation of the size of population i in the next time step,

$F_{mated(t,i)}$ is the latent number of mated females in population i at time t, p is the probability of an

individual being female, R_{Ei} is the latent density-independent population growth rate for

environment E, α is the egg cannibalism rate, $N_{t,i}$ is the total size of population i at time t, k_E

represents environmental stochasticity, R_0 is the density-independent population growth rate for

the average environment, $N_{migrants(t,i)}$ is the total number of migrants to population i at time t, and

$N_{residents(t,i)}$ is the total number of residents in population i at time t.

We imposed a mating function such that a population would deterministically go extinct if it

comprised only non-migrant females. In cases with an all female population and a mixture of

residents and migrants, we only included the number of migrant females in the density-

dependent effect of the demographic heterogeneity term $k_D*F_{mated(t,i)}$ because this effect

manifests via the number of eggs laid by females and only migrant females, being pre-mated,

would lay eggs. Weakly regularizing gamma priors were taken from Melbourne and Hastings

(2008). The expected equilibrium population size for the model is:

$$K = \frac{\log(R_0)}{\alpha}$$

(Eq. 2)

For each combination of the 4 introduction regimes and 2 recipient environment types (described

above), we simulated 500,000 replicate populations for 7 generations. All simulations were

performed in R (R Core Team 2016). This model captures the stochastic and deterministic

ecological dynamics of the *Tribolium* system well, but does not include evolutionary processes

(Melbourne and Hastings 2008, Hufbauer et al. 2015).

To parameterize the model, we censused 125 *Tribolium* populations one generation after

establishing them at various, known densities (between 5 and 200 individuals) on the novel

growth medium following the rearing procedure described in "Microcosm Experiment" below.

We fit the hierarchical model (Eq. 1) to these data in a Bayesian framework to generate posterior

distributions for each of the parameters using the nimble package in R (de Valpine et al. 2016).

We used a Metropolis-Hastings random walk sampler with 3 chains having 50,000 samples each

(including 10,000 samples that were removed for burn in). The chains converged (multivariate $\hat{R}$

for the 4 key parameters =1.01) and produced a sufficient number of effective samples (R_0: 757,

α: 548, k_E: 4103, k_D: 1703).

A single novel growth medium mixture was used for these 125 populations (mixture 5,

Supplementary Table 1), so we treated the estimated environmental stochasticity parameter, k_E,

as the expectation for a stable environment. We parameterized a fluctuating environment model

by increasing the standard deviation of the density-independent per capita population growth rate

by approximately 10% for any given population at any given time step. This was accomplished

by multiplying all of the samples from the posterior distribution of the environmental

stochasticity parameter, k_E, by 100/121 (derived by moment matching for the gamma

distribution). By using random sets of parameters drawn from the samples of the posterior

distributions, we were able to propagate the model uncertainty through the simulations and

combine it with the deterministic and stochastic dynamics represented by the model itself.

Microcosm Experiment

We founded 842 *Tribolium* populations with different introduction regimes (20 individuals in the

first generation, 10 individuals in the first 2 generations, 5 individuals in the first 4 generations,

or 4 individuals in the first 5 generations) and environments (stable or fluctuating) with between

96 and 120 replicate populations per treatment combination (Supplemental Table 3). Each

population was reared in a 4cm x 4cm x 6cm plastic box (AMAC Plastic Products) with 2

tablespoons (approximately 15 g) of freshly prepared growth medium that had been humidified

for at least 24 hours. The growth medium used for the source population (the natal environment

of the colonists) comprised 95% wheat flour (Pillsbury Co. or Gold Medal Products Co.) and 5%

brewers' yeast (Sensient Flavors). Colonists were introduced into a novel growth medium

comprising a small percentage of natal medium mixed with corn flour (Bob's Red Mill). All

populations were reared in one of two dark incubators at 31° and approximately 70% relative

humidity (standard conditions) and were haphazardly rotated between incubators weekly.

For each population in each generation, a known number of adults laid eggs for 24 hours in fresh

medium, and were then removed. Offspring were given 35 days to develop, and adults were then

censused. Censused adults laid eggs on freshly prepared growth medium for 24 hours,

completing their laboratory life cycle. In censusing trials, we estimated the maximum

observation error to be 4.6% (median: 0%, mean: 0.26%) with no detectable difference across

observers or populations with different sizes.

Each replicate population experienced a novel environment that was either stable or randomly

fluctuating through time. The same novel growth medium mixture containing 99.05% novel corn

flour and 0.95% natal medium (mixture 5, Supplementary Table 1) was used for the stable

environment for the duration of the experiment, which preliminary results indicated would yield

a population growth rate of $\lambda = 1.2$ compared to a mean population growth rate of 3.36 on 100%

natal medium. To create the fluctuating environment, we randomly selected a novel growth

medium mixture from one of 9 possible media mixtures for each population in each generation.

Each population in the fluctuating environment therefore experienced a unique series of

environmental conditions. The 9 possible media mixtures represented a gradient of corn flour to

natal medium ratios, and were designed to yield expected population growth rates between 0.88

and 1.33 (Supplementary Table 1). We chose this range to mimic environmental stochasticity

measured in nature (Sæther and Engen 2002) while remaining within the bounds of biologically

realistic population growth (λ between 0.5 and 1.5) (Morris et al. 2008). The random choice was

constrained such that the expected geometric mean population growth rate for each population

resembled expected growth of populations in the stable environment ($\lambda_{expected} = 1.2 \pm 0.05$).

We estimated the amount of environmental stochasticity that we achieved in the fluctuating

environment as the difference in mean total stochasticity between populations in the fluctuating

and stable environments (Sæther and Engen 2002). We assumed that total stochasticity was a

combination of demographic and environmental stochasticity for populations in the fluctuating

environment, and that demographic stochasticity was the sole contributor to total stochasticity

for populations in the stable environment. Total stochasticity (demographic plus environmental)

of each population that did not experience extinction (n=667) was calculated as the variance of

the natural logarithms of its population growth rates through 7 generations:

$$s_{total} = s_{demographic} + s_{environmental} = var(\log(\lambda_t))$$

(Eq. 3)

where, for a particular population, s_{total} is its total stochasticity, $s_{demographic}$ is its demographic

stochasticity, $s_{environmental}$ is its environmental stochasticity (assumed to be 0 for populations

in the stable environment), and λ_t is its per capita population growth rate between generation *t-1*

and generation *t* (*t=1, 2, ... , 7*). We only calculated total stochasticity for populations that did

not experience any extinction in order to capture the full temporal extent of environmental

fluctuations and because extinctions would have an infinite effect on this measure of

stochasticity.

Statistical analyses

We evaluated how our environment treatment affected variability in population growth rate (total

stochasticity from Eq. 3) using a linear mixed effects model with environment (stable or

fluctuating) as a fixed effect and block as a random intercept effect.

We used a mixed effects logistic regression with a logit link to predict the binary response of

establishment, and a mixed effects Poisson regression with a log link to analyze population size.

In both models, introduction regime, environment treatment, and their interaction were treated as

fixed effects, and block was treated as a random intercept effect.

We assessed the effect of temporary extinctions on the establishment probability and mean

population size by fitting the generalized linear mixed effects models described above to data

from the multiple introduction regimes (i.e. not the 20x1 regime) and with additional predictor

variables. To assess the effect of the presence of a temporary extinction, we included an

additional Boolean predictor for whether a population went temporarily extinct or not. To assess

the role of the total propagule pressure, we included a numeric predictor representing the number

of beetles introduced after the latest temporary extinction.

Group-level significance of fixed effects was tested using likelihood ratio tests on nested models,

and least-squares contrasts were used to compare levels of the fixed effects. All statistical

analyses were performed in R, version 3.3.2 (R Core Team 2016). Generalized linear mixed

models were fit using the lme4 package, version 1.1-12 (Bates et al. 2015) and pairwise

comparisons were made using the lsmeans package, version 2.25 (Lenth 2016).

Data availability

The raw experiment data, simulated population trajectories, R code for the simulations, R code

fitting the NBBg model, samples from the posterior distributions of the NBBg parameters, and R

code fitting the mixed effects models are available as supplemental material, on Figshare

(https://doi.org/10.6084/m9.figshare.4648865.v1), and on GitHub

(https://github.com/mikoontz/ppp-establishment).

Note on egg contamination

Laboratory procedures after generation 3 resulted in occasional egg contamination between

replicate populations of the same treatment. Some populations of 1 individual persisted when

they should have deterministically gone extinct, so we manually edited those population

trajectories such that they went extinct in the generation after having only 1 individual.

RESULTS

Simulations

Summary statistics for the posterior distributions of the four NBBg model parameters are given

in Supplementary Table 2.

Our simulations showed introduction regimes with more introduction events were more likely

establish a population by the seventh generation (Figure 1). Introductions into a stochastically

fluctuating environment resulted in slightly lower population establishment for all introduction

regimes (difference of ~1%), but did not favor any particular regime (not shown). The mean

population sizes for each introduction regime/environment combination were approximately

equal in simulations. Mean population sizes only incorporated extant populations, so they were

slightly larger than the expectation for the equilibrium population size (Figure 2).

Microcosm experiment

Mean environmental stochasticity of populations in the fluctuating environment was 0.052 (95%

CI = 0.0073 to 0.0966; p = 0.023). This value is near the median value of 0.055 measured in

nature by Sæther and Engen (2002) in a metaanalysis of 35 avian populations, indicating that we

achieved biologically realistic fluctuations in population growth rate.

We found no evidence that the probability of establishment was affected by a main effect of

environment (χ^2=0.72, df=1, p=0.40), nor by an interaction between environment and

introduction regime (χ^2=3.49, df=3, p=0.32). However, there was strong support for an effect of

introduction regime on establishment probability (χ2=59.76, df=3, p<0.0001). Pairwise

comparisons of the different introduction regimes averaged across the environment treatments

revealed that the 4x5 regime was the most likely to establish populations, with a probability of

about 0.98, whereas the 20x1 and 10x2 regimes were the least likely to establish populations,

with a probability reduced to about 0.8 (Figure 1).

The mean size of populations that persisted until generation 7 was shaped by significant effects

of introduction regime (χ^2=91.65, df=3, p<0.0001), environment treatment (χ^2=117.83, df=3,

p<0.0001), and their interaction (χ^2=44.62, df=3, p<0.0001). Populations established via more

introduction events were larger when averaged across the environment treatments. Extant

populations in the stable environment were larger than those in the fluctuating environment when

averaged across the introduction regimes. The interaction manifests as the benefit of a stable

environment increases with more, smaller introduction events (Figure 2).

Extinctions accumulated regularly throughout the experiment period, with 101 out of 842

populations (12.0%) going extinct by generation 7 (Supplementary Figure 2). The additional

introductions that some populations received often restored a population that had temporarily

gone extinct. Out of 602 populations that received more than 1 introduction (i.e. not the 20x1

introduction regime), 104 of them (17.3%) temporarily went extinct at least once before being

replenished by additional colonizing individuals. Twelve populations were rescued in this way at

least twice, and one population was rescued in this way three times.

Temporary extinctions significantly affected colonization success. The presence of a temporary

extinction significantly reduced average establishment probability from 92.4% to 82.1%

(difference = -1.13 on logit scale, 95% CI = -0.22 to -2.04, χ^2 = 5.44, df = 1, p = 0.020) and mean

population size from 47.8 to 45.4 (difference = -0.052 on log scale, 95% CI = -0.02 to -0.09, χ^2 =

9.42, df = 1, p=0.0021). Each additional colonist contributing to a population after the latest

temporary extinction significantly increased the mean population size (estimate = 0.005 on the

log scale, 95% CI = 6.4e-05 to 0.01, χ^2 = 3.94, df = 1, p=0.047).

Results from the simulations are expected to represent the dynamics in the microcosm

experiment if the assumption of the model on which the simulations were built holds—that only

demographic processes are responsible for the dynamics observed in the experiment. However,

establishment probability at generation 7 was equal to or greater than that expected from the

simulations (Figure 1) and mean population sizes were much larger in the experiment than in the

simulations (Figure 2), together suggesting an influence of non-demographic mechanisms.

DISCUSSION

We assessed how the number and size of introduction events through time drive colonization

success in a novel environment when the total number of individuals introduced to a site is fixed.

We considered novel environments that were either stable or randomly fluctuating in quality

through time, and evaluated populations through 7 discrete generations. We approached this

question in two ways: 1) stochastic simulations of a demographic population dynamics model

parameterized with empirical data, and 2) a highly replicated laboratory microcosm experiment.

By coupling these approaches, we were able to expand and test the theoretical understanding of

how the introduction regime affects colonization in stable and fluctuating environments as well

as to develop new avenues for research when observations did not align with predictions. We

found that several, small introductions increase colonization success and that demographic

processes alone are insufficient to explain the dynamics observed in the experiment.

We found minimal to no effect of a biologically realistic level of environmental stochasticity on

establishment probability in demographic simulations. This finding is inconsistent with the

prediction made by Grevstad (1999) who found with simulations that several, small introductions

would produce an especially high establishment probability compared to a single, large

introduction in a variable environment. The difference in findings is perhaps due to a difference

in the kinds of introduction regimes modeled: Grevstad (1999) simulated multiple introductions

distributed in space, while we focused on introductions separated in time. Additionally, we may

not have detected the same effect of environmental stochasticity that Grevstad (1999) noted

because of differences in the magnitude of environmental variation imposed. At the highest

levels of simulated environmental variation imposed in Grevstad's study, over 65% of

environments would result in population decline and over 40% of environments would result in

population loss of greater than 90%. This kind of environmental variability might be

characterized as random catastrophes—major events causing immediately disastrous effects to

the population, rather than a level of environmental stochasticity leading to more moderate

fluctuations in growth rates (Lande 1993). Our simulations and microcosm results do corroborate

the results from simulations by Cassey et al. (2014) who also used introductions through time

and found a minimal effect of environmental stochasticity on establishment probability. Cassey

et al. (2014) further found a minimal effect of random, infrequent catastrophes and bonanzas,

suggesting that increasing environmental variability through time in the broadest sense (i.e.,

encompassing environmental stochasticity, random catastrophes, and bonanzas) does not

magnify the benefit of several, small introductions through time in the same way that Grevstad

(1999) found with several, small introductions across space.

There also did not appear to be a strong effect of environmental stochasticity on population size

in the demographic simulations. All treatments in the simulations reached similar mean

population size by generation 7, which was slightly higher than the equilibrium size due to our

measure only including extant populations. This suggests that strong density dependence in the

demographic simulations was the primary influence on population size.

We observed striking differences in the measures of colonization success between the

demographic simulations and the microcosm experiment. We found that establishment

probability increased with the number of introduction events (as in the simulations), but that all

microcosm establishment probabilities equaled or exceeded expectations from simulations. In the

microcosm, mean population size in stable environments was greater than in fluctuating

environments. There was also an interaction between environment and introduction regime,

whereby the mean population size was increasingly greater in stable compared to fluctuating

environments as the number of introductions increased. Furthermore, populations grew larger by

generation 7 in the microcosm experiment than in the simulations.

Departures from the simulation expectations in the microcosm experiment suggest that the

demographic processes captured by the model do not account for all of the biological processes

that occurred in the microcosm. Recent work shows that adaptation to the novel, harsh

environment from standing variation is possible in this species (Hufbauer et al. 2015), and likely

explains the greater establishment probability and population sizes in the microcosm compared

to expectations derived from demographic simulations which do not include adaptation.

The rescue effect of immigration can act demographically by increasing the size of populations

(Brown and Kodric-Brown 1977). Certainly, demographic rescue played a critical role for the

104 populations that went extinct temporarily until another introduction event revived them.

Those temporary extinctions had lasting effects on colonization success. Colonization success

declined for populations that experienced a temporary extinction, and the mean population size

significantly increased if more colonists contributed to the population after a temporary

extinction. These results reflect the overarching importance of total propagule pressure

regardless of introduction regime.

The rescue effect can also act genetically by increasing the fitness of populations (Brown and

Kodric-Brown 1977, Frankham 2015, Whiteley et al. 2015). The immigrants all came from the

same large external source population, so it is unlikely that gene flow from migrants produced

novel genetic combinations on which selection could act (Novak 2007). Thus, a more likely

mechanism by which immigration increased mean population size was by relieving inbreeding

depression or counteracting drift-induced allele loss. Small populations are more prone to

experiencing increased homozygosity and inbreeding depression, which can reduce population

growth rates and increase extinction risk (McCauley and Wade 1981, Saccheri et al. 1998,

O'Grady et al. 2006, Szűcs et al. 2014, 2017). Even small amounts of gene flow can alleviate

these effects, so the additional small introductions of mated individuals from the external source

population were well-suited to bring about longer-term relief (Slatkin 1985, Mills and Allendorf

1996). However, Cassey et al. (2014) found that simulated inbreeding depression was especially

detrimental for several, small introductions through time, so other mechanisms might be at play.

Sustained immigration may affect adaptation of the incipient population. Introductions to a harsh,

novel habitat can result in adaptive evolution with the right amount of gene flow if additional

immigrants to a declining population prevent extinction long enough to allow for adaptation to

occur (Holt and Gomulkiewicz 1997). However, too much gene flow can lead to genetic

swamping whereby the homogenizing effects of gene flow overpowers ongoing local adaptation

(Mills and Allendorf 1996, Lenormand 2002). This may have been the case for populations in

the 10x2 introduction regime, which had the highest average migration rate, lowest establishment

probability, and lowest mean population size. Alternatively, negative density dependence may

have reduced population fitness when migration rates were high, reducing population growth

rates and hampering the spread of adaptive alleles (Holt and Gomulkiewicz 1997). Though not

significant, the lower establishment probability and mean population size in the 10x2

introduction regime warrants further work to assess whether they exemplify the yet-unseen

scenario described by Blackburn et al. (2015) in which maladaptive gene flow from multiple

introductions hampers population establishment in a novel range. More broadly, further study is

necessary to evaluate how immigration affected adaptation in this system, if at all

(Gomulkiewicz and Holt 1995, Boulding and Hay 2001).

CONCLUSION

Our experimental results suggest that several, small introductions through time lead to greater

colonization success in a novel habitat, and that introductions into a stable recipient environment

lead to larger population sizes, but not greater establishment probability. Further, introductions to

a stable recipient environment are especially beneficial to populations established with more

introduction events. These results defied our expectations derived from parallel simulations of a

model that included demographic processes but not evolutionary ones, so we suspect a genetic

mechanism might be at work. Genetic mechanisms are rarely incorporated when simulating the

effect of introduction regime on colonization (but see Cassey et al. 2014), and our multi-

generation microcosm is unique in bringing evolutionary processes to bear on parsing two key

components of propagule pressure in an experimental setting.

For invasions, our results highlight the importance of preventing further introductions to the

same location, even for established species. For conservation and biological control, our results

suggest that emphasis should be placed on increasing the number of introductions or re-

introductions to a location, rather than increasing the size of those events if Allee effects are

weak. Sustained introduction efforts should also bring about concomitant benefits in the form of

longer-term monitoring, increased data collection, and more opportunities for experimentation

and adaptive management (Lockwood et al. 2005, Godefroid et al. 2011).

ACKNOWLEDGMENTS

We thank Charlotte Hoover, Kaitlyn Adkins, Mary Catherine Cochran, Don Waters, Stacy

Endriss, Marianna Szűcs, and Jenny Rickford for help with data collection. Funding was

provided by NSF Graduate Research Fellowship Grants #DGE-1321845 Amend. 3 (to MJK) and

#DGE-1106400 (to MFO), NSF Grant DEB-0949619 (to RAH), and NSF Grant DEB-0949595

(to BAM). RAH also acknowledges the support of the USDA via the Colorado Experiment

Station. We thank the lab groups of Ruth Hufbauer, Paul Ode, Andrew Norton, Andrew Latimer,

Truman Young, and Jenny Gremer for comments on the project and manuscript.

REFERENCES

Van Allen, B. G., and V. H. W. Rudolf. 2013. Ghosts of habitats past: Environmental carry-over

effects drive population dynamics in novel habitat. The American Naturalist 181:596–608.

Bates, D., M. Maechler, B. Bolker, and S. Walker. 2015. Fitting linear mixed-effects models

using lme4. Journal Of Statistical Software 67:1–48.

Berggren, A. 2001. Colonization success in Roesel's bush-cricket *Metrioptera roeseli*: The

effects of propagule size. Ecology 82:274–280.

Blackburn, T. M., and R. P. Duncan. 2001. Establishment patterns of exotic birds are constrained

by non-random patterns in introduction. Journal of Biogeography 28:927–939.

Blackburn, T. M., J. L. Lockwood, and P. Cassey. 2015. The influence of numbers on invasion

success. Molecular Ecology:1–12.

Boulding, E. G., and T. Hay. 2001. Genetic and demographic parameters determining population

persistence after a discrete change in the environment. Heredity 86:313–24.

Brown, J. H., and A. Kodric-Brown. 1977. Turnover rates in insular biogeography: Effect of

immigration on extinction. Ecology 58:445–449.

Cassey, P., T. M. Blackburn, R. P. Duncan, and S. L. Chown. 2005. Concerning invasive

species: Reply to Brown and Sax. Austral Ecology 30:475–480.

Cassey, P., T. A. A. Prowse, and T. M. Blackburn. 2014. A population model for predicting the

successful establishment of introduced bird species. Oecologia 175:417–428.

Colautti, R. I., I. A. Grigorovich, and H. J. MacIsaac. 2006. Propagule pressure: A null model for

biological invasions. Biological Invasions 8:1023–1037.

Drake, J. M., P. Baggenstos, and D. M. Lodge. 2005. Propagule pressure and persistence in

experimental populations. Biology Letters 1:480–3.

Drolet, D., and A. Locke. 2016. Relative importance of propagule size and propagule number for

establishment of non-indigenous species: A stochastic simulation study. Aquatic Invasions

11.

Fauvergue, X., E. Vercken, T. Malausa, and R. A. Hufbauer. 2012. The biology of small,

introduced populations, with special reference to biological control. Evolutionary

Applications 5:424–43.

Frankham, R. 2015. Genetic rescue of small inbred populations: Meta-analysis reveals large and

consistent benefits of gene flow. Molecular Ecology 24:2610–2618.

Godefroid, S., C. Piazza, G. Rossi, S. Buord, A.-D. Stevens, R. Aguraiuja, C. Cowell, C. W.

Weekley, G. Vogg, J. M. Iriondo, I. Johnson, B. Dixon, D. Gordon, S. Magnanon, B.

Valentin, K. Bjureke, R. Koopman, M. Vicens, M. Virevaire, and T. Vanderborght. 2011.

How successful are plant species reintroductions? Biological Conservation 144:672–682.

Gomulkiewicz, R., and R. D. Holt. 1995. When does evolution by natural selection prevent

extinction? Evolution 49:201–207.

Grevstad, F. S. 1999. Factors influencing the chance of population establishment: Implications

for release strategies in biocontrol. Ecological Applications 9:1439–1447.

Grevstad, F. S., E. M. Coombs, and P. B. McEvoy. 2011. Revisiting release strategies in

biological control of weeds: Are we using enough releases? XIII International Symposium

on Biological Control of Weeds:368–376.

Haccou, P., and Y. Iwasa. 1996. Establishment probability in fluctuating environments: A

branching process model. Theoretical Population Biology 50:254–280.

Haccou, P., and V. Vatutin. 2003. Establishment success and extinction risk in autocorrelated

environments. Theoretical Population Biology 64:303–314.

Hedge, L. H., W. A. O'Connor, and E. L. Johnston. 2012. Manipulating the intrinsic parameters

of propagule pressure: Implications for bio-invasion. Ecosphere 3:art48.

Holt, R. D., and R. Gomulkiewicz. 1997. How does immigration influence local adaptation? A

reexamination of a familiar paradigm. The American Naturalist 149:563–572.

Hopper, K. R., and R. T. Roush. 1993. Mate finding, dispersal, number released, and the success

of biological control introductions. Ecological Entomology 18:321–331.

Hufbauer, R. A., M. Szűcs, E. Kasyon, C. Youngberg, M. J. Koontz, C. Richards, T. Tuff, and B.

A. Melbourne. 2015. Three types of rescue can avert extinction in a changing environment.

Proceedings of the National Academy of Sciences 112:10557–10562.

Lande, R. 1988. Genetics and demography in biological conservation. Science 241:1455–1460.

Lande, R. 1993. Risks of population extinction from demographic and environmental

stochasticity and random catastrophes. The American Naturalist 142:911–927.

Lenormand, T. 2002. Gene flow and the limits to natural selection. Trends in Ecology and

Evolution 17:183–189.

Lenth, R. V. 2016. Least-squares means: The R package lsmeans. Journal Of Statistical Software

69:1–33.

Lockwood, J. L., P. Cassey, and T. Blackburn. 2005. The role of propagule pressure in

explaining species invasions. Trends in Ecology and Evolution 20:223–8.

Lodge, D. M. 1993. Biological invasions: Lessons for ecology. Trends in Ecology and Evolution

8:133–137.

Mack, R. N., D. Simberloff, W. M. Lonsdale, H. Evans, M. Clout, and F. A. Bazzaz. 2000.

Biotic invasions: Causes, epidemiology, global consequences, and control. Ecological

Applications 10:689–710.

McCauley, D. E., and M. J. Wade. 1981. The populational effects of inbreeding in *Tribolium*.

Heredity 46:59–67.

Melbourne, B. A., and A. Hastings. 2008. Extinction risk depends strongly on factors

contributing to stochasticity. Nature 454:100–3.

Memmott, J., P. G. Craze, H. M. Harman, P. Syrett, and S. V Fowler. 2005. The effect of

propagule size on the invasion of an alien insect. Journal of Animal Ecology 74:50–62.

Mills, L. S., and F. W. Allendorf. 1996. The one-migrant-per-generation rule in conservation and

management. Conservation Biology 10:1509–1518.

Morris, W. F., C. A. Pfister, S. Tuljapurkar, C. V Haridas, C. L. Boggs, M. S. Boyce, E. M.

Bruna, D. R. Church, T. Coulson, D. F. Doak, S. Forsyth, J.-M. Gaillard, C. C. Horvitz, S.

Kalisz, B. E. Kendall, T. M. Knight, C. T. Lee, and E. S. Menges. 2008. Longevity can

buffer plant and animal populations against changing climatic variability. Ecology 89:19–25.

Novak, S. J. 2007. The role of evolution in the invasion process. Proceedings of the National

Academy of Sciences 104:3671–2.

O'Grady, J. J., B. W. Brook, D. H. Reed, J. D. Ballou, D. W. Tonkyn, and R. Frankham. 2006.

Realistic levels of inbreeding depression strongly affect extinction risk in wild populations.

Biological Conservation 133:42–51.

R Core Team. 2016. R: A language and environment for statistical computing. http://www.r-

project.org/. Vienna, Austria.

Ricciardi, A. 2007. Are modern biological invasions an unprecedented form of global change?

Conservation Biology 21:329–36.

Saccheri, I., M. Kuussaari, M. Kankare, P. Vikman, and I. Hanski. 1998. Inbreeding and

extinction in a butterfly metapopulation. Nature 392:491–494.

Sæther, B.-E., and S. Engen. 2002. Pattern of variation in avian population growth rates.

Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences

357:1185–95.

Sakai, A. K., F. W. Allendorf, J. S. Holt, D. M. Lodge, J. Molofsky, K. A. With, S. Baughman, R.

J. Cabin, J. E. Cohen, N. C. Ellstrand, D. E. McCauley, P. O. Neil, I. M. Parker, J. N.

Thompson, and S. G. Weller. 2001. The population biology of invasive species. Annual

Review of Ecology and Systematics 32:305–332.

Schreiber, S. J., and J. O. Lloyd-Smith. 2009. Invasion dynamics in spatially heterogeneous

environments. The American Naturalist 174:490–505.

Simberloff, D. 1989. Which insect introductions succeed and which fail? Pages 61–75 *in* J. A.

Drake et al., editor. Biological Invasions: A Global Perspective. John Wiley & Sons Ltd.

Simberloff, D. 2009. The role of propagule pressure in biological invasions. Annual Review of

Ecology, Evolution, and Systematics 40:81–102.

Sinclair, J. S., and S. E. Arnott. 2015. Strength in size not numbers: Propagule size more

important than number in sexually reproducing populations. Biological Invasions.

Slatkin, M. 1985. Gene flow in natural populations. Annual Review of Ecology and Systematics

16:393–430.

Szűcs, M., B. A. Melbourne, T. Tuff, and R. A. Hufbauer. 2014. The roles of demography and

genetics in the early stages of colonization. Proceedings of the Royal Society B Biological

Sciences 281:1–7.

Szűcs, M., B. A. Melbourne, T. Tuff, C. Weiss-Lehman, and R. A. Hufbauer. 2017. Genetic and

demographic founder effects have long-term fitness consequences for colonising

populations. Ecology Letters.

Taylor, S. S., I. G. Jamieson, and D. P. Armstrong. 2005. Successful island reintroductions of

New Zealand robins and saddlebacks with small numbers of founders. Animal Conservation

8:415–420.

de Valpine, P., C. Paciorek, D. Turek, C. Anderson-Bergman, and D. Temple Lang. 2016.

nimble: Flexible BUGS-compatible system for hierarchical statistical modeling and

algorithm development. R package version 0.6-2.

Whiteley, A. R., S. W. Fitzpatrick, W. C. Funk, and D. A. Tallmon. 2015. Genetic rescue to the

rescue. Trends in Ecology and Evolution 30:42–49.

Zenni, R. D., and M. A. Nuñez. 2013. The elephant in the room: The role of failed invasions in

understanding invasion biology. Oikos 122:801–815.

Figure 1. Establishment probabilities for each introduction regime assessed at generation 7.

Triangles represent results from simulations. Dot-whiskers represent estimates and 95%

confidence intervals from the mixed effects logistic regression model fit to data from the

microcosm experiment. Different letters over dot-whiskers represent introduction regimes with

significantly different establishment probabilities.

Figure 2. Mean sizes of extant populations for each introduction regime/environment

combination in generation 7. Triangles represent results from simulations. Dot-whiskers

represent estimates and 95% confidence intervals from the mixed effects Poisson regression

model fit to data from the microcosm experiment. Different letters over dot-whiskers represent

introduction regimes with significantly different mean population sizes. The dashed line

represents the theoretical equilibrium population size derived from the NBBg model, which

includes extinction and is therefore lower than the mean population size from the simulations.