Parsing propagule pressure: Number, not size, of introductions drives colonization

success in a novel environment

Michael J Koontz [a,b,c,1], Meagan F Oldfather [d,2], Brett A Melbourne[e,3], Ruth A Hufbauer[b,c,4]

a. Graduate Group in Ecology, University of California, Davis, CA 95616

b. Graduate Degree Program in Ecology, Colorado State University, Fort Collins, CO 80523

c. Department of Bioagricultural Science and Pest Management, Colorado State University, Fort

Collins, CO 80523

d. Department of Integrative Biology, University of California, Berkeley, CA 94720

e. Department of Ecology and Evolutionary Biology, University of Colorado, Boulder, CO

80309

Running header: Parsing propagule pressure

[1] Email: mkoontz@ucdavis.edu; ORCID: orcid.org/0000-0002-8276-210X
[2] Email: meagan_oldfather@berkeley.edu; ORCID: orcid.org/0000-0003-3256-4786
[3] Email: brett.melbourne@colorado.edu
[4] Email: hufbauer@colostate.edu

ABSTRACT

Predicting whether individuals will colonize a novel habitat is of fundamental ecological interest

and is crucial to both conservation efforts and invasive species management. A consistently

supported predictor of colonization success is the number of individuals introduced, also called

propagule pressure. Propagule pressure increases with the number of introductions and the

number of individuals per introduction (the size of the introduction), but it is unresolved which

process is a stronger driver of colonization success. Furthermore their relative importance may

depend upon the environment, with multiple introductions potentially enhancing colonization of

fluctuating environments. To evaluate the relative importance of the number and size of

introductions and its dependence upon environmental variability, we paired demographic

simulations with a microcosm experiment. Using *Tribolium* flour beetles as a model system, we

introduced a fixed number of individuals into replicated novel habitats of stable or fluctuating

quality, varying the number of introductions through time and size of each introduction. We

evaluated establishment probability and the size of extant populations after 7 generations. In the

simulations and microcosms, we found that establishment probability increased with more,

smaller introductions, but was not affected by biologically realistic fluctuations in environmental

quality. Population size was not significantly affected by environmental variability in the

simulations, but populations in the microcosms grew larger in a stable environment, especially

with more introduction events. In general, the microcosm experiment yielded higher

establishment probability and larger populations than the demographic simulations. We suggest

that genetic mechanisms likely underlie these differences and thus deserve more attention in

efforts to parse propagule pressure. Our results highlight the importance of preventing further

introductions of undesirable species to invaded sites, and suggest conservation efforts should

focus on increasing the number of introductions or re-introductions of desirable species rather

than increasing the size of those introduction events.

KEYWORDS: invasion, reintroduction, propagule pressure, biocontrol, conservation,

microcosm, population dynamics, stochasticity, simulation

INTRODUCTION

Colonization is the ecologically fundamental process of population establishment in an

unoccupied site, and it underlies the past, present, and future distributions of species.

Colonization occurs naturally, but is increasingly prevalent due to anthropogenic influences

(Sakai et al. 2001, Cassey et al. 2005, Ricciardi 2007). Incipient populations often face

environments that are entirely novel, which is especially likely in the case of anthropogenic

colonization (Cassey et al. 2005, Ricciardi 2007). Regardless of whether colonization events to

novel habitats are natural (e.g., range expansion) or human-mediated (e.g., biological invasions,

reintroductions of rare species, release of biological control agents), their successes or failures

have significant implications for natural resource managers and society (Mack et al. 2000).

Most introductions to novel habitats fail, and colonization success can be difficult to predict

(Lockwood et al. 2005, Zenni and Nuñez 2013). Incipient populations are commonly small, and

face threats from environmental, demographic, and genetic stochasticity (Lande 1988, 1993,

Fauvergue et al. 2012). Furthermore, the success of any given population can be idiosyncratic

with respect to taxonomy and geography (Lodge 1993, Lockwood et al. 2005). Thus, it is crucial

to understand more general features of the colonization process beyond the particular invading

organism or the particular invaded environment (Lockwood et al. 2005). Propagule pressure is

one such general feature that is a consistent predictor of colonization success in novel habitats

(Lockwood et al. 2005, Colautti et al. 2006, Simberloff 2009, Jeschke 2014).

Propagule pressure is the total number of potentially reproductive individuals (e.g., adults, eggs,

seeds, vegetative material) introduced to an area (Novak 2007). It is often described in this broad

sense, which belies its complexity. Two important components of propagule pressure are the

number of introduction events— sometimes termed propagule number, and the number of

individuals per introduction event— sometimes termed propagule size (*sensu* Fauvergue et al.

2012). Here, we use "introduction regime" to refer to different combinations of the number of

introduction events and the number of individuals introduced per event. The same total

propagule pressure, N, is realized by different introduction regimes depending on how those N

individuals are distributed in time or space (Haccou and Iwasa 1996). An introduction regime of

N individuals lies on a continuum bounded by maximizing the number of individuals introduced

per event (all N individuals introduced in one event to the same location) and maximizing the

number of introduction events (one individual introduced in each of N sequential events through

time or to each of N unique locations).

Colonization success increases with total propagule pressure, but it is unclear whether the

correlation is driven by the number of individuals introduced per event or the number of

introduction events (Lockwood et al. 2005, Colautti et al. 2006, Simberloff 2009). Historical data

suggest that multiple introductions can facilitate population establishment, but conclusions from

these studies are limited by the inability to control for the total number of individuals introduced

(Hopper and Roush 1993, Blackburn and Duncan 2001, Grevstad et al. 2011, Fauvergue et al.

2012). Models that hold the total propagule pressure constant agree that multiple, small

introductions distributed across space will lead to greater establishment probability compared to

a single, large introduction when Allee effects are weak (Haccou and Iwasa 1996, Grevstad 1999,

Schreiber and Lloyd-Smith 2009). However, both modeling and empirical approaches have

generated conflicting views on how an introduction regime affects colonization success when

introductions are distributed through time, a situation in which individuals from later

introductions interact both demographically and genetically with individuals from previous

introductions.

There is evidence from both models and experiments that colonization success can increase with

more, smaller introduction events through time. Branching process models show that, in the long

run, several small introductions will be more likely to successfully establish a population than a

single large introduction (Haccou and Iwasa 1996, Haccou and Vatutin 2003). This finding was

corroborated by simulations with no Allee effects (Drolet and Locke 2016). In a *Daphnia*

microcosm experiment, increasing introduction frequency (proportional to the number of

introductions) positively affected population growth, but the number of individuals per

introduction had no detectable effect (Drake et al. 2005). Establishment probability was only

affected by total propagule pressure, and did not increase with increasing introduction frequency

(Drake et al. 2005). However, Drake et al. (2005) did not continue scheduled introductions if a

population went extinct, denying those populations one of the main benefits of repeated

introductions and creating variability in the total propagule pressure. In a more recent experiment,

several small introductions through time led to a 65% increase in abundance of successfully

colonizing invasive Pacific Oyster (*Crassostrea gigas*) compared to a single large introduction

(Hedge et al. 2012).

There is contrasting evidence from both models and experiments that colonization success

increases with fewer, larger introduction events through time. In simulations of bird invasions, a

single large introduction event always led to the greatest establishment probability (Cassey et al.

2014). When Drolet and Locke (2016) included Allee effects in their simulations, their

previously observed pattern reversed and colonization success was instead more likely with

fewer, larger introductions than with more, smaller introductions. In a field experiment with the

psyllid biocontrol agent, *Arytainilla spartiophila*, the number of individuals per introduction

event was a better predictor of establishment success than the number of introduction events

(Memmott et al. 2005). In this case however, the introduction regimes with the most individuals

per event also had the highest total propagule pressure (Memmott et al. 2005). In an experiment

that controlled total propagule pressure, a single, large introduction of the non-native mysid,

Hemimysis anomala, led to larger populations and greater survival probabilities compared to

several, small introductions through time (Sinclair and Arnott 2016).

Environmental stochasticity in the recipient environment may also affect which introduction

regime is optimal for colonization. Branching process models show that a more variable

environment reduces the probability of population establishment for all introduction regimes

(Haccou and Iwasa 1996, Haccou and Vatutin 2003), and simulations of introductions distributed

in space suggest that greater environmental variability will magnify the benefit of multiple

introductions (Grevstad 1999). However, simulations of introductions through time by Cassey et

al. (2014) did not support either of these outcomes— a single, large introduction was most likely

to establish a population even with extreme levels of environmental stochasticity.

Thus, modeling and empirical approaches have not resolved how different introduction regimes

with a fixed total propagule pressure will affect colonization success when introductions are

distributed through time. Further, there has been no experimental test of whether variability in

the recipient environment interacts with the introduction regime to affect colonization. We paired

demographic simulations with a laboratory microcosm experiment using the red flour beetle,

Tribolium castaneum, to reconcile conflicts in the literature and test how different introduction

regimes implemented through time affect colonization success in novel habitats. We explicitly

manipulated whether the novel habitat was stable or randomly fluctuating in quality to assess

how the success of different introduction regimes may depend upon variability in the recipient

environment. With the total number of individuals introduced held constant at 20, we varied the

size and number of introduction events used to distribute those individuals in four different

introduction regimes. We evaluated establishment probability and population size over 7 discrete

generations to ask: 1) does colonization success increase with more introduction events or with

more individuals per introduction event?, and 2) does the effect of the introduction regime on

colonization success depend on whether the recipient novel environment is stable or fluctuating

through time?

METHODS

General Framework

In simulations and a microcosm experiment, we evaluated the outcome of introducing 20 total

individuals to one of two environmental contexts (a stable or fluctuating novel environment),

varying the number of introduction events used to distribute those individuals through time. This

total propagule pressure was low enough to allow some population extinction within an

observable timeframe, but high enough to be representative of documented introductions in the

literature (Simberloff 1989, Simberloff 2009; Grevstad 1999; Berggren 2001; Taylor et al. 2005;

Drake et al. 2005). The introduction regimes were: 20 individuals introduced in the first

generation, 10 individuals introduced in each of the first 2 generations, 5 individuals introduced

in each of the first 4 generations, and 4 individuals introduced in each of the first 5 generations.

To create the fluctuating environment, we imposed a magnitude of variability corresponding to

environmental stochasticity in nature, which leads to frequent, mild to moderate perturbations in

population growth rate due to external forces (Lande 1993). Populations were tracked for seven

generations following the initial introduction.

Establishment probability and the size of established populations were used as measures of

colonization success. Populations were deemed 'established' for any time step in which they

were extant and population size was noted for all extant populations in every time step.

Establishment probability and mean population size were assessed in generation 7.

Study System

Our simulations and microcosm experiment use *Tribolium castaneum* (red flour beetles) to

model the life history of organisms with discrete, non-overlapping generations (e.g., annual

insects and fishes) following Melbourne and Hastings (2008). Individual simulated colonists

were randomly sourced from a randomly mating, infinitely large population. Individual

experimental colonists came from a thoroughly mixed source population maintained at 800

individuals in each of the 4 temporal blocks over which the experiment was replicated. To obtain

colonists, beetles from the source population were mixed freely on a plate, selected from many

sections of the plate, and introduced to subsets of the experiment populations in a random order

(about 24 subsets per block). All migrant females were assumed to arrive mated from the source

population. The strong maternal effects exhibited by *Tribolium* flour beetles were reduced by

rearing individuals from the source population on novel growth medium for 1 generation prior to

using them as colonists (Van Allen and Rudolf 2013, Hufbauer et al. 2015).

Simulations

Population dynamics were simulated with a Negative Binomial-Binomial gamma (NBBg) Ricker

model (Melbourne and Hastings 2008) with an additional mating function of the form:

$$N_{t+1,i} \sim NegBinom\left(\mu_i, k_D F_{mated_{(t,i)}}\right)$$

$$\mu_i = \frac{F_{mated_{(t,i)}}}{p} R_{E_i} e^{-\alpha N_{t,i}}$$

$$R_{E_i} \sim Gamma\left(k_E, \frac{R_0}{k_E}\right)$$

$$N_{t,i} = N_{migrants_{(t,i)}} + N_{residents_{(t,i)}}$$

$$F_{mated_{(t,i)}} = \begin{cases} F_{migrants_{(t,i)}}, & F_{migrants_{(t,i)}} + F_{residents_{(t,i)}} = N_{t,i} \\ F_{migrants_{(t,i)}} + F_{residents_{(t,i)}}, & F_{migrants_{(t,i)}} + F_{residents_{(t,i)}} < N_{t,i} \end{cases}$$

$$F_{migrants_{(t,i)}} \sim Binom\left(N_{migrants_{(t,i)}}, p\right)$$

$$F_{residents_{(t,i)}} \sim Binom(N_{residents_{(t,i)}}, p)$$

$$p = 0.5$$

$$R_0 \sim Gamma(2.6, 1)$$

$$\alpha \sim Gamma(0.0037, 1)$$

$$k_E \sim Gamma(17.6, 1)$$

$$k_D \sim Gamma(1.07, 1)$$

(Eq. 1)

Where $N_{t+1,i}$ is the size of population i at time $t+1$, k_D represents stochasticity due to

demographic heterogeneity, μ_i is the expectation of the size of population i in the next time step,

$F_{mated(t,i)}$ is the latent number of mated females in population i at time t, p is the probability of an

individual being female, R_{Ei} is the latent density-independent population growth rate for

environment E, α is the egg cannibalism rate, $N_{t,i}$ is the total size of population i at time t, k_E

represents environmental stochasticity, R_0 is the density-independent population growth rate for

the average environment, $N_{migrants(t,i)}$ is the total number of migrants to population i at time t, and

$N_{residents(t,i)}$ is the total number of residents in population i at time t.

We imposed a mating function such that a population would deterministically go extinct if it

comprised only non-migrant females. In cases with an all female population and a mixture of

residents and migrants, we only included the number of migrant females in the density-

dependent effect of the demographic heterogeneity term $k_D*F_{mated(t,i)}$ because this effect

manifests via the number of eggs laid by females and only migrant females, being pre-mated,

would lay eggs. Weakly regularizing gamma priors were taken from Melbourne and Hastings

(2008). The expected equilibrium population size for the model is:

$$K = \frac{\log(R_0)}{\alpha}$$
                  (Eq. 2)

For each combination of the 4 introduction regimes and 2 recipient environment types (described

above), we simulated 500,000 replicate populations for 7 generations. All simulations were

performed in R (R Core Team 2016). This model captures the stochastic and deterministic

ecological dynamics of the *Tribolium* system well, but does not include evolutionary processes

(Melbourne and Hastings 2008, Hufbauer et al. 2015).

To parameterize the model, we censused 125 *Tribolium* populations one generation after

establishing them at various, known densities (between 5 and 200 individuals) on the novel

growth medium following the rearing procedure described in "Microcosm Experiment" below.

We fit the hierarchical model (Eq. 1) to these data in a Bayesian framework to generate posterior

distributions for each of the parameters using the nimble package in R (de Valpine et al. 2016).

We used a Metropolis-Hastings random walk sampler with 3 chains having 50,000 samples each

(including 10,000 samples that were removed for burn in). The chains converged (multivariate $\hat{R}$

for the 4 key parameters =1.01) and produced a sufficient number of effective samples (R_0: 757,

α: 548, k_E: 4103, k_D: 1703).

A single novel growth medium mixture was used for these 125 populations (mixture 5,

Supplementary Table 1), so we treated the estimated environmental stochasticity parameter, k_E,

as the expectation for a stable environment. We parameterized a fluctuating environment model

by increasing the standard deviation of the density-independent per capita population growth rate

by approximately 10% for any given population at any given time step. This was accomplished

by multiplying all of the samples from the posterior distribution of the environmental

stochasticity parameter, k_E, by 100/121 (derived by moment matching for the gamma

distribution). By using random sets of parameters drawn from the samples of the posterior

distributions, we were able to propagate the model uncertainty through the simulations and

combine it with the deterministic and stochastic dynamics represented by the model itself.

Microcosm Experiment

We founded 842 *Tribolium* populations with different introduction regimes (20 individuals in the

first generation, 10 individuals in the first 2 generations, 5 individuals in the first 4 generations,

or 4 individuals in the first 5 generations) and environments (stable or fluctuating) with between

96 and 120 replicate populations per treatment combination (Supplemental Table 3). Each

population was reared in a 4cm x 4cm x 6cm plastic box (AMAC Plastic Products) with 2

tablespoons (approximately 15 g) of freshly prepared growth medium that had been humidified

for at least 24 hours. The growth medium used for the source population (the natal environment

of the colonists) comprised 95% wheat flour (Pillsbury Co. or Gold Medal Products Co.) and 5%

brewers' yeast (Sensient Flavors). Colonists were introduced into a novel growth medium

comprising a small percentage of natal medium mixed with corn flour (Bob's Red Mill). All

populations were reared in one of two dark incubators at 31° and approximately 70% relative

humidity (standard conditions) and were haphazardly rotated between incubators weekly.

For each population in each generation, a known number of adults laid eggs for 24 hours in fresh

medium, and were then removed. Offspring were given 35 days to develop, and adults were then

censused. Censused adults laid eggs on freshly prepared growth medium for 24 hours,

completing their laboratory life cycle. We estimated the maximum observation error during

census to be 4.6% (median: 0%, mean: 0.26%) with no detectable difference across observers or

populations with different sizes.

Each replicate population experienced a novel environment that was either stable or randomly

fluctuating through time. The same novel growth medium mixture containing 99.05% novel corn

flour and 0.95% natal medium (mixture 5, Supplementary Table 1) was used for the stable

environment for the duration of the experiment, which preliminary results indicated would yield

a population growth rate of $\lambda = 1.2$ compared to a mean population growth rate of 3.36 on 100%

natal medium. To create the fluctuating environment, we randomly selected a novel growth

medium mixture from one of 9 possible media mixtures for each population in each generation.

Each population in the fluctuating environment therefore experienced a unique series of

environmental conditions. The 9 possible media mixtures represented a gradient of corn flour to

natal medium ratios, and were designed to yield expected population growth rates between 0.88

and 1.33 (Supplementary Table 1). We chose this range to mimic environmental stochasticity

measured in nature (Sæther and Engen 2002) while remaining within the bounds of biologically

realistic population growth (λ between 0.5 and 1.5) (Morris et al. 2008). The random sequence of

growth media was constrained such that the expected geometric mean population growth rate for

each population resembled expected growth of populations in the stable environment ($\lambda_{expected}$ =

1.2±0.05).

We estimated the amount of environmental stochasticity that we achieved in the fluctuating

environment as the difference in mean total stochasticity between populations in the fluctuating

and stable environments (Sæther and Engen 2002). We assumed that total stochasticity was a

combination of demographic and environmental stochasticity for populations in the fluctuating

environment, and that demographic stochasticity was the sole contributor to total stochasticity

for populations in the stable environment. Total stochasticity (demographic plus environmental)

of each population that did not experience extinction (n=667) was calculated as the variance of

the natural logarithms of its population growth rates through 7 generations:

$$s_{total} = s_{demographic} + s_{environmental} = var(\log(\lambda_t))$$

(Eq. 3)

where, for a particular population, s_{total} is its total stochasticity, $s_{demographic}$ is its demographic

stochasticity, $s_{environmental}$ is its environmental stochasticity (assumed to be 0 for populations

in the stable environment), and λ_t is its per capita population growth rate between generation *t-1*

and generation *t* (*t=1, 2, … , 7*). We only calculated total stochasticity for populations that did

not experience any extinction in order to capture the full temporal extent of environmental

fluctuations and because extinctions would have an infinite effect on this measure of

stochasticity.

Statistical analyses

We evaluated how our environment treatment affected variability in population growth rate (total

stochasticity from Eq. 3) using a linear mixed effects model with environment (stable or

fluctuating) as a fixed effect and block as a random intercept effect.

We used a mixed effects logistic regression with a logit link to predict the binary response of

establishment, and a mixed effects Poisson regression with a log link to analyze population size.

In both models, introduction regime, environment treatment, and their interaction were treated as

fixed effects, and block was treated as a random intercept effect.

We assessed the effect of temporary extinctions on the establishment probability and mean

population size by fitting the generalized linear mixed effects models described above to data

from the multiple introduction regimes (i.e. not the 20x1 regime) and with additional predictor

variables. To assess the effect of the presence of a temporary extinction, we included an

additional Boolean predictor for whether a population went temporarily extinct or not. To assess

the role of the total propagule pressure, we included a numeric predictor representing the number

of beetles introduced after the latest temporary extinction.

Group-level significance of fixed effects was tested using likelihood ratio tests on nested models,

and least-squares contrasts were used to compare levels of the fixed effects. All statistical

analyses were performed in R, version 3.3.2 (R Core Team 2016). Generalized linear mixed

models were fit using the lme4 package, version 1.1-12 (Bates et al. 2015) and pairwise

comparisons were made using the lsmeans package, version 2.25 (Lenth 2016).

Data availability

The raw experiment data, simulated population trajectories, R code for the simulations, R code

fitting the NBBg model, samples from the posterior distributions of the NBBg parameters, and R

code fitting the mixed effects models are available as supplemental material, on Figshare

(https://doi.org/10.6084/m9.figshare.4648865.v1), and on GitHub

(https://github.com/mikoontz/ppp-establishment).

Note on egg contamination

Laboratory procedures after generation 3 resulted in occasional egg contamination between

replicate populations of the same treatment. Some populations of 1 individual persisted when

they should have deterministically gone extinct, so we manually edited those population

trajectories such that they went extinct in the generation after having only 1 individual.

RESULTS

Simulations

Summary statistics for the posterior distributions of the four NBBg model parameters are given

in Supplementary Table 2.

Our simulations showed introduction regimes with more introduction events were more likely

establish a population by the seventh generation (Figure 1). Simulated introductions into a

stochastically fluctuating environment resulted in slightly lower population establishment for all

introduction regimes (difference of ~1%), but did not favor any particular regime (not shown).

The mean population sizes for each introduction regime/environment combination were

approximately equal in simulations. Mean population sizes only incorporated extant populations,

so they were slightly larger than the expectation for the equilibrium population size (Figure 2).

Microcosm experiment

Mean environmental stochasticity of populations in the fluctuating environment was 0.052 (95%

CI = 0.0073 to 0.0966; p = 0.023). This value is near the median value of 0.055 measured in

nature by Sæther and Engen (2002) in a metaanalysis of 35 avian populations, indicating that we

achieved biologically realistic fluctuations in population growth rate.

We found no evidence that the probability of establishment was affected by a main effect of

environment (χ^2=0.72, df=1, p=0.40), nor by an interaction between environment and

introduction regime (χ^2=3.49, df=3, p=0.32). However, there was strong support for an effect of

introduction regime on establishment probability (χ^2=59.76, df=3, p<0.0001). Pairwise

comparisons of the different introduction regimes averaged across the environment treatments

revealed that the 4x5 regime was the most likely to establish populations, with a probability of

about 0.98, whereas the 20x1 and 10x2 regimes were the least likely to establish populations,

with a probability reduced to about 0.8 (Figure 1, Figure 3).

The size of populations that persisted until generation 7 was shaped by significant effects of

introduction regime (χ^2=91.65, df=3, p<0.0001), environment treatment (χ^2=117.83, df=3,

p<0.0001), and their interaction (χ^2=44.62, df=3, p<0.0001). Populations established via more

introduction events were larger when averaged across the environment treatments. Extant

populations in the stable environment were larger than those in the fluctuating environment when

averaged across the introduction regimes. The interaction manifests as the benefit of a stable

environment increases with more, smaller introduction events (Figure 2, Figure 4).

Extinctions accumulated regularly throughout the experiment period, with 101 out of 842

populations (12.0%) going extinct by generation 7 (Figure 3). The additional introductions that

some populations received often restored a population that had temporarily gone extinct. Out of

602 populations that received more than 1 introduction (i.e. not the 20x1 introduction regime),

104 of them (17.3%) temporarily went extinct at least once before being replenished by

additional colonizing individuals. Twelve populations were rescued in this way at least twice,

and one population was rescued in this way three times.

Temporary extinctions significantly affected colonization success. The presence of a temporary

extinction significantly reduced average establishment probability from 92.4% to 82.1%

(difference = -1.13 on logit scale, 95% CI = -0.22 to -2.04, χ^2 = 5.44, df = 1, p = 0.020) and mean

population size from 47.8 to 45.4 (difference = -0.052 on log scale, 95% CI = -0.02 to -0.09, χ^2 =

9.42, df = 1, p = 0.0021). Each additional colonist contributing to a population after the latest

temporary extinction significantly increased the mean population size (estimate = 0.005 on the

log scale, 95% CI = 6.4e-05 to 0.01, χ^2 = 3.94, df = 1, p = 0.047).

Results from the simulations are expected to represent the dynamics in the microcosm

experiment if the assumption of the model on which the simulations were built holds—that only

demographic processes are responsible for the dynamics observed in the experiment. However,

establishment probability at generation 7 was equal to or greater than that expected from the

simulations (Figure 1) and mean population sizes were much larger in the experiment than in the

simulations (Figure 2), together suggesting an influence of non-demographic mechanisms.

DISCUSSION

We assessed how the number and size of introduction events through time drive colonization

success in a novel environment when the total number of individuals introduced to a site is fixed.

We considered novel environments that were either stable or randomly fluctuating in quality

through time, and evaluated populations through 7 discrete generations. We approached this

question in two ways: 1) stochastic simulations of a demographic population dynamics model

parameterized with empirical data, and 2) a highly replicated laboratory microcosm experiment.

By coupling these approaches, we were able to expand and test the theoretical understanding of

how the introduction regime affects colonization in stable and fluctuating environments as well

as to develop new avenues for research when observations did not align with predictions. We

found that several, small introductions increase colonization success and that demographic

processes alone are insufficient to explain the dynamics observed in the experiment.

We found minimal to no effect of a biologically realistic level of environmental stochasticity on

establishment probability in demographic simulations or in the microcosm experiment. Our

results corroborate those of Cassey et al. (2014) who simulated introductions through time and

also found a minimal effect of environmental stochasticity on establishment probability. Cassey

et al. (2014) further found a minimal effect of random, infrequent catastrophes and bonanzas,

suggesting that increasing temporal environmental variability in the broadest sense (i.e.,

encompassing environmental stochasticity, random catastrophes, and bonanzas) does not

magnify the benefit of several, small introductions through time. A minimal role of

environmental stochasticity contrasts with the results of Grevstad (1999) who found with

simulations that several, small introductions would produce an especially high establishment

probability compared to a single, large introduction in a variable environment. The difference in

findings is perhaps due to a difference in the kinds of introduction regimes modeled: Grevstad

(1999) simulated multiple introductions in space with strong environmental stochasticity often

leading to catastrophic mortality, while we focused on introductions separated in time with less

extreme environmental stochasticity leading to moderate fluctuations in growth rates (Lande

1993).

There also did not appear to be a strong effect of environmental stochasticity on population size

in the demographic simulations. All treatments in the simulations reached similar mean

population size by generation 7, which was slightly higher than the equilibrium size due to our

measure only including extant populations. This suggests that strong density dependence in the

demographic simulations was the primary influence on population size.

Population growth rate may interact with introduction regime to affect colonization success.

Cassey et al. (2014) suggested that their lower simulated mean population growth rates, where R

was between 1.0 and 1.38, compared to that of Grevstad (1999), where R was equal to 2.0,

explained why they found that a single large introduction always led to a greater establishment

probability, while Grevstad (1999) found several small introductions to be more successful.

Further, Wittmann et al. (2014) show that population establishment occurs faster with fewer

larger introductions in populations with growth rates below one or with growth rates that

fluctuate above and below one. Wittmann et al. (2014) also show that population establishment is

faster with several smaller introductions when population growth rate is consistently greater than

one. However, our expected mean population growth rate was relatively low ($R = 1.132$;

Supplementary Table 2) and we still found that several small introductions had the greatest

colonization success. Also, several small introductions performed best in both the fluctuating

environment (population growth fluctuated above and below one; Supplementary Table 1) and in

the stable environment (population growth consistently greater than one; Supplementary Table 1)

(Figure 1, Figure 2). Thus, our results contradict the notion that net reproductive rate is the key

control on how introduction regime affects colonization success, which merits further

investigation.

We observed striking differences in the measures of colonization success between the microcosm

experiment and the demographic simulations. We found that establishment probability increased

with the number of introduction events in both the experiment and the simulations, but that all

experiment establishment probabilities equaled or exceeded expectations from simulations. In

the experiment, mean population size in stable environments was greater than in fluctuating

environments, and there was an interaction between environment and introduction regime

whereby the mean population size was increasingly greater in stable compared to fluctuating

environments as the number of introductions increased. Furthermore, populations grew larger by

generation 7 in the experiment than in the simulations.

Differences between the results of the simulations and of the microcosm experiment suggest that

the demographic processes captured by the model do not account for all of the biological

processes that occurred in the microcosm. Recent work shows that adaptation to the novel, harsh

environment from standing variation is possible in this species (Hufbauer et al. 2015), and likely

explains the greater establishment probability and population sizes in the microcosm compared

to expectations derived from demographic simulations which do not include adaptation.

The rescue effect of immigration can act demographically by increasing the size of populations

(Hufbauer et al. 2015). Certainly, demographic rescue played a critical role for the 104

populations that went extinct temporarily until another introduction event revived them. Those

temporary extinctions had lasting effects on colonization success. Colonization success declined

for populations that experienced a temporary extinction, and the mean population size

significantly increased if more colonists contributed to the population after a temporary

extinction. These results reflect the overarching importance of total propagule pressure

regardless of introduction regime.

The rescue effect can also act genetically by increasing the fitness of populations (Frankham

2015, Hufbauer et al. 2015, Whiteley et al. 2015). The experimental immigrants all came from

the same source population, so it is unlikely that gene flow from migrants united previously

separated alleles into high-fitness genotypes (Novak 2007). Thus, a more likely mechanism by

which immigration increased mean population size beyond expectations was by relieving

inbreeding depression or counteracting drift-induced allele loss. Small populations are more

prone to experiencing increased homozygosity and inbreeding depression, which can reduce

population growth rates and increase extinction risk (McCauley and Wade 1981, O'Grady et al.

2006, Szűcs et al. 2017). Even small amounts of gene flow can alleviate these effects, so the

additional small introductions of mated individuals from the external source population were

well-suited to bring about longer-term relief (Slatkin 1985, Hufbauer et al. 2015). However,

Cassey et al. (2014) found that simulated inbreeding depression was especially detrimental for

several, small introductions through time, so other mechanisms are likely at play.

One such mechanism may be adaptation of the incipient population, which is affected by

sustained immigration. Introductions to a harsh, novel habitat can result in adaptive evolution

with the right amount of gene flow if additional immigrants to a declining population prevent

extinction long enough to allow for adaptation to occur (Holt and Gomulkiewicz 1997). However,

too much gene flow can lead to genetic swamping whereby the homogenizing effects of gene

flow overpowers ongoing local adaptation (Lenormand 2002). This may have been the case for

populations in the 10x2 introduction regime, which had the highest average migration rate,

lowest establishment probability, and lowest mean population size. Alternatively, negative

density dependence may have reduced population fitness when migration rates were high,

reducing population growth rates and hampering the spread of adaptive alleles (Holt and

Gomulkiewicz 1997). Though not significant, the lower establishment probability and mean

population size in the 10x2 introduction regime warrants further work to assess whether they

exemplify the yet-unseen scenario described by Blackburn et al. (2015) in which maladaptive

gene flow from multiple introductions hampers population establishment in a novel range. More

broadly, further study is necessary to evaluate how immigration affected adaptation in this

system, if at all (Gomulkiewicz and Holt 1995, Boulding and Hay 2001).

CONCLUSION

Our experimental results suggest that several, small introductions through time lead to greater

colonization success in a novel habitat, and that introductions into a stable recipient environment

lead to larger population sizes, but not greater establishment probability. Further, introductions to

a stable recipient environment are especially beneficial to populations established with more

introduction events. These results defied our expectations derived from parallel simulations of a

model that included demographic processes but not evolutionary ones, so we suspect a genetic

mechanism might be at work. Genetic mechanisms are rarely incorporated when simulating the

effect of introduction regime on colonization (but see Cassey et al. 2014), and our multi-

generation microcosm is unique in bringing evolutionary processes to bear on parsing two key

components of propagule pressure in an experimental setting.

For invasions, our results highlight the importance of preventing further introductions to the

same location, even for established species. For conservation and biological control, our results

suggest that emphasis should be placed on increasing the number of introductions or re-

introductions to a location, rather than increasing the size of those events if Allee effects are

weak. Sustained introduction efforts should also bring about concomitant benefits in the form of

longer-term monitoring, increased data collection, and more opportunities for experimentation

and adaptive management (Lockwood et al. 2005, Godefroid et al. 2011).

ACKNOWLEDGMENTS

We thank Charlotte Hoover, Kaitlyn Adkins, Mary Catherine Cochran, Don Waters, Stacy

Endriss, Marianna Szűcs, and Jenny Rickford for help with data collection. Funding was

provided by NSF Graduate Research Fellowship Grants #DGE-1321845 Amend. 3 (to MJK) and

#DGE-1106400 (to MFO), NSF Grant DEB-0949619 (to RAH), and NSF Grant DEB-0949595

(to BAM). RAH also acknowledges the support of the USDA via the Colorado Experiment

Station. We thank the lab groups of Ruth Hufbauer, Paul Ode, Andrew Norton, Andrew Latimer,

Truman Young, and Jenny Gremer for comments on the project and manuscript.

REFERENCES

Bates, D. et al. 2015. Fitting linear mixed-effects models using lme4. - J. Stat. Softw. 67: 1–48.

Berggren, A. 2001. Colonization success in Roesel's bush-cricket Metrioptera roeseli: The
effects of propagule size. - Ecology 82: 274–280.

Blackburn, T. M. and Duncan, R. P. 2001. Establishment patterns of exotic birds are constrained
by non-random patterns in introduction. - J. Biogeogr. 28: 927–939.

Blackburn, T. M. et al. 2015. The influence of numbers on invasion success. - Mol. Ecol.: 1–12.

Boulding, E. G. and Hay, T. 2001. Genetic and demographic parameters determining population

persistence after a discrete change in the environment. - Heredity (Edinb). 86: 313–24.

Cassey, P. et al. 2005. Concerning invasive species: Reply to Brown and Sax. - Austral Ecol. 30:

475–480.

Cassey, P. et al. 2014. A population model for predicting the successful establishment of

introduced bird species. - Oecologia 175: 417–428.

Colautti, R. I. et al. 2006. Propagule pressure: A null model for biological invasions. - Biol.

Invasions 8: 1023–1037.

de Valpine, P. et al. 2016. Programming with models: writing statistical algorithms for general

model structures with NIMBLE. - J. Comput. Graph. Stat. 8600: 1–20.

Drake, J. M. et al. 2005. Propagule pressure and persistence in experimental populations. - Biol.

Lett. 1: 480–3.

Drolet, D. and Locke, A. 2016. Relative importance of propagule size and propagule number for

establishment of non-indigenous species: A stochastic simulation study. - Aquat. Invasions

11: 101–110.

Fauvergue, X. et al. 2012. The biology of small, introduced populations, with special reference to

biological control. - Evol. Appl. 5: 424–43.

Frankham, R. 2015. Genetic rescue of small inbred populations: Meta-analysis reveals large and

consistent benefits of gene flow. - Mol. Ecol. 24: 2610–2618.

Godefroid, S. et al. 2011. How successful are plant species reintroductions? - Biol. Conserv. 144:

672–682.

Gomulkiewicz, R. and Holt, R. D. 1995. When does evolution by natural selection prevent

extinction? - Evolution (N. Y). 49: 201–207.

Grevstad, F. S. 1999. Factors influencing the chance of population establishment: Implications

for release strategies in biocontrol. - Ecol. Appl. 9: 1439–1447.

Grevstad, F. S. et al. 2011. Revisiting release strategies in biological control of weeds: Are we

using enough releases? - XIII Int. Symp. Biol. Control Weeds: 368–376.

Haccou, P. and Iwasa, Y. 1996. Establishment probability in fluctuating environments: A

branching process model. - Theor. Popul. Biol. 50: 254–280.

Haccou, P. and Vatutin, V. 2003. Establishment success and extinction risk in autocorrelated

environments. - Theor. Popul. Biol. 64: 303–314.

Hedge, L. H. et al. 2012. Manipulating the intrinsic parameters of propagule pressure:

Implications for bio-invasion. - Ecosphere 3: 1–13.

Holt, R. D. and Gomulkiewicz, R. 1997. How does immigration influence local adaptation? A

reexamination of a familiar paradigm. - Am. Nat. 149: 563–572.

Hopper, K. R. and Roush, R. T. 1993. Mate finding, dispersal, number released, and the success

of biological control introductions. - Ecol. Entomol. 18: 321–331.

Hufbauer, R. A. et al. 2015. Three types of rescue can avert extinction in a changing

environment. - Proc. Natl. Acad. Sci. 112: 10557–10562.

Jeschke, J. M. 2014. General hypotheses in invasion ecology. - Divers. Distrib. 20: 1229–1234.

Lande, R. 1988. Genetics and demography in biological conservation. - Science 241: 1455–1460.

Lande, R. 1993. Risks of population extinction from demographic and environmental

stochasticity and random catastrophes. - Am. Nat. 142: 911–927.

Lenormand, T. 2002. Gene flow and the limits to natural selection. - Trends Ecol. Evol. 17: 183–

189.

Lenth, R. V 2016. Least-squares means: The R package lsmeans. - J. Stat. Softw. 69: 1–33.

Lockwood, J. L. et al. 2005. The role of propagule pressure in explaining species invasions. -

Trends Ecol. Evol. 20: 223–8.

Lodge, D. M. 1993. Biological invasions: Lessons for ecology. - Trends Ecol. Evol. 8: 133–137.

Mack, R. N. et al. 2000. Biotic invasions: Causes, epidemiology, global consequences, and

control. - Ecol. Appl. 10: 689–710.

McCauley, D. E. and Wade, M. J. 1981. The populational effects of inbreeding in Tribolium. -

Heredity (Edinb). 46: 59–67.

Melbourne, B. A. and Hastings, A. 2008. Extinction risk depends strongly on factors

contributing to stochasticity. - Nature 454: 100–3.

Memmott, J. et al. 2005. The effect of propagule size on the invasion of an alien insect. - J. Anim.

Ecol. 74: 50–62.

Morris, W. F. et al. 2008. Longevity can buffer plant and animal populations against changing

climatic variability. - Ecology 89: 19–25.

Novak, S. J. 2007. The role of evolution in the invasion process. - Proc. Natl. Acad. Sci. 104:

3671–2.

O'Grady, J. J. et al. 2006. Realistic levels of inbreeding depression strongly affect extinction risk

in wild populations. - Biol. Conserv. 133: 42–51.

R Core Team 2016. R: A language and environment for statistical computing. http://www.r-

project.org/. in press.

Ricciardi, A. 2007. Are modern biological invasions an unprecedented form of global change? -

Conserv. Biol. 21: 329–36.

Sæther, B.-E. and Engen, S. 2002. Pattern of variation in avian population growth rates. - Philos.

Trans. R. Soc. Lond. B. Biol. Sci. 357: 1185–95.

Sakai, A. K. et al. 2001. The population biology of invasive species. - Annu. Rev. Ecol. Syst. 32:

305–332.

Schreiber, S. J. and Lloyd-Smith, J. O. 2009. Invasion dynamics in spatially heterogeneous

environments. - Am. Nat. 174: 490–505.

Simberloff, D. 1989. Which insect introductions succeed and which fail? - In: Drake et al., J. A.

(ed), Biological Invasions: a Global Perspective. John Wiley & Sons Ltd, pp. 61–75.

Simberloff, D. 2009. The role of propagule pressure in biological invasions. - Annu. Rev. Ecol.

Evol. Syst. 40: 81–102.

Sinclair, J. S. and Arnott, S. E. 2016. Strength in size not numbers: Propagule size more

important than number in sexually reproducing populations. - Biol. Invasions 18: 497–505.

Slatkin, M. 1985. Gene flow in natural populations. - Annu. Rev. Ecol. Syst. 16: 393–430.

Szűcs, M. et al. 2017. Genetic and demographic founder effects have long-term fitness

consequences for colonising populations. - Ecol. Lett.: 1–7.

Taylor, S. S. et al. 2005. Successful island reintroductions of New Zealand robins and

saddlebacks with small numbers of founders. - Anim. Conserv. 8: 415–420.

Van Allen, B. G. and Rudolf, V. H. W. 2013. Ghosts of habitats past: Environmental carry-over

effects drive population dynamics in novel habitat. - Am. Nat. 181: 596–608.

Whiteley, A. R. et al. 2015. Genetic rescue to the rescue. - Trends Ecol. Evol. 30: 42–49.

Wittmann, M. J. et al. 2014. Decomposing propagule pressure: the effects of propagule size and

propagule frequency on invasion success. - Oikos 123: 441–450.

Zenni, R. D. and Nuñez, M. A. 2013. The elephant in the room: The role of failed invasions in

understanding invasion biology. - Oikos 122: 801–815.

Figure 1. Establishment probabilities for each introduction regime assessed at generation 7.

Triangles represent results from simulations. Dot-whiskers represent estimates and 95%

confidence intervals from the mixed effects logistic regression model fit to data from the

microcosm experiment. Different letters over dot-whiskers represent introduction regimes with

significantly different establishment probabilities in the microcosm experiment.

Figure 2. Mean sizes of extant populations for each introduction regime/environment

combination in generation 7. Triangles represent results from simulations. Dot-whiskers

represent estimates and 95% confidence intervals from the mixed effects Poisson regression

model fit to data from the microcosm experiment. Different letters over dot-whiskers represent

introduction regimes with significantly different mean population sizes. The dashed line

represents the theoretical equilibrium population size derived using Eq. 2, which includes

extinction and is therefore lower than the mean population size from the simulations.

Figure 3. Percent of microcosm populations that were established in each generation for the 4

different introduction regimes. Data are pooled across the two environmental variability

treatments. The introduction regimes are as follows: 20x1 = 20 individuals in the first generation,

10x2 = 10 individuals in each of the first two generations, 5x4 = 5 individuals in each of the first

4 generations, and 4x5 = 4 individuals in each of the first 5 generations.

Figure 4. Population trajectories for 842 populations of *Tribolium* in the microcosm experiment. The first panel represents all populations in stable environments (n = 418) and the second panel represents all populations in fluctuating environments (n = 424). Vertical steps in the population size represent additional migrant individuals in introduction regimes with more than 1 introduction. Light grey lines represent individual population trajectories, and colored lines represent the mean size of extant populations for each of the 4 introduction regimes. The introduction regimes are as follows: 20x1 = 20 individuals in the first generation, 10x2 = 10 individuals in each of the first two generations, 5x4 = 5 individuals in each of the first 4 generations, and 4x5 = 4 individuals in each of the first 5 generations. Shading around the colored lines represents the mean size of extant populations plus and minus one standard error.