A first-in-class inhibitor of apicomplexan FtsH1 disrupts plastid biogenesis in human pathogens

Katherine Amberg-Johnson[1,3], Sanjay B. Hari[4], Suresh M. Ganesan[5], Hernan A. Lorenzi[6], Robert T. Sauer[4], Jacquin C. Niles[5], Ellen Yeh[1,2,3,*]

[1]Department of Biochemistry, [2]Pathology, and [3]Microbiology and Immunology, Stanford Medical School
[4]Department of Biology, Massachusetts Institute of Technology
[5]Department of Biological Engineering, Massachusetts Institute of Technology
[6]Department of Infectious Disease, The J. Craig Venter Institute
*Corresponding author

The malaria parasite *Plasmodium falciparum* and related apicomplexan pathogens contain an essential plastid organelle, the apicoplast, which is a key anti-parasitic target. Apicoplast biogenesis depends on novel, but largely cryptic, mechanisms for protein/lipid import and organelle inheritance during parasite replication. These critical pathways present untapped opportunities to discover new parasite-specific drug targets. We used an innovative screen to identify the natural product actinonin as a first-in-class antimalarial compound inhibiting apicoplast biogenesis. Resistant mutation, chemical-genetic interaction, and biochemical inhibition demonstrate that the unexpected target of actinonin in *P. falciparum* and *Toxoplasma gondii* is FtsH1, a homolog of a bacterial membrane AAA+ metalloprotease. *Pf*FtsH1 is the first novel factor required for apicoplast biogenesis identified in an unbiased screen. Our findings demonstrate that FtsH1 is a novel and, importantly, druggable antimalarial target. Development of FtsH1 inhibitors will have significant advantages over existing apicoplast-targeting compounds with improved drug kinetics and multistage efficacy against multiple human parasites.

Introduction

There is an urgent need for antimalarials with novel mechanisms-of-action to combat resistance to frontline drugs. The apicoplast is a plastid organelle uniquely found in *Plasmodium* parasites that cause malaria and related apicomplexan pathogens[1,2]. Though non-photosynthetic, the apicoplast is required for the biosynthesis of essential metabolites in the organisms that retained it[3–9]. Acquired by a secondary eukaryote-eukaryote endosymbiosis, it utilizes bacterial and organellar pathways distinct from those of mammalian host cells. Given its unique and essential biology, the apicoplast is an ideal source of parasite-specific drug targets that minimize host off-target toxicity.

Despite its biomedical potential, broadly effective antimalarials targeting the apicoplast have been elusive. The first class of apicoplast inhibitors to be identified was drugs like doxycycline, clindamycin, and chloramphenicol that block prokaryotic protein synthesis[10,11]. The antimalarial effects of these clinically-approved antibiotics were noticed well before they were shown to inhibit the bacterial-like ribosome in the apicoplast[12]. Unfortunately translation inhibitors cause a characteristic "delayed death" *in vitro* in which parasite growth inhibition occurs in the second replication cycle after drug treatment. The delayed effect is related to their mechanism-of-action and is also seen with

DNA replication and transcription inhibitors that block apicoplast gene expression. Delayed death manifests as a slow onset-of-action of translation inhibitors that limits their antimalarial efficacy and clinical use.

The next druggable targets in the apicoplast were several prokaryotic metabolic pathways. Fosmidomycin, an inhibitor of MEP isoprenoid precursor biosynthesis in the apicoplast, causes parasite growth inhibition in a single replication cycle *in vitro* and rapid parasite clearance in human clinical trials[3,13–15]. Unfortunately initial parasite clearance is followed by recrudescent infections in 50% of patients. These treatment failures have brought into question the clinical utility of fosmidomycin and other MEP inhibitors. Yet despite the expectation that the apicoplast would serve many essential functions, we showed that only isoprenoid precursor biosynthesis is required during the symptomatic blood stage of *Plasmodium*[8]. The apicoplast's limited function in blood-stage *Plasmodium* precludes opportunities to target alternative metabolic pathways.

Clearly the pipeline for essential and druggable apicoplast targets has run dry. New approaches are needed to identify drug targets that avoid the known liabilities of targeting apicoplast gene expression and metabolism. Until now, identification of drug targets in the apicoplast has been based on conserved bacterial and primary chloroplast pathways whose functions can be inferred from model organisms. However, many essential pathways for protein/lipid import into the apicoplast and organelle inheritance during parasite replication will be unique to secondary plastids and evolutionarily divergent from model eukaryotic biology[16,17]. Drug targets in these organelle biogenesis pathways are likely to be 1) essential to parasite survival, even in the context of the apicoplast's drastically reduced metabolic function, 2) required for all proliferative stages of the parasite life cycle, and 3) conserved across apicomplexan parasites. Thus apicoplast biogenesis represents a promising but unexplored frontier in antimalarial drug discovery.

While forward genetic screens have been extremely powerful in uncovering novel cellular pathways in model organisms, they are not currently tractable in *Plasmodium*. To circumvent these limitations, we took a chemical-genetic approach from phenotypic screening to target identification. We first identified the natural product actinonin as a novel inhibitor of apicoplast biogenesis. We then uncovered the essential role of the membrane metalloprotease FtsH1 in organelle biogenesis in apicomplexan pathogens. FtsH1 is a druggable target in apicoplast biogenesis that offers advantages over the mechanisms-of-action of existing apicoplast inhibitors and has a ready hit compound to pursue drug development.

Results

Identification of a novel apicoplast biogenesis inhibitor

We screened >400 growth-inhibitory antimalarial compounds with unknown mechanisms-of-action to identify novel inhibitors of apicoplast biogenesis (Extended Data Table 1)[18–22]. Our selection criteria aimed to identify compounds that 1) caused parasite growth inhibition (essential target) in a single replication cycle (avoiding delayed death), 2) were specific in their disruption of the apicoplast, and 3) resulted in loss of the apicoplast during parasite replication. This unique inhibition phenotype distinguished

92   selected compounds from known apicoplast gene expression or metabolism inhibitors,
93   ensuring discovery of a novel class of inhibitors.
94   A single natural product antibiotic, actinonin, matched our criteria[23]. Consistent
with its reported antimalarial activity, actinonin caused *P. falciparum* growth inhibition
in a single replication cycle (EC_{50} = 3.2 μM; 95% CI 2.5-4.1)[19,24]. Because isoprenoid
precursor biosynthesis is the only essential function of the apicoplast in blood-stage *P.*
falciparum, any disruption of the apicoplast, including complete loss of the organelle, can
be rescued by addition of the isoprenoid precursor, isopentenyl pyrophosphate (IPP), in
the growth media[8]. Indeed actinonin's EC_{50} shifted nearly 20-fold in the presence of IPP
to 61 μM (95% CI 50-75). Although the translation inhibitor chloramphenicol also
caused growth inhibition rescued by IPP, parasite growth inhibition and resulting
apicoplast biogenesis defects were delayed by one replication cycle following drug
treatment (Extended Data Fig. 1b and 2a,c). Compared to chloramphenicol, actinonin
clearly had more rapid antimalarial activity. These results demonstrate that actinonin
specifically inhibits the apicoplast without a delayed death phenotype (Fig. 1a; Extended
Data Fig. 1a; Extended Data Table 1).
Previously actinonin was observed to cause a block in the morphologic
development of the apicoplast during parasite replication[24]. However, these defects were
observed without IPP growth rescue when parasite replication was also inhibited. Absent
IPP rescue, the MEP inhibitor fosmidomycin also showed a block in apicoplast
development. Based on the observed similarities in their inhibition phenotype, the authors
concluded that actinonin, like fosmidomycin, inhibited apicoplast metabolism. We sought
to clarify these earlier observations using IPP rescue to distinguish between specific
apicoplast defects caused by the inhibitor and nonspecific defects observed in a non-
replicating parasite. Using quantitative PCR (qPCR) to detect the apicoplast genome and
microscopy to localize an apicoplast-targeted GFP, we show that actinonin-treated *P.*
falciparum, rescued for growth by IPP, no longer replicated their apicoplast genome and
produced daughter parasites lacking apicoplasts, consistent with a defect in apicoplast
biogenesis (Fig. 1b-c, Extended Data Fig. 2a,e)[8]. In contrast, both the copy number of the
apicoplast genome and apicoplast development were unaffected by fosmidomycin
treatment under IPP rescue, indicating that the previously reported apicoplast biogenesis
defects were nonspecific (Extended Data Fig 1b and 2a,d). Taken together, actinonin's
inhibition phenotype distinguishes it from known inhibitors that disrupt apicoplast gene
expression and metabolism and indicates that it has a novel mechanism-of-action in
organelle biogenesis (Extended Data Fig 1b and 2).
127
Actinonin is unlikely to inhibit the peptide deformylase in *Plasmodium* parasites
To identify actinonin's molecular target, we first took a candidate-based
approach. In bacteria and mammalian mitochondria, actinonin potently inhibits a peptide
deformylase (PDF) which catalyzes the co-translational removal of the formyl group
from nascent chains[25]. The apicoplast translation machinery is prokaryotic in origin and
therefore contains a PDF. Previously actinonin was shown to inhibit the enzymatic
activity of purified *P. falciparum* PDF with a reported IC_{50}= 2.5 μM[26]. The potency of
this inhibition is 10^{2-3} fold less than those reported for *E. coli* or human PDF[25,27,28]. Given
the role of *Pf*PDF in apicoplast translation, its inhibition is expected to give a delayed
death phenotype as observed with other translation inhibitors, whereas we clearly

observed growth inhibition in a single replication cycle (Figure 1). Puzzled by this
inconsistency, we used both genetic and chemical approaches to ascertain whether the
*Pf*PDF is the target of actinonin.
141   In bacteria, resistance to actinonin is conferred by overexpression of PDF[29]. We
constructed a *P. falciparum* strain containing a second copy of *Pf*PDF-myc in which
expression was regulated by tetR-DOZI-binding aptamer sequences[30]. As expected,
*Pf*PDF-myc expression was observed in the presence of anhydrotetracycline, which
disrupts the tetR-DOZI repressor-aptamer interaction (Extended Data Fig. 3A). We
assessed the import of overexpressed *Pf*PDF-myc into the apicoplast by determining its
apicoplast-dependent protein cleavage. Lumenal apicoplast proteins, including *Pf*PDF,
contain an *N*-terminal transit peptide sequence that is required for trafficking to the
apicoplast and is cleaved upon import[16]. In normal parasites, *Pf*PDF-myc was detected as
a majority processed and minority unprocessed protein (Extended Data Fig 3A-B). In
IPP-rescued parasites lacking their apicoplast, *Pf*PDF-myc accumulates as only the
unprocessed protein (Extended Data Fig 3B)[8]. These results indicate that *Pf*PDF-myc is
properly trafficked. Contrary to what is seen in bacteria, *Pf*PDF-myc overexpression did
not confer actinonin resistance indicating that *Pf*PDF may not be the target of actinonin
(Extended Data Fig. 3C). Because the lack of actinonin resistance may also have been
due to insufficient overexpression or nonfunctional *Pf*PDF-myc, we attempted to
generate a strain with *pdf* expression regulated at the endogenous loci but were unable
obtain viable parasites in two separate transfections performed in parallel with successful
positive controls.
160   We also assessed whether the actinonin target functions downstream of the
ribosome, as is expected for *Pf*PDF. In mitochondria, translation inhibitors suppress the
effects of actinonin since a block in translation supercedes a block in PDF activity[31].
Similarly, we expected that if PDF was the target of actinonin leading to growth
inhibition in a single replication cycle, then co-treatment with chloramphenicol would
result in a delayed death phenotype, whereby translation inhibition would mask the
effects of PDF inhibition. Surprisingly, chloramphenicol did not suppress the effects of
actinonin (Extended Data Fig. 3D). Taken together, our results indicate that 1) actinonin
does not cause delayed death as would be expected for *Pf*PDF inhibition, 2)
overexpression of *Pf*PDF did not confer actinonin resistance, and 3) the effect of
actinonin was not suppressed by inhibition of the ribosome upstream of *Pf*PDF. These
results are inconsistent with *Pf*PDF being the target of actinonin.
172
173   **Actinonin-resistant *T. gondii* contain a mutation in *Tg*FtsH1**
174   Because the target of actinonin in *P. falciparum* did not appear to be conserved in
model organisms, we turned to an unbiased approach. Our collaborators and we
independently attempted to isolate actinonin-resistant *P. falciparum* but failed using
multiple selection methods, including chemical mutagenesis, that successfully selected
resistance against other compounds (Prof. David Fidock, personal communication)[21,22].
Therefore, we turned to *Toxoplasma gondii,* a related apicomplexan parasite, which
contains an apicoplast of the same evolutionary origin, because it is easier to grow to
large numbers and to genetically modify.
182   To determine whether actinonin disrupts apicoplast biogenesis in *T. gondii*, we
characterized its inhibition phenotype. Curiously, *T. gondii* also shows a delayed death

phenotype with apicoplast translation inhibition, whereby drug treatment in the first lytic
cycle results in parasite growth inhibition in the second cycle[10]. However, the mechanism
underlying delayed death in *T. gondii* is distinct from that in *Plasmodium*. Delayed death
in *T. gondii* is associated with defects in apicoplast biogenesis and formation of parasites
that lack apicoplasts during the first lytic cycle[32–34]. Though these "apicoplast-minus"
parasites are viable in the first lytic cycle, they cannot replicate in the second lytic cycle.
(In contrast, *Plasmodium* parasites treated with translation inhibitors exhibit apicoplast
biogenesis defects in the second replication/lytic cycle, wherein loss of the apicoplast
immediately blocks parasite replication[11].) Consistent with an apicoplast biogenesis
defect in *T. gondii*, actinonin showed delayed death growth inhibition (EC$_{50}$= 14 μM;
95% CI 13-14) with formation of apicoplast-minus parasites during drug treatment (Fig.
2a and Extended Fig. 4a)[32].
Because actinonin disrupted apicoplast biogenesis in both *T. gondii* and *P.*
falciparum, suggesting a common target, we selected actinonin-resistant *T.gondii* and
determined the whole-genome sequences for eight independently selected clones (Fig.
2b). Remarkably, five clones harbored the same mutation encoding a N805S variant in
the metalloprotease domain of the membrane-bound AAA+ protease *TgFtsH1*
(TGGT1_259260) (Fig. 2d; Extended Data Table 2). *Tg*FtsH1 was compelling as the
actinonin target for two reasons. First, *Tg*FtsH1 localizes to the apicoplast[35]. Second,
actinonin is a peptide mimetic containing a metal-binding hydroxamic acid, a class of
molecules that typically binds metalloproteases[25,36]. Replacement of the endogenous
TgFtsH1 locus with the allele encoding *TgFtsH1(N805S)* caused a 3.5-fold shift in the
actinonin EC$_{50}$, similar to the resistance observed in the actinonin-resistant clones (Fig.
2c). Notably growth inhibition at this higher actinonin concentration did not result in
delayed death, suggesting that actinonin no longer bound FtsH1 and growth inhibition
was the result of secondary targets (Extended Data Fig. 4b). In contrast, no increase in the
actinonin EC$_{50}$ was observed when the endogenous locus was replaced with the WT allele
(Fig. 2c). Taken together, the known metalloprotease binding of actinonin, the predicted
metalloprotease activity of TgFtsH1, and the validated actinonin-resistant mutation in
TgFtsH1 support FtsH1 as the target of actinonin in *T. gondii*, providing a strong
candidate for validation in *P. falciparum*.
215
C-terminal cleavage of the *P. falciparum* FtsH1 homolog is dependent on the
apicoplast
The *P. falciparum* genome contains three FtsH homologs. One of these, *Pf*FtsH1
(Pf3D7_1239700) is most closely related to *Tg*FtsH1 by phylogenetic analysis[37]. Unlike
*Tg*FtsH1, *Pf*FtsH1 was previously reported to localize to mitochondria, not the
apicoplast[37]. However the same study reported that, like *Tg*FtsH1, *Pf*FtsH1 undergoes
processing. The mitochondrial localization was most clearly demonstrated for the *C*-
terminally tagged fragment, but the localization of the *N*-terminal fragment containing the
ATPase and protease domains was unclear. Proteins required for apicoplast biogenesis
may localize to various cellular compartments outside of the apicoplast. For example,
both the dynamin DrpA and Atg8 required for apicoplast biogenesis are cytoplasmic
proteins, whereas apicoplast protein trafficking machinery is likely to exist in the ER[32,34].
Close association of the mitochondria and apicoplast has also been observed and may be
important for organelle biogenesis[38,39].

To clarify its localization and function, we constructed a *P. falciparum* strain in
which the endogenous *PfFtsH1* locus was modified with a *C*-terminal FLAG epitope and
3' UTR tetR-DOZI-binding aptamer sequences for regulated expression[30]. We first
sought to localize *Pf*FtsH1 with the *C*-terminal FLAG epitope. However only the FLAG-
tagged Cas9 expressed in this strain was detected with no visible band corresponding to
the size of *Pf*FtsH1, consistent with the previous report that the *C*-terminus of the protein
is cleaved (Extended Data Fig. 5a). *Tg*FtsH1 was proposed to be *C*-terminally cleaved in
the apicoplast[40]. We hypothesized that if cleavage of *Pf*FtsH1 is a result of apicoplast
localization, then IPP-rescued parasites lacking an apicoplast would retain an intact
*Pf*FtsH1-FLAG. Indeed, *Pf*FtsH1-FLAG was detected in these apicoplast-minus
parasites, indicating that *C*-terminal cleavage of *Pf*FtsH1 is dependent on the apicoplast
(Extended Data Fig. 5). This evidence strongly suggests that *Pf*FtsH1 traffics to the
apicoplast, as has been shown for *Tg*FtsH1.
243
Knockdown of *Pf*FtsH1 disrupts apicoplast biogenesis
Using the endogenously tagged and regulated *Pf*FtsH1 strain, we determined
whether *Pf*FtsH1 is essential for apicoplast biogenesis. As expected, *Pf*FtsH1 expression
was down-regulated when the tetR-DOZI repressor bound the aptamer sequences, and
restored when anhydrotetracycline, which disrupts this interaction, was added (Fig. 3a,
Extended Data 5B)[30]. Upon downregulation of *Pf*FtsH1, we observed a nearly 4-fold
decrease in parasitemia after 3 replication cycles, compared to parasites expressing
normal levels of *Pf*FtsH1 (Fig 3b, Extended Data Fig. 6a). This growth defect observed
over multiple replication cycles reflected the slower kinetics of genetic regulation
(compared to chemical inhibition) and partial knockdown, which has been reported using
this regulation system (personal communication, Prof. Dan Goldberg). Significantly,
growth of *Pf*FtsH1 knockdown parasites was restored by addition of IPP (Fig 3b,
Extended Data 6a). Finally, we confirmed that loss of *Pf*FtsH1 led to apicoplast loss.
Using qPCR, we observed a steady decrease in the apicoplast:nuclear genome ratio upon
knockdown of *Pf*FtsH1 under IPP rescue compared to control parasites, consistent with
loss of the apicoplast[8] (Fig. 3c). These results demonstrate that *Pf*FtsH1 is required for
apicoplast biogenesis and is the first molecular player in apicoplast biogenesis identified
in an unbiased screen.
262
Knockdown of FtsH1 results in specific hypersensitivity to actinonin
Because knockdown of *Pf*FtsH1 phenocopies the apicoplast biogenesis defect of
actinonin, we sought to determine whether actinonin functionally inhibits *Pf*FtsH1.
Although we identified an actinonin-resistant mutation in *Tg*FtsH1 encoding an N805S
variant, this residue is not conserved between the *P. falciparum* and *T. gondii* homologs
and does not allow the same resistant mutation to be tested[35]. As an alternative, we down-
regulated *Pf*FtsH1 expression and observed a >50-fold decrease in the actinonin EC_{50}
compared to control parasites (Fig. 3d). To ensure that the decrease in *Pf*FtsH1 did not
cause confounding effects, growth inhibition was measured over a single replication
cycle during which *Pf*FtsH1 downregulation did not significantly affect parasite growth
(Fig. 3b, Extended Data Fig. 6b). Thus reduced levels of *Pf*FtsH1 caused hypersensitivity
to actinonin, analogous to drug-induced haploinsufficiency which is commonly used in
yeast to identify drug mechanism-of-action[41,42]. Importantly, *Pf*FtsH1 downregulation did

not alter the EC_{50} of fosmidomycin, which blocks apicoplast metabolism, or
chloramphenicol, which causes delayed biogenesis defects. These negative controls
indicate that the functional interaction of *Pf*FtsH1 with actinonin is specific and is not
observed with other inhibitors that disrupt apicoplast metabolism or biogenesis (Extended
Data Fig. 6c-d). This specific and strong chemical-genetic interaction indicates that
*Pf*FtsH1 is required for actinonin's mechanism-of-action.
282
Actinonin inhibits *Pf*FtsH1 activity *in vitro*
After our results revealed a specific and biologically relevant functional
interaction between actinonin and *Pf*FtsH1 in intact parasites, we sought to determine
whether actinonin directly inhibits *Pf*FtsH1 enzymatic activity *in vitro*. $Pf\text{FtsH1}_{91\text{-}612}$
lacking its transmembrane domain was expressed as a soluble fusion protein and purified
using tandem affinity tags (Figure 4a; Extended figure 7a). As expected, the purified
enzyme showed ATPase and protease activity (Extended Data Fig. 7a; Figure 4b-c).
Specific mutations reported to inactivate the ATPase (E249Q) or protease (D493A)
domains in other AAA+ proteases[43,44] abolished these respective activities in the *Pf*FtsH1
construct (Figure 4a-c)[43,44]. Notably, actinonin inhibited *Pf*FtsH1 protease activity with
an $IC_{50} \leq 0.6$ μM (Figure 4d) (sensitivity limitations of the assay precluded a more
accurate number). The metal-binding hydroxamate group of actinonin is known to be
very important for inhibition of peptide deformylase[25]. To determine the contribution of
the hydroxamate group to the inhibition of *Pf*FtsH1, we tested its inhibition by
actinamide, an analog in which the hydroxamate is replaced with an amide. Indeed,
actinamide was at least 10-fold less potent against *Pf*FtsH1 (IC_{50}=7.3 μM; Fig. 4e)[27]. The
decreased enzymatic inhibition correlated with a substantial decrease in parasite growth
inhibition compared to actinonin, although growth inhibition by actinamide was still
dependent on *Pf*FtsH1 levels (Extended Data Fig 7b, Fig. 4d). Hence, the direct inhibition
of *Pf*FtsH1 enzymatic activity by actinonin demonstrates that *Pf*FtsH1 is its target in *P.*
falciparum.
304
Discussion
We identified actinonin as a first-in-class antimalarial based on its distinct
apicoplast inhibition phenotype compared to known inhibitors. Because the phenotypic
screen and the mechanism-of-action elucidation were unbiased, the identification of
apicomplexan FtsH1 as actinonin's molecular target was surprising in a number of ways.
First, the actinonin target in bacteria and mammalian mitochondria is PDF, and *Pf*PDF
was presumed to be the target in *Plasmodium* parasites[25,27]. However, our functional
assays did not support this hypothesis. Identification of antimalarial drug targets based on
homology with known drug targets in distantly related organisms can be misleading. For
example, the misidentification of fatty acid enzyme *Pf*FabI as the target of triclosan,
based on its inhibition of bacterial homologs, led to failed drug development programs
when fatty acid biosynthesis was later shown to be dispensable in blood-stage
Plasmodium[5,6,45].
Second, though *P. falciparum* and *T. gondii* share conserved apicoplast biogenesis
pathways, it was uncertain whether *T. gondii*'s tractable culture system and genetics
could be leveraged to identify the actinonin target in *P. falciparum*. Fortunately,
actinonin caused apicoplast biogenesis defects in both *P. falciparum* and *T. gondii*

supporting a common mechanism-of-action. Specifically, actinonin-treated *P. falciparum*
formed "apicoplast-minus parasites" with IPP rescue while *T. gondii* formed them
spontaneously[8,46]. Importantly we show that, unlike in *T. gondii*, disruption of apicoplast
biogenesis does not necessarily lead to delayed death in *P. falciparum.* Based on our
experience, drug resistance selection in *T. gondii* to aid in elucidating antimalarial
mechanism-of-action, where resistant *P. falciparum* cannot be selected, is a powerful
approach. The evidence for FtsH1 as the common actinonin target in these two related
organisms greatly strengthens the target identification.
Third, we identified an actinonin-resistant mutation in *Tg*FtsH1 but, since *P.*
falciparum contains 3 FtsH homologs, it was not initially clear which of them was the
*Tg*FtsH1 ortholog. Almost all eukaryotes contain at least two FtsH homologs located in
the mitochondrial inner membrane: The i-AAA has a single transmembrane domain with
catalytic domains facing the intermembrane space, while the m-AAA has two
transmembrane domains and faces the matrix[47]. PF3D7_1119600 is the only homolog
predicted to have two transmembrane domains and phylogenetically clusters with human
and yeast m-AAA[37]. Of the two homologs with a single transmembrane domain,
PF3D7_1464900 is more closely related to mitochondrial i-AAA from human and yeast,
leaving *Pf*FtsH1 as the apparent ortholog of *Tg*FtsH1[37]. However this pairing
contradicted localization studies assigning *Tg*FtsH1 to the apicoplast and *Pf*FtsH1 to the
mitochondria[35,37]. *Tg*FtsH1 was shown to undergo *C*-terminal processing associated with
its trafficking to the apicoplast[40]. Similarly, we showed that *Pf*FtsH1 undergoes *C*-
terminal processing dependent on the presence of the apicoplast. This shared functional
phenotype strongly suggests that *Pf*FtsH1 also traffics to the apicoplast. Therefore, the
most parsimonious assignment for the three *Plasmodium* FtsH homologs is
PF3D7_1119600 and PF3D7_1464900 are mitochondrial and PfFtsH1 is in the
apicoplast. Unfortunately, we were unable to generate endogenously-tagged knockdown
strains of either PF3D7_1119600 or PF3D7_1464900 to perform localization and
functional studies (see Methods).
Finally, there are multiple potential metalloproteases in *P. falciparum* and *T.*
gondii that may be inhibited by actinonin. However, our evidence demonstrates that the
primary target of actinonin in these parasites, associated with its disruption of apicoplast
biogenesis, is FtsH1. Using IPP rescue to gauge apicoplast specificity, we assayed for
secondary targets in *P. falciparum* and detected a non-IPP rescuable target at an actinonin
concentration 20-fold higher than that for its apicoplast-specific target (Extended figure
1A). Similarly, in *T. gondii*, the loss of "delayed death" growth inhibition (and its
associated formation of apicoplast-minus parasites) signified actinonin's inhibition of a
secondary target at a concentration 3.5-fold greater than its specific inhibition of *Tg*FtsH1
(Extended figure 4b). Moreover, actinonin preferentially targets metalloproteases from
MEROPS clan MA composed of structurally homologous proteins with a conserved
metal-binding HEXXH motif and catalytic Glu[48]. Out of six metalloproteases predicted to
localize to the apicoplast, only FtsH and PDF are in clan MA[26,37,49–52]. Curiously, FtsH1
is the first FtsH homolog shown to be inhibited by actinonin or any small molecule.
Overall, we conclude that apicomplexan FtsH1 is the target of actinonin based on
1) an actinonin-resistant N805 variant in *Tg*FtsH1, 2) disruption of apicoplast biogenesis
upon *Pf*FtsH1 knockdown, phenocopying actinonin's biogenesis defect, 3) specific
actinonin-induced sensitization to *Pf*FtsH1 knockdown, and 4) *in vitro* inhibition of

*Pf*FtsH1 protease activity by actinonin. The unpredicted outcome underscores the power
of unbiased screens to uncover novel drug targets in unique but poorly characterized
cellular pathways. As the first screen for drug targets in essential apicoplast biogenesis
pathways, the approach highlighted in our study provides a framework for similar screens
in the future.
Our finding also presents an exciting opportunity for antimalarial drug discovery.
FtsH1 inhibitors will have significant advantages over existing antimalarials that target
apicoplast metabolism or gene expression. Whereas metabolic needs vary throughout the
parasite lifecycle and even between the same stage of different *Plasmodium* species[53,54],
apicoplast biogenesis is required at every proliferative stage of the parasite lifecycle and
is highly conserved among apicomplexan parasites. For example, apicoplast translation
inhibitors have broad clinical application as malaria prophylaxis targeting liver-stage
Plasmodium spp and as a partner drug, in combination with faster-acting compounds, for
acute parasitic diseases targeting blood-stage *Plasmodium* spp*, Toxoplasma gondii,* and
Babesia spp. In fact, the utility of these antibiotics as antiparasitics would be greater if
not for their slow activity. Inhibition of FtsH1 retains all the benefits of targeting
apicoplast biogenesis with no delay in the onset-of-action. Moreover, our inability to
select actinonin-resistant *Plasmodium* contrasts with the ready selection of *in vitro*
resistance against antibiotics and MEP inhibitors and indicates a lower likelihood of
clinical resistance to FtsH1 inhibitors[21,55–57]. Finally, several clinical lead candidates
based on the actinonin scaffold (e.g. GSK1322322, LBM-415, BB83698) have advanced
into human clinical trials as bacterial PDF inhibitors, setting precedence that actinonin
analogs can meet strong clinical entry criteria including selectivity for *Pf*FtsH1 over
human metalloproteases[58]. Taken together, FtsH1 inhibitors have potential for rapid
onset, multi-stage efficacy against multiple parasitic infections, and minimal clinical
resistance and routes to identifying a clinical lead candidate are readily accessible.
To pursue this goal, optimization of actinonin's antimalarial activity and drug
properties to identify a lead candidate suitable for testing in mouse malarial models will
be imperative. Both the *Pf*FtsH1 activity assay and *P. falciparum* FtsH1 knockdown
strain reported in our study may be adapted for high-throughput screening of actinonin
derivatives with increased potency and specificity. Structures of *Pf*FtsH1+/- actinonin
bound will aid in design of analogs. Importantly, optimization of *Pf*FtsH1 inhibitors will
benefit enormously from a "piggyback" strategy to access compound libraries,
counterscreens for off-target activity against human metalloproteases, and knowledge
about drug properties of actinonin-based scaffolds, which have already been established
for bacterial PDF inhibitor programs[58].
Finally, apicomplexan FtsH1 is the first novel molecular player in apicoplast
biogenesis identified in an unbiased screen. Acquired by secondary endosymbiosis of an
alga, the apicoplast is evolutionarily distinct. FtsH1's role in organelle biogenesis is not
conserved in homologs found in mitochondria or primary chloroplasts and likely
represents a novel pathway unique to secondary endosymbionts in this parasite
lineage[17,32,59,60]. FtsH homologs have broad substrate specificity to perform general
degradation of misfolded membrane proteins[47]. However they have also been shown to
catalyze the proteolysis of native cytosolic proteins under specific conditions critical for
cellular regulation[61]. For example, the essential function of *E. coli* FtsH is its regulated
proteolysis of LpxC, the key enzyme in lipopolysaccharide biosynthesis[62]. In *Bacillus*

subtilis, FtsH-dependent degradation of SpoIVFA regulates spore differentiation[63].
Similarly, *Ec*FtsH degrades a CII transcription factor required for the lysis-lysogeny
decision by phage lambda[64]. Based on these regulatory functions of bacterial FtsH
homologs, we propose that apicomplexan FtsH1 regulates the proteolysis of key
apicoplast membrane protein(s) during parasite replication. Identification of these
substrates will be critical to understanding its molecular mechanism in apicoplast
biogenesis. FtsH1 offers a rare foothold into a novel apicoplast biogenesis pathway
evolved from secondary endosymbiosis and will yield deeper insight into the molecular
mechanisms of eukaryogenesis.
423
424 **References**
425

1. McFadden, G. I., Reith, M. E., Munholland, J. & Lang-Unnasch, N. Plastid in human

parasites. *Nature* **381,** 482 (1996).

2. Köhler, S. *et al.* A plastid of probable green algal origin in Apicomplexan

parasites. *Science* **275,** 1485–1489 (1997).

3. Jomaa, H. *et al.* Inhibitors of the nonmevalonate pathway of isoprenoid

biosynthesis as antimalarial drugs. *Science* **285,** 1573–1576 (1999).

4. Mazumdar, J., H Wilson, E., Masek, K., A Hunter, C. & Striepen, B. Apicoplast fatty

acid synthesis is essential for organelle biogenesis and parasite survival in

Toxoplasma gondii. *Proc. Natl. Acad. Sci. U. S. A.* **103,** 13192–13197 (2006).

5. Vaughan, A. M. *et al.* Type II fatty acid synthesis is essential only for malaria

parasite late liver stage development. *Cell. Microbiol.* **11,** 506–520 (2009).

6. Yu, M. *et al.* The Fatty Acid Biosynthesis Enzyme FabI Plays a Key Role In the

Development of Liver Stage Malarial Parasites. *Cell Host Microbe* **4,** 567–578

(2008).

7. Nair, S. C. *et al.* Apicoplast isoprenoid precursor synthesis and the molecular

basis of fosmidomycin resistance in Toxoplasma gondii. *J. Exp. Med.* (2011).

doi:10.1084/jem.20110039

8.   Yeh, E. & DeRisi, J. L. Chemical Rescue of Malaria Parasites Lacking an Apicoplast

Defines Organelle Function in Blood-Stage Plasmodium falciparum. *PLoS Biol* **9,**

e1001138 (2011).

9.   Ke, H. *et al.* The heme biosynthesis pathway is essential for Plasmodium

falciparum development in mosquito stage but not in blood stages. *J. Biol. Chem.*

289, 34827–34837 (2014).

10. Fichera, M. E. & Roos, D. S. A plastid organelle as a drug target in apicomplexan

parasites. *Nature* **390,** 407–409 (1997).

11. Dahl, E. L. *et al.* Tetracyclines specifically target the apicoplast of the malaria

parasite Plasmodium falciparum. *Antimicrob. Agents Chemother.* **50,** 3124–3131

(2006).

12. Geary, T. G. & Jensen, J. B. Effects of antibiotics on Plasmodium falciparum in

vitro. *Am. J. Trop. Med. Hyg.* **32,** 221–225 (1983).

13. Oyakhirome, S. *et al.* Randomized controlled trial of fosmidomycin-clindamycin

versus sulfadoxine-pyrimethamine in the treatment of Plasmodium falciparum

malaria. *Antimicrob. Agents Chemother.* **51,** 1869–1871 (2007).

14. Lanaspa, M. *et al.* Inadequate Efficacy of a New Formulation of Fosmidomycin-

Clindamycin Combination in Mozambican Children Less than Three Years Old

with Uncomplicated Plasmodium falciparum Malaria. *Antimicrob. Agents*

Chemother. **56,** 2923–2928 (2012).

15. Guggisberg, A. M. *et al.* Whole-Genome Sequencing to Evaluate the Resistance

Landscape Following Antimalarial Treatment Failure With Fosmidomycin-

Clindamycin. *J. Infect. Dis.* **214,** 1085–1091 (2016).

16. Waller, R. F. *et al.* Nuclear-encoded proteins target to the plastid in Toxoplasma

gondii and Plasmodium falciparum. *Proc. Natl. Acad. Sci.* **95,** 12352–12357

(1998).

17. Vaishnava, S. & Striepen, B. The cell biology of secondary endosymbiosis--how

parasites build, divide and segregate the apicoplast. *Mol. Microbiol.* **61,** 1380–

1387 (2006).

18. Spangenberg, T. *et al.* The Open Access Malaria Box: A Drug Discovery Catalyst

for Neglected Diseases. *PLoS ONE* **8,** e62906 (2013).

19. Wiesner, J., Sanderbrand, S., Altincicek, B., Beck, E. & Jomaa, H. Seeking new

targets for antiparasitic agents. *Trends Parasitol.* **17,** 7–8 (2001).

20. Dahl, E. L. & Rosenthal, P. J. Multiple Antibiotics Exert Delayed Effects against the

Plasmodium falciparum Apicoplast. *Antimicrob Agents Chemother* **51,** 3485–

3490 (2007).

21. Wu, W. *et al.* A chemical rescue screen identifies a Plasmodium falciparum

apicoplast inhibitor targeting MEP isoprenoid precursor biosynthesis.

Antimicrob. Agents Chemother. AAC.03342-14 (2014). doi:10.1128/AAC.03342-

14

22. Gisselberg, J. E., Herrera, Z., Orchard, L., Llinas, M. & Yeh, E. A Specific Non-

Bisphosphonate Inhibitor Of The Bifunctional Farnesyl/Geranylgeranyl

Diphosphate Synthase In Malaria Parasites. *bioRxiv* 134338 (2017).

doi:10.1101/134338

23. Gordon, J. J., Kelly, B. K. & Miller, G. A. Actinonin: an antibiotic substance

produced by an actinomycete. *Nature* **195,** 701–702 (1962).

24. Goodman, C. D. & McFadden, G. I. Ycf93 (Orf105), a small apicoplast-encoded

membrane protein in the relict plastid of the malaria parasite Plasmodium

falciparum that is conserved in Apicomplexa. *PloS One* **9,** e91178 (2014).

25. Chen, D. Z. *et al.* Actinonin, a naturally occurring antibacterial agent, is a potent

deformylase inhibitor. *Biochemistry (Mosc.)* **39,** 1256–1262 (2000).

26. Bracchi-Ricard, V. *et al.* Characterization of an eukaryotic peptide deformylase

from Plasmodium falciparum. *Arch. Biochem. Biophys.* **396,** 162–170 (2001).

27. Lee, M. D. *et al.* Human mitochondrial peptide deformylase, a new anticancer

target of actinonin-based antibiotics. *J. Clin. Invest.* **114,** 1107–1116 (2004).

28. Kumar, A. *et al.* Crystals of peptide deformylase from Plasmodium falciparum

reveal critical characteristics of the active site for drug design. *Struct. Lond. Engl.*

1993 **10,** 357–367 (2002).

29. Margolis, P. S. *et al.* Peptide Deformylase in Staphylococcus aureus: Resistance to

Inhibition Is Mediated by Mutations in the Formyltransferase Gene. *Antimicrob.*

Agents Chemother. **44,** 1825–1831 (2000).

30. Ganesan, S. M., Falla, A., Goldfless, S. J., Nasamu, A. S. & Niles, J. C. Synthetic RNA-

protein modules integrated with native translation mechanisms to control gene

expression in malaria parasites. *Nat. Commun.* **7,** 10727 (2016).

31. Richter, U. *et al.* A Mitochondrial Ribosomal and RNA Decay Pathway Blocks Cell

Proliferation. *Curr. Biol.* **23,** 535–541 (2013).

32. van Dooren, G. G. *et al.* A novel dynamin-related protein has been recruited for

apicoplast fission in Toxoplasma gondii. *Curr. Biol. CB* **19,** 267–276 (2009).

33. Jacot, D., Daher, W. & Soldati-Favre, D. Toxoplasma gondii myosin F, an essential

motor for centrosomes positioning and apicoplast inheritance. *EMBO J.* **32,**

1702–1716 (2013).

34. Lévêque, M. F. *et al.* Autophagy-Related Protein ATG8 Has a Noncanonical

Function for Apicoplast Inheritance in Toxoplasma gondii. *mBio* **6,** e01446-1415

(2015).

35. Karnataki, A., Derocher, A. E., Coppens, I., Feagin, J. E. & Parsons, M. A membrane

protease is targeted to the relict plastid of toxoplasma via an internal signal

sequence. *Traffic Cph. Den.* **8,** 1543–1553 (2007).

36. Ganji, R. J. *et al.* Structural basis for the inhibition of M1 family aminopeptidases

by the natural product actinonin: Crystal structure in complex with E. coli

aminopeptidase N. *Protein Sci. Publ. Protein Soc.* **24,** 823–831 (2015).

37. Tanveer, A. *et al.* An FtsH protease is recruited to the mitochondrion of

Plasmodium falciparum. *PloS One* **8,** e74408 (2013).

38. van Dooren, G. G. *et al.* Development of the endoplasmic reticulum,

mitochondrion and apicoplast during the asexual life cycle of Plasmodium

falciparum. *Mol. Microbiol.* **57,** 405–419 (2005).

39. Stanway, R. R. *et al.* Organelle segregation into Plasmodium liver stage

merozoites. *Cell. Microbiol.* **13,** 1768–1782 (2011).

40. Sequential processing of the Toxoplasma apicoplast membrane protein FtsH1 in

topologically distinct domains during intracellular trafficking. Available at:

http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2817949/. (Accessed: 22nd

June 2016)

41. Giaever, G. *et al.* Genomic profiling of drug sensitivities via induced

haploinsufficiency. *Nat. Genet.* **21,** 278–283 (1999).

42. Baetz, K. *et al.* Yeast genome-wide drug-induced haploinsufficiency screen to

determine drug mode of action. *Proc. Natl. Acad. Sci. U. S. A.* **101,** 4525–4530

(2004).

43. Hersch, G. L., Burton, R. E., Bolon, D. N., Baker, T. A. & Sauer, R. T. Asymmetric

interactions of ATP with the AAA+ ClpX6 unfoldase: allosteric control of a

protein machine. *Cell* **121,** 1017–1027 (2005).

44. Bieniossek, C. *et al.* The molecular architecture of the metalloprotease FtsH.

Proc. Natl. Acad. Sci. U. S. A. **103,** 3066–3071 (2006).

45. Surolia, N. & Surolia, A. Triclosan offers protection against blood stages of

malaria by inhibiting enoyl-ACP reductase of Plasmodium falciparum. *Nat. Med.*

7, 167–173 (2001).

46. He, C. Y. *et al.* A plastid segregation defect in the protozoan parasite Toxoplasma

gondii. *EMBO J* **20,** 330–339 (2001).

47. Janska, H., Kwasniak, M. & Szczepanowska, J. Protein quality control in

organelles — AAA/FtsH story. *Biochim. Biophys. Acta BBA - Mol. Cell Res.* **1833,**

381–387 (2013).

48. MEROPS - the Peptidase Database. Available at: http://merops.sanger.ac.uk/cgi-

bin/smi_summary?mid=J16.404. (Accessed: 10th May 2017)

49. van Dooren, G. G., Su, V., D'Ombrain, M. C. & McFadden, G. I. Processing of an

Apicoplast Leader Sequence inPlasmodium falciparum and the Identification of a

Putative Leader Cleavage Enzyme. *J. Biol. Chem.* **277,** 23612–23619 (2002).

50. Mallari, J. P., Oksman, A., Vaupel, B. & Goldberg, D. E. Kinase-associated

endopeptidase 1 (Kae1) participates in an atypical ribosome-associated complex

in the apicoplast of Plasmodium falciparum. *J. Biol. Chem.* **289,** 30025–30039

(2014).

51. Ponpuak, M. *et al.* A role for falcilysin in transit peptide degradation in the

Plasmodium falciparum apicoplast. *Mol. Microbiol.* **63,** 314–334 (2007).

52. Chen, X. *et al.* Inhibitors of Plasmodium falciparum methionine aminopeptidase

1b possess antimalarial activity. *Proc. Natl. Acad. Sci. U. S. A.* **103,** 14548–14553

(2006).

53. Srivastava, A. *et al.* Stage-Specific Changes in Plasmodium Metabolism Required

for Differentiation and Adaptation to Different Host and Vector Environments.

PLoS Pathog. **12,** e1006094 (2016).

54. Shears, M. J., Botté, C. Y. & McFadden, G. I. Fatty acid metabolism in the

Plasmodium apicoplast: Drugs, doubts and knockouts. *Mol. Biochem. Parasitol.*

199, 34–50 (2015).

55. Dharia, N. V. *et al.* Use of high-density tiling microarrays to identify mutations

globally and elucidate mechanisms of drug resistance in Plasmodium falciparum.

Genome Biol. **10,** R21 (2009).

56. Guggisberg, A. M. *et al.* A sugar phosphatase regulates the methylerythritol

phosphate (MEP) pathway in malaria parasites. *Nat. Commun.* **5,** 4467 (2014).

57. Sidhu, A. B. S. *et al.* In vitro efficacy, resistance selection, and structural modeling

studies implicate the malarial parasite apicoplast as the target of azithromycin. *J.*

Biol. Chem. **282,** 2494–2504 (2007).

58. Sangshetti, J. N., Khan, F. A. K. & Shinde, D. B. Peptide deformylase: a new target

in antibacterial, antimalarial and anticancer drug discovery. *Curr. Med. Chem.* **22,**

214–236 (2015).

59. Moore, R. B. *et al.* A photosynthetic alveolate closely related to apicomplexan

parasites. *Nature* **451,** 959–963 (2008).

60. Spork, S. *et al.* An Unusual ERAD-Like Complex Is Targeted to the Apicoplast of

Plasmodium falciparum. *Eukaryot. Cell* **8,** 1134–1145 (2009).

61. Bittner, L.-M., Arends, J. & Narberhaus, F. When, how and why? Regulated

proteolysis by the essential FtsH protease in Escherichia coli. *Biol. Chem.* **398,**

625–635 (2017).

62. Ogura, T. *et al.* Balanced biosynthesis of major membrane components through

regulated degradation of the committed enzyme of lipid A biosynthesis by the

AAA protease FtsH (HflB) in Escherichia coli. *Mol. Microbiol.* **31,** 833–844

(1999).

63. Rudner, D. Z. & Losick, R. A sporulation membrane protein tethers the pro-σK

processing enzyme to its inhibitor and dictates its subcellular localization. *Genes*

Dev. **16,** 1007–1018 (2002).

64. Shotland, Y. *et al.* Proteolysis of the phage lambda CII regulatory protein by FtsH

(HflB) of Escherichia coli. *Mol. Microbiol.* **24,** 1303–1310 (1997).

599
Methods
Chemicals
Fosmidomycin was purchased from Santa Cruz Biotechnology and 10mM aliquots were
prepared in water. Chloramphenicol was purchased from Sigma Aldrich and 50mM
aliquots were prepared in 100% ethanol. Actinonin was purchased from Sigma Aldrich
and 25mM aliquots were prepared in 100% ethanol. Anhydrotetracycline was purchased
from Sigma and 2.5mM aliquots prepared in 100% ethanol and used at a final

concentration of 0.5uM. Actinamide was a gift from Drs. David Scheinberg and Ouathek
Ouerfelli and 25mM aliquots were prepared in 100% ethanol.
609
610 Enoxacin, ciprofloxacin, levofloxacin, norfloxacin, novobiocin, coumeramycin,
611 mericitabine, 2'deoxy-2-F-cytidine, gemcitabine, ADEP1a, beta-lactone 4, beta-lactone
612 7, and rifampin were acquired and solubilized as noted in Supplementary Table 1.
613
614 Isopentenyl pyrophosphate (IPP) was purchased from Isoprenoids LC and stored at 2
615 mg/mL in 70% methanol, 30% 10mM ammonium hydroxide at -80C. To prevent
616 methanol toxicity, aliquots of IPP were dried in the speed vacuum centrifuge before
617 adding to cultures. All drugs were stored at -20C and resuspended just prior to use.
618
***Plasmodium falciparum* culture and transfections**
P. falciparum D10 (MRA-201), and D10 ACP$_L$-GFP (MRA-568) were obtained from
MR4. *P. falciparum* NF54[attB] was a gift from David Fidock (Columbia University). NF54
[attB] strain constitutively expressing Cas9 and T7 Polymerase, generated previously[63], was
used in this study. Parasites were maintained in human erythrocytes (2% hematocrit) in
RPMI 1640 media supplemented with 0.25% Albumax II (GIBCO Life Technologies), 2
g/L sodium bicarbonate, 0.1 mM hypoxanthine, 25 mM HEPES (pH 7.4), 50 µg/L
gentamycin, and 0.4% glucose at 37°C, 5% O_2, and 5% CO_2. All cell lines tested
negative for mycoplasma contamination during routine checks.
628
Parasites were transfected using methods already published[30]. Briefly, we used 50 ug of
plasmid per 200 uL packed red blood cells (RBCs) adjusted to 50% hematocrit. We used
a Bio-Rad Gene Pulser II to preload uninfected RBCs using eight square-wave pulses of
365 V for 1 ms, separated by 100 ms. Preloaded RBCs were resealed for 1 hour at 37C
and washed twice in RPMI to remove lysed cells. Schizont stage parasites at 0.5%
parasitemia were then allowed to invade half of the preloaded RBCs during two
sequential reinvasions. Media was changed daily for the first 12 days and every other day
thereafter. Parasites were split 1:1 into fresh blood every 4 days until parasites were
visible by Giemsa smear. To select for integration of the pFtsH1 into *P. falciparum*
NF54[attB-pPfCRISPR] parasites, transfected parasites were maintained in media containing 5
nM WR99210 and 0.5 uM anhydrotetracycline (Sigma) and then selected with 2.5 mg/l
Blasticidin S (Sigma) beginning 4 days after transfection.
641
***Toxoplasma gondii* culture and transfection**
T. gondii RH and *T. gondii* RH Δ*ku80*Δ*hxgprt* strains were a gift from Matthew Bogyo
(Stanford University) and maintained by passage through confluent monolayers of human
foreskin fibroblasts (HFFs) host cells. HFFs were cultured in DMEM (Invitrogen)
supplemented with 10% FBS (Fetal Plex Animal Serum from Gemini), 2mM L-
glutamine (Gemini), and 100 ug penicillin and 100 ug streptomycin per mL (Gibco Life
Technologies), maintained at 37 C and 5% CO_2. Parasites were harvested for assays by
syringe lysis of infected HFF monolayers.
650
For transfection of *T. gondii* Δ*ku80*Δ*hxgprt,* 15ug of the pTgCRISPR plasmid[65] was
combined with 3 ug of the pFtsH1N805S or pFtsH1WT that had been linearized by NotI

digestion. Approximately 10^7 parasites were released from host cells using syringe lysis
and washed into 400uL of cytomix containing both plasmids. Parasites were
eletroporated (BTX Electro Cell Manipulator 600) with 1.2-1.4 kV, 2.5kV/resistance, R2
(24 ohm) and then allowed to recover in the cuvette at room temperature for 10 mins
before adding to host cells. After 24 hours, media containing 25ug/mL mycophenolic
acid (Sigma) and 50 ug/mL xanthine (Sigma) was added to select for transfectants. After
a week, plaques were observed and single clones were isolated using limiting dilution.
660
661   ***Toxoplasma gondii* genome sequencing and SNP identification**
662   Actinonin resistant and susceptible *T. gondii* were grown on 15 cm dishes containing
663   confluent HFF monolayers until spontaneous lysis of the monolayer was achieved.
664   Released parasites were collected and filtered through 5 micrometer syringe filters
665   (Millipore) before isolating DNA (Qiagen DNAeasy Blood & Tissue).
666
T. gondii genomic DNA isolated from either the parental *T. gondii* RH strain
(SRR3666219) or any of its derived mutants (SRR3666219, SRR3666222, SRR3666224,
SRR3666792, SRR3666794, SRR3666796, SRR3666798, SRR3666799, SRR3666801)
was sequenced in an Illumina NextSeq apparatus using 2x150bp reads at an average
sequencing depth of 35x. Sequencing reads were quality trimmed and remnants of
sequencing adaptors removed with *trimmomatic* (PMID:24695404). Next, reads were
mapped to the reference nuclear assembly of the *T. gondii* GT1 strain (ToxoDB v13.0)
and the apicoplast genome assembly from the RH strain (NC_001799) with the program
bowtie2 (PMID:25621011). Duplicated aligned reads were removed with *picard tools*
(**http://broadinstitute.github.io/picard**) and reads spanning InDels were realigned with
GATK (PMID:20644199). Afterwards, allelic variants were called with *samtools mpileup*
(PMID:19505943) followed by *bcftools call* with –p set to 0.05 (PMID:26826718).
Finally, classification of mutations was performed with *snpEff* (PMID:22728672).
680
Growth inhibition assays
For *P. falciparum* EC_{50} calculations, growth assays were performed in 96 well plates
containing serial dilution of drugs in triplicate. Media was supplemented with 200 uM
IPP as indicated. Growth was initiated with ring-stage parasites (synchronized with 5%
sorbitol treatment) at 1% parasitemia and 1% hematocrit. To calculate growth, cultures
were incubated for 72h and growth was then terminated by incubation with 1%
formaldehyde (Electron Microscopy Services) for 30 minutes at room temperature.
Parasitized cells were stained with 50 nM YOYO-1 (Invitrogen) overnight at room
temperature and the parasitemia was determined by flow cytometry (BD Accuri C6
Sampler). Data were analyzed by BD Accuri C6 Sampler software. *P. falciparum* growth
assays measuring the actinonin EC_{50} were repeated in the laboratory > 10 times for WT
parasites and >3 times upon knockdown of FtsH1.
693
For *T. gondii* EC_{50} calculations, plaque assays were performed in 24 well plates
containing confluent HFF monolayers serial dilutions of drugs in duplicate.
Approximately 50 parasites were counted using flow cytometry and added to each well.
After incubating for 6 days, infected monolayers were washed, fixed with methanol for
10 minutes, stained with 2% crystal violet (Sigma) for 30 minutes, and then washed

again. Plaques were visualized as non-stained areas. The area of each plaque in a given
well was measured and summed using ImageJ as a proxy for growth and normalized to
the vehicle only control. *T. gondii* plaque assays measuring the actinonin EC_{50} were
repeated in the laboratory > 2 times for resistant parasites that arose from actinonin
selection, > 10 times for WT parasites, and > 3 times for TgFtsH(N805S).
704

For measuring the growth inhibition of *P. falciparum* during the time course, 10 uM
actinonin, 10 uM fosmidomycin, 30 uM chloramphenicol, 200 uM IPP, and 0.5 uM of
anhydrotetracycline was used as necessary. For comparison of growth between different
treatment conditions, cultures were carried simultaneously and handled identically with
respect to media changes and addition of blood cells. Daily samples were collected and
fixed with 1% formaldehyde for 30 minutes at RT. At the end of the time course, all
samples were stained with 50 nM YOYO-1 and parasitemia was measured using flow
cytometry. All growth curves were plotted using GraphPad Prism. *P. falciparum* time
course experiments were repeated in the laboratory > 2 times for WT parasites and > 2
times upon knockdown of FtsH1.
715

For measuring the growth inhibition of *T. gondii* during the time course, 6-well plates
were set up with no drug, 40 uM actinonin, 25 nM clindamycin, and 4 uM
pyrimethamine. *T. gondii* was added at a MOI = 3. Every 12 hours, parasites were
released from HFFs using syringe lysis and counted using flow cytometry (BD Accuri C6
Sampler). After 36 hours, spontaneous lysis of the monolayer was observed and parasites
were counted using flow cytometry and then added back to fresh monolayers at MOI = 3
in the absence of drug and parasites were counted every 12 hours as before. *T. gondii*
time course experiments were repeated in the laboratory > 2 times.
724

725   For measuring growth inhibition using tandem-tomato expression as a proxy for growth,
726   tandem-tomato *T. gondii* were seeded onto a black, clear bottom 96 well plate (Costar
727   3603) at 2000 parasites per well. Parasites were treated at a range of drug concentrations
728   and fluorescence was measured daily in a plate reader (BioTek, Synergy) for 5 days using
729   the bottom read function.
730

For co-treatments of *P. falciparum* with actinonin and chloramphenicol, 96 well plates
containing ring stage parasites at 1% parasitemia and 1% hematocrit were treated with
serial dilutions of both actinonin and chloramphenicol alone and in combination. To
determine the effect on growth after one lytic cycle, parasites were fixed at 72h and
parasitemia was measured by flow cytometry as above. To determine the effect on
growth after two lytic cycles, 75% of the media was exchanged at 72 hours and plates
were incubated for an additional 48 hours following fixation and flow cytometry as
above. Media was supplemented with 200 uM IPP as a separate control to insure
specificity of the drug at the concentrations used. Co-treatments with actinonin and
chloramphenicol were performed a single time in the laboratory using three technical
replicates.
742

Quantitative Real-Time PCR

Parasites from 1 mL of *P. falciparum* culture at ring stage were isolated by saponin lysis
followed by two washes with PBS. DNA was purified using DNAeasy Blood & Tissue
(Qiagen). Primers were designed to target genes found on the apicoplast or nuclear
genome: tufA (apicoplast) 5'-GATATTGATTCAGCTCCAGAAGAAA-3' and CHT1
(nuclear) 5'-TGTTTCCTTCAACCCCTTTT-3' /5'-TGTTTCCTTCAACCCCTTTT-3'.
Reactions contained template DNA, 0.15 uM of each primer, and 1x SYBR Green I
Master mix (Roche). qPCR reactions were performed at 56C primer annealing and 65C
template extension for 35 cycles on a Applied Biosystem 7900HT system. Relative
quantification of target genes was determined[64]. For each time point, the
apicoplast:nuclear genome ratio was calculated relative to the appropriate control
collected at the same time. The apicoplast:nuclear genome ratio was measured by qPCR
> 5 times for WT parasites treated with actinonin and > 2 times upon knockdown of
FtsH1.
757
Fluorescence Microscopy
P. falciparum D10 ACP(L)-GFP parasites diluted to 0.05% hematocrit were settled on a
Lab-Tek II Chambered Coverglass (Thermo Fisher) and incubated in 2 ug/mL Hoescht
33342 stain for 15 minutes at 37C. Widefield epifluorescence live cell images were
acquired with an Olympus IX70 microscope. The microscope was outfitted with a
Deltavision Core system (Applied Precision) using an Olympus x60 1.4NA Plan Apo
Lens, a Sedat Quad filter set (Semrock) and a CoolSnap HQ CCD Camera
(Photometrics). The microscope was controlled and images were deconvolved via
softWoRx 4.1.0 software. ImageJ software was used to analyze resulting images.
767
Live microscopy of *T. gondii* RH FNR-RFP parasites[68] was performing using Lab-Tek II
Chambered Coverglasses containing confluent HFF monolayers. Parasites were added at
an MOI = 1. After 36 hours of incubation, parasites were incubated with 2 ug/mL of
Hoescht for 15 minutes. Widefield epifluorescence live cell images were acquired with a
Nikon Eclipse Ti inverted fluorescence microscope with a NA 1.40 oil-immersion
objective (Nikon Instruments) and controlled using MicroManager v1.4. An iXon3 888
EMCCD camera (Andor) was used for fluorescence imaging and an a Zyla 5.5 sCMOS
camera (Andor) was used for phase contrast imaging. ImageJ software was used to
analyze the resulting images. ACP-GFP parasites treated with actinonin were imaged > 4
different times in the laboratory.
778
Immunoblot
Parasites from 9 mL of *P. falciparum* culture were isolated by saponin lysis, washed with
PBS and resuspended in 1 x NuPAGE LDS sample buffer (Invitrogen). Proteins were
separated by electrophoresis on 4-12% Bis-Tris gel (Invitrogen) and transferred to a
nitrocellulose membrane. After blocking, membranes were probed with 1:2000
monoclonal mouse anti-FLAG M2 (Sigma) and 1:10,000 IRDye 680RD goat anti-mouse
IgG (LiCor Bioscience) for anti-FtsH1 immunoblots. For anti-PDF immunoblots,
membranes were probed with 1:2000 rabbit monoclonal anti-MYC (Cell Signaling
Technology 2278S), followed by 1:20,000 rabbit polyclonal anti-PfAldolase (Abcam
ab207494) and 1:10,000 donkey anti-rabbit 800 (LiCor Biosciences). Fluorescence
antibody-bound proteins were detected with Odyssey Imager (LiCor Biosciences). When

antibodies of the same species were used, membranes were probed and imaged
sequentially. Immunoblots of FtsH-FLAG and PFD-myc were repeated in the laboratory
> 2 times.
793
***Toxoplasma gondii* resistance selection**
Approximately 2 x 10^6 *T. gondii* RH parasites were added to T25s containing a
confluent HFF monolayer and allowed to grow for 24 hours. To mutagenize, between
500 uM – 2 mM N-ethyl N-nitrosourea (ENU) diluted in DMSO was added to flasks and
incubated for 2 hours at 37 C. Cultures were then washed twice with 10mLs of cold PBS
and then released from host cells using syringe lysis. A quarter of the resulting parasites
were passaged to T25s containing a fresh monolayer of HFFs. After two passages,
parasites were treated with 40uM actinonin (the minimum inhibitory concentration of
sensitive *T. gondii*). After one passage under actinonin selection, a severe bottleneck was
observed. Plaques of resistant parasites could be observed after one week of constant
actinonin pressure with periodic media changes. Finally, single clones were isolated
using limiting dilution. Actinonin resistance selection was repeated in the laboratory
twice.
807
***Plasmodium falciparum* construct generation**
The primers used for generating different fragments are listed in the Extended Data Table
3. A construct for inducing PfPDF (PF3D7_0907900) expression in *P. falciparum* was
generated from the parental pMG96 plasmid. PDF gene was amplified from pUC57-Amp
plasmid with the codon optimized PDF gene (PF3D7_0907900) using primers SMG374
and SMG375. The amplicon was cloned in PfCAM base plasmid (pMG96), which
contains single aptamer at 5'UTR and 10x aptamer at 3'UTR. The restriction sites MScI
and BstEII were used for cloning thus encoding PDF containing c-myc tag at the c-
terminal end. The transfection was carried out as discussed previously[69]. A construct for
regulating expression of FtsH1 (PF3D7_1239700) in *P. falciparum* (pFtsH1) was
generated from the parental pSN054, a modified pJazz linear plasmid (Extended Data
Table 3). The left homology region was amplified from parasite genomic DNA using
primers SMG476 and SMG 477 and was cloned using FseI and AsisI restriction sites.
FtsH1 protein coding nucleotides 2348- 2643 were recoded using gene block (IDT) to
remove the PAM site. The right homology region was amplified from parasite genomic
DNA using primers SMG501 and SMG502. These fragments were cloned using the I-
SceI restriction site. Targeting guide RNA was generated by klenow reaction using
primers SMG514 and SMG515 and was inserted using the AflII site. All sequences were
ligated into the parent plasmid using Gibson assembly. Constructs for regulating
PF3D7_1464900 were generated similarly (SMG505 and SMG506 for the right
homology region; SMG495 and SMG496 for the left homology region; SMG518 and
SMG519 for the gRNA klenow reaction). While this construct could be generated, no
parasites emerged after two transfections. Constructs for regulating PF3D7_1119600
using primers SMG503 and SMG504 for the right homology region and SMG481 and
SMG507 for the left homology region were unable to be generated because of
unsuccessful PCR.
834
***Toxoplasma gondii* construct generation**

A construct for knocking in the FtsH1$_{N805S}$ allele into the endogenous loci of FtsH1
(pFtsHN805S) was generated from the parental pTKO2 vector. Briefly, a ~800 bp
sequence upstream of the TGGT1_259260 start codon was amplified as the left
homology region (using primers KAJ1 and KAJ2). The FtsH$_{N805S}$ sequence was then
amplified from actinonin resistant cDNA (using primers KAJ3 and KAJ4). The HXGPRT
resistance cassette was amplified off of the pTKO2 cassette (using primers KAJ5 and
KAJ6). A ~800 bp sequence downstream of the TGGT1_259260 stop codon was
amplified as the right homology region (using primers KAJ7 and KAJ8). To insert the
right homology region, the pTKO2 plasmid was cut with HindIII and HpaI and the 800
bp sequence was inserted using infusion (Clontech). To insert the left homology region
and the FtsH1$_{N805S}$ allele, the pTKO2 plasmid containing the downstream homology
region was cut with NotI and EcoRI and the two PCR products were inserted also using
infusion (Clontech). The resulting colonies were tested using a diagnostic HpaI digest
and correct clones were subjected to Sanger sequencing of the inserts. To revert the
pFtsHN805S construct to pFtsHWT, we used Q5 mutagenesis (NEB) and primers KAJ9
and KAJ10.
852
853   To increase the transfection efficiency and specificity, CRISPR-Cas9 was used to insert a
854   double stranded break at the site of insertion (the endogenous TGGT1_259260 allele).
855   Briefly, pSAG1::Cas9-U6::sgUPRT[65] was modified to contain a guide sequence specific
856   to a FtsH1 intron using the Q5 mutagenesis kit (NEB) and primers KAJ11 and KAJ12.
857   Sanger sequencing of the guide was used to verify the resulting plasmid.
858
Recombinant *Pf*FtsH1 expression and purification
His$_6$-SUMO-*Pf*FtsH1$_{91-612}$-GST was cloned into a pET22-based vector by Gibson
assembly[71]. Point mutations in this gene were constructed by site-directed mutagenesis.
Liquid cultures of T7 Express cells (NEB) harboring these plasmids were grown to log
phase, cooled to 20 °C, and induced with 0.5 mM IPTG plus 0.1% benzyl alcohol
(Sigma). Cultures were incubated for an additional 3 h at 20 °C before harvesting and
freezing. Cell pellets were resuspended in lysis buffer (50 mM HEPES, pH 7.5, 200 mM
NaCl, 10% glycerol, 2 mM β-mercaptoethanol [βME], 20 mM imidazole), mixed with
lysozyme (1 mg/mL) and Benzonase (Sigma), and lysed by sonication. Cleared lysates
were incubated with Ni-NTA resin (G-Biosciences) for 1 h at 4 °C. After extensive
washing with lysis buffer, bound proteins were eluted from the resin with lysis buffer
plus 300 mM imidazole. The eluent was applied to a GSTrap column (GE Healthcare)
and washed with storage buffer (50 mM HEPES, pH 7.5, 200 mM NaCl, 10% glycerol, 2
mM βME) before eluting with storage buffer plus 10 mM reduced glutathione. The eluent
was dialyzed overnight against storage buffer, concentrated, snap-frozen, and stored in
aliquots at –80 °C.
875
PfFtsH1 enzymatic assays
Rates of ATP hydrolysis by FtsH were measured using a coupled spectrophotometric
assay[72] in PD buffer (25 mM HEPES, pH 7.5, 200 mM NaCl, 5 mM MgSO4, 10 µM
ZnSO4, 10% glycerol) with 3% dimethyl sulfoxide (DMSO) at 37 °C. Protein
degradation rates were measured by incubating FtsH (1 µM) with FITC-labeled casein (4
µM, Sigma, Type III) in PD buffer plus 3% DMSO. Reactions were started by adding

ATP (4 mM) with a regeneration system (16 mM creatine phosphate and 75 µg/mL
creatine kinase), and degradation was followed by measuring the fluorescence intensity
(excitation 485 nm; emission 528 nm) at 37 °C. For activity-inhibition assays, reactions
were pre-incubated with inhibitor in DMSO for 10 min before adding ATP. Addition of
inhibitor provided the 3% DMSO found in the other assays. For determining specific
activities, the total concentration of casein was 10 µM (2 µM of FITC-labeled casein plus
8 µM unlabeled casein).
889
Statistical analysis
When applicable, data was analyzed using Graph Pad Prism software and expressed as
mean values +/- standard error of the mean (SEM). Basic experiments were repeated at
least twice including both positive and negative controls. Biological replicates were
performed on different days or on independent cultures while technical replicates were
performed using cells from the same culture. Experiments were not blinded. All new
reagents were validated prior to use. All generated strains were checked for integration
using PCR, sequencing of the modified locus, western blot, and microscopy when
applicable. All qPCR primers were assessed for single amplicon.
899
900 **Data Availability**
901 Our research resources, including methods, cells, reagents and protocols, are available
902 upon request. Source data for the whole genome sequencing analysis is provided in
903 Extended Data Table 2.
904

905 63. Sidik, S. M. *et al.* A Genome-wide CRISPR Screen in Toxoplasma Identifies
906 Essential Apicomplexan Genes. *Cell* **166,** 1423–1435.e12 (2016).
907
908 64. Pfaffl, M. W. A new mathematical model for relative quantification in real-time RT-
909 PCR. *Nucleic Acids Res.* **29,** e45 (2001).
910
911 65. Shen, B., Brown, K. M., Lee, T. D. & Sibley, L. D. Efficient gene disruption in
912 diverse strains of Toxoplasma gondii using CRISPR/CAS9. *mBio* **5,** e01114-01114
913 (2014).
914
915 66. Camps, M. *et al.* A rRNA mutation identifies the apicoplast as the target for
916 clindamycin in Toxoplasma gondii. *Mol Microbiol.* **5**, 1309-18 (2002).
917
918 67. Dahl, E.L., Rosenthal, P.J. Multiple antibiotics exert delayed effects against the
919 Plasmodium falciparum apicoplast. *Antimicrob Agents Chemother.* **51**, 3485-90
920 (2007).
921
922 68. Striepen, B. *et al.* The plastid of *Toxoplasma gondii* is divided by association with
923 the centrosomes. *J Cell Biol.* **151**, 1423-1424 (2000).
924
925 69. Nkrumah L.J. *et al.* Efficient site-specific integration in *Plasmodium falciparum*
926 chromosomes mediated by mycobacteriophage Bxb1 integrase. *Nature Methods.*
927 **8,** 615-21 (2006).

928
929 70. Gubbels M.J. *et al.* High-throughput growth assay for *Toxoplasma gondii* using
930   yellow fluorescent protein. *Antimicrob Agents Chemother.* **47,** 309-16. (2003).
931
932
933 71. Gibson D.G. *et al.* Enzymatic assembly of DNA molecules up to several hundred
934   kilobases. *Nature Methods.* **6(5),** 343-5. (2009).
935
936 72. Norby, J.G. Coupled assay of Na+, K+-ATPase activity. *Methods Enzymol.* **156**, 116-
937   9. (1988).
938

Acknowledgements
We thank Katrina Hong for assistance in drug assays, Dr. Boris Striepen for providing the
T. gondii FNR-RFP as well as tandem tomato strains, Drs. David Scheinberg and
Ouathek Ouerfelli for providing actinamide, and Dr. Saman Habib for providing *Pf*FtsH1
antibody. This project has been funded with federal funds from the NIAID, NIGMS, and
Director's Fund, National Institutes of Health, Department of Health and Human Services
under Award Numbers 1K08AI097239 (EY), 1DP5OD012119 (EY), U19AI110819
(HAL), 1DP2OD007124 (JCN), P50 GM098792 (JCN), AI016892 (RTS),
F32GM116241 (SBH), and T32GM007276 (KAJ). Funding was also provided by the
Burroughs-Wellcome Fund (EY), Bill and Melinda Gates Foundation (OPP1069759;
JCN), and the Stanford Bio-X SIGF William and Lynda Steere Fellowship (KAJ).
950

951 **Author Information**
952 **Affiliations**
953 Department of Biochemistry, Stanford Medical School, Stanford, California 94305
954 Katherine Amberg-Johnson, Ellen Yeh
955
956 Department of Pathology, Stanford Medical School, Stanford, California 94305
957 Ellen Yeh
958
959 Department of Microbiology & Immunology, Stanford Medical School, Stanford,
960 California, 94305
961 Katherine Amberg-Johnson, Ellen Yeh
962
963 Department of Biology, Massachusetts Institute of Technology, Cambridge,
964 Massachusetts, 02139
965 Sanjay B. Hari, Robert T. Sauer
966
967 Department of Biological Engineering, Massachusetts Institute of Technology,
968 Cambridge, Massachusetts, 02139
969 Suresh M. Ganesan, Jacquin C. Niles
970
971 Department of Infectious Disease, The J. Craig Venter Institute, Rockville, Maryland,
972 20850
973 Hernan A. Lorenzi

Contributions

K.A-J. and E.Y. conceived and designed experiments. K.A-J. performed the majority of the experiments. S.B.H. and R.T.S. conceived and designed the enzymatic experiments. S.B.H purified PfFtsH1 and performed all the enzymatic assays. S.M.G. and J.C.N. designed and generated the FtsH1 knockdown and PDF induction constructs. H.A.L. performed the whole-genome sequencing and variant analysis. K.A-J. and E.Y. analyzed the data and wrote the manuscript. All authors discussed and edited the manuscript.

Competing financial interests

The authors declare no competing financial interests.

Corresponding author

Correspondence to: Ellen Yeh

Fig 1: Actinonin inhibits apicoplast
biogenesis in *P. falciparum*
(a) Time course of parasite growth
during actinonin treatment with or
without IPP, normalized to control
cultures with or without IPP as
appropriate. Error bars represent the
SEM of two biological replicates.
(b) Time course of the
apicoplast:nuclear genome ratio
measured by quantitative PCR (qPCR)
using primers for the apicoplast and
nuclear genomes during treatment
with actinonin and IPP. Genome ratios
were normalized to control parasites
grown with IPP only. Error bars as in
a.
(c) Representative images of the
apicoplast of IPP-rescued control and
actinonin treated parasites 24 hours
after treatment during the schizont
stage. The apicoplast is visualized
using the *P. falciparum* reporter strain
D10 ACP-GFP in which GFP is
targeted to the apicoplast and the
nucleus is stained with Hoescht
33342. During *Plasmodium*
replication, the apicoplast starts as a
single small spherical organelle (ring
stage) which branches and divides into
multiple apicoplasts (schizont stage).
A punctate apicoplast that does not
branch indicates a defect in apicoplast
biogenesis.
1023
1024

Fig 2: A mutation in the protease domain of *ftsH1* is sufficient to confer resistance to
actinonin in *T. gondii*
(a) Representative images of the apicoplast of control and actinonin treated parasites 36
hours after infection. The apicoplast is visualized using the *T. gondii* reporter strain RH
FNR-RFP in which RFP is targeted to the apicoplast and the nucleus is stained with
Hoescht 33342. Each parasite contains one apicoplast, except during cell division when
there may be two. White arrows point at examples of *T. gondii* parasites missing an
apicoplast.
(b) Dose-dependent parasite growth inhibition upon treatment with actinonin for the
actinonin-sensitive parent strain (RH) compared with 3 independent clones following
selection for actinonin resistance (resistant 1, resistant 2, resistant 3). These three
resistant clones are representative of the eight clones submitted for whole genome
sequencing. Growth was measured via summed areas of the plaques formed during
plaque assays and normalized to untreated controls. Error bars represent the SEM of two
biological replicates.
(c) Dose-dependent parasite growth inhibition upon treatment with actinonin for *ftsH1WT*
compared with *ftsH1(N805S)* parasites in RH Δku80 strain. Data was measured and
analyzed as in 2b.
(d) Schematic of TgFtsH1. This protein contains a *N*-unique region containing a putative
transmembrane domain, an AAA ATPase domain used for unfolding proteins, a
peptidase domain with a zinc co-factor in the catalytic site, and a *C*-unique region. The
resistance-conferring variant FtsH(N805S) is found in the peptidase domain near the
catalytic site.
1048

1049

Fig 3: Knockdown of *ftsH1* in *P. falciparum* leads to apicoplast loss and hypersensitivity to actinonin

(a) Schematic of the endogenous knockdown strategy. When aTC is present in the media, the tet-repressor binds aTC and does not bind the 10x-aptamer sequence, which relieves translational repression, allowing PfFtsH1 to be expressed. When aTC is washed out of the media, the tet-repressor binds the 10x-aptamer and prevents expression of PfFtsH1.

(b) Time course of parasite growth without aTC and in the presence or absence of IPP in the media, normalized to the untreated or IPP-rescued parental strain as appropriate. Error bars represent the SEM of two biological replicates.

(c) Time course of the apicoplast:nuclear genome ratio measured by quantitative PCR (qPCR) using primers for the apicoplast and nuclear genomes during treatment with or without aTC. All samples contained IPP to rescue parasite growth. Genome ratios were normalized to respective parental cultures also grown with IPP. Error bars as in **c.**

(d) Dose-dependent parasite growth inhibition by actinonin in the absence or presence of aTC. Error bars as in **c.**

Figure 4: Actinonin inhibits *Pf*FtsH1 *in vitro*.
(a) Schematic of *Pf*FtsH1 constructs used for biochemical assays. Amino acids 91-612 of
the endogenous protein (PfFtsH1 endogenous), which include the AAA+ ATPase and
peptidase domains, were placed between His$_6$-SUMO and GST domains to aid in
purification and solubility. WT is the parent construct, E249Q is an inactivating mutation
in the AAA+ ATPase domain, and D493A is an inactivating mutation in the peptidase
domain.
(b) ATP hydrolysis by *Pf*FtsH1 WT and E249Q measured using a coupled
spectrophotometric assay[72].
(c) ATP-dependent proteolysis of FITC-labeled casein by *Pf*FtsH1 WT and D493A.
(d) Dose-dependent parasite growth inhibition by actinonin (black) or actinamide (red)
with and without knockdown of *Pf*FtsH1. Error bars represent the SEM of two biological
replicates.
(e) Dose-dependent proteolytic inhibition of FITC-labeled casein by *Pf*FtsH1 WT. Error
bars represent the SEM of 3 replicates.
1108

1109
Extended Data 1: Actinonin
specifically inhibits the apicoplast of
P. falciparum **leading to parasite**
death after a single replication cycle
(a) Dose-dependent parasite growth
inhibition by actinonin in the absence
or presence of IPP. Error bars
represent the standard error of the
mean (SEM) of two biological
replicates.
(b) Time course of parasite growth
during treatment with actinonin,
chloramphenicol, or fosmidomycin in
the absence or presence of IPP.
Growth is normalized to untreated or
IPP-rescued controls as appropriate.
Error bars as in **a.** In contrast to other
antimalarials, which cause growth
inhibition in a single replication cycle,
"delayed death" in *P. falciparum* is
associated with inhibitors of apicoplast
gene expression and has previously been described in detail[67]. It is characterized by a
normal apicoplast and cell division during drug treatment for one replication cycle,
followed by halted apicoplast and cell division in daughter parasites of drug-treated
parasites.
1135
1136
Extended Data Fig 2: Actinonin has a distinct inhibition phenotype compared to
inhibitors of apicoplast metabolism and translation
(a) Time course of the apicoplast:nuclear genome ratio measured by quantitative PCR
(qPCR) for targets in the apicoplast and nuclear genome during treatment with actinonin
(black), chloramphenicol (blue), or fosmidomycin (green). While actinonin treatment
blocks apicoplast genome replication in a single replication cycle, this effect is not
observed until the second replication cycle of chloramphenicol treatment and not at all
during fosmidomycin treatment. This is consistent with actinonin blocking apicoplast
biogenesis, chloramphenicol blocking apicoplast translation leading to a delayed
apicoplast biogenesis defect, and fosmidomycin blocking apicoplast metabolic function
but not biogenesis. All samples were grown in IPP and genome ratios were normalized to
the untreated control cultures also containing IPP. Error bars represent the SEM of at
least 2 biological replicates.
(b-e) Representative images of the apicoplast during schizont stage of three successive
replication cycles in untreated **(b),** chloramphenicol **(c)**, fosmidomycin **(d)** and actinonin
(e) treated cultures all grown with IPP. The apicoplast is visualized using the *P.*
falciparum reporter strain D10 ACP-GFP in which GFP is targeted to the apicoplast and
the nucleus is stained with Hoescht 33342. In this case, a branched apicoplast indicates

successful apicoplast development while punctate apicoplasts (observed in replication
cycle 1 for actinonin treatment (e) and replication cycle 2 for chloramphenicol treatment
(c)) represents an apicoplast that has failed to develop. The apicoplast is no longer
present after replication cycle 2 of actinonin treatment and upon replication cycle 3 of
chloramphenicol treatment, which leads to complete mislocalization of the GFP.
1160

1161

Extended Data Fig 3: Actinonin is unlikely to inhibit the peptide deformylase of *P.*
falciparum
(A) Western blot of the parental line (NF54attB-pCRISPR) and the PDF-myc parasites
grown with our without aTC for 24 hours. Induction of the second copy of PfPDF (PDF-
myc) with 4uM aTC results in two bands, the top lighter band representing unprocessed
PDF-myc, and the bottom darker band representing processed PDF-myc. PfAldolase is
used as a loading control. Induction with 0.125-4uM aTC results in similar amount of
PDF-myc induction (data not shown).
(B) Western blot for PDF-myc of parasites with or without their apicoplast. An
accumulation of unprocessed PDF-myc is observed when the apicoplast is missing, due
to loss of the transit peptide cleavage that usually occurs upon import to the apicoplast.
This has been shown previously for apicoplast-resident proteins and is consistent
apicoplast localization[8].
(C) Dose dependent parasites growth inhibition by actinonin in the presence of 4uM aTC
does not change the actinonin EC_{50}. This experiment was also performed under IPP
rescue conditions, to confirm apicoplast specificity of actinonin and with a range of aTC
concentrations (0.125-4uM) to insure max expression of PDF-myc (data not shown).
Error bars represent the SEM of 3 biological replicates.
(D) Parasite growth after one or two replication cycles after treatment with actinonin,
chloramphenicol, or both actinonin and chloramphenicol normalized to growth of an
untreated control. Treatment with actinonin alone inhibited growth after the first
replication cycle, whereas treatment with chloramphenicol alone inhibited growth after
the second replication cycle. Co-treatment with chloramphenicol, which targets
apicoplast translation, did not suppress effects of actinonin treatment, which was
inconsistent with actinonin targeting the peptide deformylase (PDF) of the apicoplast.
This experiment was tried using a range of concentrations of actinonin and
chloramphenicol to insure the data was not the result of partial inhibition. All
concentrations that lead to apicoplast-specific death gave this phenotype (data not
shown).
1191

1192
1193

1194

Extended Data Fig 4: Actinonin
treatment causes "delayed death" in
T. gondii **associated with apicoplast**
loss

(A) Time course of parasite growth of
untreated (blue), clindamycin (green),
pyrimethamine (black), or actinonin
(red) treated parasites over the course
of two lytic cycles. Unlike *P.*
falciparum, T. gondii undergoes
multiple replication cycles in the host
cell before lysis, thus each *T. gondii*
lytic cycle represents multiple parasite
replication cycles. Pyrimethamine
inhibits parasite dihydrofolate
reductase and was used as a control for
a non-apicoplast targeting drug that
inhibits growth in a single lytic cycle.
As previously reported[66], clindamycin,
an apicoplast translation inhibitor,
gave a "delayed death" phenotype in *T.*
gondii characterized by completed cell
divisions and host re-invasion during
drug treatment in the first lytic cycle,
followed by halted cell division in the
second lytic cycle. Actinonin also led
to growth inhibition in the second lytic
cycle, suggesting that it also targets the
apicoplast. Error bars represent the

SEM of two biological replicates. It is important to distinguish between apicoplast-
associated "delayed death" in *P. falciparum* and that in *T. gondii*. In *T. gondii*, apicoplast
loss occurs in the first lytic cycle and is temporally separate from defects in parasite cell
division and growth inhibition observed in the second lytic cycle; whereas in *P.*
falciparum defects in apicoplast biogenesis, parasite cell division, and growth inhibition
all occur in the second replication cycle. "Delayed death" in *T. gondii* therefore appears
more broadly associated with disruption of apicoplast biogenesis, where "delayed
death"in *P. falciparum* appears more specific to disruption of apicoplast gene expression
that leads to delayed biogenesis defects.
(B) To determine dose-dependent drug target specificity, we used a previously described
fluorescent growth assay to quantify the kinetics of death in *T. gondii*[70]. Briefly, *T. gondii*
expressing tandem-tomato are treated with drug and fluorescence is quantified daily as a
proxy of parasite replication. While 40uM actinonin leads to delayed-death kinetics
similar to clindamycin, 80uM actinonin results in immediate death kinetics similar to
pyrimethamine, indicating that at this concentration, actinonin hits a second, non-
apicoplast target.

1240
Extended Data Fig 5: ***C*-terminal**
cleavage of PfFtsH1 is dependent
on the presence of the apicoplast
and efficiency of *Pf*FtsH1 knock-
down can be assessed in parasites
missing their apicoplast.

(A) Western blot of *Pf*FtsH-FLAG
levels in parasites with and without
an apicoplast and with and without
aTC induction. Consistent with the
previous report of *Pf*FtsH1 *C*-
terminal processing, we were unable
to detect full-length *Pf*FtsH1-FLAG
using anti-FLAG in parasites with
intact apicoplasts (lane 1)[37,40].

However, in parasites missing apicoplasts, full-length *Pf*FtsH1-FLAG was detectable,
suggesting that *Pf*FtsH1 processing requires the apicoplast (lanes 2). *Pf*FtsH1-FLAG is
not observed in parasites with or without apicoplasts if there is no aTC induction,
indicating that the band representing FtsH-FLAG is specific and not an artifact of missing
apicoplasts (lanes 3-4 respectively). All samples contain IPP to rescue growth and Cas9-
FLAG is used as a loading control. Each sample was taken at the trophozoite stage.
(B) To assess the knockdown efficiency of *Pf*FtsH1, we used a western blot comparing
*Pf*FtsH1-FLAG levels in the presence (lanes 1-4) or absence of aTC (lanes 5-8) in IPP-
rescued parasites missing their apicoplast. Each sample was taken at the trophozoite stage
and cycle 0 indicates 24 hours after the removal of aTC. Lanes 9 and 10 are samples from
the parental strain that do not contain the FLAG-tag or the aptamer sequence in the 3'
UTR of *Pf*FtsH1. In each case, Cas9-FLAG was used as a loading control. *Pf*FtsH1-
FLAG levels were reduced to undetectable levels at 24 hours after aTC removal,
validating our knockdown strategy.
1270
1271
1272
1273

Extended Data Fig 6: Knockdown of *Pf*FtsH1 specifically disrupts the apicoplast and leads to specific hypersensitivity to actinonin

(**a**) Time course of parasite growth with or without aTC and with or without IPP in the media. IPP rescues the growth defect observed in upon *Pf*FtsH1 downregulation, indicating that *Pf*FtsH1 is essential for an apicoplast-specific function. Growth is shown normalized to the untreated or IPP-rescued parental strain as appropriate. Error bars represent the SEM of two biological replicates.

(**b**) Dose-dependent parasite growth inhibition by actinonin with or without aTC for the parental (red) and *Pf*FtsH (black) strain. The EC_{50} of the parental strain is unchanged by the removal of aTC. Error bars as in **a.**

(**c**) Dose-dependent parasite growth inhibition by fosmidomycin with or without aTC and with or without IPP. The fosmidomycin EC_{50} is unchanged by regulating levels of *Pf*FtsH1, indicating that the observed hypersensitivity to actinonin upon knockdown of *Pf*FtsH1 is specific to actinonin and does not occur for all apicoplast drug. Error bars represent the SEM of three technical replicates.

(**d**) Dose-dependent parasites growth inhibition by chloramphenicol during the second lytic cycle (120 h) with or without aTC. Error bars represent the SEM of three technical replicates. The chloramphenicol EC_{50} is unchanged by regulating levels of PfFtsH1, indicating that the observed hypersensitivity to actinonin upon knockdown of PfFtsH1 is specific to actinonin and does not occur for all apicoplast drugs.

1316 **Extended Data Fig 7: Purification of *Pf*FtsH1.** Purified samples were separated by
1317 SDS-PAGE and stained with Coomassie. Full-length product is 97 kDa.
1318

1319