The Stochastic Early Reaction, Inhibition, and Late Action (SERIA) Model for Antisaccades

SERIA - A model for errors and reaction times in the antisaccade task

Eduardo A. Aponte[1,*], Dario Schoebi[1], Klaas E. Stephan[1,2], Jakob Heinzle[1*]

[1]Translational Neuromodeling Unit, Institute for Biomedical Engineering, University of Zurich & Swiss Institute of Technology Zurich, Zurich, Switzerland

[2]Wellcome Trust Centre for Neuroimaging, University College London, London, United Kingdom

*Corresponding authors

aponte@biomed.ee.ethz.ch

heinzle@biomed.ee.ethz.ch

Abstract

The antisaccade task is a classic paradigm used to study the voluntary control of eye movements. It requires participants to suppress a reactive eye movement to a visual target and to concurrently initiate a saccade in the opposite direction. Although several models have been proposed to explain error rates and reaction times in this task, no formal model comparison has yet been performed. Here, we describe a Bayesian modeling approach to the antisaccade task that allows us to formally compare different models on the basis of their evidence. First, we provide a formal likelihood function of actions (pro- and antisaccades) and reaction times based on previously published models. Second, we introduce the *Stochastic Early Reaction, Inhibition, and late Action model* (SERIA), a novel model postulating two different mechanisms that interact in the antisaccade task: an early GO/NO-GO race decision process and a late GO/GO decision process. Third, we apply these models to a data set from an experiment with three mixed blocks of pro- and antisaccade trials. Bayesian model comparison demonstrates that the SERIA model explains the data better than competing models that do not incorporate a late decision process. Moreover, we show that the race decision processes postulated by the SERIA model are, to a large extent, insensitive to the cue presented on a single trial. Finally, we use parameter estimates to demonstrate that changes in reaction time and error rate due to the probability of a trial type (prosaccade or antisaccade) are best explained by faster or slower inhibition and the probability of generating late voluntary prosaccades.

Author summary

One widely replicated finding in schizophrenia research is that patients tend to make more errors in the antisaccade task, a psychometric paradigm in which participants are required to look in the opposite direction of a visual cue. This deficit has been suggested to be an endophenotype of schizophrenia, as first order relatives of patients tend to show similar but milder deficits. Currently, most models applied to experimental findings in this task are limited to fit average reaction times and error rates. Here, we propose a novel statistical model that fits experimental data from the antisaccade task, beyond summary statistics. The model is inspired by the hypothesis that antisaccades are the result of several competing decision processes that interact nonlinearly with each other. In applying this model to a relatively large experimental data set, we show that mean reaction times and error rates do not fully reflect the complexity of the processes that are likely to underlie experimental findings. In the future, our model could help to understand the nature of the deficits observed in schizophrenia by providing a statistical tool to study their biological underpinnings.

Introduction

In the antisaccade task ([1]; for reviews, see [2,3]), participants are required to saccade in the contralateral direction of a visual cue. This behavior is thought to require both the inhibition of a reflexive saccadic response towards the cue and the initiation of a voluntary eye movement in the opposite direction. A failure to inhibit the reflexive response leads to an erroneous saccade towards the cue (i.e., a prosaccade), which is often followed by a corrective eye movement in the opposite direction (i.e., an antisaccade). As a probe of inhibitory capacity, the antisaccade task has been widely used to study psychiatric and neurological diseases [3]. Notably, since the initial report [4], studies have consistently found an increased number of errors in patients with schizophrenia when compared to healthy controls, independent of medication and clinical status [5-8]. Moreover, there is evidence that an increased error rate constitutes an endophenotype of schizophrenia, as antisaccade deficits are also present in non-affected, first-degree relatives of diagnosed individuals (for example [5,7]; but for negative findings see for example [9][10]).

Unfortunately, the exact nature of the antisaccade deficits and their biological origin in schizophrenia remain unclear. One path to improve our understanding of these experimental findings is to develop generative models of their putative computational and/or neurophysiological causes [11]. Generative models that capture the entire distribution of responses can reveal features of the data that are not apparent when only considering summary statistics such as mean error rate (ER) and reaction time (RT) [12-15]. Additionally, this type of model can potentially relate behavioral findings in humans to their biological substrate.

Here, we apply a generative modeling approach to the antisaccade task. First, we introduce a novel model of this paradigm based on previous proposals [16-20]. For this, we formalize the ideas introduced by Noorani and Carpenter [17] and extend them into what we refer to as the *Stochastic Early Response, Inhibition and late Action* (SERIA) model. Second, we apply both models to an experimental data set of three mixed blocks of pro- and antisaccades trials with different trial type probability using formal Bayesian inference. More specifically, we compare several models using Bayesian model comparison. Third, we use the parameter estimates from the best model to investigate the effects of our experimental manipulation. We found that there was positive evidence

in favor of the SERIA model when compared to our formalization of the model proposed

in [17]. Moreover, the parameters estimated through model inversion revealed the

complexity of the decision processes underlying the antisaccade task that is not obvious

from mean RT and ER.

This paper is organized as follows. First, we formalize the model developed in [17] and

introduce the SERIA model. Second, we describe our experimental setup. Third, we

present our behavioral findings in terms of summary statistics (mean RT and ER), the

comparison between different models, and the parameter estimates. Finally, we review

our findings, discuss other recent models, potential future developments, and

translational applications.

Materials and methods

Ethics statement

All participants gave written informed consent before the study. All experimental procedures were approved by the local ethics board (Kantonale Ethikkomission Zürich, KEK-ZH-Nr.2014-0246).

Race models for antisaccades

In this section, we derive a formal description of the models evaluated in this paper. We start with a formalized version of the model proposed by Noorani and Carpenter in [17] and proceed to extend it. Their approach resembles the model originally proposed by Camalier and colleagues [21] to explain RT and ER in the double step and search step tasks, in which participants are either asked to saccade to successively presented targets or to saccade to a target after a distractor was shown. Common to all these tasks is that subjects are required to inhibit a prepotent reaction to an initial stimulus and then to generate an action towards a secondary goal. Briefly, Camalier and colleagues [21] extended the original 'horse-race' model [16] by including a secondary action in countermanding tasks. In [17], Noorani and Carpenter used a similar model in combination with the LATER model [22] in the context of the antisaccade task by postulating an endogenously generated inhibitory signal. Note that this model, or variants of it, have been used in several experimental paradigms (reviewed in [20]). Here, we limit our discussion to the antisaccade task.

The pro, stop, and antisaccade model (PROSA)

Following [17], we assume that the RT and the type of saccade generated in a given trial are caused by the interaction of three competing processes or units. The first unit u_p represents a command to perform a prosaccade, the second unit u_s represents an inhibitory command to stop a prosaccade, and the third unit u_a represents a command to perform an antisaccade. The time t required for unit u_i to arrive at threshold s_i is given by:

$$s_i = r_i t, \qquad (1)$$

$$\frac{s_i}{r_i} = t, \qquad (2)$$

where r_i represents the slope or increase rate of unit u_i, s_i represents the height of the
threshold, and t represents time. We assume that, on each trial, the increase rates are
stochastic and independent of each other.

The time and order in which the units reach their thresholds s_i determines the action and
RT in a trial. If the prosaccade unit u_p reaches threshold before any other unit at time t, a
prosaccade is elicited at t. If the antisaccade unit arrives first, an antisaccade is elicited at
t. Finally, if the stop unit arrives before the prosaccade unit, an antisaccade is elicited at
the time when the antisaccade unit reaches threshold. It is worth mentioning that,
although this model is motivated as a race-to-threshold model, actions and RTs depend
only on the arrival times of each of the units and ultimately no explicit model of increase
rates or thresholds is required. Thus, for the sake of clarity, we refer to this approach as
a 'race' model, in contrast to 'race-to-threshold' models that explicitly describe increase
rates and thresholds.

Formally (but in a slight abuse of language), the two random variables of interest, the
reaction time $T \in [0, \infty[$ and the type of action performed $A \in \{pro, anti\}$, depend only
on three further random variables: the arrival times $U_p, U_s, U_a \in [0, \infty[$ of each of the
units. The probability of performing a prosaccade at time t is given by the probability of
the prosaccade unit arriving at time t, and the stop and antisaccade unit arriving
afterwards:

$$p(A = pro, T = t) = p(U_p = t)p(U_a > t)p(U_s > t). \qquad (3)$$

143   The probability of performing an antisaccade at time t is given by

$$p(A = anti, T = t) = \qquad (4)$$

$$p(U_a = t)p(U_p > t)p(U_s > t) + p(U_a = t) \int_0^t p(U_s = \tau)p(U_p > \tau)d\tau.$$

144   The first term on the right side of Eq. 4 corresponds to the unlikely case that the
145   antisaccade unit arrives before the prosaccade and the stop unit. The second term
146   describes trials in which the stop unit arrives before the prosaccade unit. It can be
147   decomposed into two terms:

$$p(U_a = t) \int_0^t p(U_s = \tau)p(U_p > \tau)d\tau \qquad (5)$$

$$= p(U_a = t) \left(p(U_s < t)p(U_p > t) + \int_0^t p(U_s = \tau)p(\tau < U_p < t)d\tau \right)$$

$$= p(U_a = t) \left(p(U_s < t)p(U_p > t) + \int_0^t p(U_s < \tau)p(U_p = \tau)d\tau \right). \qquad (6)$$

148   The term $p(U_a = t) \int_0^t p(U_s < \tau)p(U_p = \tau)d\tau$ describes the condition in which the

149   prosaccade unit is inhibited by the stop unit allowing for an antisaccade. Note that if the

150   prosaccade unit arrives later than the antisaccade unit, the arrival time of the stop unit is

151   irrelevant. That means that we can simplify Eq. 4 to

$$p(A = anti, T = t) = p(U_a = t) \left(p(U_p > t) + \int_0^t p(U_s < \tau)p(U_p = \tau)d\tau \right). \qquad (7)$$

Eq. 3 and 7 constitute the likelihood function of a single trial, defining the joint probability

of an action and the corresponding RT. We refer to this likelihood function as the PRO-

Stop-Antisaccade (PROSA) model. It shares the central assumptions of [17], namely: (i)

the time to reach threshold of each of the units is assumed to depend linearly on the rate

r, (ii) it includes a stop unit whose function is to inhibit prosaccades and (iii) there is no

lateral inhibition between the different units. Finally, (iv) RTs are assumed to be equal to

the arrive-at-threshold times. Note that the RT distributions are different from the arrival

time distributions because of the interactions between the units described above. The

main difference of this model compared to [17] is that we do not exclude *a priori* the

possibility of the antisaccade unit arriving earlier than the other units. Aside from this,

both models are conceptually equivalent.

The Stochastic Early Reaction, Inhibition, and Late Action Model (SERIA)

The PROSA model is characterized by a strict association between units and action types.

In other words, the unit u_p leads unequivocally to a prosaccade, whereas the unit u_a

always triggers an antisaccade. This implies that if the distribution of the arrival times of

the units is unimodal and strictly positive, the PROSA model cannot predict voluntary

slow prosaccades with a late peak, or in simple words, the PROSA model cannot account

for slow, voluntary prosaccades that have been postulated in the antisaccade task [23].

Similarly, it has been argued that prosaccade RT can be described by the mixture of two

distributions (for example [2,22]).To account for this, we introduce the Stochastic Early

Reaction, Inhibition and Late Action (SERIA) model.

According to this model, and in analogy to the PROSA model, an early reaction takes place

at time t if the early unit u_e arrives before the late and inhibitory units, u_l and u_i,

respectively. If the inhibitory or late unit arrives before the early unit, a late response is

triggered at the time the late unit reaches threshold. Crucially, both early and late

responses can trigger pro- and antisaccades with a certain probability. Thus, in parallel

to the race processes which determine RTs, an independent, secondary decision process

is responsible for which reaction is generated. Fig. 1 shows the structure of the SERIA

model.

Fig 1. Layout of the SERIA model.

The presentation of a visual cue (a green bar) triggers the race of three independent units. The inhibitory unit can stop an early response. Importantly, both early and late responses can trigger pro- and antisaccades. Note that the PROSA model is a special case of the SERIA model in which $\pi_e = 1$ and $\pi_l = 0$, i.e. all early responses are prosaccades, whereas all late responses are antisaccades.

182 To formalize the concept of early and late responses, we introduce a new unobservable

183 random variable that represents the type of response $R \in \{early, late\}$. The distribution

184   of the RTs is analogous to the PROSA-model, such that, for instance, the probability of an

185   early response at time t is given by

$$p(R = early, T = t) = p(U_e = t)p(U_i > t)p(U_l > t) \qquad (8)$$

where U_e, U_i, and U_l represent the arrival times of the early, inhibitory, and late units

respectively. The fundamental assumption of the SERIA model is that a secondary

decision process, beyond the race between early, inhibitory, and late units, decides the

action generated in a single trial. An initial approach to model this secondary decision

process is to assume that the action type (pro- or antisaccade) is conditionally

independent of the RT given the response type (early or late). Hence, the distribution of

RTs is not *a priori* coupled to the saccade type anymore; RT distributions for both pro-

and antisaccades could in principle be bimodal, consisting of both fast reactive and slow

voluntary saccades.

Formally, the conditional independency assumption can be written down as

$$p(A, T|R) = p(A|R)p(T|R), \qquad (9)$$

$$p(A, T|R)p(R) = p(A|R)p(T|R)p(R), \qquad (10)$$

$$p(A, T, R) = p(A|R)p(T, R). \qquad (11)$$

196   The term $p(A|R)$ is simply the probability of an action, given a response type. We denote

197   it as

$$p(A = pro|R = early) = \pi_e \in [0,1], \qquad (12)$$

$$p(A = anti|R = early) = 1 - \pi_e, \qquad (13)$$

$$p(A = pro|R = late) = \pi_l \in [0,1], \qquad (14)$$

$$p(A = anti|R = late) = 1 - \pi_l. \qquad (15)$$

198   Since the type of response R is not observable, it is necessary to marginalize it out in Eq.

199   [11] to obtain the likelihood of the SERIA model:

$$p(A, T) = p(A, T, R = early) + p(A, T, R = late). \qquad (16)$$

200 The complete likelihood of the model is given by substituting the terms in Eq. [16]:

$$p(A = pro, T = t) = \pi_e p(U_e = t)p(U_i > t)p(U_l > t) + \quad (17)$$

$$\pi_l p(U_l = t)\left(p(U_e > t) + \int_0^t p(U_e = \tau)p(U_i < \tau)d\tau\right),$$

$$p(A = anti, T = t) = (1 - \pi_e)p(U_e = t)p(U_i > t)p(U_l > t) + \quad (18)$$

$$(1 - \pi_l)p(U_l = t)\left(p(U_e > t) + \int_0^t p(U_e = \tau)p(U_i < \tau)d\tau\right).$$

201 It is worth noting here that the PROSA model is a special case of the SERIA model, namely,
202 it corresponds to the assumption that $\pi_e = 1$ and $\pi_l = 0$. The SERIA model allows for
203 bimodal distributions, as both early and late responses can be pro- and antisaccades.
204 Importantly, one prediction of the model is that late prosaccades have the same
205 distribution as late antisaccades.

Late race competition model for saccade type

Until now, we have assumed that the competition that leads to late pro- and antisaccades
does not depend on time in the sense that late actions are conditionally independent of
RT. This assumption can be weakened by postulating a secondary race between late
responses; this leads us to a modified version of the SERIA model, that we refer to as the
late race SERIA model (SERIA$_{lr}$). The derivation proceeds similarly to the SERIA model,
except that we postulate a fourth unit that generates late prosaccades instead of assuming
that the late decision process is time insensitive.

This version of the SERIA model includes an early unit u_e that, for simplicity, we assume
produces only prosaccades, an inhibitory unit that stops early responses u_i, a late unit
that triggers antisaccades u_a, and a further unit that triggers late prosaccades u_p. As
before, if the early unit reaches threshold before any other unit, a prosaccade is generated
with probability

$$p(U_e = t)p(U_i > t)p(U_a > t)p(U_p > t). \quad (19)$$

219 If any of the late units arrive first, the respective action is generated with probability:

$$\text{Antisaccade: } p(U_a = t)p\big(U_p > t\big)p(U_e > t)p(U_i > t). \quad (20)$$

$$\text{Prosaccade: } p(U_p = t)p(U_a > t)p(U_e > t)p(U_i > t). \qquad (21)$$

220   Finally, if the inhibitory unit arrives first, either a late pro- or antisaccade is generated

221   with probability

$$\text{Antisaccades: } p(U_a = t)p(U_p > t)\left(\int_0^t p(U_i = \tau)p(U_e > \tau)d\tau\right), \qquad (22)$$

$$\text{Prosaccades: } p(U_p = t)p(U_a > t)\left(\int_0^t p(U_i = \tau)p(U_e > \tau)d\tau\right). \qquad (23)$$

Implicit in the last two terms is the competition between the late units, which are

assumed again to be independent of each other. Formally, this competition is expressed

as the probability of, for example, the late antisaccade unit arriving before a late

prosaccade $p(U_a = t)p(U_p > t)$. A schematic representation of the model is shown in Fig.

2. This late race is similar to the Linear Ballistic Accumulation model proposed by [24],

although in that model decisions are seen as the result of a race of ballistic accumulation

processes with fixed threshold but stochastic base line and increase rate. Here we only

assume that the late decision process is a GO-GO race [21].

Early response

early

Early prosaccade
at time rt_e

inhib.

late pro

Faster unit defines
response type and time rt_l

late anti

Late response

Fig 2. Layout of the SERIA$_{Ir}$ model.

The presentation of a visual cue (a green bar) triggers the race of four independent
units. The inhibitory unit can stop an early response. The late decision process is
triggered by the competition between two further units.

230   The likelihood of an action is given by summing over all possible outcomes that lead to
231   that action:

$$p(A = pro, T = t) = p(U_e = t)p(U_i > t)p(U_a > t)p(U_p > t) + \qquad (24)$$

$$p(U_p = t)p(U_a > t)p(U_i > t)p(U_e > t) +$$

$$p(U_p = t)p(U_a > t)\left(\int_0^t p(U_i = \tau)p(U_e > \tau)d\tau \right),$$

$$(A = anti, T = t) = p(U_a = t)p(U_p > t)p(U_i > t)p(U_e > t) + \qquad (25)$$

$$p(U_a = t)p(U_p > t)\left(\int_0^t p(U_i = \tau)p(U_e > \tau)d\tau \right).$$

232 We have left out some possible simplifications in Eq. 24 and 25 for the sake of clarity.

233 The conditional probability of a late antisaccade is given by the interaction between the
234 late units, such that

$$p(U_a < U_p) = \int_0^\infty p(U_a = t)p(U_p > t)dt = 1 - p(U_p < U_a), \tag{26}$$

is analogous to the probability of a late antisaccade $1 - \pi_l$ in the SERIA model. This
observation shows that the main difference between the SERIA and SERIA$_{lr}$ model is that
the former postulates that the distribution of late pro- and antisaccades are equal and
conditionally independent of the action performed, whereas the latter constrains the
probability of a late antisaccade to be a function of the arrival times of the late units.

The expected *response time* of late pro and antisaccade actions is given by

$$\frac{1}{p(U_p < U_a)} \int_0^\infty t\, p(U_p = t)p(U_a > t)dt, \tag{27}$$

$$\frac{1}{p(U_a < U_p)} \int_0^\infty t\, p(U_a = t)p(U_p > t)dt. \tag{28}$$

241 We will refer to these terms as the mean *response time* of pro- and antisaccade actions, in
242 contrast to the mean arrival times, which are the expected value of any single unit.

Non-decision time

The models above can be further finessed to account for non-decision times δ by
transforming the RT t to $t_\delta = t - \delta$. The delay δ might be caused by, for example,
conductance delays from the retina to the cortex. In addition, the antisaccade or late units
might include a constant delay δ_a, which is often referred to as the antisaccade cost [1].
Note that the model is highly sensitive to δ because any RT below it has zero probability.
In order to relax this condition and to account for early outliers, we assumed that
saccades could be generated before δ at a rate $\eta \in [0,1]$ such that the marginal likelihood
of an outlier is

$$p(T < \delta) = p(T_\delta < 0) = \eta. \tag{29}$$

252 For simplicity, we assume that outliers are generated with uniform probability in the
253 interval $[0, \delta]$:

$$p(T = t) = \frac{\eta}{\delta} \text{ if } t < \delta. \tag{30}$$

254 Furthermore, we assume that the probability of an early outlier being a prosaccade was
255 approximately 100 times higher than being an antisaccade. Because of the new parameter
256 η, the distribution of saccades with a RT larger than δ needs to be renormalized by the
257 factor $1 - \eta$. In the case of the PROSA model, for example, this means that the joint
258 distribution of action and RT is given by the conditional probability

$$p(A = pro, T = t_\delta | t_\delta > 0) = \quad p(U_p = t_\delta)p(U_a > t_\delta - \delta_a)p(U_s > t_\delta), \tag{31}$$

$$p(U_a < 0) = 0, \tag{32}$$

$$p(A = anti, T = t_\delta | t_\delta > 0) = \tag{33}$$

$$p(U_a = t_\delta - \delta_a)\left(p(U_p > t_\delta) + \int_0^{t_\delta} p(U_p = \tau)p(U_s < \tau)d\tau\right).$$

259 A similar expression holds for the SERIA models. However, in the PROSA model a unit-
260 specific delay is equal to an action-specific delay. By contrast, in the SERIA model both
261 early and late responses can generate pro- and antisaccades. Thus, δ_a, represents a delay
262 in the late actions that affects both late pro- and antisaccades.

Parametric distributions of the increase rate

The models discussed in the previous sections can be defined independently of the
distribution of the rate of each of the units. In order to fit experimental data, we
considered four parametric distributions with positive support for the rates: gamma [13],
inverse gamma, lognormal [25] and the truncated normal distribution (similarly to [22]
and [24]). Table 1 and Fig 3 summarize these distributions, their parameters, and the
corresponding arrival time densities. We considered five different configurations: 1) all
units were assigned *inverse gamma* distributed rates, 2) all units were assigned *gamma*
distributed rates, 3) the increase rate of the pro and stop units (or early and the inhibitory
units) was *gamma distributed* but the antisaccade (late) unit's increase rate was *inverse*
gamma distributed, 4) all the units were assigned *lognormal* distributed rates or 5) all
units were assigned *truncated normal* distributed rates.

Table 1: Parametric density functions of the increase rates.

Name	Parameters	Rate p.d.f.	Arrival time p.d.f.
Gamma	k, θ	$\dfrac{\theta^{-k}}{\Gamma(k)} e^{-r/\theta} r^{k-1}$	$\dfrac{\theta^{k}}{\Gamma(k)} e^{-\theta/t} t^{-k-1}$
Inv. gamma	k, θ	$\dfrac{\theta^{k}}{\Gamma(k)} e^{-\theta/r} r^{-k-1}$	$\dfrac{\theta^{-k}}{\Gamma(k)} e^{-t/\theta} t^{k-1}$
Log normal	μ, σ^2	$\dfrac{1}{\sqrt{2\pi}\sigma r} e^{-\frac{1}{2}\left(\frac{\ln r - \mu}{\sigma}\right)^2}$	$\dfrac{1}{\sqrt{2\pi}\sigma t} e^{-\frac{1}{2}\left(\frac{\ln t + \mu}{\sigma}\right)^2}$
T. normal	μ, σ^2	$\dfrac{1}{Z} e^{-\frac{1}{2}\left(\frac{r - \mu}{\sigma}\right)^2}$	$\dfrac{1}{Zt^2} e^{-\frac{1}{2}\left(\frac{t^{-1} - \mu}{\sigma}\right)^2}$

Z is the normalization constant $Z = \int_0^\infty exp\left(-\dfrac{(r-\mu)^2}{2\sigma^2}\right) dr$.

Fig 3. Illustration of probability distributions used to model increase rates.

Left: Distribution of the rates based on different probability density functions: Normal (red), gamma (blue), inverse gamma (green), and log-normal (cyan). All distributions were matched to have equal mean and variance. Center: Probit plots of the same distributions. While the gamma and lognormal distributions are very close to the straight line induced by the normal distribution, the inverse gamma distribution diverges slightly more from linearity. Right: Arrival time distribution (scaled to ms).

All the parametric distributions considered here can be fully characterized by two parameters which we generically refer to as k and θ. Hence, the PROSA model is characterized by the parameters for each unit $k_p, k_a, k_s, \theta_p, \theta_a, \theta_s$. The SERIA model can be characterized by analogous parameters $k_e, k_l, k_i, \theta_e, \theta_l, \theta_i$ and the probabilities of early and late prosaccades π_e and π_l. In the case of the SERIA$_{lr}$ model, the probability of a late

280   prosaccade is replaced by the parameters of a late prosaccade unit k_p, θ_p. In addition to

281   the unit parameters, all models included the non-decision time δ, the antisaccade (or late

282   unit) cost δ_a, and the marginal rate of early outliers η.

Experimental procedures

In this section, we describe the experimental procedures, statistical methods, and inference scheme used to invert the models above. The data is from the placebo condition of a larger pharmacological study that will be reported elsewhere.

Participants

Fifty-two healthy adult males naïve to the antisaccade task were invited to a screening session through the recruitment system of the Laboratory of Social and Neural Systems Research of the University of Zurich. During screening, and after being debriefed about the experiment, subjects underwent an electrocardiogram, a health survey, a visual acuity test, and a color blindness test. Subjects were excluded if any of the following criteria were met: age below 18 or above 40 years, regular smoking, alcohol consumption the day before the experiment, any possible interaction between current medication and levodopa or benserazide, pulse outside the range 55-100bpm, recreational drug intake in the past 6 months, history of serious mental or neurological illness, or if the medical doctor supervising the experiment deemed the participant not apt. All subjects gave their written informed consent to participate in the study and received monetary compensation.

Procedure

Each subject was invited to two sessions. During both visits, the same experimental protocol was followed. After arrival, placebo or levodopa (Madopar® DR 250, 200mg of levopa + 50 mg benserazide) was orally administered in the form of shape- and color-matched capsules. The present study is restricted to data from the session in which subjects received placebo. Participants and experimenters were not informed about the identity of the substance. Immediately afterwards subjects were introduced to the experimental setup and to the task through a written document. This was followed by a short training block (see below).

The experiment started 70 minutes after substance administration. Subjects participated in three blocks of 192 randomly interleaved pro- and antisaccade trials. The percentages of prosaccade trials in the three blocks were 20%, 50%, or 80%. This yielded three *prosaccade probability* (PP) conditions: PP20, PP50, and PP80. Thus, in the PP20 block, subjects were presented a prosaccade cue in 38 trials, while in all other 154 trials an antisaccade cue was shown. The order of the trials was randomized in each block, but the

315   same order was used in all subjects and sessions. The order of the conditions was
316   counterbalanced across subjects.

Stimulus and apparatus

During the experiment, subjects sat in front of a CRT monitor (Philipps 20B40, distance
eye-screen: $\simeq 60cm$, refresh rate: 75Hz). The screen subtended a horizontal visual angle
of 38 degrees of visual angle (dva). Eye movements were recorded using a remote
infrared camera (Eyelink II, SR-Research, Canada). Participants' head was stabilized with
a chin rest. Data were stored at a sampling rate of 500 Hz.

During the task, two red dots (0.25dva) that constituted the saccadic targets were
constantly displayed at an eccentricity of $\pm$12dva. Displaying the saccadic target before
the execution of an antisaccade has been reported to affect saccadic velocity and accuracy,
but not RTs [26], and arguably decreases the need for sensorimotor transformations [27].
At the beginning of each trial, a gray fixation cross ($0.6 \times 0.6\ dva$) was displayed at the
center of the screen. After a random fixation interval (500 to 1000 ms), the cross
disappeared, and the cue instructing either a pro- or an antisaccade trial (see below) was
shown centered on either of the red dots. As mentioned above, in each block, subjects
were presented with a prosaccade cue in either 20, 50, or 80 percent of the trials. The
order of the presentation of the cues was randomized. The cue was a green rectangle
($3.48 \times 0.8 dva$) displayed for 500ms in either horizontal (prosaccade) or vertical
orientation (antisaccade). Once the cue was removed and after 1000ms, the next trial
started.

Subjects were instructed to saccade in the direction of the cue when a horizontal bar was
presented (prosaccade trial) and to saccade in the opposite direction when a vertical bar
was displayed (antisaccade trial, see Fig. 3). See [28,29] for similar task designs.

Fig 4. Task design.

After a variable fixation period of 500-1000ms (top) the cue (green rectangle) appeared on the screen for 500 ms. The orientation of the cue (horizontal or vertical) indicated the required response (prosaccade or antisaccade).

Prior to the main experiment, participants were trained on the task in a block of 50

prosaccade trials, immediately followed by 50 antisaccade trials. During the training,

subjects were automatically informed after each trial whether their response was correct

or not (see below), or whether they had failed to produce a saccade within 500ms after

cue presentation (CP). No feedback was given during the main experimental blocks.

Data preparation

Data were parsed and preprocessed using the Python programming language (2.7).

Saccades were detected using the algorithm provided by the eyetracker manufacturer

(SR Research), which uses a velocity and acceleration threshold of $22dva/s$ and

$3800dva/s^2$ [30]. We only considered saccades with a magnitude larger than $2dva$. RT

was defined as the time between CP and the first saccade larger than $2dva$. A prosaccade

trial was considered correct if the end position of the saccade was ipsilateral to the cue

and, conversely, an antisaccade trial was considered correct if the end position of the

saccade was contralateral to the cue.

Trials were excluded from further analysis if a) data were missing, b) a blink occurred

between CP and the main saccade, c) the trial was aborted by the experimenter, d)

subjects failed to fixate in the interval between fixation detection and CP, e) if a saccade

was detected only later than 800ms after CP, f) if the RT was below 50ms, and in the case

of an antisaccade if it was below 110ms. Corrective antisaccades were defined as saccades

that a) followed a prosaccade error, b) occurred no later than $900ms$ after CP, and c) had

less than $3dva$ horizontal error from the red circle contralateral to the cue.

Besides the fitted non-decision time δ we assumed a fixed non-decision time of $50ms$ for

all participants [17]. This was implemented by subtracting $50ms$ of all saccades before

being entered into the model. In order to avoid numerical instabilities, RT were rescaled

from millisecond to tenths of a second during all numerical analysis. All results are

presented in ms.

Modeling

We aimed to answer three questions with the models considered here. First, we

investigated which model (i.e. PROSA, SERIA or SERIA$_{lr}$) explained the experimental data

best, and whether all important qualitative features of the data were captured by this

model. We did not have a strong hypothesis regarding the parametric distribution of the

data and hence, comparisons of parametric distributions were only of secondary interest

in our analysis. Second, we investigated whether reduced models that kept certain

parameters fixed across trial types were sufficient to model the data. Third, we

investigated how the probability of a trial type in a block affected the parameters of the

model.

Model space

Initially, we considered 15 different models as shown in Table 2. Each model was fitted

independently for each subject and condition. Since our experimental design included

mixed blocks, we allowed for different parameters in pro- and antisaccade trials, i.e.,

different increase-rate distributions depending on the *trial type* (TT). Under this

hypothesis, the PROSA model had 12 free parameters (6 for each TT), whereas the SERIA

model required 4 further parameters (π_e and π_l in each TT). The late race SERIA$_{lr}$ model

included 16 parameters for the units (8 for each TT). We did not investigate the case that

early reactions could trigger antisaccades but rather fixed the probability of an early

antisaccade $1 - \pi_e$ to 10^{-3}. The rationale behind this was that if early reactions are a

priori assumed to never trigger antisaccades, rare but possible early antisaccades might

cause large biases when fitting a model.

Regarding the non-decision time δ, antisaccade cost δ_a, and rate of outliers η, we assumed

equal parameters in both TT. Consequently, the full PROSA model had 15 free parameters

whereas the full SERIA and SERIA$_{lr}$ models had both 19 free parameters.

Table 2. Model families with the respective increase-rate distributions.

Model	PROSA		
	Prosaccade/ stop units	Anti. unit	# Param. full/const.
m_1/m_1^c	Inv. gamma	Inv. gamma	15/13
m_2/m_2^c	Gamma	Gamma	15/13
m_3/m_3^c	Gamma	Inv. gamma	15/13
m_4/m_4^c	Lognorm.	Lognorm.	15/13
m_5/m_5^c	T. norm.	T. norm.	15/13
	SERIA		
	Early/stop units	**Late unit**	
m_6/m_6^c	Inv. gamma	Inv. gamma	19/13
m_7/m_7^c	Gamma	Gamma	19/13
m_8/m_8^c	Gamma	Inv. gamma	19/13
m_9/m_9^c	Lognorm.	Lognorm.	19/13
m_{10}/m_{10}^c	T. norm.	T. norm.	19/13
	SERIA$_{lr}$		
	Early/stop units	**Late pro./anti. units**	
m_{11}/m_{11}^c	Inv. gamma	Inv. gamma	19/15
m_{12}/m_{12}^c	Gamma	Gamma	19/15
m_{13}/m_{13}^c	Gamma	Inv. gamma	19/15
m_{14}/m_{14}^c	Lognorm.	Lognorm.	19/15
m_{15}/m_{15}^c	T. norm.	T. norm.	19/15

Models with parameters constrained to be equal across trial types are referred through the superscript c.

In addition to the full models, we evaluated restricted versions of each of them by
constraining some parameters to be equal across TT. In the case of the SERIA model, we
hypothesized that the parameters of all units were equal, irrespective of TT (i.e., that the
rate of the units was not affected by the cue presented in a trial). However, we assumed
that the probability that an early or late response was a prosaccade was different in pro-
and antisaccade trials. Therefore, in the case of the SERIA model, instead of 12 unit
parameters (6 per TT), the restricted model had only 6 parameters for the units' rates.
The parameters π_e and π_l were allowed to differ in pro and antisaccade trials. In the case
of the restricted SERIA$_{lr}$ model, the units that underlie the late decision process were
allowed to vary across TT, yielding a restricted model with 4 parameters for the early and
inhibitory units, and 8 for the late decision process, half of them for each trial type. In the

case of the PROSA model, similarly to [17], it is possible to assume that the parameters of

the prosaccade unit remain constant across TT, and that the parameters of the stop and

antisaccade units depend on TT, yielding 10 parameters for the units.

Prior distributions for model parameters

To complete the definition of the models, the prior distribution of the parameters was

specified. This distribution reflects beliefs that are independent of the data and provides

a form of regularization when inverting a model. In order to avoid any undesired bias

regarding the parametric distributions considered here, we reparametrized all but the

truncated normal distribution in terms of their mean and variance. We then assumed that

the log of the mean and variance of the rate of the units were equally normally distributed

(see Table 3). Therefore, the parametric distributions had the same prior in terms of their

first two central moments. In the case of the truncated normal distribution, instead of an

analytical transformation between its first two moments and its natural parameters μ

and σ^2, we defined the prior distribution as a density of μ and $\ln \sigma^2$. To ensure that μ was

positive with high likelihood (96%) we assumed that $\mu \sim N(0.55, 0.09)$. The variance term

was distributed as displayed in Table 3. As a further constraint, we restricted the

parameter space to enforce that the first two moments of the distributions of rates and

RTs existed. We relaxed this constraint for the late units of the SERIA$_{lr}$ in order to allow

for 'flat' distributions with possibly infinite mean and variance. This can describe a case

in which the increase rate of one of the late units is extremely low.

Table 3. Prior probability density functions.

Parameter	Probability density function	Expected value	Variance
μ_r	$\mathcal{N}(\ln \mu_r\,; -1.08, 0.97)$	0.55	0.5
σ_r^2	$\mathcal{N}(\ln \sigma_r^2\,; -2.64, 0.69)$	0.1	0.01
δ	$\mathcal{N}(\ln \delta\,; -1.58, 1.79)$	0.5	1.25
δ_a	$\mathcal{N}(\ln \delta_a\,; -0.87, 1.17)$	0.75	1.25
π_e	$Beta(\pi_e; 0.5, 0.5)$	0.5	0.145
π_l	$Beta(\pi_e; 0.5, 0.5)$	0.5	0.145
η	$Beta(\eta; 0.5, 0.5)$	0.5	0.145

For the non-decision time δ and the antisaccade cost δ_a, the prior of their log transform
was a normal distribution, consistent across all models. Note that the scale of the
parameters δ and $_a$ in Table 3 is tenths of a second. The fraction of early outliers η, and
early and late prosaccades π_e and π_l were assumed to be Beta distributed, with
parameters 0.5 and 0.5. Thus, for example, the prior probability of an early outlier is given
by

$$p(\eta) \propto \eta^{0.5}(1-\eta)^{0.5}. \tag{34}$$

427 This parametrization constitutes the minimally informative prior distribution, as it is the
428 Jeffrey's prior of η, π_e and π_l. Table 3 displays the parameters used for the prior
429 distributions.

Bayesian inference

Inference on the model parameters was performed using the Metropolis-Hastings
algorithm [31]. To increase the efficiency of this sampling scheme, we iteratively modified
the proposal distribution during an initial 'burn-in' phase as proposed by [32]. Moreover,
we extended this method by drawing from a set of chains at different temperatures and
swapping samples across chains. This method, called population MCMC or parallel
tempering, increases the statistical efficiency of the Metropolis-Hasting algorithm [33]
and has been used in similar contexts before [34]. We simulated 16 chains with a 5-th
order temperature schedule [35]. For all but the models including a truncated normal
distribution, we drew a total of 4.1×10^4 samples per chain, from which the first 1.6×10^4

samples were discarded as part of the burn-in phase. When a truncated normal distribution was included (models m_5, m_{10}, and m_{15}), the total number of samples was increased to 6×10^4, from which 2×10^4 were discarded. The convergence of the algorithm was assessed using the Gelman-Rubin criterion [33,36] such that the $\tilde{R}$ statistic of the parameters of the model was aimed to be below 1.1. When a simulation did not satisfy this criterion, it was repeated until 99.5 percent of all inversions satisfied it.

Models were scored using their log marginal likelihood or log model evidence (LME). This is defined as the log probability of the data given a model after marginalizing out all its parameters. When comparing different models, the LME corresponds to the log posterior probability of a model under a uniform prior on model identity. Thus, for a single subject with data y, the posterior probability of model k, given models 1 to n is

$$p(m_k|y) = \frac{p(y|m_k)p(m_k)}{\sum_{i=1}^{n} p(y|m_i)\,p(m_i)} = \frac{p(y|m_k)}{\sum_{i=1}^{n} p(y|m_i)}. \tag{35}$$

Importantly, this method takes into account not only the accuracy of the model but also its complexity, such that overparameterized models are penalized [37]. A widely used approximation to the LME the Bayesian Information Criterion (BIC) which, although easy to compute, has limitations (for discussion, see [38]). Here, we computed the LME through sampling using thermodynamic integration [33,39]. This method provides robust estimates and can be easily computed using samples obtained through population MCMC.

One important observation here is that the LME is sensitive to the prior distribution, and thus can be strongly influenced by it [40]. We addressed this issue in two ways: On one hand and as mentioned above, we defined the prior distribution of the increase rates of all models in terms of the same mean and variance. This implies that the priors were equal up to their first two moments, and hence all models were similarly calibrated. On the other hand, we complemented our quantitative analysis with qualitative posterior checks [33] as shown in the results section.

Besides comparing the evidence of each model, we also performed a hierarchical or random effects analysis described in [38,41]. This method can be understood as a form of soft clustering in which each subject is assigned to a model using the LME as assignment criterion. Here, we report the expected probability of the model r_i, which

469 represents the percentage of subjects that is assigned to the cluster representing model
470 *i*. This hierarchical approach is robust to population heterogeneity and outliers, and
471 complements reporting the group-level LME. Finally, we compared families of models
472 [42] based on the evidence of each model for each subject summed across conditions.

Classical statistics

In addition to a Bayesian analysis of the data, we used classical statistics to investigate
the effect of our experimental manipulation on behavioral variables (mean RT and ER)
and the parameters of the models. We have suggested previously [11,43,44] that
generative models can be used to extract hidden features from experimental data that
might not be directly captured by, for example, standard linear methods or purely data
driven machine learning techniques. In this sense, classical statistical inference can be
boosted by extracting interpretable data features through Bayesian techniques.

Frequentist analyses of RT, ER, and parameter estimates were performed using a mixed
effects generalized linear model with independent variables *subject* (SUBJECT),
prosaccade probability (PP) with levels PP20, PP50 and PP80, and when pro- and
antisaccade trials were analyzed together, *trial type* (TT). The factor SUBJECT was always
entered as a random effect, whereas PP and TT were treated as categorical fixed effects.
In the case of ER, we used the probit function as link function.

Analyses were conducted with the function *fitglme.m* in MATLAB 9.0. The significance
threshold α was set to 0.05.

Implementation

All likelihood functions were implemented in the *C* programming language using the GSL
numerical package (v.1.16). Integrals without an analytical form or well-known
approximations were computed through numerical integration using the Gauss-Kronrod-
Patterson algorithm [45] implemented in the function *gsl_integration_qng*. The sampling
routine was implemented in MATLAB (v. 8.1) and is available as a module of the open
source software package TAPAS (www.translationalneuromodeling.org/tapas).

Results

Behavior

Forty-seven subjects (age: 23.8 ± 2.9) completed all blocks and were included in further analyses. A total of 27072 trials were recorded, from which 569 trials (2%) were excluded (see Table 4).

Table 4. <u>Summary of trials per subject</u>

	Valid	Blink	Missing	Aborted	FE	Late S.	Early S.	Total
Total	26503	188	60	42	249	0	30	27072
Mean	563.9	4.0	1.3	0.9	5.3	0.0	0.6	576
Std.	9.9	5.1	2.5	1.5	5.0	0.0	1.3	-
Min.	536	0	0	0	0	0	0	-
Max.	576	22	15	6	19	0	8	-

FE: Fixation errors. Late saccades are saccades elicited after 800ms. Early saccades are prosaccades elicited before 50ms after CP or antisaccades elicited before 110ms after CP.

Both ER and RT showed a strong dependence on PP (Fig 5 and Table 5). Individual data is included in the S1 File and is displayed in S2 Figure. The mean RT of correct pro- and antisaccade trials was analyzed independently with two ANOVA tests with factors SUBJECT and PP. We found that in both pro- ($F_{2,138} = 46.9, p < 10^{-5}$) and antisaccade trials ($F_{2,138} = 37.3, p < 10^{-5}$) the effect of PP was significant; with higher PP, prosaccade RT decreased, whereas the RT of correct antisaccades increased. On a subject-by-subject basis, we found that between the PP20 and PP80 conditions, 91% of the participants showed increased RT in correct antisaccade trials, while 81% demonstrated the opposite effect (a decrease in RT) in correct prosaccade trials. Similarly, there was a significant effect of PP on ER in both prosaccade ($F_{2,138} = 376.1, p < 10^{-5}$) as well as in antisaccade ($F_{2,138} = 347.0, p < 10^{-5}$) trials. This effect was present in all but one participant in antisaccade trials and in all subjects in prosaccade trials. Exemplary RT data of one subject in the PP50 condition is displayed in Fig 6.

Fig 5. Error rate and reaction times as a function of prosaccade trial probability (PP).

Left panel: Mean error rates for pro and antisaccades. Right panel: Mean reaction time in *ms*. Error bars indicate standard errors of the mean. Only correct responses are displayed.

Table 5. Summary of mean RTs and ERs.

Trial type	Action	Reaction times [ms]		
		PP 20	PP 50	PP80
Pro.	Pro.	330(72)	319(67)	284(59)
Pro.	Anti.	326(68)	329(46)	336(57)
Anti.	Anti	354(60)	378(57)	389(61)
Anti.	Pro	234(50)	231(47)	225(31)
		Error rates [%]		
Pro.		26(15)	11(8)	4(4)
Anti.		23(17)	35(21)	51(20)

Standard deviations are shown in brackets.

Fig 6. Exemplary RT histogram of one subject in the PP50 condition.

Prosaccade trials are displayed in the upper half plane and antisaccade trials in the lower (negative) half plane. Prosaccade actions are depicted in red color, whereas antisaccade actions are shown in blue. Errors in prosaccade trials are antisaccades that for this subject occurred after the first peak of early prosaccades. Errors in antisaccade trials (lower half plane) occurred at a similar latency as early prosaccades in prosaccade trials. The histograms have been normalized to have unit probability mass, i.e., the sum of the area of all bars is one.

Modeling

Model comparison results

Initially, we considered the models outlined in Table 2. The LME over all participants (fixed effects analysis) and the posterior probability of all models and all subjects are presented in Fig 7. Independently of the particular parametric distribution of the units, the SERIA$_{lr}$ models had higher evidence compared to the PROSA and SERIA models. A random effects, family-wise model comparison [42] resulted in an expected frequency of $r = 87\%$ for the SERIA$_{lr}$ family, $r = 11\%$ for the SERIA family, and $r=2\%$ for the PROSA family. In addition, constraining the parameters to be equal across trial types increased the model evidence irrespective of the parametric distribution assigned to the units (Fig 7). Here, the family-wise model comparison showed that models with constrained parameters had an expected frequency of $r = 98\%$. Over all 30 models, m_{13}^c (SERIA$_{lr}$ with constrained parameters, early and inhibitory increase rates gamma distributed, and late units' rate inverse gamma distributed) showed the highest LME with $\Delta LME > 200$ compared to all other models. Following [40], a difference in LME larger than 3 corresponds to strong evidence.

To verify that the SERIA_lr family was not preferred simply because the probability of early

prosaccades was fixed, we considered models in the SERIA family with the same property

(not displayed). We found that although fixing this value increased the LME of the SERIA

family, there was still a difference of $\Delta LME > 90$ when comparing the best model of the

SERIA_lr family and the best model of the SERIA family with a fix probability of early

prosaccades.

Fig 7. Summary of model comparison.

Top: Summed LME of all subjects for all 30 models. White bars show models with all parameters free, grey bars models with constrained parameters. LMEs are normalized by subtracting the lowest LME (m_5). Model m_{13}^c (constrained SERIA_lr) exceeded all other models (ΔLME>200). Bottom: Illustration of model probability for all subjects. The posterior model probabilities for all subjects are shown as black dots. In white shading are models with all parameters free, grey bars represent models with restricted parameters. Note that in nearly all subjects, the SERIA_lr models with restricted parameters showed high model probabilities.

Fits of four subjects using the posterior samples of the best PROSA (m_1), SERIA (m_8^c), and

SERIA_lr (m_{13}^c) models are depicted in Fig 8. Although model m_1 was the best model in the

PROSA family, it clearly did not explain the apparent bimodality of the prosaccade RT

distributions. Instead, RTs were explained through wider distributions. No obvious

difference could be observed between the SERIA and SERIA_lr models. We further

examined the model fits in Fig 9 and Fig 10 by plotting the weighted fits and cumulative

density functions of the reciprocal RT in the probit scale (reciprobit plot [22]) collapsed

across subjects for the best model of each family. The histograms of RTs clearly show a

large number of late prosaccades whose distribution is similar to the distribution of

antisaccade RTs. The most pronounced – but still small - difference between the SERIA

and SERIA$_{lr}$ models was visible in prosaccade trials in the PP20 condition (left panel,

upper half plane), in which antisaccade errors displayed lower RT than correct late

prosaccades.

Fig 8. Fits of best PROSA (m_1), SERIA (m_8^c) and SERIA$_{lr}$ (m_{13}^c) models

Columns display the normalized histogram of the RTs of pro- (red) and antisaccades (blue) in each of the conditions. Rows correspond to individual subjects named S1 to S4 for display purpose. As in Fig. 6, prosaccade trials are displayed on the upper half plane, whereas antisaccade trials are displayed in the lower half plane. The predicted RT distributions based on the samples from the posterior distribution are displayed in solid (SERIA$_{lr}$), broken (SERIA), and dash-dotted (PROSA) lines. Note that data from subject 3 in the PP50 condition is the same as shown in Fig. 6. Early outliers are not displayed.

Fig 9. Fits from the best models in each family (m_1, m_8^c, m_{13}^c).

Model fits and RT histograms for each condition collapsed across subjects. For more details see Fig. 6 and 8.

Fig 10. Reciprobit of best models.

Predicted and empirical cumulative density function of the reciprocal RT in the probit scale for each condition and model collapsed across all subjects. The data shown are the same as in Fig 9, but split for trial types and illustrated as cumulative distributions. Note that the y-axis is in the probit scale and that nearly all differences between the model and the data occur at very small probability values of 5% or below.

554   *Corrective antisaccades*

555   The RTs of antisaccades that follow an error prosaccade were not directly modeled.

556   However, we hypothesized that corrective antisaccades are delayed late antisaccade

557   actions, whose distribution is given by the *response time* distribution of late antisaccades

$$\frac{1}{p(U_a < U_p)} \int_0^\infty p(U_a = t)p(U_p > t)dt. \tag{36}$$

A total of 2989 corrective antisaccades were included in the analysis. The mean (±std)

end time of the erroneous prosaccades was 268(±63)ms. The mean RT of corrective

antisaccades was 447(±103)ms, and the weighted mean arrival time of the late

antisaccade unit was 367ms. Fig 11 displays the histogram of the end time of all

prosaccade errors, the RT of all corrective antisaccades and the time shifted (+80ms)

predicted response time of late antisaccades. Since we did not have a strong hypothesis

regarding the magnitude of the delay of the corrective antisaccades, we selected the time

shift to be the difference between the mean corrective antisaccade RT and the mean

predicted response time of late antisaccades. Visual inspection strongly suggests that the

distribution of corrective antisaccade RTs is well approximated by the distribution of the

late responses. The short difference between corrective antisaccades' RT and the

expected response time of the late antisaccade unit (80ms) favors the hypothesis that the

plan for a corrective antisaccade is initiated before the incorrect prosaccade had finished.

Fig 11. Empirical and predicted RT of corrective antisaccades.

Left: End time of erroneous prosaccades, RTs of corrective antisaccades, and time shifted predicted response time distribution of late antisaccades. The time shift was selected to be the difference between the empirical and predicted mean response time. Center: Quantile-quantile plot of the predicted and empirical distribution of corrective antisaccades, and a linear fit to the central 98% quantiles. There is a small deviation only at the tail of the distribution. Right: Reciprobit plot of the empirical and predicted cumulative density functions of the RT of corrective antisaccades. The scale of the horizontal axis is proportional to the reciprocal RT. The vertical axis is in the probit scale.

571 *Effects of prosaccade probability on model parameters*

572 The effect of PP on the parameters of the model was investigated by examining the

573 expected value of the parameters of the best scoring model (m_{13}^c). Initially, we considered

574 the question of whether the mean arrival or response time of each of the units changed

575 as a function of PP. For arrival times, this corresponds to

$$\frac{1}{N}\sum_{j=1}^{N}\left(E\left[U_i\middle|k_j^i,\theta_j^i\right]+\delta_j^i\right) \tag{37}$$

where i is an index over the units, j is an index over N samples collected using MCMC, and

δ_j^i is the estimated delay. In the case of the late units, we considered only the response

time of correct actions. Fig 12 left displays the mean arrival and response times. These

were submitted to four separate ANOVA tests, which revealed that PP had a significant

effect on all four units: early unit ($F_{2,138} = 9.2, p < 10^{-3}$), late antisaccade ($F_{2,138} =$

$26.6, p < 10^{-3}$), late prosaccade ($F_{2,138} = 19.6, p < 10^{-3}$), and inhibitory unit ($F_{2,138} =$

$30.9, p < 10^{-3}$). We then explored the differences across conditions through planned

post hoc tests on each condition for each of the units (Table 6). The arrival times of the

early unit did not change significantly between condition PP20 and PP50, but decreased

significantly in the PP80 condition as compared to the PP50 block. The response times of

late antisaccades increased significantly between the PP20 and the PP50 conditions but

not so between the PP50 and PP80. Late prosaccades followed the opposite pattern,

showing only a significant decrease in response time between the PP50 and PP80

conditions. Finally, the inhibitory unit changed significantly across all conditions.

Fig 12. Model parameters.

Left: Mean arrival or response time and standard error of the early and inhibition units and late pro- and antisaccades. Right: Probability of a late antisaccade $p(U_a > U_p)$ in prosaccade (red) and antisaccade (blue) trials in each condition in the probit scale.

	Early			Inhib.			Late pro.			Late anti.		
	Mean [ms]	t_{138}	p	Mean [ms]	t_{138}	p	Mean [ms]	t_{138}	p	Mean [ms]	t_{138}	p
PP20-PP50	4	0.6	0.50	-30	-4.0	<0.001*	3	0.5	0.55	-24	-36	<0.001*
PP20–PP80	26	3.9	<0.001*	-58	-7.8	<0.001*	32	5.7	<0.001*	-32	-5.4	<0.001*
PP50–PP80	22	3.3	0.001*	-28	-3.8	<0.001*	29	5.1	<0.001*	-7	-1.5	0.13

Effect of PP on the mean arrival time for each of the early, inhibitory units, and the late pro and antisaccade units in the corresponding trial type.

Finally, we examined how the probability of a late antisaccade $p(U_a > U_p)$ (Fig 12, right) depended on PP and TT. The estimated parameters for both pro- and antisaccade trials were analyzed with a model with factors SUBJECT, TT, PP and the interaction between TT and PP. An ANOVA test demonstrated that both PP ($F_{2,276} = 51.2, p < 10^{-3}$) and TT ($F_{1,276} = 985.0, p < 10^{-3}$) had a significant effect, but there was no evidence for an interaction between the two factors ($F_{2,276} = 1.5, p = 0.23$), suggesting that PP affected the probability of a late antisaccade equally in pro- and antisaccade trials.

Subject specific parameters

Finally, we investigated how some of the parameters of the model were related to each other across subjects. Because it has been commonly reported that schizophrenia is related with higher ER, but also with increased antisaccade RT, an interesting question is whether higher late-action response times are correlated with the percentage of late errors and inhibition failures, i.e., early saccades that are not stopped. We found that the response time of late pro ($F_{1,135} = 13.6, p < 0.001$) and antisaccades ($F_{1,135} = 7.1, p < 0.01$) was negatively correlated with the probability of a late error (Fig 13), but no significant interaction between PP and response time was found (pro: $F_{2,135} = 1.7, p = 0.19$; anti $F_{2,135} = 0.3, p = 0.76$). Hence, late responders tended to make fewer late errors, suggesting a speed/accuracy trade-off in addition to the main effect of PP. We further considered the question whether the percentage of inhibition failures was correlated with the expected arrival time of the late antisaccade unit in antisaccade trials (Fig. 13 right). Note that the number of inhibition failures is the same in both trial types

612 in a constrained model, but inhibition failures are errors in antisaccade trials and correct

613 early reactions in prosaccade trials. We found that these parameters were not

614 significantly correlated $(F_{2,135} = 1.2, p = 0.26)$. This was also the case when we

615 considered the expected response time of late prosaccades in prosaccade trials (not

616 displayed; $F_{2,135} = 0.0, p = 0.98$).

Fig 13. Correlation between late arrival times and errors

Left: Percentage of late errors against late antisaccades' response times in antisaccade trials. Center: Percentage of late errors against late prosaccades' response time in prosaccade trials. Left: Percentage of inhibitory failures against late antisaccades' response time in antisaccade trials. The horizontal axis is in the probit scale.

Fig 14 illustrates the posterior distribution of late errors and inhibition failures of two

representative subjects as estimated using MCMC. Clearly, PP induced strong differences

in the percentage of inhibition failures and late errors in prosaccade trials in both

subjects. The effect of PP is less pronounced in late errors in antisaccade trials. The

posterior distributions also illustrate how the SERIA_lr model can capture individual

differences: For example, the percentage of late prosaccade errors in the PP80 condition

and the percentage of inhibition failures across all conditions are clearly different in each

subject.

Fig 14. Posterior distribution of late errors and inhibition failures

The posterior distribution of the percentage of late and inhibitory failures of two exemplary subjects (see Fig 8 and 13). Samples from the posterior distribution were obtained using MCMC. Histograms display the distributions of the samples in in the probit scale (horizontal axis). For these two subjects, the posterior distribution of late prosaccade and inhibitory failures clearly discriminates between the three PP conditions.

Discussion

In this study, we provided a formal treatment of error rates (ER) and reaction times (RT) in the antisaccade task using probabilistic models. We applied these models to data from an experiment consisting of 3 mixed blocks with different probabilities of pro- and antisaccades trials. Model comparison showed that a novel model that allows for late pro- and antisaccades explains our experimental findings better than a model in which all late responses are assumed to be antisaccades. The parameter estimates of the hidden units of the model showed that changes in the inhibition unit and a late decision process explained most of the overt changes in behavior caused by our experimental manipulation, i.e., differences in trial type probability. Moreover, we found that all units were sensitive to the PP in a block, although late responses tended to plateau when the corresponding trial type was not highly frequent.

Our main finding is that two decision processes are necessary to properly model the antisaccade task: on one hand, an early race between a prepotent response towards a target and an endogenously generated signal to cancel this action, and, on the other hand, a secondary late race between two units encoding the cue-action mapping. Although the late decision process can be closely approximated by assuming that RT and actions are independent (at least in our experimental design), Bayesian model comparison demonstrated that late decisions are more accurately described by a race between two units representing different actions. The two decision processes are the sources of early errors –fast prosaccades in antisaccade trials- and late errors –late actions incongruent with the cue presented. The late decision process displays a speed/accuracy tradeoff and is biased by the probability of a trial type in a block. Moreover, this decision process predicts the RT distribution of corrective antisaccades that follow early errors. Because the extra latency of these corrective antisaccades (80ms) is much lower than the average time by which the erroneous prosaccade is terminated, it is unlikely that corrective antisaccades are due to a restart in the decision process, but rather these are late actions that overwrite early errors.

Influence of trial type probability on reaction times and error rates

Our results show that both RT and ER depend on PP. While this was a highly significant factor in our study, there are mixed findings in previous reports. ER in antisaccade trials was found to be correlated with TT probability in several studies [29,46,47]. However,

this effect might depend on the exact implementation of the task [47,48]. Changes in

prosaccade ER similar to our study have been reported by [29] and [48]. Studies in which

the type of saccade was signaled at fixation prior to the presentation of the peripheral cue

do not always show this effect [47]. The results on RTs are less consistent in the literature.

Our findings of increased anti- and decreased prosaccade RTs with higher PP are in line

with the overall trend in [29], and with studies in which the cue was presented centrally

[29,47]. Often, there is an additional increase in RT in the PP50 condition [29,47], which

was visible in our data as a slight increase in RT in the PP50 condition on top of the linear

effect of PP. Overall, RTs in our study were relatively slow compared to studies in which

the TT cue was separated from the spatial cue [46,47]. However, a study with a similar

design and added visual search reported even slower RTs in both pro- and antisaccades

[29].

Interpretation of model comparison results

Formal comparison of generative models can offer insight into the mechanisms

underlying eye movement behavior [11] and might be relevant in translational

neuromodeling applications, such as computational psychiatry [49-53]. Here, we have

presented what is, to our knowledge, the first formal statistical comparison of models of

the antisaccade task. For this, we formalized the model introduced in [17] and proceeded

to develop a novel model that relaxes the one-to-one association of early and late

responses with pro- and antisaccades, respectively. All models and estimation techniques

presented here are openly available under the GPLv3.0 license as part of the open source

package TAPAS (www.translationalneuromodeling.org/tapas).

Bayesian model comparison yielded four conclusions at the family level. First, the SERIA

models were clearly favored when compared to the PROSA models. Second, including a

late race between actions representing late pro- and antisaccades (SERIA$_{lr}$) resulted in an

increase in model evidence, compared to a model not including a late race (SERIA). Third,

models in which the race parameters of the early and inhibitory unit were constrained to

be equal across TT had a higher LME than models in which all parameters were free.

Hence, the effect of the cue in a single trial was limited to the late action, and did not affect

the race between an early and inhibitory process. This constitutes an important external

validation, as it means that model comparison does favor a model which respects the

temporal order of the experiment: Information about TT is only available after the

stimulus was presented and, thus, it is unlikely to have an impact of the fast reactive

responses. Fourth, early responses were nearly always prosaccades. Crucially, these four

conclusions are based on family-wise comparison across all parametric distribution of

the increase rate of the units.

A further consequence of our findings is that two independent and qualitatively different

decision processes lead to an antisaccade: the race process between early and inhibitory

units, and the secondary decision process that generates late responses. A separation of

decisions into a 'where' and a 'when' component has been proposed by [54], but mainly

in conceptual terms. However, model comparison showed that these two components

('where' and 'when') cannot be completely dissociated and that time plays a role in the

late decision. Nevertheless, the assumption that action type and arrival time of late

responses were independent yielded a good fit to this particular data set, suggesting that

it is, in many cases, an acceptable approximation to assume a time-independent late

decision process. The most obvious difference between the SERIA and SERIA$_{lr}$ can be

observed in prosaccade trials in the PP20 condition (left panel, upper half plane Fig. 9),

in which late prosaccades are slower than antisaccades. We discuss this point in more

detail later.

Parametric distribution of reaction times

The parametric distribution of oculomotor RTs has been discussed in great detail in the

literature (e.g., [13,55]). Here, we did not aim at determining the most suitable

distribution, but rather opted for a practical approach by evaluating different models

with a reduced number of parametric distributions. We then based our conclusions on

the model with the highest LME. Nevertheless, one can consider the connection of the

models presented here with other families of parametric distributions. In particular, the

linear relationship

$$\frac{s_i}{r_i} = t \tag{38}$$

could be seen as to be formally inconsistent with the observation that RT are likely to be

explained by stochastic accumulation processes (see for example [56,57], but [58]). This

is a weaker constraint than one would expect, because under low noise conditions, for

example, the linear relationship can be a good approximation of neural activity. Even if

the relationship is not linear, for any continuous function ϕ with an inverse function ϕ^{-1},
the model can be recasted as [59]:

$$s_i = \phi(tr_i), \tag{39}$$

$$\phi^{-1}(s_i)/r_i = t. \tag{40}$$

721 In any case, linear accumulation models have been shown to yield similar conclusions to
722 stochastic accumulation models [58].

723 More generally, it can be shown that if RTs follow a generalized inverse normal
724 distribution (GIN) of the form

$$GIN(t; \lambda, \kappa, \psi) = \frac{(\psi/\kappa)^\lambda}{2K_\lambda(\sqrt{\kappa\psi})} t^{\lambda-1} \exp\left(-\frac{1}{2}(\kappa t^{-1} + \psi t)\right) \tag{41}$$

725 where $\lambda \leq 0$, and K_λ is a modified Bessel function of the second kind, there exists a
726 continuous diffusion process whose first hit distribution (FHD) follows the GIN [60]. A
727 particular case of this distribution is the Wald distribution for which $\lambda = -\frac{1}{2}, \kappa = 0$. It is
728 the FHD of the Brownian diffusion process with drift

$$X_t = -\sqrt{\sigma}\psi t + \sigma W_t \tag{42}$$

729 where W_t denotes a Wiener process, $x_0 > 0$, and the absorbing boundary a is zero. More
730 relevant here, when $\psi = 0$ the distribution reduces to an inverse gamma distribution, the
731 FHD of the process

$$X_t = \sqrt{\sigma}(2\lambda - 1) t^{-1} + \sigma W_t \tag{43}$$

with $x_0 > 0$ and boundary $a = 0$ (for a detailed mathematical treatment see [60]). Thus,
if the rates of a ballistic, linear processes are assumed to be gamma distributed, the RTs
follow a distribution that is formally equivalent to a first hit model with stochastic
updates and fixed rates. While the model presented here can be seen as a ballistic
accumulation model, this equivalence suggests that it is *compatible* with a diffusion
process with infinitesimal mean change proportional to t^{-1}.

Other antisaccade models

In broad terms, three families of antisaccade models can be distinguished (reviewed in

[61]). The first set of models is based on a race process with independent saccadic and

stop units. These models build on the seminal work of [16] on the stop-signal paradigm.

According to this model, a 'GO' signal triggers a stochastic 'race' process that generates a

response once it reaches threshold. Critically, a stop signal triggers a second process that

inhibits the first 'GO' response if it is the first to reach threshold. Importantly, the pace of

both units is mutually independent. This model was further extended for the antisaccade

task by [17] (but see [14,21], and the review in [20]), who included a third unit such that

an antisaccade is generated when a reflexive prosaccade is inhibited by an endogenously-

triggered stop process. Note that the original 'horse-race' model has also been modified

[62] to account for different competing response actions, similarly as in the antisaccade

task. The models proposed here belong to this family.

A second type of model relies on lateral or mutual inhibition of competing pro- and

antisaccade units. In this direction, Cutsuridis and colleagues [61,63,64] proposed that

lateral inhibition is implemented by inhibitory connections in the intermediate layers of

the superior colliculus. Thus, saccades are the result of accumulation processes, but these

are not independent of each other. Crucially, no veto-like stop signal is required. Although

no formal model-fitting has been proposed for this model, qualitative agreement with

data suggests that it might capture behavioral patterns relevant in translational

applications [64,65]. Since no probabilistic version of this model is available, it is not yet

possible to decide on the grounds of model comparison whether mutually dependent or

independent race processes best explain current behavioral findings.

Finally, several models that incorporate detailed physiological mechanisms have been

proposed [23,66-68]. These models cannot be easily assigned to one of the above

categories, as they often employ both an inhibitory mechanism that stops or withholds

the reactive responses as well as competition between actions. In addition, while more

realistic models possess a more fine-grained representation of the underlying

neurobiology, they rely on a large number of parameters and for this reason, it is difficult

to fit them to behavioral data (for discussion, see [11]).

Regarding neurobiologically realistic models, the model proposed by [23] is the most

similar to the SERIA model. It posits two different mechanisms that interact in the

generation of antisaccades: an action-selection module and a remapping module that controls the cue-action mapping. As a consequence, this model allows for the generation of late errors that follow a similar RT distribution as correct antisaccades. Consistent with this observation, the SERIA model can quantitatively distinguish between inhibition and late cue-action mapping errors (Fig 15, left panel). A less obvious similarity between the SERIA model and [23] is that different cues do not lead *directly* to different dynamics in the action module, but only in the so-called 'remapping' module. Furthermore, the incorporation of a late race is conceptually close to the approach proposed by [23], which includes a winner take all competition in what we have referred here as late responses. Similarly, our model comparison results show that different cues (i.e., trial types) do not affect the GO/NO-GO process but only the late cue-action mapping.

Fig 15. Error sources and the correlation of response times

Left: Error rate (black line) split into the two causes predicted by the model. Inhibition errors are early actions that always trigger prosaccades. Similarly as described by [23], late errors occur when a late response leads to a prosaccade. Right: Correlation between correct antisaccade and late prosaccades response times according to the best SERIA$_{lc}$ model. The best linear fit is depicted as a solid line. The mean ratio of pro- and antisaccade response times (s) is displayed on the right. Although late pro – and antisaccade response times are highly correlated, their ratio is different in each condition (interaction PP and late prosaccade response time $F = 9.2, p < 0.001$).

Parameter changes across trial types

One of the most salient results presented here is that models in which the parameters of the units were constrained to be equal across trial types had a larger LME than models in which all the parameters were free, suggesting that the early and inhibitory race units were not affected by the cue presented on a single trial. While visual inspection of the predicted likelihood under the posterior parameters showed that most of the prominent

characteristics of the data were explained correctly, some more subtle effects were not

captured accurately by the SERIA model. This is particularly clear in the PP20 condition,

in which the SERIA model displays a large bias in prosaccades trials in the PP20 condition.

One possible explanation is that restricting the parameters across trial types made the

model too rigid to capture this effect. Fig 16 compares the fitted RT distributions for

models m_8 (SERIA) and m_{13} (SERIA$_{lr}$), in which no constraint on the parameters was

imposed. Both models are qualitatively almost identical, although as shown in Fig 7, the

LME favored the SERIA$_{lr}$ model. Thereby, the qualitative similarity between both models

indicates that, in our experiment, the RT of late decisions is only weakly dependent on

time. In conclusion, although removing the constraint on the parameters did improve the

fit, the differences are marginal and, thus did not justify the additional model complexity.

As mentioned above this is consistent with the notion that the information about trial

type is only available to the subject once the peripheral stimulus (green bar) has been

processed. Presumably tens of milliseconds after the stimulus onset. In fact, this example

illustrates the protection against overfitting provided by the LME, as this is a case in

which simpler models were preferred over more complex models despite of slightly less

accurate fits.

Fig 16: Comparison between unconstrained SERIA and SERIA$_{lr}$ models.

Comparison between models m_8 (broken lines; SERIA model) and m_{13} (solid lines; late race SERIA$_{lr}$ model.).

804   Arguably, the constrained SERIA model fails to fully capture the RT of late prosaccade in

805   the PP20 and PP50 conditions because of the assumption that late prosaccades have the

806   same arrival time as late antisaccades. As shown in Fig 15, although the response time of

807 late pro- and antisaccades are strongly correlated, the average ratio of the response times

808 changes across conditions.

The effect of trial type probability

It is far from obvious why TT probability affects RT and ER in the antisaccade task. One

possible explanation is that increased probability leads to higher preparedness for either

pro- or antisaccades. Such a theory posits an intrinsic trade-off between preparations for

one of the two action types that leads to higher RTs and ERs in low probability trials.

Thus, a trade-off theory predicts that the arrival times of early and late responses should

be negatively correlated. Although this hypothesis can explain our behavioral findings in

terms of summary statistics, our model suggests a more complicated picture.

The main explanation of our results is the effect of TT probability on the inhibitory unit

and the probability of a late prosaccade. A higher probability of antisaccade trials leads

to faster inhibition and to a higher number of late prosaccades. This resulted in higher

mean RT in prosaccade trials when PP is low. In the case of antisaccades, although the

mean arrival times of the late unit increased in the PP50 condition, the increased arrival

time of the inhibitory unit on the PP80 condition skewed the antisaccade distribution

towards higher RTs. Nevertheless, the SERIA$_{lr}$ implies the anticorrelation of late pro- and

antisaccades in a single trial type, as these are the results of a GO-GO race.

Action inhibition

The biological implementation of action inhibition in the antisaccade and other

countermanding tasks has received a lot of attention and is still debated [69-73]. Our

work adds evidence to the theory that the antisaccade task requires a process that

inhibits prepotent responses and is independent of the initiation of a late action [20].

Recent evidence from electrophysiological recordings in the rat brain ([74] reviewed by

[71]) suggests that the hypothesized race between go and inhibitory responses might be

implemented by different pathways in the basal ganglia [68]. In addition to the basal

ganglia, microstimulation of the supplementary eye fields tends to facilitate inhibition of

saccades in the countermanding task [75].

Corrective antisaccades

Although not a primary goal of our model, we considered the question of predicting

corrective antisaccades. This problem has received some attention recently

[18,61,65,76], as more sophisticated models of the antisaccade task have been developed.

We speculated that corrective antisaccades are generated by the same mechanism as late

responses. Thus, their RT distribution should follow a similar distribution. Our results

strongly suggest that this is the case (see Fig 11). Moreover, the time delay of the

corrective antisaccades indicates that, on average, these actions are not the result of the

late unit being restarted at the end time of the erroneous prosaccade, as this would lead

to much higher RTs. Rather, the planning of a corrective antisaccade might be started

much before the end of the execution of an erroneous prosaccade, in accordance with the

parallel planning model of the antisaccade task [46] and the 'GO – STOP+GO' model in

[21].

Translational applications

Despite the large number of studies of clinical patients using the antisaccade task, an

important question remains open: What are the causes of the errors in different

neurological and psychiatric conditions? For example [77,78] argued that errors in

schizophrenia might be explained, at least partially, by a failure to generate a secondary

late action based on several modifications of the antisaccade task. However, it was also

proposed that the increased ER in schizophrenia is due to high tonic dopamine levels in

the basal ganglia, that lead to decreased inhibition of early responses [68]. More

generally, different neurological and psychiatric diseases, or even patients with the same

condition, might be characterized by a different source of errors. For example, there is

intriguing evidence [79] that patients with different diseases such as attention deficits

disorders [80], Parkinons' disease [81], and amyotrophic lateral sclerosis [82] might be

characterized by different ratios of early and late errors. An interesting experimental

finding in our study related to this is the considerable amount of erroneous antisaccades

in prosaccade trials. An increased number of such errors could be caused by reduced

cognitive flexibility leading to impaired shifting between tasks as observed for example

in obsessive compulsive disorder [83]. The ability to quantify different types of errors

through computational modeling might help to further characterize these diseases.

Summary

Here we have presented a novel model of the antisaccade task. While the basic structure

of the model follows the layout of a previous model [17], we have introduced two crucial

advancements. First, we postulated that late responses could trigger both pro- and

antisaccades, which are selected by an independent decision process. Second, the
generative nature of our model allows for Bayesian model inversion, which enables the
comparison of different models and families of models on formal grounds. To our
knowledge this has not been done for any of the previous models of the antisaccade task,
which is of relevance for translational applications that aim at better understanding
psychiatric diseases by means of computational modeling.

The application of the model to a large data set yielded several novel results. First, the
early and inhibitory race processes triggered by different cues are almost identical.
Moreover, different PP had very different effects on the individual units, which was not
obvious from the linear analysis of the mean RT and ER. Crucially, our modeling approach
allowed us to look at a mechanistic explanation or the effects of PP by examining the
individual units. In future work we aim to disentangle the mechanisms of behavioral
differences caused by neuromodulatory drugs and psychiatric illnesses using formal
Bayesian inference.

Acknowledgments

We thank Saee Paliwal for her remarks, and Jeffrey D. Schall and an anonymous reviewer for their helpful comments on an earlier version of this manuscript.

Supporting Information

Supporting information 1

S1_Dataset.zip. Table of Data. Spreadsheet including all reaction times, actions and errors that entered the analysis. More details are included directly in the file.

Supporting information 2

S2: Reaction time and error rate in all conditions for each subject.

Mean reaction times and error rates are displayed as solid lines.

References

1. Hallett PE. Primary and secondary saccades to goals defined by instructions. Vision Res. 1978;18: 1279–1296.

2. Munoz DP, Everling S. Look away: the anti-saccade task and the voluntary control of eye movement. Nat Rev Neurosci. 2004;5: 218–228.

3. Hutton SB, Ettinger U. The antisaccade task as a research tool in psychopathology: a critical review. Psychophysiology. 2006;43: 302–313.

4. Fukushima J, Fukushima K, Chiba T, Tanaka S, Yamashita I, Kato M. Disturbances of voluntary control of saccadic eye movements in schizophrenic patients. Biol Psychiatry. 1988;23: 670–677.

5. Curtis CE, Calkins ME, Grove WM, Feil KJ, Iacono WG. Saccadic disinhibition in patients with acute and remitted schizophrenia and their first-degree biological relatives. Am J Psychiatry. 2001;158: 100–106.

6. Harris MS, Reilly JL, Keshavan MS, Sweeney JA. Longitudinal studies of antisaccades in antipsychotic-naive first-episode schizophrenia. Psychol Med. 2006;36: 485–494.

7. Reilly JL, Frankovich K, Hill S, Gershon ES, Keefe RS, Keshavan MS, et al. Elevated antisaccade error rate as an intermediate phenotype for psychosis across diagnostic categories. Schizophr Bull. 2014;40: 1011–1021.

8. Radant AD, Millard SP, Braff DL, Calkins ME, Dobie DJ, Freedman R, et al. Robust differences in antisaccade performance exist between COGS schizophrenia cases and controls regardless of recruitment strategies. Schizophr Res. 2015;163: 47–52.

9. Crawford TJ, Sharma T, Puri BK, Murray RM, Berridge DM, Lewis SW. Saccadic eye movements in families multiply affected with schizophrenia: the Maudsley Family Study. Am J Psychiatry. 1998;155: 1703–1710.

10. Radant AD, Dobie DJ, Calkins ME, Olincy A, Braff DL, Cadenhead KS, et al. Antisaccade performance in schizophrenia patients, their first-degree biological relatives, and community comparison subjects: data from the COGS study. Psychophysiology. 2010;47: 846–856.

11. Heinzle J, Aponte EA, Stephan KE. Computational models of eye movements and their application to schizophrenia. Current Opinion in Behavioral Sciences. 2016;11: 21–29. doi:http://dx.doi.org/10.1016/j.cobeha.2016.03.008

12. Wolfe JM, Palmer EM, Horowitz TS. Reaction time distributions constrain models of visual search. Vision Res. 2010;50: 1304–1311.

13. Palmer EM, Horowitz TS, Torralba A, Wolfe JM. What are the shapes of response time distributions in visual search? J Exp Psychol Hum Percept Perform. 2011;37: 58–71.

14. Noorani I, Carpenter RH. Full reaction time distributions reveal the complexity of
neural decision-making. Eur J Neurosci. 2011;33: 1948–1951.

15. Huys QJ, Maia TV, Frank MJ. Computational psychiatry as a bridge from
neuroscience to clinical applications. Nat Neurosci. 2016;19: 404–413.

16. Logan GD, Cowan WB, Davis KA. On the ability to inhibit simple and choice
reaction time responses: a model and a method. J Exp Psychol Hum Percept
Perform. 1984;10: 276–291.

17. Noorani I, Carpenter RH. Antisaccades as decisions: LATER model predicts
latency distributions and error responses. Eur J Neurosci. 2013;37: 330–338.

18. Noorani I, Carpenter RH. Re-starting a neural race: anti-saccade correction. Eur J
Neurosci. 2014;39: 159–164.

19. Noorani I. LATER models of neural decision behavior in choice tasks. Front Integr
Neurosci. 2014;8: 67.

20. Noorani I. Towards a unifying mechanism for cancelling movements. Philos Trans
R Soc Lond, B, Biol Sci. 2017;372.

21. Camalier CR, Gotler A, Murthy A, Thompson KG, Logan GD, Palmeri TJ, et al.
Dynamics of saccade target selection: race model analysis of double step and
search step saccade production in human and macaque. Vision Res. 2007;47:
2187–2211.

22. Carpenter RH, Williams ML. Neural computation of log likelihood in control of
saccadic eye movements. Nature. 1995;377: 59–62.

23. Lo CC, Wang XJ. Conflict Resolution as Near-Threshold Decision-Making: A
Spiking Neural Circuit Model with Two-Stage Competition for Antisaccadic Task.
PLoS Comput Biol. 2016;12: e1005081.

24. Brown SD, Heathcote A. The simplest complete model of choice response time:
linear ballistic accumulation. Cogn Psychol. 2008;57: 153–178.

25. Gorea A, Rider D, Yang Q. A unified comparison of stimulus-driven, endogenous
mandatory and "free choice" saccades. PLoS ONE. 2014;9: e88990.

26. Edelman JA, Valenzuela N, Barton JJ. Antisaccade velocity, but not latency, results
from a lack of saccade visual guidance. Vision Res. 2006;46: 1411–1421.

27. Zhang M, Barash S. Neuronal switching of sensorimotor transformations for
antisaccades. Nature. 2000;408: 971–975.

28. Sato TR, Schall JD. Effects of stimulus-response compatibility on neural selection
in frontal eye field. Neuron. 2003;38: 637–648.

29. Chiau HY, Tseng P, Su JH, Tzeng OJ, Hung DL, Muggleton NG, et al. Trial type
probability modulates the cost of antisaccades. J Neurophysiol. 2011;106: 515–

526.

30.   Stampe D. Heuristic filtering and reliable calibration methods for video-based
pupil-tracking systems. Behavior Research Methods, Instruments, \& Computers.
Springer-Verlag; 1993;25: 137–142. doi:10.3758/BF03204486

31.   Robert C, Casella G. Monte Carlo statistical methods. Springer Science \& Business
Media; 2013.

32.   Shaby B, Wells MT. Exploring an adaptive Metropolis algorithm. Durham, NC,
USA: Department of statistical science. Duke Universitiy; 2010.

33.   Gelman A, Carlin JB, Stern HS, Rubin DB. Bayesian Data Analysis. Chapman and
Hall/CRC; 2003.

34.   Aponte EA, Raman S, Sengupta B, Penny WD, Stephan KE, Heinzle J. mpdcm: A
toolbox for massively parallel dynamic causal modeling. J Neurosci Methods.
2016;257: 7–16.

35.   Ben Calderhead, Girolami MA. Estimating Bayes factors via thermodynamic
integration and population MCMC. Computational Statistics \& Data Analysis.
2009;53: 4028–4045. doi:10.1016/j.csda.2009.07.025

36.   Gelman A, Rubin DB. Inference from iterative simulation using multiple
sequences. Statistical Science. JSTOR; 1992;: 457–472.

37.   MacKay DJC. Information Theory, Inference, and Learning Algorithms. Cambridge
University Press; 2003.

38.   Stephan KE, Penny WD, Daunizeau J, Moran RJ, Friston KJ. Bayesian model
selection for group studies. Neuroimage. 2009;46: 1004–1017.

39.   Gelman A, Meng XL. Simulating Normalizing Constants: From Importance
Sampling to Bridge Sampling to Path Sampling. Statistical Science. Institute of
Mathematical Statistics; 1998;13: 163–185. doi:10.2307/2676756

40.   Kass RE, Raftery AE. Bayes factors. Journal of the american statistical association.
Taylor \& Francis Group; 1995;90: 773–795.

41.   Rigoux L, Stephan KE, Friston KJ, Daunizeau J. Bayesian model selection for group
studies - revisited. Neuroimage. 2014;84: 971–985.

42.   Penny WD, Stephan KE, Daunizeau J, Rosa MJ, Friston KJ, Schofield TM, et al.
Comparing families of dynamic causal models. PLoS Comput Biol. 2010;6:
e1000709.

43.   Brodersen KH, Schofield TM, Leff AP, Ong CS, Lomakina EI, Buhmann JM, et al.
Generative embedding for model-based classification of fMRI data. PLoS Comput
Biol. 2011;7: e1002079.

44.   Stephan KE, Schlagenhauf F, Huys QJ, Raman S, Aponte EA, Brodersen KH, et al.

Computational neuroimaging strategies for single patient predictions.
Neuroimage. 2017;145: 180–199.

45. Patterson TNL. The optimum addition of points to quadrature formulae.
Mathematics of Computation. 1968;22: 847–856.

46. Massen C. Parallel programming of exogenous and endogenous components in
the antisaccade task. Q J Exp Psychol A. 2004;57: 475–498.

47. Pierce JE, McDowell JE. Effects of preparation time and trial type probability on
performance of anti- and pro-saccades. Acta Psychol (Amst). 2016;164: 188–194.

48. Pierce JE, McDowell JE. Modulation of cognitive control levels via manipulation of
saccade trial-type probability assessed with event-related BOLD fMRI. J
Neurophysiol. 2016;115: 763–772.

49. Montague PR, Dolan RJ, Friston KJ, Dayan P. Computational psychiatry. Trends
Cogn Sci (Regul Ed). 2012;16: 72–80.

50. Wang XJ, Krystal JH. Computational psychiatry. Neuron. 2014;84: 638–654.

51. Stephan KE, Mathys C. Computational approaches to psychiatry. Curr Opin
Neurobiol. 2014;25: 85–92.

52. Paulus MP, Huys QJ, Maia TV. A Roadmap for the Development of Applied
Computational Psychiatry. Biological Psychiatry: Cognitive Neuroscience and
Neuroimaging. Elsevier; 2016.

53. Huys QJ, Maia TV, Paulus MP. Computational Psychiatry: From Mechanistic
Insights to the Development of New Treatments. Biological Psychiatry: Cognitive
Neuroscience and Neuroimaging. Elsevier; 2016;1: 382–385.

54. Findlay JM, Walker R. A model of saccade generation based on parallel processing
and competitive inhibition. Behav Brain Sci. 1999;22: 661–674.

55. Feng G. Is there a common control mechanism for anti-saccades and reading eye
movements? Evidence from distributional analyses. Vision Res. 2012;57: 35–50.

56. Gold JI, Shadlen MN. The neural basis of decision making. Annu Rev Neurosci.
2007;30: 535–574.

57. Ratcliff R, Smith PL, Brown SD, McKoon G. Diffusion Decision Model: Current
Issues and History. Trends Cogn Sci (Regul Ed). 2016;20: 260–281.

58. Donkin C, Brown S, Heathcote A, Wagenmakers EJ. Diffusion versus linear ballistic
accumulation: different models but the same conclusions about psychological
processes? Psychon Bull Rev. 2011;18: 61–69.

59. Moscoso del Prado Martin F. A theory of reaction time distributions. 2008.

60. Barndorff-Nielsen O, Blæsild P, Halgreen C. First hitting time models for the

generalized inverse Gaussian distribution. Stochastic Processes and their
Applications. Elsevier; 1978;7: 49–54.

61.   Cutsuridis V. Behavioural and computational varieties of response inhibition in
eye movements. Philos Trans R Soc Lond, B, Biol Sci. 2017;372.

62.   Logan GD, Van Zandt T, Verbruggen F, Wagenmakers EJ. On the ability to inhibit
thought and action: general and special theories of an act of control. Psychol Rev.
2014;121: 66–95.

63.   Cutsuridis V, Smyrnis N, Evdokimidis I, Perantonis S. A neural network model of
decision making in an antisaccade task by the superior colliculus. Neural
Networks. 2007;20: 690–704.

64.   Cutsuridis V, Kumari V, Ettinger U. Antisaccade performance in schizophrenia: a
neural model of decision making in the superior colliculus. Front Neurosci.
2014;8: 13.

65.   Cutsuridis V. Neural competition via lateral inhibition between decision processes
and not a STOP signal accounts for the antisaccade performance in healthy and
schizophrenia subjects. Front Neurosci. 2015;9: 5.

66.   Brown JW, Bullock D, Grossberg S. How laminar frontal cortex and basal ganglia
circuits interact to control planned and reactive saccades. Neural Netw. 2004;17:
471–510.

67.   Heinzle J, Hepp K, Martin KA. A microcircuit model of the frontal eye fields. J
Neurosci. 2007;27: 9341–9353.

68.   Wiecki TV, Frank MJ. A computational model of inhibitory control in frontal
cortex and basal ganglia. Psychol Rev. 2013;120: 329–355.

69.   Carpenter R, Noorani I. Movement suppression: brain mechanisms for stopping
and stillness. Philos Trans R Soc Lond, B, Biol Sci. 2017;372.

70.   Boucher L, Palmeri TJ, Logan GD, Schall JD. Inhibitory control in mind and brain:
an interactive race model of countermanding saccades. Psychol Rev. 2007;114:
376–397.

71.   Schmidt R, Berke JD. A Pause-then-Cancel model of stopping: evidence from basal
ganglia neurophysiology. Philos Trans R Soc Lond, B, Biol Sci. 2017;372.

72.   Bissett PG. The countermanding task revisited: mimicry of race models. J
Neurosci. 2013;33: 12150–12151.

73.   Schall JD, Palmeri TJ, Logan GD. Models of inhibitory control. Philos Trans R Soc
Lond, B, Biol Sci. 2017;372.

74.   Schmidt R, Leventhal DK, Mallet N, Chen F, Berke JD. Canceling actions involves a
race between basal ganglia pathways. Nat Neurosci. 2013;16: 1118–1124.

75.   Stuphorn V, Schall JD. Executive control of countermanding saccades by the
1077   supplementary eye field. Nat Neurosci. 2006;9: 925–931.

1078   76.   Pouget P, Murthy A, Stuphorn V. Cortical control and performance monitoring of
1079   interrupting and redirecting movements. Philos Trans R Soc Lond, B, Biol Sci.
1080   2017;372.

1081   77.   Reuter B, Rakusan L, Kathmanna N. Poor antisaccade performance in
1082   schizophrenia: an inhibition deficit? Psychiatry Res. 2005;135: 1–10.

1083   78.   Reuter B, Jager M, Bottlender R, Kathmann N. Impaired action control in
1084   schizophrenia: the role of volitional saccade initiation. Neuropsychologia.
1085   2007;45: 1840–1848.

1086   79.   Coe BC, Munoz DP. Mechanisms of saccade suppression revealed in the anti-
1087   saccade task. Philos Trans R Soc Lond, B, Biol Sci. 2017;372.

1088   80.   Hakvoort Schwerdtfeger RM, Alahyane N, Brien DC, Coe BC, Stroman PW, Munoz
1089   DP. Preparatory neural networks are impaired in adults with attention-
1090   deficit/hyperactivity disorder during the antisaccade task. Neuroimage Clin.
1091   2012;2: 63–78.

1092   81.   Cameron IG, Pari G, Alahyane N, Brien DC, Coe BC, Stroman PW, et al. Impaired
1093   executive function signals in motor brain regions in Parkinson's disease.
1094   Neuroimage. 2012;60: 1156–1170.

1095   82.   Witiuk K, Fernandez-Ruiz J, McKee R, Alahyane N, Coe BC, Melanson M, et al.
1096   Cognitive deterioration and functional compensation in ALS measured with fMRI
1097   using an inhibitory task. J Neurosci. 2014;34: 14260–14271.

1098