RESEARCH ARTICLE

Genetic Architecture and Molecular Networks Underlying Leaf Thickness in Desert-Adapted Tomato *Solanum pennellii*

Viktoriya Coneva[1], Margaret H. Frank[1], Maria A. de Luis Balaguer[2], Mao Li[1], Rosangela Sozzani[2], Daniel H. Chitwood [1,3,4]

[1] *Donald Danforth Plant Science Center, 975 North Warson Road, St. Louis, MO 63132, USA*
[2] *Department of Plant and Microbial Biology, North Carolina State University, 2552A Thomas Hall, Raleigh, NC 27695, USA*
[3] *Current address: Independent Researcher, Santa Rosa, CA 95409, USA*
[4] *Corresponding author: Daniel H. Chitwood, dhchitwood@gmail.com*

Author Contributions
VC, MHF and DHC designed the research. VC, MF, MADLB, RS, and ML conducted the experiments and analyzed the data. VC wrote the manuscript with contributions from other authors.

One-Sentence Summary
We identified QTL for leaf thickness in desert-adapted tomato and characterized the anatomic and transcriptional alterations associated with this trait using a set of introgression lines.

Funding information
This work was supported by funds from the Donald Danforth Plant Science Center. RS is supported by an NSF CAREER grant (MCB-1453130). MHF is supported by an NSF NPGI post-doctoral fellowship (IOS-1523668).

Abstract

Thicker leaves allow plants to grow in water-limited conditions. However, our understanding of the genetic underpinnings of this highly functional leaf shape trait is poor. We used a custom-built confocal profilometer to directly measure leaf thickness in a set of introgression lines (ILs) derived from the desert tomato species *Solanum pennellii*, and identified quantitative trait loci (QTL). We report evidence of a complex genetic architecture of this trait and roles for both genetic and environmental factors. Several ILs with thick leaves have dramatically elongated palisade mesophyll cells and, in some cases, increased leaf ploidy. We characterized thick ILs 2-5 and 4-3 in detail and found increased mesophyll cell size and leaf ploidy levels, suggesting that endoreduplication underpins leaf thickness in tomato. Next, we queried the transcriptomes and inferred Dynamic Bayesian Networks of gene expression across early leaf ontogeny in these lines to compare the molecular networks that pattern leaf thickness. We show that thick ILs share *S. pennellii*-like expression profiles for putative regulators of cell shape and meristem determinacy, as well as a general signature of cell cycle related gene expression. However, our network data suggest that leaf thickness in these two lines is patterned by at least partially distinct mechanisms. Consistent with this hypothesis, double homozygote lines combining introgression segments from these two ILs show additive phenotypes including thick leaves, higher ploidy levels and larger palisade mesophyll cells. Collectively, these data establish a framework of genetic, anatomical, and molecular mechanisms that pattern leaf thickness in desert-adapted tomato.

Introduction

Leaves are the primary photosynthetic organs of land plants. Quantitative leaf traits have important connections to their physiological functions, and ultimately, to whole plant productivity and survival. While few aspects of leaf morphology have been unambiguously determined as functional (Nicotra et al., 2011), clear associations between leaf traits and variations in climate have been drawn (Wright et al., 2004). Leaf thickness, the distance between the upper (adaxial) and lower (abaxial) leaf surfaces, has been shown to correlate with environmental variables such as water availability, temperature and light quantity. Thus, on a global scale, across habitats and land plant diversity, plants adapted to arid environments tend to have thicker leaves (Wright et al., 2004; Poorter et al., 2009).

Leaf thickness is a continuous, rather than a categorical, trait. Thus, it is important to distinguish between thickness in the context of "typical" leaf morphology, generally possessing clear dorsiventrality (adaxial/abaxial flattening) in comparison to extremely thick leaves, described as "succulent", which are often more radial. While the definition of succulence is eco-physiological, rather than morphological (Ogburn and Edwards, 2010), at the cellular level it is broadly associated with increased cell size and relative vacuole volume (Gibson, 1982; von Willert et al., 1992). These cellular traits promote the capacity to store water and to survive in dry environments (Becker, 2007). Allometric studies across land plant families have shown that leaf thickness scales specifically with the size of palisade mesophyll cells - the adaxial layer of photosynthetic cells in leaves (Garnier and Laurent, 1994; Roderick et al., 1999; Sack and Frole, 2006; John et al., 2013). Increased palisade cell height leads to increased area of contact with the intercellular space and thereby to improved uptake of carbon dioxide (CO_2) into mesophyll cells (Oguchi et al., 2005; Terashima et al., 2011), possibly offsetting the increased CO_2 diffusion path in thicker leaves. At the organismal level, thicker leaves present a tradeoff between rapid growth versus drought and heat tolerance (Smith et al., 1997). This idea is supported by global correlations between leaf mass per area (LMA), a proxy for leaf thickness, and habits associated with slower growth (Poorter et al., 2009).

Although leaf thickness is a highly functional trait, mechanistic understanding of how it

is patterned during leaf ontogeny is poor. The main cellular events that underpin leaf

development are the establishment of adaxial/abaxial polarity, followed by cell division,

directional expansion, and differentiation (Effroni et al., 2008). Changes in the relative

timing (heterochrony) and duration of these events can impact leaf morphology,

including thickness. Several mutants have been identified that show clear alterations in

leaf thickness. These include the Arabidopsis *angustifolia* and *rotundifolia3* (Tsuge et al.,

1996), as well as *argonaute1*, *phantastica*, and *phabulosa* (Bohmert et al., 1998), which

have aberrations in the polarity of cell elongation and the establishment of adaxial/abaxial

polarity, respectively, as well as the *N. sylvestris fat* and *lam-1* (McHale, 1992, 1993),

which affect the extent of periclinal cell division in leaves. However, these

developmental mutants do not necessarily inform us of the mechanisms by which natural

selection acts to pattern quantitative variation in leaf thickness.

Efforts to understand the genetic basis of leaf thickness in the context of natural variation

face several important challenges. First, direct measurement of leaf thickness at a scale

that would allow the investigation of Quantitative Trait Loci (QTL) for the trait is not

trivial. Because of the difficulty in measuring leaf thickness directly, LMA is most often

used as a proxy for this trait (Poorter et al., 2009; Muir et al., 2014). Second, in addition

to genetic components, leaf thickness is environmentally plastic – it is responsive to both

the quantity and quality of light (Pieruschka and Poorter, 2012). Finally, because leaf

thickness varies on a continuous spectrum and is not associated with any particular

phylogenetic lineage or growth habit, mechanistic questions regarding its patterning need

to be addressed in a taxon-specific manner.

With these considerations in mind, we used two members of the tomato clade (*Solanum*

sect. *Lycopersicon*)**,** which are closely related, morphologically distinct, and occupy

distinct environments (Nakazato et al., 2010) to study the genetic basis and

developmental patterning of leaf thickness. The domesticated tomato species *S.*

lycopersicum inhabits a relatively wide geographic range characterized by warm, wet

conditions with little seasonal variation. By contrast, the wild species *S. pennellii* is

endemic to coastal regions of the Atacama desert of Peru, a habitat characterized by

extremely dry conditions (Nakazato, et al., 2010). The leaves of *S. pennellii* plants,

therefore, exhibit morphological and anatomical features that are likely adaptations to dry

conditions (McDowell et al., 2011; Haliński et al., 2015), including thick leaves (Koenig

et al., 2013). Moreover, a set of homozygous introgression lines (ILs) harboring defined,

partially overlapping segments of the *S. pennellii* genome in an otherwise *S. lycopersicum*

background (Eshed and Zamir 1995) has been used to successfully map a number of

QTL, including fruit metabolite concentrations (Fridman et al., 2004; Schauer et al.,

2006), yield (Semel et al., 2006), and leaf shape (Chitwood et al., 2013). Here, we used a

custom-built dual confocal profilometer to obtain precise measurements of leaf thickness

across the IL panel and identified QTL for this trait in tomato. Leaf thickness correlates

with other facets of leaf shape, as well as a suite of traits associated with desiccation

tolerance and lower productivity. We investigated the anatomical manifestations of

thickness in tomato and found a prominent increase in palisade cell height in many thick

ILs. Finally, we inferred comparative gene regulatory networks of early leaf development

(plastochron stages P1-P4) in two thick lines using organ-specific RNA-Seq and

identified molecular networks that pattern *S. pennellii*-like desert-adapted leaves.

132   Results

133   Complex genetic architecture of leaf thickness across *S. pennellii* ILs

To investigate the genetic architecture and patterning of leaf thickness in the *S. pennellii*

IL panel, we used a custom-built dual confocal profilometer device (Fig. S1), which

generates precise thickness measurements throughout the leaflet lamina at a range of

resolutions ($0.1 - 1.0 \ mm^2$) and at high-throughput. The device makes use of two

confocal lasers positioned on either side of the sample and calculates thickness by

measuring the distance between each of the sample's surfaces and the corresponding laser

probe. Finally, we visualize thickness as a heatmap of thickness values across the surface

of the leaf lamina (Fig. 1A).

We first compared leaflet thickness in *S. lycopersicum* var. M82 and its desert relative *S.*

pennellii LA0716. Our confocal profilometer measurements showed that *S. pennellii*

leaflets are thicker than those of domesticated tomato, as previously reported (Fig. 1,

Koenig et al., 2013), demonstrating the capacity of this device to quantitatively detect

fine differences in leaf lamina thickness. We compared dynamic growth patterns of the

two species under water limited conditions and show that, unlike the domesticated

species, *S. pennellii* is unaffected by drought (Fig. 1C). This observation highlights the

importance of understanding the patterning of developmental traits in this species, such

as leaf thickness, which may contribute to drought tolerance. We proceeded to measure

leaf thickness across the *S. pennellii* introgression line panel in field conditions.

We used mixed linear regression models to compare each of the introgression lines to the

domesticated parent M82 (Dataset S1) and found that 31 ILs had significantly thicker

leaflets than the M82 parent, while 5 had transgressively thinner leaflets. The overall

broad-sense heritability for leaflet thickness is 39.1% (Fig. 2). The lines with thickest

leaflets are IL5-4, IL5-3, IL8-1, IL4-3, IL8-1-1 (contained within IL8-1), and IL2-5,

while IL4-1-1, IL2-6-5, IL9-1-3, IL12-4-1, and IL2-1 have thinner leaves than the M82

parent.

Based on the observation that the heritability value for leaf thickness is 39.1 %, we

reasoned that environmental factors are likely to play a role in modulating leaf thickness.

We thus compared our field experiment with leaf thickness data for vegetative leaves of

greenhouse-grown plants. We selected 20 ILs, which were highly significant for leaf

thickness differences from M82 in field conditions ($p < 0.001$) and observed that only

some of these lines are also significantly thicker than the domesticated parent in

greenhouse conditions ($p < 0.05$, Fig. S2A). Finally, our observations suggest that leaf

thickness varies across the shoot of a number of our select thick leaf ILs with post-

flowering leaves having thicker leaves than vegetative leaves (Fig. S2B).

For each leaflet in our field experiment, we also quantified leaf mass per unit area

(LMA), which reflects both thickness and density, and is traditionally used as a proxy for

leaf thickness. Although the heritability for LMA is similar to that for thickness (33.2%

and 39.1%, respectively), significant QTL for these two traits do not consistently overlap

(Dataset S1).

Leaf thickness and LMA are correlated with distinct suites of traits in tomato

We generated pairwise correlations between leaflet thickness, LMA, and a suite of other

previously published traits including metabolite (MET), morphological (MOR),

enzymatic activity in fruit pericarp (ENZ), seed-related (SED), developmental (DEV),

and elemental profile-related (ION) (Datasets S2-4, Chitwood et al., 2013 and references

therein). Spearman's correlation coefficients with significant q-values (q < 0.050) are

reported in Fig.2B. Leaf thickness and LMA are correlated (rho = 0.423, q = 0.003). Leaf

thickness also correlates with leaf shape parameters, such as roundness (rho = 0.328, q =

0.044), aspect ratio (rho = -0.327, q = 0.045), and the first two principal components of

the elliptical Fourier descriptors of leaflet shape (EFD.PC1 rho = 0.414, q = 0.004 and

EFD.PC2 rho = 0.406, q = 0.005). Thickness is negatively correlated with several

reproductive traits, including yield (rho = -0.337, q = 0.037), seed weight (rho = -0.342, q

= 0.033) and seed number per plant (rho = -0.339, q = 0.036). Moreover, leaf thickness is

negatively correlated with leaf stomatal ratio, the relative density of stomata on the

abaxial and adaxial sides of the leaf (rho=-0.352, q = 0.031), and positively with

glutamate dehydrogenase activity (rho = 0.367, q = 0.017) and seed galactinol content

(rho = 0.342, q = 0.048).

Leaf mass per area is associated with a distinct suite of traits from leaf thickness. In

addition to a positive correlation with the content of some enzymes (GAPDH and

Shikimate DE) and metabolites (Glutamate), LMA is significantly negatively correlated

with the accumulation of Na and Mg in all leaflets tested. LMA, but not leaf thickness, is

also significantly positively correlated with total plant weight, reflecting vegetative

biomass accumulation.

Thick IL leaves have elongated palisade parenchyma cells

Leaf cross-sections of field-grown M82 and select ILs with increased leaf thickness, as

well as greenhouse-grown *S. pennellii* (Sp) leaves were stained with propidium iodide to

assess the anatomical changes that lead to increased leaf thickness. We observed that,

relative to the M82 parent, the Sp parent and several ILs, have an elongated palisade

mesophyll cell layer corresponding to the adaxial layer of photosynthesizing cells in

tomato leaves (Fig. 3). Palisade parenchyma elongation is especially dramatic for IL1-3,

IL2-5, IL4-3, and IL10-3. Both leaf thickness and palisade elongation phenotypes are

attenuated for vegetative leaves of greenhouse-grown plants (Fig. S2, Fig. S3A).

Anatomy and early leaf development in select ILs with thick leaves

To capture an overall view into the core mechanisms of leaf thickness patterning, we

further analyzed lines IL2-5 and IL4-3. We selected IL2-5 due to its dramatic anatomy in

field conditions (Fig. 3) and its lack of other characterized leaf morphology phenotypes

(Chitwood et al., 2013), while IL4-3 leaflets are both significantly thicker and less

serrated than those of the domesticated parent (Fig. 2; Dataset S1, circularity - the ratio

between leaflet area and the square of its perimeter - reflects lobing and serration). To

further investigate the relationships between genetic determinants of leaf thickness in

these ILs, we generated a double homozygous line combining the entire *S. pennellii*

segments of IL2-5 and IL4-3.

Double homozygotes (IL2-5/IL4-3) have significantly thicker leaves than M82 at both

vegetative (Fig. 4A, p = 0.019) and post-flowering stages (Fig. S2B) in greenhouse

conditions. Additionally, IL2-5/IL4-3 plants have significantly smoother margins than

either of the IL parents (Fig. 4B), suggesting additive genetic interactions for both of

these traits. We next compared the dimensions of the mesophyll cell layers in each IL and

the double homozygote line to determine the contributions each cell layer makes to the

observed increase in leaflet thickness. We found that palisade mesophyll cells are

significantly larger in IL2-5/IL4-3 than in M82 leaves (Fig. S4). Further, the ratios of

palisade cell length to both total leaf thickness and to the length of the spongy mesophyll

are significantly larger in IL2-5/IL4-3 than in M82 leaves (Fig. S4). IL2-5 shows similar

albeit less pronounced trends as the double homozygote line, while in IL4-3 both spongy

and palisade mesophyll cell layers are longer than in M82, with the spongy mesophyll

layer making the most significant contribution to leaf thickness.

Since increases in cell size are often driven by endopolyploidy, we performed flow

cytometry on fully expanded vegetative leaves of each genotype and observed increased

ploidy profiles in all lines relative to the domesticated parent (Fig. 4C). Notably, the

double homozygote line exhibited higher ploidy levels than both single ILs and the *S.*

penellii parent (Fig. 4C, Fig. S4). Notably, we also observed a trend to increased ploidy

in several greenhouse-grown thick ILs (IL7-4-1, IL8-1) (Fig. S3B).

To understand if alterations in leaf size occur during early stages of leaf ontogeny in

these lines, we quantified P3 organ dimensions and compared them with the M82

parental line. For this, we assembled 3D confocal reconstructions of vegetative shoot

apices, calculated the surface mesh, extracted P3 leaf primordia, and quantified their total

volume, length, and mean diameter. We found that IL4-3 P3 leaf primordia are

significantly larger than M82 in terms of overall volume (p = 0.0179), as well as both

length (p = 0.0035) and diameter (p = 0.0230). In IL2-5 P3 volume (not statistically

significant) and diameter (p = 0.0116) are increased, while length is comparable to M82.

Although P3 primordia of double homozygote plants were statistically indistinguishable

from those of M82 plants except for shorter arc length (p = 0.0411) (Fig. 4D) our

observations also suggest that double homozygote leaves increase in size dramatically

between P3 and P4 stages (Fig. S5).

Transcriptomic signatures of early leaf development in thicker ILs

To investigate the molecular events that define the patterning of IL2-5 and IL4-3 leaves,

we isolated leaf primordia from each IL and the two parents (M82 or Sl and Sp) at four

successive stages of development: P1 (containing the shoot apical meristem, SAM, and

the youngest leaf primordium), P2, P3 (characterized by leaflet emergence) and P4

(typically the onset of cell differentiation) (Fig. 5A). For *S. pennellii*, P1 samples were

comprised of the SAM, P1, and P2, since these organs were not separable by hand

dissection. Thus, the Sp transcriptomic dataset includes samples designated as P1, P3,

and P4. Principal Component Analysis (PCA) of the resulting RNA-Seq data, after

normalization and filtering, shows that samples group clearly by organ stage (PC2) (Fig.

5B). In addition, PC1 separates *S. pennellii* samples from all other genotypes. To

investigate how IL leaves are similar to the Sp parent, we looked for genes that are

differentially expressed (DEGs) between corresponding stages of each IL and the M82

parent, while also being differentially expressed between M82 and Sp. In other words, we

identified the set of DEG for each organ stage that is common to each IL and Sp relative

to M82. For P2 we considered only the comparison with M82, as our Sp dataset did not

include independently dissected P2 stage primordium samples (Fig. S6, Dataset S5).

We identified a total of 812 DEGs across P1-P4 stages in IL2-5, and of these, 544 are up-

regulated in at least one organ stage, while 269 are down-regulated (Fig.5C). In IL4-3, we

detected 632 DEG, 361 of which are up-regulated and 271 are down-regulated in the IL

(Fig. 5C). Many of the DEGs are differentially expressed at more than one stage (Fig. 5C,

Dataset S5). Additionally, based on tomato transcription factor (TF) annotation by Suresh

et al. (2014), we identified putative transcription factor-encoding genes among each IL's

DEG sets. Myb-related, Ethylene Responsive, MADS, and WRKY are the abundant

classes of TF-encoding DEGs in IL2-5, while in IL4-3 TFs belonging to bZIP and Myb-

related are highly represented families (Fig. S7).

We identified differentially expressed TF-encoding genes that are common to the two ILs

and the Sp parent (Fig. 6), reasoning that some of these can be regulators of leaf

thickness. Five of the seven shared TF-encoding genes are up-regulated in the ILs relative

to M82. A MADS-box TF (Solyc12g087830) is up-regulated at all stages in both ILs,

while two additional inflorescence meristem-related transcription factors, LFY-like

(Solyc03g118160) and AP2-like (Solyc07g049490) are differentially expressed at

corresponding stages in both ILs. The SHORTROOT-like (SHR-like) GRAS TF

Solyc08g014030 is up-regulated at P2 in both ILs, while its expression increases at each

progressive stage and peaks at P4 in all genotypes. A putative JASMONATE ZIM-

domain protein (JAZ1, Solyc12g009220) is also up-regulated at P2 in both ILs, while a

LIM-domain protein (Solyc04g077780) is up-regulated in the ILs at P3 (in IL4-3) and P4

(both ILs) (Fig. 6A).

Next, we compared the expression profiles of genes known to be involved in tomato leaf

development (Ichihashi et al., 2014). We selected only genes that are differentially

expressed in the same direction in each IL and Sp relative to the domesticated parent

M82 and highlighted genes that are common to both thick ILs to arrive at a set of entities

that may be core to the patterning of leaf thickness (Fig. 6B). A gibberellin 20-oxidase

encoding gene (GA 20-ox, Solyc03g006880) is up-regulated at P3 in both ILs and

throughout the P1-P3 interval in IL4-3. A set of two closely related ULTRAPETALA1

genes (ULT1, Solyc12g010360 and Solyc12g010370) is down-regulated at all leaf

developmental stages in both ILs. A number of leaf development regulators are

additionally differentially expressed in either of the ILs. Some noteworthy classes include

entities related to auxin metabolism or transport (auxin efflux carrier, IAA-

carboxymethyltransferase, YUCCA-like monooxygenase), leaf complexity, lobing and

serrations (three BEL1-like TFs, CUC2-like and BOP2-like), meristem maintenance or

patterning (two BAM1-like receptor kinases and an AP2-like TF), and cell division and

expansion (GRF1 and ROT3-like TFs).

Similarly, we also queried DEG sets for entities annotated as cell cycle or

endoreduplication to assess whether these two thick ILs share a common trajectory of

cellular events during leaf ontogeny (Fig. 6C). Overall, we observed distinct expression

profiles for these genes in IL2-5 and IL4-3.

Finally, to broadly characterize the types of processes that may regulate the molecular

networks of early leaf development in the ILs we applied GO enrichment analysis

(agriGO, Du et al., 2010) (Dataset S6) and identified statistically enriched promoter

motifs among the organ-specific DEG sets (Dataset S7). Importantly, we observed that at

P4, the set of up-regulated genes in IL2-5 is enriched for biological process terms relating

to "photosynthesis" (GO:0015979) and "translation" (GO:0006412), while down-

regulated genes at this stage are enriched for terms relating to "DNA binding"

(GO:0003677). Our promoter motif analysis showed that motifs associated with

regulation by abiotic factors such as light, circadian clock, water availability, and

temperature are prominent among IL2-5 DE genes. In addition, binding sites for

developmental regulators, hormone-associated promoter motifs, and a cell cycle regulator

are among the list of significant motifs. Among development-associated motifs, CArG

(MADS-box), BEL1-like (BELL) and SBP-box transcription factor binding sites are also

significantly enriched in both IL 2-5 and 4-3 DEG sets. (Fig. S8, Dataset S7).

Gene regulatory networks of early leaf development in thick ILs

To detect regulators of early leaf development that each IL (IL2-5 and IL4-3) shares with

the *S. pennellii* parent, we inferred Dynamic Bayesian Networks (DBN) using the IL and

Sp overlapping DEG sets described in the previous section (de Luis Balaguer et al.,

2017). Additionally, we only allowed putative transcription factor-encoding genes

(Suresh et al., 2014) as "source" nodes (genes that control the expression of other co-

expressed genes). First, we constructed individual networks for each leaf developmental

stage, for which an overlap with Sp data is available (P1, P3, P4), and then combined the

results to visualize the overall *S. pennellii*-like leaf developmental networks (Fig. 7,

Dataset S8). The IL2-5 network (Fig. 7A) contains two major regulators, which are

central to more than one developmental stage: a SQUAMOSA promoter-binding protein-

like domain gene (SBP-box 04g, Solyc04g064470) and a CONSTANS-like Zinc finger

(Zn-finger CO-like 05g, Solyc05g009310) (Dataset S8). Similarly, the IL4-3 network

(Fig. 7C) features two central regulators: a BEL1-like homeodomain transcription factor

gene (BEL1 04g, Solyc04g080780) and a MADS-box domain-containing gene (MADS-

box 12g, Solyc12g087830) (Dataset S8). Importantly, few nodes are shared between the

organ-specific networks of IL2-5 and IL4-3. We surveyed each network for shared

differentially expressed leaf development genes and found that GA 20-ox 03g

(Solyc03g006880) is present in both networks but is regulated by different sets of

transcription factors in each IL (Fig. 7B, D).

We also inferred a second set of networks for each of the ILs by identifying DEGs using

similar criteria as above. However, in contrast to the previous set of networks, where

genes were separated into organ stages based on differential expression at each discrete

stage, we used a clustering approach to group regulators and select co-expressed gene

sets according to expression profiles. For these analyses, we also included P2 DEGs (IL

vs M82) to ensure continuity of expression profiles (Dataset S9). This approach allowed

us to examine a more dynamic view of early developmental processes. The resulting

networks (Dataset S9) feature a putative auxin responsive TF AUX/IAA 12g

(Solyc12g096980) for both ILs (Fig. 7E, F). Moreover, the AUX/IAA 12g sub-network

or IL2-5 includes the SHR-like GRAS domain TF that is up-regulated during leaf

development in both ILs (GRAS 08g, Solyc08g014030) (Fig. 6A, Fig. 7E).

Discussion

Leaf thickness has a complex genetic architecture in desert-adapted tomato and is

associated with overall leaf shape, desiccation tolerance, and decreased yield

While extensive progress has been made dissecting the molecular-genetic patterning of

two-dimensional leaf morphology, relatively little is known about the third dimension of

leaf shape – thickness. Here, we used a custom-built dual confocal profilometer to obtain

direct measurements of leaf thickness across the *S. pennellii* x *S. lycopersicum* IL panel

(Eshed and Zamir, 1995) (Fig. 1, Fig. S1) and identified QTL for this trait (Fig. 2A). We

found that nearly half of the ILs have significantly thicker leaves than the domesticated

parent M82, while a small number have transgressively thinner leaves. The broad-sense

heritability for leaf thickness in this experiment is moderate (39%). Collectively, these

observations point to a complex genetic basis for this trait. A previous quantitative

genetic analysis of a suite of desert-adaptive traits in the same *S. pennellii* IL panel found

fewer significantly thicker lines and lower heritability (12%) for this trait (Muir et al.,

2014). However, the previous study estimated thickness as the ratio of LMA to leaflet dry

matter content, while we measured thickness directly. Further, our study was conducted

in field conditions, while Muir et al. (2014) measured the trait using greenhouse-grown

plants. Given that environment significantly affects the magnitude of this trait (Fig. S2) it

is not surprising that these studies report only partially overlapping outcomes.

In order to understand how variation in leaf thickness relates to other traits, particularly to

leaf mass per area, we calculated pairwise correlation coefficients among all leaf shape

and elemental profile traits, as well as a collection of previously published traits

(summarized in Chitwood et al., 2013; Datasets S3, S4). As expected, leaf thickness and

LMA are significantly correlated across the IL panel. However, the two traits have

distinct sets of significant trait correlations (Fig. 2B). Collectively, these data suggest that

thickness and LMA are likely patterned by separate mechanisms and that direct

measurements of leaf thickness are necessary to further dissect the genetic basis of this

trait.

Leaf thickness is significantly correlated with leaf shape traits such as aspect ratio and the

first two principal components of elliptical Fourier descriptors of overall shape. However,

our data do not establish whether this correlation reflects a common patterning

mechanism or developmental and/or mechanical constraints among these traits.

Alternatively, the relatively modest correlations (rho values between 0.33 - 0.41) could

reflect independent variation in these traits resulting in considerable flexibility in final

leaf morphology, as suggested by Muir et al. (2016).

Leaf thickness is negatively correlated with yield-related traits, which suggests a trade-off

between investments in vegetative and reproductive biomass that is further substantiated

by the positive correlation between LMA and plant weight (Fig.2B). Some studies

support the hypothesis of a tradeoff between leaf mass per area and rapid growth (Smith

et al., 1997; Poorter et al., 2009), while others find poor coordination between growth

rate and LMA (Muir et al., 2016). Finally, leaf thickness is significantly correlated with

leaf stomatal ratio, glutamate dehydrogenase activity, and galactinol content in seeds, a

suite of traits associated with desiccation tolerance in plants (Taji et al., 2002; Lightfoot

et al., 2007). We also observed negative correlations between LMA and the accumulation

of several elements in leaves, most notably Na and Mg (Fig. 2C). This finding supports

the idea that LMA and thickness are distinct traits, and that LMA reflects the material

composition of leaves, while leaf thickness is a developmentally patterned trait.

Thicker *S. pennellii* IL leaves have elongated palisade mesophyll cells

The observed elongated palisade mesophyll cells in the leaves of several field-grown ILs

with significantly thicker leaves (Fig. 3A), as well as in the desert-adapted *S. pennellii*

parent suggest that dorsiventral expansion of palisade mesophyll cells contributes most

prominently to increased leaf thickness. This hypothesis is supported by the fact that

palisade cell height increases more significantly than the total height of the spongy

mesophyll in thick leaves of double homozygous IL2-5/IL4-3 lines (Fig. S4). Palisade

cell height is positively correlated with photosynthetic efficiency (Niinemets et al., 2009;

Terashima et al., 2011) and water storage capacity in succulent CAM (Crassulacean Acid

Metabolism) plants (Nelson et al., 2005). Our data also indicate that the magnitudes of

palisade cell elongation, as well as overall leaf thickness are modulated by environmental

inputs (Fig. 2, Fig. S2). High light has been shown to mediate increased leaf thickness

(Poorter et al., 2009; Wuyts et al., 2012; Kalve et al., 2014), as well as specifically

palisade cell elongation (Kozuka et al., 2011) in Arabidopsis, while elongated palisade

cells promote a more efficient distribution of direct light throughout the photosynthetic

mesophyll compared with shorter cells (Brodersen et al., 2008; Brodersen and Vogelman

2010). Thus, thicker leaves composed of elongated palisade cells may be an adaptation to

desert-like dry, direct light environments, whereby the magnitude of these traits is

responsive to these environmental cues. Consistent with this hypothesis, we observed that

IL2-5 DEG promoters are enriched in motifs that reflect sensitivity to abiotic stimuli,

prominently light and water status (Fig. S8, Dataset S7).

Mechanisms of cell enlargement in thick ILs: increased ploidy and alterations in cell

cycle related gene expression

We compared the size of palisade mesophyll cells in leaf cross sections of thick ILs 2-5,

4-3, and a homozygous line combining both introgression segments and observed larger

palisade cells compared to M82 (Fig. S4), suggesting a link between leaf thickness and

cell size in tomato. Further, we showed significantly higher ploidy levels in the leaves of

these lines relative to the domesticated parent (Fig. 4C), indicating that increased

endoreduplication may underpin larger cells, and ultimately, thicker leaves. A partially

overlapping series of cell division, cell expansion, and cell differentiation events underlie

leaf development (Effroni et al., 2008). These processes are tightly coordinated to buffer

perturbations in overall organ shape and size (Tsukaya 2003; Beemster et al., 2003).

Thus, the relative timing and duration of any of these events can impact leaf size and

morphology. Additionally, different tissue types in the leaf can have distinct schedules of

cellular events during leaf ontogeny; for example, in Arabidopsis palisade mesophyll

cells have a shorter window of cell division compared to epidermal cells, and thus an

earlier onset of cell expansion and endoreduplication, resulting in differences in cell

volumes and geometry (Wuyts et al., 2012; Kalve et al., 2014). Given the prominent

contribution of specific cell types to leaf thickness (palisade mesophyll cells in IL2-5, for

example, vs both palisade and spongy mesophyll cells in IL4-3 (Fig. S4)), kinematic

studies to capture the timing and extent of tissue-specific cell division and

endoreduplication are needed to fully address the dynamic cellular basis of leaf thickness

patterning. The observed increase in P3 organ volume and thickness in IL4-3 and, to a

lesser extent, IL2-5 relative to M82 (Fig. 4D) support the notion that differences in the

trajectory of cellular events during early leaf ontogeny may underpin leaf thickness.

Comparative gene expression profiles of early leaf ontogeny in ILs 2-5 and 4-3 show

evidence of *S. pennellii*-like alterations in cell proliferative activity in these thick ILs.

Specifically, among a small set of shared differentially expressed genes is a GRAS-

domain TF GRAS 08g (Solyc08g014030) up-regulated at P2 in both lines (Fig. 6A,

Dataset S5). This gene is closely related to the Arabidopsis gene encoding SHORTROOT

(SHR) (Huang et al., 2015), which together with another GRAS-domain TF,

SCARECROW (SCR), regulates the duration of cell proliferation in leaves (Dhondt et

al., 2010). Moreover, consistent with previous reports, IL2-5 and IL4-3 DEGs are

enriched for E2F binding site motifs (Dataset S7, Ranjan et al., 2016). E2F transcription

factors act downstream of SHR and SCR to regulate progression through the S-phase of

the cell cycle (Dhondt et al., 2010). These data support the notion that the extent and/or

duration of cell proliferation underpin increased thickness in these lines. Another set of

DEGs that distinguish the thick ILs and the Sp parent from domesticated tomato include

three genes with predicted functions in regulating the cell cycle and cell expansion

activities: a LIM-domain protein (Solyc04g077780), a JAZ1 TF (Solyc12g009220), and a

GA 20-oxidase (Solyc03g006880) (Fig. 6). LIM-domain proteins have been implicated in

a variety of functions including regulation of the cell cycle and organ size in Arabidopsis

(Li et al., 2008). GA 20-oxidase encodes a key GA biosynthetic enzyme, which acts to

promote cell elongation (Hisamatsu et al., 2005; De Lucas et al., 2008) and thus,

determinacy during leaf morphogenesis of compound leaves, such as those of tomato

(Hay et al., 2002). Moreover, JAZ proteins act as transcriptional repressors and are a

central hub in the signaling circuit that integrates environmental cues, such as light

quality, to balance growth and defense (reviewed in Hou et al., 2013). Finally, it is

noteworthy to highlight that abiotic cues such as light quality and ABA have been shown

to interact and modulate the activity of GA 20-ox and JAZ, and the Arabidopsis LIM-

domain protein DA1, respectively, thereby establishing a conceptual means of

environmental regulation of leaf thickness patterning. Taken together with higher

endopolyploidy levels, the shared expression patterns for these genes between both thick

ILs and the Sp parent suggests that leaf thickness results from an alteration in the

trajectory of cellular events during leaf ontogeny, specifically, the duration of cell

proliferation, and the timing and extent of cell expansion.

Gene expression networks point to distinct leaf ontogeny in ILs 2-5 and 4-3

Since we observed a set of shared DEGs in lines 2-5 and 4-3, we hypothesized that

general patterns of leaf ontogeny may also be shared between these lines, suggesting a

core shared trajectory of leaf thickness patterning. However, we found that Dynamic

Bayesian Networks of gene co-expression in ILs 2-5 and 4-3 are largely distinct (Datasets

S8, S9; Fig. 7A, D).

For example, central to the organ-specific network of IL2-5 is an SBP-box domain gene,

SBP 04g (SQUAMOSA promoter binding protein, Solyc04g064470) which is highly

expressed throughout leaf development in IL2-5 (Fig. S6, Fig. 7A,B). SBP transcription

factors regulate various aspects of plant growth by controlling the rate and timing of

developmental events, including leaf initiation rate (reviewed in Preston and Hileman,

2013). Further, the promoters of IL2-5 DEGs are enriched for SBP motifs (Dataset S7)

supporting the central role of this group of transcription factors during IL2-5 leaf

ontogeny. Interestingly, GO terms for "photosynthesis" and "translation" are enriched

among P4 up-regulated genes. This observation suggests that processes associated with

cell differentiation (i.e. photosynthetic gene function and protein translation) are

precociously activated in IL2-5 relative to domesticated tomato and supports a hypothesis

whereby the overall schedule of leaf developmental events may be hastened in IL2-5.

In contrast, a central node in the IL4-3 co-expression network is a BEL1-like 04g

(Solyc04g080780). BEL1-like homeodomain proteins interact with class I KNOX

transcription factors to pattern the SAM and lateral organs, including leaf complexity

(Kimura et al., 2008; Hay and Tsiantis, 2010) and the extent of lobing and serrations

(Kumar et al., 2007). Like *S. pennellii*, IL4-3 leaflets have significantly smoother margins

(fewer serrations) than M82, as reflected in increased circularity (Fig. 4B; Holtan and

Hake, 2003; Chitwood et al., 2013).

These distinct dynamic patterns of leaf ontogeny that each IL shares with the desert-

adapted parent may reflect aspects of leaf development unrelated to the patterning of leaf

thickness, such as the patterning of leaf complexity and leaflet shape in IL4-3.

Alternatively, it is also possible that the core mechanism of leaf thickness patterning is

achieved by regulation of the timing and extent of cellular activities, such as the balance

between cell proliferation and the onset of cell expansion and endoreduplication, with a

number of potential molecular networks needed to accomplish these roles. An

observation supporting this model is the fact that IL2-5 and IL4-3 have non-overlapping

sets of cell cycle related DEGs. This hypothesis is consistent with the additive

phenotypes of IL2-5/IL4-3 double homozygotes (Fig. 4, Fig. S4), whereby IL-specific

regulators may converge on a common set of targets to regulate cell size and shape, and

ultimately leaf thickness.

Materials and Methods

Plant material and growth conditions

Seeds for 76 *S. pennellii* introgression lines (LA4028-LA4103; Eshed and Zamir, 1995)

and the *S. lycopersicum* domesticated variety M82 (LA3475) were obtained either from

Dr. Neelima Sinha (University of California, Davis) or from the Tomato Genetics

Resource Center (University of California, Davis). All seeds were treated with 50%

bleach for 3 min, rinsed with water and germinated in Phytatrays (P1552, Sigma-

Aldrich). Seeds were left in the dark for 3 days, followed by 3 days in light, and finally

transferred to greenhouse conditions in 50-plug trays. Hardened plants were transplanted

to field conditions at the Bradford Research Station in Columbia, MO (May 21, 2014)

with 3 m between rows and about 1 m spacing between plants within rows. A non-

experimental M82 plant was placed at both ends of each row, and an entire row was

placed at each end of the field to reduce border effects on experimental plants. The final

design had 15 blocks, each consisting of 4 rows with 20 plants per row. Each of the 76

ILs and 2 experimental M82 plants were randomized within each block. IL6-2 was

excluded from final analyses due to seed stock contamination. For the analysis of leaf

primordia by confocal microscopy and RNA-Seq, IL2-5, IL4-3, M82, and *S. pennellii*

seeds were germinated as above and transferred to pots in controlled growth chamber

conditions: irradiance at 400 μmol/m^2/s, 23 $^\circ$C, 14-hour days. Growth conditions for the

drought phenotyping experiment were irradiance of 200 μmol/m^2/s at a daytime

temperature of 22 $^\circ$C and 18 $^\circ$C at night.

Whole-plant phenotyping under drought

The LemnaTec Scanalyzer plant phenotyping facility at the Donald Danforth Plant

Science Center (LemnaTec GmbH, Aachen, Germany) was used to phenotype

approximately 3-week old *S. lycopersicum* and *S. pennellii* plants (n = 8/genotype)

subjected to one of three watering regimes: 40 % field capacity, 20 % field capacity, and

no watering (0 % field capacity). Top view images of each plant taken every second night

over 16 nights were analyzed using custom pipelines in Lemna Launcher (LemnaTec

software) to extract total plant pixel area (a proxy for biomass).

Trait measurements

After flowering (July 2014), four fully expanded adult leaves were harvested from each

plant; the adaxial (upper) surfaces of distal lateral leaflets harvested from the left side of

the rachis were scanned with a flatbed scanner to obtain raw JPG files. The middle

portion of each leaflet was then attached on a custom-build dual confocal profilometer

device (Fig. S1) and the thickness of each leaflet was measured across the leaflet surface

at a resolution of 1 mm^2. Median thickness was calculated across each leaflet using

values in the range (0 mm < thickness < 2 mm) and these median values were averaged

across four leaflets per plant to arrive at a single robust metric of leaf thickness. Finally,

entire leaflets were dried and their dry mass used to calculate leaf mass per area (LMA)

for each leaflet. Leaflet outline scans were processed using custom macros in Image J

(Abramoff et al., 2004) to segment individual leaflets and to threshold and binarize each

leaflet image. Shape descriptors area, aspect ratio, roundness, circularity, and solidity

(described in detail in Chitwood et al., 2013) were extracted from binary images.

Additionally, elliptical Fourier descriptors (EFDs) for leaflet outlines were determined

using SHAPE (Iwata and Ukai, 2002). For this analysis 20 harmonics with 4 coefficients

each were used to derive principal components (PC) that describe major trends in the

shape data.

Elemental profiling (ionomics)

Distal lateral leaflets of fully expanded young (Y) and old (O) leaves of the same plants

as above were collected from five individuals of each genotype. Whole leaflets were

weighed and digested in nitric acid at 100 ^{o}C for 3 hours. Elemental concentrations were

measured using an inductively coupled plasma mass spectrometer (ICP-MS, Elan DRC-e,
Perkin Elmer) following the procedure described in Ziegler et al. (2012). Instrument
reported concentrations were corrected for losses during sample preparation and changes
in instrument response during analysis using Yttrium and Indium internal standards and a
matrix-matched control run every tenth sample. Final concentrations were normalized to
sample weight and reported in mg analyte per kilogram tissue.
598
Statistical analyses and data visualization

599
All statistical analysis and visualization was carried out using R packages (R Core Team,
2013). QTL were identified using the mixed effect linear model packages lme4 (Bates et
al., 2014) and lmerTest (Kuznetsova et al., 2015) with M82 as intercept, IL genotype as a
fixed effect, and field position attributes (block, row, and column) as random effects.
Only effects with significant variance ($p < 0.05$) were included in the final models. For
elemental composition data, leaf age ("young" and "old") was also included as a random
effect unless the variance due to age was the greatest source of variance; in such cases,
young and old samples were analyzed separately. Heritability values represent the
relative proportion of variance due to genotype. For the quantification of organ volume
parameters and photosynthesis measurements, linear models were used to test the effect
of genotype. All plots were generated with the package ggplot2 (Wickham, 2009).
611
Trait correlations and hierarchical clustering

612
For trait correlation analyses we included all traits reported in this study and measured on
the same set of field-grown IL individuals (leaf thickness, LMA, leaflet shape traits,
elemental profiles). We also included several sets of meta-data detailed in Dataset S3,
including DEV (developmental), MOR (morphological), MET (fruit pericarp metabolite
content), ENZ (enzyme activity), and SED (seed metabolite content) related traits (from
Chitwood et al., 2013 and references within). Spearman correlation coefficients (rho)
were calculated between each pair of traits using the rcorr function in Hmisc (Harrell et
al., 2015) and p-values for the correlations were corrected for False Discovery Rate using
Benjamini Hochberg (Dataset S4). Hierarchical clustering and visualization of significant

correlation (q < 0.05) of leaf thickness and LMA were clustered (hierarchical "ward"

algorithm) and visualized using pheatmap (Kolde, 2015).

Estimation of nuclear size profiles by flow cytometry

Distal lateral leaflets were harvested from the 7[th] leaf of greenhouse-grown 6-week-old

plants and immediately chopped in 1 mL of ice-cold buffer LB01 as in Doležel et al.

(2008). The resulting fine homogenate was filtered through a 30 um Partec CellTrics

filter (5004-004-2326) and incubated with 50 ug/mL propidium iodide (Thermo Fisher,

P21493) and 50 ug/mL RNase A (Qiagen, 19101) for 20 min on ice. Fluorescence scatter

data was collected without gating using a BD Acuri CS6 instrument (BD Biosciences).

Plots of event count as a function of fluorescence area were used to estimate the

proportion of nuclei of sizes corresponding to 2C, 4C, and 8C in each genotype.

Confocal microscopy, 3D-reconstructions, and organ volume quantification

For mature leaf cross-sections, field-grown leaves were fixed in FAA (4 % formaldehyde,

5 % glacial acetic acid, 50 % ethanol), vacuum infiltrated, dehydrated through an ethanol

series, rehydrated to 100 % water, stained in 0.002 % propidium iodide (Thermo Fisher,

P21493) for 2 hours, dehydrated to 100 % ethanol, and finally cleared in 100 % methyl

salicylate (Sigma, M6752) for 7 days. Hand-sections were visualized with a Leica SP8

laser scanning confocal microscope using white light laser excitation at 514 nm with a

20X objective. Two partially overlapping images were captured for each cross-section

and merged into a single image using the "Photomerge" function in Adobe Photoshop CC

2014 (Adobe Systems Incorporated). For the quantification of P3 leaf primordium

dimensions, shoot apices (shoot apical meristem and P1-P4) of 14 day-old seedlings

grown in controlled conditions were excised, fixed, processed, and stained as detailed for

leaf cross sections above. Confocal stacks were obtained at software-optimized intervals,

and exported as TIFF files. Raw stack files were imported into MorphoGraphX (Reulle et

al., 2015). After Gaussian filtering, the marching cubes surface reconstruction function

was used (cube size = 5 μm and threshold = 7,000). The resulting surface mesh was

smoothed and subdivided twice and exported as a PLY file. To minimize the effects of

trichomes on P3 volume, all meshes were trimmed in MeshLab (Cignoni et al., 2008).

Volume, length, and diameter of processed P3 meshes were calculated using custom

scripts in MatLab (MathWorks, Inc.). Briefly, first, we detected the boundary of each

hole and calculated its centroid point. We connected boundary points of each hole to its

centroid and filled the triangle faces. After filling all the holes, 3D mesh represents the

closed surface. Then we calculated the volume based on the divergence theorem, which

makes use of the fact that the inside fluid expansion equals the flux ($\vec{F}$) of the fluid out of

the surface (S). When the flux is $\vec{F} = (x, 0,0)$, the volume is $V = \oiint (\vec{F} \cdot \vec{n})\, dS$, where $\vec{n}$

is normal vector. Thus, for each triangle, we computed the normal vector $\vec{n} =$

(x_n, y_n, z_n), the area A, and the centroid point $P = (x_p, y_p, z_p)$. The volume V is the

summation of $Ax_n x_p$ for all triangles. To estimate organ arch length we made use of the

fact that the Laplace-Beltrami eigenfunctions are deformation invariant shape descriptor

(Rustamov, 2007). We thus employed its first eigenfunction, which is associated with the

smallest positive eigenvalue and discretized the eigenfunction values into 50 sets to

compute the centroid point to each set. We fit a cubic function by fixing two end-point

constraints to those centroid points to get a smooth principle median axis. Note that the

two end points were manually adjusted to correct for artifacts. The length of this axis is

used to quantify the length of the organ. Finally, we calculated mean organ diameter as

$d = 2\sqrt{\dfrac{V}{\pi L}}$.

RNA-Seq library preparation and sequencing

Apices of fourteen day-old IL2-5, IL4-3, M82, and *S. pennellii* (Sp) plants grown in a

randomized design under controlled growth conditions were hand-dissected under a

dissecting microscope to separate plastochrons P4, P3, P2, and P1+SAM organs

corresponding approximately to leaves L5 – L8. For Sp plants we were not able to

separate P2 primordia from the apex and so we obtained P4, P3, and SAM+P1+P2

samples. Dissected organs were removed from the apex in less than 60 seconds and

immediately fixed in 100 % ice-cold acetone to preserve the integrity of RNA in the

sample. Each biological replicate is a pool of 10 individuals, and a total of 5 biological

replicates were obtained for each genotype/organ combination. RNA was extracted using

PicoPure RNA Isolation Kit (Thermo Fisher Scientific, MA, USA) according to the

manufacturer's protocol with the optional on-column DNase treatment. RNA integrity

(RIN) was assessed by running all samples on an Agilent RNA 6000 Pico chip (Agilent

Technologies, CA, USA) and three biological replicates with RIN > 7.0 were selected for

further processing. Double stranded cDNA amplified using Clontech SMARTer PCR

cDNA synthesis kit (634926, TaKaRa Bio USA) was fragmented for 15 min using

Fragmentase (M0348, New England Biolabs) and processed into Illumina sequencing

libraries as follows: the ends of 1.5X AMPure XP bead (A63880, Agencourt) purified

fragmented DNA was repaired with End Repair Enzyme Mix (E6050 New England

Biolabs) and Klenow DNA Polymerase (M0210, NEB), followed by dA-tailing using

Klenow 3'-5' exonuclease (M0212, NEB). The Illumina TruSeq universal adapter dimer

was ligated to library fragments with rapid T4 DNA Ligase (L6030-HC-L, Enzymatics)

followed by 3 rounds of 1X AMPure XP bead purification to remove unligated adapter.

Finally, libraries were enriched and indexed by PCR using Phusion HiFi Polymerase mix

(M0531, NEB). Illumina libraries were quantified using a nanodrop, pooled to a final

concentration of 20 nM and sequenced as single end 100 bp reads on Illumina HiSeq2500

at the Washington University in St. Louis School of Medicine Genome Technology

Access Center (https://gtac.wustl.edu/).

RNA-Seq data analysis

Adapters and low quality bases were removed using Trimmomatic (Bolger et al., 2014)

with default parameters. Trimmed reads were mapped to the ITAG2.3 *Solanum*

lycopersicum genome

(https://solgenomics.net/organism/Solanum_lycopersicum/genome; The Tomato Genome

Consortium, 2012) using bowtie2 (Langmead and Salzberg, 2012) to obtain SAM files.

After sorting and indexing of SAM files, BAM files files were generated using samtools

commands (Li and Handsaker et al., 2009). The BEDtools multicov tool (Quinlan and

Hall, 2010) was then used to obtain read counts per annotated gene for each sample.

Subsequent analysis was done with the R package edgeR (Robinson et al., 2010). After

normalization for library size 20,231 genes with at least one count per million reads

across three samples were retained for further analysis. Lists of Differentially Expressed

Genes (DEGs) were generated between pairwise sample combinations with q-value <

0.05. For IL2-5 and IL4-3 at P1, P3, and P4 stages, we identified genes that are

differentially expressed relative to M82 in both the IL and the Sp parent to interrogate

Sp-like changes in gene expression in the ILs. For P2, the list of DEG in each IL reflects

changes relative to M82 only (Dataset S5).

Gene Ontology, Mapman, and promoter motif enrichment analyses

Lists of IL organ-specific DEGs were interrogated for enrichment of Gene Ontology

terms using agriGO (http://bioinfo.cau.edu.cn/agriGO/; Du et al., 2010) with default

parameters (Fisher's exact significance test and Yekutieli FDR adjustment at $q < 0.05$).

We further divided DEG gene lists into IL up-regulated and down-regulated genes and

report significant terms in Dataset S6. We tested IL organ-specific DEGs for enrichment

of annotated promoter motifs using a custom R script (Dr. Julin Maloof). Briefly,

functions in the Bioconductor Biostrings package (Pages et al., 2016) were implemented

to count the frequency of 100 known motifs in the promoters of DEGs (1000 bp upstream

sequence) and calculate p-values for enrichment based on these counts. We report exact

matches of known motifs and motifs with up to 1 mismatch in IL up-regulated and down-

regulated organ-specific gene sets (Dataset S7).

IL organ-specific gene network inference

To infer IL organ-specific networks (Fig. 7A-D, Dataset S8), we selected DEGs between

IL2-5/M82 (IL4-3/M82) and Sp/M82 for each organ (P1, P3, P4) (q value < 0.05). Since

co-expression analysis can inform the likelihood that genes interact, or participate in the

same functional pathway, the selected genes for each IL (IL2-5 or IL4-3) and each organ

were clustered based on their co-expression across genotypes. To perform clustering, the

Silhouette index (Rousseeuw, 1987) followed by K-means (MacQueen, 1967) were

applied. After clustering, networks were inferred as in de Luis Balaguer et al. (2017).

Briefly, for each DEG, we identified a set of potential regulators and measured the

likelihood of gene-target regulation using a Bayesian Dirichlet equivalence uniform (Boutine, 1991). Genes that had the highest value of the Bayesian Dirichlet equivalence uniform were chosen as regulators, and of these only transcription factors (as annotated by Suresh et al., 2014) were further considered as regulatory (source) nodes. To obtain the final IL2-5 and IL4-3 organ-specific networks, the networks for each cluster were connected. For this, we found regulations among the cluster hubs (node of each individual network with the largest degree of edges leaving the node) by using the same Bayesian Dirichlet equivalence uniform metric. In addition, we implemented a score to estimate whether the inferred interactions were activations or repressions. The score was calculated for each edge and it measured the ratio between i) the conditional probability that a gene is expressed given that its regulator was expressed in the prior time point, and ii) the conditional probability that a gene is expressed given that its regulator was not expressed in the prior time point. If the first conditional probability is larger than the second one, then the parent was found to be an activator and *vice versa*. In the case of a tie, the edge was found to have an undetermined sign. Networks for each organ were jointly visualized in Cytoscape (Shannon et al., 2003).

Dynamic IL network construction

To infer dynamic IL networks (Fig. 7E-F, Dataset S9), we selected DEGs between IL2-5/M82 or IL4-3/M82 and Sp/M82 for each organ (P1, P3, P4) (q value < 0.05 or (FC > 2.0 and q value < 0.2)). All DEG in the IL2-5 or IL4-3 were clustered in four groups, corresponding to the four developmental stages: each gene was assigned to the developmental stage where it showed the maximum expression. A network was then inferred for each developmental stage as described for the IL organ-specific networks. To ensure that all potential regulators of each gene were considered, genes from the preceding developmental stage were included in the inference of the network of each developmental stage. The final network for each IL was visualized in Cytoscape (Shannon et al., 2003).

Accession Numbers

An NCBI SRA accession number will be provided upon publication.

Supporting Data

Supplemental Figure S1. Dual confocal profilometer device used to measure leaf

thickness.

Supplemental Figure S2. Comparison of leaf thickness of select ILs as a function of

shoot position and field vs. greenhouse conditions.

Supplemental Figure S3. Representative leaf cross-sections and flow cytometry of leaf

6/7 for 10 ILs harboring leaf thickness QTLs grown in greenhouse conditions.

Supplemental Figure S4. Mean dimensions of palisade and spongy mesophyll cell

layers in select thick leaf ILs. Representative flow cytometry histograms of leaf 7 and

post-flowering leaves from each genotype.

Supplemental Figure S5. Representative shoot apex reconstructions highlighting the

appearance of early and late stage leaf primordia for each genotype in Fig. 4

Supplemental Figure S6. Summary of differentially expressed genes in IL2-5 and IL4-3.

Supplemental Figure S7. Expression profiles of differentially expressed putative

transcription factors in ILs 2-5 and 4-3.

Supplemental Figure S8. Summary of enriched promoter motifs among differentially

expressed genes in ILs 2-5 and 4-3.

Supplemental Dataset S1. Trait value estimates and heritability for leaf thickness, LMA,

and leaflet shape across the IL panel.

Supplemental Dataset S2. Trait value estimates and heritability for elemental

concentration across the IL panel.

Supplemental Dataset S3. Summary of all measured and meta-data traits used in

correlation matrix.

Supplemental Dataset S4. Pairwise trait correlation matrix including significance

values.

Supplemental Dataset S5. List of differentially expressed genes ($q < 0.05$) in each organ

(P1 – P4) for the comparison: (M82/IL) overlapping with (M82/*S. pennellii*).

Supplemental Dataset S6. List of significantly enriched ($q < 0.05$) Gene Ontology (GO)

terms for gene sets listed in Dataset S5.

Supplemental Dataset S7. List of enriched (q < 0.05) promoter motifs for gene sets in

Dataset S5.

Supplemental Dataset S8. List of organ-specific (P1, P3, P4) gene interactions for IL2-5

and IL4-3.

Supplemental Dataset S9. List of dynamic gene interactions for IL2-5 and IL4-3.

Acknowledgments

S. pennellii introgression line panel seeds were provided by Dr. Neelima Sinha
(University of California, Davis) and the Tomato Genetics Resource Center (University
of California, Davis). We would like to thank Dr. Ivan Baxter and Dr. Greg Ziegler
(Danforth Plant Science Center) for generating elemental profile data, and Dr. Julin
Maloof (University of California, Davis) for sharing custom promoter enrichment
analysis scripts. We acknowledge the advice and assistance of Dr. Noah Fahlgren, Dr.
Malia Gehan, and Melinda Darnell (Danforth Plant Science Center) with drought
phenotyping experiments. We thank Dr. Elizabeth Kellogg for insightful discussions and
comments on the manuscript. This work was supported by funds from the Donald
Danforth Plant Science Center. RS is supported by an NSF CAREER grant (MCB-
1453130). MHF is supported by an NSF NPGI post-doctoral fellowship (IOS-1523668).

Literature Cited

Abràmoff, M. D., Magalhães, P. J., & Ram, S. J. (2004). Image processing with ImageJ. *Biophotonics International* 11(7), 36-42.

Becker, B. (2007). Function and evolution of the vacuolar compartment in green algae and land plants (Viridiplantae). *International Review of Cytology* 264, 1-24.

John, G. P., Scoffoni, C., & Sack, L. (2013). Allometry of cells and tissues within leaves. *American Journal of Botany* 100(10), 1936-1948.

Bohmert, K., Camus, I., Bellini, C., Bouchez, D., Caboche, M., & Benning, C. (1998). AGO1 defines a novel locus of Arabidopsis controlling leaf development. *The EMBO Journal* 17(1), 170-180.

Brodersen, C. R., & Vogelmann, T. C. (2010). Do changes in light direction affect absorption profiles in leaves?. *Functional Plant Biology*, *37*(5), 403-412.

Brodersen, C. R., Vogelmann, T. C., Williams, W. E., & Gorton, H. L. (2008). A new paradigm in leaf□level photosynthesis: direct and diffuse lights are not equal. *Plant, cell & environment*, *31*(1), 159-164.

Buntine, W. 1991. Theory refinement on Bayesian networks. In *Proceedings of the Seventh conference on Uncertainty in Artificial Intelligence*, 52-60. Morgan Kaufmann Publishers Inc.

Chitwood, D. H., Kumar, R., Headland, L. R., Ranjan, A., Covington, M. F., Ichihashi, Y., ... & Sinha, N. R. (2013). A quantitative genetic basis for leaf morphology in a set of precisely defined tomato introgression lines. *The Plant Cell* 25(7), 2465-2481.

Cignoni, P., Callieri, M., Corsini, M., Dellepiane, M., Ganovelli, F., & Ranzuglia, G. (2008, July). Meshlab: an open-source mesh processing tool. In *Eurographics Italian Chapter Conference* 2008, 129-136.

De Lucas, M., Daviere, J. M., Rodriguez-Falcon, M., Pontin, M., Iglesias-Pedraz, J. M., Lorrain, S., ... & Prat, S. (2008). A molecular framework for light and gibberellin control of cell elongation. *Nature*, *451*(7177), 480-484.

Dhondt, S., Coppens, F., De Winter, F., Swarup, K., Merks, R. M., Inzé, D., ... & Beemster, G. T. (2010). SHORT-ROOT and SCARECROW regulate leaf growth in

Arabidopsis by stimulating S-phase progression of the cell cycle. *Plant Physiology*

154(3), 1183-1195.

Doležel, J., Greilhuber, J., & Suda, J. (2007). Estimation of nuclear DNA content in

plants using flow cytometry. *Nature protocols*, *2*(9), 2233-2244.

Du, Z., Zhou, X., Ling, Y., Zhang, Z., & Su, Z. (2010). agriGO: a GO analysis toolkit for

the agricultural community. *Nucleic Acids Research* 38, W64-W70.

Eggli, U., & Nyffeler, R. (2009). Living under temporarily arid conditions–succulence as

an adaptive strategy. *Bradleya* 27, 13-36.

Eshed, Y., & Zamir, D. (1995). An introgression line population of Lycopersicon

pennellii in the cultivated tomato enables the identification and fine mapping of yield-

associated QTL. *Genetics* 141(3), 1147-1162.

Fridman, E., Carrari, F., Liu, Y. S., Fernie, A. R., & Zamir, D. (2004). Zooming in on a

quantitative trait for tomato yield using interspecific introgressions. *Science* 305(5691),

1786-1789.

Garnier, E., & Laurent, G. (1994). Leaf anatomy, specific mass and water content in

congeneric annual and perennial grass species. *New Phytologist* 128(4), 725-736.

Gibson, A. C. (1982). The anatomy of succulence. In *Crassulacean acid metabolism*. J.

Proc. V ann. Symp. Bot., Univ. Calif., Riverside. *J. Amer. Soc. Plant Physiologists*:

Rockville, USA, 1-17.

Haliński, Ł. P., Kalkowska, M., Kalkowski, M., Piorunowska, J., Topolewska, A., &

Stepnowski, P. (2015). Cuticular wax variation in the tomato (Solanum lycopersicum L.),

related wild species and their interspecific hybrids. *Biochemical Systematics and Ecology*

60, 215-224.

Hay, A., Kaur, H., Phillips, A., Hedden, P., Hake, S., & Tsiantis, M. (2002). The

gibberellin pathway mediates KNOTTED1-type homeobox function in plants with

different body plans. *Current Biology*, *12*(18), 1557-1565.

Hay, A., & Tsiantis, M. (2010). KNOX genes: versatile regulators of plant development

and diversity. *Development* 137(19), 3153-3165.

Hisamatsu, T., King, R. W., Helliwell, C. A., & Koshioka, M. (2005). The involvement

of gibberellin 20-oxidase genes in phytochrome-regulated petiole elongation of

Arabidopsis. *Plant Physiology*, *138*(2), 1106-1116.

Holtan, H. E., & Hake, S. (2003). Quantitative trait locus analysis of leaf dissection in

tomato using Lycopersicon pennellii segmental introgression lines. *Genetics* 165(3),

1541-1550.

Hou, X., Ding, L., & Yu, H. (2013). Crosstalk between GA and JA signaling mediates

plant growth and defense. *Plant cell reports*, *32*(7), 1067-1074.

Huang, W., Xian, Z., Kang, X., Tang, N., & Li, Z. (2015). Genome-wide identification,

phylogeny and expression analysis of GRAS gene family in tomato. *BMC Plant Biology*

15(1), 209.

Ichihashi, Y., Aguilar-Martínez, J. A., Farhi, M., Chitwood, D. H., Kumar, R., Millon, L.

V., ... & Sinha, N. R. (2014). Evolutionary developmental transcriptomics reveals a gene

network module regulating interspecific diversity in plant leaf shape. *Proceedings of the*

National Academy of Sciences 111(25), E2616-E2621.

Iwata, H., & Ukai, Y. (2002). SHAPE: a computer program package for quantitative

evaluation of biological shapes based on elliptic Fourier descriptors. *Journal of Heredity*

93(5), 384-385.

John, G. P., Scoffoni, C., & Sack, L. (2013). Allometry of cells and tissues within leaves.

American Journal of Botany 100(10), 1936-1948.

Kalve, S., Fotschki, J., Beeckman, T., Vissenberg, K., & Beemster, G. T. (2014). Three-

dimensional patterns of cell division and expansion throughout the development of

Arabidopsis thaliana leaves. *Journal of Experimental Botany* 65, 6385-6397.

Kimura, S., Koenig, D., Kang, J., Yoong, F. Y., & Sinha, N. (2008). Natural variation in

leaf morphology results from mutation of a novel KNOX gene. *Current Biology* 18(9),

672-677.

Koenig, D., Jiménez-Gómez, J. M., Kimura, S., Fulop, D., Chitwood, D. H., Headland, L.

R., ... & Tohge, T. (2013). Comparative transcriptomics reveals patterns of selection in

domesticated and wild tomato. *Proceedings of the National Academy of Sciences*

110(28), E2655-E2662.

Kolde, R. (2015). pheatmap: Pretty Heatmaps. R package version 1.0. 2.

Kozuka, T., Kong, S. G., Doi, M., Shimazaki, K. I., & Nagatani, A. (2011). Tissue-

autonomous promotion of palisade cell development by phototropin 2 in

Arabidopsis. *The Plant Cell*, *23*(10), 3684-3695.

Kumar, R., Kushalappa, K., Godt, D., Pidkowich, M. S., Pastorelli, S., Hepworth, S. R.,

& Haughn, G. W. (2007). The Arabidopsis BEL1-LIKE HOMEODOMAIN proteins

SAW1 and SAW2 act redundantly to regulate KNOX expression spatially in leaf

margins. *The Plant Cell* 19(9), 2719-2735.

Kuznetsova, A., Christensen, R. H., Bavay, C., & Brockhoff, P. B. (2015). Automated

mixed ANOVA modeling of sensory and consumer data. *Food Quality and Preference*

40, 31-38.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., ... & Durbin, R.

(2009). The sequence alignment/map format and SAMtools. *Bioinformatics* 25(16),

2078-2079.

Li, Y., Zheng, L., Corke, F., Smith, C., & Bevan, M. W. (2008). Control of final seed and

organ size by the DA1 gene family in Arabidopsis thaliana. *Genes &*

Development, *22*(10), 1331-1336.

Lightfoot, D. A., Mungur, R., Ameziane, R., Nolte, S., Long, L., Bernhard, K., ... &

Young, B. (2007). Improved drought tolerance of transgenic Zea mays plants that express

the glutamate dehydrogenase gene (gdhA) of E. coli. *Euphytica* 156(1-2), 103-116.

de Luis Balaguer, M. A., Fisher, A. P., Clark, N. M., Fernandez-Espinosa, M. G., Moller,

B., Weijers, D., Lohmann, J. U., Williams, C., Lorenzo, O. and Sozzani, R. (2017). An

inference approach combines spatial and temporal gene expression data to predict gene

regulatory networks in Arabidopsis stem cells. *Pre-print* doi:

https://doi.org/10.1101/140269.

McDowell, E. T., Kapteyn, J., Schmidt, A., Li, C., Kang, J. H., Descour, A., ... & Jones,

A. D. (2011). Comparative functional genomic analysis of Solanum glandular trichome

types. *Plant Physiology* 155(1), 524-539.

McHale, N. A. (1992). A nuclear mutation blocking initiation of the lamina in leaves of

Nicotiana syhestris. *Planta* 186(3), 355-360.

McHale, N. A. (1993). LAM-1 and FAT genes control development of the leaf blade in

Nicotiana sylvestris. *The Plant Cell* 5(9), 1029-1038.

MacQueen, J. (1967, June). Some methods for classification and analysis of multivariate

observations. In *Proceedings of the fifth Berkeley symposium on mathematical statistics*

and probability 1(14), 281-297.

Muir, C. D., Conesa, M. À., Roldán, E. J., Molins, A., & Galmés, J. (2016). Weak

coordination between leaf structure and function among closely related tomato species.

New Phytologist.

Muir, C. D., Pease, J. B., & Moyle, L. C. (2014). Quantitative genetic analysis indicates

natural selection on leaf phenotypes across wild tomato species (Solanum sect.

Lycopersicon; Solanaceae). *Genetics* 198(4), 1629-1643.

Nakazato, T., Warren, D. L., & Moyle, L. C. (2010). Ecological and geographic modes of

species divergence in wild tomatoes. *American Journal of Botany* 97(4), 680-693.

Nelson, E. A., Sage, T. L., & Sage, R. F. (2005). Functional leaf anatomy of plants with

crassulacean acid metabolism. *Functional Plant Biology* 32(5), 409-419.

Nicotra, A. B., Leigh, A., Boyce, C. K., Jones, C. S., Niklas, K. J., Royer, D. L., &

Tsukaya, H. (2011). The evolution and functional significance of leaf shape in the

angiosperms. *Functional Plant Biology* 38(7), 535-552.

Niinemets, Ü., Díaz-Espejo, A., Flexas, J., Galmés, J., & Warren, C. R. (2009). Role of

mesophyll diffusion conductance in constraining potential photosynthetic productivity in

the field. *Journal of Experimental Botany* 60(8), 2249-2270.

Ogburn, R., & Edwards, E. J. (2010). The ecological water-use strategies of succulent

plants. *Advances in Botanical Research* 55, 179-225.

Oguchi, R., Hikosaka, K., & Hirose, T. (2005). Leaf anatomy as a constraint for

photosynthetic acclimation: differential responses in leaf anatomy to increasing growth

irradiance among three deciduous trees. *Plant, Cell & Environment* 28(7), 916-927.

Pieruschka, R., & Poorter, H. (2012). Phenotyping plants: genes, phenes and machines.

Functional Plant Biology 39(11), 813-820.

Poorter, H., Niinemets, Ü., Poorter, L., Wright, I. J., & Villar, R. (2009). Causes and

consequences of variation in leaf mass per area (LMA): a meta-analysis. *New Phytologist*

182(3), 565-588.

Preston, J. C., & Hileman, L. (2013). Functional evolution in the plant SQUAMOSA-

PROMOTER BINDING PROTEIN-LIKE (SPL) gene family. *Frontiers in Plant Science*

4, 80.

Quinlan, A. R., & Hall, I. M. (2010). BEDTools: a flexible suite of utilities for comparing

genomic features. *Bioinformatics* 26(6), 841-842.

Ranjan, A., Budke, J., Rowland, S. D., Chitwood, D. H., Kumar, R., Carriedo, L. G., ... &

Sinha, N. (2016). eQTL regulating transcript levels associated with diverse biological

processes in tomato. *Plant Physiology* 172, 328-340.

de Reuille, P. B., Routier-Kierzkowska, A. L., Kierzkowski, D., Bassel, G. W.,

Schüpbach, T., Tauriello, G., ... & Burian, A. (2015). MorphoGraphX: a platform for

quantifying morphogenesis in 4D. *Elife* 4, e05864.

Robinson, M. D., McCarthy, D. J., & Smyth, G. K. (2010). edgeR: a Bioconductor

package for differential expression analysis of digital gene expression data.

Bioinformatics 26(1), 139-140.

Roderick, M. L., Berry, S. L., Noble, I. R., & Farquhar, G. D. (1999). A theoretical

approach to linking the composition and morphology with the function of leaves.

Functional Ecology 13(5), 683-695.

Rousseeuw, P. J. (1987). Silhouettes: a graphical aid to the interpretation and validation

of cluster analysis. *Journal of Computational and Applied Mathematics* 20, 53-65.

Rustamov, R. M. (2007, July). Laplace-Beltrami eigenfunctions for deformation invariant

shape representation. In *Proceedings of the fifth Eurographics symposium on Geometry*

processing, 225-233. Eurographics Association.

Sack, L., & Frole, K. (2006). Leaf structural diversity is related to hydraulic capacity in

tropical rain forest trees. *Ecology* 87(2), 483-491.

Schauer, N., Semel, Y., Roessner, U., Gur, A., Balbo, I., Carrari, F., ... & Willmitzer, L.

(2006). Comprehensive metabolic profiling and phenotyping of interspecific

introgression lines for tomato improvement. *Nature Biotechnology* 24(4), 447-454.

Semel, Y., Nissenbaum, J., Menda, N., Zinder, M., Krieger, U., Issman, N., ... & Zamir,

D. (2006). Overdominant quantitative trait loci for yield and fitness in tomato.

Proceedings of the National Academy of Sciences 103(35), 12981-12986.

Shannon, P., Markiel, A., Ozier, O., Baliga, N. S., Wang, J. T., Ramage, D., ... & Ideker,

T. (2003). Cytoscape: a software environment for integrated models of biomolecular
interaction networks. *Genome Research* 13(11), 2498-2504

Smith, S. D., Monson, R. K., & Anderson, J. E. (1997). Plant processes and responses to
stress. In *Physiological Ecology of North American Desert Plants,* 45-71. Springer Berlin
Heidelberg.

Suresh, B. V., Roy, R., Sahu, K., Misra, G., & Chattopadhyay, D. (2014). Tomato
genomic resources database: an integrated repository of useful tomato genomic
information for basic and applied research. *PLoS One* 9(1), e86387.

Taji, T., Ohsumi, C., Iuchi, S., Seki, M., Kasuga, M., Kobayashi, M., ... & Shinozaki, K.
(2002). Important roles of drought☐and cold☐inducible genes for galactinol synthase in
stress tolerance in Arabidopsis thaliana. *The Plant Journal* 29(4), 417-426.

Terashima, I., Hanba, Y. T., Tholen, D., & Niinemets, Ü. (2011). Leaf functional
anatomy in relation to photosynthesis. *Plant Physiology* 155(1), 108-116.

Tomato Genome Consortium. (2012). The tomato genome sequence provides insights
into fleshy fruit evolution. *Nature* 485(7400), 635-641.

Tsuge, T., Tsukaya, H., & Uchimiya, H. (1996). Two independent and polarized
processes of cell elongation regulate leaf blade expansion in Arabidopsis thaliana (L.)
Heynh. *Development* 122(5), 1589-1600.

Tsukaya, H. (2003). Organ shape and size: a lesson from studies of leaf morphogenesis.
Current Opinion in Plant Biology 6(1), 57-62.

Wickham, H. (2009). ggplot2: elegant graphics for data analysis. Springer New York
1(2), 3.

von Willert, D. J. (1992). *Life strategies of succulents in deserts: with special reference*
to the Namib Desert. CUP Archive.

Wright, I. J., Reich, P. B., Westoby, M., Ackerly, D. D., Baruch, Z., Bongers, F., ... &
Flexas, J. (2004). The worldwide leaf economics spectrum. *Nature* 428(6985), 821-827.

Wuyts, N., Massonnet, C., Dauzat, M., & Granier, C. (2012). Structural assessment of the
impact of environmental constraints on Arabidopsis thaliana leaf growth: a 3D approach.
Plant, Cell & Environment 35(9), 1631-1646.

Ziegler, G., Terauchi, A., Becker, A., Armstrong, P., Hudson, K., & Baxter, I. (2013).

Ionomic screening of field-grown soybean identifies mutants with altered seed elemental
composition. *The Plant Genome* 6(2).

 Figure Legends

Figure 1. Desert-adapted tomato plants have thicker leaves than domesticated

tomato and are resistant to drought. (**A**) Thickness across leaflet blades of

domesticated (*S. lycopersicum,* M82) and desert-adapted (*S. pennellii*) tomatoes

measured with a custom-built dual confocal profilometer device (Supplemental Figure 1).

Median thickness of the *S. lycopersicum* leaflet shown here is 211 μm, and 294 μm for *S.*

pennellii. (**B**) Confocal images of propidium iodide stained leaflet cross-sections; scale

bar is 200 μm. (**C**) Total shoot area normalized by taking the square root of pixels from

top view phenotyping images over 16 days in three water treatments (n=8). Gray shading

reflects standard error.

1042

Figure 2. Quantitative Trait Loci for leaf thickness in tomato. (**A**) Leaflet thickness

values across the *S. pennellii* introgression line panel. Colors indicate level of

significance in comparisons of each IL with M82 (arrow). (**B**) Significant correlations

(Spearman's rho) between leaf thickness ("Thickness"), or leaf mass per area ("LMA")

and a suite of other traits across the *S. pennellii* IL panel (q < 0.05). Traits are grouped by

type: ION, elemental profile; MOR, morphological; DEV, developmental; ENZ, enzyme

activity; SED, seed metabolite content (Datasets S3 and S4).

1050

1051 **Figure 3. Anatomical manifestations of thicker leaves.** (**A**) Confocal images of

1052 propidium iodide stained cross-sections of field-grown M82, select ILs and S. pennellii

1053 grown in greenhouse conditions; scale bars are 50 μm**.** (**B**) Representative leaf thickness

1054 plots and (C) leaflet binary images of field-grown plants as for (A).

1055

Figure 4. Leaf morphology and ploidy of IL2-5/IL4-3 double homozygote plants. (**A**)

Representative propidium iodide-stained leaflet cross-sections (left) and thickness

measurements (right) for the 7th leaf of greenhouse-grown M82 and double homozygous

IL2-5/IL4-3 plants (n = 10). Scale bars are 200 μm; ** p < 0.01. (**B**) Circularity (ratio of

area to the square of the perimeter) of distal lateral leaflets as in (A). Silhouettes of

representative M82 and IL2-5/IL4-3 leaflets are shown above bars. Letters indicate

statistical significance in each pairwise genotype comparison ($p < 0.05$). **(C)** Distribution

of relative nuclear sizes reflecting endoreduplication in leaflets as in (A) and (B) (n = 5).

Letters denote statistical significance between pairwise genotype comparisons at each

ploidy level. (D) Leaf plastochron P3 dimensions calculated from 3D surface

reconstructions of vegetative shoot apices (n = 9; * $p < 0.05$, ** $p < 0.01$ relative to

M82).

1068

Figure 5. Comparative transcriptomics of leaf development in two thick ILs and

their parents. (A) Successive stages of leaf development (plastochrons P1-P4 colored as

in legend in (B)) were dissected from M82, *S. pennellii* (Sp) and thick ILs 2-5 and 4-3.

(B) Principal Components Analysis (PCA) of normalized RNA-Seq read counts.

(C) Venn diagrams (not to scale) depict an overview of differentially expressed genes

(DEGs, $q < 0.05$) that are shared in each IL and the Sp parent relative to M82. The

number of DEGs unique to each organ is shown within elipses and those common to all

organs, in the center. The total number of DEGs at each plastochron stage is shown

outside ellipses.

1078

Figure 6. Comparative expression profiles of genes in three functional categories

across leaf development (P1 – P4) in thick ILs 2-5 and 4-3: (A) Transcription factors

common to both ILs. **(B)** Genes involved in leaf development in tomato (as in Ichihashi

et al., 2014), and **(C)** Gene annotated to encode components of the cell cycle or ubiquitin

protesaome pathway (contain one of the terms "cell cycle", "cyclin", "ubiquitin", "E2F",

"mitosis", "mitotic", "SKP"). Plastochron stages with statistically significant DE ($q <$

0.05) relative to M82 are marked with an asterisk. Genes, which are differentially

expressed in at least one stage in both ILs are marked in bold.

1087

1088   **Figure 7. Select leaf development gene regulatory sub-networks for (A-C) IL2-5 and**

1089   **(D-F) IL4-3.** Sub-networks for regulators central to more than one plastochron stage are

1090   shown in **(A)** and **(D)**. A GA 20-oxidase gene (GA 20-ox 03g, Solyc03g006880) and its

1091   regulators in each IL **(B)** and **(E)**. Sub-networks of dynamic gene regulatory networks,

showing interactions of an AUX/IAA TF (AUX/IAA 12g, Solyc12g096980) with other source nodes **(C)** and **(F)**. Gene IDs of highlighted nodes: SBP-box 04g, Solyc04g064470; BEL1-like 04g, Solyc04g080780; MADS-box 12g, Solyc12g087820; MYB TF 05g, Solyc05g007710; bHLH TF 04g, Solyc04g074810; GRAS 08g, Solyc08g014030; Myb 07g, Solyc07g052490; WRKY 05g, Solyc05g015850; AUX/IAA 06g, Solyc06g008580; Myb 03g, Solyc03g005570; Myb 08g, Soly08g005870; WRKY 02g, Solyc02g080890. Nodes and edges are colored according to legend.

Figure 1. Desert-adapted tomato plants have thicker leaves than domesticated tomato and are resistant to drought. (A) Thickness across leaflet blades of domesticated (*S. lycopersicum*) and desert-adapted (*S. pennellii*) tomatoes measured with a custom-built dual confocal profilometer device (Fig. S1). Median thickness of the *S. lycopersicum* leaflet shown here is 211 μm, and 294 μm for *S. pennellii*. **(B)** Confocal images of propidium iodide stained leaflet cross-sections; scale bar is 200 μm. **(C)** Total shoot area normalized by taking the square root of pixels from top-view phenotyping images taken over 16 days in three water treatments (n = 8). Gray shading around each colored line reflects standard error.

Figure 2. Quantitative Trait Loci for leaf thickness in tomato. (A) Leaflet thickness values across the *S. pennellii* introgression line panel. Colors indicate level of significance in comparisons of each IL with M82 (arrow). **(B)** Significant correlations (Spearman's rho) between leaf thickness ("Thickness"), or leaf mass per area ("LMA") and a suite of other traits across the *S. pennellii* IL panel (q < 0.05). Traits are grouped by type: ION, elemental profile; MOR, morphological; DEV, developmental; ENZ, enzyme activity; SED, seed metabolite content (Datasets S3 and S4).

Figure 3. Anatomical manifestations of thicker leaves. **(A)** Confocal images of propidium iodide-stained cross-sections of field-grown M82, select ILs and *S. pennellii* grown in greenhouse conditions; scale bars are 50 μm. **(B)** Representative leaf thickness plots and **(C)** leaflet binary images of field-grown plants as for (A).

Figure 4. Leaf morphology and ploidy of IL2-5/IL4-3 double homozygote plants. (A) Representative propidium iodide-stained leaflet cross-sections (left) and thickness measurements (right) for the 7th leaf of greenhouse-grown M82 and double homozygous IL2-5/IL4-3 plants (n = 10). Scale bars are 200 μm; ** p < 0.01. **(B)** Circularity (ratio of area to the square of the perimeter) of distal lateral leaflets as in (A). Outlines of representative M82 and IL2-5/IL4-3 leaflets are shown above bars. Letters indicate statistical significance in each pairwise genotype comparison (p < 0.05). **(C)** Distribution of relative nuclear sizes reflecting endoreduplication in leaflets as in (A) and (B) (n = 5). Letters denote statistical significance between pairwise genotype comparisons at each ploidy level. **(D)** Leaf plastochron P3 dimensions calculated from 3D surface reconstructions of vegetative shoot apices (n = 18 for M82, n = 9 for each IL; n = 8 for double homozygotes; * p < 0.05, ** p < 0.01, *** p < 0.001 relative to M82).

Figure 5. Comparative transcriptomics of leaf development in two thick ILs and their parents. (A) Successive stages of leaf development (plastochrons P1-P4 colored as in legend in (B)) were dissected from M82, *S. pennellii* (Sp) and thick ILs 2-5 and 4-3. **(B)** Principal Components Analysis (PCA) of normalized RNA-Seq read counts. **(C)** Venn diagrams (not to scale) depict an overview of differentially expressed genes (DEGs, q < 0.05) that are shared in each IL and the Sp parent relative to M82. The number of DEGs unique to each organ is shown within elipses and those common to all organs, in the center. The total number of DEGs at each plastochron stage is shown outside ellipses.

Figure 6. Comparative expression profiles of genes in three functional categories across leaf development (P1 - P4) in thick ILs 2-5 and 4-3: (A) Transcription factors common to both ILs. **(B)** Genes involved in leaf development in tomato (as in Ichihashi et al., 2014), and **(C)** Gene annotated to encode components of the cell cycle or ubiquitin protesaome pathway (contain one of the terms "cell cycle", "cyclin", "ubiquitin", "E2F", "mitosis", "mitotic", "SKP"). Plastochron stages with statistically significant DE (q < 0.05) relative to M82 are marked with an asterisk. Genes, which are differentially expressed in at least one stage in both ILs are marked in bold.

Figure 7. Select leaf development gene regulatory sub-networks for (A-C) IL2-5 and (D-F) IL4-3. Sub-networks for regulators central to more than one plastochron stage are shown in **(A)** and **(D)**. A GA 20-oxidase gene (GA 20-ox 03g, Solyc03g006880) and its regulators in each IL **(B)** and **(E)**. Sub-networks of dynamic gene regulatory networks, showing interactions of an AUX/IAA TF (AUX/IAA 12g, Solyc12g096980) with other source nodes **(C)** and **(F)**. Gene IDs of highlighted nodes: SBP-box 04g, Solyc04g064470; BEL1-like 04g, Solyc04g080780; MADS-box 12g, Solyc12g087820; MYB TF 05g, Solyc05g007710; bHLH TF 04g, Solyc04g074810;GRAS 08g, Solyc08g014030; Myb 07g, Solyc07g052490; WRKY 05g, Solyc05g015850; AUX/IAA 06g, Solyc06g008580; Myb 03g, Solyc03g005570; Myb 08g, Soly08g005870; WRKY 02g, Solyc02g080890. Nodes and edges are colored according to legend.