TITLE:

Evaluating the clinical validity of gene-disease associations: an evidence-based

framework developed by the Clinical Genome Resource

AUTHORS:

Natasha T. Strande*[1], Erin Rooney Riggs*[2], Adam H. Buchanan[3], Ozge Ceyhan-Birsoy[6-9], Marina DiStefano[6], Selina S. Dwight[4], Jenny Goldstein[1], Rajarshi Ghosh[5], Bryce A. Seifert[1], Tam P. Sneddon[4], Matt W. Wright[4], Laura V. Milko[1], J. Michael Cherry[4], Monica A. Giovanni[3], Michael F. Murray[3], Julianne M. O'Daniel[1], Erin M. Ramos[10], Avni B. Santani[11-12], Alan F. Scott[13], Sharon E. Plon[5], Heidi L. Rehm[6-9], Christa L. Martin#[2,3], Jonathan S. Berg#[1]

AFFILIATIONS:

[1] Department of Genetics, School of Medicine, University of North Carolina at Chapel Hill, Chapel Hill, North Carolina, USA.

[2] Autism & Developmental Medicine Institute, Geisinger Health System, Danville, Pennsylvania, USA.

[3] Genomic Medicine Institute, Geisinger Health System, Danville, Pennsylvania, USA.

[4] Department of Genetics, Stanford University, Stanford, California 94305, USA.

[5] Department of Pediatrics, Baylor College of Medicine, Houston, Texas, USA.

[6] Laboratory for Molecular Medicine, Partners Personalized Medicine, Boston, Massachusetts, USA.

[7] The Broad Institute of Harvard and MIT, Cambridge, Massachusetts, USA.

[8] Harvard Medical School, Boston, Massachusetts, USA.

[9] Department of Pathology, Brigham & Women's Hospital, Boston, Massachusetts.

[10]National Human Genome Research Institute, National Institutes of Health, Bethesda,

Maryland, USA.

[11]Department of Pathology and Laboratory Medicine, Perelman School of Medicine,

University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA.

[12]Division of Genomic Diagnostics, Children's Hospital of Philadelphia, Philadelphia,

Pennsylvania 19104, USA.

[13]McKusick-Nathans Institute of Genetic Medicine, Johns Hopkins University School of

Medicine, Baltimore, MD 21287, USA.

JOINT AUTHORSHIP: *, # Authors contributed equally

CORRESPONDING AUTHOR: Jonathan S. Berg (jonathan_berg@med.unc.edu)

ABSTRACT

With advances in genomic sequencing technology, the number of reported gene-

disease relationships has rapidly expanded. However, the evidence supporting these

claims varies widely, confounding accurate evaluation of genomic variation in a clinical

setting. Despite the critical need to differentiate clinically valid relationships from less

well-substantiated relationships, currently no standard guidelines for such evaluation

exist. Thus the NIH-funded Clinical Genome Resource (ClinGen) has developed a

framework to define and evaluate the clinical validity of gene-disease pairs across a

variety of Mendelian disorders. Relevant genetic and experimental evidence supporting

or contradicting a gene-disease relationship is evaluated semi-quantitatively and

assigned a preliminary classification: "Definitive", "Strong", "Moderate", "Limited", "No

Reported Evidence" or "Conflicting Evidence." Classifications are reviewed and

confirmed or adjusted based on clinical expertise of appropriate disease experts. This

evidence-based, systematic method to assess the strength of gene-disease

relationships will facilitate more knowledgeable utilization of genomic variants in clinical

and research settings.

INTRODUCTION

INTRODUCTION

The human genome comprises approximately 20,000 protein-coding genes[1], of which

about 3,000 have been reported in association with at least one Mendelian disease[2].

Roughly half[2] of these gene-disease relationships have been identified over the last

decade, as technological advances have made it possible to use sequence information

from small families or even single individuals to discover new candidate gene-disease

relationships[3; 4]. However, there is substantial variability in the level of evidence

supporting these claims, and a systematic method for curating and assessing evidence

is needed.

Despite this variability, clinical laboratories may include genes with preliminary evidence

of a gene-disease relationship on disease-targeted panels, or in results returned from

exome/genome sequencing. Some of the gene-disease relationships are either unable

to be confirmed for many years or are ultimately proven wrong[5]. Evaluating the clinical

impact of variants identified in genes with an unclear role in disease is exceedingly

difficult, and could lead to an incorrect diagnosis for the patient, preventing further

evaluations and/or resulting in errant management of the patient and family. This

scenario highlights the need for a standardized method to evaluate the evidence

implicating a gene in disease and thereby determine the clinical validity[3] of a gene-

disease relationship.

The NIH-funded Clinical Genome Resource (ClinGen)[6] is creating an open-access

resource to better define clinically relevant genes and variants based on standardized,

transparent evidence assessment for use in precision medicine and research. Our

group has developed a method that 1) qualitatively defines gene-disease clinical validity

using a classification scheme based on the strength of evidence supporting the

relationship, and 2) provides a standardized semi-quantitative approach to evaluate

available evidence and arrive at such a classification. Currently, this framework is

optimized for genes associated with monogenic disorders following autosomal

dominant, autosomal recessive, or X-linked inheritance. Future iterations will expand the

framework to consider other modes of inheritance, such as mitochondrial, and diseases

with more complex genomic etiologies, including oligogenic or multifactorial conditions.

Our approach is neither intended to define multifactorial disease risk, nor to be a

substitute for well-established statistical thresholds used for genome-wide association

studies[7; 8].

This novel framework classifies gene-disease relationships by the quantity and quality

of the evidence supporting such a relationship. It builds on efforts to catalog gene-

disease associations, such as the Online Mendelian Inheritance in Man (OMIM)[1] and

OrphaNet, by systematically organizing the supporting and refuting evidence, and

categorizing the strength of evidence supporting these relationships. The resulting

clinical validity classifications are valuable to both clinicians and clinical laboratories.

First, they provide insight into the strength of clinical associations for clinicians

interpreting genetic test results for patient care. Second, they serve to guide clinical

genetic testing laboratories as they develop disease-specific clinical genetic testing

panels or interpret genome-scale sequencing tests. By including only those genes with

established clinical validity, the possibility of returning ambiguous, incorrect, or

uninformative results is reduced, improving the quality of interpretation of genomic data.

QUALITATIVE DESCRIPTION: CLINICAL VALIDITY CLASSIFICATIONS

The ClinGen Gene Curation Working Group (GCWG) is comprised of medical geneticists, clinical laboratory diagnosticians, genetic counselors, and biocurators with broad experience in both clinical and laboratory genetics. Over the course of three years, this group convened bi-monthly to develop the described framework for assessing gene-disease clinical validity through expert opinion and working group consensus (additional details in Supplemental Methods). We first defined six classes to qualitatively describe the strength of evidence supporting a gene-disease association (Figure 1). The amount and type of evidence required for each clinical validity classification builds upon that of the previous classification level. Evidence used within this framework to assign a classification to a gene-disease pair is divided into two main types: genetic evidence and experimental evidence (described below). As evidence is likely to change over time, any given classification is only representative of the level of evidence at the time of curation.

The classification "No Reported Evidence" is used for genes that have not yet been asserted to have a causal relationship with a human monogenic disorder, but may have some experimental data (e.g., model system data) suggesting a potential role for that gene in disease. The "Limited" classification requires at least one variant, asserted to be disease-causing, to have plausible genetic evidence to support the association with human disease with or without gene-level experimental data. "Moderate" classification encompasses additional clinical evidence (eg. multiple unrelated probands harboring variants with potential roles in disease) and supporting experimental evidence, all of which may be provided by multiple studies or a single robust study. Replication of the

gene-disease association in subsequent independent publications and additional

substantial genetic and experimental data are critical factors for the "Strong"

classification. Finally, the hallmark of a "Definitive" gene-disease association is that, in

addition to the accumulation of convincing genetic and experimental evidence, the

relationship has been replicated, and ample time has passed since the initial publication

(in general, greater than three years) for any conflicting evidence to emerge. It is

important to highlight that these classifications do not reflect the effect size or relative

risk attributable to variants in a particular gene, but instead the strength of the evidence.

For example, a definitive gene-disease association does not imply that a pathogenic

variant in that gene confers 100% penetrance of the phenotype. This metric is not

intended to assess the penetrance or risk to develop a disease outcome.

A gene-disease relationship can be determined to have one of the above classifications

provided no substantial relevant and valid contradictory evidence exists to call the gene-

disease relationship into question. If such evidence emerges, then the relationship is

described as "Conflicting Evidence Reported." Types of contradictory evidence may

come from population studies (such as ExAC[9]), attempts to experimentally validate the

gene-disease association, or re-analysis of the original family or cohort that was

previously studied. Although the role of a specific *variant* in a given disease may be

called into question by new evidence, this may not be sufficient to invalidate the role of

the *gene* in that disease. Thorough evaluation by experts in the particular disease area

is recommended to determine whether the contradictory evidence outweighs the

existing supportive evidence to classify a gene into either a "Disputed" or "Refuted"

category (see Figure 1 for additional details).

METHODS: SEMI-QUANTITATIVE ASSESSMENT OF EVIDENCE

Assigning a clinical validity classification to a gene-disease pair requires assessment of the evidence supporting the association. We developed a semi-quantitative approach to evaluate both genetic (Figure 2) and experimental evidence (Figure 3) in a standardized manner that promotes consistent collection and weighting of evidence. Defined sub-categories of genetic and experimental evidence are given a suggested default "score." However, given that evidence of the same general type may vary in its strength (particularly when considering different diseases), the scoring system also allows these scores to be adjusted within a set range of points, with final approval by experts within the particular disease domain. Finally, the maximum number of points allowed for the various types of genetic and experimental evidence is capped to prevent a preponderance of weak evidence from inappropriately inflating the gene-disease classification. Similarly, certain evidence categories are provided higher maximum scores, allowing key pieces of stronger evidence to proportionately influence the classification of a gene-disease pair.

Genetic Evidence

For the purposes of scoring, genetic evidence is divided into two categories: case-level data and case-control data (Figure 2). Studies describing individuals or families with genetic variants are scored as case-level data, while studies using statistical analyses to compare variants in cases and controls are scored as case-control data. When case-level and case-control data are present in a single publication, points can be assigned in each category, but the same piece of evidence should not be counted more than once.

For example, an individual case that is also included within a case-control cohort should not be given points in both the "case-level data" and "case-control data" categories. In this scenario, points should be assigned to the most compelling and informative evidence.

Assessing case-level data requires consideration of the inheritance pattern and evaluation of the individual variants identified in each case. Within this framework, a case should only be counted towards supporting evidence if the reported variant has some indication of a potential role in disease (e.g., impact on gene function, recurrence in affected individuals, etc.) and does not have evidence that would contradict pathogenicity (e.g., population allele frequency). Unless otherwise noted, the term "qualifying variant" implies that these criteria are met. In addition, points are assigned separately for segregation data to reflect the statistical probability that the locus is implicated in the disease. Figure 2 and Figure S1 provide guidance on the number of points that should be considered for segregation evidence by LOD score; if a LOD score is not provided within the publication being evaluated, an estimated LOD score may be calculated in certain scenarios, as described in the Supplemental Methods.

Each study categorized as "case-control data" should be independently assessed to evaluate the quality of the study design (see Figure 2 and Supplemental Methods). Consultation with a clinical domain expert group (such as those affiliated with ClinGen, https://www.clinicalgenome.org/working-groups/clinical-domain/) is recommended. For the purposes of this framework, studies are classified based on whether they include single variant analysis or aggregate variant analysis. Single variant analyses are those in which individual variants are evaluated for statistical enrichment in cases compared

to controls. More than one variant may be analyzed, but the variants have been

independently assessed with appropriate statistical correction for multiple testing.

Aggregate variant analyses are those in which the total number of variants is assessed

for enrichment in cases compared with controls. This comparison is typically

accomplished by sequencing the entire gene in both cases and controls and

demonstrating an increased "burden" of variants of one or more types.

Experimental Evidence

The experimental data scoring system is presented in Figure 3. The gene-level

experimental data used in this framework to assess a gene-disease association are

consistent with those proposed by MacArthur and colleagues to implicate a gene in

disease[10]. The following experimental evidence types are used: biochemical function,

experimental protein interactions, expression, functional alteration, phenotypic rescue

and model systems (Figure 3 bottom panel). These categories capture the most

relevant types of experimental information necessary to determine whether the function

of the gene product is at least consistent with the disease with which it is associated, if

not causally implicated.

Contradictory Evidence

While curators are encouraged to seek out and document (via qualitative description)

conflicting evidence, no specific points are assigned to this category. The types of valid

contradictory evidence and their relative weights will be unique to each gene-disease

pair, and it would be misleading to attempt to uniformly quantify this type of negative

evidence against the reported positive evidence. If there is substantial conflicting

evidence, manual review and expert input is required to evaluate the strength of the

contradictory evidence, determine whether it outweighs any available supporting

evidence, and, if so, decide whether the gene-disease association should be classified

as "Disputed" or "Refuted".

Summary & Final Matrix

The scores assigned to both genetic and experimental evidence are tallied to generate

a total score (ranging from 1-18) that corresponds to a preliminary clinical validity

classification (Figure 4). The system provides a transparent method for summarizing

and assessing all curated evidence for a gene-disease pair, encouraging consistency

between curators. While the summary matrix facilitates a preliminary assessment of the

gene-disease relationship, the initial curator or expert reviewer may adjust the

classification, supplying a specific rationale for the change. Final classifications are

determined in collaboration with disease experts, who review the preliminary

classification and supporting evidence and work to come to a consensus with the

preliminary curators. In the event that the disease experts and preliminary curators

disagree on a final classification, a senior member of the ClinGen Gene Curation

Working Group may be brought in to facilitate a final classification, erring towards the

more conservative classification if consensus cannot be achieved. It should be noted

that experimental data alone cannot justify a clinical validity classification beyond "No

Reported Evidence," and at least one human genetic variant with a plausible causal

association must be present to attain "Limited" classification. The difference between

"Limited," "Moderate," and "Strong" gene-disease classifications is justified by the

quality and quantity of evidence; it is expected that valid gene-disease associations will

1 gradually accumulate enough supporting evidence and be replicated over time to attain

2 a "definitive" classification. This framework relies on evidence obtained primarily from

3 published literature; however, if necessary, unpublished information available from

4 publicly accessible resources, such as variant databases[11; 12], may be used as long as

5 some supporting evidence is provided.

7 *RESULTS: VALIDATION OF METHOD*

Using this framework we evaluated 33 gene-disease pairs representing a variety of

disease domains and spanning the spectrum of clinical validity classifications (see

Figure 5 and Supplemental Methods). To assess the reproducibility of our scoring

metric, each gene-disease pair was evaluated by two independent curators; paired

curators reached concordant clinical validity classifications in 29 of the 31 (93.5%)

gene-disease pairs with available published evidence (Figure 5; associations classified

as "No Reported Evidence" were excluded). Each gene-disease pair was subsequently

reviewed by clinical domain experts; experts agreed with the preliminary classifications

for 87.1% (27/31) of the gene-disease pairs with published evidence (Figure 5). The

four discrepancies between the expert and curator classifications were each different by

only a single category (e.g. limited versus moderate). Of note, the original classifications

for *HNRNPK* and *SMARCA1* were at the border between limited and moderate (6.5

points); in each case, the preliminary curators' lack of specific clinical expertise led to

uncertainty regarding the scoring of evidence requiring such knowledge. Consulting with

clinical experts in the disease resolved these issues resulting in both genes being

upgraded to moderate. In the case of *WRAP53,* the expert was aware of additional

published experimental evidence that when included increased the classification from

limited to moderate. Upon reviewing the curated evidence for *RAD51D* and breast

cancer, the domain expert upgraded the classification from disputed to limited (with the

approval of the GCWG) due to the specificity of the experimental evidence and

insufficient power of the current studies to rule out a role for *RAD51D* in breast cancer

(Figure 5). Details and references for each curation are provided in Supplemental

Appendix.

8
DISCUSSION

The evidence-based framework described here qualitatively defines clinical validity

classifications for gene-disease associations in monogenic conditions and provides a

systematic framework for evaluating key criteria required for these classifications. This

method is intentionally flexible to accommodate curation of a wide spectrum of genes

and conditions by curators with varying levels of expertise. The semi-quantitative

scoring system combined with the qualitative classification scheme guides curators

through the preliminary decision-making process, while the expert-level review provides

disease-specific experience to weigh in on the final classification.

This effort to create a generalized framework may result in some specific challenges

due to the heterogeneity of genetic conditions, in both phenotype and prevalence. For

example, ultra-rare disorders may have a relatively small number of probands described

in the medical literature, thus limiting their potential to achieve a high genetic evidence

score within this matrix. This obstacle is mostly circumvented by allowing compelling

pieces of genetic evidence to score the maximum number of points (for example, see

CD3E and severe combined immunodeficiency, detailed in the Supplemental Appendix). When substantial experimental evidence is also available, these conditions can attain a "Strong" or "Definitive" classification.

On the opposite end of the spectrum are conditions that occur commonly in the general population, such as cancer, where the predominant etiology is multifactorial rather than monogenic. In the less common Mendelian cancer predisposition syndromes, incomplete penetrance is a typical feature that can lead to confounding factors in family genetic studies such as apparently non-penetrant family members who carry a disease-associated variant and phenocopies among family members without a disease-associated variant. For such conditions, case-control data may provide more compelling evidence to support the gene-disease association (see the curation of *PALB2* and hereditary breast cancer in the Supplemental Appendix as an example).

One limitation of any such system is the challenge of balancing thorough literature curation and practical time commitment. This system can accommodate an exhaustive literature review, but in most cases will only require curating the amount of information sufficient to reach the maximum number of points in the matrix. In some scenarios this method may fail to include pertinent information, which could impact the classification (e.g., omission of contradictory evidence). Another potential limitation is the subjective nature of certain evidence types (e.g., experimental), which may lead to variability between different groups assessing evidence. However, due to the transparency of the evidence base, the incorporation of expert review, and the ability to reassess classifications over time, such drawbacks are likely to be self-limiting.

ClinGen's ultimate goal is to enhance the incorporation of genomic information into patient care, an important component of the Precision Medicine Initiative[13]. The implementation of this framework will be supported by an open-access ClinGen curation interface (under development), which will provide a platform for extension to the community. In essence, this framework aims to provide a systematic, transparent method to evaluate a gene-disease relationship in an efficient and consistent manner suitable for a diverse set of users. A detailed standard operating procedure for this framework is available on the ClinGen website. All curated evidence, including clinical validity assessments, will be made readily accessible to clinical laboratories, clinicians, patients, researchers, and others via our website.

Carefully evaluated gene-disease clinical validity classifications, as provided by this framework, will be useful to clinical laboratories as they evaluate genes for inclusion on disease-targeted panels, or as they decide how to categorize, prioritize, and return results from exome/genome sequencing. Clinicians may choose to use these types of gene-disease classifications as they interpret laboratory results for their patients; for instance, they may choose not to adjust medical management based on variants in genes of limited clinical validity. Researchers could also utilize this framework to evaluate the clinical validity of their own newly discovered associations and identify promising target genes for future work in order to augment the currently available evidence and attain a "Strong" or "Definitive" classification. In addition, professional societies and regulatory bodies may utilize these clinical validity assessments when making recommendations or guidelines for clinical genetic testing. Ultimately, our

1   systematic, evidence-based method for evaluating gene-disease associations will

2   provide a strong foundation for genomic medicine.

DESCRIPTION OF SUPPLEMENTAL DATA:

The Supplemental file includes methods, footnotes for Figure 2, one figure, an appendix

with curated evidence for each example presented in Figure 5 and a list of references.

ACKNOWLEDGEMENTS:

This work was supported by grants from the National Human Genome Research

Institute (NHGRI), through the following three grants: U41 HG006834-01A1, U01

HG007437-01, U01 HG007436-01, as well as from the National Cancer Institute (NCI)

through the following contract: HHSN261200800001E. ClinGen also receives funding

through the Eunice Kennedy Shriver National Institute of Child Health and Human

Development (NICHD) and ClinVar is supported by the Intramural Research Program of

the NIH, National Library of Medicine. We would like to thank the following groups and

individuals for contributing their disease expertise to review the examples included in

this manuscript: Alan Beggs Ph.D.; Alison Bertuch, M.D., Ph.D.; Rebecca H. Buckley,

M.D.; Eugene Chung, M.D.; Bill Craigen M.D., Ph.D.; Jennifer M. Puck, M.D.; Sharon A.

Savage, M.D.; Fergus J. Couch Ph.D. and the ClinGen Hereditary Breast and Ovarian

Cancer gene curation working group; Birgit H. Funke Ph.D. and the ClinGen

Cardiomyopathy gene curation working group; and the ClinGen RASopathy curation

working group. Input on the framework was also provided by the ClinGen Hereditary

Breast and Ovarian Cancer gene curation working group and Ray Hershberger, Mike

Gollob and the ClinGen Channelopathy gene curation working group. We would also

like to thank Scott Goehringer for his invaluable help in preparing the curated examples

for the ClinGen website and appendix.

WEB RESOURCES:

Clinical Genome Resource: www.clinicalgenome.org

Gene Curation Working Group members: http://clinicalgenome.org/about/working-

groups/gene-curation/

Link to the standard operating procedure for the clinical validity framework described in

this manuscript: http://bit.ly/2clingenGCSOP

REFERENCES:

1. Online Mendelian Inheritance in Man, OMIM®. In. (Baltimore, MD, McKusick-Nathans Institute of Genetic Medicine, Johns Hopkins University

2. Chong, J.X., Buckingham, K.J., Jhangiani, S.N., Boehm, C., Sobreira, N., Smith, J.D., Harrell, T.M., McMillin, M.J., Wiszniewski, W., Gambin, T., et al. (2015). The Genetic Basis of Mendelian Phenotypes: Discoveries, Challenges, and Opportunities. American journal of human genetics 97, 199-215.

3. Haddow, J., Palomacki, G. (2003). ACCE: A Model Process for Evaluating Data on Emerging Genetic Tests. In Human Genome Epidemiology: A Scientific Foundation for Using Genetic Information to Improve Health and Prevent Disease, M. Khoury, Little, J., Burke, W., ed. (Oxford University Press), pp 217-233.

4. Wilfert, A.B., Chao, K.R., Kaushal, M., Jain, S., Zollner, S., Adams, D.R., and Conrad, D.F. (2016). Genome-wide significance testing of variation from single case exomes. Nat Genet 48, 1455-1461.

5. Eisenberger, T., Di Donato, N., Baig, S.M., Neuhaus, C., Beyer, A., Decker, E., Murbe, D., Decker, C., Bergmann, C., and Bolz, H.J. (2014). Targeted and genomewide NGS data disqualify mutations in MYO1A, the "DFNA48 gene", as a cause of deafness. Human mutation 35, 565-570.

6. Rehm, H.L., Berg, J.S., Brooks, L.D., Bustamante, C.D., Evans, J.P., Landrum, M.J., Ledbetter, D.H., Maglott, D.R., Martin, C.L., Nussbaum, R.L., et al. (2015). ClinGen--the Clinical Genome Resource. The New England journal of medicine 372, 2235-2242.

7. Lander, E., and Kruglyak, L. (1995). Genetic dissection of complex traits: guidelines for interpreting and reporting linkage results. Nat Genet 11, 241-247.

8. Sham, P.C., and Purcell, S.M. (2014). Statistical power and significance testing in large-scale genetic studies. Nature reviews Genetics 15, 335-346.

9. Lek, M., Karczewski, K.J., Minikel, E.V., Samocha, K.E., Banks, E., Fennell, T., O'Donnell-Luria, A.H., Ware, J.S., Hill, A.J., Cummings, B.B., et al. (2016). Analysis of protein-coding genetic variation in 60,706 humans. Nature 536, 285-291.

10. MacArthur, D.G., Manolio, T.A., Dimmock, D.P., Rehm, H.L., Shendure, J., Abecasis, G.R., Adams, D.R., Altman, R.B., Antonarakis, S.E., Ashley, E.A., et al. (2014). Guidelines for investigating causality of sequence variants in human disease. Nature 508, 469-476.

11. Landrum, M.J., Lee, J.M., Riley, G.R., Jang, W., Rubinstein, W.S., Church, D.M., and Maglott, D.R. (2014). ClinVar: public archive of relationships among sequence variation and human phenotype. Nucleic acids research 42, D980-985.

12. Fokkema, I.F., Taschner, P.E., Schaafsma, G.C., Celli, J., Laros, J.F., and den Dunnen, J.T. (2011). LOVD v.2.0: the next generation in gene variant databases. Human mutation 32, 557-563.

13. Collins, F.S., and Varmus, H. (2015). A new initiative on precision medicine. The New England journal of medicine 372, 793-795.

 FIGURE TITLES AND LEGENDS:

 Figure 1: ClinGen clinical validity classifications and qualitative descriptions. The

 suggested minimum criteria needed to obtain a given classification are described for

 each clinical validity classification. The types of evidence comprising these criteria are

 described in the text. The default classification for genes without a convincing human

 disease-causing variant is "No Reported Evidence." The level of evidence needed for

 each supportive gene-disease association category builds upon the previous category

 (i.e. "Limited" builds upon "Moderate"). Gene-disease associations classified as

 "Contradictory" likely have supporting evidence as well as opposing evidence, but are

 described separately from the classifications for supportive gene-disease associations.

 Figure 2: Classes of genetic evidence and their relative weights used in the

 ClinGen clinical validity framework. Genetic evidence is separated into two main

 categories: case-level data and case-control data. While a single publication may

 include both case-level and case-control data, individual cases should NOT be included

 in both categories. Each category is assigned a range of points with a maximum score

 that can be achieved. *Case-Level Data* is derived from studies describing individuals

 and/or families with qualifying variants in the gene of interest. Points should be assigned

 to each case based on the variant's inheritance pattern, molecular consequence and

 evidence of pathogenicity in disease. In addition to variant evidence points, a gene-

 disease pair may also receive points in the segregation evidence category for

 compelling segregation analysis (see Figure S1). *Case-Control Data:* Studies utilizing

 statistical analysis to evaluate variants in cases compared to controls. Case-control

 studies can be classified as either single variant analysis or aggregate variant analysis,

however the number of points allowable for either category is the same. Points should be assigned to case-control studies according to the overall quality of each study based on these criteria: variant detection methodology, power, bias and confounding factors, and statistical power. Additional details included in Supplemental Methods. Note that the maximum total scores allowed for different types of Case-Level data are not intended to add up to the total points allowed for Genetic Evidence as a whole. This permits different combinations of evidence types to achieve the maximum total score.

Figure 3: Types of gene-level experimental evidence and their relative weights used in the ClinGen clinical validity framework. Experimental evidence types used in the ClinGen gene curation framework are modified from MacArthur, et al. 2014. Evidence types are divided into three categories based on their relative contribution to the overall clinical validity of a gene-disease pair giving more weight to *in vivo* data. Each category is assigned a range of points with a maximum score that can be achieved, allowing more weight to be given to *in vivo* data (e.g. Models & Rescue) over *in vitro* experimental data. Evidence within the function category is given the least weight and is comprised of the following types of evidence: biochemical function, interactions, and expression. Functional alteration experiments in cells from patients carrying candidate pathogenic variants are given more weight than the function category. Finally model systems and phenotypic rescue experiments are given the most weight in our framework. Note that the maximum total scores allowed for different categories of Experimental Evidence are not intended to add up to the total allowable points. This permits different combinations of evidence types to achieve the maximum total score.

Figure 4. Final summary matrix used to provisionally classify gene-disease

associations. A summary matrix was designed to generate a "provisional" clinical

validity assessment using a point system consistent with the qualitative descriptions of

each classification. *Genetic Evidence:* total number of points (not exceeding 12)

obtained using the scoring metric in Fig. 2. If no human mutations have been found

within the literature, then the default classification is "No Reported Evidence."

Experimental Evidence: total number of points (not exceeding 6) derived from each of

the experimental categories in Fig. 3. *Replication Over Time* – Yes, if more than three

years has passed since the publication of the first paper reporting the gene-disease

relationship AND more than two publications with human mutations exist. *Contradictory*

Evidence – No points are assigned to this category. Instead, the curator should provide

a summary of contradictory information. *Scoring* - The sum of the quantified evidence

from each category can be used to determine a "provisional" classification using the

scale at the bottom of the figure. If a curator does not agree with this classification,

he/she may provide a different suggested classification along with appropriate

justification.

Figure 5. Comparison of provisional clinical validity classifications and

associated matrix scores for selected gene-disease pairs evaluated by multiple

curators. Of the 33 gene-disease pairs (y-axis) curated to validate the clinical validity

curation framework, 31 were classified using the summary matrix (2 gene-disease pairs,

PMS2:pancreatic cancer and *ARSD*:chondrodysplasia punctata, were classified as "No

evidence reported" and are not shown). Genetic evidence (grey bars) and experimental

evidence (black bars) were evaluated by two independent curators (C1-C9) to arrive at

a provisional classification (x-axis). Gene-disease relationships scoring between 12-18

points can be "Strong" or "Definitive," depending on whether the association has been

replicated over time (indicated by the squared "r/t"), in which case the preliminary

classification is "Definitive". Clinical validity classifications that were discordant between

preliminary curators are represented with a dashed background. Gene-disease pairs in

which conflicting evidence was reported are represented by diagonal lines through the

evidence bars and a grey background. The letter "C" in a triangle indicates that the

curators classified the gene-disease pair as "Conflicting Evidence Reported". Each

gene-disease pair was ultimately evaluated by an expert in the field for a final

classification (far right column). Final expert classifications that differed from the

preliminary classification are indicated by italics and asterisks.

Figure 1

Evidence Level		Evidence Description
Supportive Evidence	**DEFINITIVE**	The role of this gene in this particular disease has been repeatedly demonstrated in both the research and clinical diagnostic settings, and has been upheld over time (in general, at least 3 years). No convincing evidence has emerged that contradicts the role of the gene in the specified disease.
	STRONG	The role of this gene in disease has been independently demonstrated typically in at least two separate studies providing **strong** supporting evidence for this gene's role in disease, usually including both of the following types of evidence: • Strong variant-level evidence demonstrating numerous unrelated probands with variants that provide convincing evidence for disease causality[1] **as well as** • Compelling gene-level evidence from different types of supporting experimental data[2]. In addition, no convincing evidence has emerged that contradicts the role of the gene in the noted disease.
	MODERATE	There is **moderate** evidence to support a causal role for this gene in this disease, typically including both of the following types of evidence: • Several probands with variants that provide convincing evidence for disease causality[1] • Moderate experimental data[2] supporting the gene-disease association The role of this gene in disease may not have been independently reported, but no convincing evidence has emerged that contradicts the role of the gene in the noted disease.
	LIMITED	There is **limited** evidence to support a causal role for this gene in this disease, such as: • Fewer than three observations of variants that provide convincing evidence for disease causality[1] OR • Variants have been observed in probands, but none have sufficient evidence for disease causality. • Limited experimental data[2] supporting the gene-disease association The role of this gene in disease may not have been independently reported, but no convincing evidence has emerged that contradicts the role of the gene in the noted disease.
NO REPORTED EVIDENCE		Evidence for a causal role in disease has not been reported. These genes might be "candidate" genes based on linkage intervals, animal models, implication in pathways known to be involved in human diseases, etc., but no reports have directly implicated the gene in human disease cases.

Contradictory Evidence	**CONFLICTING EVIDENCE REPORTED**	Although there has been an assertion of a gene-disease association, conflicting evidence for the role of this gene in disease has arisen since the time of the initial report indicating a disease association. Depending on the quantity and quality of evidence disputing the association, the association may be further defined by the following two sub-categories:
		1. **Disputed**
		  a. Convincing evidence *disputing* a role for this gene in this disease has arisen since the initial report identifying an association between the gene and disease.
		  b. Refuting evidence need not outweigh existing evidence supporting the gene:disease association.
		2. **Refuted**
		  a. Evidence refuting the role of the gene in the specified disease has been reported and significantly outweighs any evidence supporting the role.
		  b. This designation is to be applied at the discretion of clinical domain experts after thorough review of available evidence

NOTES		
[1]Variants that disrupt function and/or have other strong genetic and population data (e.g. *de novo* occurrence, absence in controls, strong linkage to a small genomic interval, etc.) are considered convincing of disease causality in this framework. [2]Examples of appropriate types of supporting experimental data based on those outlined in MacArthur et al. 2014.		

Figure 2

	Evidence Type		Case Information		Suggested Points/Case		Points Given	Max Score
					Default	Range		
Case-Level Data[1]	Variant Evidence	Autosomal Dominant OR X-Linked Disorder[2]	Variant is de novo[3]		2	0-3		**12**
			Proband with predicted or proven null variant[4]		1.5	0-2		**10**
			Proband with other variant type with some evidence of gene impact[5]		0.5	0-1.5		**7**
		Autosomal Recessive	Two variants in trans and at least one de novo[3] or a predicted/proven null variant[4]		2	0-3		**12**
			Two variants (not predicted/proven null) with some evidence of gene impact[5] in trans		1	0-1.5		
	Segregation Evidence		Evidence of segregation in one or more families[6]	LOD Score Examples: 3 / 2 / 1.5 / 1 → 5 / 4 / 3 / 1.5		0-7		**7**

	Case-Control Study Type[8]	Case-Control Quality Criteria[9]	Suggested Points/Study	Points Given	Max Score
Case-Control Data[7]	Single Variant Analysis[8a]	• Variant Detection Methodology[9a] • Power[9b]	0-6		12
	Aggregate Variant Analysis[8b]	• Bias and Confounding Factors[9c] • Statistical Significance[9d]	0-6		

TOTAL ALLOWABLE POINTS for Genetic Evidence	12

General Notes
- All variants under consideration should be rare enough in the general population to be consistent with disease.
- Cohorts/cases should not be double counted. For example, individual cases included as part of case-control studies should not be given points from both the "Case Level Data" and "Case-Control Data" categories.
- Case-Level Data includes studies describing individuals or families with variation in the gene of interest
- Case-Control studies are those in which statistical analysis is used to evaluate variation in cases compared to controls.
- Numbered footnotes are included in the Supplemental Material.

Figure 3

Evidence Category	Evidence Type	Suggested Points		Points Given	Max
		Default	Range		
Function	Biochemical Function		0-2		
	Protein Interaction	0.5	0-2		**2**
	Expression		0-2		
Functional Alteration	Patient cells	1	0-2		**2**
	Non-patient cells	0.5	0-1		
Models & Rescue	Animal model	2	0-4		
	Cell culture model system	1	0-2		**4**
	Rescue in animal model	2	0-4		
	Rescue in engineered equivalent	1	0-2		
Total Allowable Points for Experimental Evidence					**6**

Figure 4

Clinical Validity Summary Matrix

GENE/DISEASE PAIR:				
Assertion criteria	**Genetic Evidence (0-12 points)**	**Experimental Evidence (0-6 points)**	**Total Points (0-18)**	**Replication Over Time (Y/N)**
Description	Case-level, family segregation, or case-control data that support the gene-disease association	Gene-level experimental evidence that support the gene-disease association	Sum of Genetic & Experimental Evidence	> 2 pubs w/ convincing evidence over time (>3 yrs)
Assigned Points				
CALCULATED CLASSIFICATION		**LIMITED**	**1-6**	
		MODERATE	**7-11**	
		STRONG	**12-18**	
		DEFINITIVE	**12-18 & Replicated Over Time**	
Valid contradictory evidence (Y/N)*	**List PMIDs and describe evidence:**			
CURATOR CLASSIFICATION				
FINAL CLASSIFICATION				

Figure 5