Adaptation of the *FADS* gene family in Europe: Variation across time, geography and
subsistence

Kaixiong Ye[1], Feng Gao[1], David Wang[1], Ofer Bar-Yosef[2], Alon Keinan[1]*

[1] Department of Biological Statistics and Computational Biology, Cornell University, Ithaca,
NY, USA

[2] Department of Anthropology, Harvard University, Cambridge, MA, USA

*corresponding author: alon.keinan@cornell.edu

Abstract: The *FADS* gene family encodes rate-limiting enzymes for the biosynthesis of omega-6
and omega-3 long chain polyunsaturated fatty acids (LCPUFAs), which is essential for
individuals subsisting on LCPUFAs-poor diets (e.g. plant-based). Positive selection on *FADS*
genes has been reported in multiple populations, but its presence and pattern in Europeans
remain elusive. Here, with analyses of ancient and modern DNA, we demonstrated positive
selection acted on variants centered on *FADS1* and *FADS2* both before and after the advent of
farming in Europe, but adaptive alleles in these two periods are opposite. Selection signals in
recent history also vary geographically, with the strongest in Southern Europe. We showed that
adaptive alleles in recent farmers are associated with expression of *FADS* genes, enhanced
LCPUFAs biosynthesis and reduced risk of inflammatory bowel diseases. Thus, the adaptation of
FADS genes in Europe varies across time and geography, probably due to varying diet and
subsistence.

Identifying genetic adaptations to local environment, including historical dietary practice, and elucidating their implications on human health and disease are of central interest in human evolutionary genomics[1]. The fatty acid desaturase (*FADS*) gene family consists of *FADS1*, *FADS2* and *FADS3*, which evolved by gene duplication[2]. *FADS1* and *FADS2* encode rate-limiting enzymes for the endogenous synthesis of omega-3 and omega-6 long-chain polyunsaturated fatty acids (LCPUFAs) from shorter-chain precursors from plants (Supplementary Fig. 1). LCPUFAs are indispensable for proper human brain development, cognitive function and immune response[3,4]. While omega-3 and omega-6 LCPUFAs can be consumed from animal-based diets, their endogenous synthesis is essential to compensate for their absence from plant-based diets. Adaptation (positive selection) acting on the *FADS* locus, a 100 kilobase (kb) region containing all three genes (Supplementary Fig. 2), has been identified in multiple populations[5-9]. Our recent study showed that a 22 bp insertion-deletion polymorphism (indel, rs66698963) within *FADS2*, which is associated with *FADS1* expression[10], has been adaptive in Africa, South Asia and parts of East Asia, possibly driven by local historical plant-based diets[8]. We further supported this hypothesis by the functional association of the adaptive insertion allele with more efficient endogenous synthesis[8]. In Greenlandic Inuit, who have traditionally subsisted on a LCPUFAs-rich marine diet, adaptation signals were also observed on the *FADS* locus, with adaptive alleles associated with less efficient endogenous synthesis[9].

In Europeans, positive selection on the *FADS* locus has only been reported recently in a study based on ancient DNA (aDNA)[11]. Evidence of positive selection from modern DNA (mDNA) is still lacking even though most of the above studies also performed similarly-powered tests in Europeans[5-8]. Moreover, although there are well-established differences in the Neolithization process and in dietary patterns across Europe[12-14], geographical differences of selection signals within Europe have not been investigated before. Furthermore, before the advent of farming, pre-Neolithic hunter-gatherers throughout Europe had been subsisting on animal-based diets with significant aquatic contribution[15-17], in contrast to the plant-heavy diets of recent European farmers[18-20]. We hypothesized that these drastic differences in subsistence strategy and dietary practice before and after the Neolithic revolution within Europe exert different selection pressures on the *FADS* locus. In this study, we combined analyses on ancient and modern DNA to investigate the geographical and temporal differences of selection signals on the *FADS* locus in Europe. We further interpreted the functional significance of adaptive alleles with analysis of expression quantitative trait loci (eQTLs) and genome-wide association studies (GWAS), as well as with anthropological findings within Europe.

Results

Evidence of recent positive selection in Europe from both ancient and modern DNA

To systematically evaluate the presence of positive selection on the *FADS* locus in Europe, we performed an array of selection tests using both ancient and modern samples. We first generated a uniform set of variants across the locus in a variety of aDNA data sets (Supplementary Table S1) via imputation (Methods). For all these variants, we conducted an aDNA-based test for recent positive selection (Methods)[11]. This test includes three groups of ancient European samples and four groups of modern samples. The three ancient groups represent the three major

ancestry sources of most present-day Europeans: Western and Scandinavian hunter-gatherers
(WSHG), early European farmers (EF), and Steppe-Ancestry pastoralists (SA)[11,21-23]. The four
groups of modern samples were drawn from the 1000 Genomes Project (1000GP), representing
Tuscans (TSI), Iberians (IBS), British (GBR) and additional northern Europeans (CEU). Each
modern population has been modelled as a linear mixture of the three ancestral sources with
relative proportions estimated with genome-wide single nucleotide polymorphisms (SNPs)[11].
The frequencies of a neutral SNP in the four modern populations are expected to be the linear
combinations of its frequencies in the three ancient sources (the null hypothesis H_0), while
significant deviation from this expectation (the alternative hypothesis H_1) serves as a signal for
the presence of positive selection during recent history of Europe (not more ancient than 8500
years ago)[11]. Our results confirmed the presence of significant selection signals on many SNPs in
the *FADS* locus (Fig. 1A), including the previously identified peak SNP rs174546 ($p = 1.04e$-
21)[11]. We observed the most significant signal in the locus for an imputed SNP, rs174594 ($p =$
$1.29e$-24), which was not included in the original study[11]. SNP rs174570, one of the top adaptive
SNPs reported in Greenlandic Inuit[9], also carries significant signal ($p = 7.64e$-18) while indel
rs66698963 has no evidence of positive selection ($p = 3.62e$-3, but see Supplementary Text).
Overall, the entire peak of selection signals coincides with a linkage disequilibrium (LD) block
(henceforth referred to as the *FADS1-FADS2* LD block) in Europeans, which extends over a long
genomic region of 85 kb, covering the entirety of *FADS1* and most of the much longer *FADS2*
(Supplementary Figs. 2 and 3). This suggests that the large number of SNPs showing genome-
wide significant signals is likely the result of one causal variant targeted by strong selection and
extensive hitchhiking of nearby SNPs.

We next performed several selection tests solely based on mDNA in European populations.
Considering the five European populations from 1000GP, including samples of Finns (FIN) and
the four samples described above, two haplotype-based selection tests, iHS[24] and nSL[25], revealed
positive selection on the derived allele of the peak SNP from the aDNA-based test, rs174594, as
well as many other SNPs in the *FADS1-FADS2* LD block (Fig. 1B, Supplementary Figs. 4 and
5). The normalized nSL values are significant in all five populations and the signal exhibits a
gradient of being stronger in southern Europeans and weaker in northern Europeans, as per the
following order (Fig. 1B): TSI ($p = 0.00044$), IBS ($p = 0.0020$), CEU ($p = 0.0039$), GBR ($p =$
0.0093), and FIN ($p = 0.017$). The iHS value is only significant in TSI ($p = 0.026$). Repeating the
analysis in two whole-genome sequencing cohorts of British ancestry from the UK10K project
revealed consistently significant positive selection on the derived allele (Supplementary Fig. 6).
The other three variants of potential interest (rs174546, rs174570, and rs66698963, colored in
Fig. 1A) exhibit no or borderline selection signals, with only rs174570 showing significant
normalized nSL values in the two southernmost populations (TSI: $p = 0.022$; IBS: $p = 0.050$, Fig.
1B). Interestingly, it is the ancestral allele of rs174570 that was under positive selection, while its
derived allele has been shown to be targeted by positive selection in Greenlandic Inuit[9].

We further applied two site frequency spectrum (SFS)-based selection tests that consider
frequencies of all variants in a tested region (Methods). One of these two tests, Fay and Wu's
H[26], consistently revealed a cluster of significant signals spanning a 14 kb genomic region
surrounding the peak SNP rs174594 in all five 1000GP European populations ($p < 0.05$, Fig. 1C,

Supplementary Fig. 7). Similar results were observed with UK10K cohorts (Supplementary Fig.
8). Local peaks of H values also surround rs174546 and rs66698963, though they do not reach
significance level (Fig. 1C, Supplementary Fig. 8). In all these tests, whether significant or not,
the signals are gradually stronger towards Southern populations.

Taken together, standard tests on mDNA (Fig. 1B and 1C) support the results based on aDNA
(Fig. 1A) of recent positive selection on the *FADS* locus and, specifically, on the *FADS1-FADS2*
LD block. Moreover, all mDNA- and aDNA-based tests unanimously point to the same genomic
region as the peak of selection signals (Fig. 1). The mDNA results additionally reveal a South-
North gradient correlating with more pronounced selection signals in the South. As these tests
across 1000GP populations are of comparable power (N=91-107 individuals), these results
highlight an interesting possibility of stronger positive selection in Southern compared to
Northern Europe.

Geographical differences of recent positive selection signals across Europe

To further rigorously evaluate geographical differences of recent positive selection signals on the
FADS locus across Europe, we revisited the aDNA-based selection test[11]. We started by
decomposing the original test for four representative SNPs (Fig. 2A) and then performed a
revised version of the test separately in Northern and Southern Europeans for all variants in the
FADS locus (Fig. 2B; Methods). Our first analysis included four SNPs, three of which
(rs174594, rs174546, and rs174570) are top SNPs from this and previous studies[9,11] and are
highlighted in all our analyses, while the fourth SNP (rs4246215) is the one showing the biggest
difference in the upcoming South-North comparison analysis (Fig. 2B). The indel rs66698963
was not highlighted in this and all upcoming analyses because it has no significant selection
signals in Europe (Fig. 1). The original aDNA-based test evaluates the frequencies of a tested
allele in three ancient samples and four modern 1000GP samples under two hypotheses (H_0 and
H_1). Under H_1, maximum likelihood estimates (MLEs) of frequencies in all samples are
constrained only by observed allele counts and thus equivalent to the direct observed frequencies
(Fig. 2A; blue bars). Among the four modern samples, the observed adaptive allele frequencies
for all four SNPs exhibit a South-North gradient with the highest in Tuscans and the lowest in
Finns, consistent with the gradient of selection signals observed before based on mDNA (Fig. 1).
Among the three ancient samples, the observed allele frequencies, equivalent to the frequencies
upon admixture (Fig. 2A, orange bars for ancient groups), are always the lowest and often zero
in the WSHG sample.

Under H_0, the MLEs of frequencies are constrained by the observed allele counts and an
additional assumption that an allele's frequencies in the four modern samples are each a linear
combination of its frequencies in the three ancient samples. Considering the later assumption
alone, we can predict the frequencies of adaptive alleles right after admixture for each modern
population. Because the admixture contribution of WSHG, as estimated genome-wide, is higher
towards the North, constituting of 0%, 0%, 19.6%, and 36.2% for TSI, IBS, CEU, and GBR,
respectively[11], the predicted adaptive allele frequencies upon admixture for these four modern
populations are usually lower in the North (Fig. 2A; orange bars in modern populations),
suggesting higher starting frequencies in the South at the onset of selection. Further taking into
account observed allele counts in modern populations, we obtained the MLEs of frequencies
under H_0 (Fig. 2A; yellow bars in modern populations). As expected, the predicted allele

frequencies are higher in the South. But more importantly, the differences between H_0 and H_1
estimates in modern populations (Fig. 2A; indicated frequency differences between yellow and
orange bars) are still higher in the South, suggesting more recent factors, in addition to higher
starting frequencies, might contribute to the observation of stronger selection signals in the
South.

We systematically evaluated geographical differences of selection signals for all SNPs in the
FADS locus by applying the aDNA-based selection test separately for the two Southern and the
two Northern populations (Methods). All SNPs across the *FADS1-FADS2* LD block that were
significant in the combined analyses (Fig. 1A) were also significant in each of the two separate
analyses, but many exhibited much stronger signals in the analysis with Southern populations
(Fig. 2B; Supplementary Fig. 9). The maximum difference was found for SNP rs4246215, whose
p value in Southern populations is 12 orders of magnitude stronger than that in Northern
populations. SNP rs174594, rs174546 and rs174570 also have signals that are several orders (7,
11, and 10 respectively) of magnitude stronger in the South. A further decomposition of the
selection test and comparison of maximum likelihoods under the null and alternative hypotheses
between South and North revealed that a stronger deviation under the null hypothesis in the
South is driving the signal (Supplementary Fig. S10). This is also manifested by the bigger
differences between H_0 and H_1 estimates of adaptive allele frequencies in modern Southern
Europeans for the four representative SNPs (Fig. 2A, indicated frequency differences between
yellow and blue bars). It is noteworthy that the pattern of stronger signal in the South is observed
only for some but not all SNPs, excluding the possibility of systemic bias and pointing at SNP-
specific properties, likely for SNPs that are in LD with an underlying causal variant. Indeed, the
most common haplotype (referred to as haplotype D; Methods) within the *FADS1-FADS2* LD
block, also exhibits frequency patterns that are consistent with adaptive alleles of the four
representative SNPs: higher frequencies in the South among modern European populations,
while lowest frequency in WSHG among ancient groups (Fig. 2C). Hence, these results
demonstrated stronger selection signals on the *FADS1-FADS2* LD block in Southern Europeans.

Opposite selection signals in pre-Neolithic European hunter-gatherers

Motivated by the very different diet of pre-Neolithic European hunter-gatherers, we set to test
the action of natural selection on the *FADS* locus before the Neolithic revolution. The
availability of aDNA for pre-Neolithic European hunter-gatherers over a long historic period
offers this unprecedented opportunity. To this end, we started by examining the frequency
trajectory of haplotype D, the candidate adaptive haplotype in recent European history after the
Neolithic revolution. As noted above, its frequency increase drastically during recent European
history (Fig. 2C, the contrast between orange and blue bars). In stark contrast, it shows a clear
trajectory of decreasing frequency over time among pre-Neolithic hunter-gatherers[27] (Fig. 3A):
starting from 32% in the ~30,000-year-old (yo) "Věstonice cluster", through 21% in the ~15,000
yo "El Mirón cluster", to 13% in the ~10,000 yo "Villabruna cluster", and to being practically
absent in the ~7,500 yo WSHG group. We hypothesized that there was positive selection on
alleles opposite to the recently adaptive alleles that are associated with haplotype D.

To search for SNPs with evidence of positive selection during the pre-Neolithic period, we
considered the allele frequency time series for all SNPs around the *FADS* locus. We applied to
each SNP a rigorous, recently-published Bayesian method[28] to infer selection coefficients from

time series data while taking into account the European demographic history[29] (Methods). The
test highlighted two SNPs (rs174570 and rs2851682) within the *FADS1-FADS2* LD block
carrying suggestive evidence for the presence of positive selection during the pre-Neolithic
period tested, approximately 30,000-7,500 years ago (Supplementary Fig. 11). The derived
alleles of rs174570 and rs2851682 have similar frequency trajectories during this period,
increasing from 35.7% to 77.8% (Fig. 3B). It is noteworthy that the derived allele of rs174570
has also been shown to be targeted by positive selection in modern Greenlandic Inuit[9]. Moreover,
the ancestral alleles of rs174546 and rs174594 also experienced frequency increase from about
65% to almost fixation (Fig. 3B). However, presumably due to the high starting frequencies,
results from the time series test are not significant for these two SNPs. Importantly, for each of
these four SNPs, the allele experiencing frequency increase is opposite to the allele associated
with haplotype D, with the latter allele experiencing extreme frequency increase after the
Neolithic revolution.

We inferred selection coefficients concurrently with allele age for the derived alleles of rs174570
and rs2851682[28] (Methods). For rs1745470, the marginal maximum *a posteriori* (MAP)
estimates of selection coefficients (s_1 and s_2 respectively) for heterozygote and homozygote are
0.28% (95% credible interval (CI): -0.025% – 1.3%) and 0.38% (95% CI: 0.038% – 0.92%)
while the joint MAP of (s_1, s_2) is (0.24%, 0.34%). The age of the mutation giving rise to the
derived allele is estimated to be 57,380 years (95% CI: 157,690 – 41,930 years) (Fig. 3C,
Supplementary Fig. 12). For the derived allele of rs2851682, the marginal MAP for s_1 and s_2 are
0.31% (95% CI: 0.033% – 1.65%) and 0.40% (95% CI: 0.028% – 1.12%), while the joint MAP
is (0.26%, 0.35%) and its allele age is 53,440 years (95% CI: 139,620 – 39,320 years) (Fig. 3D,
Supplementary Fig. 13). As the observed allele frequency time series for the derived alleles of
these two SNPs fall well within the 95% CI of the posterior distribution (Figs. 3C and 3D), these
results support the presence of positive selection on these two alleles since their first appearance.
For both SNPs, s_2 is larger than s_1, suggesting there was directional selection and the presence of
derived allele was always beneficial. Additionally, we identified another haplotype in the
FADS1-FADS2 LD block, referred to as haplotype M2 (Methods, Supplementary Table 2), that
appears in modern Europeans at frequency of 10% but is much more common in Eskimos (Fig.
4A). 99% of chromosomes carrying haplotype M2 in modern Europeans also carries the derived
allele of rs174570, indicating a strong association between these two. Consistent with positive
selection on the derived allele of rs174570, haplotype M2 exhibits increasing frequency over
time in pre-Neolithic hunter-gatherers (Supplementary Table 2), suggesting that the causal allele
is associated with this haplotype.

The temporal and global evolutionary trajectory of *FADS* haplotypes

Thus far we have revealed haplotype M2 and D in the *FADS1-FADS2* LD block as the candidate
adaptive haplotype within Europe before and after the Neolithic revolution, respectively. To
further reveal a more complete picture of the evolutionary trajectories of haplotypes in that long
LD block, we performed more detailed analyses with global ancient and modern samples.
Specifically, we conducted a haplotype network analysis (Fig. 4A, Supplementary Table 2) with
450 ancient haplotypes (422 from ancient European samples included in the two previous
aDNA-based selection tests and additional 28 from representative ancient samples worldwide,

such as Neanderthal[30], Denisovan[31], Ust'-Ishim[32], Anzick[33], and Kennewick[34]) and 4,358 modern
haplotypes (4,314 from 1000GP and 44 from modern Eskimos). Moreover, we examined the
geographical frequency distribution of the resulting haplotypes in 29 previously-defined ancient
Eurasian groups[11,27,35] with 600 haplotypes from 300 ancient samples (Fig. 4B, Supplementary
Table 3; 422 haplotypes from ancient European samples included in the two previous aDNA-
based tests and additional ones from the Middle East[35]) and also in 27 modern groups from
1000GP and modern Eskimos (Fig. 4C, Supplementary Table 4; 5,008 haplotypes from 1000GP
and 44 from Eskimos).

The top five haplotypes in modern Europeans, designated as D, M1, M2, M3 and M4 from the
most to the least common (63.4%, 15.3%, 10.2%, 4.7%, 4.3%, respectively), were all observed
in aDNA and in modern Africans. They account for more than 95% of haplotypes in any extant
non-African populations, but only for 42% in extant African populations and for 64% in the
ancient samples (Fig. 4A, Supplementary Table 2). The difference between Africans and non-
Africans is consistent with the general Out-of-Africa dispersal carrying with it only a subset of
African haplotypes[36]. The additional difference between ancient and modern European samples
is consistent with the action of positive selection as already illustrated for haplotype D and M2,
which reduces haplotype diversity[37]. Among 450 aDNA haplotypes included in the haplotype
network analysis, the most common haplotypes are M2 (22%), D (17%), and M1 (16%).

Haplotype D has a frequency of 32% in the oldest European hunter-gatherer group, the ~30,000
yo "Věstonice cluster", and a frequency of 42% in the ~14,000 yo Epipalaeolithic Natufian
hunter-gatherers in the Levant (Fig. 4B, Supplementary Table 3), suggesting that it was of
relatively high frequency of ~35% in the Out-of-Africa ancestors. This number is also similar to
those in modern-day African populations (35% - 44%, Fig. 4C, Supplementary Table 4). As we
have shown above among pre-Neolithic European hunter-gatherers, the D frequency decreased
over time such that it was essentially absent by the advent of farming, possibly as a result of
positive selection on haplotype M2. In addition to its absence in WSHG, D was not observed in
the three ~7,500-year-old Eastern hunter-gatherers (EHG, Fig. 4B). D was re-introduced into
Europe with the arrival of farmers and Steppe-Ancestry pastoralists. Since the admixture of the
three ancient groups in Europe, the frequency of D has increased dramatically as a result of
positive selection, possibly driven by the dietary changes associated with farming. At the same
time, globally D also experienced dramatic frequency increase in South Asia and parts of East
Asia (Fig. 4C). However, D was absent in modern-day Eskimos.

Haplotype M2 has frequencies of 29% in the "Věstonice cluster" and of 25% in Natufian hunter-
gatherers, suggesting a medium frequency of ~27% at the time of Out-of-Africa dispersal
(Supplementary Table 3). However, this number is much higher than its current frequencies in
present-day Africans (0% - 3%, Supplementary Table 4), which might be a result of recent
positive selection on other haplotypes[5,6,8]. During the pre-Neolithic period, M2 increased in
frequency from 29% in the "Věstonice cluster" to 56% in WSHG and 50% in EHG
(Supplementary Table 3). After Neolithic revolution, the frequency of M2 decreased
dramatically to 10% among all present-day Europeans. There is also a South-North frequency
gradient for M2: TSI (4%), IBS (7%), CEU (9%), GBR (10%), and FIN (22%). It is noteworthy
that these two trends are opposite to those of haplotype D. Globally, in addition to its low
frequency in Africa, M2 has low frequency in South Asia (1% - 5%) but high frequency in
southern parts East Asia (44% - 53%, Supplementary Table 4). Its frequency in Eskimos is 27%.

Haplotype M1 has frequencies of 11% in the "Věstonice cluster" and of 8% in Natufian hunter-
gatherers, suggesting a low frequency of ~10% at the time of Out-of-Africa dispersal
(Supplementary Table 3). Similar to M2, this frequency is much higher than that in present-day
Africans (0% - 6%, Supplementary Table 4). In contrast to D and M2, M1 had little frequency
change during the pre-Neolithic period, maintaining at ~11% from the "Věstonice cluster" to
WSHG (Supplementary Table 3). It also had little frequency change over time in Europe, with a
frequency of ~15% in modern Europeans. Globally, M1 has overall low frequencies (<20%)
except for Eskimos and American populations (Supplementary Table 4). With a frequency of
73%, it dominates the haplotypes observed in Eskimos, making it the candidate adaptive
haplotype in this seafood-eating population[9].

The global frequency patterns of representative variants within the *FADS1-FADS2* LD block
(rs174570, rs66698963, rs174594, rs174546, and rs2851682; Fig. 4D, Supplementary Figs. 15-
19) mostly mirror those of key haplotypes, but with discrepancies that provide insights into
casual variants and allele ages. One major discrepancy was found in Africa. The derived alleles
of rs174570 and rs2851682 remains almost absent in Africa (Fig. 4D, Supplementary Figs. 15
and 19), consistent with their allele age estimates of ~55,000 years (Figs. 3C and 3D) and ruling
out their possible involvement in the positive selection on *FADS* genes in Africa[5,6,8]. Considering
the poor LD structure of the *FADS* locus in Africa (Supplementary Fig. 20), it is possible that
selection in Africa may be on haplotypes and causal variants that are different from those in
Europe.

Functional analyses of adaptive variants

Previous studies on adaptive evolution of the *FADS* locus suggested that alleles targeted by
positive selection are also associated with expression levels of *FADS* genes[5,6,8]. To test this
possibility in the context of this large-sale analysis, we considered data from the Genotype-
Tissue Expression (GTEx) project[38]. Our results point to many SNPs on the *FADS1-FADS2* LD
block being eQTLs of *FADS* genes. Out of a total of 44 tissues, these eQTLs at genome-wide
significance level are associated with the expression of *FADS1*, *FADS2*, and *FADS3* in 12, 23,
and 4 tissues, respectively, for a total of 27 tissues (Supplementary Figs. 21-23). Considering the
peak SNP rs174594 alone, nominally significant associations with these three genes were found
in 29, 28 and 4 tissues, respectively. More importantly, out of these tissues with association
signals, the adaptive allele in recent European history is associated with higher expression of
FADS1, lower expression of *FADS2* and higher expression of *FADS3* in 28, 27 and 4 tissues,
respectively. The general trend that recently adaptive allele is associated with higher expression
of *FADS1* but lower expression of *FADS2* was also observed for other representative SNPs
(rs174546, rs174570, and rs2851682) in the *FADS1-FADS2* LD block.

Genome-wide association studies (GWAS) have revealed 178 association signals with 44
different traits in the 85 kb *FADS1-FADS2* LD block, as recorded in the GWAS catalog
(Supplementary Tables 5-9)[39]. All effects reported in the following are based on GWAS
conducted with individuals of European ancestry, while some are also replicated in other ethnic
groups. Dissecting different associations, (1) the most prominent group of associated traits are
polyunsaturated fatty acids (PUFAs, Supplementary Fig. 1), including LCPUFAs and their
shorter chain precursors. Alleles on haplotype D are associated with higher levels of arachidonic

acid (20:4n-6, AA)[40-42], adrenic acid (22:4n-6, AdrA)[40,42-44], eicosapentaenoic acid (20:5n-3,
EPA)[42,45] and docosapentaenoic acid (22:5n-3, DPA)[42,43,45], but with lower levels of dihomo-
gamma-linolenic acid (20:3n-6, DGLA)[40-43], all of which suggest increased activity of delta-5
desaturase encoded by *FADS1*[42,46]. This is consistent with the association of recently adaptive
alleles with higher *FADS1* expression. Surprisingly, alleles on haplotype D are associated with
higher levels of gamma-linolenic acid (18:3n-6, GLA)[40,41,43] and stearidonic acid (18:4n-3,
SDA)[42], but with lower levels of linoleic acid (18:2n-6, LA)[40,41,43,47] and alpha- linolenic acid
(18:3n-3, ALA)[41,43,45], suggesting increased activity of delta-6 desaturase encoded by *FADS2*[41].
However, the above eQTL analysis suggested that adaptive alleles tend to be associated with
lower *FADS2* expression. Some of these association signals have been replicated across
Europeans[40,42-47], Africans[45], East Asians[41,45], and Hispanic/Latino[45]. (2) Besides PUFAs,
recently adaptive alleles on haplotype D are associated with decreased cis/trans-18:2 fatty
acids[48], which in turn is associated with lower risks for systemic inflammation and cardiac
death[48]. Consistently, adaptive alleles are also associated with decreased resting heart rate[49,50],
which reduces risks of cardiovascular disease and mortality. (3) With regards to other lipid
levels, adaptive alleles have been associated with higher levels of high-density lipoprotein
cholesterol (HDL)[51-56], low-density lipoprotein cholesterol (LDL)[51-53,57] and total cholesterol[51-53],
but with lower levels of triglycerides[51,52,55,56]. (4) In terms of direct association with disease risk,
adaptive alleles are associated with lower risk for inflammatory bowel diseases (IBD), both
Crohn's disease[58-60] and ulcerative colitis[60].

Going beyond known associations from the GWAS catalog, we analyzed data from the two
sequencing cohorts of the UK10K study. Focusing on the peak SNP rs174594, we confirmed the
association of the recently adaptive allele with higher levels of TC, LDL, and HDL. We further
revealed that its adaptive allele is associated with higher levels of additional lipids, Apo A1 and
Apo B (Supplementary Fig. 24). Taken together, adaptive alleles in the *FADS1-FADS2* LD
block, beyond their direct association with fatty acid levels, are associated with factors that are
mostly protective against inflammatory and cardiovascular diseases, and indeed also show direct
association with decreased risk of a type of inflammatory autoimmune diseases.

Discussion

Evidence for positive selection on *FADS* genes in Europe. For the first time, we revealed that
patterns of positive selection on *FADS* genes within Europe vary geographically, between the
North and the South, and temporally, before and after the Neolithic revolution. Positive selection
on *FADS* genes within Europe was initially reported in a recent aDNA-based study[11]. Here, we
repeated the aDNA-based analysis with much higher density of variants and confirmed the
presence of positive selection. Moreover, we strengthened this discovery by providing
independent evidence based on mDNA analyses. Both aDNA and mDNA results consistently
pointed to the region surrounding SNP rs174594 as the peak of signals, suggesting the possibility
of a causal variant in that region. Overall, selection signals revealed by both aDNA and mDNA
analyses coincide with an 85 kb LD block covering *FADS1* and *FADS2*. Within this LD block,
the most common haplotype in current Europeans, haplotype D, is the candidate adaptive
haplotype. With regards to the timing of the selection event underlying these signals, because the
aDNA-based analysis specifically models the frequency change from ancient to current samples,

the onset of selection must have occurred after the first admixture between early farmers and
northwestern hunter-gatherers which was around 8,500 years ago[11]. One of the top adaptive
SNPs reported in Greenlandic Inuit (rs174570)[9], also locates in the *FADS1-FADS2* LD block and
carries adaptive signals in Europeans based on our aDNA-based analysis and haplotype-based
test on mDNA. Interestingly, while its derived allele is adaptive Inuit[9], it is its ancestral allele
that is adaptive in Europeans, suggesting the presence of opposite selection pressures, possibly
because of very different diets in these two populations. The indel rs66698963, previously
reported to be adaptive in Africans, South Asians, and parts of East Asians, does not carry
significant adaptive signals in Europeans. However, there is a caveat that the imputation quality
for this indel might not be good enough. This indel is also a copy number variation and has a
very complex sequence context (Supplementary Text). 1000GP, the reference panel for
imputation, consists of known genotype calling errors for this indel[8]. Both of our aDNA-based
test and haplotype-based test revealed little signals for this indel, but the SFS-based test (Fay and
Fu's H) unraveled a local peak around the indel, although not reaching genome-wide
significance. Inaccurate imputation might explain this pattern because the first two tests are
single-variant-based test while the third one draws information from all SNPs within a 5 kb
window and thus is less affected by imputation inaccuracy of a single variant. Besides the
FADS1-FADS2 LD block, additional selection signals were detected with mDNA analyses (Figs.
1B and 1C) around the beginning of *FADS3*. Detailed analyses on this region are beyond the
scope of this study and will be published separately.

For the first time, we demonstrated geographical differences of positive selection on *FADS* genes
within Europe. The possibility of geographical differences was first suggested in our mDNA
analyses (nSL, iHS, and Fay and Wu's H), with the strongest signals always observed in
Southern Europeans, especially Tuscans. To formally evaluate the presence of geographical
differences, we used four SNPs as examples and dissected different layers of forces, either
demographic or selection, contributing to their final adaptive allele frequencies in current
European populations. We revealed three layers of forces. First, among the three ancient
samples, adaptive alleles always have the lowest frequencies or are even absent in western and
Scandinavian hunter-gatherers (Fig. 2A). This is consistent with our observation that opposite
selection forces operated in pre-Neolithic European hunter-gatherers and in more recent
European farmers. Second, there are differential admixture proportions of ancient sources for
Northern and Southern Europeans. The contribution of hunter-gatherers is higher towards the
North, while the contribution of early farmers is higher towards the South. As a result, the
predicted frequencies right after admixture are already higher in the South (Fig. 2A). Third, with
a null model taking into account the first two layers and also observed allele counts in modern
populations, we predicted current allele frequencies under neutrality. They are still lower than
observed allele frequencies, calculated directly from observed allele counts, indicating the
presence of positive selection as already detected in the aDNA-based test (Fig. 1A). More
importantly, the bigger differences in the two Southern European populations compared to the
two Northern populations suggest still stronger selection signals in the South (Fig. 2A), which
might be a result of stronger selection pressure or earlier onset of selection in Southern Europe.
These detailed analyses on the four SNPs were further confirmed by a global analysis on all
SNPs in the region with aDNA-based tests separately applied on Northern and Southern
Europeans. As the selection signal detected by the aDNA-based only describes the period
starting from the ancient admixture to present and the exact timing of ancient admixture could be
different for different populations, it is possible that ancient admixture finished earlier in

Southern Europe and there was a longer time for the action of selection, resulting in the stronger
signals we detected. The other possibility is stronger selection pressure in the South, which is
consistent with the dietary differences between Southern and Northern European farmers as
discussed in the next section.

We also unraveled a novel discovery regarding the temporal differences of positive selection
signals within Europe before and after the Neolithic revolution. Haplotype D in the *FADS1-*
FADS2 LD block, the candidate adaptive haplotype during recent European history, exhibits
gradual frequency decrease over time among four groups of pre-Neolithic Hunter-gatherers, from
approximately 30,000-7,500 years ago. With a recently-published Bayesian method[28] for
inferring selection coefficients from allele frequency time series data, we identified two SNPs
(rs174570 and rs2851682) with evidence of positive selection during this period. The ages of the
derived alleles for these two SNPs are similar, about 55,000 years, after the Out-of-Africa
dispersal. This is consistent with the near absence of these two alleles in modern Africans (Fig.
4D, Supplementary Figs. 15 and19). Although the trend of increasing frequency over time was
also observed for other SNPs in the region (e.g. rs174546 and rs174594), the formal test did not
reveal significant signals for them. Several factors could potentially contribute to reduced power
of the test, including the higher starting frequencies for some SNPs, the small sample size for
each group, and the use of samples of different ages in the same group. Future studies with much
bigger sample size are needed to refine the selection signal for this pre-Neolithic period.
Additionally, it will be of interest in the future to explore potential geographical differences
among hunter-gatherers, especially considering the dietary differences between Northern and
Southern pre-Neolithic hunter-gatherers, which are discussed in the next section.

Interpretation of positive selection signals in light of anthropological findings. The dispersal
of the Neolithic package into Europe that began some 8,500 years ago caused a sharp dietary
shift from an animal-based diet with significant aquatic contribution to a terrestrial plant-heavy
diet including dairy products[15-20]. Before the Neolithic revolution, consumption of aquatic food
had been prominent in diets of pre-Neolithic European hunter-gatherers[61]. The significant role of
aquatic food, either marine or freshwater, has been established in sites along the Atlantic
coast[17,62-64], around the Baltic sea[17], and along the Danube river[65]. The content of LCPUFAs are
usually the highest in aquatic foods, lower in animal meat and milk, and almost negligible in
most plants[66]. Consistent with the subsistence strategy and dietary pattern in pre-Neolithic
hunter-gatherers, positive selection on *FADS* genes during this period was on alleles that are
associated with less efficient endogenous synthesis of LCPUFAs, possibly compensating for the
high dietary input. In addition to optimal absolute levels of LCPUFAs, maintaining a balanced
ratio of omega-6 to omega-3 is also critical for human health[67]. It is also possible that positive
selection on *FADS* genes in hunter-gatherers was in response to an unbalanced omega-6 to
omega-3 ratio (e.g. too much omega-3 LCPUFAs). Similar selection signals on *FADS* genes
have been observed in modern Greenlandic Inuit, who subsist on a seafood diet[9]. Specifically,
the derived allele of SNP rs174570 carries positive selection signals in both pre-Neolithic
European hunter-gatherers and modern Greenlandic Inuit. More generally, haplotype M2, the
candidate adaptive haplotype during the pre-Neolithic period in Europe, is also common in the
modern Eskimo samples examined in our study. It is noteworthy that aquatic food was less
prevalent among pre-Neolithic hunter-gatherers around the Mediterranean basin, possibly due to
the low productivity of the Mediterranean Sea[68-70]. It would be interesting to examine the
geographical differences of selection signals among different European groups of pre-Neolithic

hunter-gatherers. However, aDNA from pre-Neolithic hunter-gatherers is still scarce and under-
represented around the Mediterranean basin, prohibiting such an analysis at present.

The Neolithization of Europe[12,71,72] started in the Southeast region around 8,500 years ago when
farming and herding spread into the Aegean and the Balkans. It continued in spite of a few
temporary stops into central and northern Europe following the Danube River and its tributaries,
and along the Mediterranean coast. It arrived at the Italian Peninsula about 8,000 years ago and
shortly after reached the Iberia by 7,500 years ago. While farming rapidly spread across the loess
plains of Central Europe and reach the Paris Basin by 7,000 years ago, it took another 1,000 or
more years before it spread into Britain and Northern Europe around 6,000 years ago. From that
time on, European farmers relied heavily on their domesticated animals and plants. Compared to
pre-Neolithic hunter-gatherers, European farmers consumed much more plants but less aquatic
foods[18-20,73]. Consistent with the lack of LCPUFAs in plant-based diets, positive selection on
FADS genes during recent European history has been on alleles that are associated with
enhanced endogenous synthesis of LCPUFAs from plant-derived precursors (LA and ALA).
Positive selection for enhanced LCPUFAs was also observed before in Africans, South Asians
and some East Asians, possibly driven by the local traditional plant-based diets[8].

Despite the overall trend of relying heavily on domesticated plants, there are geographical
differences of subsistence strategies and dietary patterns among European farmers. In addition to
the 2,000-year-late arrival of farming at Northern Europe, animal husbandry and the
consumption of animal milk became gradually important as Neolithic farmers spread to the
Northwest[18,72,74-76]. Moreover, similar to their pre-Neolithic predecessors, Northwestern
European farmers close to the Atlantic Ocean or the Baltic Sea still consumed some marine food,
more so than their Southern counterparts in the Mediterranean basin[77,78]. It is noteworthy that
historic dairying practice in Northwestern Europe has driven the adaptive evolution of lactase
persistence in Europe to reach the highest prevalence in this region[75]. In this study, we observed
stronger positive selection signals on *FADS* genes during recent history in Southern than in
Northern Europeans, even after considering the later arrival of farming and the lower starting
allele frequencies in the North. The higher aquatic contribution and stronger reliance on animal
meat and milk might be responsible for the weaker selection pressure in the North, although the
possibilities of other environmental factors could not be ruled out.

Interpretation of eQTLs and GWAS results. Although liver is the primary site for the
endogenous synthesis of LCPUFAs, the action of the pathway has been observed in a wide range
of tissues[79,80], including heart[81], brain[81-83], both white and brown adipose tissues[84]. Moreover,
while the synthesis rate and relevant enzyme levels in liver are regulated by dietary fatty acid
inputs, they are not affected in other tissues[81], indicating that identifying eQTLs for *FADS* genes
in the liver might need extra control for dietary inputs. Based on data from the GTEx project,
eQTLs within the *FADS1-FADS2* LD block for the three *FADS* genes were identified in multiple
tissues and in general recently adaptive alleles are associated with higher *FADS1* expression but
lower *FADS2* expression. No genome-wide significant eQTLs for *FADS1* and *FADS2* were
found in the liver, probably due to the complication of dietary inputs, which were not available to
be controlled for during analysis. However, an apparent cluster of elevated association signals
with *FADS1* was observed in the liver, although they do not reach genome-wide significance
level (Supplementary Fig. 21). Furthermore, for the recently adaptive allele of peak SNP
rs174594, the directions of association with *FADS1* and *FADS2* in the liver, although not
significant, are consistent with the general trend – higher *FADS1* but lower *FADS2* expression.

The exact causal regulatory variants and the underlying mechanisms are still unknown, but
variants disrupting the sterol response element (SRE) are among the most likely candidates[2].

GWAS revealed several potential beneficial effects of the recently adaptive alleles: enhanced
efficiency of the overall LCPUFAs synthesis, lower risks of systemic inflammation,
inflammatory bowel diseases, and cardiovascular diseases. The directions of association with
PUFAs along the synthesis pathway (Supplementary Fig. 1) reflect the relative efficiency of rate-
limiting enzymes, delta-5 and delta-6 desaturases: enhanced detal-5 desaturase activity is
expected to reduce levels of its precursors, LA and ALA, but to increase levels of its products,
GLA and SDA, while similarly enhanced delta-6 desaturase activity is expected to reduce DGLA
and ETA, but to increase levels of AA, AdrA, EPA and DPA. While GWAS results are
consistent with eQTLs analysis in revealing increased *FADS1* expression and enhanced delta-5
desaturase activity, they seem contradictory for *FADS2*: recently adaptive alleles are associated
with lower *FADS2* expression but enhanced delta-6 desaturase activity. There are several
possible explanations. First, the *FADS2* expression level might not directly correlate with the
final delta-6 desaturase level because of post-transcriptional regulation. Second, the direction of
FADS2 eQTLs might be different in the liver from other tissues. Currently, there are marginal
association signals for *FADS1* in the liver but no signals for *FADS2*. Additional analysis for
FADS2 is needed in the liver with proper control for dietary inputs. Third, there may be
alternative splicing in addition to expression level change. Further experiments are needed to
address this discrepancy and to unravel the underlying molecular mechanisms. Besides PUFAs,
GWAS also revealed an overall trend that recently adaptive alleles are protective against
inflammatory conditions, especially inflammatory bowel diseases. But there are exceptions:
these alleles were also found to be associated with increased risk of rheumatoid arthritis[85] and
colorectal cancer[86]. Because LCPUFAs-derived signaling molecules have both pro-inflammatory
and anti-inflammatory effects (Supplementary Fig. 1), elucidating the effects of these adaptive
alleles on specific diseases will require case-by-case analysis with special consideration of the
relative contributions of omega-6 and omega-3 LCPUFAs. The effort in understanding the
clinical significance of genetic variants in *FADS* genes might also reveal additional selection
pressures beyond diet acting on these genes.

Conclusions

In summary, we demonstrated that in Europe an extended LD block covering *FAD1* and *FADS2*
of the *FADS* gene family has been under strong recent positive selection both before and after the
Neolithic revolution. During the recent history, positive selection also varies geographically,
with selection signals and adaptive allele frequencies gradually increasing from Northern
towards Southern Europe. The plant-heavy diet of European farmers, with its lack of LCPUFAs,
is one possible environmental factor contributing to the recent positive selection. The higher
consumption of aquatic resources and animal milk among Northwestern European farmers might
contribute to the weaker selection signals observed in the North. Consistently, many alleles on
the recently adaptive haplotype are eQTLs that increase *FADS1* expression, thereby increasing
the efficacy of LCPUFAs synthesis. Additional evidence comes from a multitude of GWAS
showing recently adaptive alleles associated with enhanced LCPUFAs biosynthesis. Before the
advent of farming, the recently adaptive haplotype showed dramatic decrease in frequency across
pre-Neolithic hunter-gatherers. While this could have been due to negative selection affecting
alleles on the haplotype, time series analysis showed that it was driven by positive selection on

alleles opposite to those on the recently adaptive haplotype. Considering that pre-Neolithic
hunter-gatherers subsisted on animal-based diets with significant aquatic contribution, limiting
the rate of endogenous LCPUFAs synthesis by decreasing *FADS1* expression might be beneficial
and contributed to that ancient adaptation. This discovery of subsistence-based temporal and
geographical variations of selection in Europe supports and completes the global picture of the
local adaptation of *FADS* genes: positive selection on alleles enhancing LCPUFAs biosynthesis
in populations traditionally subsisting on plant-based diets[5,6,8], but positive selection on opposite
alleles in populations subsisting on a LCPUFAs-rich marine diet[9]. This opposite pattern of
positive selection in different dietary environment highlights the potential of matching diet to
genome in the future nutritional practice. Finally, the vast number of traits associated with the
adaptive region in the *FADS* genes, while raising the possibility of additional selection forces
beyond diet, stresses the clinical and nutritional significance of understanding the evolutionary
forces shaping the *FADS* gene family and other diet-related genes.

Methods

Data sets. The ancient DNA (aDNA) data set included in this study was compiled from two
previous studies[27,35], which in turn were assembled from many other studies[11,21,22,30-34,87-96], in
addition to new sequenced samples. These two data sets were downloaded from
https://reich.hms.harvard.edu/datasets and were merged by removing overlapping samples. In
total, there are 325 ancient samples included in this study (Supplementary Table 1). For the
aDNA-based test for recent selection in Europe, a subset of 178 ancient samples were used and
clustered into three groups as in the original study[11], representing the three major ancestral
sources for most present-day European populations. These three groups are: West and
Scandinavian hunter-gatherers (WSHG, N=9), early European farmers (EF, N=76), and
individuals of Steppe-pastoralist Ancestry (SA, N=93). Three samples in the EF group in the
original study were excluded from our analysis because they are genetic outliers to this group
based on additional analysis[35]. For aDNA-based test for ancient selection in pre-Neolithic
European hunter-gatherers, a subset of 42 ancient samples were used and four groups were
defined. In addition to the WSHG (N=9), the other three groups were as originally defined in a
previous study[27]: the "Věstonice cluster", composed of 14 pre-Last Glacial Maximum
individuals from 34,000-26,000 years ago; the "El Mirón cluster", composed of 7 post-Last
Glacial Maximum individuals from 19,000-14,000 years ago; the "Villabruna cluster", composed
of 12 post-Last Glacial Maximum individuals from 14,000-7,000 years ago. There were three
Western hunter-gatherers that were originally included in the "Villabruna cluster"[27], but we
included them in WSHG in the current study because of their similar ages in addition to genetic
affinity[11]. In haplotype network analysis, all aDNAs included in the two aDNA-based selection
tests were also included in this analysis. In addition, we included some well-known ancient
samples, such as the Neanderthal, Denisovan, and Ust'-Ishim. In total, there were 225 ancient
samples (450 haplotypes). For geographical frequency distribution analysis, a total of 300
ancient samples were used and classified into 29 previously defined groups[11,27,35] based on their
genetic affinity, sampling locations and estimated ages.

Data for the 1000 Genomes Project (1000GP, phase 3)[7] were downloaded from the official FTP
site (ftp://ftp.1000genomes.ebi.ac.uk/vol1/ftp/release/20130502/). There are in total 2,504
individuals from 5 continental regions and 26 global populations. There are 7 populations of
African ancestry (AFR, N=661): Yoruba in Ibadan, Nigeria (YRI, N=108), Luhya in Webuye,
Kenya (LWK, N=99), Gambian in Western Divisions in the Gambia (GWD, N=113), Mende in
Sierra Leone (MSL, N=85), Esan in Nigeria (ESN, N=99), Americans of African Ancestry in
SW USA (ASW, N=61), African Caribbeans in Barbados (ACB, N=96); 5 populations of
European ancestry (EUR, N=503): Utah Residents with Northern and Western European
Ancestry (CEU, N=99), Toscani in Italia (TSI, N=107), Finnish in Finland (FIN, N=99), British
in England and Scotland (GBR, N=91), Iberian Population in Spain (IBS, N=107); 5 populations
of East Asian ancestry (EAS, N=504): Han Chinese in Beijing, China (CHB, N=103), Japanese
in Tokyo, Japan (JPT, N=104), Southern Han Chinese (CHS, N=105), Chinese Dai in
Xishuangbanna, China (CDX, N=93), Kinh in Ho Chi Minh City, Vietnam (KHV, N=99); 5
populations of South Asian ancestry (SAS, N=489): Gujarati Indian from Houston, Texas (GIH,
N=103), Punjabi from Lahore, Pakistan (PJL, N=96), Bengali from Bangladesh (BEB, N=86),
Sri Lankan Tamil from the UK (STU, N=102), Indian Telugu from the UK (ITU, N=102), and 4
populations of American ancestry (AMR=347): Mexican Ancestry from Los Angeles USA
(MXL, N=64), Puerto Ricans from Puerto Rico (PUR, N=104), Colombians from Medellin,
Colombia (CLM, N=94), Peruvians from Lima, Peru (PEL, N=85).

The data set for Human Genome Diversity Project (HGDP)[97] was downloaded from
http://www.hagsc.org/hgdp/files.html. There were ~650K SNPs in 939 unrelated individuals
from 51 populations. The data from the Population Reference Sample (POPRES)[98] were
retrieved from dbGaP with permission. Only 3,192 Europeans were included in our analysis. The
country of origin of each sample was defined with two approaches. Firstly, a "strict consensus"
approach was used: an individual's country of origin was called if and only if all four of his/her
grandparents shared the same country of origin. Secondly, a more inclusive approach was used to
further include individuals that had no information about their grandparents. In this case, their
countries of birth were used. Both approaches yielded similar results and only results from the
inclusive approach are reported. The 22 Eskimo samples were extracted from the Human Origins
dataset[22].

The two sequencing cohorts of UK10K were obtained from European Genome-phenome
Archive with permission[99]. These two cohorts, called ALSPAC and TwinsUK, included low-
depth whole-genome sequencing data and a range of quantitative traits for 3,781 British
individuals of European ancestry (N=1,927 and 1,854 for ALSPAC and TwinsUK,
respectively)[99].

Imputation for ancient and modern DNA. Genotype imputation was performed using Beagle
4.1[100] separately for the data sets of aDNA, HGDP and POPRES. The 1000GP phase 3 data were
used as the reference panel[7]. Imputation was performed for a 5-Mb region surrounding the *FADS*
locus (hg19:chr11: 59,100,000-64,100,000), although most of our analysis was restricted to a 200
kb region (hg19:chr11:61,500,000-61,700,000). For most of our analysis (e.g. estimated allele
count or frequency for each group), genotype probabilities were taken into account without

setting a specific cutoff. For haplotype-based analysis (e.g. estimated haplotype frequency for
each group), a cutoff of 0.8 was enforced and haplotypes were defined with missing data (if the
genotype does not reach the cutoff) following the phasing information from imputation.

Genotype imputation for aDNA has been shown to be desirable and reliable[88]. We also evaluated
the imputation quality for aDNA by comparing with the two modern data sets (Supplementary
Fig. 25). Overall, the imputation accuracy for ungenotyped SNPs, measured with allelic R^2 and
dosage R^2, is comparable between aDNA and HGDP, but is higher in aDNA when compared
with POPRES. Note that the sample sizes are much larger for HGDP (N=939) and POPRES
(N=3,192), compared to aDNA (N=325). The comparable or even higher imputation quality in
aDNA was achieved because of the higher density of genotyped SNPs in the region.

Linkage disequilibrium and haplotype network analysis. Linkage disequilibrium (LD)
analysis was performed with the Haploview software (version 4.2)[101]. Analysis was performed
on a 200-kb region (chr11:61,500,000-61,700,000), covering all three *FADS* genes. Variants
were included in the analysis if they fulfilled the following criteria: 1) biallelic; 2) minor allele
frequency (MAF) in the sample not less than 5%; 3) with rsID; 4) *p* value for Hardy-Weinberg
equilibrium test larger than 0.001. Analysis was performed separately for the combined UK10K
cohort and each of the five European populations in 1000G.

Haplotype network analysis was performed with the R software package, pegas[102]. To reduce the
number of SNPs and thus the number of haplotypes included in the analysis, we restricted this
analysis to part of the 85 kb *FADS1-FADS2* LD block, starting 5 kb downstream of *FDAS1* to
the end of the LD block (a 60-kb region). To further reduce the number of SNPs, in the analysis
with all 1000GP European samples, we applied an iterative algorithm[103] to merge haplotypes that
have no more than three nucleotide differences by removing the three corresponding SNPs. The
algorithm stops when all remaining haplotypes are more than 3 nucleotides away. With this
procedure, we were able to reduce the number of total haplotypes from 81 to 12, with the number
of SNPs decreased from 88 to 34 (Supplementary Fig. 26). This set of 34 representative SNPs
was used in all haplotype-based analysis in aDNA, 1000GP, HGDP and POPRES. Missing data
(e.g. from a low imputation genotype probability) were included in the haplotype network
analysis.

Of note, for the 12 haplotypes identified in 1000GP European samples, only five of them have
frequency higher than 1% (Supplementary Table 2). These five haplotypes were designated as D,
M1, M2, M3 and M4, from the most common to the least.

Ancient DNA-based test for recent selection in Europe. The ancient DNA-based selection test
was performed as described before[11]. Briefly, most European populations could be modelled as a
mixture of three ancient source populations at fixed proportions. The three ancient source
populations are West or Scandinavian hunter-gatherers (WSHG), early European farmers (EF),
and Steppe-Ancestry pastoralist (SA) (Supplementary Table 1). For modern European
populations in 1000G, the ancestral proportions of these three populations estimated at genome-
wide level are (0.196, 0.257, 0.547) for CEU, (0.362, 0.229, 0.409) for GBR, (0, 0.686, 0.314)
for IBS, and (0, 0.645, 0.355) for TSI. FIN was not used because it does not fit this three-

population model[11]. Under neutrality, the frequencies of a SNP (e.g. reference allele) in present-
day European populations are expected to be the linear combination of its frequencies in the
three ancient source populations. This serves as the null hypothesis: $p_{mod} = Cp_{anc}$, where
p_{mod} is the frequencies in A modern populations (A is always 3 in our test), p_{anc} is the
frequencies in B ancient source populations while C is an AxB matrix with each row
representing the estimated ancestral proportions for one modern population. The alternative
hypothesis is that p_{mod} is unconstrained by p_{anc}. The frequency in each population is modelled
with binomial distribution: $L(p; D) = B(X, 2N, p)$, where X is the number of designated allele
observed while N is the sample size. In ancient populations, X is the expected number of
designated allele observed, taking into account uncertainty in imputation. We write $\ell(p; D)$ for
the log-likelihood. The log-likelihood for SNP frequencies in all three ancient populations and
four modern populations are: $\ell(\vec{p}; \vec{D}) = \sum_{i=1}^{A} \ell(p_i; D_i) + \sum_{j=1}^{B} \ell(p_j; D_j)$. Under the null
hypothesis, there are A parameters in the model, corresponding to the frequencies in A ancient
populations. Under the alternative hypothesis, there are A+B parameters, corresponding to the
frequencies in A ancient populations and B modern populations. We numerically maximized the
likelihood separately under each hypothesis and evaluate the statistic (twice the difference in log-
likelihood) with the null χ_B^2 distribution. Inflation was observed with this statistic in a previous
genome-wide analysis and a $\lambda = 1.38$ was used for correction in the same cases of three ancient
source populations and four present-day European populations [11]. Following this, we applied the
same factor in correcting the p values in our analysis. For genotyped SNPs previously tested,
similar scales of statistical significance were observed as in the previous study (Supplementary
Fig. 27). We note that for the purpose of refining the selection signal with imputed variants, only
relative significance levels across variants are informative.

In addition to combining signals from four present-day European populations, we further
performed tests separately in the two South European populations (IBS and TSI) and in the two
North European populations (CEU and GBR). In these two cases, B = 2 and the null distribution
is χ_2^2. No genomic correction was performed for these two cases.

Ancient DNA-based test for ancient selection in pre-Neolithic European hunter-gatherers.
A Bayesian method[28] was applied to infer natural selection from allele frequency time series
data. The software was downloaded from https://github.com/Schraiber/selection. This method
models the evolutionary trajectory of an allele under a specified demographic history and
estimates selection coefficients (s_1 and s_2) for heterozygote and homozygote of the allele under
study. This method has two modes, with or without the simultaneous estimation of allele age
(with or without "-a" in the command line). Without the estimation of allele age, this method
models the frequency trajectory only between the first and last time points provided and its
estimates of selection coefficients describe the selection force during this period only. With the
simultaneous estimation of allele age, this method models the frequency trajectory starting from
the first appearance of the allele to the last time point provided. In this case, the selection
coefficients describe the selection force starting from the mutation of the allele, which therefore
should be the derived allele. For demographic history, we used the model with two historic
epochs of bottleneck and recent exponential growth[29]. However, the recent epoch of exponential

growth does not have an impact on our analysis because for our analysis the most recent sample, WSHG, had an age estimate of around 7500 years ago, predating the onset of exponential growth (3520 years ago, assuming 25 years per generation). Four groups of pre-Neolithic European hunter-gatherers were included in our test: the Věstonice cluster (median sample age: 30,076 yo), the El Mirón cluster (14,959 yo), the Villabruna cluster (10,059 yo) and WSHG (7,769 yo).

The use of allele frequency time series data in this Bayesian method makes several assumptions, including 1) all samples are from a randomly mating population with continuity of genetic ancestry; and 2) samples are drawn at different time points[28]. Although there was population structure among pre-Neolithic hunter-gatherers, the four groups used in our study were clustered mainly based on their genetic affinity with additional filtering based on their archaeological contexts[27], therefore population structure in each group was minimized. There is also demonstrated shared genetic ancestry among these groups[27]. Each of the four groups includes samples of different ages and the median sample age was used to represent the sampling time of the group. This approach might introduce noise into the time series and thus reduce the power of the method, making the test conservative. Overall, our time series data do not deviate from these assumptions.

To identify SNPs with evidence of positive selection during the historic period covered by available ancient samples (from Věstonice to WSHG), we first ran the software for most SNPs in the *FADS* locus without the simultaneous estimation of allele age. SNPs with small frequency difference ($< 5\%$) between the first (Věstonice) and last (WSHG) time points were not included in the analysis. For each tested SNP, the allele under analysis was the one showing increasing frequency at the last time point compared to the first. Allele frequency time series data, for each tested SNP, were provided to the software as the expected number of the allele (calculated based on genotype probability) and the sample size. Each software run generated 1,000 Markov chain Monte Carlo (MCMC) samples out of 1,000,000 MCMC simulations with a sampling frequency of every 1,000. The effective sample size of these 1,000 samples were evaluated with the R package, coda[104]. Only runs with effective sample size larger than 50 for four parameters (the sampling likelihood, path likelihood, α_1 estimate, and α_2 estimate) were used[28]. A maximum of 100 runs were attempted for each SNP until a run with sufficient effective sample size was achieved. Otherwise, the SNPs were discarded in our analysis. Visual examination of the observed frequency trajectory for multiple failed SNPs revealed that none of them showed increasing frequency over time and therefore they were unlikely to be under selection. For SNPs with successful software runs, the maximum *a posteriori* (MAP) estimates and the 90% credible intervals (CI) for s_1 and s_2 were calculated. Suggestive evidence for positive selection was called if the 90% CI does not overlap with 0. Second, for the two candidate SNPs (rs174570 and rs2851682) identified in the unbiased global analysis, we further ran the software with the simultaneous estimation of derived allele age. The inference results were plotted with R scripts accompanying the software and additional customized scripts (available upon request).

Modern DNA-based selection tests. We performed two types of selection tests for modern DNAs: site frequency spectrum (SFS)-based and haplotype-based tests. These tests were performed separately in each of the five European populations from 1000G and each of the two cohorts from UK10K. For SFS-based tests, we calculated genetic diversity (π), Tajima's D[105],

and Fay and Wu's H[26], using in-house Perl scripts (available upon request). We calculated these
three statistics with a sliding-window approach (window size = 5 kb and moving step = 1 kb).
Statistical significance for these statistics were assessed using the genome-wide empirical
distribution. Haplotype-based tests, including iHS[24] and nSL[25], were calculated using software
selscan (version 1.1.0a)[106]. Only common biallelic variants (Minor allele frequency > 5%) were
included in the analysis. Genetic variants without ancestral information were further excluded.
These two statistics were normalized in their respective frequency bins (1% interval) and the
statistical significance of the normalized iHS and nSL were evaluated with the empirical
genome-wide distribution. The haplotype bifurcation diagrams and EHH decay plots were drawn
using an R package, rehh[107].

Geographical frequency distribution analysis. For plots of geographical frequency
distribution, the geographical map was plotted with R software package, maps (https://CRAN.R-
project.org/package=maps) while the pie charts were added with the mapplots package
(https://cran.r-project.org/web/packages/mapplots/index.html). Haplotype frequencies were
calculated based on haplotype network analysis with pegas[102], which groups haplotypes while
taking into account missing data. SNP frequencies were either the observed frequency, if the
SNP was genotyped, or the expected frequency based on genotype probability, if the SNP was
imputed.

Targeted association analysis for peak SNP rs174594 in UK10K. We performed association
analysis for rs174594 in two UK10K datasets – ALSPAC and TwinsUK[99]. For both datasets, we
analyzed height, weight, BMI and lipid level related traits including total cholesterol (TC), low
density lipoprotein (LDL), very low density lipoprotein (VLDL), high density lipoprotein
(HDL), Apolipoprotein A-I (APOA1), Apolipoprotein B (APOB) and triglyceride (TRIG). We
performed principal components analysis using smartpca from EIGENSTRAT software[108] with
genome-wide autosomal SNPs and we added top 4 principal components as covariates for all
association analysis. We also used age as a covariate for all association analysis. Sex was added
as a covariate only for ALSPAC dataset since all individuals in TwinsUK dataset are female. For
all lipid-related traits, we also added BMI as a covariate.

Data availability.

Ancient DNA: https://reich.hms.harvard.edu/datasets

1000 Genomes Project: ftp://ftp.1000genomes.ebi.ac.uk/vol1/ftp/release/20130502/

Human Genome Diversity Project (HGDP): http://www.hagsc.org/hgdp/files.html

Population Reference Sample (POPRES): dbGaP Study Accession: phs000145.v4.p2

UK10K: https://www.uk10k.org/data_access.html

Code availability. Most analyses were conducted with available software and packages as
described in the corresponding subsections of Methods. Customized Perl and R scripts were used
in performing SFS-based selection test, and for general plotting purposes. All these scripts are
available upon request (Contact K.Y. at ky279@cornell.edu).

References

Fan, S., Hansen, M. E. B., Lo, Y. & Tishkoff, S. A. Going global by adapting local: A review of recent human adaptation. *Science* **354**, 54-59 (2016).

Nakamura, M. T. & Nara, T. Y. Structure, function, and dietary regulation of delta6, delta5, and delta9 desaturases. *Annu Rev Nutr* **24**, 345-376 (2004).

Raphael, W. & Sordillo, L. M. Dietary polyunsaturated fatty acids and inflammation: the role of phospholipid biosynthesis. *Int J Mol Sci* **14**, 21167-21188 (2013).

Bazinet, R. P. & Laye, S. Polyunsaturated fatty acids and their metabolites in brain function and disease. *Nat Rev Neurosci* **15**, 771-785 (2014).

Mathias, R. A. *et al.* Adaptive evolution of the FADS gene cluster within Africa. *PLoS One* **7**, e44926 (2012).

Ameur, A. *et al.* Genetic adaptation of fatty-acid metabolism: a human-specific haplotype increasing the biosynthesis of long-chain omega-3 and omega-6 fatty acids. *Am J Hum Genet* **90**, 809-820 (2012).

The 1000 Genomes Project Consortium *et al.* A global reference for human genetic variation. *Nature* **526**, 68-74 (2015).

Kothapalli, K. S. *et al.* Positive Selection on a Regulatory Insertion-Deletion Polymorphism in FADS2 Influences Apparent Endogenous Synthesis of Arachidonic Acid. *Mol Biol Evol* **33**, 1726-1739 (2016).

Fumagalli, M. *et al.* Greenlandic Inuit show genetic signatures of diet and climate adaptation. *Science* **349**, 1343-1347 (2015).

Reardon, H. T. *et al.* Insertion-deletions in a FADS2 intron 1 conserved regulatory locus control expression of fatty acid desaturases 1 and 2 and modulate response to simvastatin. *Prostaglandins Leukot Essent Fatty Acids* **87**, 25-33 (2012).

Mathieson, I. *et al.* Genome-wide patterns of selection in 230 ancient Eurasians. *Nature* **528**, 499-503 (2015).

Bar-Yosef, O. in *On Human Nature: Biology, Psychology, Ethics, Politics, and Religion* (eds M. Tibayrenc & F. J. Ayala) Ch. 19, 297-331 (Academic Press, 2017).

Coward, F., Shennan, S., Colledge, S., Conolly, J. & Collard, M. The spread of Neolithic plant economies from the Near East to northwest Europe: a phylogenetic analysis. *Journal of Archaeological Science* **35**, 42-56 (2008).

Bogaard, A. *et al.* Crop manuring and intensive land management by Europe's first farmers. *Proc Natl Acad Sci U S A* **110**, 12589-12594 (2013).

Richards, M. P. in *The Evolution of Hominin Diets: Integrating Approaches to the Study of Palaeolithic Subsistence* (eds J. J. Hublin & M. P. Richards) 251-257 (Springer Science; Business Media, 2009).

Richards, M. P., Schulting, R. J. & Hedges, R. E. Archaeology: sharp shift in diet at onset of Neolithic. *Nature* **425**, 366 (2003).

Richards, M. P., Price, T. D. & Koch, E. Mesolithic and Neolithic Subsistence in Denmark:New Stable Isotope Data. *Current Anthropology* **44**, 288-295 (2003).

Fraser, R. A., Bogaard, A., Schäfer, M., Arbogast, R. & Heaton, T. H. E. Integrating botanical, faunal and human stable carbon and nitrogen isotope values to reconstruct land use and palaeodiet at LBK Vaihingen an der Enz, Baden-Württemberg. *World Archaeology* **45**, 492-517 (2013).

Knipper, C. *et al.* What is on the menu in a Celtic town? Iron Age diet reconstructed at Basel-Gasfabrik, Switzerland. *Archaeological and Anthropological Sciences* (2016).

20   López-Costas, O., Müldner, G. & Martínez Cortizas, A. Diet and lifestyle in Bronze Age Northwest
Spain: the collective burial of Cova do Santo. *Journal of Archaeological Science* **55**, 209-218
(2015).
21   Haak, W. *et al.* Massive migration from the steppe was a source for Indo-European languages in
Europe. *Nature* **522**, 207-211 (2015).
22   Lazaridis, I. *et al.* Ancient human genomes suggest three ancestral populations for present-day
Europeans. *Nature* **513**, 409-413 (2014).
23   Patterson, N. *et al.* Ancient admixture in human history. *Genetics* **192**, 1065-1093 (2012).
24   Voight, B. F., Kudaravalli, S., Wen, X. & Pritchard, J. K. A map of recent positive selection in the
human genome. *PLoS Biol* **4**, e72 (2006).
25   Ferrer-Admetlla, A., Liang, M., Korneliussen, T. & Nielsen, R. On detecting incomplete soft or
hard selective sweeps using haplotype structure. *Mol Biol Evol* **31**, 1275-1291 (2014).
26   Fay, J. C. & Wu, C. I. Hitchhiking under positive Darwinian selection. *Genetics* **155**, 1405-1413
(2000).
27   Fu, Q. *et al.* The genetic history of Ice Age Europe. *Nature* **534**, 200-205 (2016).
28   Schraiber, J. G., Evans, S. N. & Slatkin, M. Bayesian Inference of Natural Selection from Allele
Frequency Time Series. *Genetics* **203**, 493-511 (2016).
29   Gazave, E. *et al.* Neutral genomic regions refine models of recent rapid human population
growth. *Proc Natl Acad Sci U S A* **111**, 757-762 (2014).
30   Prufer, K. *et al.* The complete genome sequence of a Neanderthal from the Altai Mountains.
Nature **505**, 43-49 (2014).
31   Meyer, M. *et al.* A high-coverage genome sequence from an archaic Denisovan individual.
Science **338**, 222-226 (2012).
32   Fu, Q. *et al.* Genome sequence of a 45,000-year-old modern human from western Siberia.
Nature **514**, 445-449 (2014).
33   Rasmussen, M. *et al.* The genome of a Late Pleistocene human from a Clovis burial site in
western Montana. *Nature* **506**, 225-229 (2014).
34   Rasmussen, M. *et al.* The ancestry and affiliations of Kennewick Man. *Nature* **523**, 455-458
(2015).
35   Lazaridis, I. *et al.* Genomic insights into the origin of farming in the ancient Near East. *Nature*
536, 419-424 (2016).
36   Tishkoff, S. A. & Verrelli, B. C. Patterns of human genetic diversity: implications for human
evolutionary history and disease. *Annu Rev Genomics Hum Genet* **4**, 293-340 (2003).
37   Fu, W. & Akey, J. M. Selection and adaptation in the human genome. *Annu Rev Genomics Hum*
Genet **14**, 467-489 (2013).
38   GTEx Consortium. The Genotype-Tissue Expression (GTEx) pilot analysis: multitissue gene
regulation in humans. *Science* **348**, 648-660 (2015).
39   Welter, D. *et al.* The NHGRI GWAS Catalog, a curated resource of SNP-trait associations. *Nucleic*
Acids Res **42**, D1001-1006 (2014).
40   Guan, W. *et al.* Genome-wide association study of plasma N6 polyunsaturated fatty acids within
the cohorts for heart and aging research in genomic epidemiology consortium. *Circ Cardiovasc*
Genet **7**, 321-331 (2014).
41   Dorajoo, R. *et al.* A genome-wide association study of n-3 and n-6 plasma fatty acids in a
Singaporean Chinese population. *Genes Nutr* **10**, 53 (2015).
42   Shin, S. Y. *et al.* An atlas of genetic influences on human blood metabolites. *Nat Genet* **46**, 543-
550 (2014).

43   Tintle, N. L. *et al.* A genome-wide association study of saturated, mono- and polyunsaturated
red blood cell fatty acids in the Framingham Heart Offspring Study. *Prostaglandins Leukot Essent*
Fatty Acids **94**, 65-72 (2015).
44   Xie, W. *et al.* Genetic variants associated with glycine metabolism and their role in insulin
sensitivity and type 2 diabetes. *Diabetes* **62**, 2141-2150 (2013).
45   Lemaitre, R. N. *et al.* Genetic loci associated with plasma phospholipid n-3 fatty acids: a meta-
analysis of genome-wide association studies from the CHARGE Consortium. *PLoS Genet* **7**,
e1002193 (2011).
46   Gieger, C. *et al.* Genetics meets metabolomics: a genome-wide association study of metabolite
profiles in human serum. *PLoS Genet* **4**, e1000282 (2008).
47   Kettunen, J. *et al.* Genome-wide association study identifies multiple loci influencing human
serum metabolite levels. *Nat Genet* **44**, 269-276 (2012).
48   Mozaffarian, D. *et al.* Genetic loci associated with circulating phospholipid trans fatty acids: a
meta-analysis of genome-wide association studies from the CHARGE Consortium. *Am J Clin Nutr*
101, 398-406 (2015).
49   Eijgelsheim, M. *et al.* Genome-wide association analysis identifies multiple loci related to resting
heart rate. *Hum Mol Genet* **19**, 3885-3894 (2010).
50   den Hoed, M. *et al.* Identification of heart rate-associated loci and their effects on cardiac
conduction and rhythm disorders. *Nat Genet* **45**, 621-631 (2013).
51   Global Lipids Genetics Consortium *et al.* Discovery and refinement of loci associated with lipid
levels. *Nat Genet* **45**, 1274-1283 (2013).
52   Teslovich, T. M. *et al.* Biological, clinical and population relevance of 95 loci for blood lipids.
Nature **466**, 707-713 (2010).
53   Aulchenko, Y. S. *et al.* Loci influencing lipid levels and coronary heart disease risk in 16 European
population cohorts. *Nat Genet* **41**, 47-55 (2009).
54   Zabaneh, D. & Balding, D. J. A genome-wide association study of the metabolic syndrome in
Indian Asian men. *PLoS One* **5**, e11961 (2010).
55   Kathiresan, S. *et al.* Common variants at 30 loci contribute to polygenic dyslipidemia. *Nat Genet*
41, 56-65 (2009).
56   Waterworth, D. M. *et al.* Genetic variants influencing circulating lipid levels and risk of coronary
artery disease. *Arterioscler Thromb Vasc Biol* **30**, 2264-2276 (2010).
57   Sabatti, C. *et al.* Genome-wide association analysis of metabolic traits in a birth cohort from a
founder population. *Nat Genet* **41**, 35-46 (2009).
58   Franke, A. *et al.* Genome-wide meta-analysis increases to 71 the number of confirmed Crohn's
disease susceptibility loci. *Nat Genet* **42**, 1118-1125 (2010).
59   Liu, J. Z. *et al.* Association analyses identify 38 susceptibility loci for inflammatory bowel disease
and highlight shared genetic risk across populations. *Nat Genet* **47**, 979-986 (2015).
60   Jostins, L. *et al.* Host-microbe interactions have shaped the genetic architecture of inflammatory
bowel disease. *Nature* **491**, 119-124 (2012).
61   Richards, M. P., Jacobi, R., Cook, J., Pettitt, P. B. & Stringer, C. B. Isotope evidence for the
intensive use of marine foods by Late Upper Palaeolithic humans. *J Hum Evol* **49**, 390-394
(2005).
62   Richards, M. P. & Hedges, R. E. M. Stable Isotope Evidence for Similarities in the Types of Marine
Foods Used by Late Mesolithic Humans at Sites Along the Atlantic Coast of Europe. *Journal of*
Archaeological Science **26**, 717-722 (1999).
63   Lubell, D., Jackes, M., Schwarcz, H. & Knyf, M. The Mesolithic-Neolithic Transition in
Portugal:Isotopic and Dental Evidence of Diet. *Journal of Archaeological Science* **21**, 201-216
(1994).

64 Richards, M. P. & Mellars, P. A. Stable isotopes and the seasonality of the Oronsay middens.
Antiquity **72**, 178-184 (1998).
65 Bonsall, C. *et al.* Mesolithic and Early Neolithic in the Iron Gates: A Palaeodietary Perspective.
Journal of European Archaeology **5**, 50-92 (1997).
66 Abedi, E. & Sahari, M. A. Long-chain polyunsaturated fatty acid sources and evaluation of their
nutritional and functional properties. *Food Sci Nutr* **2**, 443-463 (2014).
67 Simopoulos, A. P. Evolutionary aspects of diet: the omega-6/omega-3 ratio and the brain. *Mol*
Neurobiol **44**, 203-215 (2011).
68 Mannino, M. A., Thomas, K. D., Leng, M. J., Di Salvo, R. & Richards, M. P. Stuck to the shore?
Investigating prehistoric hunter-gatherer subsistence, mobility and territoriality in a
Mediterranean coastal landscape through isotope analyses on marine mollusc shell carbonates
and human bone collagen. *Quaternary International* **244**, 88-104 (2011).
69 Mannino, M. A. *et al.* Origin and diet of the prehistoric hunter-gatherers on the mediterranean
island of Favignana (Egadi Islands, Sicily). *PLoS One* **7**, e49802 (2012).
70 Lightfoot, E., Boneva, B., Miracle, P. T., Šlaus, M. & O'Connell, T. C. Exploring the Mesolithic and
Neolithic transition in Croatia through isotopic investigations. *Antiquity* **85**, 73-86 (2015).
71 Bocquet-Appel, J.-P., Naji, S., Vander Linden, M. & Kozlowski, J. Understanding the rates of
expansion of the farming system in Europe. *Journal of Archaeological Science* **39**, 531-546
(2012).
72 Rowley-Conwy, P. Westward Ho! The Spread of Agriculture from Central Europe to the Atlantic.
Current Anthropology **52**, S431-S451 (2011).
73 Vigne, J.-D. in *The Neolithic Demographic Transition and its Consequences* (eds J.-P. Bocquet-
Appel & O. Bar-Yosef) 179-205 (Springer Science+Business Media B.V., 2008).
74 Cramp, L. J. *et al.* Immediate replacement of fishing with dairying by the earliest farmers of the
Northeast Atlantic archipelagos. *Proc Biol Sci* **281**, 20132372 (2014).
75 Curry, A. Archaeology: The milk revolution. *Nature* **500**, 20-22 (2013).
76 Salque, M. *et al.* Earliest evidence for cheese making in the sixth millennium BC in northern
Europe. *Nature* **493**, 522-525 (2013).
77 Lidén, K., Eriksson, G., Nordqvist, B., Götherström, A. & Bendixen, E. "The wet and the wild
followed by the dry and the tame" – or did they occur at the same time? Diet in Mesolithic –
Neolithic southern Sweden. *Antiquity* **78**, 23-33 (2004).
78 Rottoli, M. & Castiglioni, E. Prehistory of plant growing and collecting in northern Italy, based on
seed remains from the early Neolithic to the Chalcolithic (c. 5600–2100 cal b.c.). *Vegetation*
History and Archaeobotany **18**, 91-103 (2008).
79 Cho, H. P., Nakamura, M. & Clarke, S. D. Cloning, expression, and fatty acid regulation of the
human delta-5 desaturase. *J Biol Chem* **274**, 37335-37339 (1999).
80 Cho, H. P., Nakamura, M. T. & Clarke, S. D. Cloning, expression, and nutritional regulation of the
mammalian Delta-6 desaturase. *J Biol Chem* **274**, 471-477 (1999).
81 Rapoport, S. I., Igarashi, M. & Gao, F. Quantitative contributions of diet and liver synthesis to
docosahexaenoic acid homeostasis. *Prostaglandins Leukot Essent Fatty Acids* **82**, 273-276
(2010).
82 DeMar, J. C., Jr., Ma, K., Chang, L., Bell, J. M. & Rapoport, S. I. alpha-Linolenic acid does not
contribute appreciably to docosahexaenoic acid within brain phospholipids of adult rats fed a
diet enriched in docosahexaenoic acid. *J Neurochem* **94**, 1063-1076 (2005).
83 DeMar, J. C., Jr. *et al.* Brain elongation of linoleic acid is a negligible source of the arachidonate
in brain phospholipids of adult rats. *Biochim Biophys Acta* **1761**, 1050-1059 (2006).
84 Qin, X. *et al.* Brown but not white adipose cells synthesize omega-3 docosahexaenoic acid in
culture. *Prostaglandins Leukot Essent Fatty Acids* **104**, 19-24 (2016).

85   Okada, Y. *et al.* Genetics of rheumatoid arthritis contributes to biology and drug discovery.
Nature **506**, 376-381 (2014).
86   Zhang, B. *et al.* Large-scale genetic study in East Asians identifies six new loci associated with
colorectal cancer risk. *Nat Genet* **46**, 533-542 (2014).
87   Allentoft, M. E. *et al.* Population genomics of Bronze Age Eurasia. *Nature* **522**, 167-172 (2015).
88   Gamba, C. *et al.* Genome flux and stasis in a five millennium transect of European prehistory.
Nat Commun **5**, 5257 (2014).
89   Olalde, I. *et al.* Derived immune and ancestral pigmentation alleles in a 7,000-year-old
Mesolithic European. *Nature* **507**, 225-228 (2014).
90   Raghavan, M. *et al.* Upper Palaeolithic Siberian genome reveals dual ancestry of Native
Americans. *Nature* **505**, 87-91 (2014).
91   Jones, E. R. *et al.* Upper Palaeolithic genomes reveal deep roots of modern Eurasians. *Nat*
Commun **6**, 8912 (2015).
92   Fu, Q. *et al.* An early modern human from Romania with a recent Neanderthal ancestor. *Nature*
524, 216-219 (2015).
93   Seguin-Orlando, A. *et al.* Paleogenomics. Genomic structure in Europeans dating back at least
36,200 years. *Science* **346**, 1113-1118 (2014).
94   Gunther, T. *et al.* Ancient genomes link early farmers from Atapuerca in Spain to modern-day
Basques. *Proc Natl Acad Sci U S A* **112**, 11917-11922 (2015).
95   Olalde, I. *et al.* A Common Genetic Origin for Early Farmers from Mediterranean Cardial and
Central European LBK Cultures. *Mol Biol Evol* **32**, 3132-3142 (2015).
96   Gallego Llorente, M. *et al.* Ancient Ethiopian genome reveals extensive Eurasian admixture
throughout the African continent. *Science* **350**, 820-822 (2015).
97   Li, J. Z. *et al.* Worldwide human relationships inferred from genome-wide patterns of variation.
Science **319**, 1100-1104 (2008).
98   Novembre, J. *et al.* Genes mirror geography within Europe. *Nature* **456**, 98-101 (2008).
99   The UK10 Consortium *et al.* The UK10K project identifies rare variants in health and disease.
Nature **526**, 82-90 (2015).
100   Browning, B. L. & Browning, S. R. Genotype Imputation with Millions of Reference Samples. *Am J*
Hum Genet **98**, 116-126 (2016).
101   Barrett, J. C., Fry, B., Maller, J. & Daly, M. J. Haploview: analysis and visualization of LD and
haplotype maps. *Bioinformatics* **21**, 263-265 (2005).
102   Paradis, E. pegas: an R package for population genetics with an integrated-modular approach.
Bioinformatics **26**, 419-420 (2010).
103   Dannemann, M., Andres, A. M. & Kelso, J. Introgression of Neandertal- and Denisovan-like
Haplotypes Contributes to Adaptive Variation in Human Toll-like Receptors. *Am J Hum Genet* **98**,
22-33 (2016).
104   Plummer, M., Best, N., Cowles, K. & Vines, K. CODA: convergence diagnosis and output analysis
for MCMC. *R News* **6**, 7-11 (2006).
105   Tajima, F. Statistical method for testing the neutral mutation hypothesis by DNA polymorphism.
Genetics **123**, 585-595 (1989).
106   Szpiech, Z. A. & Hernandez, R. D. selscan: an efficient multithreaded program to perform EHH-
based scans for positive selection. *Mol Biol Evol* **31**, 2824-2827 (2014).
107   Gautier, M. & Vitalis, R. rehh: an R package to detect footprints of selection in genome-wide SNP
data from haplotype structure. *Bioinformatics* **28**, 1176-1177 (2012).
108   Price, A. L. *et al.* Principal components analysis corrects for stratification in genome-wide
association studies. *Nat Genet* **38**, 904-909 (2006).

109   Wang, J. *et al.* Factorbook.org: a Wiki-based database for transcription factor-binding data
generated by the ENCODE consortium. *Nucleic Acids Res* **41**, D171-176 (2013).

Acknowledgements

We thank Montgomery Slatkin and Joshua Schraiber for their help in running their software, David Reich and Iain Mathieson for making their data publicly available, Leonardo Arbiza, Charles Liang, Daniel (Alex) Marburgh, Kumar Kothapalli, Tom Brenna, and all members of the Keinan lab for helpful discussion and comments on the manuscript. This work was supported by the National Institutes of Health (Grants R01HG006849 and R01GM108805 to AK) and the Edward Mallinckrodt, Jr. Foundation (AK).

Author contributions

K.Y. and A.K. conceived and designed the project. K.Y. performed the vast majority of data analysis with help from F.G. and D.W.. K.Y. and A.K. interpreted the results, with contribution from O.B.Y. in interpretation from an anthropological perspective. K.Y. and A.K. wrote the manuscript. All authors read, edited and approved the final version of the manuscript.

Competing interests

The authors declare no competing interests.

1047

Fig. 1. Recent positive selection on the *FADS* locus in Europeans. (**A**) Ancient DNA-based selection test. The overall pattern is consistent with that previously described[11] (Supplementary Fig. S27). Four variants are highlighted: the most significant SNP (purple); the top SNP reported by Mathieson *et al.*[11] (blue); one of the top adaptive SNPs reported in Greenlandic Inuit[9] (orange); the indel reported to be targeted by positive selection in populations with historical plant-based diets[8] (red). (**B**) Haplotype-based selection test (nSL[25]) in modern Europeans. Tests were performed separately for each of the five 1000GP European groups. Only variants with significant values, beyond the 5% genome-wide significance cutoff (magenta dashed lines) are shown with population-specific colors, with exceptions for the four highlighted variants (positions indicated with vertical dashed lines, colored as in A). Consistent patterns were detected with iHS (Supplementary Fig. 4) and in UK10K cohorts (Supplementary Fig. 6). (**C**) Site frequency spectrum (SFS)-based selection test (Fay and Wu's H[26]) in modern Europeans. Tests were performed separately for each of the five 1000GP European groups. Positions for the four highlighted variants are indicated as in (B). The additional significant result observed around the transcription starting site of *FADS3* is beyond the scope of this paper, and detailed

results will be published separately (Ye, Keinan *et al.*). Similar patterns were observed in UK10K cohorts (Supplementary Fig. 8). At the bottom are the representative transcript models for the three *FADS* genes and the four transcription factor binding sites (TFBS) for SREBF1 from generated from ENCODE[109] (blue) and another previous study[10] (red). The 5% significance cutoffs in (B) and (C) are the most extreme ones among the five populations (but p values reported in the main text were based on population-specific empirical distributions). The five 1000 GP European populations are: CEU – Utah Residents (CEPH) with Northern and Western Ancestry; FIN – Finnish in Finland; GRB – British in England and Scotland; IBS – Iberian Population in Spain; TSI – Toscani in Italia.

Fig. 2. Varying selection and frequency patterns between Southern and Northern Europe.
(**A**) South-North frequency gradient for adaptive alleles of four representative SNPs under
different scenarios. AAF refers to adaptive allele frequency. Orange bars represent frequencies
upon admixture, which were directly observed in ancient groups and predicted for extant
populations based on linear mixture of frequencies in ancient groups. Yellow bars represent
frequencies estimated under the null hypothesis. Estimates for ancient groups were not shown
because they are not relevant here. Blue bars represent frequencies estimated under the
alternative hypothesis, whose only constraint is the observed data and therefore the MLEs are
just the observed means. The estimates for ancient groups are the same as their frequencies upon
admixture and are omitted on the plot. The absolute difference between H_0 and H_1 estimates are
indicated above the corresponding bars. Please note that the frequences upon admixture in

WSHG are 0 for rs174594, rs174546 and rs4246215 and no bars were plotted. (**B**) Comparison of aDNA-based selection signals between Southern and Northern Europe. aDNA-based selection tests were performed separately for Southern (TSI and IBS) and Northern (CEU and GBR) Europeans. For each variant, the *p* values from these two tests were compared at a $-\log_{10}$ scale (y axis). SNPs of interest were colored as indicated. (**C**) South-North frequency gradient for the adaptive haplotype in extant populations. The adaptive haplotype is referred to as haplotype D. The two frequency types are just as in (A). The frequency upon admixture for WSHG is 0. In (A) and (C), FIN has only observed values. If values are not shown or not available, signs of "//" are indicated at corresponding positions. Error bars stand for standard errors.

Fig. 3. Temporal frequency pattern and selection signals in pre-Neolithic European hunter-gatherers. (**A**) The frequency of haplotype D over time in four groups of hunter-gatherers. Frequency for each group is plotted as a black point at the median age of samples. The horizontal box surrounding the point represents the medians of lower- and upper-bound estimates of sample ages. The error bar is standard error. Group names are indicated next to their frequencies. (**B**) Allele frequencies for four SNPs. It has similar format as in (A) except that small arbitrary values were added on their x coordinates in order to visualize all SNPs, which were colored as indicated in the legend. The alleles chosen are the ones increasing over time. They are derived alleles for rs174570 and rs2851682, and ancestral alleles for rs174546 and rs174594. (**C**) and (**D**) Posterior distribution on the derived allele frequency path for rs174570 and rs2851682, respectively. The sampled frequencies are indicated with black points, which are the same point estimates as in (B). The median, 25% and 75% quantiles, and 5% and 95% quantiles of the posterior distribution are indicated respectively with black, red and green lines. The posterior distribution on the age of derived allele is shown with a blue line, with values on the right y axis.

1110

Fig. 4. Haplotype network and geographical frequency distribution. (**A**) Haplotype network for 1000G samples (2,157 individuals, excluding admixed American samples), 22 modern Eskimos and 225 aDNAs. Haplotype are defined on the *FADS1-FADS2* LD block. Each pie chart represents one unique haplotype and its size is proportional to $\log_2$(# of chromosomes carrying the haplotype) plus a minimum size so as to visualize the rare haplotypes. The sections in the pie provide the breakdown of the haplotype representation amongst populations with population-specific colors. Sample size for each group (# of haplotypes) is indicated in the legend. Because of the small sample size of some groups, it can be difficult to visualize their proportions in this plot. Detailed haplotype frequencies for each group could be found in Supplementary Table 2. The edges connecting haplotypes are of arbitrary length. Haplotypes for some well-known ancient samples are indicated with their names and arrows pointing at the corresponding pies. The top five haplotypes in modern Europeans, referred to as D, M1, M2, M3, and M4 from the most to least frequent common, are indicated with their names in blue font right next to the haplotypes. (**B**) Frequency of haplotype D in Eurasian ancient DNAs. Each pie represents one sampled group and is placed at the sampling location or nearby with a line pointing at the sampling location. The color of the pie chart border and its associated line indicates the archaeological period of the sample. If multiple samples of different periods were collected at the same geographical location, these samples are ordered vertically with the older samples at the bottom. Hunter-gatherer groups are indicated with black arrows and pastoralist groups with gray arrows, while others are farmers. Geographical locations for some hunter-gatherer groups (e.g. the Věstonice, El Mirón and Villabruna clusters) are only from representative samples. Detailed frequencies could be found in Supplementary Table 3. Pre-N: Pre-Neolithic; N: Neolithic; EN: Early Neolithic; MN: Mid-Neolithic; ChL: Chalcolithic; LN/BA: Late Neolithic/Bronze Age. (**C**) Frequency of haplotype D in present-day global populations. Detailed frequencies are given in Supplementary Table 4. (**D**) Frequency of SNP rs174570 in present-day global populations. All 26 populations from 1000GP and one Eskimo group are included. The color of the pie chart border represents the genetic ancestry. It is noteworthy that there are two samples in America that are actually of African ancestry. Similar global patterns were observed with HGDP samples (Supplementary Figs. 14 and 15).