RNA Stores Tau Reversibly in Complex Coacervates

Xuemei Zhang[1,4], Yanxian Lin[3], Neil A. Eschmann[2], Hongjun Zhou[1,4], Jennifer Rauch[1,4], Israel Hernandez[1,4], Elmer Guzman[1,4], Kenneth S. Kosik[*,1,4], Songi Han[*,2,5]

[1] Molecular, Cell and Developmental Biology, University of California Santa Barbara, Santa Barbara, CA 93106, USA.

[2] Department of Chemistry and Biochemistry, University of California Santa Barbara, Santa Barbara, CA 93106, USA.

[3] Biomolecular Science and Engineering, University of California Santa Barbara, Santa Barbara, CA 93106, USA.

[4] Neuroscience Institute, University of California Santa Barbara, Santa Barbara, CA 93106, USA.

[5] Department of Chemical Engineering, University of California Santa Barbara, Santa Barbara, CA 93106, USA.

*Correspondence to: kenneth.kosik@lifesci.ucsb.edu; songi@chem.ucsb.edu

Competing interests

The authors declare no competing financial interests.

Contributions

K.S.K. and S.H. designed the experiments; X.Z. did the PAR-iCLIP experiments and the *in vitro* RNA-protein binding assays; E.G. sequenced the DNA libraries; H.Z. performed the

bioinformatics analysis; J.R. did the ITC experiments; N.A.E., Y.L. and J.R. did the RNA-tau *in*

vitro droplet formation; N.A.E. did the EPR and DEER experiments; I.H. did the sarkosyl

fractionation; Y.L. and H.Z. contributed to the manuscript preparation; X.Z., K.S.K. and S.H.

wrote the manuscript.

Abstract

Non-membrane-bound organelles that behave like liquid droplets are widespread among

eukaryotic cells. Their dysregulation appears to be a critical step in several neurodegenerative

conditions. Here we report that tau protein, the primary constituent of Alzheimer neurofibrillary

tangles, can form liquid droplets and therefore has the necessary biophysical properties to

undergo liquid-liquid phase separation (LLPS) in cells. Consonant with the factors that induce

LLPS, tau is an intrinsically disordered protein that complexes with RNA to form droplets.

Uniquely, the pool of RNAs to which tau binds in living cells are tRNAs. This phase state of tau

is held in an approximately 1:1 charge balance across the protein and the nucleic acid

constituents, and can thus be maximal at different RNA:tau mass ratios depending on the

biopolymer constituents involved. This feature is characteristic of complex coacervation. We

furthermore show that the LLPS process is directly and sensitively tuned by salt concentration

and temperature, implying it is modulated by both electrostatic interactions between the involved

protein and nucleic acid constituents, as well as net changes in entropy. Despite the high protein

concentration within the complex coacervate phase, tau is locally freely tumbling and capable of

diffusing through the droplet interior. In fact, tau in the condensed phase state does not reveal

any immediate changes in local protein packing, local conformations and local protein dynamics

from that of tau in the dilute solution state. In contrast, the population of aggregation-prone tau

as induced by the complexation with heparin is accompanied by large changes in local tau

conformations and irreversible aggregation. However, prolonged residency within the droplet

state eventually results in the emergence of detectable β-sheet structures according to thioflavin-

T assay. These findings suggest that the droplet state can incubate tau and pre-dispose the protein

toward the formation of insoluble fibrils.

50 Introduction

Inclusions consisting of the tau protein occur in many neurological conditions with

Alzheimer's disease the most prominent among them. Normally, tau is in a dynamic equilibrium

between a microtubule-bound and free state. Under disease conditions tau self-assembles into

fibrils that eventually lead to highly insoluble polymeric inclusions known as neurofibrillary

tangles. The underlying biophysical basis for the transition of tau from a microtubule-associated

protein to an insoluble fibril is unknown. However, a clue comes from the observation that

polyanions, such as heparin, promote tau fibrillization [1]. Although less effectively, RNA can

also induce tau fibrillization [2, 3], and unlike heparin, RNA is present intracellularly, making it

accessible to interact with tau.

Our experiments began with the finding that tau can bind RNA in living cells.

Interestingly tau-RNA binding showed selectivity for tRNAs. This observation along with the

known categorization of tau as intrinsically disordered and its ability to spread from cell to cell in

a manner that resembles prions [4, 5] suggested that tau might share additional properties with

other RNA-binding proteins involved in neurodegeneration. These proteins include FUS [6-8],

TDP-43 [9], C9ORF72 [10, 11], hnRNPA2B1 and hnRNPA1 [12-14], all of which can undergo

liquid-liquid phase-separation (LLPS) from the surrounding aqueous medium into droplets *in*

vitro. These highly protein-dense structures, also known in the literature as complex coacervates

[15, 16], establish a separated liquid phase typically associated with (1) exceptionally high

protein concentration [17], (2) tunability with salt concentration and temperature [18], and (3)

multivalent electrostatic interactions involving polyelectrolytes, including RNA, single-stranded

DNA and intrinsically disordered proteins (IDPs) [19]. A consensus property of a complex

coacervate fluid is low interfacial tension that promotes fusion and coating, and is associated

with low cohesive energy between hydrated polyelectrolyte complexes and weakly bound water

constituents, consistent with high internal fluid dynamics [16, 20]. Complex coacervate

chemistry has been implicated in bio-inspired coating, wet adhesion and engulfment [21-23].

RNA-based coacervates are an important organizing principle for biomolecular condensates in

cell biology [17, 19, 24-26].

Here we show that tau-RNA complexation can lead to complex coacervation. When

multiple tau molecules weakly bind RNA, and overall charge matching is achieved between the

polycation, tau, and polyanion, RNA, tau undergoes reversible condensation and liquid-liquid

phase separation into micrometer-sized droplets. Remarkably, within this liquid phase-separated

state tau maintains high internal segment mobility and a locally compact conformation that

protects the core region of tau known as PHF6(*), as found in dilute solution state, despite the

molecular crowding associated with coacervation. In contrast, this region experiences full

extension that exposes the PHF6(*) region to the solvent and stacks into β-sheets in the presence

of a different polyanion, heparin, that induces irreversible fibrillization of tau [27]. The

spontaneous and reversible droplet formation suggests that tau is held in a low energy-barrier

fluid state between dilute solution and complex coacervate condensate, with the free energy

difference toggled by interactions mediated by ions and hydration water. In fact, a systematic

study of complex coacervation as a function of temperature verified the process to be entropy-

driven that is likely toggled by the release of counterions and/or hydration water that reduces the

net excluded volume of the hydrated biopolymer constituents. However, prolonged residence in

this phase state begins to induce β-sheet formation, suggesting that the highly condensed phase

state of tau can be a precursor to fibril formation.

Results

Tau binds RNA in living cells

RNA binding to tau in living cells was examined by PAR-iCLIP (individual-nucleotide

resolution cross-linking and immunoprecipitation) using the human tau-specific antibody, HJ 8.5.

Human embryonic kidney (HEK) 293T cells expressing wild type full-length human tau (4R2N),

mutant tau (P301L-4R2N) or mutant tau fused to CFP (P301L, 4R1N) were cross-linked, and tau

immunoprecipitated with the tau antibody (Fig. 1a-b and Fig. S2a-c). The tau constructs and tau

mutants used in this study are shown in Fig S1. Wild type human-induced pluripotent stem cell

(hiPSC)-derived neurons (Fig 1c), as well as lines harboring P301L tau and a risk variant for

progressive supranuclear palsy, A152T were cross-linked and immunoprecipitated (data not

shown). A retinoic acid-differentiated neuroblastoma line (SH-SY5Y) that expresses both tau

and the short isoform of MAP2 called MAP2c [28] was also cross-linked and

immunoprecipitated (Fig. S2d). [32]P-labeled RNA bands correspond to immunoprecipitated cross-

linked tau-RNA complexes (Fig. 1a-c, lanes 2; Fig. S2a, lane 2; Fig. S2d, lane 2), with strong

binding of tau to RNA was observed in all these experiments. Tau-RNA complexes were

observed regardless of the tau genotype. PAR-iCLIP experiments with varying RNase

concentrations did not shift the [32]P-labeled band, nor change its intensity (Fig. S2b, lanes 2-3).

The radioactive RNA band ran close to the tau protein itself, in contrast to that of most known

RNA-binding proteins in the literature [29, 30] that run at a range of higher molecular weights.

This radioactive tau band was cut from the membrane, and when sized on an RNA gel, ran in the

range of 30 to 100 nucleotides (data not shown). This confirms the presence of RNA, and

suggests that tau bound predominantly to small RNAs or RNA fragments. Selectivity was further

confirmed by showing that MAP2 did not bind RNA despite its proline rich and microtubule

binding domains that are highly homologous to that of tau (Fig. S2d, lane 3).

Because small RNAs are abundant and may engage in non-specific interactions, we

performed multiple confirmatory controls, including immunoprecipitation in the absence of tau

expression, the use of non-tau antibodies rather than HJ 8.5 to rule out non-specific binding, and

without UV exposure to eliminate the possibility that tau-RNA complexes formed *in vitro* after

cell lysis (Fig. 1a, b lanes 1; Fig. 1c lanes 1, 3; Fig. S2b lanes 1, 4, S2c lanes 3, 4, and Fig. S2d,

lanes 1, 3, and 4).

To identify the types of RNA crosslinked to tau, DNA libraries were prepared from the

immunoprecipitated radiolabeled bands and sequenced. We analyzed the distribution of tau-

bound RNA from the human genome by defining eight regions: exons, introns, lincRNAs,

snRNAs, rRNAs, miRNAs, tRNAs and intergenics. In tau-expressing HEK cells, tRNAs were

overwhelmingly the highest category of RNA crosslinked to tau (Fig. 1d). Endogenous tau in

hiPSC-derived neurons also bound tRNAs; however, background RNA from introns and

intergenics were relatively abundant as well (Fig. 1e). Background RNA sequences are common

in all CLIP studies, particularly for atypical RNA binding proteins [31]. Despite the abundance

of tRNAs in cells, however, background reads from CLIP experiments consistently show a

paucity of tRNAs [29, 32-35]. Correcting for background reads by dividing the percentage of

tau-bound RNA by the percentage of the nucleotides of each category in the genome

demonstrated a selective enrichment of tRNAs that bind to tau in both HEK cells (Fig. 1f) and

hiPSC-derived neurons (Fig. 1g). Furthermore, in all experiments, the non-tau controls

consistently showed relatively few total reads and very few uniquely aligned reads (Fig. S2e-f).

The specific tRNA species crosslinked to tau overlapped extensively between the HEK

and hiPSC-derived neuron samples. Of 625 annotated tRNA loci in the human genome, 462 of

the tRNA genes crosslinked to tau in HEK cells, with 79% of them observed in all four tau CLIP

samples and 94% in at least two samples. In the hiPSC-derived neurons, all 231 tRNA genes

identified were also observed in the HEK cells and 119 of these were verified in at least two tau

CLIP samples. The distribution of the tRNAs cross-linked to tau in HEK and hiPSC-derived

neurons differed markedly from the total endogenous tRNA distribution. This difference

indicated that the tRNAs selected by the CLIP experiment were non-randomly drawn from the

total tRNA pool (Fig. S3a-b). Among the most differentially selected tRNAs by CLIP was

tRNA^{Arg} (Fig. S3c).

PAR-iCLIP identifies the cross-linked sites in the tRNAs (Fig. 2a-b). We found the tRNA

sequences extend from the most 3' nucleotide of the tRNA to the covalently cross-linked site

where the sequencing terminates (Fig. S4). In both HEK cells and hiPSC-derived neurons, the

crosslink site was predominantly located within the anti-codon loop followed by the T loop and

in the D-loop, but the latter two with far smaller frequencies (Fig. 2b). The observed preference

for single-stranded segments of tRNA as crosslink sites to tau, is likely due to the crosslinking

between the nucleobases in this region and aromatic rings on the protein. We conclude that tau

selectively binds RNA, and the predominant RNA specie bound is tRNA. CLIP procedures are

not highly quantitative, and therefore cannot resolve differences in tau-RNA binding among the

tau variants tested, P301L or A152T. Therefore, we are unable to conclude whether tau

mutations affect its RNA binding.

161
Tau-RNA binding affinity and stoichiometry

A gel shift assay using recombinant wild type full-length human tau (4R2N) induced a

shift in unacetylated tRNA^{Lys} (Fig. 3a), yielding a dissociation constant (K_d) for 4R2N tau

binding to tRNA of 460 ± 47 nM (Fig. 3b). The derived Hill coefficient [36] was 2.8, implying

cooperative binding of multiple tau proteins to tRNA. Isothermal titration calorimetry (ITC)

experiments independently confirmed the affinity of tau binding to tRNA to yield K_d = 735 ±

217 nM and 372 ± 9 nM for 4R2N and K18 tau, respectively (Fig. 3c and Fig. S5a). The

dissociation constant for 4R2N binding to a random 43 nucleotide RNA sequence still yielded a

K_d = 832 ± 94 nM with a Hill coefficient of 2.6 according to a gel shift assay (data not shown),

suggesting that tau effectively and non-specifically binds RNA *in vitro*, although there may be

differences in binding affinity.

The gel shift assay showed multiple bands corresponding to different tau:tRNA

stoichiometries corresponding to high molecular weight protein-RNA complexes (Fig. 3d,

similar gel shift with the tau fragment K18 shown in Fig S5b). The fraction of bound tRNA to

4R2N tau (from the low and high bands) was plotted as a function of tau:tRNA molar ratios (Fig.

3e), and compared to theoretical binding saturation curves representing a range of

stoichiometries from 1:1 to 6:1 (see Fig. 3f). The theoretical curves serve the purpose of showing

that multiple tau molecules bind tRNA with increasing tau concentrations. The model is not

intended for data fitting; rather it shows that multiple populations with different tau:tRNA ratios

coexist. However, the data suggests that at lower tau:tRNA molar ratios, tau and RNA

predominantly form protein dimer-RNA (P_2R) complexes, while at higher tau:tRNA molar ratios,

tau and tRNA form larger protein multimer-RNA ($P_{2n}R$) complexes with $2n$ reaching as high as

6. This implies that tRNA is capable of binding multiple tau proteins in a multi-step process.

Interestingly, when the tau:tRNA ratio was decreased by increasing the tRNA concentration

relative to tau, the higher order $P_{2n}R$ complexes dissociated to P_2R, maintaining a tau dimer

bound to RNA in the presence of excess (7.5 fold) RNA (Fig. S5c). Such higher stoichiometric

signatures were not observed in the ITC measurement, which is less sensitive to binding events

associated with small changes in heat. However, ITC titration experiments showed that both

4R2N and K18 tau interacted with tRNA with the stoichiometry of a protein dimer (Fig. S5a n =

0.49 ± 0.03 and Fig. 3c, 0.52 ± 0.04). We conclude that the formation of $P_{2n}R$ complexes with *$2n$*

exceeding 2 must be relying on weak interactions between multiple tau proteins and tRNA. A

two stage model, in which a protein dimer-RNA (P_2R) complex forms first followed by the

formation of a protein multimer-RNA ($P_{2n}R$) complex, is in fact consistent with the model for

RNA binding to the protein AUF1 [37] and hnRNP A1 [38].

Tau phase-separates in the presence of RNA

Mixing of 4R2N or Δtau187 (similar to K18, see Fig. S1) with tRNA (25 kDa), poly(A)

RNA (66~660 kDa) or poly(U) RNA (800~1000 kDa) reliably produced a turbid solution under

a wide range of tau:RNA mass ratios and salt concentrations. According to bright-field

microscopy (Fig. 4a), droplets formed and phase separated from the bulk aqueous phase with a

clearly visible and highly spherical boundary. Tau droplets were capable of merging into a single

droplet with the complete and nearly instantaneous loss of any boundary at the fusion interface,

indicating that the droplets are fluidic with a relatively low interfacial tension (a series of

snapshots capturing the fusion of two droplets are shown in Fig. 4a). Confocal microscopy

images of fluorescence-labeled tau verified that tau was predominantly contained within the

droplet (Fig. 4b). Depending on the specific condition, we were able to observe droplet

formation with total Δtau187 protein concentration ranging between 2-400 μM, as will be

discussed in greater detail next.

Tau-RNA droplets form a complex coacervate phase

The amount of tau-RNA droplet was quantified as % coverage from the microscopy images, as

well as verified against turbidity measurements relying on light scattering at $\lambda = 500$ nm. Droplet

formation follows spontaneously from the mixing of two oppositely charged biopolymers, tau

and RNA, at a given range of pH, salt and protein concentration, as well as temperature (Fig. 5).

This process is fully reversible and reproducible, characteristic of equilibrium states. We

highlight the effect of salt concentration first. At a given total protein concentration and

temperature, the systematic increase in salt concentration decreased the amount of tau-RNA

droplet. Specifically, at room temperature, a total protein concentration of 80 μM and in the

presence of tRNA, the droplets reproducibly disappeared when the salt concentration was

increased to 100 mM or higher (see Fig. 5a). This general trend was observed repeatedly under a

range of experimental conditions, including the use of different RNA types (Fig. 5b, Fig. S6a left

panel). Interestingly, droplets formed in the presence of poly(A) and poly(U) tolerated higher

than 100 mM salt concentration compared to tRNA. The amount of tau-RNA droplet was also

found to be sensitively and reversibly tunable by temperature (Fig. 5d). A sharp dependence of

droplet formation on temperature can be seen in the series of images shown in Fig. 5d, where

droplets dissolve below 22 °C and appear at and above 23 °C. The observation of a clear increase

in the amount of droplet with increasing temperature signifies an entropy-driven process.

Consequently, when the temperature was increased to physiological conditions (37 °C), droplet

formation was observed at protein concentrations as low as 2.5-5 μM, in the presence of salt

concentrations as high as 100 mM and glycerol added as a crowding reagent to mimic the

intracellular environment (Fig. S6b). This observation is illustrated in a series of images

presented in Fig. 5e—droplets are observed at NaCl concentration as high as 100 mM and tau

concentrations as low as 2.5 μM when the temperature is elevated to 37°C, while maintaining the

tau:RNA molar ratios. The quantity of droplets decreased with decreasing total tau concentration,

and disappeared when tau concentration dropped from 2.5 to 1 μM (Fig. 5e, panel iv). Given that

the intracellular concentration of tau is approximately 2-4 μM in neurons [39, 40], conditions

under which droplets were observed correspond to protein concentration, salt concentration and

temperature resembling physiological conditions.

Droplet formation was maximal, as measured by % coverage from the microscopy images,

with Δtau187:tRNA molar ratios of 8:1, Δtau187:poly(A) RNA molar ratios from 33:1 to 330:1

and Δtau187:poly(U) RNA ratios from 267:1 to 333:1 (estimated from the molecular weight of

RNAs, see Fig. S6c). By computing the charge (see Supporting Information) and the mass (as

measured by the molarity) of the RNA and protein components in the droplet samples, we found

that these different molar ratios converge to approximately a 1:1 charge ratios regardless of the

type of RNA used, including poly(A) and tRNA species (Fig. 5c) and poly(U) (Fig. S6a, right),

and that at different total mass concentration of tau and RNA. Droplet formation was also

observed with full length tau 4R2N and RNA at room temperature, but was less robust than with

Δtau187, possibly due to the additional negative charges at the N-terminus of tau 4R2N that

would diminish the electrostatic association between tau and RNA. However, at pH 6 where the

net charge of tau 4R2N is similar to Δtau187, droplets reliably formed (Fig. S6d), albeit with a

significantly lower yield (Fig. S6e)—note the % coverage with tau 4R2N-derived droplets of

order < 3% compared to that with Δtau187-derived droplets of order 5-30%.

The observations that droplet formation is directly, sensitively and reversibly tunable by salt

concentration, tau:RNA charge ratios, as well as temperatures demonstrate that this process is

modulated and toggled by both electrostatic interactions and changes in net entropy, and the

balance between these contributions. Based on these findings, we conclude that tau-RNA droplet

formation follows a complex coacervation mechanism, as initiated through non-specific and

relatively weak complexation of oppositely charged and polyelectrolytes, and driven by the

further association of these polyelectrolyte complexes to form a macroscopic fluid phase.

Tau in condensed droplets assumes solution state properties

Within a condensed complex coacervate fluid, held together by non-specific and weak

electrostatic interactions, we expect the polyelectrolyte constituents to maintain their hydration

layer and remain dynamic [20, 41]. However, this assumption needs experimental verification.

To spectroscopically track tau exclusively from within the condensed phase, we first verified by

confocal fluorescence imaging that Δtau187 was predominantly localized within the droplet (see

Fig. 4b). This allowed us to characterize the droplet-internal protein properties using spin labeled

Δtau187. We carried out continuous wave electron paramagnetic resonance (cw EPR) spectral

line shape analysis of singly spin labeled Δtau187 at a cysteine site 322, Δtau187/322C-SL,

diluted with diamagnetically labeled Δtau187/322C-DL (see Methods) to compare the protein

side chain dynamics and degrees of freedom of the tethered spin label of Δtau187 in dilute

solution state (Fig. 6 red, a-c), in the droplet state associated with poly(A) RNA (blue, a) or

tRNA (green, b), and upon addition of the tau aggregation inducer, heparin (black, c).

Remarkably, the EPR lineshape of Δtau187/322C-SL within the droplet phase was

indistinguishable from that in the dilute solution state (e.g. see red vs blue trace in Fig. 6a). In

contrast, the EPR line shape dramatically broadened within minutes of adding heparin—a highly

effective aggregation inducer of tau [1, 42] (e.g. see red vs black trace in Fig. 6c). This finding

experimentally demonstrates that the condensation of Δtau187 to high protein concentration, as

found within droplets, alone is insufficient to induce dehydration and aggregation of tau, and that

tau retains the protein dynamical properties as found in solution state, despite forming long-range

associations with RNA in a highly concentrated fluid phase. From this, we infer that tau

maintains its protein dynamics and hydration shell as in dilute solution state.

To determine whether tau assumes altered protein conformations in the complex

coacervate phase compared to the dilute solution phase, we compared the intra-protein distances

flanking the PHF6 and PHF6* hexapeptide regions of tau under these conditions. These regions

of the tau sequence pack into the β-sheet core once fibrils are formed of tau [43]. Recently, we

identified that both the PHF6 and PHF6* hexapeptide regions of tau undergo a dramatic opening

from a compact to a fully extended local conformation, well before fibril formation is observed,

and remain extended as they pack into β-sheet structures as part of insoluble fibrils [27]. These

local distance measurements of the regions flanking the PHF6$^{(*)}$ regions offer a powerful tool to

compare the conformational state of tau in dilute solution, droplet and aggregation-induced states.

We prepared a Δtau187 G272C/S285C-SL$_2$ construct that was doubly MTSL-labeled at sites 272

and 285 (see Methods) to access the distances across the PHF6* hexapeptide region by double

electron-electron resonance (DEER) spectroscopy. Surprisingly, the mean distance flanking the

PHF6* region remained unchanged from dilute solution state to when tau was condensed into a

concentrated complex coacervate phase in association with poly(A) RNA and tRNA (compare

blue vs red trace in Fig. 6d, and compare green vs red trace in Fig. 6e). This contrasts with the

effect of heparin on tau that markedly extended the mean distance between the labels (from ~3 to

~4 nm) within minutes of heparin addition as recently reported [27], corresponding to the

extended conformation that the PHF6[(*)] segment adopts when neatly stacked in β-sheets

(compare black vs red trace in Fig. 6f).

Interestingly, we observed a low-level Thioflavin T (ThT) fluorescence under tau-poly(U)

RNA droplet forming conditions that gradually increased over 15 hours (see Δtau187 + poly(U)).

The ThT fluorescence, however, was nearly eliminated at increased salt concentration that

corresponded to conditions under which tau-RNA associations are weakened and droplets are

dispersed. However, even after 15 hours of incubation, the ThT fluorescence intensity from the

tau-RNA droplet samples was significantly less (less than 5%) compared to what was observed

in the presence of the aggregation-inducing heparin, under similar charge ratio and mass

concentration and following a brief incubation time of less than 20 minutes (see Fig. S6f, the

histogram for Δtau187 + poly(U) vs heparin is not to scale, as indicated with a split y axis).

Since ThT fluorescence is commonly used as an assay to detect β-sheet content, we suggest that

droplet formation through association with RNA increases the aggregation propensity of tau *in*

vitro, even when the tau-RNA complexes are held together by reversible and weak interactions

between intact RNA and the hydrated tau constituents.

Exogenous tRNA can induce sarkosyl insolubility of tau

To determine whether tau-RNA complexes have the potential for pathological

interactions *in vivo*, hiPSC-derived neurons with a P301L mutation or wild type were transfected

with 48 μg tRNA per 1.2 million cells. The uptake of tRNA was demonstrated with tRNA[Phe]-

fluorescein (Fig. S7). Cell lysates (input) were prepared in a high salt/high sucrose buffer,

followed by fractionation in a 1% sarkosyl buffer. Transfection in the absence of nucleic acids

(mock) or addition of tRNA to the mock lysate (mock + tRNA) were used as controls. Cells

transfected with tRNA accumulated sarkosyl-insoluble tau as detected with the PHF-1 tau

antibody; whereas cells without added tRNA, or when tRNA was added to the lysis buffer did

not increase the tau population in the insoluble fractions (Fig. 7a). The increase in sarkosyl-

insoluble tau populations occurred in both P301L mutation and wild-type tau cells (Fig. 7b-c)

when infected with tRNA.

Discussion and Conclusion

Many years ago Wolde and Frenkel [44] have proposed that condensed liquid droplets

serve as a "metastable crucible" to lower free energy barriers for crystal nucleation, while the

discovery of the involvement of nucleic acids in coacervation-driven liquid-liquid phase

separation (LLPS) dates back to 1949 by Bungenberg de Jong [21]. Here, we identify tau to

undergo complex coacervation upon association with RNA. We find that tau can bind RNA in

two stages, mediated by (i) strong binding at a tau:RNA molar ratio of 2:1 with nanomolar

dissociation constants and (ii) weak association between tau multimers and RNA to form higher

order complexes. Although tau lacks a recognizable RNA-binding motif, it selectively bound

tRNA as its dominant partner *in vivo*. Furthermore, tau was found to spontaneously phase

separate upon non-specific complexation with RNA into dense protein droplets, whose formation

was reversibly and sensitively tunable by salt concentrations, tau:RNA ratios, as well as

temperature. Crucially, the optimal tau:RNA molar ratios range from 8:1 to 330:1, which varied

among different RNA species. Remarkably, all higher order complexes converge to a common

tau:RNA charge ratio of ~1:1. This finding together with the sensitivity of droplet formation to

salt concentration led us to conclude that tau-RNA droplet formation is driven by complex

coacervation, in which oppositely charged polyelectrolytes associate into extended assemblies

held together by electrostatic interactions. Significantly, EPR spectroscopy verified that tau

remained locally dynamic, and maintained local conformations as found in solution state. This is

consistent with an entropically-driven complex coacervation process between tau and RNA, in

which contribution from attractive tau-RNA interactions and changes in the biopolymer

conformational entropy are small. We rationalize weak tau-RNA interaction to water-mediated

electrostatic attraction between hydrated tau and RNA molecules, that is hence weakened. The

total entropy of the complex coacervated system can be increased through the release of counter-

ions and/or hydration water, provided that the entropy of the biopolymer constituents themselves

remains relatively constant. That the release of hydration water may be one of the potential

drivers of entropically-driven complex coacervation is not contradictory to our statement that

fully hydrated biopolymers constitute the complex coacervate phase. Even sharing the hydration

shell between proteins in a crowded solution alone will increase the net entropy of the total

system's solvent, if the protein concentration within the coacervation is very high and the protein

preserves its dynamics and hydration shell as in dilute solution state. Notably, there are several

reports in the literature on the spontaneous complex coacervation of synthetic polyelectrolyte and

protein (e.g. β-lactoglobulin) constituents to be entropy-driven [6, 18, 24, 45, 46], but no studies

combine macroscopic thermodynamic measurements with studies of local protein dynamics and

conformations. We show that the majority of tau complexed with RNA in the highly condensed

droplet state maintains locally the dynamical conformations of free tau as populated in dilute

solution state. In other words, tau can reversibly switch between the dilute solution and the

dense droplet state involving minimal rearrangement of hydration water and protein

conformations, as tuned by salt concentration, pH and temperature—i.e. physiologically viable

"knobs".

Still, a low-level ThT fluorescence was observed in the tau-RNA droplet samples after

prolonged incubation, but with a significantly reduced amplitude compared to tau incubated with

heparin. Consistent with this finding, tRNA increased sarkosyl insolubility of tau *in vivo*. We

conclude that dynamic and free tau species are stored in a concentrated droplet state that can

spontaneously and reversibly dissolves into solution at increased salt concentration and/or

depressed temperature, while also being predisposed to converting to fibrils. This previously

unrecognized droplet phase state contrasts with that of tau following incubation with the

polyanion, heparin, in which a rapid conformational change occurs in solution state followed by

an irreversible fibril formation. The distinguishing effects between RNA (charge density

~3.0/nm, estimated from the nucleic acid double helix [47], a polyphosphate) and animal-derived

heparin (charge density ~4.0/nm, estimated from a disaccharide crystal structure [48], a

polysulfate) on tau—coacervation versus fibrillization—may be an operational principle of tau-

polyanion association.

Linking the *in vitro* observations of tau-RNA coacervation to *in vivo* structures that share

many of the properties we describe here remains a challenge for the field. RNA granules [49-51],

stress granules [52], germ line granules [53], synaptic vesicles [54], chemical reaction centering

at the origin of life [55] all lack taxonomy, and yet share features that resemble *in vitro* droplets.

These structures are cohesively motile in the absence of any surrounding membrane, contain

IDPs which often contain RNA-binding proteins and serve as an RNA transport vehicle in which

translation is silenced [51, 56, 57]. TDP43, FUS, C9ORF72, hnRNPA2B1 and hnRNPA1 and

TIA-1 [6-14] are all IDPs and RNA binding proteins, but establishing a connection between *in*

vivo systems and the biophysics of droplets has been challenging. Brangwynne et al have

recently employed optogenetic control of Cry2 to trigger the association between IDPs tethered

to Cry2 as a mechanism to induce phase separation involving the IDPs, presumably by

increasing the local protein concentration. Our study not only adds tau to this prominent list of

proteins that undergo LLPS upon association with RNAs, but also establishes a new phase state

of tau in which soluble tau is reversibly stored in concentrated complex coacervate droplets,

readily bioavailable for cellular functions or vulnerable to pathological aggregation depending on

the environmental cues at hand, and offers guidance for the physiologically tunable factors that

may promote LLPS *in vivo* – increased temperature, lower salt concentration, and exceeding a

local threshold protein concentration.

Materials and Methods

Cell culture

A total of eight samples were used for PAR-iCLIP studies. Four samples were human embryonic

kidney (HEK 293T) cells that expressed either 4R2N (residues 1-441) wild type *tau* (one

sample), 4R2N *tau* harboring the P301L mutation (one sample), and 4R1N *tau* harboring the

P301L mutation fused to CFP (two samples). Four samples were neurons derived from human

induced pluripotent stem cells (hiPSCs) obtained by reprogramming dermal fibroblasts with

virally transduced Yamanaka factors [58] that expressed wild type *tau*, two harboring a P301L

tau mutation and one harboring the A152T variant [59, 60]. The hiPSC lines were karyotyped by

Cell Line Genetics. Both wild type and P301L were from normal males (46XY). A152T

displayed a balanced three-way translocation of chromosomes 1, 13 and 7, which most likely

occurred during the reprogramming process (46XY, t(1;13;7) (q31.2;q21;q36.3). Pluripotent

cells were maintained under feeder-free conditions and cultured in BD Matrigel (BD Biosciences)

coated six-well plates and fed with mTSER daily (Stemcell Technologies). Neuroectoderm

differentiation utilized dorsomorphin and SB431542 (Sigma-Aldrich) for a week [61]. When

neural-rosettes were clearly observable, the media was gradually replaced with neuronal

induction media containing Knockout DMEM F12 with N2, and Glutamax 1X supplemented

with laminin (Sigma-Aldrich) and maintained for an additional six days with every other day

feeding. Neuro-rosettes were microdissected, and grown as neurospheres in the neuronal

induction media supplemented with B27 supplement. Neurons were differentiated from

neurospheres by dissociating them into individual cells enzymatically with a 1/1 mixture of 0.25%

trypsin / StemPro accutase (Thermo Fisher), and plated on Poly-L-Ornithin/Laminin coated six

well plates at a density of 200,000 cells per well. Neuron differentiation and maturation were

stimulated by the addition of 10 mM each of NT3, BDNF, and GDNF (Preprotech) to the

neuronal growth media containing Neurobasal, N2 supplement, B27 supplement and Glutamax.

Neurons were fed twice per week by replacing half of the conditioned media with pre-warmed

supplemented fresh media, and underwent maturation for at least five weeks. At this

developmental stage, hiPSC-derived neurons predominantly express the 3R tau isoform.

Neuroblastoma (SH-SY5Y) cells were plated as monolayers in DMEM/F12 medium with

10% FBS at $1x\ 10^6$ per 10 cm dish, and the cells switched to neurobasal medium the next day

and differentiated for seven days using 10 μM retinoic acid.

Antibodies

The tau antibodies used were Tau-5 for probing total tau in western blot (1 mg/ml, Abcam);

human tau-specific HJ 8.5 and mouse tau-specific HJ 9.2 (4.5 mg/ml and 2.8 mg/ml, both HJ

antibodies are gifts from Dave Holtzman, Washington University) for CLIP; PHF-1 (gift from

Peter Davis, Albert Einstein College of Medicine) for sarkosyl-insoluble tau. Other antibodies

used are MAP2 antibody (0.3 mg/ml, Proteintech[TM]); CDK5 rabbit antibody (0.2 mg/ml, Santa

Cruz); GFP rabbit antibody (5 mg/ml, Abcam); β-Actin (Sigma-Aldrich); mouse IgG_1 Alexa 680,

(2 mg/ml, Invitrogen); mouse IgG Alexa 680 (2 mg/ml, Invitrogen); rabbit IgG 800 (1 mg/ml,

Odyssey).

PAR-iCLIP

PAR-iCLIP (PhotoActivatable Ribonucleoside-enhanced Individual-nucleotide resolution UV

Cross-Linking ImmunoPrecipitation and high-throughput sequencing) was applied to detect

specific RNAs bound to tau in living cells [30, 32]. Cells were treated with 4-thiouridine (4SU)

(Sigma-Aldrich) for 1h at a final concentration of 500 μM at 37°C, rinsed with ice-cold 1x PBS

and irradiated one time with 400 mJ/cm^2 of 365 nm UV light on ice. 4-SU can enhance the cross-

linking efficiency especially for proteins in the cytoplasm [29, 30, 32]. The cells were

centrifuged and the pellet stored at -80°C. The major steps of PAR-iCLIP are listed in Fig. S8

and detailed in the supporting information. CLIP experiments require an antibody, which can

effectively immunoprecipitate tau under stringent high salt wash conditions. HJ 8.5 [62] raised

against human tau efficiently depleted tau. With a dissociation constant of 0.3 pM, HJ 8.5 pulled

down tau under the high-stringency purification conditions of CLIP and remained bound to tau

throughout the procedure. Control experiments with GFP or CDK5 antibody for the lysate

expressing those proteins, or with HJ 8.5 antibody for lysates without expressing tau were always

done in parallel to rule out false positive binding caused by the beads. After immunoprecipitation,

the cross-linked RNA was radiolabeled, the protein separated on an SDS-PAGE gel, and

transferred to nitrocellulose membrane and blotted. The RNA-protein complexes from CLIP

experiments (Fig. 1a-c) were cut from the blot, and the RNA extracted followed by reverse

transcription for library preparation [30]. The ^{32}P-labeled RNA band was run on a

polyacrylamide Tris-Borate Urea (TBU) denaturing gel to demonstrate the confirm the size of

the complex.

Library preparation, deep sequencing and bioinformatics analysis of iCLIP

The iCLIP libraries contained an experimental and a random barcode, which allowed

multiplexing and the removal of PCR duplicates. After the barcodes were introduced, sample and

control from one set of experiments were mixed to remove batch-to-batch variation. Libraries

were sequenced on an Ion Torrent Proton (Thermo Fisher). Fastx collapser from FASTX-Toolkit

was used to collapse reads and filter replicates resulting from the PCR based on the random

barcode. Reads were separated into samples by the barcodes at the 5` ends of reads. The reverse

transcription primers sequenced at both ends of the reads were trimmed with Cutadapt [63]. After

trimming the barcodes, reads with 18 bps or more were kept and counted as total unique reads,

and aligned to the human genome (hg19) by Bowtie2 [64]. RseQC [65] was used to evaluate the

quality of sequencing and mapping reads. Alignments with scores equal to or greater than ten

were kept for downstream analysis. Reads from these RNA pools were clustered by their

alignments, and Pyicos tools [66] were used to identify the significant clusters. Clusters with at

least five reads were retained and considered to contain target sites for RNA-tau crosslinking.

Gene models for RefSeq mRNAs, tRNAs, rRNAs were downloaded from the UCSC genome

browser. miRNAs were downloaded from miRBase (release 20) and other categories of RNAs

were download from Ensemble (release 73). Cross-link sites were identified as the termination

site of the sequencing based on the iCLIP protocol [67]. Individual tRNA genes from the UCSC

genome browser were predicted by using tRNAscan-SE v.1.23. The secondary structure for each

tRNA was obtained from GtRNAdb (http://gtrnadb.ucsc.edu/).

Deep sequencing and analyses of small RNA expression in HEK cells and hiPSC-derived

neurons

Small RNAs were extracted from HEK cells and hiPSC-derived neurons using miRNA isolation

kit (mirVana^{TM}). Library preparation was adapted from the Ion RNA-seq v2 (Thermo Fisher)

protocol. cDNAs with size range from 30-100 bp were selected and sequenced. Reads were

aligned to both human genomes and tRNA sequences. When mapping reads to tRNA sequences,

the Bowtie read aligner [68] protocol was used, in which a maximum of two mismatches were

allowed. Reads aligned to tRNAs were counted and analyzed with a custom Perl scripts.

Recombinant tau and tau fragments

Full length recombinant human 4R2N tau, N-terminal truncated, microtubule binding domain

containing, K18 tau (residues 244-372) and Δtau187 (residues 255-441 with a His-tag at the N-

terminus) were used for *in vitro* studies. Methods for expression and purification of all

recombinant tau variants are detailed in the supplementary text. Two variants of Δtau187 were

prepared via site-direct mutagenesis: Δtau187/322C contains a C291S mutation, leaving only one

cysteine at site 322, and Δtau187G272C/S285C contains C291S, C322S, G272C and S285C

mutations, leaving two cysteines at sites 272 and 285.

RNA gel mobility shift assay

The gel shift assay was performed with recombinant full length 4R2N and K18 tau and

chromatographically purified unacetylated yeast tRNA^{Lys} (tRNA Probes) in 100 mM sodium

acetate buffer at pH 7.0. The molar concentration of tRNA^{Lys} was accurately re-measured with

UV spectrophotometry after base-hydrolization to account for the hyperchromic effect from the

secondary and tertiary structure of tRNA [36]. RNA43 was purchased in a kit (Pierce) with the

sequence of 5'-CCUGGUUUUUAAGGAGUGUCGCCAGAGUGCCGCGAAUGAAAAA-3'.

The hydrolyzed tRNA and RNA43 samples were then quantified with UV spectrophotometry at

260 nm using an extinction coefficient of 0.025 $(\mu g/ml)^{-1}cm^{-1}$. For the gel shift assay, protein

was incubated with tRNA at 37℃ for 10 minutes in the presence of 0.5 mM EDTA, 0.5 mM

MgCl$_2$, 2 U SUPERase• In™ RNase Inhibitor (Thermo Fisher), 0.01% IGEPAL CA-630 (Sigma-

Aldrich), and then applied to a TBE 8% Polyacrylamide Gel (Thermo Fisher). After gel

separation, tRNA was stained with SYBR Gold II (Thermo Fisher). For quantitative analysis, the

fraction of free and bound tRNA was quantified in ImageJ2 (National Institute of Health [69]).

Isothermal Titration Calorimetry (ITC) experiments

Full length tau or K18 tau were dialyzed overnight into a specified buffer for ITC (20 mM

ammonium acetate, pH 7). tRNA (from Baker's yeast, Sigma-Aldrich) was re-suspended in the

ITC buffer and concentration determined using a Nanodrop 1000 (Thermo Scientific).

Experiments were run on a Nano ITC (TA Instruments), in which 300 μM tRNA was titrated (5

μl injections) into a 1 ml protein solution of 30 μM tau. Data was analyzed using the

NanoAnalyze v3.6 software (TA Instruments). After subtracting the heat generated by tRNA

titration into an empty buffer, the experimental data was fitted well with an independent binding

model, but not with a cooperative nor to an independent two-site model. Fitting to a cooperative

or to an independent two-site model gave the statistical error being over 100% for the first Ka

value. This could be due to the low enthalpy governing the binding event that is observed in the

ITC data, and could possibly be resolved with increased concentrations of the reactants, here tau

and RNA. However, solubility limitations likely preclude these experiments from being

realistically achievable. Of note, the second Ka value produced by the two site model is very

similar to the values obtained from an independent model, suggesting that the predominant

interaction is indeed the one we described. Experiments were repeated in triplicates and standard

error of the mean reported.

Spin labeling of Δtau187

To achieve labeling with paramagnetic or diamagnetic probes, the protein was dissolved in 6 M

guanidinium hydrochloride and labeled overnight at 4°C using a 20-fold molar excess of the spin

label (1-oxyl-2,2,5,5-tetramethylpyrroline-3-methyl) methanethiosulfonate (MTSL, Toronto

Research Chemicals) or the diamagnetic analog of MTSL (1-Acetoxy-2,2,5,5-tetramethyl-δ-3-

pyrroline-3-methyl) methanethiosulfonate (Toronto Research Chemicals). Excess label was

removed using a PD-10 desalting column (GE Healthcare) equilibrated in a 20 mM ammonium

acetate buffer at pH 7.0. The protein was concentrated using a 3 kDa centrifugal filter (Amicon

UFC800396). The final protein concentration was determined by UV-Vis absorption at 274 nm

using an extinction coefficient of 2.8 $cm^{-1}mM^{-1}$, calculated from extinction coefficient of

Tyrosine [70]. The two variants Δtau187/322C and Δtau187G272C/S285C were spin labeled

with paramagnetic MTSL probes at the one or two cysteine sites, and are respectively referred to

as Δtau187/322C-SL and Δtau187G272C/S285C-SL_2. In order to achieve spin dilution,

Δtau187/322C and Δtau187G272C/S285C were also labeled with the diamagnetic analogue of

MTSL probes, and are referred to as Δtau187/322C/322C-DL and Δtau187G272C/S285C-DL$_2$.

Images shown in Fig. 4a-b were taken using 400 μM tau, 800 μg/ml poly(A) RNA and 30%

glycerol in 20 mM ammonium acetate at pH 7.

Preparation of tau-RNA complex coacervate

Droplets were formed in the 20 mM ammonium acetate buffer with NaCl concentration varying

between 0 and 100 mM and glycerol concentration from 0 to 50% v/v. Unless explicitly

indicated for measurements at various temperatures, all samples were prepared and characterized

at room temperature. Solutions of Δtau187 or full length 4R2N tau was mixed with tRNA

(Baker's yeast, Sigma-Aldrich), poly(A) RNA or poly(U) RNA (Sigma-Aldrich) at varying

protein, RNA, NaCl and glycerol concentrations in a 0.6 ml Eppendorf tube. RNAs were

weighted out as powder and the mass concentration was calculated. The tau:RNA mass ratio, the

total concentration of tau and RNA, as well as the NaCl salt concentration were optimized to

maximize droplet formation, while choosing a total biopolymer density to avoid overlapping of

droplets in the images to simplify the calculation of the droplet coverage (%). Microscopy

images were acquired at 10 minutes after mixing—the droplets form spontaneously and are

expected to be in equilibrium with the dilute supernatant. A concentration of 19% v/v for

glycerol/water was determined to be an optimal concentration to ensure cryoprotection for DEER

measurements carried out at ~80 K, while also ensuring maximal droplet formation at room

temperature (Fig. S6b).

Bright-field microscopy of tau-RNA droplets

Immediately after mixing tau in a 0.6 ml Eppendorf tube with RNA under droplet forming

conditions (established above) and ensuring thorough mixing, 1 µl of this mixture was applied to

a microscope slide that is closed with a cover slide gapped by two layers of double-side sticky

tape to generate a liquid sample region with consistent thickness. The microscope slide was kept

at room temperature for 10 minutes with the cover slide facing down, during which the particles

within the liquid sample region settled down onto the surface of the cover slide. Images were

taken with a 12-bit CCD camera across the entire sample liquid region near the surface of the

cover slide using an inverted compound microscope (Olympus IX70). Before imaging, Köhler

illumination was applied and the focus close to the surface of the cover slide optimized to

enhance the contrast between the dark droplets and the bright background.

Confocal microscopy of tau-RNA droplets

For confocal microscopy, Δtau187/322C was fluorescence-labeled (Δtau187/322C-FL) with

Alexa Fluor 488 C_5 Maleimide (Thermo Fisher) at the same 322 site as Δtau187/322C-SL. 50

µM of Δtau187/322C-FL was mixed with 350 µM Δtau187/322C-SL at a 1:7 molar ratio in order

to prevent saturation. 800 µg/ml of poly(A) RNA was further added to this tau solution, resulting

in droplet formation in the 20 mM ammonium acetate buffer and in the presence of 20 mM NaCl.

10 µl of the mixture was pipetted and put on a microscope slide with a cover slide gapped by

double-sided sticky tape. Confocal images were acquired using a spectral confocal microscope

(Olympus Fluoview 1000).

Droplet quantification from image analysis

To quantify the amount of droplet formed under the given experimental condition of interest, images were taken by a 12-bit CCD camera of an inverted compound microscope (Olympus IX70), and recorded in TIF format. With illumination and focus optimized, droplets settling on the cover slide have lower intensity than their surrounding on the images. An image of the 20 mM ammonium acetate buffer was taken to calculate the average intensity to set as threshold in order to classify different parts of the image into droplets and buffer. For each image, the area of the droplets was divided by the total area of the image, generating a % droplet coverage value on the cover slide. Droplets with eccentricity above 0.9 or equivalent diameter below $1\mu m$ μm were filtered out in order to reduce false reading. The MATLAB code is made available as supplementary files on the internet (https://github.com/yanxianUCSB/DropletAnalysis).

Continuous wave (cw) Electron Paramagnetic Resonance (EPR)

Cw EPR relies on the anisotropy of nitroxide radical's Larmor frequency and hyperfine coupling that makes its lineshape highly sensitive to the local dynamics, orientation and confinement of a nitroxide-based spin label tethered to the protein. Cw EPR measurements were carried out with Δtau187/322C-SL using a X-band spectrometer operating at 9.8 GHz (EMX, Bruker Biospin) and a dielectric cavity (ER 4123D, Bruker Biospin). Samples were prepared by either mixing 200 μM Δtau187/322C-SL with 300 μM unlabeled Δtau187/322C (to generate 40% spin labeled sample) or by using 500 μM Δtau187/322C-SL (100% labeled). Viscogen was added to the sample to achieve either 19% v/v glycerol (for the droplet samples) or 30% v/v sucrose (for the aggregated samples) matching the DEER conditions. tau samples under droplet forming condition were prepared by adding 1.5 mg/ml RNA, and tau samples under aggregation-inducing

conditions prepared by adding 125 μM heparin (11 kDa average MW, Sigma-Aldrich). A sample

of 3.5 μl volume was loaded into a quartz capillary (VitroCom, CV6084) and sealed at one end

with critoseal and the other with beeswax, and then placed in the dielectric cavity for

measurements. Cw EPR spectra were acquired using 2 mW of microwave power, 0.3 gauss

modulation amplitude, 150 gauss sweep width and 25 scans for signal averaging.

Double Electron Electron Resonance (DEER)

DEER measurements were performed on a Q-band pulsed EPR spectrometer operating at 32

GHz (E580, Bruker Biospin) equipped with a QT2 resonator (measurements done by courtesy of

Bruker Biospin). Samples were prepared by mixing 50 μM Δtau187G272C/S285C-SL$_2$ with 500

μM analog-labeled Δtau187G272C/S285C-DL$_2$ at a 1:10 molar ratio to achieve spin-dilution and

avoid artifacts from unwanted inter-protein spin distances. For DEER, tau samples under droplet

forming condition were prepared by adding 1.65 mg/ml RNA and ensuring 19% v/v glycerol

concentration, and tau samples under aggregation-inducing conditions prepared by adding 137.5

μM heparin and ensuring 30% v/v sucrose concentration. 40 μL samples containing 550 μM

concentration of tau were loaded into a quartz capillary (2.4 mm od x 2 mm id) and flash frozen

in liquid nitrogen after 20 minutes of incubation at room temperature and the specific conditions

listed for the data. DEER measurements were conducted using the dead-time free four-pulse

DEER sequence at 80 K, using 22 ns (π/2) and 44 ns (π) observe pulses and a 30 ns (π) pump

pulse. The raw DEER data was processed using Gaussian fitting via DeerAnalysis2013 [71].

Quantification of fibril using Thioflavin T assay

160 μM Δtau187 was mixed with 480 μg/ml poly(U) RNA or 40 μM heparin and incubated at

room temperature in the presence of 4 μM Thioflavin T for over 15 h (Fig. S6f). The

fluorescence intensity at 485 nm was measured using a plate reader (Tecan Infinite 200 Pro).

Transfer RNA transfection

Transfection of RNA in neuron was first validated using tRNA[Phe]-fluorescein transfected in

primary mouse neuron at 14 days *in vitro* using lipofectamine 2000 transfection reagent (Thermo

Fisher) following the manufacturer's protocol (Fig. S7) Cell nuclei was stained with DAPI before

fluorescence microscopic imaging.. For sarkosyl experiments hiPSC neuronal cultures

containing wild type or mutant tau were transfected with 48 μg of bovine liver tRNA (Sigma-

Aldrich) per six well plate. Control cells were transfected in equal conditions in the absence of

nucleic acid (Mock transfection). Mock transfected cells were lysed as described below, with or

without 48 μg tRNA added to the lysis buffer.

Sarkosyl insolube tau isolation and western blotting

Separation of sarkosyl insoluble tau was done as described in the literature [72, 73]. Briefly,

adherent neuronal cell cultures were lysed with an ice-cold high salt/high sucrose Tris HCl buffer

(0.8 M NaCl, 10% Sucrose, 10 mM Tris HCl pH 7.4) containing a 1X Protease Inhibitor Cocktail,

and a 1X Phosphatase Inhibitor Cocktail. Lysis proceeded at 4°C for 30 min before detachment

with a cell scraper, followed by mechanical dissociation using a micropipette. Immediately

afterwards, the lysates were centrifuged at 4°C 3000 x g for 15 min in a microcentrifuge. The

clear supernatants were collected and sampled (Input). Sodium lauroyl sarcosinate (Sigma-

Aldrich) was then added to the supernatants to a final concentration of 1%, followed by a brief

vortexing. Samples were incubated at 4°C with continuous rocking. Samples were then

centrifuged at 4°C for 2 h at 170000 x g in a Beckman Coulter 70.1 Ti rotor. Sarkosyl soluble

supernatants (Sol.) were collected. Sarkosyl insoluble (Ins.) pellets were resuspended in 2X

sample loading buffer (250 mM Tris HCl pH 6.8, 10% Glycerol, 10% SDS, 0.5% bromophenol

blue, 20 mM DTT) and heated to 95°C for 10 min. The previously collected input and the

sarkosyl soluble fractions were diluted with the same sample loading buffer, and heated under

similar conditions. Proteins were separated on a 10% SDS-PAGE, transferred to Nitrocellulose

membranes, and blotted with either PHF-1 or β–actin antibody. Western blot signal was detected

with a LI-COR Odyssey imaging system (LI-COR Biosciences) and quantified in ImageJ2

(National Institute of Health [69].

Data availability

The data that support the findings of this study are available from the corresponding authors

upon request.

Acknowledgments

We are grateful to the tau consortium for financial support to K.S.K. and S.H. Thanks to D. M.

Holtzman, Washington University School of Medicine for the generous gift of HJ 8.5 and HJ 9.2

antibodies and P. Davies, Albert Einstein College of Medicine for PHF-1 antibody. This work

was supported by the NIH grant (#R01AG05605)) to S.H and K.S.K, the 2011 NIH Director New

Innovator Award to S.H and made use of the Material Research Laboratory (MRL) Central

Facilities supported by the National Science Foundation (NSF) through the Materials Research

Science and Engineering Center under Grant DMR 1121053. We acknowledge the use of the

NRI-MCDB Microscopy Facility and the Spectral Laser Scanning Confocal supported by the

Office the Director, National Institutes of Health (NIH) under Award # S10OD010610. The

authors declare no competing financial interests.

Fig. 1: Tau PAR-iCLIP in tau expressing HEK cells and hiPSC-derived neurons. Phosphor

images in the blue frames (a-c) show ^{32}P-labelled RNA crosslinked to tau protein in HEK cells

expressing tau (a-b) and in hiPSC-derived neurons with endogenous tau (c). PAR-iCLIP in HEK

cells expressing wild type tau (a, lane 2) or tau P301L-CFP (b, lane 2; *nota bene*: the fused CFP

retards the migration of tau). (c) PAR-iCLIP of endogenous wild type tau in hiPSC-derived

neurons (lane 2 in c). The antibodies anti-tau HJ 8.5, anti-GFP and anti-CDK5 were used for

protein precipitation. No RNase was added, unless specified. PAR-iCLIP with GFP as a control

(lane 1 in a-b), with CDK5 as a control (lane 3 in c) and no UV control (lane 1 in c). A small

RNA signal was visible in the absence of cross-linking (no UV control) in hiPSC-derived

neurons (lane 1, c), suggesting a small portion of RNA may associate with tau *in vitro* after cell

lysis. The RNA-protein complexes from CLIP marked by rectangles were cut from the blot for

DNA library preparation. Note that two regions of GFP and CDK5 were cut out as sequencing

controls in which the lower MW band corresponds to GFP or CDK5. (d-e) % of tau-CLIP reads

that are mapped to eight human genome regions in HEK cells (d) and hiPSC-derived neurons (e).

(f-g) Enrichment of tRNA in tau-CLIP of HEK cells (f) and hiPSC-derived neurons (g) as

discussed in text.

Figure 2

Fig. 2: Enrichment of tRNA in PAR-iCLIP data. (a) CLIP cDNA reads from tau expressed in hiPSC-derived neurons (neuron CLIP) and from tau expressed in HEK cells (HEK cell CLIP) that aligned to the chr15.tRNA4-ArgTCG tRNA were found in all CLIP samples, and demonstrated a similar pattern of crosslinking. (b) Analysis of crosslinked sites along tRNA secondary structure demonstrates the anticodon preference, where the anticodon (colored red) region is designated as position 1-3 for alignment purpose. The colored illustration of tRNA secondary structure is displayed as an inset, and below the x-axis in 1-D dimension.

Figure 3

a

No protein 20 nM [tau] 2 μM

Bound
Free

b

[tRNA^{Lys}] = 26 nM
K_d of tau = 460 ± 47 nM
Hill coefficient = 2.8

Fraction bound (y-axis, -0.2 to 1)
Tau concentration, μM (x-axis, 0.01 to 10)

c

Time (s)

Corrected Heat Rate (μJ/s)

Normalized Fit (kJ/mol)

4R2N-tRNA
K_d = 735 ± 217 nM
n = 0.49 ± 0.03

tRNA:tau molar ratio

d

No protein 0.16 [tau]:[RNA] 6.4

Bound high
Bound low
Free

e

Fraction (y-axis, 0 to 1)
Tau:tRNA molar ratio (x-axis, 0 to 7)

◆ Free
■ All bound
■ Bound low
■ Bound high

f

1:1 2:1 4:1 6:1

Fraction bound (y-axis, 0 to 1.2)
Tau:tRNA molar ratio (x-axis, 0 to 7)

■ All bound
■ Bound high

Fig. 3: Tau tRNA binding by gel shift assay and ITC. (a) Direct titration experiment shows 4R2N tau induces a mobility shift in tRNA^{Lys}. To determine the K_d value, direct titration experiment was done, which requires trace RNA concentrations to meet the assumption used in the equation that derives the K_d value. The assumption is that the total protein concentration is approximately the free protein concentration at equilibrium, and that protein binding to RNA is negligible. In the direct titration experiments 26 nM tRNA was used and protein concentration spans from 20 nM to 2 µM. (b) The fraction of bound tRNA by 4R2N tau plotted as a function of the monomeric tau concentration and fit to the Hill equation, $y = 1 / [1 + (K_d / x)^n]$. (c) Yeast tRNA was titrated into solutions of 4R2N tau in an ITC experiment. The top panel in (c) show the raw incremental-titration. The area under each peak is integrated and plotted against the tRNA:tau molar ratio and fitted to an independent binding model (the bottom panel), as discussed in methods. (b-c) Standard error of the mean (SEM) is reported from n=3. (d) The stoichiometric binding experiments were performed by varying the tau:RNA molar ratio at a constant 2.6 uM concentration of tRNA, which approximates the saturation concentration (more than five times the K_d of 460 nM). The representative data of three independent experiments is shown. (e) Fraction of bound tRNA from the different bands in (d) is plotted over the molar tau:tRNA ratio. (f) Fraction of bound tau plotted as a function of tau:tRNA molar ratios and compared to the theoretical saturation binding curves (dotted lines) with protein:RNA molar stoichiometries of 1:1 to 6:1. The theoretical curves serve the purpose showing that multiple tau molecules bind tRNA with increasing tau concentrations, while the model is not meant to fit the data, given that multiple populations with different tau:tRNA ratios will coexist.

Figure 4

Fig. 4: Tau and RNA forms droplet *in vitro*. (a) Bright field snapshots of droplets from 400

μM Δtau187 and 800 μg/ml poly(A) showing two droplets seamlessly fusing (highlighted with

red circle). (b) Confocal microscopy image of Δtau187 labeled with Alexa-488 of the droplets

from the same sample as in (a). 50 μM Alexa-488 labeled Δtau187 mixed with 350 μM MTSL

labeled Δtau187 of droplets formed immediately after adding 800 μg/ml poly(A) RNA. Both (a)

and (b) were sampled at room temperature and without added NaCl.

Figure 5

a

| [NaCl] = 0 mM | 30 mM | 50 mM | 100 mM |

b

Coverage [%] vs NaCl [mM]
- ○ poly(A)
- ◇ tRNA

c

Coverage [%] vs Δtau187:RNA charge ratio
- □ tRNA 4 mg/ml
- ○ tRNA 2 mg/ml
- △ poly(A) 2 mg/ml

d

| T = 21 °C | 22 °C | 23 °C | 24 °C |

e

| i | ii | ii | iv |

Fig. 5: Tau-RNA droplets form a complex coacervate phase. (a) Representative bright-field

images of tau-tRNA droplet samples at room temperature with varying [NaCl]. (b) Droplet

coverage (in %) with poly(A) or tRNA in room temperature with varying [NaCl]. Δtau187:RNA

in a-b were maintained at a mass ratio of 7:1 (corresponding to a charge ratio of 1.2:1) and a total

mass concentration of 2 mg/ml. (c) Droplet coverage with poly(A) or tRNA with varying

Δtau187: RNA charge ratios. The total mass concentrations are indicated in the legends. Samples

made by mixing of 80 μM Δtau187 with 240 μg/ml poly(A)/tRNA or 160 μM Δtau187 with 480

μg/ml tRNA gave the highest droplet coverage (%), which correspond to charge ratio of 1.3-1.2

between tau and RNA. Error bars in (b) and (c) represent the standard deviation from n = 3. (d)

Representative bright-field images of tau-RNA droplets as a function of incubation temperature.

To record these images, the temperature was ramped from 19 to 25°C at 1°C/min to acquire

confocal images with bright field illumination. The samples for these images are generated from

100 μM tau mixed with poly(U) at approximately 1:1 charge ratio in the presence of 30 mM

NaCl. (e) Representative bright-field images of tau-RNA droplet samples incubated at 37 °C

under otherwise different sample conditions. The concentration for tau, poly(U) RNA and NaCl

are i) 5 μM, 15 μg/ml, 0 mM; ii) 5 μM, 15 μg/ml, 100 mM; iii) 2.5 μM, 7.5 μg/ml, 100 mM; iv) 1

μM, 3 μg/ml, 100 mM. Arrows highlight some of the droplets in images ii and iii. Through Fig. 5,

scale bars are for 50 μm and all samples were prepared with Δtau187/322C in 20 mM

ammonium acetate at pH 7.0.

Figure 6

Fig. 6: Tau in condensed droplets assumes solution state properties. (a-c) Cw EPR spectra obtained at room temperature of 500 µM Δtau187-SL in droplets formed with 1.5 mg/ml poly(A) RNA (blue in a) and 1.5 mg/ml tRNA (green in b) is unaltered from solution before adding RNA (red in a and b). Cw EPR line shape upon adding 125 µM heparin (black in c) show dramatic line broadening (compared to red in c). DEER of Δtau187-SL$_2$ in droplets formed with 1.5 mg/ml poly(A) RNA (blue in d) and 1.5 mg/ml tRNA (green in e), as well as upon incubation with 137.5 µM heparin (black in f) compared to Δtau187-SL$_2$ in solution (red in d-f).

Figure 7

Fig. 7: tRNA transfection accumulates sarkosyl insoluble tau in hiPSC derived neurons. (a)

Representative western blot of neurons harboring a P301L *tau* mutation transfected with tRNA,

but not cells transfected in the absence of nucleic acids (Mock) present evidence for

accumulation of sarkosyl insoluble tau (Ins.) as seen in the intense band in lane 9 labeled with

PHF-1 antibody. Addition of tRNA to the lysis buffer is insufficient to increase the tau present in

the insoluble fractions (compare Mock + tRNA, lane 6 to tRNA, lane 9). (b-c) Quantification of

PHF-1 tau and β-actin level for both neurons harboring a P301L tau mutation (b) and neurons

expressing wild type tau (c) are shown. Error bar represents standard error of the mean. * $p <$

0.05, *** $p < 0.001$, n=5.

References

1.   Goedert M, Jakes R, Spillantini MG, Hasegawa M, Smith MJ, Crowther RA. Assembly of microtubule-associated protein tau into Alzheimer-like filaments induced by sulphated glycosaminoglycans. Nature. 1996;383(6600):550-3.

2.   Kampers T, Friedhoff P, Biernat J, Mandelkow EM, Mandelkow E. RNA stimulates aggregation of microtubule-associated protein tau into Alzheimer-like paired helical filaments. FEBS Letters. 1996;399(3):344-9. doi: http://dx.doi.org/10.1016/S0014-5793(96)01386-5.

3.   Wang X, Wang D, Zhao J, Qu M, Zhou X, He H, et al. The Proline-Rich Domain and the Microtubule Binding Domain of Protein Tau acting as RNA Binding Domains. Protein & Peptide Letters. 2006;13:679-85.

4.   Sanders David W, Kaufman Sarah K, DeVos Sarah L, Sharma Apurwa M, Mirbaha H, Li A, et al. Distinct Tau Prion Strains Propagate in Cells and Mice and Define Different Tauopathies. Neuron. 2014;82(6):1271-88. doi: http://dx.doi.org/10.1016/j.neuron.2014.04.047.

5.   Clavaguera F, Bolmont T, Crowther RA, Abramowski D, Frank S, Probst A, et al. Transmission and spreading of tauopathy in transgenic mouse brain. Nature Cell Biology. 2009;11(7). doi: 10.1038/ncb1901. PubMed PMID: WOS:000267603100022.

6.   Patel A, Lee Hyun O, Jawerth L, Maharana S, Jahnel M, Hein Marco Y, et al. A Liquid-to-Solid Phase Transition of the ALS Protein FUS Accelerated by Disease Mutation. Cell. 2015;162(5):1066-77. doi: 10.1016/j.cell.2015.07.047.

7.   Burke Kathleen A, Janke Abigail M, Rhine Christy L, Fawzi Nicolas L. Residue-by-Residue View of In Vitro FUS Granules that Bind the C-Terminal Domain of RNA Polymerase II. Molecular Cell. 2015;60(2):231-41. doi: http://dx.doi.org/10.1016/j.molcel.2015.09.006.

8.   Shin Y, Berry J, Pannucci N, Haataja MP, Toettcher JE, Brangwynne CP. Spatiotemporal Control of Intracellular Phase Transitions Using Light-Activated optoDroplets. Cell. 2017;168(1–2):159-71.e14. doi: http://dx.doi.org/10.1016/j.cell.2016.11.054.

9.   Kato M, Han TW, Xie S, Shi K, Du X, Wu LC, et al. Cell-free formation of RNA granules: Low complexity sequence domains form dynamic fibers within hydrogels. Cell. 2012;149(4):753-67. doi: 10.1016/j.cell.2012.04.017.

10.   Lee K-H, Zhang P, Kim Hong J, Mitrea DM, Sarkar M, Freibaum BD, et al. C9orf72 Dipeptide Repeats Impair the Assembly, Dynamics, and Function of Membrane-Less Organelles. Cell. 2016;167(3):774-88.e17. doi: http://dx.doi.org/10.1016/j.cell.2016.10.002.

11.   Lin Y, Mori E, Kato M, Xiang S, Wu L, Kwon I, et al. Toxic PR Poly-Dipeptides Encoded by the C9orf72 Repeat Expansion Target LC Domain Polymers. Cell. 2016;167(3):789-802.e12. doi: http://dx.doi.org/10.1016/j.cell.2016.10.003.

12.   Molliex A, Temirov J, Lee J, Coughlin M, Kanagaraj Anderson P, Kim Hong J, et al. Phase Separation by Low Complexity Domains Promotes Stress Granule Assembly and Drives Pathological Fibrillization. Cell. 2015;163(1):123-33. doi: http://dx.doi.org/10.1016/j.cell.2015.09.015.

13.   Kim HJ, Kim NC, Wang YD, Scarborough EA, Moore J, Diaz Z, et al. Mutations in prion-like domains in hnRNPA2B1 and hnRNPA1 cause multisystem proteinopathy and ALS. Nature. 2013;495(7442):467-73. doi: 10.1038/nature11922.

14.   Lin Y, Protter David SW, Rosen Michael K, Parker R. Formation and Maturation of Phase-Separated Liquid Droplets by RNA-Binding Proteins. Molecular Cell. 2015;60(2):208-19. doi: http://dx.doi.org/10.1016/j.molcel.2015.08.018.

15.   Pak Chi W, Kosno M, Holehouse Alex S, Padrick Shae B, Mittal A, Ali R, et al.
Sequence Determinants of Intracellular Phase Separation by Complex Coacervation of a
Disordered   Protein.   Molecular   Cell.   2016;63(1):72-85.   doi:
http://dx.doi.org/10.1016/j.molcel.2016.05.042.
16.   Jho Y, Yoo HY, Lin Y, Han S, Hwang DS. Molecular and structural basis of low
interfacial energy of complex coacervates in water. Advances in Colloid and Interface Science.
2016. doi: http://dx.doi.org/10.1016/j.cis.2016.07.003.
17.   Keating CD. Aqueous Phase Separation as a Possible Route to Compartmentalization of
Biological Molecules. Accounts of Chemical Research. 2012;45(12):2114-24. doi:
10.1021/ar200294y.
18.   Priftis D, Megley K, Laugel N, Tirrell M. Complex coacervation of poly(ethylene-
imine)/polypeptide aqueous solutions: Thermodynamic and rheological characterization. Journal
of   Colloid   and   Interface   Science.   2013;398:39-50.   doi:
http://dx.doi.org/10.1016/j.jcis.2013.01.055.
19.   Nott Timothy J, Petsalaki E, Farber P, Jervis D, Fussner E, Plochowietz A, et al. Phase
Transition of a Disordered Nuage Protein Generates Environmentally Responsive Membraneless
Organelles. Molecular Cell. 57(5):936-47. doi: 10.1016/j.molcel.2015.01.013.
20.   Huang K-Y, Yoo HY, Jho Y, Han S, Hwang DS. Bicontinuous Fluid Structure with Low
Cohesive Energy: Molecular Basis for Exceptionally Low Interfacial Tension of Complex
Coacervate Fluids. ACS Nano. 2016;10(5):5051-62. doi: 10.1021/acsnano.5b07787.
21.   Bungenberg de Jong HG. Colloid Science. In: Kruyt HR, editor. II. Amsterdam: Elsevier
Publishing Co. Inc; 1949. p. 335-432.
22.   Hwang DS, Zeng H, Srivastava A, Krogstad DV, Tirrell M, Israelachvili JN, et al.
Viscosity and interfacial properties in a mussel-inspired adhesive coacervate. Soft Matter.
2010;6(14):3232-6. doi: 10.1039/C002632H.
23.   Overbeek JT, Voorn MJ. Phase separation in polyelectrolyte solutions; theory of
complex coacervation. J Cell Physiol Suppl. 1957;49(Suppl 1):7-22; discussion, -6. PubMed
PMID: 13449108.
24.   Aumiller WM, Pir-Cakmak F, Davis BW, Keating CD. RNA-based coacervates as a
model for membraneless organelles: Formation, properties, and interfacial liposome assembly.
Langmuir. 2016. doi: 10.1021/acs.langmuir.6b02499.
25.   Shorter J. Membraneless organelles: Phasing in and out. Nat Chem. 2016;8(6):528-30.
doi: 10.1038/nchem.2534.
26.   Banani SF, Lee HO, Hyman AA, Rosen MK. Biomolecular condensates: organizers of
cellular biochemistry. Nat Rev Mol Cell Biol. 2017;advance online publication. doi:
10.1038/nrm.2017.7
http://www.nature.com/nrm/journal/vaop/ncurrent/abs/nrm.2017.7.html#supplementary-
information.
27.   Eschmann NA, Georgieva ER, Ganguly P, Borbat PP, Rappaport MD, Akdogan Y, et al.
Signature of an aggregation-prone conformation of tau. Scientific Reports. 2017;7:44739. doi:
10.1038/srep44739
https://www.nature.com/articles/srep44739#supplementary-information.
28.   Dehmelt L, Halpain S. The MAP2/Tau family of microtubule-associated proteins.
Genome Biol. 2005;6(1):204.
29.   Tollervey JR, Curk T, Rogelj B, Briese M, Cereda M, Kayikci M, et al. Characterizing
the RNA targets and position-dependent splicing regulation by TDP-43. Nat Neurosci.

2011;14(4):452-8.   doi:
http://www.nature.com/neuro/journal/v14/n4/abs/nn.2778.html#supplementary-information.
30.   Huppertz I, Attig J, D'Ambrogio A, Easton LE, Sibley CR, Sugimoto Y, et al. iCLIP:
Protein–RNA interactions at nucleotide resolution. Methods. 2014;65(3):274-87. doi:
http://dx.doi.org/10.1016/j.ymeth.2013.10.011.
31.   Friedersdorf M, Keene J. Advancing the functional utility of PAR-CLIP by quantifying
background binding to mRNAs and lncRNAs. Genome Biology. 2014;15(1):R2.
32.   Hafner M, Landthaler M, Burger L, Khorshid M, Hausser J, Berninger P, et al.
Transcriptome-wide identification of RNA-binding protein and microRNA target sites by PAR-
CLIP. Cell. 2010;141(1):129-41. doi: 10.1016/j.cell.2010.03.009. PubMed PMID: 20371350;
PubMed Central PMCID: PMC2861495.
33.   Ascano M, Mukherjee N, Bandaru P, Miller JB, Nusbaum JD, Corcoran DL, et al. FMRP
targets distinct mRNA sequence elements to regulate protein expression. Nature.
2012;492(7429):382-6.   doi:
http://www.nature.com/nature/journal/v492/n7429/abs/nature11737.html#supplementary-
information.
34.   Hoell JI, Larsson E, Runge S, Nusbaum JD, Duggimpudi S, Farazi TA, et al. RNA targets
of wild-type and mutant FET family proteins. Nat Struct Mol Biol. 2011;18(12):1428-31. doi:
http://www.nature.com/nsmb/journal/v18/n12/abs/nsmb.2163.html#supplementary-information.
35.   Kenny PJ, Zhou H, Kim M, Skariah G, Khetani RS, Drnevich J, et al. MOV10 and
FMRP regulate AGO2 association with microRNA recognition elements. Cell reports.
2014;9(5):1729-41. doi: 10.1016/j.celrep.2014.10.054. PubMed PMID: PMC4268400.
36.   Ryder SP, Recht MI, Williamson JR. Quantitative analysis of protein-RNA interactions
by gel mobility shift. Methods in molecular biology (Clifton, NJ). 2008;488:99-115. doi:
10.1007/978-1-60327-475-3_7. PubMed PMID: PMC2928675.
37.   Zucconi BE, Ballin JD, Brewer BY, Ross CR, Huang J, Toth EA, et al. Alternatively
Expressed Domains of AU-rich Element RNA-binding Protein 1 (AUF1) Regulate RNA-binding
Affinity, RNA-induced Protein Oligomerization, and the Local Conformation of Bound RNA
Ligands. Journal of Biological Chemistry. 2010;285(50):39127-39.
38.   Okunola HL, Krainer AR. Cooperative-Binding and Splicing-Repressive Properties of
hnRNP A1. Molecular and Cellular Biology. 2009;29(20):5620-31.
39.   Gamblin TC, Berry RW, Binder LI. Modeling Tau Polymerization in Vitro: A Review
and Synthesis†. Biochemistry. 2003;42(51):15009-17. doi: 10.1021/bi035722s.
40.   Reynolds MR, Berry RW, Binder LI. Site-Specific Nitration Differentially Influences τ
Assembly in Vitro. Biochemistry. 2005;44(42):13997-4009. doi: 10.1021/bi051028w.
41.   Ortony JH, Hwang DS, Franck JM, Waite JH, Han S. Asymmetric Collapse in
Biomimetic Complex Coacervates Revealed by Local Polymer and Water Dynamics.
Biomacromolecules. 2013;14(5):1395-402. doi: 10.1021/bm4000579.
42.   Pérez M, Valpuesta JM, Medina M, Montejo de Garcini E, Avila J. Polymerization of τ
into Filaments in the Presence of Heparin: The Minimal Sequence Required for τ - τ Interaction.
Journal of Neurochemistry. 1996;67(3):1183-90. doi: 10.1046/j.1471-4159.1996.67031183.x.
43.   von Bergen M, Barghorn S, Biernat J, Mandelkow E-M, Mandelkow E. Tau aggregation
is driven by a transition from random coil to beta sheet structure. Biochimica et Biophysica Acta
(BBA) - Molecular Basis of Disease. 2005;1739(2–3):158-66. doi:
http://dx.doi.org/10.1016/j.bbadis.2004.09.010.

44.   Wolde PRt, Frenkel D. Enhancement of Protein Crystal Nucleation by Critical Density
Fluctuations. Science. 1997;277(5334):1975.
45.   Aberkane L, Jasniewski J, Gaiani C, Scher J, Sanchez C. Thermodynamic
Characterization of Acacia Gum−β-Lactoglobulin Complex Coacervation. Langmuir.
2010;26(15):12523-33. doi: 10.1021/la100705d.
46.   Xu AY, Melton LD, Jameson GB, Williams MAK, McGillivray DJ. Structural
mechanism of complex assemblies: characterisation of beta-lactoglobulin and pectin interactions.
Soft Matter. 2015;11(34):6790-9. doi: 10.1039/C5SM01378J.
47.   Blackburn GM. Nucleic Acids in Chemistry and Biology: Royal Society of Chemistry;
2006.
48.   Carlström D. THE CRYSTAL STRUCTURE OF α-CHITIN (POLY-N-ACETYL-D-
GLUCOSAMINE). The Journal of Biophysical and Biochemical Cytology. 1957;3(5):669-83.
PubMed PMID: PMC2224123.
49.   Knowles RB, Sabry JH, Martone ME, Deerinck TJ, Ellisman MH, Bassell GJ, et al.
Translocation of RNA granules in living neurons. Journal of Neuroscience. 1996;16(24):7812-20.
50.   Rook MS, Lu M, Kosik KS. CaMKIIalpha 3' untranslated region-directed mRNA
translocation in living neurons: visualization by GFP linkage. J Neurosci. 2000;20(17):6385-93.
PubMed PMID: 10964944.
51.   Krichevsky AM, Kosik KS. Neuronal RNA granules: a link between RNA localization
and stimulation-dependent translation. Neuron. 2001;32. doi: 10.1016/S0896-6273(01)00508-6.
52.   Wheeler JR, Matheny T, Jain S, Abrisch R, Parker R. Distinct stages in stress granule
assembly and disassembly. eLife. 2016;5:e18413. doi: 10.7554/eLife.18413.
53.   Smith J, Calidas D, Schmidt H, Lu T, Rasoloson D, Seydoux G. Spatial patterning of P
granules by RNA-induced phase separation of the intrinsically-disordered protein MEG-3. eLife.
2016;5:e21337. doi: 10.7554/eLife.21337. PubMed PMID: PMC5262379.
54.   Milovanovic D, De Camilli P. Synaptic Vesicle Clusters at Synapses: A Distinct Liquid
Phase? Neuron. 93(5):995-1002. doi: 10.1016/j.neuron.2017.02.013.
55.   Zwicker D, Seyboldt R, Weber CA, Hyman AA, Julicher F. Growth and division of
active droplets provides a model for protocells. Nat Phys. 2017;13(4):408-13. doi:
10.1038/nphys3984
http://www.nature.com/nphys/journal/v13/n4/abs/nphys3984.html#supplementary-information.
56.   Kedersha N, Panas MD, Achorn CA, Lyons S, Tisdale S, Hickman T, et al. G3BP–
Caprin1−USP10 complexes mediate stress granule condensation and associate with 40S subunits.
The Journal of Cell Biology. 2016;212(7):845-60. doi: 10.1083/jcb.201508028. PubMed PMID:
PMC4810302.
57.   Parker R, Sheth U. P bodies and the control of mRNA translation and degradation. Mol
Cell. 2007;25. doi: 10.1016/j.molcel.2007.02.011.
58.   Takahashi K, Yamanaka S. Induction of Pluripotent Stem Cells from Mouse Embryonic
and Adult Fibroblast Cultures by Defined Factors. Cell. 126(4):663-76. doi:
10.1016/j.cell.2006.07.024.
59.   Coppola G, Chinnathambi S, Lee JJ, Dombroski BA, Baker MC, Soto-Ortolaza AI, et al.
Evidence for a role of the rare p.A152T variant in MAPT in increasing the risk for FTD-
spectrum and Alzheimer's diseases. Human Molecular Genetics. 2012;21(15):3500-12.
60.   Fong H, Wang C, Knoferle J, Walker D, Balestra Maureen E, Tong Leslie M, et al.
Genetic Correction of Tauopathy Phenotypes in Neurons Derived from Human Induced

Pluripotent Stem Cells. Stem Cell Reports. 2013;1(3):226-34. doi: 10.1016/j.stemcr.2013.08.001. PubMed PMID: PMC3849235.

61.   Mak SK, Huang YA, Iranmanesh S, Vangipuram M, Sundararajan R, Nguyen L, et al. Small molecules greatly improve conversion of human-induced pluripotent stem cells to the neuronal lineage. Stem Cells Int. 2012;2012:140427. doi: 10.1155/2012/140427. PubMed PMID: 22567022; PubMed Central PMCID: PMCPMC3339118.

62.   Yanamandra K, Kfoury N, Jiang H, Mahan TE, Ma S, Maloney SE, et al. Anti-tau antibodies that block tau aggregate seeding in vitro markedly decrease pathology and improve cognition in vivo. Neuron. 2013;80(2):402-14. doi: 10.1016/j.neuron.2013.07.046. PubMed PMID: PMC3924573.

63.   Martin M. Cutadapt removes adapter sequences from high-throughput sequencing reads. EMBnet journal. 2011;17(1):10-2.

64.   Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nat Meth. 2012;9(4):357-9. doi: 10.1038/nmeth.1923 http://www.nature.com/nmeth/journal/v9/n4/abs/nmeth.1923.html#supplementary-information.

65.   Wang L, Wang S, Li W. RSeQC: quality control of RNA-seq experiments. Bioinformatics. 2012;28(16):2184-5. doi: 10.1093/bioinformatics/bts356.

66.   Althammer S, González-Vallinas J, Ballaré C, Beato M, Eyras E. Pyicos: a versatile toolkit for the analysis of high-throughput sequencing data. Bioinformatics. 2011;27(24):3333-40.

67.   Konig J, Zarnack K, Rot G, Curk T, Kayikci M, Zupan B, et al. iCLIP reveals the function of hnRNP particles in splicing at individual nucleotide resolution. Nat Struct Mol Biol. 2010;17(7):909-15.   doi: http://www.nature.com/nsmb/journal/v17/n7/abs/nsmb.1838.html#supplementary-information.

68.   Langmead B, Trapnell C, Pop M, Salzberg S. Ultrafast and memory-efficient alignment of short DNA sequences to the human genome. Genome Biology. 2009;10(3):R25.

69.   Schneider CA, Rasband WS, Eliceiri KW. NIH Image to ImageJ: 25 years of image analysis. Nat Meth. 2012;9(7):671-5.

70.   Fasman GD, editor. Handbook of Biochemistry and Molecular Biology. 3rd ed. Cleveland, OH.: CRC Press; 1976.

71.   Jeschke G, Chechik V, Ionita P, Godt A, Zimmermann H, Banham J, et al. DeerAnalysis2006—a comprehensive software package for analyzing pulsed ELDOR data. Applied Magnetic Resonance. 2006;30(3):473-98. doi: 10.1007/BF03166213.

72.   DeTure M, Ko L-w, Easson C, Yen S-H. tau Assembly in Inducible Transfectants Expressing Wild-Type or FTDP-17 tau. The American Journal of Pathology. 2002;161(5):1711-22. PubMed PMID: PMC1850799.

73.   Carrettiero DC, Hernandez I, Neveu P, Papagiannakopoulos T, Kosik KS. The Co-chaperone BAG2 Sweeps PHF Insoluble Tau from the Microtubule. Journal of Neuroscience. 2009;29(7):2151-61. doi: 10.1523/JNEUROSCI.4660-08.2009. PubMed PMID: PMC2768429.