Community selection increases biodiversity effects

Sofia J. van Moorsel[1*], Terhi Hahl[1*], Cameron Wagg[1], Gerlinde B. De Deyn[2], Dan F.B.

Flynn[1], Varuna Yadav[1], Debra Zuppinger-Dingley[1] and Bernhard Schmid[1†]

[1] Department of Evolutionary Biology and Environmental Studies, University of Zürich,

Winterthurerstrasse 190, 8057 Zürich, Switzerland

[2] Department of Environmental Sciences, Wageningen University, Droevendaalsesteeg 4,

6708 PB Wageningen, the Netherlands

* Shared first authorship

[†] Corresponding author: Bernhard Schmid

Email: bernhard.schmid@ieu.uzh.ch

Emails of other authors (in order):

sofia.vanmoorsel@ieu.uzh.ch

terhi.hahl@ieu.uzh.ch

cameron.wagg@ieu.uzh.ch

gerlinde.dedeyn@wur.nl

danfbflynn@gmail.com

varuna.yadav@ieu.uzh.ch

debra.zuppinger@ieu.uzh.ch

Type of article: Letter

Species extinctions from local communities can negatively affect ecosystem functioning[1]. Statistical and ecological mechanisms underlying these impacts are well studied[2-4] but the role of evolutionary mechanisms has rarely been addressed[5,6]. In particular, it is not known to what extent local populations are co-adapted, not only to their environment but also to each other, because such a co-adaption would suggest that species extinctions can result in a larger than expected decline in ecosystem functioning. We used a long-term field biodiversity experiment with 52 plant species to determine if natural selection at the level of entire communities intensified the effects of biodiversity on ecosystem functioning. We re-assembled communities with 8-year co-selection histories adjacent to communities with identical species composition but no history of co-selection on native soil from the experiment and on novel soil. Over four years, selected plant communities were more productive and expressed stronger biodiversity effects than naïve communities. Novel soil increased productivity initially but not in the longer term. Our findings suggest that plant community selection can lead to increased community-level functioning, regardless of the soil in which they were assembled. As a consequence, it may take many years to reconstruct well-functioning communities if one reassembles them with populations lacking a common selection history.

A large number of experiments have shown positive effects of species richness on ecosystem functioning, in particular plant biomass production[1,3,4,7-10]. These effects have been explained by statistical sampling, increasing the chances of including productive species in diverse communities[2,3], or by ecological complementary of species, which allows plant mixtures to extract resources more efficiently from the environment[2,4]. Furthermore, species-rich communities may host more diverse communities of enemies thereby diluting specialist pathogens commonly found in species-poor communities[11].

Functional complementarity between co-occurring species increases over time[8-10],

likely through selection for niche differentiation[6]. However, the possibility that evolutionary

processes shape the diversity–productivity relationship has rarely been tested, and only at the

level of individual species[6] rather than at the level of entire communities. Community

selection may be defined as genetically based changes among species constituting the

community, which alter species performances and interactions[12]. Such changes may occur via

genetic recombination, mutations[13], or sorting-out from standing genetic variation through

differential mortality and growth of individuals[14]. Just as Darwinian natural selection can lead

to changes in gene frequencies in populations within species, selection at the level of

communities can lead to correlated changes in gene frequencies in multiple species[12] in

response to one another or to co-varying environmental conditions. However, empirical

evidence for community selection so far has only been found in planktonic and bacterial

communities[15,16].

Recent evidence suggests that selection of particular genotypes from the total genetic

pool of a species may affect ecosystem functioning in field experiments[6,17-22]. We propose

that selection at the level of entire communities is even more likely to affect ecosystem

functions, because of non-random niche or trait changes in response to other phenotypes in

the community that result in filling niche space more fully or evenly. Hence, we expect that a

community of species with a shared selection history will show a stronger biodiversity effect

in comparison with communities assembled from species without such a shared community

selection history. Our specific expectation applies to plant communities, but their interactions

with other (living) components of the ecosystem may also play a role. For example, plant–

soil feedback experiments have shown that soil biota change in response to different plant

species, which can in turn modify the composition and productivity of plant communities[23,24].

Here we compare the diversity–productivity relationships of plant communities that

have been intact communities for eight years in a long-term grassland biodiversity

experiment (Jena Experiment[25]), and hence likely have evolved together as a community,

with naïve plant communities without such a shared selection history. We used experimental

plant communities of 1, 2, 4 or 8 species, and with 12 unique species compositions within

each species richness level. To connect above- and belowground interactions we factorially

combined two community components under selection, plants and soils. We grew selected

and naïve plant communities for four years in native soil with a common selection history

with the selected plant communities or in novel soil (sterilized soil or neutral soil) without a

common selection history with either selected or naïve plant communities (see Methods and

Extended Data Fig. 1). In 2010, seeds for selected communities were produced from plants

collected in Jena in an experimental garden in Zurich. Seeds for naïve communities were

purchased from the original supplier of seeds for the Jena Experiment. Seedlings for selected

and naïve communities were propagated in the glasshouse in Zurich and transplanted into the

plots of the Jena Experiment in early spring 2011. Community-level plant productivity was

measured each spring from 2012–2015 by collecting species-specific aboveground biomass

(see Methods). The shape of the diversity–productivity relationship was quantified as a

function of the logarithm of species richness using a second-degree polynomial.

Overall, for each doubling of species richness, community aboveground biomass

increased by 100 $g \cdot m^{-2} \cdot y^{-1}$. However, between monocultures and 2- to 4-species mixtures

this increase in productivity was steeper for selected plant communities than for naïve plant

communities (Fig. 1; $F_{1,\,132.3} = 7.219$, $P = 0.008$ for the interaction between community

selection treatment and the quadratic term of species richness; Extended Data Table 1).

Selected plant communities were also more productive on average than naïve communities

($F_{1,132.4} = 22.12$, $P < 0.001$). Furthermore, productivity increased with time in selected

communities but decreased in naïve ones ($F_{1,135.1} = 5.461$, $P = 0.021$). This was related to

increased performance of 2- and 4-species mixtures over time in selected communities and

the deterioration of monocultures in naïve communities (Fig. 1 and Extended Data Fig. 2).

We plotted the diversity–productivity relationships of the first four years (2003–2006, i.e.

before selection) of the Jena Experiment (Extended Data Fig. 3), which clearly resembled

such relationships in naïve communities but not in selected communities from 2012–2015

(Fig. 1).

Community productivity initially increased in novel soil and at high diversity (Fig. 2),

reflected in an overall main effect of soil treatment ($F_{1,89.6} = 23.91$, $P < 0.001$), a significant

soil treatment by year interaction ($F_{1,90.8} = 8.591$, $P = 0.004$) and a significant three-way

interaction of the latter with species richness ($F_{1,90.5} = 8.287$, $P = 0.005$). In part this might

have been caused by a nutrient flush from gamma-sterilization of the soil[26], which affected all

novel but only half of the native soil treatments (see Methods). However, the use of native vs.

novel soils did not alter the differences in biodiversity effects between selected and naïve

plant communities ($F_{1,131.2} = 0.001$, $P = 0.973$ and $F_{1,132.0} = 0.566$, $P = 0.453$ for the three-way

interactions of soil treatment with community selection treatment and log-linear and

quadratic species richness terms, respectively).

Our results show that eight years of selection in plant communities in a biodiversity

experiment can increase the slope of diversity–productivity relationships, suggesting

community selection may at least in part explain why biodiversity effects commonly increase

over time in such experiments[8-10]. Our community selection treatment increased the

biodiversity effect on community productivity especially from monocultures to 2- and 4-

species mixtures. In a previous study, we showed that eight years of selection led to increased

biodiversity effects in 2-species mixtures compared with monocultures due to character

displacement between individual species outside their community context[6]. It is conceivable

that such processes become less effective or take more time to evolve when communities of

more than 2–4 species are considered. In addition, selection for increased growth in mixture

may be compromised by reduced pathogen defence[27]. This may explain why 8-species

mixtures were initially much more productive in novel than in native soil (Fig. 2 and

Extended Data Fig. 2). Within-species variance in specific leaf area (SLA) decreased in

selected and increased in naïve communities with increasing species richness (Extended Data

Fig. 4, $F_{1,69.2} = 4.87$, $P = 0.031$). The narrowing of within-species variance with increasing

diversity in selected communities may thus underlie character displacement between species.

In contrast, species in monocultures may have been selected for niche expansion resulting in

increased within-species variance. Species in naïve communities had not yet responded to

different diversity treatments with a similar adjustment in within-species variance in the four

years of this study; their higher variance at high diversity may stem from a more

heterogeneous biotic environment.

The focus of our study was community selection; however, our results are consistent

with the few studies concerning species selection in biodiversity experiments[6,21,22]. Our

community selection treatments led to an increased species-level productivity for most of our

species (Fig. 3). Eight years of selection increased biodiversity effects in selected

communities, naïve communities however did not catch up in the four years of growth of our

study, giving us some insights into the duration of the selection process: even if community

selection may occur mainly through sorting out from initial standing variation within plant

species, it may not occur immediately during seedling establishment and growth. The

inability of naïve communities to adapt could most clearly be seen in the deterioration of

monocultures, potentially due to pathogen accumulation[11], both in native and with some

delay in novel soil (Extended Data Fig. 2).

The generally lower productivity in native soil, and with time in novel soil, may have

occurred through nutrient depletion or stronger plant-pathogenic effects compared with plant-

mutualistic soil organisms[28]. Previous studies in the context of biodiversity–ecosystem

functioning research reported negative plant–soil feedbacks on home soils[24,29], possibly

through the accumulation of specialized pathogenic microbes[11]. The relatively rapid approach

of the slope and shape of the diversity–productivity relationship in novel soil to the

relationship in native soil (Fig. 2) may have been due to the large population sizes and short

generation times of most soil organisms, allowing for fast re-assembly and evolution of soil

communities[19].

Changes have been observed in the performance of individual species selected in

different species diversity and tested under experimental abiotic or biotic conditions in

previous studies[6,21,22]. In our study we demonstrated for the first time that changes in the

performance of entire plant communities through time may depend on a shared community

selection history of the assembled species. An ecosystem with individuals adapted to

optimise the use of the local resources by reducing interspecific competition, will be a well-

functioning and sustainable system as a whole[30]. Our new findings suggest that it is not

sufficient to preserve species outside a community context for the conservation of

biodiversity and its beneficial effects on ecosystem functioning and services. To protect

species interactions and ecosystem functioning more efficiently, novel strategies should

consider the conservation of entire communities or at least subsets of these.

Online content Methods, along with additional Extended Data display items and Source

Data, are available in the online version of the paper; references unique to these sections

appear only in the online paper.

1.   Cardinale, B.J. *et al.* Biodiversity loss and its impact on humanity. *Nature* **486,** 59-67

(2012).

2.   Huston, M.A. Hidden treatments in ecological experiments: re-evaluating the

ecosystem function of biodiversity. *Oecol.* **110,** 449-460 (1997).

3.   Tilman, D., Lehman, C.L. & Thomson, K.T. Plant diversity and ecosystem

productivity: theoretical considerations. *PNAS* **94,** 1857-1861 (1997).

4.   Loreau, M. & Hector, A. Partitioning selection and complementarity in biodiversity

experiments. *Nature* **412,** 72-76 (2001).

5.   Thorpe, A.S., Aschehoug, E.T., Atwater, D.Z. & Callaway, R.M. Interactions among

plants and evolution: plant interactions and evolution. *J. Ecol.* **99,** 729-740 (2011).

6.   Zuppinger-Dingley, D. *et al.* Selection for niche differentiation in plant communities

increases biodiversity effects. *Nature* **515,** 108-111 (2014).

7.   Balvanera, P. *et al.* Quantifying the evidence for biodiversity effects on ecosystem

functioning and services. *Ecol. Let.* **9,** 1146-1156 (2006).

8.   Reich, P.B. *et al.* Impacts of biodiversity loss escalate through time as redundancy

fades. *Science* **336,** 589-592 (2012).

9.   Cardinale, B.J. *et al.* Impacts of plant diversity on biomass production increase

through time because of species complementarity. *PNAS* **104,** 18123-18128 (2007).

10.   Meyer, S.T. *et al.* Effects of biodiversity strengthen over time as ecosystem

functioning declines at low and increases at high biodiversity. *Ecosphere* **7,** e01619

(2016).

11.   Schnitzer, S.A. *et al.* Soil microbes drive the classic plant diversity–productivity

pattern. *Ecology* **92,** 296-303 (2011).

12.   Whitham, T.G. *et al.* A framework for community and ecosystem genetics: from

genes to ecosystems. *Nat. Rev. Genet.* **7,** 510-523 (2006).

13.   Anderson, J.T., Willis, J.H. & Mitchell-Olds, T. Evolutionary genetics of plant

adaptation. *Trends Genet.* **27,** 258-266 (2011).

14.   Fakheran, S. *et al.* Adaptation and extinction in experimentally fragmented

landscapes. *PNAS* **107,** 19120-19125 (2010).

15.   Yoshida, T., Jones, L.E., Ellner, S.P., Fussmann, G.F. & Hairston, N.G. Rapid

evolution drives ecological dynamics in a predator–prey system. *Nature* **424,** 303-306

(2003).

16.   Fiegna, F., Scheuerl, T., Moreno-Letelier A., Bell, T. & Barraclough, T.G. Saturating

effects of species diversity on life-history evolution in bacteria. *Proc. R. Soc. B.* **282,**

20151794 (2015).

17.   Strauss, S.Y., Lau, J.A., Schoener, T.W. & Tiffin, P. Evolution in ecological field

experiments: implications for effect size. *Ecol. Lett.* **11,** 199-207 (2008).

18.   Agrawal, A.A., Hastings, A.P., Johnson, M.J.T., Maron, J.L. & Salminen, J.-P. Insect

herbivores drive real-time ecological and evolutionary change in plant populations.

Science **338,** 113-116 (2012).

19.   Lau, J.A. & Lennon, J.T. Rapid responses of soil microorganisms improve plant

fitness in novel environments. *PNAS* **35,** 14058–14062 (2012).

20.   Lawrence, D. *et al.* Species interactions alter evolutionary responses to a novel

environment. *PLoS Biol.* **10,** e1001330 (2012).

21.   Kleynhans, E.J., Otto, S.P., Reich, P.B. & Vellend, M. Adaptation to elevated CO_2 in

different biodiversity contexts. *Nat. Comm.* **7,** 12358 (2016).

22.   Rottstock, T., Kummer, V., Fischer, M. & Joshi, J. Rapid transgenerational effects in

Knautia arvensis in response to plant-community diversity. *J. Ecol.* (in press).

23.   Kardol, P., Cornips, N.J., van Kempen, M.M.L., Bakx-Schotman, J.M.T. & van der

Putten, W.H. Microbe-mediated plant–soil feedback causes historical contingency

effects in plant community assembly. *Ecol. Monogr.* **77,** 147-162 (2007).

24. Klironomos, J. Feedback with soil biota contributes to plant rarity and invasiveness in

communities. *Nature* **417,** 67-70 (2002).

25. Roscher, C. *et al.* The role of biodiversity for element cycling and trophic

interactions: an experimental approach in a grassland community. *Basic Appl. Ecol.*

5, 107-121 (2004).

26. Gebremikael, M.T., De Waele, J., Buchan, D., Soboksa, G.E. & De Neve, S. The

effect of varying gamma irradiation doses and soil moisture content on nematodes,

the microbial communities and mineral nitrogen. *Appl. Soil Ecol.* **92,** 1–13 (2015).

27. Lemmermeyer, S., Lörcher, L., van Kleunen, M. & Dawson, W. Testing the plant

growth-defense hypothesis belowground: Do faster-growing herbaceous plant species

suffer more negative effects from soil biota than slower-growing ones? *Am. Nat.* **186,**

264-271 (2015).

28. Zuppinger-Dingley, D., Flynn, D.F.B., De Deyn, G.B., Petermann, J.S. & Schmid, B.

Plant selection and soil legacy enhance long-term biodiversity effects. *Ecology* **97,**

918-928 (2016).

29. Petermann, J., Fergus, A.J.F., Turnbull, L. & Schmid, B. Janzen-Connell effects are

widespread and strong enough to maintain diversity in grasslands. *Ecology* **89,** 2399-

2406 (2008).

30. Pedruski, M.T., Fussman, G.F. & Gonzalez, A. A network approach reveals surprises

about the history of the niche. *Ecosphere* **7,** e01266 (2016).

Supplementary Information is available in the online version of the paper.

Acknowledgements Thanks to the Jena Experiment for providing infrastructure and help, to

D. Trujillo and M. Furler for technical assistance and to H. Martens for all the lab work for

the T-RFLP analyses. This study was supported by the Swiss National Science Foundation

(grant numbers 130720, 147092 and 166457 to B.S.) and the University Research Priority

Program Global Change and Biodiversity of the University of Zurich.

Author contributions B.S., D.B.F. and G.B.D.D. conceived the project; D.Z-D. and V.Y. set

up the experiment; S.J.V.M., T.H., D. Z-D. and V.Y. carried out the experiment; B.S., C.W.,

S.J.V.M. and T.H. analysed the data; D.B.F. analysed the TRFLP data; B.S., S.J.V.M., T.H.

and C.W. wrote the first draft of the manuscript. All authors contributed to the final

manuscript.

Author information The authors declare no competing financial interests. Correspondence

and requests for materials should be addressed to S.J.V.M. (sofia.vanmoorsel@ieu.uzh.ch) or

B.S. (bernhard.schmid@ieu.uzh.ch).

261
262

METHODS

Study site. The present study was conducted at the Jena Experiment field site (Jena,

Thuringia, Germany, 51˚N, 11˚E, 135m a.s.l.) from 2011 to 2015. The Jena Experiment is a

long-term biodiversity field experiment located in the floodplain of the river Saale where 60

Central European grassland species have been grown in a number of species combinations

since 2002[25].

Plant community selection. The 48 experimental plant communities of this study included

twelve monocultures, twelve 2-species mixtures, twelve 4-species mixtures and twelve 8-

species mixtures. We used two community selection treatments; plants with eight years of

shared community selection in the Jena Experiment (selected communities) and plants

without a common selection history in the Jena Experiment (naïve communities). The plant

seeds without a common selection history were obtained from the same commercial seed

supplier (Rieger Hofmann GmbH, in Blaufelden-Raboldshausen, Germany) as the seeds used

for the establishment of the original experimental plant communities. This supplier collected

plants of the different species at field sites in Germany and propagated them for at least 5

years in monoculture, reseeding them every year. Seeds of selected communities were

produced in an experimental garden in Zurich, Switzerland, from cuttings that had been made

in the Jena experiment and were then planted in Zurich in the original species combination in

plots fenced with plastic netting to reduce pollination between communities. To obtain

sufficient numbers of seeds from selected communities a small number was additionally

collected directly in the plots of the Jena Experiment. All these seeds were thus offspring of

plant populations that had been sown in 2002 and grown until 2010 in plots of the Jena

Experiment.

The seeds of selected and naïve communities were germinated on potting soil (BF4,

De Baat; Holland) in mid-January 2011 in a glasshouse in Zurich. In March 2011, the

seedlings were transported back to the Jena Experiment field site within 2 x 2 m subplots of

the original plots (Extended Data Fig. 1). There were four 1 x 1 m quadrats with different soil

types in each subplot (see next section). Each quadrat was split into two 1 x 0.5 m halves. We

transplanted seedlings of selected communities into one half and seedlings of naïve

communities into the other half of each quadrat at a density of 210 plants per m^2 with a 6-cm

distance between individuals in a hexagonal pattern (Extended Data Fig. 1). Species were

planted in equal proportions, but if a species was no longer present in an original plot of the

Jena Experiment it was excluded from both selected and naïve communities. Five plant

species were excluded in total. The seedlings received water every second day for 6 weeks

after transplanting to ensure the plants established.

Soil treatment. Within each 2 x 2 m subplot of the 48 plots of the Jena Experiment used for

the present study, the original plant cover was removed in September 2010 (and used for the

plant propagation in the experimental garden in Zurich, see above), and the soil was

excavated to a depth of 0.35 m and sieved. To minimize exchange of soil components

between quadrats within subplots and with the surrounding soil, two 5-cm layers of sand

were added to the bottom of the plots and separated with a 0.5 mm mesh net. The borders of

the quadrats and the subplots were separated by plastic frames (Extended Data Fig. 1). Using

the excavated original soil from each of the plots, four soil treatments were prepared. First,

half of the soil (approximately 600 kg per plot) was gamma-sterilized to remove the original

soil community. Half of the gamma-sterilized soil was then inoculated with 4 % (by weight)

of live sugar-beet soil and 4 % of sterilized original soil of the corresponding plot ("novel

soil" obtained by inoculation). Live sugar-beet soil was added to create a natural, but neutral

soil community and was previously collected in an agricultural sugar-beet field not associated

with the Jena Experiment, but with comparable soil properties. The other half of the gamma-

sterilized soil was inoculated with 4 % (by weight) of live sugar-beet soil and 4 % of live

original soil of the corresponding plot ("native soil" obtained by inoculation). The other half

of the soil was unsterilized and used for the other two soil treatments. Half of this soil was

filled back into one quadrat of the corresponding plot ("native soil"). The other half of the

unsterilized soil was mixed among all 48 plots and filled into the remaining quadrats. This

fourth soil treatment was abandoned after two years because the plant community was

excavated for another experiment. Therefore, this treatment is not included in the present

study.

Before the soils were added into the quadrats in December 2010, they were rested in

the field in closed bags to allow for the soil chemistry to equalize and to encourage soil biota

of the inocula to colonize the sterilized soil before planting. After the soil was added, all

quadrats were covered with a net and a water permeable black sheet to avoid spilling between

quadrats until the seedlings were transplanted in March 2011.

Maintenance and data collection. We maintained the test communities by weeding three

times a year and by cutting the plants to 3 cm above ground twice a year at typical grassland

harvest times (late May and August) in central Europe. To measure above ground biomass,

we harvested plant material 3 cm above ground from a 50 x 20 cm area in the center of each

half-quadrat, sorted it into species, dried it at 70°C and weighed the dry biomass.

T-RFLP assay. Terminal restricted fragment length polymorphism (T-RFLP) targeting the

16S RNA was used to characterize the composition of the soil bacterial communities[31]. In

April 2011, four soil samples per quadrat were extracted and pooled to assess the

establishment of soil microbial communities and to test whether soil treatments were distinct.

In 2012, a further set of soil samples was taken and analysed to confirm the establishment of

different soil biotic treatments. T-RFLP soil analyses revealed that bacterial communities of

the soil treatments remained distinct: each soil treatment had a characteristic bacterial

composition both one and two years after planting, with some overlap (Extended Data Fig. 5,

Extended Data Table 2).

Statistical analysis. For the present study, we analysed the data from four spring harvests

(2012, 2013, 2014 and 2015), which correspond to peak aboveground plant biomass values.

We analysed plant biomass (g/m^2) as a function of the design variables using mixed models

and summarized results in analyses of variance (ANOVA) tables (Extended Data Table 1).

Significance tests were based on approximate F-tests using appropriate error terms and

denominator degrees of freedom.

345   The fixed terms in the model were log-transformed species richness of the original

plots of the Jena Experiment (linear and quadratic), year of harvest (linear contrast and factor:

lH and fH, respectively), soil treatment (native vs. novel soil: SH), plant community selection

treatment (selected vs. naïve communities: PH) and interactions of these. The random terms

were plot, quadrat, half-quadrat and their interactions with time. Statistical analyses were

conducted using the software product R, version 3.2.3[32]. Mixed models using residual

maximum likelihood (REML) were fitted using the package ASReml for R[33].

352   The calculation of operational taxonomic units (OTUs) from the T-RFLP raw data

353   (restriction enzyme products) was done using the T-RFLP processing software T-REX[34] for

354   each soil treatment and year separately and the soil-specific outputs were then compared with

355   an analysis of similarities (anosim() function of the vegan package[35]). R-values and P-values

356   were extracted and the results were visualized using NMDS ordination[36].

358   31.   Liu, W.T., Marsh, T.L., Cheng, H. & Forney, L.J. Characterization of microbial

359   diversity by determining terminal restriction fragment length polymorphisms of

360   genes encoding 16S rRNA. *Appl. Environ. Microbiol.* **63,** 4516-4522 (1997).

32. R Core Team. R: A language and environment for statistical computing. R

Foundation for Statistical Computing, Vienna, Austria. URL https://www.R-

project.org/ (2015).

33. Butler, D. asreml: asreml() fits the linear mixed model. R package version 3.0.

www.vsni.co.uk (2009).

34. Culman, S.W., Bukowski, R., Gauch, H.G., Cadillo-Quiroz, H. & Buckley, D.H. T-

REX: software for the processing and analysis of T-RFLP data. *BMC Bioinf.* **10,** 171

(2009).

35. Oksanen, J., Blanchet, G.F., Kindt, R., Legendre, P., Minchin, P.R., O'Hara, R.B.,

Simpson, G.L., Solymos, P., Stevens, M.H.H. & Wagner, H. vegan: Community

Ecology Package. R package version 2.3-4. https://CRAN.R-

project.org/package=vegan (2016).

36. Shepard, R.N. The analysis of proximities: Multidimensional scaling with an

unknown distance function. I and II. *Psychometrika* **27,** 125-140 (1962).

Figure 1 | Selected plant communities show stronger biodiversity effects than naïve

plant communities. In selected plant communities, the increase in productivity from

monocultures to 2- and 4-species mixtures was stronger than in naïve plant communities.

Main panel: Quadratic response curves are derived from a mixed model. Points are predicted

means and standard errors derived from a mixed model including fixed-effects terms for

species richness and community selection history and random-effects terms for plot, quadrat

and half-quadrat (see Methods). Data are across four consecutive years. Inset panel: separate

regression lines for years, with darker colours representing later years.

385
386

Figure 2 | Diversity–productivity relationships are initially enhanced in novel soil but

over time approach those in native soil. In native soil, community productivity was lower

in comparison to novel soil but the diversity–productivity relationships remained stable over

time. Over time, the relationships in novel soil approached those in native soil (see inset

panel). Main panel: Quadratic response curves are derived from a mixed model. Points are

predicted means and standard errors derived from a mixed model including fixed-effects

terms for species richness and soil treatment and random-effects terms for plot, quadrat and

half-quadrat (see Methods). Data are across four consecutive years. Inset panel: separate

regression lines for years, with darker colours representing later years.

Figure 3 | Log-transformed species biomass ratios between selected and naïve plant

communities. The majority of plant species attained greater aboveground biomass in selected

plant communities compared to naïve plant communities. The studied plant species belong to

three different functional groups: grasses (white bars), herbs (light grey bars) and legumes

(dark grey bars). Data are for each species across all four experimental years, across soil

treatments and across species richness levels (n = 32–352). Four species with n < 32 were

excluded from the analysis. The stars represent P-values < 0.05 for species tested separately.