Community evolution increases plant productivity at low diversity

Sofia J. van Moorsel[1†], Terhi Hahl[1†], Cameron Wagg[1], Gerlinde B. De Deyn[2], Dan F.B.

Flynn[1], Debra Zuppinger-Dingley[1] and Bernhard Schmid[1*]

[1] URPP Global Change and Biodiversity and Department of Evolutionary Biology and

Environmental Studies, University of Zürich, Winterthurerstrasse 190, 8057 Zürich,

Switzerland

[2] Department of Environmental Sciences, Wageningen University, Droevendaalsesteeg 4,

6708 PB Wageningen, the Netherlands

[†] Shared first authorship

*Correspondence: E-mail: bernhard.schmid@ieu.uzh.ch

Emails of other authors (in order):

sofia.vanmoorsel@ieu.uzh.ch

terhi.hahl@ieu.uzh.ch

cameron.wagg@ieu.uzh.ch

gerlinde.dedeyn@wur.nl

danfbflynn@gmail.com

debra.zuppinger@ieu.uzh.ch

Running title: Community evolution in grasslands
Type of contribution: Letter
Key words: biodiversity, community evolution, co-selection, ecosystem functioning,
grassland species, Jena Experiment, plant productivity, soil organisms
Manuscript information: # words abstract: 149, # words main text: 5000, # references: 50,
figures: 5, # tables: 1

ABSTRACT

Species extinctions from local communities negatively affect ecosystem functioning.

Ecological mechanisms underlying these impacts are well studied but the role of evolutionary

processes is rarely assessed. Using a long-term field experiment, we tested whether natural

selection in plant communities increased biodiversity effects on productivity. We re-

assembled communities with 8-year co-selection history adjacent to communities with

identical species composition but no history of co-selection ("naïve communities").

Monocultures and in particular mixtures of two to four co-selected species were more

productive than their corresponding naïve communities over four years in soils with or

without co-selected microbial communities. At the highest diversity level of eight plant

species, no such differences were observed. Our findings suggest that plant community

evolution can lead to rapid increases in ecosystem functioning at low diversity but may take

longer at high diversity. This effect was not modified by treatments simulating co-

evolutionary processes between plants and soil organisms.

INTRODUCTION

A large number of experiments have shown that species richness positively influences ecosystem functioning, in particular plant biomass production (Tilman *et al.* 1997; Balvanera *et al.* 2006; Cardinale *et al.* 2007, 2012; Reich *et al.* 2012; Meyer *et al.* 2016). These biodiversity effects have been explained by selection effects that increase the chance of including productive species in diverse communities or by complementary effects between species, which allow mixtures to extract resources from the environment more efficiently (Loreau and Hector 2001; Roscher *et al.* 2008; Mueller *et al.* 2013). Furthermore, diversity-dependent reductions in soil fertility (Fornara & Tilman 2008) or density-dependent accumulations of specialist pathogens over time (Schnitzer *et al.* 2011) have been shown to contribute to decreasing productivity at low plant diversity and in plant monocultures.

In contrast to selection effects, complementarity effects between co-occurring species have been shown to increase over time (Cardinale *et al.* 2007; Fargione *et al.* 2007; Reich *et al.* 2012; Meyer *et al.* 2016). Evidence that this might be due to evolutionary processes in plant communities has been found in a glasshouse experiment comparing the community-level performance of plants selected in monocultures vs. plants selected in multi-species mixtures in newly assembled monocultures vs. two-species mixtures (Zuppinger-Dingley *et al.* 2014). This suggests that community evolution may shape diversity–productivity relationship more generally, which could be tested if entire communities of co-selected plant species would be compared with communities of the same plant species but without co-selection history ("naïve communities"). Here, the distinction between the terms "selected" and "co-selected" indicates that in the latter case the outcome of selection on a set of co-occurring species is assessed in the context of a test community formed by these species.

Evolution leading to community-level responses, i.e. changed species performances and interactions due to genetic changes in all or some of the species of the community, has

been referred to as community evolution (Goodnight 1990). Later, community evolution was

defined as genetically based changes among species constituting the community, which alter

species performances and interactions (Wilson 1997; Whitham *et al.* 2006). Such

evolutionary changes may occur via genetic recombination, mutations (Anderson *et al.*

2011), or a sorting-out from standing genetic variation through differential survival and

growth of individuals (Fakheran *et al.* 2010) and may be due to natural selection or random

genetic drift (Wilson & Bossert 1971). In particular, natural selection can lead not only to

changes in gene frequencies in populations within species, but selection at the level of

communities can in addition lead to correlated changes in gene frequencies in multiple

species (Whitham *et al.* 2006) in response to one another or to co-varying environmental

conditions. Community evolution can be detected by measurable differences in community-

level properties, such as productivity, between communities of co-selected species and naïve

communities (Swenson *et al.* 2000; Lawrence *et al.* 2012; Fiegna *et al.* 2014). But empirical

evidence for community evolution so far has only been demonstrated in bacterial

communities (Lawrence *et al.* 2012; Fiegna *et al.* 2014, 2015) and not yet in higher plants.

Here we report results from a field experiment where we tested whether plant community

evolution influences plant community productivity.

Recent evidence suggests that selection of particular genotypes from the total genetic

pool of a species may affect ecosystem functioning in field experiments (Strauss *et al.* 2008;

Lipowsky *et al.* 2011; Lau & Lennon 2012; Kleynhans *et al.* 2016; Rottstock *et al.* 2017). We

propose that co-selection of genotypes from the gene pool of entire communities may affect

ecosystem functioning if non-random niche or trait changes in response to other phenotypes

in the community result in more specialized niches with reduced overlap and a more

complete use of biotope space (Dimitrakopoulos & Schmid 2004; Jousset *et al.* 2011), thus

leading to increased plant community productivity. We therefore compared the productivity

of plant communities assembled from plants that have co-occurred for eight years in a long-

term grassland biodiversity experiment (the Jena Experiment, see Roscher *et al.* 2004) with

the productivity of plant communities of identical species composition, but without any co-

occurrence history ("naïve communities"). The naïve plants were obtained from the seed

supplier of the original seeds used to establish the Jena Experiment. We used experimental

plant monocultures and 2-, 4- or 8-species mixtures with twelve different species

compositions for each diversity level.

To explore if (co-)selection of plant species led to changes in trait distributions within

species we used plant height and specific leaf area (SLA) as representative functional traits

(Diaz *et al.* 2016), which have been used in previous biodiversity experiments (Roscher *et al.*

2015; Zuppinger-Dingley *et al.* 2014; Cadotte 2017). Variation in these two traits has been

shown to affect plant community productivity in contrasting ways: whereas low variation in

height is related to asymmetric competition for light and increased productivity via selection

effects, high variation in SLA is related to differential light-use strategies and increased

productivity via complementarity effects (Roscher *et al.* 2015; Cadotte 2017). As a

consequence, we expected a narrowing/widening of the within-species variation in

height/SLA at low biodiversity due to community evolution.

Plant community evolution in the field may also depend on the local environment,

such as the soils in which co-evolution with soil microorganisms occurs. For instance, plant–

soil feedback experiments have shown that soil biota change in response to different plant

species, which can in turn modify the composition and productivity of plant communities

(Klironomos 2002; Kardol *et al.* 2007). Further, it is thought that negative plant–soil

feedbacks may also incur selection for individuals that are able to reduce antagonistic and

improve beneficial associations with soil organisms (van der Putten *et al.* 2013; Zuppinger-

Dingley *et al.* 2016). To assess whether additional co-evolutionary processes between plants

and soil organisms modified plant community evolution, we grew the communities of co-

selected plants and naïve communities in soils with co-selected soil organisms (native soil)

and with external soil organisms (neutral soil; see Methods and Fig. S1). Community-level

plant productivity was measured each year from 2012 to 2015 by collecting species-specific

aboveground biomass at the time of peak biomass in spring in each soil treatment, whereas

the traits plant height and SLA were measured once in 2015 in the neutral soil only (see

Methods).

METHODS

Study site

The present study was conducted at the Jena Experiment field site (Jena, Thuringia,

Germany, 51˚N, 11˚E, 135m a.s.l.) from 2011 to 2015. The Jena Experiment is a long-term

biodiversity experiment where 60 grassland species have been grown in different

combinations since 2002 (Roscher *et al.* 2004).

Community-evolution treatment (plant history)

The 48 experimental plant communities of this study included twelve monocultures

(of which one was removed from all analyses because it was planted with the wrong species),

twelve 2-species mixtures, twelve 4-species mixtures and twelve 8-species mixtures. We used

two community-evolution treatments (plant histories); plants with eight years of co-selection

history in different plant communities in the Jena Experiment (communities of co-selected

plants) and plants without such co-selection history (naïve communities). For convenience,

we use the term co-selection also for monocultures with a selection history of eight years in

the Jena Experiment, even though in this case the term could only apply to potential co-

selection with soil organisms. The plant seeds of naïve communities were obtained from the

same seed supplier (Rieger Hofmann GmbH, in Blaufelden-Raboldshausen, Germany) as the

seeds used for the establishment of the original communities of the Jena Experiment. This

supplier collected plants of the different species at field sites in Germany and propagated

them for at least five years in monoculture, reseeding them every year. Seeds of communities

of co-selected plants were produced in an experimental garden in Zurich, Switzerland, from

cuttings that had been made in the Jena Experiment and were then planted in Zurich in the

original species combination in plots fenced with plastic netting to reduce pollination

between communities. To obtain sufficient numbers of seeds from communities of co-

selected plants, a small number was additionally collected directly in the plots of the Jena

Experiment. All these seeds were thus offspring of plant populations that had been sown in

2002 and grown until 2010 in plots of the Jena Experiment.

The seeds of communities of co-selected plants and naïve communities were

germinated in potting soil (BF4, De Baat; Holland) in mid-January 2011 in a glasshouse in

Zurich. In March 2011, the seedlings were transported back to the field site of the Jena

Experiment and planted within 2 x 2 m subplots of the original plots (Fig. S1). There were

four 1 x 1 m quadrats with different soil treatments in each (see next section). Each quadrat

was further divided into two 1 x 0.5 m halves. The seedlings of communities of co-selected

plants were transplanted into one half and seedlings of naïve communities into the other half

of each quadrat at a density of 210 plants per m^2 (Fig. S1). Species were planted in equal

proportions, but if a species was no longer present in an original plot of the Jena Experiment

it was excluded from both communities of co-selected plants and naïve communities. Five

plant species were excluded in total.

Soil treatment

Within each 2 x 2 m subplot of the 48 plots of the Jena Experiment used for the

present study, the original plant cover was removed in September 2010 and the soil was

excavated to a depth of 0.35 m and sieved. To minimize exchange of soil components

between quadrats within subplots and with the surrounding soil, two 5-cm layers of sand

were added to the bottom of the plots and separated with a 0.5 mm mesh net. The borders of

the quadrats and the subplots were separated by plastic frames (Fig. S1). Using the excavated

original soil from each of the plots, four soil treatments were prepared. First, half of the soil

(approximately 600 kg per plot) was gamma-sterilized to remove the original soil

community. Half of the gamma-sterilized soil was then inoculated with 4 % (by weight) of

live sugar-beet soil and 4 % of sterilized original soil of the corresponding plot ("neutral soil"

obtained by inoculation). Live sugar-beet soil was added to create a natural but neutral soil

community and was previously collected in a sugar-beet field not associated with the Jena

Experiment, but with comparable soil properties. The other half of the gamma-sterilized soil

was inoculated with 4 % (by weight) of live sugar-beet soil and 4 % of live original soil of

the corresponding plot ("native soil" obtained by inoculation). The other half of the soil was

unsterilized and used for the other two soil treatments. Half of this soil was filled back into

one quadrat of the corresponding plot ("native soil"). The other half of the unsterilized soil

was mixed among all plots and filled into the remaining quadrats. This fourth soil treatment

was abandoned after two years because the plant community was excavated for another

experiment. Therefore, this treatment is not included in the present study.

Before the soils were added into the quadrats in December 2010, they were rested in

the field in closed bags to allow for the soil chemistry to equalize and to encourage soil biota

of the inocula to colonize the sterilized soil before planting. After the soil was added, all

quadrats were covered with a net and a water permeable black sheet to avoid spilling between

quadrats until the seedlings were transplanted in March 2011.

To test the effectiveness of our soil treatments, we collected soil samples in April

2011 and August 2012 for each quadrat and analysed their fungal communities using

terminal-restriction fragment length polymorphism (T-RFLP, see Supporting information).

This method allowed us to establish that fungal communities of the soil treatments remained

distinct (Table S3). Because such a method does not inform about species identity, the

analysis of species- or even clade-specific co-evolutionary interactions between plants and

their associated microbial communities was beyond the scope of this study.

Data collection

We maintained the test communities by weeding three times a year and by cutting the

plants twice a year at typical grassland harvest times (late May and August) in central

Europe. To measure productivity, we harvested plant material 3 cm aboveground from a 50 x

20 cm area in the centre of each half-quadrat, sorted it into species, dried it at 70°C and

weighed the dry biomass.

Trait measurements

At the end of the experiment, in May 2015, we measured plant height and SLA for 30

species in neutral soil. For each species, we collected up to 20 representative leaves

(depending on the leaf size of the species) from four individuals and measured the leaf area

by scanning fresh leaves with a Li-3100 Area Meter (Li-cor Inc., Lincoln, Nebraska, USA)

immediately after harvest and determining the mass of the same leaves after drying.

Statistical analysis

We analysed the data from the four spring harvests 2012–2015, which corresponded

to peak aboveground plant biomass values. We analysed plant biomass (g/m^2) as a function

of the design variables using mixed models and summarized results in analyses of variance

(ANOVA) tables (e.g. Table 1). Significance tests were based on approximate F-tests using

appropriate error terms and denominator degrees of freedom. The fixed terms in the model

were species richness of the original plots of the Jena Experiment (linear (lSR) and quadratic

contrast (qSR) of the logarithm of plant species richness or factor with 4 levels: facSR), year

of harvest (linear contrast: linHar), soil treatment (factor with 3 levels: SH), community-

evolution treatment (communities of co-selected plants vs. naïve communities: PH as

abbreviation for plant history) and interactions of these. The random terms were plot,

quadrat, half-quadrat and their interactions with year of harvest. Statistical analyses were

conducted using the software R, version 3.2.3 (R Core Team 2015). Mixed models using

residual maximum likelihood (REML) were fitted using the package ASReml for R (Butler

2009).

　　　　To compare biomass production between communities of co-selected plants vs. naïve

communities across soil treatments and years and to compare these communities from 2012–

2015 with the ancestral communities from 2003–2006, we also calculated relative

productivities in percentage of the mean productivity of communities at the highest diversity

of eight species for each soil treatment x community-evolution treatment x year combination.

We chose the 8-species richness level as reference because there were no productivity

differences between communities of co-selected plants vs. naïve communities at this diversity

level (see Results). Furthermore, we analysed effects of the community-evolution treatment,

calculated separately for each particular species composition x soil treatment x year

combination, using log-ratios of proportional productivity changes between communities of

co-selected plants and naïve communities. In biodiversity experiments, proportional changes

typically differ from absolute changes (and absolute changes expressed relative to a control

level) because of biomass overyielding at higher compared with lower diversity.

　　　　To test whether the presence of particular plant functional groups could differentially

affect community evolution, we fitted corresponding contrasts in the analyses of community

productivity. We also analysed the responses of individual species to the community-

evolution treatment. Differential responses between species could mean that community

evolution leads to changes in species abundance distributions, which in turn could lead to

changed productivities. We therefore compared the evenness of species aboveground

biomasses between communities of co-selected plants vs. naïve communities, species

richness levels, soil treatments and years 2012–2015.

Within-species variation in plant height and SLA was calculated as the within-species

variance component for each community (residual mean square after fitting species). We had

insufficient trait data to test for increased between-species variation in communities of co-

selected plants containing a mixture of species.

RESULTS

Community productivity

Overall, for each doubling of species richness community aboveground biomass

increased by 100 $g \cdot m^{-2} \cdot y^{-1}$, a typical value for grassland biodiversity experiments (Hector *et*

al. 1999) (Fig. 1, Table 1). However, between monocultures and 2- to 4-species mixtures this

increase in productivity was steeper for communities of co-selected plants than for naïve

communities of the same species composition ($P < 0.01$ for interaction qSR x PH in Table 1).

Communities of co-selected plants were also more productive than naïve communities on

average. Furthermore, productivity increased across years in communities of co-selected

plants but decreased in naïve communities (interaction PH x linHar in Table 1). Because no

differences in productivity between the two community-evolution treatments were found at

the highest species-richness level (Fig. 1), we also analysed the productivity of all

communities in relative terms as percentage of the mean productivity of 8-species mixtures

for each plant history x soil treatment x year combination (Fig. 2 and Fig. S2). This analysis

confirmed that especially 2- and 4-species mixtures of co-selected plants increased

productivity relative to 8-species mixtures (see Supporting Information).

Comparison with ancestor plant communities

To test whether the communities of co-selected plants were already particularly

productive in 2- and 4-species mixtures at the beginning of the Jena Experiment (i.e. when

they were "naïve" communities themselves), we compared the productivity data of 2003–

2006 with the data of 2012–2015. Because productivities were generally higher at the

beginning of the Jena Experiment (Fig. S3a), we used relative productivity (percentage of

mean of 8-species mixtures per year) to standardize for differences in overall productivity

between time periods (Fig. S3b). The plant communities were established on neutral soil in

2002 at the beginning of the Jena Experiment. We therefore used only data from neutral soil

for the period 2012–2015. The communities of co-selected plants were significantly different

in their response compared to the two types of naïve communities because of their increased

relative productivity in 2- and 4-species mixtures ($F_{1,46.5} = 5.73$, $P = 0.021$ for the interaction

of plant history with the contrast "2- or 4-species mixtures vs. others"; Fig. S3b). Differences

between the communities of the naïve ancestors of the co-selected plants and our current re-

assembled naïve plant communities were small and not significant ($F_{1,46.1} = 0.23$, $P = 0.637$

for the interaction of the contrast "naïve ancestors vs. current naïve communities" with the

contrast "2- or 4- species mixtures vs. others").

Influence of soil environment

Plant community productivity was initially greater in inoculated soils, which was

reflected in an overall main effect of soil treatment and a significant interaction with year

(Table 1). This was probably caused by the nutrient flush associated with gamma-sterilization

of the soil (Gebremikael *et al.* 2015). But we found no evidence that our soil treatments

modified the differences in biodiversity effects between communities of co-selected plants

and naïve communities ($P > 0.2$ for the three-way interactions of contrasts of the logarithm of

species richness or the 4-level factor of species richness with soil and community-evolution

treatments).

Proportional productivity increases of single species compositions due to community evolution

Consistent with the results of the overall analysis of productivities, the log-ratios of

productivities between communities of co-selected plants and naïve communities were on

average zero for the 8-species mixtures, that is, the community-evolution treatment did not

lead to a proportional increase in productivity at the highest level of species richness (Fig. 3).

However, whereas the productivity increases found in the overall analysis were largest in the

4-species mixtures (see Fig. 1), the proportional increases were largest in the 2-species

mixtures (Fig. 3). This was due to the overall lower productivity of 2- as compared with 4-

species mixtures. Using contrasts between the different diversity levels, we could confirm

that the three low diversity levels were significantly different from the 8-species mixtures

($F_{1,37.1} = 5.34$ and $P = 0.026$). Among the three low diversity levels, the 2-species mixtures

had significantly greater log ratios than 4-species mixtures and monocultures ($F_{1,39.2} = 4.44$, P

$= 0.042$). Comparing 2-species mixtures and monocultures, 2-species mixtures had

significantly greater log ratios than monocultures ($F_{1,41.7} = 4.247$, $P = 0.046$).

Influence of functional groups, responses of individual species and evenness

Next, we tested whether the presence of particular plant functional groups influenced

the increase in productivity in communities of co-selected plants at the 2- and 4-species

richness levels; especially as legumes are known to drive over-yielding in grasslands (Spehn

et al. 2002). The presence of legumes and other plant functional groups did not provide any

further explanation for our results. Species-level productivity within communities was higher

for the majority of plant species with a co-selection history, irrespective of functional-group

identity (Fig. 4). Naïve communities showed marginally more even species abundance

distributions (Table S2), mainly due to the lower evenness of communities of co-selected

plants in the unsterilized native soil treatment (Fig. S4). Over the course of the experiment,

evenness decreased similarly in communities of co-selected plants and naïve communities

(Table S2).

Within-species trait variation

Finally, we analysed changes in within-species trait variation along the species

richness gradient as a potential mechanism contributing to the difference in productivity

between communities of co-selected plants and naïve communities (Roscher *et al.* 2015;

Cadotte 2017). Within-species variation in plant height was marginally reduced at low

diversity in communities of co-selected plants compared with naïve communities (Fig. 5a;

$F_{1,73.3} = 3.187$, P = 0.078 for the interaction of log species richness with plant history). In

contrast, within-species variation in specific leaf area (SLA) was increased at low diversity

for communities of co-selected plants and decreased for naïve communities (Fig. 5b; $F_{1,69.2} =$

4.87, $P = 0.031$ for the interaction of log species richness with plant history).

DISCUSSION

Our results show that eight years of community evolution in a biodiversity experiment

can increase biodiversity effects on community productivity, suggesting that this may at least

in part explain why biodiversity effects commonly increase over time in such experiments

(Cardinale *et al.* 2007; Fargione *et al.* 2007; Reich *et al.* 2012; Meyer *et al.* 2016). The

greater productivity in communities consisting of co-selected plants compared with

communities consisting of naïve plants was particularly evident in communities comprised of

two or four species whereas 8-species mixtures with and without co-selection history on

average showed the same productivity. It is conceivable that selection pressure was

dampened in communities where more than four species co-occurred. For instance, during

initial establishment in a diverse community, each individual can have a very different set of

immediate neighbours that could constrain the consistency in the selection pressure on

individuals within a community. With fewer species in a mixture, the potential for the

evolution of increased complementarity between plant species should be greater, given the

relative constancy of the neighbours any given plant experiences. The greater proportional

increase of productivity in communities of co-selected plant species at the 2- than at the 4-

species richness level, and the absence of such an increase at the 8-species richness level, are

compatible with the idea that evolution for co-adaptation is stronger at low than at high

diversity. As suggested by Cardinale et al. (2012), there might be an upper limit of species

richness beyond which selection is unlikely to strengthen biodiversity effects. Additionally,

community evolution leading to increased plant growth and productivity in diverse mixtures

may be at the expense of reduced pathogen defence (Lemmermeyer *et al.* 2015).

352   The performance of the naïve communities in the current study over the four years

was comparable to the initial performance of the ancestral community of the co-selected

plants (2003–2006). This similarity supports the view that the observed results at 2- and 4-

species richness levels in communities of co-selected compared with communities of naïve

plants are likely due to diversity-dependent community evolution. Indeed, the naïve

communities did not catch up with the communities of co-selected plants during the course of

the current experiment and differences in productivity from 2012 to 2015 even increased

between the two community-evolution treatments (see Fig. 2). This suggests that in our study

community evolution was not solely due to an immediate sorting out of genotypes from

standing variation within species (Fakheran *et al.* 2010) during seedling establishment and

initial growth. However, differential responses of individual species to the community-

evolution treatments (see Fig. 4) could have caused the marginally decreased evenness of

species abundance distributions in communities of co-selected plants. Thus, abundance shifts between species could have contributed to the higher productivities of communities of co-selected plants at intermediate diversity levels or they might have contributed to avoid reduced productivities of naïve communities at the highest diversity level. A further driving force behind community evolution for greater productivity at low diversity may have been strong responses of particular functional groups or functional group combinations to co-selection (Zuppinger-Dingley *et al.* 2014). There was, however, no evidence for any functional-group specific effect typically found in other contexts of biodiversity–ecosystem functioning research (Hooper & Vitousek 1997; Spehn *et al.* 2002).

Intraspecific variation in plant height was smaller and intraspecific variation in SLA was larger at low diversity in communities of co-selected plants than in naïve communities (see Fig. 5), a result in line with previous findings regarding effects of interspecific variation in these traits on community productivity (Roscher *et al.* 2015; Cadotte 2017): because plant-height related competition for light is asymmetric variation in this trait does not increase complementarity and productivity in plant stands but variation in SLA can do so because it is related to differential light-use strategies. Our results suggest an evolutionary narrowing of height-related niches and an evolutionary broadening of SLA-related niches in monocultures and low-diversity mixtures in (co-)selected plants. At higher diversity this trend was reversed (see Fig. 5), perhaps because here intraspecific variation is relatively less important than interspecific variation. For example, the narrowing of within-species variation in SLA with increasing diversity in communities of co-selected plants is an expected consequence of character displacement between species (Zuppinger-Dingley *et al.* 2014). In contrast, the increasing within-species variation in SLA with increasing diversity in naïve communities (see Fig. 5b) may have been caused by a more heterogeneous biotic environment at high diversity. Because our assessment of plant functional traits was limited to two traits measured

at the end of the experiment, the above interpretations can only hint at some mechanisms

potentially underpinning the observed effects of our community-evolution treatment on

community productivity. As pointed out for example by Cadotte (2017), multivariate trait

variation may be more important than variation in any particular trait in particular for

complementarity effects.

Our community-evolution treatment also led to increased productivity in

monocultures, where plants could not have been co-selected with other plant species but only

with soil organisms. This would be consistent with an earlier finding from the Jena

Experiment that showed a reduction of negative plant–soil feedbacks in plants that had been

grown for 8 years in monoculture compared with plants that had been grown in mixtures

(Zuppinger-Dingley *et al.* 2016). There was a slight indication that naïve plants where

initially less negatively affected in neutral than in native soil but over time showed increased

negative effects (see Fig. 2), possibly due to accumulation of soil pathogens (Schnitzer *et al.*

2011). An alternative but not mutually exclusive explanation for the higher productivity of

monocultures of (co-)selected plants than naïve plants could be the evolution of increased

resource-uptake niches within species as indicated in the previous paragraph (Bazzaz 1996):

assuming a correlation between resource-uptake and trait-based niches (Roscher *et al.* 2015),

the increase in within-species variation in SLA (and the decrease in within-species variation

in plant height) in monocultures of selected plants would be consistent with this explanation.

Positive plant diversity–productivity relationships may not only be driven by

complementary resource use, and thus increased performance at high diversity (Roscher *et al.*

2008; Mueller *et al.* 2013), but also by pathogen accumulation in the soil and thus reduced

performance at low diversity (Schnitzer *et al.* 2011). Previous studies in the context of

biodiversity–ecosystem functioning research have reported negative plant–soil feedbacks in

native as opposed to neutral soils (Klironomos 2002; Petermann *et al.* 2008; Cortois *et al.*

2016). Consequently, an increase of biodiversity effects during community evolution could

also be due to the presence of co-selected soil biota. In our study, however, the outcome of

the community-evolution treatment in mixtures was largely independent of the presence of

co-selected soil biota. It is conceivable that co-evolution of plants with soil biota in our

experimental systems was not effective because the large population sizes and short

generation times of most soil organisms contributed to the re-assembly and fast evolution of

soil communities (Lau & Lennon 2012).

Changes in the performance of individual species selected at different species-

richness levels and tested under experimental abiotic or biotic conditions have been observed

in previous studies (Lipowsky *et al.* 2011; Zuppinger-Dingley *et al.* 2014; Kleynhans *et al.*

2016; Rottstock *et al.* 2017). Here, we demonstrated that changes in the performance of entire

plant communities over time depend on a history of co-selection among the plant species of

the assembled mixtures. We suggest that these changes are the result of community evolution

because they were maintained through seed production in an experimental garden and

propagation of seedlings in a glasshouse to the replanting of communities in the field.

However, we cannot exclude maternal carry-over and epigenetic changes (Verhoeven *et al.*

2016) as additional potential evolutionary mechanisms. The observed effects of our

community-evolution treatment could have been due to co-selection between the species

within each particular community composition. Alternatively, in a more general process,

"diffuse" co-selection may have increased the ability of species to grow with any other

species. Considering the large residual variation among the productivity responses of the

particular community compositions (see Fig. 3), a mixture of mechanisms seems likely.

Nevertheless, that the naïve plants were generally poorly adapted to grow with other species

or under the abiotic conditions at the field site would not be consistent with their equally high

community productivity as that of co-selected plants in 8-species mixtures.

Independent of the mechanism, our findings of positive effects of community

evolution on community productivity suggest that for the conservation of biodiversity and its

beneficial influence on ecosystem functioning and services it may be best to preserve plant

species in a community context rather than in separation from other plant species (and soil

biota). Where only the latter is possible, we recommend that populations from communities

of particular species compositions are nevertheless kept separate to later re-assemble these

communities with those populations that have evolved together.

ACKNOWLEDGEMENTS

Thanks to the Jena Experiment for providing infrastructure and help, to D. Trujillo and M.

Furler for technical assistance and to H. Martens for the lab work for the T-RFLP analyses.

Thanks to V. Yadav for the establishment of the plots. We thank Tim Paine, Marc Cadotte

and Mark van Kleunen for their helpful comments on the manuscript and Akira Mori and an

anonymous reviewer for their very constructive comments during the reviewing process. This

study was supported by the Swiss National Science Foundation (grant numbers 130720,

147092 and 166457 to BS), the Netherlands Organisation for Scientific Research (NWO-

ALW Vidi grant to GBDD) and the University Research Priority Program Global Change and

Biodiversity of the University of Zurich. The Jena Experiment is funded by The Deutsche

Forschungsgemeinschaft (DFG, German Research Foundation, FOR1451).

AUTHORSHIP

BS, DFBF and GBDD conceived the project; DZ-D set up the experiment; SJVM, TH and

DZ-D carried out the experiment; BS, CW, SJVM and TH analysed the data; DFBF analysed

the TRFLP data; BS, SJVM, TH and CW wrote the first draft of the manuscript. All authors

contributed to the final manuscript.

REFERENCES

467 Anderson, J.T., Willis, J.H. & Mitchell-Olds, T. (2011). Evolutionary genetics of plant
468 adaptation. *Trends Genet.*, 27, 258–266
469

470 Balvanera, P., Pfisterer, A.B., Buchmann, N., He, J.-S., Nakashizuka, T., Raffaelli, D., *et al.*
471 (2006). Quantifying the evidence for biodiversity effects on ecosystem functioning and
472 services: Biodiversity and ecosystem functioning/services. *Ecol. Lett.*, 9, 1146–1156
473

474 Bazzaz, F.A. (1996). *Plants in Changing Environments: Linking Physiological, Population,*
475 *and Community Ecology*. Cambridge University Press
476

477 Butler, D. asreml: asreml() fits the linear mixed model. R package version 3.0.
478 www.vsni.co.uk (2009)
479

480 Cadotte, M.W. (2012). Functional traits explain ecosystem function through opposing
481 mechanisms. *Ecol. Lett.*, 20, 989–996
482

483 Cardinale, B.J., Duffy, J.E., Gonzalez, A., Hooper, D.U., Perrings, C., Venail, P., *et al.*
484 (2012). Biodiversity loss and its impact on humanity. *Nature*, 486, 59–67
485

486 Cardinale, B.J., Wright, J.P., Cadotte, M.W., Carroll, I.T., Hector, A., Srivastava, D.S., *et al.*
487 (2007). Impacts of plant diversity on biomass production increase through time because of
488 species complementarity. *Proc. Natl. Acad. Sci.*, 104, 18123–18128
489

490 Cortois, R., Schröder-Georgi, T., Weigelt, A., van der Putten, W.H. & De Deyn, G.B. (2016).
491 Plant-soil feedbacks: role of plant functional group and plant traits. *J. Ecol.*, 104, 1608–1617
492

493 Diaz, S., Kattge, J., Cornelissen, J.H.C., Wright, I.J., Lavorel, S., Dray, S., *et al.* (2016). The
494 global spectrum of plant form and function. *Nature*, 529, 167–171
495

496 Dimitrakopoulos, P.G. & Schmid, B. (2004). Biodiversity effects increase linearly with
497 biotope space. *Ecol. Lett.*, 7, 574–583
498

499 Fakheran, S., Paul-Victor, C., Heichinger, C., Schmid, B., Grossniklaus, U. & Turnbull, L.A.
500 (2010). Adaptation and extinction in experimentally fragmented landscapes. *Proc. Natl.*
501 *Acad. Sci.*, 107, 19120–19125
502

503 Fargione, J., Tilman, D., Dybzinski, R., Lambers, J.H.R., Clark, C., Harpole, W.S., *et al.*
504 (2007). From selection to complementarity: shifts in the causes of biodiversity-productivity
505 relationships in a long-term biodiversity experiment. *Proc. R. Soc. B Biol. Sci.*, 274, 871–876
506

507 Fiegna, F., Moreno-Letelier, A., Bell, T. & Barraclough, T.G. (2014). Evolution of species
508 interactions determines microbial community productivity in new environments. *ISME J.*, 9,
509 1235–1245
510

511 Fiegna, F., Scheuerl, T., Moreno-Letelier, A., Bell, T. & Barraclough, T.G. (2015). Saturating

512 effects of species diversity on life-history evolution in bacteria. *Proc. R. Soc. B Biol. Sci.*,
513 282, 20151794
514
515 Fornara, D.A. & Tilman, D. (2008). Plant functional composition influences rates of soil
516 carbon and nitrogen accumulation. *J. Ecol.*, 96, 314–322
517
518 Gebremikael, M.T., De Waele, J., Buchan, D., Soboksa, G.E. & De Neve, S. (2015). The
519 effect of varying gamma irradiation doses and soil moisture content on nematodes, the
520 microbial communities and mineral nitrogen. *Appl. Soil Ecol.*, 92, 1–13
521
522 Goodnight, C.J. (1990a) Experimental Studies of Community Evolution II: The Ecological
523 Basis of the Response to Community Selection. *Evolution*, 44, 1625–1636.
524
525 Hector, A., Schmid, B., Beierkuhnlein, C., Caldeira, M.C., Diemer, M., Dimitrakopoulos,
526 P.G., *et al.* (1999). Plant diversity and productivity experiments in European grasslands.
527 *Science*, 286, 1123–1127
528
529 Hooper, D.U. & Vitousek, P.M. (1997). The effects of plant composition and diversity on
530 ecosystem processes. *Science*, 277, 1302–1305
531
532 Jousset, A., Schulz, W., Scheu, S. & Eisenhauer, N. (2011). Intraspecific genotypic richness
533 and relatedness predict the invasibility of microbial communities. *ISME J.*, 5, 1108–1114
534
535 Kardol, P., Cornips, N.J., van Kempen, M.M.L., Bakx-Schotman, J.M.T. & van der Putten,
536 W.H. (2007). Microbe-mediated plant-soil feedback causes historical contingency effects in
537 plant community assembly. *Ecol. Monogr.*, 77, 147–162
538
539 Kleynhans, E.J., Otto, S.P., Reich, P.B. & Vellend, M. (2016). Adaptation to elevated CO2 in
540 different biodiversity contexts. *Nat. Commun.*, 7, 12358
541
542 Klironomos, J.N. (2002). Feedback with soil biota contributes to plant rarity and invasiveness
543 in communities. *Nature*, 417, 67–70
544
545 Lau, J.A. & Lennon, J.T. (2012). Rapid responses of soil microorganisms improve plant
546 fitness in novel environments. *Proc. Natl. Acad. Sci.*, 109, 14058–14062
547
548 Lawrence, D., Fiegna, F., Behrends, V., Bundy, J.G., Phillimore, A.B., Bell, T., *et al.* (2012).
549 Species interactions alter evolutionary responses to a novel environment. *PLoS Biol.*, 10,
550 e1001330
551
552 Lemmermeyer, S., Lörcher, L., van Kleunen, M. & Dawson, W. (2015). Testing the plant
553 growth-defense hypothesis belowground: do faster-growing herbaceous plant species suffer
554 more negative efects from soil biota than slower-growing ones? *Am. Nat.*, 186, 264–271
555
556 Lipowsky, A., Schmid, B. & Roscher, C. (2011). Selection for monoculture and mixture
557 genotypes in a biodiversity experiment. *Basic Appl. Ecol.*, 12, 360–371
558
559 Loreau, M. & Hector, A. (2001). Partitioning selection and complementarity in biodiversity
560 experiments. *Nature*, 412, 72–76.

561
562 Meyer, S.T., Ebeling, A., Eisenhauer, N., Hertzog, L., Hillebrand, H., Milcu, A., *et al.*
563 (2016). Effects of biodiversity strengthen over time as ecosystem functioning declines at low
564 and increases at high biodiversity. *Ecosphere*, 7, e01619
565
566 Mueller, K.E., Tilman, D., Fornara, D.A. & Hobbie, S.E. (2013). Root depth distribution and
567 the diversity–productivity relationship in a long-term grassland experiment. *Ecology*, 94,
568 787–793
569
570 Petermann, J.S., Fergus, A.J., Turnbull, L.A. & Schmid, B. (2008). Janzen-Connell effects
571 are widespread and strong enough to maintain diversity in grasslands. *Ecology*, 89, 2399–
572 2406
573
574 R Core Team. R: A language and environment for statistical computing. R Foundation for
575 Statistical Computing, Vienna, Austria. URL https://www.R-project.org/ (2015).
576
577 Reich, P.B., Tilman, D., Isbell, F., Mueller, K., Hobbie, S.E., Flynn, D.F.B., *et al.* (2012).
578 Impacts of biodiversity loss escalate through time as redundancy fades. *Science*, 336, 589–
579 592
580
581 Roscher, C., Schumacher, J., Baade, J., Wilcke, W., Gleixner, G., Weisser, W.W., *et al.*
582 (2004). The role of biodiversity for element cycling and trophic interactions: an experimental
583 approach in a grassland community. *Basic Appl. Ecol.*, 5, 107–121
584
585 Roscher, C., Thein, S., Schmid, B. & Scherer-Lorenzen, M. (2008). Complementary nitrogen
586 use among potentially dominant species in a biodiversity experiment varies between two
587 years. *J. Ecol.*, 96, 477–488
588
589 Roscher, C., Schumacher, J., Schmid, B. & Schulze, E.-D. (2015). Contrasting effects of
590 intraspecific trait variation on trait-based niches and performance of legumes in plant
591 mixtures. *PloS One*, 10, e0119786
592
593 Roscher, C., Schmid, B., Kolle, O. & Schulze, E.-D. (2016). Complementarity among four
594 highly productive grassland species depends on resource availability. *Oecologia*, 181, 571–
595 582
596
597 Rottstock, T., Kummer, V., Fischer, M. & Joshi, J. (2017). Rapid transgenerational effects in
598 *Knautia arvensis* in response to plant community diversity. *J. Ecol.*, 105, 714–725
599
600 Schnitzer, S.A., Klironomos, J.N., HilleRisLambers, J., Kinkel, L.L., Reich, P.B., Xiao, K., *et*
601 *al.* (2011). Soil microbes drive the classic plant diversity–productivity pattern. *Ecology*, 92,
602 296–303
603
604 Spehn, E.M., Scherer-Lorenzen, M., Schmid, B., Hector, A., Caldeira, M.C.,
605 Dimitrakopoulos, P.G., *et al.* (2002). The role of legumes as a component of biodiversity in a
606 cross-European study of grassland biomass nitrogen. *Oikos*, 98, 205–218
607
Strauss, S.Y., Lau, J.A., Schoener, T.W. & Tiffin, P. (2008). Evolution in ecological field
experiments: implications for effect size. *Ecol. Lett.*, 11, 199–207

Swenson, W., Wilson, D.S., Elias, R. (2000). Artificial ecosystem selection. *Proc. Natl.*
Acad. Sci., 97, 9110–9114

Tilman, D., Lehman, C.L. & Thomson, K.T. (1997). Plant diversity and ecosystem
productivity: theoretical considerations. *Proc. Natl. Acad. Sci.*, 94, 1857–1861

Van der Putten, W.H., Bardgett, R.D., Bever, JD., Bezemer, T.M., Casper, B.B., Fukami, T.,
et al. (2013). Plant-soil feedbacks: the past, the present and future challenges. *J. Ecol.*, 101,
265–276

Verhoeven, K.J.F., vonHoldt, B.M. & Sork, V.L. (2016). Epigenetics in ecology and
evolution: what we know and what we need to know. *Mol. Ecol.*, 25, 1631–1638
Whitham, T.G., Bailey, J.K., Schweitzer, J.A., Shuster, S.M., Bangert, R.K., LeRoy, C.J., *et*
al. (2006). A framework for community and ecosystem genetics: from genes to ecosystems.
Nat. Rev. Genet., 7, 510–523

Wilson, E.O. & Bossert, W.H. (1971) A primer of population biology. *Sinauer Associates*

Wilson, D.S. (1997) Biological Communities as Functionally Organized Units. *Ecology*, 78,
2018–2024

Zuppinger-Dingley, D., Flynn, D.F.B., De Deyn, G.B., Petermann, J.S. & Schmid, B. (2016).
Plant selection and soil legacy enhance long-term biodiversity effects. *Ecology*, 97, 918–928

Zuppinger-Dingley, D., Schmid, B., Petermann, J.S., Yadav, V., De Deyn, G.B. & Flynn,
D.F.B. (2014). Selection for niche differentiation in plant communities increases biodiversity
effects. *Nature*, 515, 108–111

SUPPORTING INFORMATION

Additional Supporting Information may be found online in the supporting information tab for

this article. References unique to the Supporting Information appear only there.

Figure 1 Productivity of communities of co-selected plants (dashed lines, open circles) and

naïve communities (solid lines, closed circles). In communities of co-selected plants the

increase in productivity from monocultures to 2- and 4-species mixtures was stronger than in

naïve plant communities (see Table 1). Main panel: quadratic response curves, averaged over

four years. Points are predicted means and standard errors derived from a mixed model

including fixed-effects terms for species richness as 4-level factor and community-evolution

treatment and random-effects terms for plot, quadrat, half-quadrat and interactions of these

with year of harvest (see Methods). Inset panel: separate quadratic response curves for years,

with darker colors representing later years.

Figure 2 Relative productivity (% of mean of 8-species mixture) of communities of co-

selected plants (dashed lines, open circles) and naïve communities (solid lines, closed circles)

in monocultures and 2- and 4-species mixtures in **(a)** neutral soil (sterilized soil with neutral

inoculum) **(b)** native soil obtained by inoculation (sterilized soil with neutral inoculum and

inoculum of co-selected soil biota from original plots) and **(c)** native soil (unsterilized soil

with co-selected soil biota from original plots). Raw means and standard errors are shown

(for significances see Table S1).

Figure 3 Log ratio of productivity in communities of co-selected plants ($bm_{selected}$) and

productivity in naïve communities ($bm_{naïve}$) averaged over four years and three soil

treatments. In 8-species mixtures, productivity did not differ between communities of co-

selected and naïve plants (ratio=0). Especially in 2- and 4-species mixtures, but also in

monocultures, communities of co-selected plants produced more biomass than naïve

communities. Means and standard errors are shown. Raw data are plotted in the background.

Figure 4 Log-transformed species biomass ratios between co-selected and naïve plants. The majority of plant species attained greater aboveground biomass in communities of co-selected plants compared with naïve communities. The studied plant species belong to three different functional groups: grasses (white bars), herbs (light grey bars) and legumes (dark grey bars). Data are for each species averaged over four years, three soil treatments, four species-richness levels and species compositions within richness levels ($n = 32–352$). Three species with n < 32 were excluded from the analysis (*Anthriscus sylvestris*, *Campanula patula* and *Cardamine pratensis*). The stars represent *P*-values < 0.05 for species tested separately.

Figure 5 Within-species variation in plant height **(a)** and specific leaf area (SLA) **(b)** for

communities of co-selected plants and naïve communities at the end of the experiment in

2015 in neutral soil. In monocultures within-species variations in height/SLA (measured as

the within-species variance component in analysis of variance) were smaller/larger for co-

selected than for naïve plants and these differences decreased with increasing species

richness. Open circles and dashed line refer to communities of co-selected plants, closed

circles and solid line refer to naïve communities.

Table 1 Result of mixed-effects ANOVA for the aboveground biomass of the test

communities. Significant effects are shown in bold font.

Source of variation	nDf	dDF	F	P
Linear log-richness (lSR)	**1**	**43.6**	**21.47**	**< 0.001**
Quadratic log-richness (qSR)	1	43.7	1.059	0.309
Soil history (SH)	**2**	**89.6**	**19.64**	**< 0.001**
Plant history (PH)	**1**	**133.5**	**22.48**	**< 0.001**
lSR × PH	1	132.2	0.33	0.568
qSR × PH	**1**	**133.4**	**7.40**	**0.007**
Linear Harvest (linHar)	1	45.9	0.59	0.446
SH × linHar	**2**	**91.9**	**6.93**	**0.002**
PH × linHar	**1**	**140.0**	**5.489**	**0.021**

695 *Note*: nDF = numerator degrees of freedom, dDF = denominator degrees of freedom, F =

696 variance ratio, P = probability of type-I error.