Altering lipid droplet homeostasis affects *Coxiella burnetii* intracellular growth

Minal Mulye[1+], Brianne Zapata[1,2], and Stacey D. Gilk[1*]

[1] Department of Microbiology and Immunology, Indiana University School of Medicine,

Indianapolis, Indiana.

[2] Department of Biology, California State University, Dominguez Hills, Carson, California.

*Corresponding author: 635 Barnhill Rd, MS 420, Indianapolis, IN 46202. Telephone: 317-274-

8028. Fax: 317-278-3331. E-mail: sgilk@iupui.edu

+Current address: Marian University College of Osteopathic Medicine, Indianapolis, Indiana.

Running title: Lipid droplets during *Coxiella burnetii* infection.

ABSTRACT

Coxiella burnetii is an obligate intracellular bacterial pathogen and a causative agent of culture-

negative endocarditis. While *C. burnetii* initially infects alveolar macrophages, it has also been

found in lipid droplet (LD)-containing foamy macrophages in the cardiac valves of endocarditis

patients. In addition, transcriptional studies of *C. burnetii*-infected macrophages reported

differential regulation of the LD coat protein-encoding gene perilipin 2 *(plin-2)*. To further

investigate the relationship between LDs and *C. burnetii*, we compared LD numbers using

fluorescence microscopy in mock-infected and *C. burnetii*-infected alveolar macrophages. On

average, *C. burnetii*-infected macrophages contained twice as many LDs as mock-infected

macrophages. LD numbers increased as early as 24 hours post-infection, an effect reversed by

blocking *C. burnetii* protein synthesis. The observed LD accumulation was dependent on the *C.*

burnetii Type 4B Secretion System (T4BSS), a major virulence factor that manipulates host

cellular processes by secreting bacterial effector proteins into the host cell cytoplasm. To

determine the importance of LDs during *C. burnetii* infection, we manipulated LD homeostasis

and assessed *C. burnetii* intracellular growth. Surprisingly, blocking LD formation with the

pharmacological inhibitors triacsin C or T863, or knocking out acyl-CoA transferase-1 (acat-1)

in alveolar macrophages, increased *C. burnetii* growth at least 2-fold. Conversely, preventing LD

lipolysis by inhibiting adipose triglyceride lipase (ATGL) with atglistatin almost completely

blocked bacterial growth, suggesting LD breakdown is essential for *C. burnetii*. Together these

data suggest that maintenance of LD homeostasis, possibly via the *C. burnetii* T4BSS, is critical

for bacterial growth.

IMPORTANCE

Host neutral lipid storage organelles known as lipid droplets (LDs) serve as a source of energy,

nutrients, and signaling lipids. LDs are associated with infection of the intracellular bacterial

pathogen *Coxiella burnetii*, a significant cause of culture-negative endocarditis. While *C.*

burnetii was found in LD-rich foamy macrophages in endocarditis patients, little is known about

the host LD-*C. burnetii* relationship. We demonstrated *C. burnetii* Type 4B Secretion System

(T4BSS)-dependent LD accumulation in macrophages, suggesting a T4BSS-mediated regulation

of host LD homeostasis. Further, manipulating LD homeostasis significantly affected bacterial

growth, indicating LDs play an important role during *C. burnetii* infection. As *C. burnetii*

endocarditis has a 19% mortality rate even in treated patients, exploring the LD-*C. burnetii*

association might identify novel therapeutic targets.

INTRODUCTION

Lipid droplets (LDs) are dynamic cytoplasmic organelles which store cellular lipids in

eukaryotic cells. LDs are uniquely comprised of a phospholipid monolayer surrounding a

hydrophobic core of neutral lipids, primarily sterol esters and triacylglycerols (TAGs). LD

assembly begins with neutral lipid synthesis, where fatty acyl CoA synthetases generate long

chain fatty acids which are converted to sterol esters and triacyglycerols by acyl-CoA:cholesterol

acyltransferase (ACAT) and acyl-CoA:diacylglycerol acyltransferase (DGAT), respectively.

Progressive accumulation of neutral lipids in the ER leads to budding of the lipid ester globule

surrounded by the ER membrane cytoplasmic leaflet, thus forming LDs (1, 2). Inversely, adipose

triglyceride lipase (ATGL) (3) and hormone sensitive lipase (HSL) (4) mediate LD breakdown

and release of free cholesterol and fatty acids. Functionally, LDs serve as intracellular lipid

reservoirs for membrane synthesis or energy metabolism. In addition, LDs are linked to a range

of cellular functions including protein storage, protein degradation and signaling (2, 5).

LDs are emerging as important players during host-pathogen interactions. During

infection of host cells, Hepatitis C virus (HCV) (6) and Dengue virus (7) co-opt LDs as

platforms for viral assembly and replication. Even though blocking LD formation attenuates

HCV and Dengue virus replication *in vitro*, the importance of LDs during viral infection still

remains elusive (8). LD numbers increased in host cells during infection with several pathogens

including HCV (6) and Dengue virus (7), as well as the protozoan parasites *Trypanosoma cruzi*

(9), *Plasmodium berghei* (10)*, Toxoplasma gondii* (11), *Leishmania amazonensis* (12) and

Leishmania major (13). In addition, infection with the intracellular bacterial pathogens

Chlamydia spp. (14), *Mycobacterium* spp. (15-18), *Orientia tsutugamushi* (19), and *Salmonella*

typhimurium (20) also led to increased LD numbers. Since fatty acids released from host TAGs

and sterols can serve as carbon sources during infection (21), *C. trachomatis* (14, 22) and *M.*

tuberculosis (15) are proposed to use LD components as a major source of energy and nutrients.

Furthermore, in cells infected with *M. leprae* (23), *M. bovis* (16), *T. cruzi* (24), and *Leishmania*

infantum chagasi (25), LDs serve as a source of prostaglandin and leukotriene eicosanoids,

important signaling lipids which modulate inflammation and the immune response. These LD-

derived eicosanoids potentially favor intracellular pathogen survival by downregulating the

immune response (26).

LDs have been implicated during infection by *Coxiella burnetii*, a gram-negative

intracellular bacterium and the causative agent of human Q fever. Primarily spread through

aerosols, *C. burnetii* acute infection is characterized by a debilitating flu-like illness, while

chronic disease results in endocarditis. Although *in vitro* and *in vivo C. burnetii* can infect a wide

range of cells including epithelial cells and fibroblasts, the bacterium first infects alveolar

macrophages during natural infection. Inside the host cell, *C. burnetii* directs formation of a

specialized lysosome-like compartment called the parasitophorous vacuole (PV) which is

essential for *C. burnetii* survival. PV biogenesis requires the *C. burnetii* type 4B secretion system

(T4BSS), which secretes effector proteins into the host cell cytoplasm where they manipulate a

wide range of cellular processes. While not established to be T4BSS-dependent, *C. burnetii* is

thought to manipulate LDs and other components of host cell lipid metabolism (27-30). *C.*

burnetii-containing LD-filled foam cells were found in heart valves of an infected patient (31),

and LDs were observed in the *C. burnetii* PV lumen of infected human alveolar macrophages

(32). Further, two separate microarray analyses reported differential regulation of the LD coat

protein *plin-2* in *C. burnetii*-infected human macrophage-like cells (THP-1) (29, 30), suggesting

C. burnetii induced changes in host cell LDs. Intriguingly, siRNA depletion of the phospholipase

involved in LD breakdown, PNPLA2 (also known as ATGL), increased the number of *C.*

burnetii PVs in HeLa epithelial cells (33). In addition, treatment of monkey kidney epithelial

cells (Vero cells) with a broad spectrum antiviral molecule ST699 which localizes to host cell

LDs inhibited *C. burnetii* intracellular growth (34). Despite these observations, the importance of

LDs during *C. burnetii* infection is not known. In this study, we further examined the

relationship between host LDs and *C. burnetii*. We observed a T4BSS-dependent increase in LD

numbers in infected mouse alveolar and human monocyte-derived macrophage cell lines.

Furthermore, manipulation of LD homeostasis significantly altered *C. burnetii* intracellular

growth, thus strongly indicating that LDs play an important role during *C. burnetii* infection.

RESULTS

108 **LD-associated genes are upregulated in *C. burnetii*-infected alveolar macrophages.**

109 To examine the role of LDs in *C. burnetii* pathogenesis, we first analyzed the expression of LD

110 homeostasis-related genes in infected cells. As *C. burnetii* preferentially infects alveolar

111 macrophages during natural infection, we utilized a mouse alveolar macrophage cell line (MH-S)

112 previously established as a model for *C. burnetii* infection (35). Compared to mock-infected cells

113 at day 1 post-infection, *C. burnetii* infection upregulated the expression of genes *acat-1*, *acat-2*

114 and *fabp-4* by more than 1.2 fold (Figure 1). ACATs (acyl coA transferases) catalyze the

115 formation of sterol esters, while FABP-4 (fatty acid binding protein-4) facilitates fatty acid

116 transport for storage in LDs (36). Upregulation of LD-associated genes suggests LDs may

117 increase during *C. burnetii* infection.

118

119 ***C. burnetii* infection results in bacterial protein synthesis-dependent host cell LD**

120 **accumulation.**

121   Previously, at 36 and 72 hours post-infection two separate microarray analyses of *C.*

122 *burnetii*-infected THP-1 cells reported upregulation of the LD coat protein-encoding gene *plin-2*

123 (29, 30). Mahapatra et al. also reported upregulation of *fabp-4* (29). Together with our gene

124 expression analysis, this suggests LD formation may be altered during *C. burnetii* infection. To

125 test this possibility, we stained LDs for the coat protein PLIN2 and quantitated LD numbers per

126 cell in *C. burnetii*-infected cells by fluorescence microscopy. On an average, mock-infected had

127 less than 50 LDs per cell during a 4 day experiment, irrespective of the time point (Figure 2A).

128 In contrast, we observed a significant increase in LDs in infected cells, with an average of more

129 than 90 LDs/cell at 1, 2 and 4 days after *C. burnetii* infection (Figure 2A, gray circles). Notably,

130 LD accumulation occurred as early as 1 day post-infection, when the PV has not expanded and

the bacteria are not in log growth (Figure 2D-E). To determine the role of *C. burnetii* in host cell

LD accumulation, we blocked bacterial protein synthesis with chloramphenicol (29) and

quantitated LD numbers at indicated times. Chloramphenicol-treated *C. burnetii*-infected cells

contained 50 or less LDs per cell similar to mock-infected cells (Figure 2B and G). This result

suggests *C. burnetii* protein synthesis is required for LD accumulation.

Host cell LD accumulation is dependent on the *C. burnetii* Type 4B Secretion System

(T4BSS).

C. burnetii manipulates the host cell by secreting bacterial effector proteins through the

T4BSS into the host cytoplasm. To further decipher *C. burnetii's* role in LD accumulation, we

analyzed LD numbers at 1, 2, and 4 days after infection with a *C. burnetii* T4BSS *dotA* mutant

(37). While the wild-type *C. burnetii*-infected cells had more than 90 LDs per cell (Figure 2A,

closed circles; Figure 2E), T4BSS-mutant infected cells contained an average of 50 LDs, similar

to mock-infected cells (Figure 2A, triangles; Figure 2F). This suggests the LD accumulation in

C. burnetii-infected alveolar macrophages involves the *C. burnetii* T4BSS.

To confirm the T4BSS-dependent increase in LD accumulation, we analyzed LD

numbers in human macrophage-like cells (THP-1). When compared to mock- or T4BSS mutant-

infected cells, wild-type *C. burnetii*-infected THP-1 cells had increased LD numbers at 1 and 4

days post-infection (Figure 2C), similar to mouse alveolar macrophages. Interestingly, we did

not observe T4BSS-dependent LD accumulation at 2 days post-infection. Overall these results

demonstrate that *C. burnetii* induces LD accumulation in human and mouse macrophages

through a T4BSS-dependent process.

Blocking LD formation increases *C. burnetii* growth.

Given our finding that *C. burnetii* appears to actively manipulate host LDs through the

T4BSS, we next assessed the importance of LDs during *C. burnetii* infection. We first blocked

LD formation using triacsin C, a long chain fatty acyl CoA synthetase inhibitor (38). Compared

to vehicle control, triacsin C significantly reduced macrophage LDs, with <5 LDs per cell

(Figure 3A-B). We next treated macrophages with triacsin C during *C. burnetii* infection, and

quantitated bacterial growth using a fluorescent infectious focus-forming unit (FFU) assay. At

various times post-infection, we recovered bacteria from MH-S cells, replated bacteria onto a

monolayer of Vero cells, and incubated for 5 days. After staining for *C. burnetii*, we counted the

number of fluorescent foci, with 1 focus unit equivalent to 1 viable bacterium. Surprisingly,

compared to vehicle-treated cells, triacsin C treatment increased *C. burnetii* growth 5-fold at 4

days post-infection (Figure 3F).

To further confirm this finding, we used CRISPR/Cas-9 to knockout *acat-1* (Figure 3G),

a gene functionally responsible for sterol esterification in macrophages (39) and upregulated in

C. burnetii-infected cells (Figure 1). While *acat-1$^{-/-}$* LDs lack sterol esters, fluorescence

microscopy revealed similar number of LDs in wild-type and *acat-1$^{-/-}$* cells (Figure 3A,C).

Compared to wild-type cells, *C. burnetii* growth in *acat-1$^{-/-}$* cells increased 2-fold at 4 days post-

infection (Figure 3H), indicating that blocking sterol esterification favors *C. burnetii* growth. To

further deplete both TAG- and sterol ester-containing LDs, we treated *acat-1$^{-/-}$* cells with the

DGAT-1 inhibitor T863, which specifically blocks formation of TAGs (40). T863 treatment

significantly reduced LDs in *acat-1$^{-/-}$* macrophages, compared to untreated wild-type or *acat-1$^{-/-}$*

macrophages (Figure 3A, C, D). *C. burnetii* growth increased 2-fold in T863-treated *acat-1$^{-/-}$*

cells compared to vehicle-treated *acat-1$^{-/-}$* cells (Figure 3I), demonstrating that blocking both

TAG- and sterol ester-containing LDs improves *C. burnetii* growth.

Inhibiting LD breakdown blocks *C. burnetii* growth.

Because blocking LD formation appeared to benefit *C. burnetii,* we next examined *C.*

burnetii growth after inhibiting LD breakdown. When cells or tissues require fatty acids or

cholesterol, cytosolic lipases, including adipose triglyceride lipase (ATGL) (3) and hormone

sensitive lipase (HSL) (2, 4), hydrolyze TAGs and sterol esters stored in LDs. ATGL catalyzes

the initial step of lipolysis converting TAGs to DAGs whereas HSL hydrolyses DAGs to MAGs

(41). To block LD breakdown, we first inhibited HSL with a selective inhibitor CAY10499 (42).

HSL inhibitor-treated cells showed no significant difference in *C. burnetii* growth compared to

vehicle-treated cells (Figure 4A) suggesting conversion of DAGs to MAGs is not important for

bacterial intracellular growth.

To further study the importance of LD breakdown, we inhibited ATGL with the selective

and competitive inhibitor atglistatin, which binds the ATGL patatin-like phospholipase domain

(43). To eliminate the possibility of ATGL inhibiting a *C. burnetii* phospholipase, we first

measured viability of axenic *C. burnetii* cultures in the presence or absence of atglistatin (Figure

4B). Treatment for 4 days had no effect on axenic bacterial growth, indicating atglistatin does

not directly affect *C. burnetii*.

We next tested the effect of atglistatin on intracellular bacteria. After atglistatin treatment

of wild-type MH-S cells, we observed larger LDs, although the number did not significantly

increase (Figure 3E). Interestingly, *C. burnetii* intracellular growth in atglistatin-treated wild-

type MH-S cells decreased 5-fold, with essentially no growth (Figure 4C). Further, atglistatin-

treated *acat-1$^{-/-}$* cells, which contain TAG-rich LDs, also showed reduced bacterial growth

(Figure 4D) suggesting TAG hydrolysis to DAG is critical to support *C. burnetii* growth.

Together, these data demonstrate that blocking LD breakdown, particularly TAG to DAG

conversion significantly inhibits intracellular *C. burnetii* growth.

DISCUSSION

Several intracellular pathogens intercept host cell LDs to promote their growth. While

certain viruses utilize LDs as platforms for assembly, many bacteria and parasites facilitate

intracellular growth by exploiting LDs for nutrition. To understand the role of LDs during *C.*

burnetii infection, we assessed changes in LD numbers in *C. burnetii*-infected cells and

determined the effect of manipulating LD homeostasis on bacterial growth. Our studies revealed

a bacterial T4BSS-dependent increase in LD numbers in *C. burnetii*-infected macrophages.

Further, manipulating LD homeostasis significantly altered *C. burnetii* growth. While blocking

LD formation promoted *C. burnetii* growth, inhibiting LD breakdown, particularly hydrolysis of

TAGs to DAGs, dramatically decreased bacterial growth. Collectively, our data strongly suggest

LDs play an important role during *C. burnetii* intracellular growth.

LD formation requires sterol and fatty acid esterification by the enzymes ACAT and

DGAT, while LD lipolysis is catalyzed by HSL and ATGL. Once formed, LD maintenance

requires the LD-associated proteins PLINs. Previously, two separate microarray analyses

reported upregulation of *plin-2* in *C. burnetii*-infected human macrophage-like cells (THP-1)

(29, 30), suggesting *C. burnetii* induced changes in host cell LDs. Although we did not observe

changes in *plin-2* transcript levels, post-translational regulation of *plin-2* (44) could potentially

contribute to changes in LD numbers in infected macrophages. However, consistent with the

upregulation of LD-formation genes *acat* and *fabp-4,* our results demonstrated that LD numbers

increase in *C. burnetii*-infected mouse and human macrophages.

Notably, LD accumulation occurred as early as day 1 post-infection, when the PV has not

expanded and the bacteria are not in log growth. This suggests that LD accumulation is not a host

response to a large and growing PV but could be in response to bacterial ligands or due to a

process actively manipulated by *C. burnetii* itself. Previously, based on microarray analysis of *C.*

burnetii-infected cells where bacterial protein synthesis was blocked, Mahapatra *et al.* identified

36 host cell genes specifically regulated by *C. burnetii* proteins during early stages of infection.

These genes were predominantly involved in the innate immune response, cell death and

proliferation, vesicular trafficking, cytoskeletal organization and lipid homeostasis. Interestingly,

changes in *plin-2* expression level in infected cells was dependent on *C. burnetii* protein

synthesis (29) suggesting a role for bacterial proteins in LD accumulation. In agreement with this

data, we found that blocking *C. burnetii* protein synthesis prevented increased LD accumulation

in mouse alveolar macrophages at all analyzed times post-infection. These data together suggest

that *C. burnetii* actively manipulates host LDs at early stages of infection.

Beginning 1 hour post-infection in bone marrow-derived macrophages and 8 hours in

HeLa cells, *C. burnetii* secretes effector proteins via T4BSS (45) to modulate host cell functions

and promote its intracellular survival. Intriguingly, a *C. burnetii* T4BSS mutant did not increase

host cell LD numbers, suggesting a T4BSS-dependent increase in LD accumulation during wild-

type *C. burnetii* infection. The same result was observed in both mouse and human macrophages,

demonstrating that the *C. burnetii*-induced increase in LDs is independent of species and

dependent on the *C. burnetii* T4BSS. This finding suggests that one or more T4BSS effector

proteins may actively manipulate LDs, possibly by directly targeting proteins involved in LD

homeostasis. For example, the *Salmonella* Typhimurium type 3 secretion system (T3SS) effector

protein SseJ esterifies cholesterol and increases LD numbers when ectopically expressed in

epithelial and macrophage cells (20). The *C. trachomatis* secreted protein Lda3 localizes to the

LD surface and is involved in LD translocation into the *Chlamydia*-containing inclusion (22).

Thus far, none of the identified *C. burnetii* T4BSS secreted effector proteins localize to host LDs

or associate with proteins involved in LD homeostasis. It is also possible that rather than *de novo*

formation, increased LD numbers result from fission of preexisting LDs (46). As observed by

microscopy, the LD size in *C. burnetii*-infected macrophages appeared smaller than the mock- or

T4BSS mutant-infected macrophages. Smaller, more numerous LDs might result from *C.*

burnetii T4BSS-mediated fission of the existing LDs.

While *C. burnetii* T4BSS effector proteins might directly target LD homeostasis

pathways, LD accumulation may also be a host innate immune response. In other diseases, LD

accumulation occurs during the inflammatory response in macrophages in atherosclerotic lesions

(47), leukocytes from joints of patients with inflammatory arthritis (48), and eosinophils in

allergic inflammation (49). Thus, an innate immune response to the T4BSS apparatus or T4BSS

effector proteins may increase LD numbers in *C. burnetii*-infected macrophages. Additionally,

bystander response to bacterial components including TLR ligand LPS could result in increased

LD accumulation (16, 50). However, T4BSS mutant and protein-synthesis blocked *C. burnetii*

have similar surface ligands to wild-type bacterium but show no increase in LD accumulation

thus arguing against the contribution of *C. burnetii* LPS and other surface ligands. Overall, while

our data demonstrate that the *C. burnetii* T4BSS is involved in LD accumulation in both mouse

and human macrophages, the bacterial effector proteins and the specific LD processes targeted

remain unknown.

Besides T4BSS-dependent increase in LD accumulation, we found that manipulating LD

homeostasis, in particular fatty acids, significantly altered *C. burnetii* growth. Surprisingly,

blocking LD formation by pharmaceutical and genetic approaches, which increases the

availability of free fatty acids, led to improved *C. burnetii* fitness in macrophages. Conversely,

reducing fatty acid availability by inhibiting breakdown of LD-derived TAGs to DAGs blocked

C. burnetii growth. Intriguingly, DAG to MAG conversion did not affect bacterial growth,

suggesting that DAGs in particular are critical for *C. burnetii* infection. DAGs and other fatty

acids liberated during LD breakdown can be re-esterified or serve as signaling cofactors,

building blocks for membranes, or substrates for β-oxidation (2). Other bacterial pathogens are

known to target LDs as a source of lipids. For example, *M. tuberculosis* uses free fatty acids as a

source of energy and carbon (15, 51), while *C. trachomatis* is hypothesized to break down

bacterial inclusion-associated host LDs to provide lipids for bacterial growth (14). It not known

if free fatty acids or sterols liberated from LDs, either in the cytosol or possibly in PV lumen, are

used directly by *C. burnetii*.

In addition to serving as a source of free fatty acids and sterols, macrophage LDs are rich

in substrates and enzymes that generate prostaglandins and leukotrienes, which are arachidonic

acid-derived inflammatory lipid mediators (52, 53). In *M. leprae*-infected Schwann cells and *M.*

bovis BCG-infected macrophages, increased LD biogenesis correlates with increased production

of prostaglandin E2 (PGE2), linking LDs to the production of innate immune modulators (17,

54). PGE2 has a potential role in inhibiting TH1 responses important in clearance of intracellular

pathogens (24) including *C. burnetii* (55). Interestingly, elevated levels of PGE2 were observed

in *C. burnetii* endocarditis patients and linked to *C. burnetii*-mediated immunosuppression.

Koster *et al.* reported lymphocytes from chronic Q fever patients being unresponsive to *C.*

burnetii antigens, an effect reversed by PGE2 suppression with indomethacin (56). In addition,

after stimulation with *C. burnetii* antigens, monocytes from Q fever patients produced PGE2,

which in turn downregulated T lymphocyte-mediated IL-2 and IFNγ production. Interestingly,

PGE2 synthesis inhibitor Piroxicam reversed this downregulation of pro-inflammatory cytokine

production (57). Thus, while PGE2 appears to play a role in Q fever patients, the relationship

between *C. burnetii*-induced LDs and PGE2 production is not known. Considering that LD

breakdown can serve multiple functions, *C. burnetii* could use LDs either as a source of nutrients

or for production of lipid immune mediators like PGE2, which could then modulate the host cell

response to promote *C. burnetii* intracellular growth.

In summary, our data demonstrate that LD homeostasis is important for *C. burnetii*

intracellular survival. Because the *C. burnetii* T4BSS is involved in LD accumulation,

characterizing bacterial T4BSS effector proteins that target host LD homeostasis will help further

understand the role of LDs in *C. burnetii* pathogenesis.

MATERIALS AND METHODS

<u>Bacteria and mammalian cells</u>

C. burnetii Nine Mile Phase II (NMII; clone 4, RSA439) were purified from Vero cells (African

green monkey kidney epithelial cells, ATCC CCL-81; American Type Culture Collection,

Manassas, VA) and stored as previously described (58). For experiments examining T4BSS-

dependent accumulation of LDs, NMII and the *dotA* mutant (37) were grown for 4 days in

ACCM-2, washed twice with phosphate buffered saline (PBS) and stored as previously described

(59). Vero, mouse alveolar macrophages (MH-S; ATCC CRL-2019) and human monocytes

(THP-1; ATCC TIB-202) were maintained in RPMI (Roswell Park Memorial Institute) 1640

medium (Corning, New York, NY, USA) containing 10% fetal bovine serum (Atlanta

Biologicals, Norcross, GA, USA) at 37°C and 5% CO_2 and human embryonic kidney 293

(HEK293T; ATCC CRL-3216) in DMEM (Dulbecco's Modified Eagle Medium) (Corning, New

York, NY, USA) containing 10% fetal bovine serum at 37°C and 5% CO_2. THP-1 cells were

differentiated with 200 nM of phorbol 12-myristate 13-acetate (PMA) for 24 hours. PMA was

removed, and the cells rested for 48 hours prior to infection. The multiplicity of infection (MOI)

was optimized at 37°C and 5% CO_2 for each bacterial stock, cell type and infection condition for

a final infection of ~1 internalized bacterium/cell.

<u>Generating *acat-1*^{-/-} MH-S cell line</u>

The guide RNA sequence 5′TCGCGTCTCCATGGCTGCCC3′ to mouse *acat-1* was selected

using the optimized CRISPR design site crispr.mit.edu. Oligonucleotides were synthesized (IDT,

Coralville, IA, USA), annealed, and cloned into the lentiCRISPRv2 plasmid (a gift from Feng

Zhang, Addgene # 52961, Cambridge, MA, USA) (60), at the BsmBI restriction site to generate

plentiCRISPRv2-*acat-1*. To generate lentivirus, HEK293T cells were co-transfected with

plentiCRISPRv2-*acat-1* and packaging plasmids pVSVg (Addgene # 8454), pRSV-Rev

(Addgene # 12253), and pMDLg/pRRE (Addgene # 12251) using FuGENE6 reagent (Promega,

Madison, WI, USA). At 48 hours post-transfection, supernatant was collected, centrifuged at

3000xg, and then filtered with 0.45μm filter to remove cells and debris. Supernatant was

concentrated using the Lenti-X concentrator (Catalog # PT4421-2, Clontech, USA) and viral

RNA isolated using Viral RNA isolation kit (Catalog # 740956, Macherey-Nagel, Germany) to

determine viral titer using Lenti-X qRT-PCR titration kit (Catalog # PT4006-2, Clontech). Viral

titers were optimized for transduction of MH-S cells to generate stable *acat-1*^{-/-} cells.

$2x10^5$ MH-S cells were plated in a 6 well plate and transduced with 5.8×10^6 viral

particles/ml. 1 µg/ml puromycin was used for selection 48 hours post-transduction and continued

for 24 hours. The puromycin was then removed and the cells allowed to recover before isolating

individual clones by limiting dilution.

To confirm disruption of *acat-1*, clones were lysed in 2% SDS (Sigma-Aldrich, St. Louis,

MO, USA) for SDS-PAGE and immunoblotting with 1:1000 rabbit anti-mouse ACAT1-specific

antibody (Catalog # NBP189285, Novus Biologicals, Littleton, CO, USA) and 1:4000 GAPDH

loading control monoclonal antibody (Catalog # MA5-15738, ThermoFisher Scientific,

Waltham, MA, USA).

<u>Gene expression analysis of LD genes</u>

$2x10^5$ MH-S cells were infected in a 6 well plate. At day 1 post-infection, host cell RNA was

isolated using RNEasy mini kit (Catalog # 74104, Qiagen) and reverse transcribed to cDNA

using Superscript first strand synthesis system (Catalog # 11904-018, Invitrogen). Gene

expression analysis was performed using TaqMan array mouse lipid regulated genes (Catalog #

4415461, Applied Biosystems). Fold change was calculated compared to mock-infected cells in

6 independent experiments.

<u>LD quantitation</u>

$1x10^5$ MH-S cells were plated onto ibidi-treated channel µslide $VI^{0.4}$ ($3x10^3$ cells per channel;

Ibidi, Verona, WI) and allowed to adhere overnight. After infecting with *C. burnetii* for 1 hour,

cells were gently washed with phosphate buffered saline (PBS) to remove extracellular bacteria,

and incubated in 10% FBS-RPMI. For blocking bacterial protein synthesis, infected cells were

incubated with 3 µg/ml chloramphenicol containing 10% FBS-RPMI for the remainder of

experiment, replenishing media every 24 hours. At different times post-infection, infected cells

were fixed with 2.5% paraformaldehyde on ice for 15 min, then permeabilized/blocked for 15

min with 0.1% saponin and 1% bovine serum albumin (BSA) in PBS (saponin-BSA-PBS) and

stained with 1:1000 rabbit anti-mouse PLIN2 primary antibody (Catalog # PA1-16972,

ThermoFisher Scientific), 1:2000 guinea-pig anti-*C. burnetii* primary antibody (61) and 1:1000

rat anti-LAMP (Catalog # 553792, BD Biosciences) primary antibody in saponin-BSA-PBS for 1

hour. THP-1 cells were stained with 1:500 guinea-pig anti-human PLIN2 primary antibody

(Catalog # 20R-AP002, Fitzgerald Industries International, Acton, MA), 1:2000 rabbit anti-*C.*

burnetii primary antibody and 1:1000 rat anti-LAMP primary antibody in saponin-BSA-PBS for

1 hour. After three washes with PBS, cells were stained with 1:2000 AlexaFluor 488 anti-rabbit,

AlexaFluor 594 anti-guinea pig and AlexaFluor 647 anti-rat secondary antibodies (Invitrogen)

for 1 hour. ProLong Gold mount (Invitrogen) was added to the wells after washing with PBS and

slides visualized on a Leica inverted DMI6000B microscope (100X oil). The number of LDs per

cell was quantitated for 25 cells per condition in duplicate in each of three individual

experiments, with only bacteria-containing cells counted for *C. burnetii*-infected cells.

<u>Inhibitors</u>

Each LD homeostasis inhibitor used was diluted in DMSO based on manufacturer's instructions

and optimum inhibitor concentration was determined based on 100% host cell viability

determined by trypan blue staining, and changes in LD numbers per cell. The optimum

concentrations determined for each inhibitor was: Triacsin C (Enzo Life Sciences, Farmingdale,

NY, USA) – 10 µM, T863 (Sigma-Aldrich) – 10 µM, CAY10499 (Cayman Chemicals, Ann

Arbor, MI, USA) - 10 µM, Atglistatin (Cayman Chemicals) – 20 µM.

C. burnetii growth by fluorescent infectious focus-forming unit (FFU) assay.

To measure growth of *C. burnetii* in wild-type and *acat-1$^{-/-}$* MH-S cells, $5x10^4$ cells/well were

infected for 1 hour in a 48 well plate, washed with PBS, and then incubated with media

containing respective vehicle and inhibitors. At the indicated time points, the media was

removed and cells were incubated with sterile water for 5 min, pipetted up and down and the

lysate diluted 1:5 in 2% FBS-RPMI. Serial dilutions were added to 24 well plate containing

confluent monolayers of Vero cells, incubated for 5 days, fixed with methanol and stained with

rabbit anti-*C. burnetii* antibody as well as DAPI to confirm monolayer integrity. Four fields per

well were captured on an Evos automated microscope (ThermoFisher) with 4X objective and

fluorescent foci units were quantitated using ImageJ. Each experiment was done in duplicate.

Atglistatin-treatment of *C. burnetii* axenic cultures

To test bacterial sensitivity to atglistatin, ACCM-2 was inoculated at approximately $1x10^5$

bacteria/ml with *C. burnetii* NMII and grown for 3 days as previously described (59). Bacteria

(500 µl) were then incubated with DMSO or atglistatin in 24 well plates under normal *C.*

burnetii culture conditions. Media was replenished every 24 hours by centrifuging the

supernatant at 20000xg for 10 min, and bacterial pellet resuspended in new media containing

inhibitor. After 4 days, bacteria were diluted 1:10 in 2% FBS-RPMI and serial dilutions were

added to confluent Vero cell monolayers in a 96 well ibidi-treated µplate. At 5 days post-

infection, the plate was stained and fluorescent foci were determined as above. Each experiment

was done in duplicate.

<u>Statistical analysis</u>

Statistical analyses were performed using two-tailed paired t-test, ordinary one-way ANOVA or

two-way ANOVA with Tukey's or Bonferroni's multiple comparisons test in Prism (GraphPad

Software, Inc., La Jolla, CA).

FIGURE LEGENDS:

Figure 1: LD-associated genes are upregulated in *C. burnetii*-infected alveolar

macrophages.

MH-S macrophages were infected with wild-type *C. burnetii* and RNA was collected at day 1

post-infection. Gene expression analysis was performed using Applied Biosystems lipid-

regulated TaqMan Array. Fold change was calculated compared to mock-infected samples. Error

bars show the mean of 6 independent experiments +/- SEM * =$p<0.05$, *** =$p <0.001$ compared

to respective mock-infected cells as determined by two-tailed paired t-test.

Figure 2: *C. burnetii* infection leads to host cell LD accumulation and is dependent on

bacterial protein synthesis and the T4BSS.

Macrophages were infected with *C. burnetii* and at different times post-infection, cells were

stained for PLIN2 (LDs; green), *C. burnetii* (red) and nucleus (blue). LD number per cell were

quantitated by fluorescence microscopy.

A) LD numbers in wild-type *C. burnetii* and *dotA* mutant- infected MH-S macrophages.

B) LD numbers in MH-S macrophages infected with wild-type *C. burnetii* and treated with

chloramphenicol (3μg/ml).

C) LD numbers in wild-type *C. burnetii* and *dotA* mutant- infected THP-1 macrophage-like cells.

D) Images of LDs and bacteria in MH-S cells at day 1 post-infection imaged at 100X. Scale bar

= 10 μm. Error bars show the mean of 3 independent experiments +/- SEM * =p<0.05, ** =p

<0.01, *** =p <0.001 as determined by ordinary one-way ANOVA with Tukey post-hoc test.

Figure 3: Blocking LD formation increases *C. burnetii* growth.

Wild-type *C. burnetii* growth in infected MH-S cells treated with different inhibitors was

measured at 2 and 4 days post-infection by FFU assay.

A-E) Representative images for wild-type MH-S macrophages treated with inhibitors, fixed,

stained for PLIN2 (LDs; green) and *C. burnetii* (red) and imaged day 4 post-treatment at 100X.

Scale bar = 10 μm.

F) Growth while inhibiting LD formation with triacsin C (10 μM) in wild-type MH-S

macrophages. Error bars represent the mean of 4 independent experiments +/- SEM. **= p <0.01

compared to vehicle-treated cells as determined by two-way ANOVA with Bonferroni post-hoc

test.

G) ACAT1 protein expression in wild-type and *acat-1*[-/-] macrophages. Cell lysates were

immunoblotted and ACAT1 protein levels were compared with GAPDH as loading control.

H) *C. burnetii* growth in vehicle-treated wild-type and *acat-1*[-/-] MH-S macrophages and (I) T863-

treated *acat-1*[-/-] MH-S macrophages. Error bars represent the mean of at least 3 independent

experiments +/- SEM., * =p<0.05, *** =p <0.001 as determined by two-way ANOVA with

Bonferroni post-hoc test.

Figure 4: Inhibiting LD breakdown blocks *C. burnetii* growth.

Effect of HSL inhibitor CAY10499 and ATGL inhibitor atglistatin on viability of axenic and

intracellular wild-type *C. burnetii* was determined by FFU assay.

A) *C. burnetii* growth in CAY10499-treated (10 µM) and vehicle-treated wild-type MH-S

macrophages. Error bars represent the mean of 3 independent experiments +/- SEM. ns = not

significant compared to vehicle-treated cells as determined by two-way ANOVA with

Bonferroni post-hoc test.

B) Direct effect of atglistatin on wild-type *C. burnetii*. Atglistatin was added to axenic *C.*

burnetii cultures and bacterial viability was determined at day 4 using FFU assay. Error bars

represent the mean of 3 independent experiments +/- SEM. ns = not significant compared to

vehicle treatment as determined by ordinary one -way ANOVA with Tukey post-hoc test.

C) *C. burnetii* growth in atglistatin-treated (20 µM) and vehicle-treated wild-type MH-S

macrophages. Error bars represent the mean of 4 independent experiments +/- SEM. *= p <0.05,

****= p <0.0001 compared to vehicle-treated cells as determined by two-way ANOVA with

Bonferroni post-hoc test.

D) *C. burnetii* growth in atglistatin-treated (20 µM) and vehicle-treated *acat-1*[-/-] MH-S

macrophages. Error bars represent the mean of 4 independent experiments +/- SEM. **= p <0.01

compared to vehicle-treated cells as determined by two-way ANOVA with Bonferroni post-hoc

test.

ACKNOWLEDGMENTS

This research was supported by National Institutes of Health (AI21786; S.D.G.), American Heart Association (16POST27250157; M.M.) and National Institutes of Health (5R25GM07592; B.Z.). We thank Anna Justis, Tatiana Clemente and Rajshekar Gaji for critical reading of the manuscript and members of the IU Biology of Intracellular Pathogens Group for helpful suggestions.

We have no conflicts of interest to declare.

REFERENCES

1. **Pol A, Gross SP, Parton RG.** 2014. Review: biogenesis of the multifunctional lipid droplet: lipids, proteins, and sites. The Journal of Cell Biology **204:**635-646.
2. **Walther TC, Farese RV, Jr.** 2012. Lipid droplets and cellular lipid metabolism. Annual Review of Biochemistry **81:**687-714.
3. **Haemmerle G, Lass A, Zimmermann R, Gorkiewicz G, Meyer C, Rozman J, Heldmaier G, Maier R, Theussl C, Eder S, Kratky D, Wagner EF, Klingenspor M, Hoefler G, Zechner R.** 2006. Defective lipolysis and altered energy metabolism in mice lacking adipose triglyceride lipase. Science **312:**734-737.
4. **Vaughan M, Berger JE, Steinberg D.** 1964. Hormone-Sensitive Lipase and Monoglyceride Lipase Activities in Adipose Tissue. The Journal of biological chemistry **239:**401-409.
5. **Welte MA.** 2007. Proteins under new management: lipid droplets deliver. Trends in cell biology **17:**363-369.
6. **Filipe A, McLauchlan J.** 2015. Hepatitis C virus and lipid droplets: finding a niche. Trends in Molecular Medicine **21:**34-42.
7. **Samsa MM, Mondotte JA, Iglesias NG, Assuncao-Miranda I, Barbosa-Lima G, Da Poian AT, Bozza PT, Gamarnik AV.** 2009. Dengue virus capsid protein usurps lipid droplets for viral particle formation. PLoS Pathogens **5:**e1000632.
8. **Herker E, Ott M.** 2012. Emerging role of lipid droplets in host/pathogen interactions. Journal of Biological Chemistry **287:**2280-2287.
9. **D'Avila H, Freire-de-Lima CG, Roque NR, Teixeira L, Barja-Fidalgo C, Silva AR, Melo RC, Dosreis GA, Castro-Faria-Neto HC, Bozza PT.** 2011. Host cell lipid bodies triggered by *Trypanosoma cruzi* infection and enhanced by the uptake of apoptotic cells are associated with prostaglandin E(2) generation and increased parasite growth. Journal of Infectious Disease **204:**951-961.

505   10.   **Pulido-Mendez M, Finol HJ, Giron ME, Aguilar I.** 2006. Ultrastructural pathological
506   changes in mice kidney caused by *Plasmodium berghei* infection. Journal of
507   submicroscopic cytology and pathology **38:**143-148.
508   11.   **Charron AJ, Sibley LD.** 2002. Host cells: mobilizable lipid resources for the
509   intracellular parasite *Toxoplasma gondii*. Journal of cell science **115:**3049-3059.
510   12.   **Lecoeur H, Giraud E, Prevost MC, Milon G, Lang T.** 2013. Reprogramming neutral
511   lipid metabolism in mouse dendritic leucocytes hosting live *Leishmania amazonensis*
512   amastigotes. PLoS neglected tropical diseases **7:**e2276.
513   13.   **Rabhi S, Rabhi I, Trentin B, Piquemal D, Regnault B, Goyard S, Lang T,**
514   **Descoteaux A, Enninga J, Guizani-Tabbane L.** 2016. Lipid Droplet Formation, Their
515   Localization and Dynamics during *Leishmania major* Macrophage Infection. PLoS One
516   **11:**e0148640.
517   14.   **Kumar Y, Cocchiaro J, Valdivia RH.** 2006. The obligate intracellular pathogen
518   *Chlamydia trachomatis* targets host lipid droplets. Current biology **16:**1646-1651.
519   15.   **Daniel J, Maamar H, Deb C, Sirakova TD, Kolattukudy PE.** 2011. *Mycobacterium*
520   *tuberculosis* uses host triacylglycerol to accumulate lipid droplets and acquires a
521   dormancy-like phenotype in lipid-loaded macrophages. PLoS Pathogens **7:**e1002093.
522   16.   **D'Avila H, Melo RC, Parreira GG, Werneck-Barroso E, Castro-Faria-Neto HC,**
523   **Bozza PT.** 2006. *Mycobacterium bovis* bacillus Calmette-Guerin induces TLR2-mediated
524   formation of lipid bodies: intracellular domains for eicosanoid synthesis in vivo. Journal
525   of Immunology **176:**3087-3097.
526   17.   **D'Avila H, Roque NR, Cardoso RM, Castro-Faria-Neto HC, Melo RC, Bozza PT.**
527   2008. Neutrophils recruited to the site of *Mycobacterium bovis* BCG infection undergo
528   apoptosis and modulate lipid body biogenesis and prostaglandin E production by
529   macrophages. Cellular microbiology **10:**2589-2604.
530   18.   **Mattos KA, Lara FA, Oliveira VG, Rodrigues LS, D'Avila H, Melo RC, Manso PP,**
531   **Sarno EN, Bozza PT, Pessolani MC.** 2011. Modulation of lipid droplets by
532   *Mycobacterium leprae* in Schwann cells: a putative mechanism for host lipid acquisition
533   and bacterial survival in phagosomes. Cellular microbiology **13:**259-273.
534   19.   **Ogawa M, Fukasawa M, Satoh M, Hanada K, Saijo M, Uchiyama T, Ando S.** 2014.
535   The intracellular pathogen *Orientia tsutsugamushi* responsible for scrub typhus induces
536   lipid droplet formation in mouse fibroblasts. Microbes and Infection **16:**962-966.
537   20.   **Nawabi P, Catron DM, Haldar K.** 2008. Esterification of cholesterol by a type III
538   secretion effector during intracellular Salmonella infection. Molecular microbiology
539   **68:**173-185.
540   21.   **Barisch C, Soldati T.** 2017. Breaking fat! How mycobacteria and other intracellular
541   pathogens manipulate host lipid droplets. Biochimie **141:**54-61.
542   22.   **Cocchiaro JL, Kumar Y, Fischer ER, Hackstadt T, Valdivia RH.** 2008. Cytoplasmic
543   lipid droplets are translocated into the lumen of the *Chlamydia trachomatis*
544   parasitophorous vacuole. Proceedings of the National Academy of Sciences of the U S A
545   **105:**9379-9384.
546   23.   **Mattos KA, D'Avila H, Rodrigues LS, Oliveira VG, Sarno EN, Atella GC, Pereira**
547   **GM, Bozza PT, Pessolani MC.** 2010. Lipid droplet formation in leprosy: Toll-like
548   receptor-regulated organelles involved in eicosanoid formation and *Mycobacterium*
549   *leprae* pathogenesis. Journal of leukocyte biology **87:**371-384.

24. **Toledo DA, Roque NR, Teixeira L, Milan-Garces EA, Carneiro AB, Almeida MR,**
Andrade GF, Martins JS, Pinho RR, Freire-de-Lima CG, Bozza PT, D'Avila H,
Melo RC. 2016. Lipid Body Organelles within the Parasite *Trypanosoma cruzi*: A Role
for Intracellular Arachidonic Acid Metabolism. PLoS One **11:**e0160433.
25. **Araujo-Santos T, Rodriguez NE, Moura-Pontes S, Dixt UG, Abanades DR, Bozza**
PT, Wilson ME, Borges VM. 2014. Role of prostaglandin F2alpha production in lipid
bodies from *Leishmania infantum chagasi*: insights on virulence. Journal of Infectious
Disease **210:**1951-1961.
26. **Melo RC, Weller PF.** 2016. Lipid droplets in leukocytes: Organelles linked to
inflammatory responses. Experimental cell research **340:**193-197.
27. **Justis AV, Hansen B, Beare PA, King KB, Heinzen RA, Gilk SD.** 2016. Interactions
between the *Coxiella burnetii* parasitophorous vacuole and the endoplasmic reticulum
involve the host protein ORP1L. Cellular Microbiology doi:10.1111/cmi.12637.
28. **Martinez E, Allombert J, Cantet F, Lakhani A, Yandrapalli N, Neyret A, Norville**
IH, Favard C, Muriaux D, Bonazzi M. 2016. *Coxiella burnetii* effector CvpB
modulates phosphoinositide metabolism for optimal vacuole development. Proceedings
of the National Academy of Sciences of the United States of America **113:**E3260-3269.
29. **Mahapatra S, Ayoubi P, Shaw EI.** 2010. *Coxiella burnetii* Nine Mile II proteins
modulate gene expression of monocytic host cells during infection. BMC Microbiology
10:244.
30. **Ren Q, Robertson SJ, Howe D, Barrows LF, Heinzen RA.** 2003. Comparative DNA
microarray analysis of host cell transcriptional responses to infection by *Coxiella burnetii*
or *Chlamydia trachomatis*. Annals of the New York Academy of Sciences **990:**701-713.
31. **Brouqui P, Dumler JS, Raoult D.** 1994. Immunohistologic demonstration of *Coxiella*
burnetii in the valves of patients with Q fever endocarditis. The American journal of
medicine **97:**451-458.
32. **Graham JG, MacDonald LJ, Hussain SK, Sharma UM, Kurten RC, Voth DE.** 2013.
Virulent *Coxiella burnetii* pathotypes productively infect primary human alveolar
macrophages. Cellular Microbiology **15:**1012-1025.
33. **McDonough JA, Newton HJ, Klum S, Swiss R, Agaisse H, Roy CR.** 2013. Host
pathways important for *Coxiella burnetii* infection revealed by genome-wide RNA
interference screening. mBio **4:**e00606-00612.
34. **Sandoz KM, Valiant WG, Eriksen SG, Hruby DE, Allen RD, 3rd, Rockey DD.** 2014.
The broad-spectrum antiviral compound ST-669 restricts chlamydial inclusion
development and bacterial growth and localizes to host cell lipid droplets within treated
cells. Antimicrobial agents and chemotherapy **58:**3860-3866.
35. **Soltysiak KA, van Schaik EJ, Samuel JE.** 2015. Surfactant protein D binds to *Coxiella*
burnetii and results in a decrease in interactions with murine alveolar macrophages. PLoS
One **10:**e0136699.
36. **Storch J, Thumser AE.** 2000. The fatty acid transport function of fatty acid-binding
proteins. Biochimica et biophysica acta **1486:**28-44.
37. **Beare PA, Gilk SD, Larson CL, Hill J, Stead CM, Omsland A, Cockrell DC, Howe**
D, Voth DE, Heinzen RA. 2011. Dot/Icm type IVB secretion system requirements for
Coxiella burnetii growth in human macrophages. mBio **2:**e00175-00111.
38. **Igal RA, Wang P, Coleman RA.** 1997. Triacsin C blocks *de novo* synthesis of
glycerolipids and cholesterol esters but not recycling of fatty acid into phospholipid:

evidence for functionally separate pools of acyl-CoA. The Biochemical journal **324 (Pt**
2):529-534.

39.   **Rudel LL, Lee RG, Cockman TL.** 2001. Acyl coenzyme A: cholesterol acyltransferase
types 1 and 2: structure and function in atherosclerosis. Current opinion in lipidology
12:121-127.

40.   **Cao J, Zhou Y, Peng H, Huang X, Stahler S, Suri V, Qadri A, Gareski T, Jones J,**
Hahm S, Perreault M, McKew J, Shi M, Xu X, Tobin JF, Gimeno RE. 2011.
Targeting Acyl-CoA:diacylglycerol acyltransferase 1 (DGAT1) with small molecule
inhibitors for the treatment of metabolic diseases. The Journal of biological chemistry
286:41838-41851.

41.   **Zechner R, Zimmermann R, Eichmann TO, Kohlwein SD, Haemmerle G, Lass A,**
Madeo F. 2012. FAT SIGNALS--lipases and lipolysis in lipid metabolism and signaling.
Cell Metabolism **15:**279-291.

42.   **Manna PR, Cohen-Tannoudji J, Counis R, Garner CW, Huhtaniemi I, Kraemer**
FB, Stocco DM. 2013. Mechanisms of action of hormone-sensitive lipase in mouse
Leydig cells: its role in the regulation of the steroidogenic acute regulatory protein. The
Journal of Biological Chemistry **288:**8505-8518.

43.   **Mayer N, Schweiger M, Romauch M, Grabner GF, Eichmann TO, Fuchs E, Ivkovic**
J, Heier C, Mrak I, Lass A, Hofler G, Fledelius C, Zechner R, Zimmermann R,
Breinbauer R. 2013. Development of small-molecule inhibitors targeting adipose
triglyceride lipase. Nature chemical biology **9:**785-787.

44.   **Xu G, Sztalryd C, Lu X, Tansey JT, Gan J, Dorward H, Kimmel AR, Londos C.**
2005. Post-translational regulation of adipose differentiation-related protein by the
ubiquitin/proteasome pathway. The Journal of Biological Chemistry **280:**42841-42847.

45.   **Newton HJ, McDonough JA, Roy CR.** 2013. Effector protein translocation by the
Coxiella burnetii Dot/Icm type IV secretion system requires endocytic maturation of the
pathogen-occupied vacuole. PLoS One **8:**e54566.

46.   **Long AP, Manneschmidt AK, VerBrugge B, Dortch MR, Minkin SC, Prater KE,**
Biggerstaff JP, Dunlap JR, Dalhaimer P. 2012. Lipid droplet *de novo* formation and
fission are linked to the cell cycle in fission yeast. Traffic **13:**705-714.

47.   **Paul A, Chang BH, Li L, Yechoor VK, Chan L.** 2008. Deficiency of adipose
differentiation-related protein impairs foam cell formation and protects against
atherosclerosis. Circulation Research **102:**1492-1501.

48.   **Bozza PT, Payne JL, Morham SG, Langenbach R, Smithies O, Weller PF.** 1996.
Leukocyte lipid body formation and eicosanoid generation: cyclooxygenase-independent
inhibition by aspirin. PNAS **93:**11091-11096.

49.   **Vieira-de-Abreu A, Calheiros AS, Mesquita-Santos FP, Magalhaes ES, Mourao-Sa**
D, Castro-Faria-Neto HC, Bozza MT, Bandeira-Melo C, Bozza PT. 2011. Cross-talk
between macrophage migration inhibitory factor and eotaxin in allergic eosinophil
activation forms leukotriene C(4)-synthesizing lipid bodies. American Journal of
Respiratory Cell and Molecular Biology **44:**509-516.

50.   **Cao F, Castrillo A, Tontonoz P, Re F, Byrne GI.** 2007. *Chlamydia pneumoniae*--
induced macrophage foam cell formation is mediated by Toll-like receptor 2. Infection
and Immunity **75:**753-759.

51.   **Deb C, Daniel J, Sirakova TD, Abomoelak B, Dubey VS, Kolattukudy PE.** 2006. A
novel lipase belonging to the hormone-sensitive lipase family induced under starvation to

utilize stored triacylglycerol in *Mycobacterium tuberculosis*. The Journal of biological chemistry **281:**3866-3875.

52.   **Dvorak AM, Dvorak HF, Peters SP, Shulman ES, MacGlashan DW, Jr., Pyne K, Harvey VS, Galli SJ, Lichtenstein LM.** 1983. Lipid bodies: cytoplasmic organelles important to arachidonate metabolism in macrophages and mast cells. Journal of Immunology **131:**2965-2976.

53.   **Weller PF, Monahan-Earley RA, Dvorak HF, Dvorak AM.** 1991. Cytoplasmic lipid bodies of human eosinophils. Subcellular isolation and analysis of arachidonate incorporation. The American journal of pathology **138:**141-148.

54.   **Mattos KA, Oliveira VG, D'Avila H, Rodrigues LS, Pinheiro RO, Sarno EN, Pessolani MC, Bozza PT.** 2011. TLR6-driven lipid droplets in *Mycobacterium leprae*-infected Schwann cells: immunoinflammatory platforms associated with bacterial persistence. Journal of Immunology **187:**2548-2558.

55.   **Shannon JG, Heinzen RA.** 2009. Adaptive immunity to the obligate intracellular pathogen *Coxiella burnetii*. Immunologic Research **43:**138-148.

56.   **Koster FT, Williams JC, Goodwin JS.** 1985. Cellular immunity in Q fever: modulation of responsiveness by a suppressor T cell-monocyte circuit. Journal of Immunology **135:**1067-1072.

57.   **Izzo AA, Marmion BP.** 1993. Variation in interferon-gamma responses to *Coxiella burnetii* antigens with lymphocytes from vaccinated or naturally infected subjects. Clinical and Experimental Immunology **94:**507-515.

58.   **Cockrell DC, Beare PA, Fischer ER, Howe D, Heinzen RA.** 2008. A method for purifying obligate intracellular *Coxiella burnetii* that employs digitonin lysis of host cells. Journal of Microbiological Methods **72:**321-325.

59.   **Omsland A, Cockrell DC, Howe D, Fischer ER, Virtaneva K, Sturdevant DE, Porcella SF, Heinzen RA.** 2009. Host cell-free growth of the Q fever bacterium *Coxiella burnetii*. Proceedings of the National Academy of Sciences of the U S A **106:**4430-4434.

60.   **Sanjana NE, Shalem O, Zhang F.** 2014. Improved vectors and genome-wide libraries for CRISPR screening. Nature Methods **11:**783-784.

61.   **Coleman SA, Fischer ER, Howe D, Mead DJ, Heinzen RA.** 2004. Temporal analysis of *Coxiella burnetii* morphological differentiation. The Journal of Bacteriology **186:**7344-7352.

Figure 1

Figure 1: LD-associated genes are upregulated in *C. burnetii*-infected alveolar macrophages.
MH-S macrophages were infected with wild-type *C. burnetii* and RNA was collected at day 1 post-infection. Gene expression analysis was performed using Applied Biosystems lipid-regulated TaqMan Array. Fold change was calculated compared to mock-infected samples. Error bars show the mean of 6 independent experiments +/- SEM * =p<0.05, *** =p <0.001 compared to respective mock-infected cells as determined by two-tailed paired t-test.

Figure 2

Figure 2: *C. burnetii* infection leads to host cell LD accumulation and is dependent on bacterial protein synthesis and the T4BSS.

Macrophages were infected with *C. burnetii* and at different times post-infection, cells were stained for PLIN2 (LDs; green), *C. burnetii* (red) and nucleus (blue). LD number per cell were quantitated by fluorescence microscopy.

A) LD numbers in wild-type *C. burnetii* and *dotA* mutant- infected MH-S macrophages.

B) LD numbers in MH-S macrophages infected with wild-type *C. burnetii* and treated with chloramphenicol (3ug/ml).

C) LD numbers in wild-type *C. burnetii* and *dotA* mutant- infected THP-1 macrophage-like cells.

D) Images of LDs and bacteria in MH-S cells at day 1 post-infection imaged at 100X. Scale bar = 10 μm. Error bars show the mean of 3 independent experiments +/- SEM * =p<0.05, ** =p <0.01, *** =p <0.001 as determined by ordinary one-way ANOVA with Tukey post-hoc test.

Figure 3

Figure 3: Blocking LD formation increases *C. burnetii* growth.
Wild-type *C. burnetii* growth in infected MH-S cells treated with different inhibitors was measured at 2 and 4 days post-infection by FFU assay.

A-E) Representative images for wild-type MH-S macrophages treated with inhibitors, fixed, stained for PLIN2 (LDs; green) and *C. burnetii* (red) and imaged day 4 post-treatment at 100X. Scale bar = 10 μm.

F) Growth while inhibiting LD formation with triacsin C (10 μM) in wild-type MH-S macrophages. Error bars represent the mean of 4 independent experiments +/- SEM. **= p <0.01 compared to vehicle-treated cells as determined by two-way ANOVA with Bonferroni post-hoc test.

G) ACAT1 protein expression in wild-type and *acat-1*[-/-] macrophages. Cell lysates were immunoblotted and ACAT1 protein levels were compared with GAPDH as loading control.

H) *C. burnetii* growth in vehicle-treated wild-type and *acat-1*[-/-] MH-S macrophages and (I) T863-treated *acat-1*[-/-] MH-S macrophages. Error bars represent the mean of at least 3 independent experiments +/- SEM., * =p<0.05, *** =p <0.001 as determined by two-way ANOVA with Bonferroni post-hoc test.

Figure 4

Figure 4: Inhibiting LD breakdown blocks *C. burnetii* growth.
Effect of HSL inhibitor CAY10499 and ATGL inhibitor atglistatin on viability of axenic and intracellular wild-type *C. burnetii* was determined by FFU assay.
A) *C. burnetii* growth in CAY10499-treated (10 μM) and vehicle-treated wild-type MH-S macrophages. Error bars represent the mean of 3 independent experiments +/- SEM. ns = not significant compared to vehicle-treated cells as determined by two-way ANOVA with Bonferroni post-hoc test.
B) Direct effect of atglistatin on wild-type *C. burnetii*. Atglistatin was added to axenic *C. burnetii* cultures and bacterial viability was determined at day 4 using FFU assay. Error bars represent the mean of 3 independent experiments +/- SEM. ns = not significant compared to vehicle treatment as determined by ordinary one -way ANOVA with Tukey post-hoc test.
C) *C. burnetii* growth in atglistatin-treated (20 μM) and vehicle-treated wild-type MH-S macrophages. Error bars represent the mean of 4 independent experiments +/- SEM. *= p <0.05, ****= p <0.0001 compared to vehicle-treated cells as determined by two-way ANOVA with Bonferroni post-hoc test.
D) *C. burnetii* growth in atglistatin-treated (20 μM) and vehicle-treated *acat-1*[-/-] MH-S macrophages. Error bars represent the mean of 4 independent experiments +/- SEM. **= p <0.01 compared to vehicle-treated cells as determined by two-way ANOVA with Bonferroni post-hoc test.