Conflicting evolutionary histories of the mitochondrial and nuclear genomes in New World *Myotis*
bats

Roy N. Platt II[1], Brant C. Faircloth[2], Kevin A.M. Sullivan[1], Troy Kieran[3], Travis C. Glenn[3], Michael W.
Vandewege[1], Thomas E. Lee[4], Robert J. Baker[1], Richard D. Stevens[5], David A. Ray[1*]

[1] Department of Biological Sciences, Texas Tech University, Lubbock, USA
[2] Department of Biological Sciences and Museum of Natural Science, Louisiana State University, Baton
Rouge, USA
[3] Environmental Health Science, University of Georgia, Athens, USA
[4] Department of Biology, Abilene Christian University, Abilene, USA
[5] Natural Resource Management, Texas Tech University, Lubbock, USA

*Corresponding author: E-mail: david.4.ray@gmail.com

Abstract

The rapid diversification of *Myotis* bats into more than 100 species is one of the most extensive
mammalian radiations available for study. Efforts to understand relationships within *Myotis* have
primarily utilized mitochondrial markers and trees inferred from nuclear markers lacked resolution. Our
current understanding of relationships within *Myotis* is therefore biased towards a set of phylogenetic
markers that may not reflect the history of the nuclear genome. To resolve this, we sequenced the full
mitochondrial genomes of 37 representative *Myotis*, primarily from the New World, in conjunction with
targeted sequencing of 3,648 ultraconserved elements (UCEs). We inferred the phylogeny and explored
the effects of concatenation and summary phylogenetic methods, as well as combinations of markers
based on informativeness or levels of missing data, on our results. Of the 294 phylogenies generated
from the nuclear UCE data, all are significantly different from phylogenies inferred using mitochondrial
genomes. Even within the nuclear data, quartet frequencies indicate that around half of all UCE loci
conflict with the estimated species tree. Several factors can drive such conflict, including incomplete
lineage sorting, introgressive hybridization, or even phylogenetic error. Despite the degree of
discordance between nuclear UCE loci and the mitochondrial genome and among UCE loci themselves,
the most common nuclear topology is recovered in one quarter of all analyses with strong nodal
support. Based on these results, we re-examine the evolutionary history of *Myotis* to better understand
the phenomena driving their unique nuclear, mitochondrial, and biogeographic histories.

Keywords

summary tree methods, concatenation, Vespertilionidae, phylogenomics, UCE, ultraconserved elements,

reticulation, incomplete lineage sorting

The bat genus *Myotis* (Order Chiroptera, Family Vespertilionidae) comprises more than 100

species that originated during the last 10-15 million years (Stadelmann et al. 2007), making it one of the

most successful, extant, mammalian radiations. *Myotis* are distributed worldwide, excluding polar

regions, and generally share a similar ecological niche: aerial insectivory. *Myotis* species often exhibit

little morphological differentiation and, as a result, the rate of cryptic speciation within the genus is

thought to be high. For example, specimens identified as *M. nigricans* and *M. albescens* form multiple

paraphyletic lineages distributed throughout the phylogeny of Neotropical *Myotis* (Larsen et al. 2012).

Confounding matters, the morphological variation that exists is often a poor indicator of species-level

relationships. Early classifications of *Myotis* identified three major morphotypes (Findley 1972).

Subsequent phylogenetic analyses of the mitochondrial cytochrome-b (*cytb*) gene demonstrated

paraphyletic origins of each morphotype, suggesting frequent convergent evolution (Ruedi et al. 2001).

These same analyses demonstrated that geography was a better predictor of phylogenetic relationship

than morphology (Ruedi et al. 2001, Stadelmann et al. 2007).

The ability of mitochondrial markers to resolve a well-supported topology does not guarantee

that the mitochondrial tree represents the species tree (for examples see Willis et al. 2014, Li, Gang et

al. 2016, Leavitt et al. 2017). The lack of recombination and uniparental inheritance of the

mitochondrion means that it is transmitted as a single genetic unit that is susceptible to evolutionary

processes that may cause its history to diverge from the history of the species (Edwards et al. 2009). The

most widely accepted phylogenies of *Myotis* rely heavily on mitochondrial data and even phylogenies

containing nuclear data demonstrate an over reliance on mitochondrial markers for resolution, likely

swamping out potentially conflicting signals from the nuclear genome. For example alignments of the

nuclear *RAG*2 and mitochondrial *cytb* contained 162 and 560 variable characters, respectively

(Stadelmann et al. 2007). Despite these concerns, we find ourselves in a situation where our

understanding of evolutionary relationships, our basis for conservation strategies, and our

biogeographic hypotheses are all founded, almost entirely, on mitochondrial phylogenies.

Targeted sequencing of ultraconserved elements (UCEs; Faircloth et al. 2012) to collect

sequence data from thousands of loci across the nuclear genome has resolved a number of difficult

phylogenetic problems (for examples see Faircloth et al. 2012, McCormack et al. 2013, Green et al. 2014,

Faircloth et al. 2015, McGee et al. 2016). Here we used UCE baits to collect ~1.4 Mbp from ≥3,600

nuclear loci in addition to random shotgun sequencing to collect full mitochondrial genomes in 37 taxa,

primarily representing New World (NW) *Myotis*. Rather than analyzing each sequence data set once, or

a handful of times, we chose to use a range of sampling, partitioning, and inference methods to fully

explore phylogenetic tree space of the nuclear and mitochondrial genomes. We recovered 294 trees

representing 175 distinct topologies for the nuclear UCE data and 28 trees representing 14 distinct

mitochondrial topologies. Our results show that, despite the range of trees recovered from each marker

type, nuclear and mitochondrial markers occupy distinct regions in tree space. Given that the nuclear

and mitochondrial trees are distinct from one another it is necessary to reevaluate hypotheses made

based solely on the mitochondrial phylogeny.

RESULTS

78   We used targeted sequencing of UCEs in 37 individuals (Table 1) to collect sequence data from

79   3,648 nuclear loci which we assembled into concatenated alignments as large as 1.37 Mb. In addition,

80   we assembled mitochondrial genomes for most taxa from random shotgun sequencing data. We then

81   used the data to infer the phylogenetic history of New World *Myotis* using a range of locus sampling

82   strategies, alignment partitioning methods and phylogenetic inference methods.

83   *Mitochondrial assembly, alignment, and phylogenetic inference*

84   We generated an average of 5,498,440 paired reads per sample (min = 2,2370,68, max =

9,961,348) reads per sample of random DNA sequence data for mitochondrial genome assembly.

Assemblies varied in quality. Some were almost entirely complete while others were missing small

regions. We were not able to assemble through the control region for any of the samples. We found

three premature stop codons in mtDNA protein coding genes. Subsequent manual alignment and

validation suggested that these regions were miscalled by MitoBim, and we corrected the errors prior to

analysis. Sequence coverage of the mitochondrial genomes averaged 58x (range >1x - 297x).

91   The thirty-seven mitochondrial protein coding, rRNA and tRNA genes were concatenated into an

alignment of 15,520 bp containing 5,007 informative characters. rRNA and tRNA genes were

concatenated into an alignment containing 4,157 bp containing 862 parsimony informative characters.

Protein coding genes were concatenated into an alignment containing 11,363 bp or 3770 amino acids

(aa) and containing 4,145 or 509 parsimony informative characters, respectively. For the alignment

containing all protein coding, rRNA and tRNA genes, 30 samples were ≥ 90% complete, and alignments

for five samples were 68-84% complete. Only 21% and 50% of nucleotide positions were present in the

M. albescens[3] (TK 61766) and *M. levis* alignments, respectively.

99   When considering alignment type, partitioning strategy, and phylogenetic inference method we

analyzed the mitochondrial data 28 ways as described in the Methods. These analyses recovered 14

distinct topologies, the most common of which was recovered 6 times, all by analysis of the translated

protein coding genes (Figure 1A). Despite being the most common topology, bipartition frequencies and

clade probability values were lower than nucleotide based datasets, as would be expected with more

slowly evolving amino acid sequences. Analysis of RNA coding genes recovered two topologies

depending on the method of phylogenetic inference. The remainder of the analyses recovered various

trees without any pattern relating to partitioning scheme or inference method. A majority rule

consensus tree for all 28 analyses is shown in Figure 1C. The mitochondrial consensus tree reflects

previously reported mitochondrial phylogenies for *Myotis*. The New World *Myotis* are generally

monophyletic splitting into Nearctic and Neotropical clades with the Old World taxon, *M. brandtii*, sister

to the Neotropical clade. Ambiguity, represented by polytomies, is found only in terminal relationships.

UCE assembly and alignment, and phylogenetic inference

We averaged 3.29 million reads per sample after demultiplexing reads from the UCE-enriched,

sequencing pool. These reads were assembled into an average of 5,778 contigs per sample (min = 1,562

M. nyctor, max = 11,784 *M. nigricans* [3]). Recovery rates for UCE loci varied across taxa. Of the 5,500 loci

in the Amniote probe set, we successfully recovered 3,898 UCE loci, 3,648 loci from five or more

samples and 212 loci in all 37 samples (Table 2). On average, 3,332 UCE loci were recovered per sample,

ranging from 1,106 (*M. nyctor*) to 4,008 (*M. keaysi*). Repetitive sequences, identified via RepeatMasker

searches, were minimal, occupying less than 0.02% of sites across all UCE alignments.

In all, 294 different combinations of alignment, partitioning, and inference method were used to

identify 175 unique nuclear topologies. The most common tree was recovered 45 times (Figure 1B). All

nodes were highly-supported based on clade probability values and all but three nodes were highly

supported with maximum likelihood bootstrap replicates. A consensus tree (Figure 1D) of the 294

nuclear topologies resolves New World *Myotis* (and *M. brandtii*) as a single polytomy that excludes *M.

volans* and represents a departure from previous mitochondrial phylogenies, including the one

recovered herein. Comparison of the mitochondrial and nuclear consensus trees (Figures 1C and D)

reveals substantial differences in the relationships within the genus.

As noted above, we were concerned that phylogenetic error, sampling bias or other factors may

be driving the conflict between the mitochondrial and nuclear trees. We started by investigating the

effects of matrix composition (or completeness) on phylogenetic inference using UCE loci by generating

10 alignments having varying levels of matrix completeness. Nine alignments were composed of loci

with 15 to 95% (at 10% intervals) of taxa. The tenth alignment was composed of all loci which were

recovered in 100% of specimens examined. For example, in the '15%' matrix, all loci that were

represented in at least 15% of species ($n_{taxa} \geq 5$) were included in the alignment (n_{loci} = 3,648). In the

'100%' matrix, only loci which were identified in all species (n_{taxa} = 37) were included (n_{loci} = 212).

Alignments were partitioned using three schemes: unpartitioned, individually partitioned by locus, or

combined partitions. Alignment lengths, numbers of informative characters and number of partitions

identified by PartitionFinder are available in Table 2. The result was 10 alignments that were each

partitioned three ways. These were analyzed with Bayesian, maximum likelihood, and coalescent

methods using RaxML, ExaBayes, ASTRID, ASTRAL-II and SVDquartets as described in the Methods.

Bootstrap topologies stabilized in each alignment and partition combination within 150 replicates and all

Bayesian runs converged in less than ten thousand generations. Unfortunately, computational limits

forced us to abandon the individually partitioned, Bayesian analyses. In general, the same alignment

produced the same topology regardless of inference method or partitioning scheme with the only

exception being the terminal relationships of the *M. levis/M. albescens* clade in the combined vs.

unpartitioned Bayesian analysis of the 100% complete data matrix.

Trees were also generated from data matrices incorporating UCE loci of differing lengths

(Hosner et al. 2016). All 3,648 loci were grouped into ten bins based on locus length so that the first bin

contained the 365 shortest loci, the second bin contained the 366[th] to the 731[st], and so on. The number

of informative characters per bin ranged from 1,115 to 6,995 and the number of informative characters

was correlated with average locus length (Supplemental Figure 1). On average, only 2.6% of characters

in each bin were parsimony-informative (Supplemental Figure 1). Each of the ten length-based

alignments recovered slightly different topologies. Terminal relationships were generally stable across

analyses with the majority of differences between topologies found in the early bifurcations of the

ingroup (*Myotis*).

From the above results, we observed that, in general, longer alignments produced well resolved

topologies with significant nodal support regardless of the phylogenetic method or partitioning scheme

used. Analyses of smaller data sets were more likely to recover unique topologies.

Given the goal of fully exploring nuclear tree space and generating as many reasonable nuclear

UCE based topologies for comparison with the mitochondrial tree, we decided to randomly sample a

limited number of nuclear UCE loci to create small alignments that would be more likely to result in

unique topologies. We therefore randomly subsampled UCE loci to create 100 unique data sets of 365

loci. Loci were concatenated in each replicate data set and analyzed using maximum likelihood in

RAxML. Of the 100 alignments analyzed, 80 unique topologies were generated (mean Robinson-Foulds

distance = 4.3).

In addition to concatenation methods we explored phylogenetic tree space of the UCE data set

using species tree methods. Normalized quartet scores from ASTRAL-II (Mirarab et al. 2015) analyses

were consistent among all analyses, with scores ranging from 0.540 to 0.553, and the number of

induced quartet gene trees ranging from 7,745,739 (100% complete 212 gene trees) and 63,042,410

(15% 3648 loci). SVDquartets (Chifman et al. 2014) sampled all 66,045 quartets. On average, the total

weight of incompatible quartets was 2.84%. Similar to the concatenated analysis, we inferred

coalescent-based species from the same 100 subsamples of 365 loci described above. Despite being

generated from the same underlying data, summary and concatenation methods only recovered the

same tree in 1 of 100 attempts.

Finally, we used weighted and unweighted statistical binning to combine individual gene trees

into supergenes, estimate the supergene phylogeny, and then infer the species tree from the supergene

trees. The 3,648 loci were combined into 528 binned loci with 480 bins containing seven loci each and

48 bins containing six loci each. Binning methods increases the normalized quartet scores from an

average of 0.547 with gene trees to 0.672 for the binned-unweighted and 0.673 for the binned-weighted

supergene trees. Given the relatively even distribution of loci into bins the negligible difference in

quartet/species tree discordance between the unweighted and weighted supergene trees is expected.

Binning methods have been shown to recover incorrect species trees under coalescent methods by

altering gene tree frequencies. In addition, it is possible that support for incorrect trees can actually

increase with the number of input gene trees (Liu et al. 2015). With these concerns in mind, both

binning methods recovered the same topology which was the most common nuclear topology observed

across all analyses (Figure 1C).

Topological comparisons

All mitochondrial and nuclear trees are available in Supplemental File 1. When visualizing all

topologies in tree space, nuclear trees co-localized and were distinct from mitochondrial topologies

(Figure 2A). Comparisons of Robinson-Foulds symmetrical differences show 26 symmetrical differences

between the most similar nuclear and mitochondrial trees (Figure 2B). For comparison only 9 of 43,070

and 0 of 378 pairwise nuclear vs. nuclear and mitochondrial vs. mitochondrial tree comparisons, Figures

2C and D, respectively, exhibited more than 26 symmetrical differences.

Of the 45 analyses that recovered the most frequently observed topology (Figure 1D), 38 were

Bayesian and RAxML searches that varied by matrix completeness and partitioning scheme. The fact that

these analyses recovered the same topology is expected given that they are not independent. For

example, a RAxML analysis of the 25% complete data set uses 1.26 Mb of the 1.38 Mb of data present in

the 15% complete matrix, sharing 91% of the data. Analyses that directly varied the alignments and/or

sampled less data (e.g. randomly sampling loci) were more likely to generate unique topologies than the

nested analyses described above. Of the 200 analyses that randomly sampled UCE loci 164 unique

topologies were observed. This implies that when analyses of large data sets produce well-resolved

trees with significant nodal support, sampling smaller portions of the data, may provide a mechanism

for creating phylogenetic uncertainty not represented by typical tree scoring metrics.

DISCUSSION

We analyzed 3,648 UCE loci and mitochondrial genomes of 37 *Myotis* bats and outgroups. The

mitochondrial and nuclear UCE phylogenies recovered distinct topologies (Figure 1), whether comparing

the most commonly recovered or consensus topologies. Previous work with UCE loci demonstrated that

support for deep divergences varied based on the number of loci examined (McCormack et al. 2013).

Further, bootstrap replicates and clade probability values can be inaccurate metrics of nodal support

(Douady et al. 2003, Hedtke et al. 2006). We used varied the input data and phylogenetic parameters to

produce a range of reasonable nuclear and mitochondrial topologies that may be more useful than

overreliance on a tree resulting from a one or two analyses. Rather than rejecting concordance between

the two data types from a single analysis we took steps to analyze both data sets to generate as many

viable topologies as possible, possibly revealing hidden overlap between marker types that are masked

by phylogenetic error.

We recovered 175 and 14 unique nuclear and mitochondrial topologies from 294 and 28

analyses using multiple methodologies, sampling strategies, and phylogenetic parameters. By varying

our phylogenetic methods we show that recovering a consistent topology is difficult even when using

the same marker type, thus making it difficult to determine the "best" topology. Rather than choosing

single trees to represent the nuclear and mitochondrial data, we use consensus trees to represent the

ambiguity that is present through all analyses along with example phylogenies conforming to the most

common topology (Figure 1). Despite our efforts to identify as many plausible, distinct nuclear and

mitochondrial trees as possible we were unable recover overlapping topologies between marker types

suggesting that nuclear UCE loci and the mitochondrial genomes of *Myotis* have distinct evolutionary

histories.

A comparison of topologies in tree space illustrates the differenced in trees generated for each

data set (Figure 2A). Intra-marker discordance, as measured by differences in nuclear vs. nuclear or

mitochondrial vs. mitochondrial topologies is low. Pairwise comparisons of nuclear topologies contain

an average of 8.3 symmetrical differences, mitochondrial comparisons contain an average of 10.8
differences. By contrast, pairwise differences between nuclear and mitochondrial topologies contain an
average of 33.3 symmetrical differences. The two most similar nuclear and mitochondrial topologies
contain 26 symmetrical differences. Of the 43,072 pairwise comparisons of nuclear topologies, only 27
contain more than 26 symmetrical differences. Likewise of the 378 pairwise comparisons of
mitochondrial topologies only 1 contains more than 26 symmetrical differences. These observations
indicate that discordance between marker types (Figure 2B) is much greater than within marker type
(Figure 2C and 2D). All but the most divergent nuclear topologies are more similar to each other than
any nuclear vs. mitochondrial comparison.

Conflict between mitochondrial and nuclear data may be driven by error in phylogenetic
estimation or may reflect genuine discordance between the two marker types (Degnan et al. 2009,
Huang, Huateng et al. 2010). We relied on multiple tree-inference methods (e.g. summary vs.
concatenation), manipulated phylogenetic parameters (e.g. partitioning strategy), and sampling criteria
(e.g. loci sampled in all taxa) to minimize the impacts of phylogenetic error on the data sets. In most
cases, varying parameters or methodologies generated unique topologies with minor differences. In
most cases unique topologies resulted from rearrangements of a few terminal taxa. For example,
placement of *M. volans* and *M. brandtii* was often either sister to the remaining New World *Myotis* or as
an early bifurcation between the Nearctic and Neotropical clades. *M. vivesi* was at times found as sister
to the clade containing *M. lucifugus*, *occultus* and *fortidens* or as sister to the clade containing the
neotropical *Myotis*.

Summary methods failed to recover the most common nuclear topology (Figure 1D) except
when loci were binned prior to gene tree estimation. Summary methods also tended to recover more
unique topologies than concatenation methods when analyzing data from the same gene(s). For
example, the random sample analyses recovered 80 unique topologies using concatenation (RAxML) and
89 unique topologies with summary methods (ASTRAL-II). This likely has to do with the limited number
of informative characters per locus and by extension limited phylogenetic signal per gene tree
(Supplemental Figure 1). In these instances, limited phylogenetic signal per gene would likely lead to
increased opportunity for phylogenetic error in gene tree estimation. Further supporting this idea,
binning of compatible UCE loci may have indirectly increased phylogenetic signal resulting in the same
topology that many of the concatenation analyses recovered. No other summary/coalescent method
recovered this topology.

Previous studies that used primarily mitochondrial data recovered effectively the same
relationships among *Myotis* as our mitochondrial analyses (Ruedi et al. 2001, Stadelmann et al. 2007,
Roehrs et al. 2010, Larsen et al. 2012, Ruedi et al. 2013, Haynie et al. 2016). Thus, we are confident that
the mitochondrial phylogeny we recovered here, and by others, reflects the true mitochondrial tree.
However, the mitochondrial topology may not adequately reflect the species history, particularly when
considering the factors that cause incongruence between nuclear and mitochondrial gene trees. Causes
of conflicting gene trees include horizontal transfer, gene duplication, introgressive hybridization, and
incomplete lineage sorting. The rapid radiation of this clade as well as other biological factors suggests
that some of these phenomena are more likely to have influenced the *Myotis* radiation than others.

Horizontal transfer of genes is thought to be rare in eukaryotes, but, vespertilionids in general
(Thomas et al. 2011, Platt et al. 2014, Platt et al. 2016), and *Myotis* in particular (Pritham et al. 2007, Ray
et al. 2007, Ray et al. 2008, Pagan et al. 2010), have experienced horizontal transfer of DNA transposons.
These events would not be reflected in our phylogeny because repetitive sequences were removed prior
to phylogenetic analyses. More generally, gene duplications could create conflicting signal among
individual UCE markers (ex. comparing non-orthologous UCE loci), but the number of gene duplication
events would have to be very high to impact enough of the 3,648 UCE loci to confound the
mitochondrial and nuclear phylogenies. Further ruling out gene duplication events as the dominant
cause of conflicting phylogenetic signal is the fact that such events are likely depressed in *Myotis* as
evidenced by their smaller genome size (~2.2 Gb) and trend towards DNA loss (Kapusta et al. 2017)
combined with low rates of paralogy in UCEs general (Derti et al. 2006).

Introgressive hybridization and reticulation could significantly influence the phylogenies of
Myotis in a way that leads to conflicting signal between the nuclear and mitochondrial genomes (Sota
2002, Good et al. 2015). Hybridization in bats may be relatively common given their propensity to
swarm at cave entrances for breeding purposes. In European *Myotis*, swarming has allowed for high
degrees of hybridization between *M. brandtii, M. mystacinus, and M. alcathoe* (Bogdanowicz et al.
2012). Further, *M. evotis, thysanodes,* and *keeni* all experienced historical gene flow during their
divergence (Carstens et al. 2010, Morales et al. 2016). We could also explain the differences between
the mitochondrial and nuclear UCE phylogenies if *Myotis* experienced extensive incomplete lineage
sorting during their radiation. Two factors influence the rate of lineage sorting, the fixation rate and the
speciation rate (Hudson et al. 2002). Increasing the time to fixation and/or decreasing the amount of
time between cladogenic events will increase the likelihood of incomplete lineage sorting. *Myotis* are
generally long lived species (Dzeverin 2008) and underwent a rapid radiation between 5-10 MYA (Lack et

al. 2010), suggesting that *Myotis* species are likely to experience higher levels of lineage sorting. The

importance of these events - introgressive hybridization and incomplete lineage sorting - in driving the

differences between the mitochondrial and nuclear phylogenies should be further explored.

Evolutionary history of Myotis

Our understanding of relationships within *Myotis* is heavily biased by mitochondrial data

because nuclear markers were harder to collect and produced fewer informative sites (Ruedi et al. 2001,

Stadelmann et al. 2007, Lack et al. 2010, Roehrs et al. 2010, Larsen et al. 2012, Ruedi et al. 2013, Haynie

et al. 2016) or nuclear phylogenies with large numbers of makers contained limited numbers of taxa

(Platt et al. 2015). Our UCE-based results indicate that nuclear trees vary substantially from the

mitochondrial tree. Given that the nuclear and mitochondrial trees are different, we find it necessary to

re-evaluate *Myotis* in the context of the nuclear data.

Paraphyly of *M. nigricans* and *M. albescens* was inferred from previous mitochondrial

phylogenies. Larsen et al. (2012) identified a minimum of four and potentially twelve lineages in *M.*

albescens and *M. nigricans*. Our sampling included three *M. albescens* and two *M. nigricans*, compared

to Larsen's 17 and 29 samples. Despite different mitochondrial and nuclear topologies overall, our

mitochondrial and nuclear phylogeny recovered the same paraphyletic clade of three *M. albescens*

samples and *M. levis*. Close relationships between these taxa was found in previous work and is

expected. More importantly we did not find that *M. albescens* was paraphyletic across much of

neotropical *Myotis*. We also found that *M. nigricans* is monphlyletic in the nuclear tree, but paraphyletic

in the mitochondrial tree. These results from *M. nigricans* and *M. albescens* are interesting but further

inference is limited due to low sample sizes for these taxa.

The original subgeneric taxonomy of *Myotis* was based on three morphotypes that were later

shown to be the result of convergent evolution (Ruedi et al. 2001). If lineage-sorting affected the

mitochondrial phylogeny, it is possible that the morphotypes truly are monophyletic. However,

superimposing the previous subgeneric/morphological classification onto the species tree shows an

interspersed distribution of morphotypes throughout the most common and consensus nuclear

topologies (Figures 1A and B). Many strongly supported terminal relationships link species with different

morphotypes. Based on these results, it appears that the three major morphotypes in *Myotis* are indeed

a result of convergent evolution, as suggested by previous work (Ruedi et al. 2001, Stadelmann et al.

2007) despite the conflicting mitochondrial and nuclear phylogenies recovered

Among the more dramatic differences between the nuclear and mitochondrial topologies is the

placement of *M. volans* and *M. brandtii* as sister to all New World taxa by the nuclear data. Our

mitochondrial analyses place *M. volans* within a Nearctic clade and *M. brandtii* directly in-between the

Nearctic and Neotropical bifurcations (Figure 1A and C) as has been previously reported (Stadelmann et

al. 2007). By contrast the consensus UCE tree (Figure 1D) places *M. brandtii*, a species distributed

throughout the Old World, within a polytomy at the base of the New World *Myotis* radiation and *M.*

volans sister to all New World *Myotis* (including *M. brandtii*). The placement of *M. brandtii* in the nuclear

UCE consensus tree does not necessarily contradict the mitochondrial phylogeny since it is an

unresolved polytomy. However, it is worth noting that the most commonly recovered nuclear topology

(Figure 1B) places *M. brandtii* sister to all New World *Myotis* (excluding *M. volans*). This relationship

would more closely affiliate *M. brandtii* with other Old World taxa. The placement of *M. volans* as sister

to all New World taxa (including *M. brandtii*) in the most common nuclear tree is a significant departure

from previous work and, at first glance, does not make as much sense in a biogeographic framework. *M.*

volans is distributed across western and northwestern North America as far as far north as Alaska. *M.*

brandtii is distributed across much of Northern Europe and into the extreme eastern regions of Siberia.

The key may lie in understanding the biogeography and phylogenetics of a third species, *M. gracilis*, a

species that we were unfortunately unable to include.

M. gracilis, along with *M. brandtii*, are the only two *Myotis* geographically distributed in the Old

World, but phylogenetically affiliated with the New World *Myotis* (Stadelmann et al. 2007). If future

work verifies the sister relationship between *M. brandtii* and *M. gracilis*, then we can envision a scenario

where *M. gracilis, M. brandtii,* and *M. volans* are the result of cladegenic events that occurred during

the transition of *Myotis* from the Old World to the New World. It is important to remember that this

interpretation relies on a fairly dramatic departure from the currently accepted mitochondrial

relationships of *M. volans* (represented here by a single sample) to other *Myotis* species and abandons

the nuclear UCE consensus tree for the most frequently recovered UCE topology. In addition, our

taxonomic sampling excludes *Myotis* species from the OW, Ethiopian clade, the sister clade to New

World *Myotis* (Ruedi et al. 2001, Stadelmann et al. 2007, Lack et al. 2010, Ruedi et al. 2013). Taxa from

the Ethiopian clade are needed to properly root the New World *Myotis* clade and understand its

biogeographic origins. With these caveats in mind, the hypothesis presented here should be viewed as

highly speculative. Increasing the number of *Myotis* lineages sampled will shed additional light on this

hypothesis.

Other taxa with conflicting positions between marker types include *M. lucifugus* + *M. occultus*,

M. fortidens, and *M. vivesi*. In general, these relationships are characterized by very short branches

(Figure 1D) and are the most likely to be affected by incomplete lineage sorting or limited phylogenetic

information. This could explain the strong support with the mitochondrial tree compared to the nuclear
species tree, while allowing for a number of nuclear loci to disagree with the species tree, as well.

There are a number of monophyletic groups identified with nuclear data (Figure 1A) that share
general morphologies. For example, all of the long-eared bats (*septentrionalis*, *auriculus*, *evotis*,
thysanodes and *keenii*) represent a monophyletic group of higher elevation, forest-dwelling species that
glean insects off of surfaces (Fitch et al. 1979, O'Farrell et al. 1980, Warner 1982, Manning et al. 1989,
Caceres et al. 2000). The group represented by *fortidens*, *lucifugus* and *occultus* represent a relatively
long-haired form of *Myotis*. While having a distinct dental formula, *fortidens* was historically described
as a subspecies of M. *lucifugus* (Miller Jr et al. 1928) and *occultus* has alternately represented its own
species or been considered a subspecies of *lucifugus* (Hollister 1909, Valdez et al. 1999, Piaggio et al.
2002). The clade consisting of *keaysi*, *oxyotus*, *ruber*, *riparius*, and *diminutus* represents a neotropical
group of primarily woolly-haired bats (LaVal 1973). None of these relationships are monophyletic in the
consensus or most common mitochondrial topologies. If the mitochondrial genome has been subjected
to phenomena that obscure the true species tree then these species groups, along with their
synapomorphic morphological features, can be reevaluated.

Conclusions

Relationships within *Myotis*, which until now have relied heavily on mitochondrial data, served
as the basis for species identification (Puechmaille et al. 2012), evolutionary hypotheses (Simões et al.
2007), and even conservation recommendations (Boyles et al. 2007). Previous studies using nuclear data
have largely been uninformative or utilized too few samples to draw definitive conclusions. Trees
estimated from ~3,650 nuclear loci and 295 phylogenetic analyses recovered 175 topologies, none of
which are congruent with the mitochondrial phylogeny of *Myotis*. Conflict between the mitochondrial
and nuclear trees as well as among individual nuclear loci suggest that the *Myotis* radiation may have
been accompanied by high levels of incomplete lineage sorting and possible hybridization. Rather than
placing emphasis on the mitochondrial tree, it may be more appropriate to consider it for what it really
is: a single gene on par with a single UCE locus, albeit one with many more phylogenetically informative
characters. If true, then the mitochondrial genome is as likely to reflect the true species tree as any
single UCE locus chosen at random. Phenomena such as lineage sorting, reticulation, and introgression
have likely influenced the genomes of *Myotis* and should be accounted for in subsequent work. It is
possible that the *Myotis* radiation is more accurately reflected as a hard polytomy or a phylogenetic
network rather than a strictly bifurcating phylogeny.

 MATERIALS AND METHODS:

Taxon Selection

Taxa were selected to span the major phylogenetic break points with emphasis on the Nearctic

and Neotropical bifurcation as recovered in previous mitochondrial phylogenies (Stadelmann et al. 2007,

Ruedi et al. 2013). In addition, multiple individuals morphologically identified as *M. nigricans* and *M.*

albescens were included to test for paraphyly as suggested by Larsen et al. (2012). Three Old World

species of *Myotis* and the outgroup, *E. fuscus,* were included to root phylogenetic analyses. All field

identifications were confirmed from voucher museum specimens. Information for all specimens

examined is available in Table 1.

Library preparation, sequencing, and processing

Genomic DNA was extracted from 33 samples using either a Qiagen DNEasy extraction kit or a

phenol-chloroform/ethanol precipitation. DNA was fragmented using the Bioruptor UCD-300 sonication

device (Diagenode, Denville, NJ, USA). Libraries were prepared using the Kapa Library Preparation Kit

KR0453-v2.13 (Kapa Biosystems, Wilmington, MA, USA) following the manufacturer's instructions with

five minor modifications. First, we used half volume reactions. Second, subsequent to end repair, we

added Sera-Mag Speedbeads (Thermo-Scientific, Waltham, MA, USA; prepared according to (Glenn et al.

2016)) at a ratio of 2.86:1 for end repair cleanup. Third, we ligated universal iTru y-yoke adapters (Glenn

et al. 2016) onto the genomic DNA. Fourth, following adapter ligation, we performed one post-ligation

cleanup followed by Dual-SPRI size selection using 55 µL of speedbead buffer (22.5mM PEG, 1M NaCl)

and 25 µL of Speedbeads. Finally, we performed a PCR at 95 °C for 45 sec, then 14 cycles of 98 °C for 30

sec, 60 °C for 30 sec, 72 °C for 30sec, then 72 °C for a 5 minute final extension and a 12 °C hold using

iTru5 and iTru7 primers to produce Illumina TruSeqHT compatible libraries (Glenn et al. 2016).

Libraries were quantified on a Qubit 2.0 (Life Technologies) and 83 ng from each library was

added to create 5 pools of 6 or 7 libraries each. We then split the pools in two. One subsample was

enriched for UCE loci, the other was not. UCE loci in the enriched library pools were captured using

Tetrapods 5K version 1 baits from MYcroarray (Ann Arbor, MI, USA) following their MYbaits protocol v.

2.3.1 with overnight incubations (Faircloth et al. 2012). Enriched libraries were quantified with a Qubit

and pooled with other unrelated samples prior to sequencing on an Illumina HiSeq 3000 to produce

paired-end reads of ≤ 151 bases. The unenriched samples were sequenced on a separate run using a

single lane of Illumina HiSeq 2500. All samples were demultiplexed with Illumina's fastq2bcl software.

Reads were quality filtered by removing any potential adapter sequence and trimming read ends once

the average Phred quality over a four base window score dropped below 20 using the Fastx toolkit

(Gordon et al. 2010).

Assembly, annotation, and phylogenetic analysis of the mitochondrial genome

Raw reads from the unenriched libraries were used to generate mitochondrial genomes via

MitoBim (Hahn et al. 2013). This program used MIRA (Chevreux et al. 1999) to map reads to a *M.*

brandtii reference genome (Genbank accession number KT210199.1). Alternative methods of

mitochondrial genome assembly were used when MitoBim assembly failed. These taxa include *M.*

albescens (TK61766), *M. albescens* (TK 101723), *M. albescens* (RDS 7889), *M. fortidens*, M. *keeni*, M.

melanorhinus, *M. nigricans* (QCAZ 9601), *M. septentrionalis*, *M. simus*, *M. velifer*, and *M. volans*. For

these samples, we first identified reads that were mitochondrial in origin using BLAST searches against

the *M. brandtii* mitochondrial genome (KT210199.1). Those reads were assembled using Trinity v2.2.0

with the '–single' option to treat reads as unpaired. For taxa where we used NCBI genome assemblies to

recover UCE loci *in silico* mitochondrial genomes from genbank were used to in the mitochondrial

analyses GenBank as follows: *M. brandtii* (KT210199.1), *E. fuscus* (KF111725.1), *M. lucifugus*

(KP273591.1), and *M. davidii* (KM233172.1).

Once assembled, each mitogenome was annotated via MITOS (Bernt et al. 2013). Annotated

genes were manually validated via BLAST to confirm sequence identity and length. Protein coding genes

were checked for stop codons using EMBOSS's transeq program (Rice et al. 2000). When a stop codon

was found, we used the raw reads to verify the sequence. We used BWA v0.7.12 (Li, Heng et al. 2009) to

align the reads to the Mitobim assembled mitogenome to verify base calls from Mitobim. The protein

coding rRNA and tRNA genes from each assembly were aligned using MUSCLE and concatenated into

three different alignments containing only protein coding genes, only rRNA and tRNA genes, or all

protein coding and RNA genes. A fourth alignment of all protein coding genes was translated to amino

acids and concatenated into a single alignment.

Several different partitioning schemes were examined for each of the mitochondrial alignments.

Alignments were either partitioned by gene, by codon, or by gene type (rRNA and tRNA vs. protein

coding). Genes were partitioned individually except in the instances where two genes overlapped. These

regions were partitioned separately from the individual genes resulting in three partitions for the two

genes: a partition for gene A, a partition for gene B, and a partition for the overlapping nucleotides of

gene A and B.

Assembly, and phylogenetic analysis of nuclear UCEs

Quality filtered raw sequence reads from the UCE-enriched libraries were assembled into

contigs using the Trinity assembler (Grabherr et al. 2011) with a minimum kmer coverage of 2. We used

Phyluce to identify those assembled contigs that were UCE loci. We also harvested UCE loci from

Eptesicus fuscus (GCA_000308155.1), *Myotis brandtii* (GCA_000412655.1), *M. davidii*

(GCA_000327345.1), and *M. lucifugus* (GCF_000147115.1) genome assemblies using the Phyluce

package (Faircloth 2016). Once extracted from Trinity and genome assemblies, we aligned all UCE loci

MAFFT (Katoh et al. 2002) and trimmed the aligned data with gBlocks (Castresana 2000). Repetitive

sequences (i. e. transposable elements) in each alignment were identified with RepeatMasker and

trimmed where found.

Each alignment was analyzed using three different partitioning schemes. Unpartitioned

alignments were simply concatenated UCE loci treated as a single unit. These alignments are referred to

herein as "unpartitioned". Fully partitioned alignments were concatenated alignments mitochondrial

genes that were partitioned by locus. These alignments are referred to herein as "locus partitioned".

Finally, PartitionFinder v1.1.1 (Lanfear et al. 2012) was used to combine individual loci into an optimal

partitioning scheme. The combined partitioning schemes for each alignment were identified with

PartitionFinder v1.1.1 (Lanfear et al. 2012). Rather than searching for best-fit substitution models for

each locus or partition, the GTR+Γ model of sequence evolution was assigned to all loci (Darriba et al.

2015) except in the case of amino acid alignments where the MtMam model was used. Initial trees for

PartitionFinder were generated using RAxML v7.4.1 (Stamatakis 2006) with linked branch lengths.

Partitioning schemes were heuristically searched using the hcluster algorithm. Partitioning schemes

were chosen using the Bayeisan information criterion.

Finally, each alignment and partitioning scheme was analyzed using Bayesian inference and

maximum likelihood phylogenetic methods. Bayesian phylogenies were generated with the MPI version

of ExaBayes (v1.4.1) using 4 independent runs of 4 chains each. ExaBayes runs were terminated after 10

million generations only if the average standard deviation of split frequencies was less than 0.01. The

first 25% of samples were discarded after which every 1,000[th] generation was sampled. Proper sampling,

post burn-in was inspected via Tracer v1.6. (Rambaut et al. 2014) and effective sample sizes greater than

200 were considered acceptable. Posterior probability values greater than 95% were considered to be

significant. RaxML (v8.1.3) was used to estimate and score the maximum likelihood phylogeny with the

rapid bootstrapping option and 10,000 bootstrap replicates. We define strongly supported bipartitions

as those present in 95-100% of bootstrap replicates and moderately supported bipartitions are present

in 85-95% of bipartitions (Wiens et al. 2008). All mitochondrial analyses are described in Table 3.

Alignment types. – Different combinations of UCE loci were used to create unique matrices

based on 1) locus distribution, 2) locus length, or 3) random locus sampling. The first set of alignments

(locus distribution) was determined by the number of taxa represented in the UCE alignments

(phyluce_align_get_only_loci_with_min_taxa; Faircloth 2016), or degree of completeness. Matrices

were constructed using loci which were present in 100% (number of specimens = 37), 95% (n = 35), 85%

(n = 31), 75% (n = 27), 65% (n = 24), 55% (n = 20), 45% (n = 16), 35% (n = 12), 25% (n = 9), and 15% (n = 5)

of specimens examined. These 10 groups were non-exclusive, so a locus that was assembled in all

specimens (100% complete) would also be included with loci present in only 55% of specimens. On the

other hand, a locus found in only 55% of specimens would not be included in the 100% complete data

set. Each set of UCE alignments was concatenated using phyluce_align_format_nexus_files_for_raxml

and a nexus character block was created using the phyluce_align_format_nexus_files_for_raxml –

charsets option. These datasets then served as the basis for downstream phylogenetic analyses. For

example, when a partitioning methodology (discussed below) was tested, it was performed for each of

the 100%, 95%, 85%, etc. alignments. In addition to partitioning schemes, the effect of missing data was

examined using Bayesian and maximum likelihood methods.

The second alignment criteria combined UCE loci of similar sizes. Pervious coalescent analyses of

UCE data showed that sub-sampling the most informative loci can result in different topologies

(Meiklejohn et al. 2016). Rather than using coalescent based analyses we used concatenation of UCE loci

to identify different topologies based on length. Under these assumptions, UCE loci were sorted into ten

groups based on their length and the predicted correlation between length and number of informative

characters was confirmed (Supplemental Figure 1). UCE loci in the same size cohort were combined into

a single alignment, partitioned by locus, and analyzed with RAxML using the methods described below.

The final set of alignments was generated by random sampling of UCE loci. In large phylogenetic

analyses, systematic error can result in highly supported, but incorrect topologies due to compounding

of non-phylogenetic signal (Rodríguez-Ezpeleta et al. 2007). By randomly reducing the dataset and

replicating the maximum likelihood analyses, we can reduce the potential effects of compounding error.

Roughly 10% of the dataset, 365 loci, were randomly sampled, concatenated, and partitioned by locus to

create 100 new alignments, which were then analyzed with RAxML using the methods described below.

Partitioning strategies. –Alignments were analyzed using three different partitioning schemes –

single, locus, and combined- similar to the mitochondrial partitioning schemes described above.

Unpartitioned alignments were simply concatenated UCE loci treated as a single genetic unit. Rather

than searching for best-fit substitution models for each UCE locus or partition, the GTR+Γ model of

sequence evolution was assigned to all loci (Darriba et al. 2015). Initial trees for PartitionFinder were

generated using RAxML v7.4.1 (Stamatakis 2006) with linked branch lengths. Partitioning schemes were

heuristically searched using the hcluster algorithm and the best scheme was chosen using the Bayesian

information criterion.

Inference methods. –Phylogenetic trees were generated with three different phylogenetic

inference methods across five different inference implementations including concatenation and

summary tree methods.

Maximum likelihood trees were inferred for each alignment and partitioning combination using

RAxML v8.1.3 (Aberer et al. 2014). The best scoring (lowest -lnL) tree from each dataset was identified

from 100 random starting trees and bootstrapped 100 times using the GTR+Γ in both cases. The

autoMRE function in RAxML v8.1.3 was used to determine the need for additional bootstrap replicates

beyond the initial 100 (Pattengale et al. 2009). A stopping criterion was set *a priori* if the weighted

Robinson-Foulds distance was less than 5% in 95% of random permutations of computed bootstrap

replicates (Pattengale et al. 2009). If necessary, an additional 100 bootstrap replicates were computed

until the convergence stopping criteria were met. Finally, bipartition frequencies of bootstrap replicates

were drawn onto the best scoring tree from the initial RAxML searches for each of the respective data

sets. Alignments based on UCE length or randomly sampled were analyzed using slightly different

methods. In both cases UCE loci were partitioned individually and nodal support was calculated using

100 bootstrap replicates using the RAxML fast-bootstrapping option. For all maximum likelihood

analyses, we define strongly supported bipartitions as those present in 95-100% of bootstrap replicates

and moderately supported bipartitions are present in 85-95% of bipartitions (Wiens et al. 2008).

Bayesian analyses were conducted using ExaBayes v1.4.1 (Aberer et al. 2014). For all Bayesian

analyses four independent runs of four chains each were run in parallel for a minimum of one million

generations sampling every thousandth generation and applying a GTR+Γ substitution model for each

partition. After one million generations, analyses continued until the standard deviation of the split

frequency between chains was less than 0.01. The "-M 3" option was used to reduce the memory

footprint of all ExaBayes runs. Proper sampling, post burn-in, was inspected via Tracer v1.6. (Rambaut et

al. 2014). Effective sample sizes greater than 200 were considered acceptable. Posterior probability

values greater than 95% were considered significant. An extended majority rule consensus tree was

created from all trees after the first 25% of trees were discarded using TreeAnnotator v1.7.0 (Rambaut

et al. 2013) and parameter estimates across all runs were calculated with Tracer v1.6 (Rambaut et al.

2014).

Species trees were calculated from gene trees for individual UCE loci recovered in five or more

taxa using the GTR+Γ substitution model and fast bootstrapping (-f a) option in RAxML and 1,000

bootstrap replicates. In general, gene trees were classified based on the degree of completeness (i.e.

number of taxa represented) similar to the way we treated individuals as described above.

Species trees were estimated and bootstrapped using three different programs. ASTRAL-II v4.10

(Mirarab et al. 2015) was used to build a summary tree. Support values for bipartitions in the tree were

generated from 100 bootstrap replicates using site as well as site and locus resampling (Seo 2008).

Species trees were estimated from ASTRID v1.4 (Vachaspati et al. 2015) using bionj and bootstrapped

for 100 replicates. SVDquartets (Chifman et al. 2014), as implemented in PAUP v4.0a150 (Swofford

2003), was used to estimate a species trees from a random subset of 200,000 quartets and 1,000

bootstrap replicates.

UCE loci are relatively short markers with few informative characters from which to build gene

trees. Errors in gene tree estimation may reduce the accuracy of summary methods and phylogenetic

inference (Liu et al. 2009, Leaché et al. 2011, DeGiorgio et al. 2014, Mirarab et al. 2016). We used

weighted and unweighted statistical binning to combine gene into compatible supergenes, increasing

the number of informative informative characters per "locus" and reducing the phylogenetic error

(Mirarab et al. 2014, Bayzid et al. 2015). The gene trees used for the summary tree methods described

above were used rather than re-estimating trees. Bifurcations supported by more than 50% of the

bootstrap replicates were retained for each gene tree. Alignments from compatible trees were

concatenated into a single supergene alignment. Trees for supergenes were estimated using RAxML. The

best trees for each supergene, as defined by log likelihood score, were retained from 500 searches.

Bipartition support was estimated from 500 bootstrap replicates. For all analyses, the GTR+Γ model of

substitution was used and each gene in the supergene alignment was partitioned separately. The

resulting supertrees were then used for species tree estimation using ASTRAL-II. For the unweighted

analysis, all supertrees were included in the pool of trees. For the weighted analysis, supertrees were

weighted according to the number of genes combined in the supergene alignment. For example, if a

supergene was a composite of six genes, the supertree was present 6 times compared to a composite of

five genes which would be represented only five times. Support for the weighted and unweighted

species trees was estimated by site and site and locus re-sampling (Seo 2008) for 100 bootstrap

replicates in ASTRAL-II.

Topological comparisons−

Trees recovered from all analyses were compared to each other in order to quantify the

differences between topologies. Branch lengths have different meanings based on the type of analysis.

For example, ASTRAL-II branch lengths are representative of coalescent units. ASTRID doesn't even

calculate branch lengths. For accurate tree comparisons, branch lengths were stripped from all trees.

Pairwise unweighted Robinson-Foulds distances were calculated among all trees. Robinson-Foulds

distances were transformed into two dimensions using the stochastic CCA algorithm for nonlinear

dimension reduction in TreeScaper v1.09 (Huang, Wen et al. 2016). Coordinates were then visualized in

R using hexagonal binning in the hexbin library v1.27.1 (Lewin-Koh 2011). Nuclear and mitochondrial

50% majority rule consensus trees were generated with PAUP v4.0a150 (Swofford 2003).

Acknowledgments:

We would like to thank the following museums, collection managers, and collaborators for the

tissue loans necessary to complete this work: Joseph Cook (Museum of Southwestern Biology), Museum

of Vertebrate Zoology, Manuel Ruedi (Natural History Museum of Geneva), Santiago Burneo (Pontificia

Universidad Catolica del Ecuador Museo de Zoologia), Heath Garner, Robert Bradley and Caleb Phillips

(Texas Tech University Natural Science Research Laboratory), Link Olson (University of Alaska Museum

of the North), Cody Thompson and Priscilla Tucker (University of Michigan Museum of Zoology). In

addition, we would like to thank the Texas Tech HPCC (http://www.depts.ttu.edu/hpcc/) for providing

the computational resources necessary to complete this project. This work was supported by the

National Science Foundation, DEB-1355176. Additional support was provided by College of Arts and

Sciences at Texas Tech University. All sequence data is available at NCBI's Short Read Archive

(SRP095250). UCE contig assemblies are available at (PENDING ACCESSION NUMBER). Supplemental files

include: a nexus file containing all estimated species trees (supFile-all_trees_2017-06-07.nexus.txt), and

tables listing each of the phylogenetic analyses. An archived file of individual UCE alignments in fasta

format (uceLociAlignments.tgz), and an archived file containing all individual UCE gene trees in newick

format (UceGeneTrees.tgz) are available through DRYAD (PENDING ACCESSION NUMBER).

Figure and Table Captions

Figure 1. Comparison of nuclear and mitochondrial phylogenetic trees of *Myotis*. The most common

mitochondrial (A) and nuclear (B) topologies. The mitochondrial topology arises from Bayesian and

maximum likelihood analysis of the amino acid sequences portioned based on the PartitonFinder

recommendations. The nuclear topology is from the 55% complete data matrix partitioned based on the

recommendations of PartitionFinder. For clarity, bootstrap values greater than 95 and clade probability

values greater than 0.95 have been omitted from the nuclear consensus tree. Majority-rule consensus

trees from 28 mitochondrial (C) and 294 nuclear (D) topologies. Values above the branches in the

consensus trees (C and D) are bipartition frequencies for that clade across all nuclear or mitochondrial

topologies. Conflicting tips between data types (consensus nuclear vs. consensus mitochondrial) are

indicated with red lines between the topologies. Biogeographic regions are color coded, as are

subgeneric classifications based on morphotypes, as defined by Findley (1972). Species with more than

one sample are designated with a superscript that is referenced in Table 1. Specimens derived from

whole genome alignments are designated with a superscript "G".

Figure 2. Comparison of mitochondrial and nuclear topologies. (A) All trees recovered from

mitochondrial and nuclear analyses were visualized in tree space using multidimensional scaling of

Robinson-Foulds distances. Nuclear trees (green outlines) were clustered separately from mitochondrial

trees (blue outlines). Robinson-Foulds distances between nuclear and mitochondrial trees (B), nuclear

vs. nuclear trees (C) and mitochondrial vs. mitochondrial trees (D) indicate the overall distinctions

between the nuclear and mitochondrial topologies.

Table 1. Specimens examined. Collection abbreviations: Museum of Southwestern Biology (MSB),

Museum of Vertebrate Zoology (MVZ), Natural History Museum of Geneva (MHNG), Pontificia

Universidad Catolica del Ecuador Museo de Zoologia (QCAZ), Texas Tech University Natural Science

Research Laboratory (TK), University of Alaska Museum of the North (UAM), University of Michigan

Museum of Zoology (UMMZ).

Table 2. General alignment information. For a subset of analyses a series of alignments were generated

based on the number of taxa per locus. Thirty-seven taxa were examined so an alignment with all 37

taxa was considered 100% complete. Parsimony-informative characters make up a small portion of the

total alignment. The number of partitions in the combined partitioning scheme was calculated with

PartitionFinder.

Sup Fig 1. Phylogenetic content of UCE loci based on length. The length of a UCE locus is correlated with

the number of phylogenetically informative characters. UCE loci were sorted by length and ten bins of

alignments were created so that the shortest loci were combined into one alignment, the next shortest

set of loci were combined … etc. Parsimony informative characters made up a minor part of each

alignment.

Sup File 1 - supFile-all_trees_2017-06-07.nexus.txt – Trees generated from all analyses in nexus format.

Literature Cited

Aberer A. J., Kobert K., Stamatakis A. 2014. ExaBayes: Massively parallel Bayesian tree inference for the
whole-genome era. Mol Biol Evol. 31:2553-2556.
Bayzid M. S., Mirarab S., Boussau B., Warnow T. 2015. Weighted statistical binning: Enabling statistically
consistent genome-scale phylogenetic analyses. PLoS ONE. 10:e0129183.
Bernt M., Donath A., Jühling F., Externbrink F., Florentz C., Fritzsch G., Pütz J., Middendorf M., Stadler P.
F. 2013. MITOS: Improved de novo metazoan mitochondrial genome annotation. Mol Phylogenet Evol.
69:313-319.
Bogdanowicz W., Piksa K., Tereba A. 2012. Hybridization hotspots at bat swarming sites. PLoS ONE.
7:e53334.
Boyles J. G., Storm J. J. 2007. The perils of picky eating: Dietary breadth is related to extinction risk in
insectivorous bats. PLOS ONE. 2:e672.
Caceres M. C., Barclay R. M. R. 2000. *Myotis septentrionalis*. Mammalian Species:1-4.
Carstens B. C., Dewey T. A. 2010. Species delimitation using a combined coalescent and information-
theoretic approach: An example from North American *Myotis* bats. Syst Biol. 59:400-414.
Castresana J. 2000. Selection of conserved blocks from multiple alignments for their use in phylogenetic
analysis. Mol Biol Evol. 17:540-552.
Chevreux B., Wetter T., Suhai S. 1999. Genome sequence assembly using trace signals and additional
sequence information. Proceedings of the German Conference on Bioinformatics (GCB).
Chifman J., Kubatko L. 2014. Quartet inference from SNP data under the coalescent model.
Bioinformatics. 30:3317-3324.
Darriba D., Posada D. 2015. The impact of partitioning on phylogenomic accuracy. bioRxiv.
https://doi.org/10.1101/023978.
DeGiorgio M., Degnan J. H. 2014. Robustness to divergence time underestimation when inferring species
trees from estimated gene trees. Syst Biol. 63:66-82.
Degnan J. H., Rosenberg N. A. 2009. Gene tree discordance, phylogenetic inference and the multispecies
coalescent. Trends Ecol Evol. 24:332-340.
Derti A., Roth F. P., Church G. M., Wu C. t. 2006. Mammalian ultraconserved elements are strongly
depleted among segmental duplications and copy number variants. Nat Genet. 38:1216-1220.
Douady C. J., Delsuc F., Boucher Y., Doolittle W. F., Douzery E. J. P. 2003. Comparison of Bayesian and
maximum likelihood bootstrap measures of phylogenetic reliability. Mol Biol Evol. 20:248-254.
Dzeverin I. 2008. The stasis and possible patterns of selection in evolution of a group of related species
from the bat genus *Myotis* (Chiroptera, Vespertilionidae). J Mamm Evol. 15:123-142.
Edwards S., Bensch S. 2009. Looking forwards or looking backwards in avian phylogeography? A
comment on Zink and Carrowclough 2008. Mol Ecol. 18:2930-2933.
Faircloth B. C. 2016. PHYLUCE is a software package for the analysis of conserved genomic loci.
Bioinformatics. 32:786-788.
Faircloth B. C., Branstetter M. G., White N. D., Brady S. G. 2015. Target enrichment of ultraconserved
elements from arthropods provides a genomic perspective on relationships among Hymenoptera. Mol
Ecol Resour. 15:489-501.
Faircloth B. C., McCormack J. E., Crawford N. G., Harvey M. G., Brumfield R. T., Glenn T. C. 2012.
Ultraconserved elements anchor thousands of genetic markers spanning multiple evolutionary
timescales. Syst. Biol. 61:717-726.
Findley J. S. 1972. Phenetic relationships among bats of the genus *Myotis*. Syst Biol. 21:31-52.
Fitch J. H., Shump K. A. 1979. *Myotis keenii*. Mammalian Species Archive. 121:1-3.
Glenn T. C., Nilsen R., Kieran T. J., Finger J. W., Pierson T. W., Bentley K. E., Hoffberg S., Louha S., Garcia-
De-Leon F. J., Angel del Rio Portilla M., Reed K., Anderson J. L., Meece J. K., Aggery S., Rekaya R., Alabady

M., Belanger M., Winker K., Faircloth B. C. 2016. Adapterama I: Universal stubs and primers for
thousands of dual-indexed Illumina libraries (iTru & iNext). bioRxiv. https://doi.org/10.1101/049114.
Good J. M., Vanderpool D., Keeble S., Bi K. 2015. Negligible nuclear introgression despite complete
mitochondrial capture between two species of chipmunks. Evolution. 69:1961-1972.
Gordon A., Hannon G. 2010. Fastx-toolkit. http://hannonlab.cshl.edu/fastx_toolkit.
Grabherr M. G., Haas B. J., Yassour M., Levin J. Z., Thompson D. A., Amit I., Adiconis X., Fan L.,
Raychowdhury R., Zeng Q., Chen Z., Mauceli E., Hacohen N., Gnirke A., Rhind N., di Palma F., Birren B.
W., Nusbaum C., Lindblad-Toh K., Friedman N., Regev A. 2011. Full-length transcriptome assembly from
RNA-Seq data without a reference genome. Nat Biotech. 29:644-652.
Green R. E., Braun E. L., Armstrong J., Earl D., Nguyen N., Hickey G., Vandewege M. W., St. John J. A.,
Capella-Gutiérrez S., Castoe T. A., Kern C., Fujita M. K., Opazo J. C., Jurka J., Kojima K. K., Caballero J.,
Hubley R. M., Smit A. F., Platt R. N., Lavoie C. A., Ramakodi M. P., Finger J. W., Suh A., Isberg S. R., Miles
L., Chong A. Y., Jaratlerdsiri W., Gongora J., Moran C., Iriarte A., McCormack J., Burgess S. C., Edwards S.
V., Lyons E., Williams C., Breen M., Howard J. T., Gresham C. R., Peterson D. G., Schmitz J., Pollock D. D.,
Haussler D., Triplett E. W., Zhang G., Irie N., Jarvis E. D., Brochu C. A., Schmidt C. J., McCarthy F. M.,
Faircloth B. C., Hoffmann F. G., Glenn T. C., Gabaldón T., Paten B., Ray D. A. 2014. Three crocodilian
genomes reveal ancestral patterns of evolution among archosaurs. Science. 346:1254449.
Hahn C., Bachmann L., Chevreux B. 2013. Reconstructing mitochondrial genomes directly from genomic
next-generation sequencing reads—a baiting and iterative mapping approach. Nucleic Acids Res.
41:e129.
Haynie M. L., Tsuchiya M. T. N., Ospina-Garcés S. M., Arroyo-Cabrales J., Medellín R. A., Polaco O. J.,
Maldonado J. E. 2016. Placement of the rediscovered *Myotis planiceps* (Chiroptera: Vespertilionidae)
within the *Myotis* phylogeny. J Mammal. 97:701-712.
Hedtke S. M., Townsend T. M., Hillis D. M. 2006. Resolution of phylogenetic conflict in large data sets by
increased taxon sampling. Syst Biol. 55:522-529.
Hollister N. 1909. Two new bats from the southwestern United States. publisher not identified.
Hosner P. A., Faircloth B. C., Glenn T. C., Braun E. L., Kimball R. T. 2016. Avoiding missing data biases in
phylogenomic inference: An empirical study in the landfowl (Aves: Galliformes). Mol Biol Evol. 33:1110-
1125.
Huang H., He Q., Kubatko L. S., Knowles L. L. 2010. Sources of error inherent in species-tree estimation:
Impact of mutational and coalescent effects on accuracy and implications for choosing among different
methods. Syst Biol. 59:573-583.
Huang W., Zhou G., Marchand M., Ash J. R., Morris D., Van Dooren P., Brown J. M., Gallivan K. A.,
Wilgenbusch J. C. 2016. TreeScaper: Visualizing and extracting phylogenetic signal from sets of trees.
Mol Biol Evol.
Hudson R. R., Coyne J. A., Huelsenbeck J. 2002. Mathematical consequences of the genealogical species
concept. Evolution. 56:1557-1565.
Kapusta A., Suh A., Feschotte C. 2017. Dynamics of genome size evolution in birds and mammals. Proc
Natl Acad Sci U S A. 114:E1460-E1469.
Katoh K., Misawa K., Kuma K. i., Miyata T. 2002. MAFFT: a novel method for rapid multiple sequence
alignment based on fast Fourier transform. Nucleic Acids Res. 30:3059-3066.
Lack J. B., Roehrs Z. P., Stanley C. E., Ruedi M., Van Den Bussche R. A. 2010. Molecular phylogenetics of
Myotis indicate familial-level divergence for the genus *Cistugo* (Chiroptera). J Mammal. 91:976-992.
Lanfear R., Calcott B., Ho S. Y. W., Guindon S. 2012. PartitionFinder: Combined selection of partitioning
schemes and substitution models for phylogenetic analyses. Mol Biol Evol. 29:1695-1701.
Larsen R. J., Knapp M. C., Genoways H. H., Khan F. A. A., Larsen P. A., Wilson D. E., Baker R. J. 2012.
Genetic diversity of neotropical *Myotis* (Chiroptera: Vespertilionidae) with an emphasis on South
American species. PLoS ONE. 7:e46578.

LaVal R. K. 1973. A revision of the Neotropical bats of the genus *Myotis*. Natural History Museum, Los
Angeles County.
Leaché A. D., Rannala B. 2011. The accuracy of species tree estimation under simulation: A comparison
of methods. Syst Biol. 60:126-137.
Leavitt D. H., Marion A. B., Hollingsworth B. D., Reeder T. W. 2017. Multilocus phylogeny of alligator
lizards (Elgaria, Anguidae): Testing mtDNA introgression as the source of discordant molecular
phylogenetic hypotheses. Mol Phylogenet Evol.
Lewin-Koh N. 2011. Hexagon binning: an overview.
ftp://ftp.naist.jp/pub/lang/R/CRAN/web/packages/hexbin/vignettes/hexagon_binning.pdf.
Li G., Davis B. W., Eizirik E., Murphy W. J. 2016. Phylogenomic evidence for ancient hybridization in the
genomes of living cats (Felidae). Genome Res. 26:1-11.
Li H., Durbin R. 2009. Fast and accurate short read alignment with Burrows–Wheeler transform.
Bioinformatics. 25:1754-1760.
Liu L., Edwards S. V. 2015. Comment on "Statistical binning enables an accurate coalescent-based
estimation of the avian tree". Science. 350:171-171.
Liu L., Yu L., Kubatko L., Pearl D. K., Edwards S. V. 2009. Coalescent methods for estimating phylogenetic
trees. Mol Phylogenet Evol. 53:320-328.
Manning R. W., Jones J. K. 1989. *Myotis evotis*. Mammalian Species Archive. 329:1-5.
McCormack J. E., Harvey M. G., Faircloth B. C., Crawford N. G., Glenn T. C., Brumfield R. T. 2013. A
Phylogeny of birds based on over 1,500 loci collected by target enrichment and high-throughput
sequencing. PLoS ONE. 8:e54848.
McGee M. D., Faircloth B. C., Borstein S. R., Zheng J., Darrin Hulsey C., Wainwright P. C., Alfaro M. E.
2016. Replicated divergence in cichlid radiations mirrors a major vertebrate innovation. Proc R Soc Lond
B Biol Sc. 283.
Meiklejohn K. A., Faircloth B. C., Glenn T. C., Kimball R. T., Braun E. L. 2016. Analysis of a rapid
evolutionary radiation using Ultraconserved Elements: Evidence for a bias in some multispecies
coalescent methods. Syst Biol. 65:612-627.
Miller Jr G. S., Allen G. 1928. The American bats of the genus *Myotis* and *Pixonyx*. Bulletin of the United
States National Museum. 144:1-218.
Mirarab S., Bayzid M. S., Boussau B., Warnow T. 2014. Statistical binning enables an accurate coalescent-
based estimation of the avian tree. Science. 346.
Mirarab S., Bayzid M. S., Warnow T. 2016. Evaluating summary methods for multilocus species tree
estimation in the presence of incomplete lineage sorting. Syst Biol. 65:366-380.
Mirarab S., Warnow T. 2015. ASTRAL-II: Coalescent-based species tree estimation with many hundreds
of taxa and thousands of genes. Bioinformatics. 31:i44-i52.
Morales A. E., Jackson N. D., Dewey T. A., O'Meara B. C., Carstens B. C. 2016. Speciation with gene flow
in North American *Myotis* bats. Syst Biol.
O'Farrell M. J., Studier E. H. 1980. *Myotis thysanodes*. Mammalian Species. 137:1-5.
Pagan H. J. T., Smith J. D., Hubley R. M., Ray D. A. 2010. PiggyBac-ing on a Primate genome: Novel
elements, recent activity and horizontal transfer. Genome Biol Evol. 2.
Pattengale N. D., Alipour M., Bininda-Emonds O. R. P., Moret B. M. E., Stamatakis A. 2009. How many
bootstrap replicates are necessary? In: Batzoglou S editor. Research in Computational Molecular
Biology: 13th Annual International Conference, RECOMB 2009, Tucson, AZ, USA, May 18-21, 2009.
Proceedings. Berlin, Heidelberg, Springer Berlin Heidelberg, p. 184-200.
Piaggio A. J., Valdez E. W., Bogan M. A., Spicer G. S. 2002. Systematics of *Myotis occultus* (Chiroptera:
Vespertilionidae) inferred from sequences of two mitochondrial genes. J Mammal. 83:386-395.
Platt R. N., Mangum S. F., Ray D. A. 2016. Pinpointing the vesper bat transposon revolution using the
Miniopterus natalensis genome. Mob DNA. 7:12.

Platt R. N., Vandewege M. W., Kern C., Schmidt C. J., Hoffmann F. G., Ray D. A. 2014. Large numbers of
novel miRNAs originate from DNA transposons and are coincident with a large species radiation in bats.
Mol Biol Evol. 31:1536-1545.
Platt R. N., Zhang Y., Witherspoon D. J., Xing J., Suh A., Keith M. S., Jorde L. B., Stevens R. D., Ray D. A.
2015. Targeted Capture of Phylogenetically Informative Ves SINE Insertions in Genus Myotis. Genome
Biol Evol. 7:1664-1675.
Pritham E. J., Feschotte C. 2007. Massive amplification of rolling-circle transposons in the lineage of the
bat *Myotis* lucifugus. Proc Natl Acad Sci U S A. 104:1895-1900.
Puechmaille S. J., Allegrini B., Boston E. S. M., Dubourg-Savage M.-J., Evin A., Knochel A., Le Bris Y., Lecoq
V., Lemaire M., Rist D., Teeling E. C. 2012. Genetic analyses reveal further cryptic lineages within the
Myotis nattereri species complex. Mammalian Biology. 77:224-228.
Rambaut A., Drummond A. 2013. TreeAnnotator v1. 7. 0. http://beast.bio.ed.ac.uk.
Rambaut A., Suchard M., Xie D., Drummond A. 2014. Tracer v1.6. http://beast.bio.ed.ac.uk.
Ray D. A., Feschotte C., Pagan H. J. T., Smith J. D., Pritham E. J., Arensburger P., Atkinson P. W., Craig N.
L. 2008. Multiple waves of recent DNA transposon activity in the bat, *Myotis lucifugus*. Genome Res.
18:717-728.
Ray D. A., Pagan H. J. T., Thompson M. L., Stevens R. D. 2007. Bats with hATs: Evidence for recent DNA
transposon activity in genus *Myotis*. Mol Biol Evol. 24:632-639.
Rice P., Longden I., Bleasby A. 2000. EMBOSS: the European molecular biology open software suite.
Elsevier Current Trends.
Rodríguez-Ezpeleta N., Brinkmann H., Roure B., Lartillot N., Lang B. F., Philippe H. 2007. Detecting and
overcoming systematic errors in genome-scale phylogenies. Syst Biol. 56:389-399.
Roehrs Z. P., Lack J. B., Van Den Bussche R. A. 2010. Tribal phylogenetic relationships within
Vespertilioninae (Chiroptera: Vespertilionidae) based on mitochondrial and nuclear sequence data. J
Mammal. 91:1073-1092.
Ruedi M., Mayer F. 2001. Molecular systematics of bats of the genus *Myotis* (Vespertilionidae) suggests
deterministic ecomorphological convergences. Mol Phylogenet Evol. 21:436-448.
Ruedi M., Stadelmann B., Gager Y., Douzery E. J. P., Francis C. M., Lin L.-K., Guillén-Servent A., Cibois A.
2013. Molecular phylogenetic reconstructions identify East Asia as the cradle for the evolution of the
cosmopolitan genus *Myotis* (Mammalia, Chiroptera). Mol Phylogenet Evol. 69:437-449.
Seo T.-K. 2008. Calculating bootstrap probabilities of phylogeny using multilocus sequence data. Mol Biol
Evol. 25:960-971.
Simões B. F., Rebelo H., Lopes R. J., Alves P. C., Harris D. J. 2007. Patterns of genetic diversity within and
between *Myotis d. daubentonii* and *M. d. nathalinae* derived from cytochrome *b* mtDNA sequence data.
Acta Chiropt. 9:379-389.
Sota T. 2002. Radiation and reticulation: extensive introgressive hybridization in the carabid beetles
Ohomopterus inferred from mitochondrial gene genealogy. Population Ecology. 44:0145-0156.
Stadelmann B., Lin L. K., Kunz T. H., Ruedi M. 2007. Molecular phylogeny of New World *Myotis*
(Chiroptera, Vespertilionidae) inferred from mitochondrial and nuclear DNA genes. Mol Phylogenet Evol.
43:32-48.
Stamatakis A. 2006. RAxML-VI-HPC: maximum likelihood-based phylogenetic analyses with thousands of
taxa and mixed models. Bioinformatics. 22:2688-2690.
Swofford D. L. 2003. PAUP*. Phylogenetic analysis using parsimony (* and other methods). Version 4.
Thomas J., Sorourian M., Ray D., Baker R. J., Pritham E. J. 2011. The limited distribution of Helitrons to
vesper bats supports horizontal transfer. Gene. 474:52-58.
Vachaspati P., Warnow T. 2015. ASTRID: Accurate Species TRees from Internode Distances. BMC
Genomics. 16:S3.

Valdez E. W., Choate J. R., Bogan M. A., Yates T. L. 1999. Taxonomic status of *Myotis occultus*. J Mammal.
80:545-552.
Warner R. M. 1982. *Myotis auriculus*. Mammalian Species. 191:1-3.
Wiens J. J., Kuczynski C. A., Smith S. A., Mulcahy D. G., Sites J. J. W., Townsend T. M., Reeder T. W. 2008.
Branch lengths, support, and congruence: Testing the phylogenomic approach with 20 nuclear loci in
snakes. Syst Biol. 57:420-431.
Willis S. C., Farias I. P., Ortí G. 2014. Testing mitochondrial capture and deep coalescence in Amazonian
cichlid fishes (Cichlidae: *Cichla*). Evolution. 68:256-268.

Table 1 - Specimens Examined

Species	Name herein	Museum identification num.	Num. UCE Loci	UCE contig accession	Mitochondrial genome accession
Eptesicus fuscus	*fuscus* [G]	GCA_000308155.1	2,467	Pending	KF111725.1
Eptesicus fuscus	*fuscus*	TK 178736	2,849	Pending	MF143474
Myotis albescens	*albescens* [1]	RDS 7889	2,764	Pending	MF143471
Myotis albescens	*albescens* [2]	QCAZ 9157	1,185	Pending	MF143497
Myotis albescens	*albescens* [3]	TK 61766	2,872	Pending	MF143470
Myotis ater	*ater*	M4430	2,774	Pending	MF143493
Myotis auriculus	*auriculus*	MSB 40883	2,229	Pending	MF143486
Myotis brandtii	*brandtii* [G]	GCA_000412655.1	2,446	Pending	KT210199.1
Myotis californicus	*californicus*	UMMZ 175828	2,948	Pending	MF143469
Myotis davidii	*davidii* [G]	GCA_000327345.1	2,450	Pending	KM233172.1
Myotis diminutus	*diminutus*	QCAZ 9168	3,078	Pending	MF143481
Myotis dominicensis	*dominicensis*	TK 15624	2,576	Pending	MF143467
Myotis evotis	*evotis*	MSB 47323	2,586	Pending	MF143468
Myotis fortidens	*fortidens*	MSB 54941	2,791	Pending	MF143483
Myotis horsfieldii	*horsfieldii*	MHNG 1926.039	3,017	Pending	MF143494
Myotis keaysi	*keaysi*	TK 13525	3,195	Pending	MF143477
Myotis keenii	*keenii*	UAM 113849	2,723	Pending	MF143472
Myotis leibii	*leibii*	TK 24872	3,119	Pending	MF143488
Myotis levis	*levis*	RDS 7781	2,538	Pending	MF143482
Myotis lucifugus	*lucifugus* [G]	GCA_000147115.1	2,429	Pending	KP273591.1
Myotis lucifugus	*lucifugus*	MSB 46679	2,736	Pending	MF143491
Myotis melanorhinus	*melanorhinus*	TK 193888	3,177	Pending	MF143489
Myotis nigricans	*nigricans* [1]	QCAZ 9601	2,854	Pending	MF143495
Myotis nigricans	*nigricans* [2]	TK 194145	3,159	Pending	MF143484
Myotis nyctor	*nyctor*	TK 151413	856	Pending	MF143498
Myotis occultus	*occultus*	MSB 121995	2,957	Pending	MF143490
Myotis oxyotus	*oxyotus*	UMMZ RCO1013	3,106	Pending	MF143479
Myotis riparius	*riparius* [1]	TK 22688	2,924	Pending	MF143473
Myotis riparius	*riparius* [2]	TK 101723	2,990	Pending	MF143475
Myotis riparius	*riparius* [3]	TK 145199	2,890	Pending	MF143480
Myotis ruber	*ruber*	MVZ 185692	2,757	Pending	MF143478
Myotis septentrionalis	*septentrionalis*	TK 194420	2,916	Pending	MF143487
Myotis thysanodes	*thysanodes*	07LEP	2,821	Pending	MF143492
Myotis velifer	*velifer*	MSB 70877	2,704	Pending	MF143499
Myotis vivesi	*vivesi*	MSB 42658	2,469	Pending	MF143476
Myotis volans	*volans*	MSB 40886	2,819	Pending	MF143496
Myotis yumanensis	*yumanensis*	TK 194144	2,589	Pending	MF143485

Collection abbreviations: Museum of Southwestern Biology (MSB), Museum of Vertebrate Zoology (MVZ), Natural History Museum of Geneva (MHNG), Pontificia Universidad Catolica del Ecuador Museo de Zoologia (QCAZ), Texas Tech University Natural Science Research Laboratory (TK), University of Alaska Museum of the North (UAM), University of Michigan Museum of Zoology (UMMZ). Samples beginning with "GCA_" represent genome assemblies available through NCBI.

Table 2. Character information

Min. num. of taxa	Percent of taxa per locus	Number of Loci	Parsimony-informative characters	Variable uninformative	Alignment length	Optimum Partitions
5	15	3,648	38,718	62,588	1,377,262	31
9	25	3,379	37,894	57,672	1,260,248	37
12	35	3,232	37,259	56,453	1,227,093	34
16	45	3,064	36,539	54,782	1,187,492	31
20	55	2,890	35,284	53,200	1,144,471	27
24	65	2,668	34,373	51,148	1,091,620	27
27	75	2,481	33,031	48,732	1,041,099	20
31	85	2,034	29,179	41,711	903,903	16
35	95	1,193	18,288	24,189	575,321	15
37	100	212	3,480	4,778	112,125	6

General alignment information. For a subset of analyses a series of alignments were generated based on the number of taxa per locus. Thirty-seven taxa were examined so an alignment with all 37 taxa was considered 100% complete. Parsimony-informative characters make up a small portion of the total alignment. The optimum partitioning scheme was calculated with PartitionFinder.