Despite reproductive interference, the net outcome of reproductive interactions

among spider mite species is not necessarily costly

Salomé H. Clemente, Inês Santos, Rita Ponce, Leonor R. Rodrigues, Susana A. M.

Varela* and Sara Magalhães*[1]

salome.hc@gmail.com, inesflsantos@gmail.com, anaritaponce@gmail.com,

leonor.rodrigues89@gmail.com, savarela@fc.ul.pt, snmagalhaes@fc.ul.pt

*co-last authorship

cE3c – Centre for Ecology, Evolution and Environmental Changes

Faculdade de Ciências da Universidade de Lisboa

Edifício C2, 3º Piso

Campo Grande

1749-016 Lisbon

Portugal

[1]: snmagalhaes@fc.ul.pt

Abstract

Reproductive interference is considered a strong ecological force, potentially leading

to species exclusion. This supposes that the net effect of reproductive interactions is

strongly negative for one of the species involved. Testing this requires a

comprehensive analysis of interspecific reproductive interactions, accounting for the

order and timing of mating events, and for their effects on either fertility or fecundity.

To this aim, we measured reproductive interactions between a focal species,

Tetranychus urticae, and an invasive (*T.evansi*) and a resident (*T. ludeni*) species,

varying the mating sequence and interval, and measuring the effect of such crosses on

fecundity and offspring sex ratio (a measure of fertility, as these species are

haplodiploid). We found that mating with heterospecifics affected fecundity and sex

ratio negatively, but also positively, depending on the species involved, and on the

order and timing of mating events. Overall, the net effect of reproductive interactions

was weak despite strong effects of particular events. In natural situations the outcome

of reproductive interactions will thus hinge upon the frequency of each event.

Keywords:

Biological invasions, sperm precedence, *Tetranychus*, reproductive interactions,

mating.

Introduction

Reproductive interference, that is, any kind of sexual interaction between two species that diminishes the fitness of at least one of them (Gröning and Hochkirch 2008, Kishi et al. 2009, Burdfield-Steel and Shuker 2011), may have severe effects on the outcome of species interactions. Indeed, theory predicts that reproductive interference may contribute to species exclusion more often than resource competition (Gröning and Hochkirch 2008, Kishi et al. 2009, Kishi and Nakazawa 2013). For example, it has been posited that reproductive interference may underlie the success of some invasive species (e.g. Nishida et al. 2012).

Most studies of reproductive interference concern the fitness outcome of interspecific matings of two species that do not produce viable hybrids (Gröning and Hochkirch 2008). In this case, the reproductive effects of the interspecific interaction will be expressed only when organisms mate with both conspecifics and heterospecifics (as mating with conspecifics alone will yield no offspring). Moreover, clearly evaluating the effects of reproductive interference on exclusion in polyandrous species necessitates measuring all possible combinations of mating order (*i.e.*, whether heterospecific matings occur before or after conspecific ones) and timing (*i.e.*, the interval between mating events) between pairs of species. It is also important to test whether reproductive interactions affect fecundity (egg production) or fertility (egg fertilization). This information can then be integrated to predict the net outcome of reproductive interactions between species. Despite the many studies on reproductive interference, none has yet applied this approach. Indeed, some studies attempt to predict how reproductive interference affects species exclusion, but do so by focussing on some sequence events only. For example, Takafuji (1997) used a Lotka-Volterra

modified model to predict the effect of reproductive interference between two mite

species (*Panonychus citri* and *P. mori*) on species exclusion, but they used only one

possible combination of mating interactions between species. In contrast, other

studies consider different orders of mating events (eg, Kyogoku and Nishida 2013), but

do not integrate this information to generate a prediction concerning the net effect of

reproductive interactions on species distributions.

Here, we aimed at testing how the outcome of different mating events among

species may affect their life-history traits, using spider mites, a group where

reproductive interference has been frequently observed (Collins and Margolies 1994;

Takafuji et al. 1997; Ben-David et al. 2009, Sato et al. 2014). Spider mites are

haplodiploid, hence the distinction between fecundity and fertilization effects can be

made given that fertilized eggs result in female offspring and unfertilized eggs in male

offspring. Thus, fertilization failures can be detected by a reduction in the proportion

of female offspring, whereas impairment of egg production is detected by a reduction

in the total number of offspring.

We used a system composed of one focal species, *Tetranychus urticae*, in

sexual heterospecific interactions with another resident species, *T. ludeni*, and an

invasive species, *T. evansi*. These three herbivorous species co-occur in the

Mediterranean region and are often found on the same host plant (Escudero and

Ferragut 2005, Boubou et al. 2012, Godinho et al. 2016). Whereas *T. urticae* and *T.*

ludeni are resident species, *T. evansi* has only recently invaded the European continent

(Boubou et al. 2012). Whereas information on the interaction between *T. urticae* and

T. ludeni is as yet lacking, heterospecific matings have been observed between *T.*

urticae and *T. evansi* (Sato et al. 2014, 2016, Clemente et al. 2016). Moreover, *T. evansi*

can exclude *T. urticae* on tomato plants (Sarmento et al. 2011a), a result that

correlates with field observations (Ferragut et al. 2013). Finally, a recent study has

shown that, in competition with *T. evansi*, the population growth of *T. urticae* is more

severely affected when plants are colonized by virgin females than when plants are

colonized by mated females, suggesting that reproductive interference may be

responsible for the species distribution patterns observed (Sato et al. 2014).

To postulate hypotheses concerning the consequences of heterospecific

matings, it is crucial to understand within-species reproductive behaviour. *T. urticae*,

the focal species, exhibits first male sperm precedence, with second matings being

sometimes effective if they occur within the 24 hours following the first (Helle 1967).

However, females that mate multiple times with conspecific males, after a 24h interval

between matings, produce fewer fertilized offspring (*i.e.*, females) (Macke et al. 2012),

suggesting that sperm displacement after 24h is possible. Here, we hypothesize that

mating order and the mating interval will affect the outcome of reproductive

interference in *T. urticae*. Also, given that *T. evansi*, the invasive species, displaces *T.

urticae*, unlike *T. ludeni*, we expect the former to exert stronger effects than the latter.

To this aim, we performed crosses between *T. urticae* and the two other species at

different time intervals and with different mating orders, and measured the

consequences for the two species involved in the cross.

Material and Methods

Stock Cultures

The mite species used in this study were collected in Carregado (39.022260, -

8.966566), Portugal, and all laboratory populations were established from an initial

pool of 300 mated females. The laboratory population of *T. urticae* was collected on

tomato plants (*Solanum lycopersicum)* in May 2010, that of *T. evansi* on *Physalis*

angulata in May 2012 and that of *T. ludeni* on tomato in September 2012. The

populations of *T. evansi* and *T. ludeni* became extinct in August 2012 and May 2013,

respectively, being subsequently replaced with populations from the same location,

both collected in *Datura stramonium* plants. Both populations of *T. evansi* and *T.*

ludeni were used in the experiments.

Species identity was confirmed through polymerase chain reaction–restriction

fragment length polymorphism (PCR–RFLP) of the ITS2 region (Hurtado et al. 2008), on

approximately 50 females of each population. Total genomic DNA was extracted from

each individual spider mite using the Sigma-Aldrich GenEluteTM Mammalian Genomic

DNA Miniprep Kit, following manufacturer´s instructions, except for the elution

volume, which we set to 20µL of RNase free water (Qiagen NV, Venlo, The

Netherlands) to increase the concentration of DNA obtained from this very small

animal (c.a. 300µm long).

Adult females from populations used in this experiment were screened for

Wolbachia using the primers *wsp* (*Wolbachia*-specific primers) 81F and 691R (Braig et

al. 1998). We did this to avoid potential cytoplasmic incompatibility as a confounding

factor in our measurements. PCR assay procedures were as described in Breeuwer

(1997). Results were positive for *Wolbachia* infection and all spider mite populations

were thus treated by placing adult females in detached bean leaves with tetracycline

(0.025% w/v) for three consecutive generations, then absence of *Wolbachia* was

confirmed using the same protocol as above.   Other endosymbionts tested

(Arsenophorous, Rickettsia, Spiroplasma and Cardinium) were absent from these

populations.

Bean (*Phaseolus vulgaris*) and tomato (*Solanum lycopersicum*) plants were planted

every week and grown in an herbivore-free greenhouse, being watered two to three

times a week. *T. urticae* populations were maintained on trays with 6-10 bean plants

whereas those of *T. evansi* and *T. ludeni* were kept on tomato plants at 25°C, both with

a 16 L: 8D photoperiod. Plant trays were changed every two weeks, placing old leaves

on top of uninfested plants. Cultures were kept inside plastic boxes (28x39x28 cm),

with an opening of 25x15 cm polyamide fabric (80 µm mesh width).

Experimental procedure

Experiments were done on the plant species from which the female tested had been

cultured. As in the literature there was no information on whether hybridization is

possible between *T. urticae* and *T. ludeni*, we studied the outcome of a single

heterospecific mating between these two species (the same analysis for *T. urticae* and

T. evansi was performed in a previous experiment (Clemente et al. 2016)).

Subsequently, we set out to study the heterospecific interactions between *T. urticae*

and the invasive *T. evansi* and the resident *T. ludeni* species for which we analysed the

outcome of mating with a heterospecific male before or after a conspecific male. Since

we focused on interactions with *T. urticae* (the focal species of our study), we

performed crosses between *T. urticae* males or females and *T. evansi* or *T. ludeni*

males or females, but not between the two latter species. All experiments were

performed in an acclimatized room at approximately 25°C.

a) The outcome of a single heterospecific mating between *T. urticae* and *T. ludeni*

To determine whether hybridization occurred between *T. urticae* and *T. ludeni*, we measured the offspring sex-ratio resulting from single heterospecific matings. Given that only females develop from fertilized eggs, a whole-male offspring would mean unsuccessful hybridization. However, even in the absence of viable hybrids, heterospecific matings could result in aborted development of heterospecifically-fertilized eggs, meaning that females would produce fewer eggs. To test this, we compared the fecundity of *T. urticae* and *T. ludeni* females that mated with a heterospecific male to that of virgin females and of females mated with a conspecific male.

Females were collected from the stock populations, isolated at the quiescent deutonymph stage (which precedes their last moult before reaching adulthood), and kept in groups of approximately 15 females on bean (*Phaseolus vulgaris*) leaf discs (2 cm^2) until emergence, to ensure their virginity. Adult males were collected from the same stock populations and kept isolated in leaf discs (2 cm^2) for at least 24 hours before the assay, to ensure sperm replenishment. Females were placed individually in leaf discs (1 cm^2) with either a conspecific or a heterospecific male and observed continuously until copulation occurred. Only matings that lasted at least 1 minute were considered effective (Boudreaux 1963). These experiments had the maximum duration of 2 hours. If no mating occurred within this time, individuals were discarded. Subsequently, females were isolated in a leaf disc (2 cm^2), then transferred to a new disc every three days until the female's death. The number of eggs laid was registered after female transfer to a new leaf disc. Eggs were left to develop until adulthood

when offspring sex-ratio could be determined. With this data, we tested whether

heterospecific matings affected (a) the mean daily fecundity and (b) offspring sex ratio

(hence the proportion of fertilized offspring).

b) The outcome of heterospecific matings that precede or follow conspecific ones

To determine the outcome of mating with a heterospecific male before or after a

conspecific male between *T. urticae* and the other two species, we compared the

fecundity and offspring sex ratio of those crosses to that of females that mated with

two conspecific males. The experimental procedure was as described above, except

that we let females mate with a conspecific or a heterospecific male, then placed the

focal females with another male. We created the following mating sequences:

conspecific-conspecific, conspecific-heterospecific and heterospecific-conspecific. The

second mating occurred either immediately after the first mating (0 hours treatment)

or 24 hours later. If no mating was observed within 2 hours, the females were

discarded. We used the 0h and 24h mating intervals because the time interval was

shown to affect the degree of sperm precedence in spider mites (Helle 1967).

Statistical analysis

All analyses were carried out using R (version 3.3.2, R Development Core Team 2016).

To analyse female fecundity within each species (*T. urticae*, *T. evansi* and *T. ludeni*), we

used linear models (LM procedure), considering the mean number of eggs per day as

the response variable (oviposition rate). To analyse offspring sex ratio within each

species, we used generalized linear models (GLM procedure) with a quasi-binomial

distribution – due to overdispersion of the data –, considering the number of female

and male offspring produced by each focal female as the response variables (analysed

together with the function cbind).

For both types of analyses, we used as fixed factors the mating order (with

three levels: the control treatment, where a female mated twice with conspecific

males; an experimental treatment where the heterospecific male was the first to mate

with the female; and another experimental treatment where the heterospecific male

was the second to mate with the female) and the mating interval (with two levels:

either 0h or 24h interval between matings). We also tested the interaction among

these fixed factors. If the interaction was non-significant, a backward stepwise

procedure was used to find the best simplified fitted model. We performed

independent analyses for each species within each species pair (i.e. for *T. urticae* and

T. evansi females in *T. urticae versus T. evansi* crosses; and for *T. urticae* and *T. ludeni*

females in *T. urticae versus T. ludeni* crosses), as shown in Table 1.

We did a first block of experiments with the populations of *T. evansi* and *T.*

ludeni collected in 2012 (block 1). For question b) we also did a second block of

experiments with populations of those species from 2013 (block 2). In block 2 we did

not repeat all treatments, but only the crosses that were not complete before the

extinction of block 1 populations, as well as their respective controls – hence, there

were no treatments that were only performed in block 2. Because of that, instead of

including the factor block in the statistical models as a covariate, we did all the

statistical analyses with block 1 only and with block 1 and block 2 together. Since the

results were qualitatively similar (Table S1), here we present the results from the

analysis with block 1 and block 2 together.

Results

a) The outcome of a single heterospecific mating between *T. urticae* and *T.*

231   ***ludeni***

Crosses between *T. ludeni* and *T. urticae* resulted in 100% male offspring, indicating

that hybrid production between these species is inexistent. The fecundity of *T. urticae*

females that mated heterospecifically was not significantly different from that of virgin

females or from that of females mated with a conspecific male ($F_{2,78}$= 1.886, P= 0.1585;

Figure 1). On the other hand, the fecundity of *T. ludeni* females that mated with

conspecifics or heterospecifics was significantly higher than that of virgin females

($F_{2,66}$= 1.886, P= 0.1585; Figure 1 and Table 1). Therefore, mating with heterospecific

males does not result in the aborted fertilization of oocytes for *T. urticae* and *T. ludeni*

females.

b) The outcome of heterospecific matings that precede or follow conspecific

243   **ones**

(i)   T. urticae vs T. evansi

The oviposition rate of *T. urticae* females that mated with either a conspecific and a

heterospecific or with two conspecific mates varied significantly according to mating

order in interaction with mating interval ($F_{2,136}$ = 6.026, P = 0.0031). Specifically, it was

higher for *T. urticae* females that mated with *T. evansi* males just before mating with a

conspecific male than for any other cross at 0h mating interval ($|t|$ = 4.964, P < 0.0001

and $|t|$ = 3.288, P = 0.0009, in comparison with double conspecific matings and with

matings with a conspecific followed by a mating with an heterospecific, respectively;

Fig. 2a). At the 24h interval, however, mating combinations did not affect this trait. The

proportion of fertilized offspring (*i.e.*, daughters) of females *T. urticae* also varied

significantly according to mating order in interaction with mating interval ($F_{2,106}$=

4.963, P= 0.0087). But in contrast to the oviposition rate, this trait was affected at the

24h interval only, in which mating with a *T. evansi* male after mating with a conspecific

male resulted in a decrease in the proportion of fertilized offspring of *T. urticae*

females, relative to other mating sequences (|t| = 5.362, P < 0.0001 and |t| = 5.103, P

< 0.0001, in comparison with double conspecific matings and with matings with an

heterospecific followed by a mating with a conspecific male, respectively; Fig. 2b).

The mating order also affected differentially the oviposition rate of *T. evansi*

females, depending on the interval between matings ($F_{2,187}$= 4.977, P= 0.0078). *T.*

evansi females that mated with *T. urticae* males immediately after conspecific mates

had reduced oviposition rate relative to other mating sequences at this time interval

(|t| = 2.841, P = 0.0050 and |t| = 2.692, P = 0.0078 in comparison with double

conspecific matings and with matings with a heterospecific followed by a mating with a

conspecific male, respectively; Fig. 2c); however, if the heterospecific cross occurred

24 hours before the conspecific cross, the oviposition rate of *T. evansi* females

increased relative to double conspecific matings at this time interval (|t| = 2.948, P =

0.0036; Fig. 2c). These crosses did not significantly affect sex ratio ($F_{2,111}$= 0.368, P=

0.6931; Fig. 2d).

(ii)   T. urticae vs T. ludeni

In crosses with the resident species (*T. ludeni*), the oviposition rate of *T. urticae*

females varied significantly according to mating order in interaction with mating

interval ($F_{2,144}$ = 3.694, P = 0.0273). Specifically, we found that, at 0h interval, females

that mated first with a conspecific then with a heterospecific male had lower

oviposition rate than females that mated first with a heterospecific then with a

conspecific male ($|t| = 2.736$, $P = 0.0070$; Fig. 3a) At the 24h interval, the oviposition

rate of females that mated first with a conspecific then with a heterospecific male was

lower than that of double conspecific crosses. ($|t| = 2.505$, $P = 0.0134$; Fig. 3a). *T.*

urticae females suffered no significant changes in offspring sex ratio from matings with

T. ludeni males ($F_{2,99} = 1.141$, $P = 0.3237$; Figure 3b).

In *T. ludeni* females, the oviposition rate and the proportion of fertilized

offspring varied significantly according to mating order in interaction with the mating

interval (($F_{2,248} = 14.098$, $P < 0.0001$ and $F_{2,152} = 10.1064$, $P < 0.0001$, for oviposition rate

and proportion of fertilized offspring respectively). Compared to the control

treatment, *T. ludeni* females had lower oviposition rate when mating with *T. urticae*

males immediately before conspecifics males ($|t| = 2.605$, $P = 0.0097$; Fig. 3c). At the

24 hour interval, the conspecific crosses yielded higher oviposition rate than all other

crosses in this time interval (($|t| = 4.646$, $P < 0.0001$ and $|t| = 3.805$, $P = 0.0002$, in

comparison with females mating with a conspecific before an heterospecific male and

females mating with an heterospecific before mating with a conspecific, respectively;

Fig. 3c). Additionally, when *T. ludeni* females mated with *T. urticae* males 24h after

conspecific matings, the proportion of fertilized offspring was significantly lower than

that of other crosses at this time interval (($|t| = 4.084$, $P < 0.0001$ and $|t| = 3.586$, $P =$

0.0005, in comparison with double conspecific matings and with females mating with a

heterospecific before mating with a conspecific, respectively; Figure 3d). The mating

sequence had no effect on the sex ratio at the 0h interval.

Discussion

In this study, we investigated the consequences of mating with heterospecifics for the

fertilization success and offspring viability in a system composed of three spider-mite

species. We found that heterospecific matings between *T. urticae* and *T. ludeni* did not

result in fertilized offspring (i.e., females), nor did it have any negative effects on egg

viability, as shown for matings between *T. urticae* and *T. evansi* (Sato et al. 2014,

Clemente et al. 2016). In fact, *T. ludeni* females that mate with *T. urticae* males

produce more (male) offspring than virgin *T. ludeni* females. Second, the effects of

heterospecific matings on the outcome of previous or subsequent matings with

conspecifics were highly dependent on the species pair involved, on the trait measured

and on the timing and order of mating events. Despite strong effects of particular

mating sequences, the results taken as a whole suggest that the net effect of

reproductive interactions between species are relatively weak.

Positive effects of interspecific reproductive interactions were found for

fecundity. This can be due to a stimulation of oogenesis by the sperm of heterospecific

males, increasing the availability of oocytes to subsequent matings with conspecifics.

Indeed, oogenesis is stimulated by conspecific sperm in several species (Qazi et al.

2003, Xu and Wang 2011). This could also be the case with heterospecific sperm. If so,

it could explain the higher fecundity found in crosses between *T. urticae* and *T. evansi*.

In fact, earlier studies have documented that interactions with heterospecific males

are not always negative. In some gynogenetic species, heterospecific mating is a

prerequisite for embryogenesis (Gumm and Gabor 2005, Schlupp 2010). Moreover, in

some invertebrate species, females receive nuptial gifts from heterospecific males

(Vahed 1998, Costa-Schmidt and Machado 2012). However, to our knowledge, this is

the first time that an increase in fecundity following a heterospecific mating is

described in the literature. Such effects may thus be rare. Still, earlier studies may have

overlooked them because they have not examined the roles of the order of mating in

the outcome of heterospecific mating interactions.

329   Nonetheless, we also detected several negative effects of mating with

heterospecifics, as found in most studies of reproductive interference (Gröning and

Hochkirch 2008, Kishi 2015). We found both a reduction in the number of eggs laid and

a decrease in fertilization success (i.e., offspring sex ratio). However, the incidence of

these two effects varied according to the species involved, the order of matings and

the time interval. Whereas effects on fecundity were found in several mating

sequences, an effect on fertilization success was found only when the heterospecific

male mated with the female 24 hours after the conspecific male. This is at odds with

expectations stemming from findings on conspecific matings, which show (a) first-male

precedence and (b) exceptions to this rule only if the second male mates immediately

after the first (Helle 1967). Therefore, the mechanisms underlying sperm displacement

by heterospecific males in spider mites should be investigated.

341   This also suggests that first male precedence found in conspecific matings

cannot be extrapolated to matings involving heterospecific sperm. This contrasts with

the recent finding that effects of heterospecific matings in *Drosophila* could be

predicted from the harmful effects of conspecific mates (Yassin and David 2015), and

that genes involved in conspecific male precedence also affect sperm precedence in

multiple matings involving heterospecifics (Civetta and Finn 2014). This indicates that

the equivalence of effects of conspecific and heterospecific sperm on the outcome of

conspecific matings is dependent on the type of effect and/or the species involved in

the interaction.

Since effects of heterospecific matings depend on the order and timing of

occurrence, the outcome of interspecific reproductive interactions will depend on the

frequency with which those different types of matings occur in nature. This, in turn,

will depend on the discrimination abilities between species. First, these interactions

will occur only if species discrimination is weak. This, indeed, has been explicitly

demonstrated for the *T. evansi*/*T. urticae* interaction (Clemente et al. 2016), but not

for *T. ludeni*/*T. urticae*. Still, these species do mate with heterospecifics under no

choice scenarios, as shown here, hence the scope for the occurrence of reproductive

interference does exist.

What then, would be the relative frequency of the mating sequences tested

here? In spider mites, conspecific males often guard quiescent females (*i.e.*, the last

larval stage before becoming adult female), to ensure mating immediately after

emergence. If males guard preferentially conspecific females, as shown in other spider

mite species pairs (Collins et al. 1993, Takafuji et al. 1997), heterospecific matings will

occur more often after rather than before conspecific ones. Moreover, we have shown

that *T. urticae* females become less receptive to both conspecific and heterospecific

matings if the first mating has occurred 24h before the second (Clemente et al. 2016).

Hence, this leads to the prediction that the most common mating sequence among

these species will be a heterospecific mating immediately following a conspecific one.

Under those circumstances the only effect of heterospecific matings is a fecundity

reduction in *T. evansi* upon mating with *T. urticae*. This would mean that the invasive

species suffers more from reproductive interference than the resident.

Even assuming that all mating combinations do occur, reproductive interactions

between *T. urticae* and *T. evansi* can be positive or negative for the two species,

depending on the mating sequence. Therefore, reproductive interference cannot be

invoked to explain the exclusion of *T. urticae* in habitats with *T. evansi* (Ferragut et al.

2013, Sarmento et al. 2011b). Other factors may contribute to this exclusion, as the

production of a dense web by *T. evansi*, which prevents heterospecifics from accessing

the surface of the leaves to feed and oviposit (Sarmento et al. 2011b). Importantly,

however, we show that the occurrence and strength of reproductive interference

cannot be assessed with the unique evaluation of the outcome of a specific type of

reproductive interaction. The different types of mating combinations – the order and

interval between matings – have great influence on the overall outcome of

heterospecific interactions and on the relative frequency of such events. This confirms

the importance of using complete experimental designs on the detection and

characterization of reproductive interference.

Acknowledgements

The authors wish to thank Yukie Sato and Maurice W. Sabelis for helpful discussions

during the early stages of this work. These experiments and Inês Santos were funded

by Portuguese National Funds through an FCT-ANR project (FCT-ANR//BIA-

EVF/0013/2012) to SM and Isabelle Olivieri. SHC and LRR had PhD fellowships funded

by FCT (SFRH/ BD/ 90156/2012 and SFRH/BD/87628/2012, respectively). SAMV and RP

had Post-doctoral fellowships from FCT, (SFRH/BPD/66042/2009) and (SFRH / BPD /

42801 /2008) respectively. The authors declare no conflict of interest.

 FIGURE LEGENDS

Figure 1 | Average daily fecundity of virgin females, and of females that have mated

with a conspecific or a heterospecific male. Tu: *T. urticae* males or females; Tl: *T. ludeni*

males or females. Grey bars: matings involving *T. urticae* females; white bars: matings

involving *T. ludeni* females. Error bars represent the standard errors of the mean.

Numbers on the bottom of bars represent the sample size for each type of mating.

Figure 2 | Average daily fecundity and estimated offspring sex ratio resulting from

interactions between *T. urticae* (a, b; grey solid bars) and *T. evansi* (c,d; striped bars)

females with conspecific and heterospecific males. In each plot, bars on the left side of

the dotted straight line correspond to treatments where second matings occurred

immediately (0h) after the first one; bars on the right side correspond to treatments

where second matings occurred 24h after the first one. "1st M": first male that mated

with the female; "2nd M": second male. The interval indicates the time of occurrence

of the second mating, i.e., if immediately after the first mating (0h) or 24h later. "Tu":

T. urticae males; "Te": *T. evansi* males. Letters above the bars indicate significant

differences among treatments (small letters: among crosses occurring with a 0h

interval; capital letters: among crosses occurring with a 24h interval). Error bars

represent the standard errors of the mean. For offspring sex ratio, we obtained the

estimates of the probability of being female and correspondent standard errors of the

mean from the statistical GLM models. This takes into account sex ratio variation

among females, as well as the quasi-binomial correction for overdispersion of the data.

Numbers on the bottom of bars represent the sample size for each type of mating.

Figure 3 | Average daily fecundity and estimated offspring sex ratio resulting from interactions between *T. urticae* (plots a, b; grey bars) and *T. ludeni* (plots c, d; white bars) females with conspecific and heterospecific males. In each plot, bars on the left side of the dotted line correspond to treatments where second matings occurred immediately (0h) after the first one; bars on the right side correspond to treatments where second matings occurred 24h after the first one. "1st M": first male that mated with the female; "2nd M": second male. The interval indicates the time of occurrence of the second mating, i.e., if immediately after the first mating (0h) or 24h later. "Tu": *T. urticae* males; "Tl": *T. ludeni* males. Letters above the bars indicate the significant differences between treatments (small letters: among crosses occurring with a 0h interval; capital letters: among crosses occurring with a 24h interval. Error bars represent the standard errors of the mean. For offspring sex ratio, we obtained the estimates of the probability of being female and correspondent standard errors of the mean from the statistical GLM models. This takes into account sex ratio variation among females, as well as the quasi-binomial correction for overdispersion of the data. Numbers on the bottom of bars represent the sample size for each type of mating.

References

Alpedrinha J. and Magalhães S. 2016. Sex Allocation: l'enfer c'est les autres? — Curr.

Biol. 26: R476-R478.

Arnqvist, G. and Rowe, L. 2005. Sexual Conflict. — Princeton University Press,

Princeton.

Ben-David, T. et al. 2009. Asymmetric reproductive interference between two closely

related spider mites: Tetranychus urticae and T. turkestani (Acari: Tetranychidae). —

Exp. Appl. Acarol. 48: 213-227.

Braig, H. R. et al.1998. Cloning and Characterization of a Gene Encoding the Major

Surface Protein of the Bacterial Endosymbiont *Wolbachia pipienti*s. — J. Bacteriol. 180:

2373–2378.

Breeuwer, J. A. 1997. *Wolbachia* and cytoplasmic incompatibility in the spider mites

Tetranychus urticae and *T.* turkestani. — Heredity79: 41–47.

Boubou, A. et al.2012. Test of colonisation scenarios reveals complex invasion history

of the red tomato spider mite *Tetranychus evansi*. — PLoS One 7: 35601.

Boudreaux, H. B. 1963. Biological aspects of some phytophagous mites. — Annu. Rev.

Entomol. 8: 137–154.

Burdfield-Steel, E. R. and Shuker, D. M. 2011. Reproductive interference. — Curr. Biol.

21: 450–451.

Civetta, A. and Finn, S. 2014. Do candidate genes mediating conspecific sperm

precedence affect sperm competitive ability within species? A test case in Drosophila.

– G3 (Bethesda) 4: 1701-1707.

Clemente, S. H. et al. 2016. Incomplete species recognition entails few costs in spider

mites, despite first-male precedence. – Behav. Ecol.Sociobiol. in press. DOI

10.1007/s00265-016-2124-0

Collins, R. D. et al. 1993. Guarding behavior and reproductive isolation in two

tetranychid mite species (Acari: Tetranychidae). – Ann. Entomol. Soc. Am. 86: 111-116.

Collins, R. and Margolies, D. 1994. The effect of interspecific mating on sex ratios in the

twospotted spider mite and the Banks grass mite (Acarina: Tetranychidae). –

J. Insect Behav., 8: 189–206.

Costa-Schmidt, L. E. and Machado, G. 2012. Reproductive interference between two

sibling species of gift-giving spiders. – Anim. Behav. 84: 1201-1211.

Crampton, W. G. et al. 2011. Reproductive character displacement and signal ontogeny

in a sympatric assemblage of electric fish. – Evolution 65: 1650-1666.

Crowder, D. W. et al. 2010. Plasticity in mating behaviour drives asymmetric

reproductive interference in whiteflies. – Anim. Behav. 79: 579-587.

Del Re, A. C. 2013. compute.es: Compute Effect Sizes. R package version 0.2-2.

http://cran.r-project.org/web/packages/compute.es.

delBarco-Trillo, J. and Johnston R. 2010. Adult female hamsters require long and

sustained exposures to heterospecific males to avoid interspecific mating. – Evol. Ecol.

25: 391–401.

Escudero, L. and Ferragut, F. 2005. Life-history of predatory mites Neoseiulus

californicus and Phytoseiulus persimilis (Acari: Phytoseiidae) on four spider mite

species as prey, with special reference to Tetranychus evansi (Acari: Tetranychidae). –

Biol. Control 32: 378-384.

Ferragut, F. et al. 2013. The invasive spider mite Tetranychus evansi (Acari:

Tetranychidae) alters community composition and host-plant use of native relatives. –

Exp. Appl. Acarol. 60: 321-341.

Godinho, D. P. et al. 2016. Down-regulation of plant defence in a resident spider mite

species and its effect upon con-and heterospecifics. – Oecologia 180: 161-167.

Gotoh, T. et al. 2015. Development and reproduction of five Tetranychus species

(Acari: Tetranychidae): Do they all have the potential to become major pests? – Exp.

Appl. Acarol. 66: 453-479.

Grbić, M. et al. 2011. The genome of Tetranychus urticae reveals herbivorous pest

adaptations. – Nature 479: 487-492.

Gröning, J. and Hochkirch, A. 2008. Reproductive interference between animal species.

– Q. Rev. Biol. 83: 257-282.

Gumm, J. M. and Gabor, C. R. 2005. Asexuals looking for sex: conflict between species

and mate-quality recognition in sailfin mollies (Poecilia latipinna). – Behav. Ecol.

Sociobiol. 58: 558–565.

Helle, W. 1967. Fertilization in the two-spotted spider mite (Tetranychus urticae:

Acari). – Entomol. Exp. Appl. 10: 103–110.

Hochkirch A., Gröning J. & Bücker A. 2007. Sympatry with the devil: reproductive

interference could hamper species coexistence. Journal of Animal Ecology 76(4): 633–

642.

Hurtado M. A. et al. 2008. Sequence analysis of the ribosomal internal transcribed

spacers region in spider mites (Prostigmata: Tetranychidae) occurring in citrus orchards

in Eastern Spain: use for species discrimination. – Ann. Appl. Biol. 153: 167-174.

Kishi, S. 2015. Reproductive interference in laboratory experiments of interspecific

competition. – Popul. Ecol. 57: 283-292.

Kishi, S. et al. 2009. Reproductive interference determines persistence and exclusion in

species interactions. – J. Anim. Ecol. 78: 1043-1049.

Kishi, S. and Nakazawa, T. 2013. Analysis of species coexistence co-mediated by

resource competition and reproductive interference. – Popul. Ecol. 55: 305-313.

Kyogoku, D. and Nishida, T. 2013. The mechanism of the fecundity reduction in

Callosobruchus maculatus caused by Callosobruchus chinensis males. – Popul. Ecol. 55:

87-93.

Kyogoku, D. 2015. Reproductive interference: ecological and evolutionary

consequences of interspecific promiscuity. – Popul. Ecol. 57: 253-260.

Lievens, E. J. et al. 2016. Maladaptive Sex Ratio Adjustment in the invasive brine

shrimp Artemia franciscana. – Curr. Biol. 26: 1463- 1467.

Macke, E. et al. 2012. Mating modifies female life history in a haplodiploid spider mite.

– Am. Nat. 179: E147-E162.

Matsuda, T. et al. 2013. DNA-based identification of spider mites: molecular evidence

for cryptic species of the genus Tetranychus (Acari: Tetranychidae). – J. Econ. Entomol.

106: 463-472.

McPeek, M.A., and Gavrilets, S. 2006. The evolution of female mating preferences:

differentiation from species with promiscuous males can promote speciation. –

Evolution 60: 1967-1980.

Nakagawa, S. and Poulin, R. 2012. Meta-analytic insights into evolutionary ecology: an

introduction and synthesis. – Evol. Ecol. 26: 1085–1099.

Navajas, M. 1998. Host plant associations in the spider mite *Tetranychus urticae* (Acari:

Tetranychidae): insights from molecular phylogeography. – Exp. Appl. Acarol. 22: 201-

214.

Navajas, M. et al. 2013. Review of the invasion of *Tetranychus evansi*: biology,

colonization pathways, potential expansion and prospects for biological control. – Exp.

Appl. Acarol. 59: 43-65.

Noriyuki, S. et al. 2012. Asymmetric reproductive interference between specialist and

generalist predatory ladybirds. – J. Anim. Ecol. 81: 1077-1085.

Qazi M. C. B. et al. 2003. The developments between gametogenesis and fertilization:

ovulation and female sperm storage in Drosophila melanogaster. – Dev. Biol. 256: 195–

211.

Sarmento, R. A. et al. 2011a. A herbivore that manipulates plant defence. – Ecol. Lett.

14: 229-236.

Sarmento, R. A. et al. 2011b. A herbivorous mite down-regulates plant defence and

produces web to exclude competitors. – PLoS One 6: 23757.

Sato, Y. et al. 2014. Testing for reproductive interference in the population dynamics of

two congeneric species of herbivorous mites. – Heredity 113: 495–502.

Sato, Y. et al. 2016. Why do males choose heterospecific females in the red spider

mite? – Exp. Appl. Acarol. 68: 21-31.

Satoh, Y. et al. 2001. Mating strategy of spider mite, *Tetranychus urticae* (Acari:

Tetranychidae) males: Postcopulatory guarding to assure paternity. – Appl. Entomol.

Zool. 36: 41– 45.

Schlupp, I. 2010. Mate Choice and the Amazon Molly: How Sexuality and Unisexuality

Can Coexist. – J. Hered. 101: S55–S61.

Servedio, M. R. and Kirkpatrick, M. 1997. The effects of gene flow on reinforcement. –

Evolution 51: 1764-1772.

Takafuji, A. 1986. Effectiveness of second mating for two incompatible types of the

citrus red mite, Panonychus citri (McGregor). – Res. Popul. Ecol. 28: 91-101.

Takafuji, A. et al. 1997. Reproductive interference and its consequences for the

competitive interactions between two closely related Panonychus spider mites. – Exp.

Appl. Acarol. 21: 379-391.

Takakura, K. I. et al. 2015. Conflicting intersexual mate choices maintain interspecific

sexual interactions. – Popul. Ecol. 57: 261-271.

Vahed, K. 1998. The function of nuptial feeding in insects: review of empirical studies.

– Biol. Rev. Camb. Philos. 73: 43–78.

Vala, F. et al. 2000. *Wolbachia*-induced hybrid breakdown in the two--spotted spider

mite *Tetranychus urticae* Koch. – P. Roy. Soc. Lond. B Bio. 267: 1931-1937.

Viechtbauer, W. 2010. Conducting meta-analyses in R with the metafor package. – J.

Stat. Softw. 36: 1-48.

Wang, P. et al. 2010. Crossing experiments and behavioral observations reveal

reproductive incompatibility among three putative species of the whitefly *Bemisia*

tabaci. – Insect Sci. 17: 508-516.

Yassin, A. and David, J. R. 2016. Within-species reproductive costs affect the

asymmetry of satyrization in Drosophila. – J. Evol. Biol. 29: 455-60.

Xu, J. and Wang, Q. 2011. Seminal fluid reduces female longevity and stimulates egg

production and sperm trigger oviposition in a moth. – J. Insect Physiol. 57: 385-390.

Supporting Information

The following Supporting Information has been made available in the online version of

this article.

Table S1 |Statistical tests and contrasts for the comparisons of fecundity and offspring

sex ratio in crosses between conspecific and heterospecific males and females, using

data from Bloc 1 or Blocs 1+2.

Table S1 | Statistical tests and contrasts for the comparisons of fecundity and offspring sex ratio in crosses between conspecific and heterospecific males and females, using data from Bloc 1 or Blocs 1+2

	Matings	Bloc	Fecundity (F test)	Sex-ratio (Chisq test)
Matings with an invasive species				
T. urticae vs *T. evansi*				
With *T. urticae* females		Blocs 1+2		
Mating order			$F_{2,136} = 7.919$, P = 0.0006	$F_{2,109} = 188.740$, P < 0.0001
Mating interval			$F_{1,136} = 0.039$, P = 0.8440	$F_{1,108} = 39.6442$, P = 0.01
Mating order x Mating interval			$F_{2,136} = 6.026$, P = 0.0031	$F_{2,106} = 57.219$, P = 0.0087
		Bloc 1		
Mating order			$F_{2,116} = 5.4688$, P = 0.0054	$F_{2,98} = 196.758$, P < 0.0001
Mating interval			$F_{1,116} = 3.3351$, P = 0.0703	$F_{1,97} = 34.031$, P = 0.0158
Mating order x Mating interval			$F_{2,116} = 2.3097$, P = 0.1038	$F_{2,95} = 58.029$, P = 0.0075
With *T. evansi* females		Blocs 1+2		
Mating order			$F_{2,187} = 4.680$, P = 0.0104	$F_{2,114} = 25.230$, P = 0.0908
Mating interval			$F_{1,187} = 2.555$, P = 0.1116	$F_{1,113} = 0.233$, P = 0.8319
Mating order x Mating interval			$F_{2,187} = 4.977$, P = 0.0078	$F_{2,111} = 3.786$, P = 0.6931
		Bloc 1		
Mating order			$F_{2,116} = 0.4240$ P = 0.6554	$F_{2,78} = 18.669$, P = 0.2401
Mating interval			$F_{1,116} = 0.3022$, P = 0.5836	$F_{1,77} = 4.2233$, P = 0.4199
Mating order x Mating interval			$F_{2,116} = 1.8729$, P = 0.1583	$F_{2,80} = 8.2368$, P = 0.5294
Matings with a resident species				
T. urticae vs *T. ludeni*				
With *T. urticae* females		Blocs 1+2		
Mating order			$F_{2,144} = 6.997$, P = 0.0013	$F_{2,102} = 23.926$, P = 0.0844
Mating interval			$F_{1,144} = 2.598$, P = 0.1092	$F_{1,101} = 3.138$, P = 0.4168
Mating order x Mating interval			$F_{2,144} = 3.694$, P = 0.0273	$F_{2,99} = 10.769$, P = 0.3237
		Bloc 1		
Mating order			$F_{2,89} = 0.7683$, P = 0.4669	$F_{2,71} = 22.610$, P = 0.1185
Mating interval			$F_{1,89} = 0.8446$, P = 0.3606	$F_{1,70} = 3.968$, P = 0.3825
Mating order x Mating interval			$F_{2,89} = 1.6405$, P = 0.1997	$F_{2,68} = 7.271$, P = 0.4962
With *T. ludeni* females		Blocs 1+2		
Mating order			$F_{2,248} = 10.534$, P < 0.0001	$F_{2,155} = 33.710$, P = 0.1204
Mating interval			$F_{1,248} = 5.180$, P = 0.0237	$F_{1,154} = 20.150$, P = 0.1112
Mating order x Mating interval			$F_{2,248} = 14.098$, P < 0.0001	$F_{2,152} = 158.690$, P < 0.0001
		Bloc 1		
Mating order			$F_{2,88} = 0.3338$, P = 0.7171	$F_{2,73} = 26.670$, P = 0.3760
Mating interval			$F_{1,88} = 3.3770$, P = 0.06949	$F_{1,72} = 32.737$, P = 0.1231
Mating order x Mating interval			$F_{2,88} = 1.0712$, P = 0.3402	$F_{2,70} = 247.099$, P = 0.0003

Legend: "Tu": matings with T. urticae males. "Te": matings with T. evansi males. "Tl": matings with T. ludeni males. "0h" and "24h" indicate the time of occurrence of the second mating, i.e., if immediately after the first mating (0h) or 24h later. TuTu means that both mating events were with a T. urticae male. TuTe means that the first mating was with a T. urticae male and the second with a T. evansi male. The same logic applies to TeTe, TeTu, TlTl, TlTu and TuTl.

Fig.2

T. urticae versus *T. evansi*

Fig.3

T. urticae versus *T. ludeni*