Negative frequency-dependent selection is frequently confounding

Dustin Brisson

University of Pennsylvania, 3451 Walnut Street, Philadelphia, PA 19104, United States of America

Corresponding author:

Dr. Dustin Brisson

Department of Biology

University of Pennsylvania

Leidy Laboratories, 209

433 South University Avenue

Philadelphia PA 19104-6018

Phone: (215) 746-1731

Fax: (215) 898-8780

Email: dbrisson@sas.upenn.edu

ABSTRACT

The existence of persistent genetic variation within natural populations presents an evolutionary problem as natural selection and genetic drift tend to erode genetic diversity. Models of balancing selection were developed to account for the high and sometimes extreme levels of polymorphism found in many natural populations. Negative frequency-dependent selection may be the most powerful selective force maintaining balanced natural polymorphisms but it is also commonly misinterpreted. The aim of this review is to clarify the processes underlying the negative frequency-dependent selection model, describe classes of natural polymorphisms that can and cannot result from these processes, and discuss observational and experimental data that can aid in accurately identifying the processes that generated or are maintain diversity in nature. Finally, I consider the importance of accurately describing the processes affecting genetic diversity within populations as it relates to research progress.

INTRODUCTION

Natural diversity - the "endless forms most beautiful and most wonderful" [1] - is an enduring focus of both evolutionary biologists and nature lovers. The evolutionary processes that have generated or are maintaining many examples of diversity in nature, however, remain obscure and can be controversial [2]. The processes that result in persistent polymorphisms within populations demand a special explanation as both directional natural selection and genetic drift should eliminate alleles and thus erode genetic diversity [3–5]. Nevertheless, many examples of persistent polymorphisms occur in nature [6–11]. Models of balancing selection - including negative frequency-dependent selection, spatial or temporal habitat heterogeneity, and heterozygote advantage - provide theoretical frameworks of the processes that can account for persistent polymorphisms within populations. A core tenet of each balancing selection model is that the selective value of an allele – whether it is beneficial or detrimental – is dependent on the environmental context [12,13]. That is, alleles are advantageous and deleterious in different ecological contexts.

Negative frequency-dependent selection has been called the most powerful selective force maintaining balanced polymorphisms [14–17] with some proposing that a large proportion of natural genetic polymorphisms are maintained by selection favoring rare alleles [18]. In models of negative frequency-dependent selection, the selective value of an allele is dependent on its relative abundance in the population such that Darwinian fitness increases as the relative abundance, or frequency, of the allele decreases [19]. Thus, negative frequency-dependent selection can maintain genetic polymorphisms within populations because relatively rare alleles have a selective advantage over more common alleles and thus tend to increase in frequency and avoid local extinction.

Numerous environmental interactions can result in a selective advantage for relatively rare alleles including sexual selection, parasite or predator preferences, and resource competition. In fact, each

of these mechanisms has been shown to create a selective advantage for rare alleles that has resulted in persistent polymorphisms in multiple natural populations [10,19–22]. In a classic example, color polymorphisms are maintained in natural populations of *Cepaea nemoralis* snails by negative frequency-dependent selection because their predators, the song thrush (*Turdus philomelos*), form a search image for the most common morph causing a much greater predation pressure on the common than the rare morph [21,23]. The rare morph can increase in frequency due to the relaxed predation pressure until it becomes common, resulting in a search image switch that now targets the new common morph, a process that maintains this polymorphism in *C. nemoralis* populations. Two luminaries in population genetics – R. Fisher and S. Wright – have also demonstrated the power of negative frequency-dependent selection to maintain diversity in natural systems. Wright famously demonstrated that self-incompatibility alleles, a genetic mechanism in plants to prevent inbreeding, are incredibly diverse because pollen containing a rare allele is more likely to find a receptive mate than pollen containing a common allele [24–26]. Thus, plants with rare alleles have a selective advantage (**Figure 1**). Similarly, Fisher's principle demonstrates that males and females are equally frequent because, if one sex were more frequent, the alternate sex would enjoy a *per capita* reproductive advantage [20,27].

The many incontrovertible demonstrations of the power of negative frequency-dependent selection to maintain polymorphisms in nature have led some to suggest that it is a "pervasive" force maintaining natural diversity [28]. The pervasiveness of negative frequency-dependent selection is further supported by the perception that "nearly every [selective agent] works in a way liable to produce frequency-dependent selection of the kind that favours rare phenotypes and hinders common ones" [28]. Although negative frequency-dependent selection may be a "powerful, perhaps a dominant, factor maintaining genetic diversity" within populations [28], many natural polymorphisms are maintained by other evolutionary processes [29–34]. Still, many natural

polymorphisms have been assumed to result from negative frequency-dependent selection even when the theoretical framework and data from the system are inconsistent with the processes of selection favoring relatively rare types. In this essay, I describe several patterns of allele dynamics that are commonly described in the literature as resulting from negative frequency-dependent selection despite data demonstrating that other explanatory processes are causative. These processes include allelic diversity resulting from directional selection within a changing ecological context, density dependent population regulation, other models of balancing selection, and aspects of community ecology. I will discuss concepts and experiments that can aid in identifying the processes underlying patterns of allele dynamics and suggest that accurately identifying the evolutionary process underlying natural patterns facilitates the development of hypotheses and future experiments to determine the ecological interactions or molecular mechanisms at the root of the process.

Directional selection attributed to negative frequency-dependent selection

As a broad concept, negative frequency-dependent selection may be the "most intuitively obvious explanation" of polymorphisms in nature [35]. However, the original concept becomes ambiguous, complex, and even controversial as a result of differing definitions and applications in both theoretical and empirical work [36]. Even some of the greatest thinkers in evolutionary biology have explained scenarios where the selective values of alleles are independent of their relative abundance through a negative frequency-dependent selection framework. A prominent example comes from a sweeping and influential essay by JBS Haldane where he suggested several "lines of thought" concerning infectious diseases as a major selective force in metazoan evolution [37]. Contrary to his assertion that "many or all" of these ideas "may prove to be sterile," most have been "followed profitably" (very profitably indeed). However, the negative frequency-dependent selection framework described in this essay appears to be one of the few unsound lines of thought. In this

framework, Haldane suggested that a host with a rare defensive phenotype has a selective advantage in the face of highly-adapted pathogens, "For just because of its rarity it will be resistant to diseases which attack the majority of its fellows." That is, the adapted pathogen has evolved mechanisms to overcome the common defensive phenotypes in host populations but cannot overcome rare defensive phenotypes. Thus, hosts expressing rare but effective defensive phenotypes enjoy a *per capita* fitness advantage over hosts expressing common but exploitable defenses.

The scenario described by Haldane, however, confounds natural selection favoring a specific (*effective*) phenotype in the current environment with a selective advantage resulting from rarity. Haldane's escape variants have a selective advantage because they cannot be subverted by the pathogen, not because they are rare. Further, the novel defensive phenotype has not yet lost its efficacy against the pathogen not because it is rare, but because it is novel. This point can be illustrated by extending this line of thought to allow migration of many individuals expressing a novel and effective defensive phenotype. These migrants would enjoy the same *per capita* selective advantage over the previously common resident phenotype, regardless of frequency of the novel phenotype in the population immediately following the mass-migration event. The evolutionary dynamics occurring in this framework do not occur because of rare advantage and, in most cases, will not result in a balanced polymorphism. These evolutionary dynamics are more likely the result of directional selection in a continuously changing environment [38–42]. These two processes - negative frequency-dependent selection and selection in a changing environment - can potentially be distinguished by artificially manipulating variant frequencies or by introducing a previously common but now extinct variant into a controlled population.

The genetic diversity of haemagglutinin (HA) glycoproteins in the influenza virus is another conspicuous example of selection in a changing environment often confounded with negative frequency-dependent selection. The dynamics of HA alleles change over time such that rare alleles

enter the population, rise to high population sizes, and subsequently decline toward extinction [43–45]. The strains expressing a numerically common allele have low fitness and decline in frequency because there are few hosts still susceptible to this strain as hosts acquire immunity to strains with which they have been previously infected [46–48]. By contrast, strains expressing numerically rare alleles have many susceptible hosts available and enjoy high rates of secondary infections per infected host causing an increase in frequency [47,48]. While there is undoubtedly strong selection at the HA locus, the selective advantage is derived not from relative rarity but from antigenic novelty [47,49–51], similar to Haldane's example. The presence or frequency of alternative HA alleles does not affect the fitness or temporal dynamics other alleles. That is, the population dynamics of a numerically rare allele is the same if the host population is already plagued by other numerically common strains (0.0001% when one novel allele enters a population of 10^6 infected hosts) and if it enters a host population in which no other influenza strain is circulating (100% when one novel allele enters a previous uninfected host population) (**Figure 2**). As the selective value of the allele is conditioned on the absolute abundance - but not the relative abundance - of the allele, it is unlikely that negative frequency-dependent selection is the evolutionary process underlying the polymorphism commonly observed at the HA locus. More likely, the common type is changing its own environment such that there are few susceptible hosts in which new infections can establish, but is not affecting the environment of alternative variants.

Density dependent fitness dynamics attributed to negative frequency-dependent selection

A recent luminary in evolutionary biology, R.C. Lewinton, suggested that negative frequency-dependent selection should be pervasive because, whenever "a genotype is its own worst enemy, its fitness will decrease as it becomes more common" [3]. As similar variants occupy similar niches and are commonly their own worst enemy, this logic suggests that negative frequency-dependent selection should indeed be pervasive. However, "common" in this case refers not to relative

abundance but absolute abundance. For example, the *per capita* fitness of individuals within a monomorphic population, one in which the frequency of a genotype is always at 100%, decreases as it "becomes more common" in absolute abundance. Further, relatively rare types suffer negative fitness effects in proportion to the absolute abundance of their numerically common competitors such that relative rarity may not provide a selective advantage.

There is an extensive literature describing the dependence of *per capita* fitness on absolute abundance [52–56]. The above scenario can be characterized using classical Logistic growth models that include competition among variants such that "a genotype is its own worst enemy" (Lotka-Volterra models) (**eq. 1**). These population growth models depend on the absolute abundance of each variant with respect to the carrying capacity (K), but are not conditioned on relative frequency.

$$\frac{dN_1}{dt} = r_1 N_1 \left(1 - \frac{N_1 + \alpha_{12} N_2}{K_1}\right)$$

$$\frac{dN_2}{dt} = r_2 N_2 \left(1 - \frac{N_2 + \alpha_{21} N_1}{K_2}\right)$$

eq. 1

It is often challenging to distinguish the effect of numerical rarity from relatively rarity on the selective value of an allele through observations of patterns of allelic diversity. Experimental manipulations of the carrying capacity (K), potentially through resource supplementation, can assuage the reductions in fitness experienced by common types that result from high densities without altering relative frequencies. In these experiments, the relative fitness of common types should increase if the effects are associated with density while the relative fitness of the common and rare types should not be altered if the allelic diversity is maintained by negative frequency-dependent selection.

Multiple niche polymorphisms attributed to negative frequency-dependent selection

In the multiple niche selection model of balancing selection, the selective value of an allele is conditioned on their ability to exploit different environmental features in a heterogeneous habitat

[57,58]. Multi-niche selection maintains multiple variants in a population if each variant has a fitness advantage in some available habitats while other variants are superior in other habitats. This idea – that environmentally variable selection can result in balanced polymorphisms – has a long history in the literature in which the foundational idea is stated by Dobzhansky [12]. Although incontrovertible examples of multi-niche selection maintaining polymorphism in natural populations are relatively rare, correct inference of the process resulting in balancing selection is necessary to generate hypotheses and controlled experiments to determine the underlying ecological interactions or molecular mechanisms causing the process.

The study of pattern, in isolation from the evolutionary processes that generated it, is not likely to advance general theories nor an understanding of a specific system [59]. However, determining the processes responsible for balanced polymorphism patterns observed in nature is a difficult task [60–62,17]. The balanced polymorphism at the outer surface protein C (*ospC*) locus in populations of *Borrelia burgdorferi,* the cause of human Lyme disease, provides a fitting example. Although the function of OspC remains unclear [63–68], the within-population diversity at this locus bears all the hallmarks of balancing selection – large numbers of alleles in all local populations; allele frequencies that are more even than expected for neutrally evolving loci; and genetic evidence of an ancient polymorphism [33,69–73].

Negative frequency-dependent selection and multi-niche selection have both been proposed as processes maintaining the *ospC* polymorphisms, and both frameworks have empirical support [69,71,74–76]. The negative frequency-dependent selection model suggests that the polymorphism can be maintained if previously infected hosts are immune to subsequent infections by the same OspC variant but susceptible to novel variants, a molecular mechanism that has been demonstrated in laboratory animals [77,78,but see ,79]. However, in this scenario the frequency or even presence of alternative OspC variants does not affect the number of susceptible hosts for the common strain,

similar to the influenza example, arguing against negative frequency-dependent selection as an evolutionary process maintaining *ospC* polymorphisms. Further, negative frequency-dependent selection is most effective when few hosts remain susceptible to the common *ospC* types, a pattern that is not observed in natural data sets [33,80–83]. Studies investigating allelic diversity at *ospC* from natural hosts consistently demonstrate that most natural reservoir hosts, those that are regularly infected with *B. burgdorferi,* are rarely infected with all of the common *ospC* alleles [33,80,82,84]. Most hosts are, however, infected with a subset of the *ospC* alleles, as expected if each host species represented a different ecological niche [33,80,82,84]. Further, host individuals of the same species, including humans, are infected by the same subset of *ospC* alleles across both time and space [33,80,82,84–87]. The collective evidence suggests that the balanced *ospC* polymorphisms are more likely maintained by multi-niche selection - with each host species representing multiple niches [88], one for each *ospC* variant by which it can be infected - than by negative frequency-dependent selection. These results suggest that the mechanisms causing the balanced polymorphism are more likely to involve genotype-by-host species interactions than to involve a memory immune response mechanism that is conserved across vertebrate species.

It has been argued that "Selection in multiple niches is not an alternative to frequency-dependent selection...but a way of generating it" [28]. However, scenarios in which balanced polymorphisms can be maintained without a selective advantage favoring relatively rare variants are not uncommon, suggesting that these are two distinct evolutionary processes in at least some cases. To illustrate this point, image two variants occupy a heterogeneous habitat where each variant has a selective advantage in one niche but is disadvantaged in another, a classical multi-niche selection model (**Figure 3**). Here we assume that the carrying capacity in niche A is much lower than the carrying capacity in niche B (K_A=10; K_B=10^5). In this scenario, variant B - which has a competitive advantage in niche B - can retain a fitness advantage (a greater *per capita* growth rate) even when it is more

common than variant A - which has a competitive advantage in niche A. For example, in a population with 90 variant B individuals and 10 variant A individuals, variant B has a rapid *per capita* rate of increase while variant A does not increase (**Figure 3**). Here, the relatively common variant B has a "selective advantage" over the relatively rare variant A due to multi-niche selection, which is independent of negative frequency-dependent selection. Depending on the parameter values in this model, a balanced polymorphism can be maintained in the absence of rare advantage.

Community diversity attributed to negative frequency-dependent selection

There is a rich empirical and theoretical history describing the causes and consequences of species diversity within ecological communities [2,89–93]. Mechanisms of coexistence function in two major ways: *equalizing* mechanisms minimize the average fitness differences between species while *stabilizing* mechanisms increase negative *intra*specific interactions relative to negative *inter*specific interactions [2]. Stabilizing mechanisms promote species coexistence and include mechanisms such as resource partitioning and frequency-dependent predation, as well as mechanisms that depend on spatial or temporal fluctuations in population densities or environmental factors. Equalizing mechanisms contribute to stable coexistence when they reduce large average fitness inequalities which might negate the effects of stabilizing mechanisms [2]. While some natural forces that affect the maintenance of community diversity have frequency dependent mechanisms, this should not be mistaken for negative frequency-dependent selection which maintains polymorphisms within populations. Applying models of natural selection to groups above the population level should be exercised only with the greatest caution [94].

The 'Killing the Winner' hypothesis is a recent endeavor to understand patterns of diversity within communities through a negative frequency-dependent selection framework [95,96]. More recent versions of the Killing the Winner hypothesis suggest that a frequency-dependent functional

response in predator populations can promote community diversity. However, the functional response in this hypothesis is often not conditioned on the frequency of the species but on the presence or absence of character traits of the species that are being targeted by predators [96–99]. The "winner" in the Killing the Winner hypothesis refers to species that invest resource into reproduction at the expense of investing in predator defenses, which may or may not correspond to the most frequent species [97]. In these cases, neither the relative nor the absolute abundance of the prey species affects the functional responses of the predator.

Conclusions

Understanding the processes that produce or maintain diversity in natural populations is a central challenge in evolutionary biology. Negative frequency-dependent selection maintains many noted and striking polymorphisms in nature [16,22,100–104], and many polymorphisms exist in the absence of a selective advantage favoring rare types [33,29–32,34]. Ideally, one could unequivocally determine the causative process through observations of the patterns of variation in nature. Unfortunately, many processes result in identical patterns, especially when those patterns are observed over short time scales. In some cases, long-term observations of allelic dynamics can distinguish polymorphisms caused by mutation-selection balance or selection in a changing environment from a stable polymorphism resulting from balancing selection (some of these? [32,105–109]). Evidence suggesting negative frequency-dependent selection - such as allelic cycles where each allele gains a selective advantage as it becomes more rare - may also be observed from long-term observational studies [22,110]. The patterns resulting from various evolutionary processes can also be tested through controlled and natural experiments such as manipulating allele frequencies in sub-populations [32,105,107,109].

Ecological and molecular mechanisms are rarely deducible from patterns [111], but accurate identification of the evolutionary processes causing the pattern can generate hypotheses about these mechanisms. For example, the northern acorn barnacle, *Semibalanus balanoides*, shows clear evidence of a balanced polymorphism at the mannose-6-phosphate isomerase (*mpi*) locus [112,113]. The pattern of *mpi* genotype frequencies among intertidal microhabitats, where one allele is common in high intertidal zones but rare in low intertidal zones, suggests that multi-niche selection maintains this polymorphism [114]. Experimental manipulations of genotypes among microhabitats confirmed that multi-niche selection is the process responsible for the allelic variation [32,115]. The molecular mechanism linking mannose utilization with survivorship in high intertidal zones, where temperature and desiccation stress is high, was subsequently elucidated through controlled laboratory experiments [106]. As this and many other examples demonstrate, the ecological interaction or molecular mechanism underlying an evolutionary process can best be understood when the evolutionary process is accurately determined.

Figure 1. A. A cartoon depiction of the self-incompatibility allele model modified from [19]. For simplicity, the plant population represented has only two alleles although most populations maintain 10s or 100s of *S*-alleles. As describe in the classical model [19], S_1S_1 homozygote plants produce only pollen containing the S_1 allele and can be pollenated only by pollen with the S_2 allele. Similarly, S_2S_2 homozygotes produce S_2-containing pollen and can be pollenated only by S_1-containing pollen. Heterozygote plants can produce pollen with either allele but cannot be pollenated by either pollen type. The allele that is relatively rare in the population has a selective advantage over the common allele as pollen containing the rare allele is much more likely to pollenate a receptive ovule. In contrast, pollen containing the common allele is likely to attempt to pollenate a plant containing the common allele and be rejected, resulting in limited breeding success. **B.** The temporal dynamics of the alleles in this system are likely to fluctuate as expected when rare alleles have a selective advantage. For example, if 81% of the plants are homozygous S_1S_1 (time 0), the S_2-containing pollen (~10%) has a high probability of finding an S_1S_1 plant and successfully breeding. By contrast, the S_1-containing pollen is highly unlikely to find a S_2S_2 homozygote (~1% of all plants), resulting in very low breeding success. Due to the limited breeding success, the S_1 allele will decrease in frequency until S_1-containing pollen is rare and becomes more likely to find a receptive mate. These dynamics occur because a pollen grain with a common allele will be limited in terms of mates, while a pollen grain with a rare allele will not. Hence, plants with rare alleles have a selective advantage in terms of mating (see model by [19]).

Figure 2. Influenza virus carrying rare HA or NA alleles do not have a selective advantage because they are relatively rare – a necessary condition of negative frequency-dependent selection – but because they are numerically rare compared to the number of susceptible hosts. **A.** The population dynamics of two influenza strains (dark and light lines) that enter a host population sequentially. Both strains increase when they are numerically rare, but not relatively rare, and decrease after they become numerically common. For example, the maximal rate of increase of the first strain occurs prior to the second strain entering the population, despite remaining at a maximal relative abundance (100%). **B.** The relative frequencies of the two influenza strains through time. If negative frequency-dependent selection were affecting the relative abundances of these strains, the common strain at time=0 (dark line) should have lower fitness than the rare strain (light line). However, the *per capita* rate of increase of the common strain remains high until it has substantially reduced the number of susceptible hosts, regardless of its relative abundance. The arrows indicate expected affect of negative frequency-dependent selection on the fitness of each strain given its relative abundance. Red arrows indicate the time periods when the expectations of negative frequency-dependent selection are not satisfied; black arrows indicate time periods when negative frequency-dependent selection expectations are satisfied. **C.** The *per capita* rate of increase and the population dynamics of each strain have the same temporal patterns in the absence of the alternative strain. Strain 1 remains at 100% frequencies throughout the time period, suggesting that relative abundance does not drive of changes in fitness.

Figure 3. Multi-niche selection is an alternative model of balancing selection that does not depend on the core assumption of negative frequency-dependent selection models that fitness is a function of relative frequency in the population. Shown is a simulation where variant A has high fitness in niche A while variant B has high fitness in niche B (Supplemental material). However, variant A has low fitness in niche B while variant B has low fitness in niche A. Additionally, the carrying capacity in niche A is much lower than in niche B (K_A =10, K_B=10000). At the start of the simulation, there are 10 variant A individuals (10% of the population) and 90 variant B individuals (10% of the population), yet the *per capita* fitness of variant A individuals much lower than for variant B individuals. In the negative frequency-dependent selection model, the frequency of variant A should increase as it is currently less frequent than variant B. Although the conditions of negative frequency-dependent selection are not satisfied, both variants can be maintained in the population due to the selective advantage each enjoys in their preferred niche. Parameters used in the simulation *per capita* growth rate = 0.35, death rate in preferred niche = 0.05, death rate in non-preferred habitat =0.25, migration among niches = 0.01.

References cited

Darwin, C. (1859) *The Origin of Species*, Penguin Books.
Chesson, P. (2000) Mechanisms of maintenance of species diversity. *Annu. Rev. Ecol. Syst.* 31, 343–66
Lewontin, R.C. (1974) *The genetic basis of evolutionary change*, Columbia University Press.
Charlesworth, B. and Hughes, K.A. (2000) The maintenance of genetic variation in life-history traits. In *Evolutionary Genetics: from Molecules to Morphology* (Singh, R. S. and Krimbas, C. B., eds), pp. 369–392, Cambridge University Press
Nielsen, R. (2005) Molecular signatures of natural selection. *Annu. Rev. Genet.* 39, 197–218
Mallet, J. and Joron, M. (1999) Evolution of diversity in warning color and mimicry: polymorphisms, shifting balance, and speciation. *Annu. Rev. Ecol. Syst.* 30, 201–233
Richman, A. (2000) Evolution of balanced genetic polymorphism. *Mol. Ecol.* 9, 1953–1963
Hedrick, P.W. (1986) Genetic-polymorphism in heterogeneous environments - a decade later. *Annu. Rev. Ecol. Syst.* 17, 535–566
Hedrick, P.W. *et al.* (2002) Major histocompatibility complex variation in red wolves: Evidence for common ancestry with coyotes and balancing selection. *Mol. Ecol.* 11, 1905–1913
Delph, L.F. and Kelly, J.K. (2014) On the importance of balancing selection in plants. *New Phytol.* 201, 45–56
Carius, H.J. *et al.* (2001) Genetic variation in a host-parasite association: Potential for coevolution and frequency-dependent selection. *Evolution* 55, 1136–1145
Dobzhansky, T. (1982) *Genetics and the origin of species*, Columbia University Press.
Clarke, B.C. *et al.* (1988) Frequency-dependent selection, metrical characters and molecular evolution. *Philos. Trans. R. Soc. B.* 319, 631–640
Turelli, M. and Barton, N.H. (2004) Polygenic variation maintained by balancing selection: Pleiotropy, sex-dependent allelic effects and G × E interactions. *Genetics* 166, 1053–1079
Ayala, F.J. and Campbell, C.A. (1974) Frequency-dependent selection. *Annu. Rev. Ecol. Syst.* 5, 115–138
Fitzpatrick, M.J. *et al.* (2007) Maintaining a behaviour polymorphism by frequency-dependent selection on a single gene. *Nature* 447, 210–212
Kazancioǧlu, E. and Arnqvist, G. (2014) The maintenance of mitochondrial genetic variation by negative frequency-dependent selection. *Ecol. Lett.* 17, 22–27
Kojima, K. and Yarbrough, K.M. (1967) Frequency-dependent selection at the *esterase-6-locus* in *Drosophila melanogaster. Proc. Natl. Acad. Sci.* 57, 645–649
Wright, S. (1939) The distribution of self-sterility alleles in populations. *Genetics* 24, 538–552
Fisher, R.A. (1930) *The genetical theory of natural selection*, Oxford University Press.
Harvey, P.H. *et al.* (1975) The effect of experience on the selective behaviour of song thrushes feeding on artificial populations of *Cepaea. Genetica* 45, 211–216
Gigord, L.D. *et al.* (2001) Negative frequency-dependent selection maintains a dramatic flower color polymorphism in the rewardless orchid *Dactylorhiza sambucina* (L.) Soo. *Proc. Natl. Acad. Sci.* 98, 6253–6255
Allen, J.A. (1988) Frequency-dependent selection by predators. *Philos. Trans. R. Soc. B.* 319, 485–503
Castric, V. and Vekemans, X. (2004) Plant self-incompatibility in natural populations: A critical assessment of recent theoretical and empirical advances. *Mol. Ecol.* 13, 2873–2889
Wright, S. (1964) The distribution of self-incompatibility alleles in populations. *Evolution* 18, 609–619

Wright, S. (1969) *Evolution and the Genetics of Populations, Vol II. The theory of gene frequencies.*, University Of Chicago Press.

Edwards, A.W.F. (1998) Natural selection and the sex ratio: Fisher's sources. *Am. Nat.* 151, 564–569

Clarke, B.C. (1979) The evolution of genetic diversity. *Proc. R. Soc. B.* 205, 453–474

Weatherall, D.J. (1997) Thalassaemia and malaria, revisited. *Ann. Trop. Med. Parasitol.* 91, 885–90

Allison, A.C. (1954) Protection afforded by sickle-cell trait against subtertian malareal infection. *Br. Med. J.* 1, 290–294

Smith, T.B. (1990) Natural selection on bill characters in the two bill morphs of the African finch, *Pyrenestes ostrinus. Evolution* 44, 832–842

Schmidt, P.S. *et al.* (2000) Environmental heterogeneity and balancing selection in the acorn barnacle, *Semibalanus balanoides. Philos. Trans. R. Soc. B.* 267, 379–384

Brisson, D. and Dykhuizen, D.E. (2004) *ospC* diversity in *Borrelia burgdorferi*: Different hosts are different niches. *Genetics* 168, 713–722

Barton, N. (1986) The maintenance of polygenic variation through a balance between mutation and stabilizing selection. *Genet. Res.* 47, 209–216

Trotter, M. V. and Spencer, H.G. (2007) Frequency-dependent selection and the maintenance of genetic variation: Exploring the parameter space of the multiallelic pairwise interaction model. *Genetics* 176, 1729–1740

Heino, M. *et al.* (1998) The enigma of frequency-dependent selection. *Trends Ecol. Evol.* 13, 367–370

Haldane, J.B.S. (1949) Disease and evolution. *La Ric. Sci. Suppl.* 19, 1–11

Collins, S. *et al.* (2007) Adaptive walks toward a moving optimum. *Genetics* 176, 1089–1099

Orr, H.A. (2005) Theories of adaptation: What they do and don't say. *Genetica* 123, 3–13

Lande, R. and Shannon, S. (1996) The role of genetic variation in adaptation and population persistence in a changing environment. *Evolution* 56, 429

Bell, G. (2010) Fluctuating selection: the perpetual renewal of adaptation in variable environments. *Philos. Trans. R. Soc. B.* 365, 87–97

Levins, R. (1968) *Evolution in changing environments*, Princeton University Press.

Andreasen, V. (2003) Dynamics of annual influenza A epidemics with immuno-selection. *J. Math. Biol.* 46, 504–536

Lin, J. *et al.* (2003) Traveling waves in a model of influenza A drift. *J. Theor. Biol.* 222, 437–445

Earn, D.J.D. *et al.* (2002) Ecology and evolution of the flu. *Trends Ecol. Evol.* 17, 334–340

Pease, C.M. (1987) An evolutionary epidemiological mechanism, with applications to type a influenza. *Theor. Popul. Biol.* 31, 422–452

Virseda, S. *et al.* (2010) Seasonal and pandemic A (H1N1) 2009 influenza vaccination coverage and attitudes among health-care workers in a Spanish University Hospital. *Vaccine* 28, 4751–4757

Stegeman, A. *et al.* (2004) Avian influenza A virus (H7N7) epidemic in the Netherlands in 2003: Course of the epidemic and effectiveness of control measures. *J. Infect. Dis.* 190, 2088–2095

Plotkin, J.B. and Dushoff, J. (2003) Codon bias and frequency-dependent selection on the hemagglutinin epitopes of influenza A virus. *Proc. Natl. Acad. Sci.* 100, 7152–7157

Cherry, J.L. *et al.* (2009) Evolutionary dynamics of N-glycosylation sites of influenza virus Hemagglutinin. *PLoS Curr.* DOI: 10.1371/currents.RRN1001

Nelson, M.I. and Holmes, E.C. (2007) The evolution of epidemic influenza. *Nat. Rev. Genet.* 8, 196–205

Emlen, J.M. (1985) The assessment of frequency- and density-dependent influences on fitness in natural populations. *Am. Nat.* 125, 507–520

Roughgarden, J. (1971) Density-dependent natural selection. *Ecology* 52, 453–468

MacArthur, R.H. *et al.* (1967) *The theory of island biogeography*, Princeton University Press.

MacArthur, R.H. (1962) Some generalized theorums of natural selection. *Genetics* 48, 1893–1897

Birch, L.C. (1955) Selection in *Drosophila pseudoobscura* in relation to crowding. *Evolution* 9, 389–399

Levene, H. (1953) Genetic equilibrium when more than one ecological niche is available. *Am. Nat.* 87, 331–333

Ravigne, V. *et al.* (2004) Implications of habitat choice for protected polymorphisms. *Evol. Ecol. Res.* 6, 125–145

Cale, W. *et al.* (1989) Inferring process from pattern in natural communities. *Bioscience* 39, 600–605

Barrett, J.A. (1988) Frequency-dependent selection in plant fungal interactions. *Philos. Trans. R. Soc. B.* 319, 473–483

Laine, A.L. *et al.* (2011) Spatial variation in disease resistance: From molecules to metapopulations. *J. Ecol.* 99, 96–112

Chaboudez, P. and Burdon, J.J. (1995) Frequency-dependent selection in a wild plant-pathogen system. *Oecologia* 102, 490–493

Onder, O. *et al.* (2012) OspC is potent plasminogen receptor on surface of *Borrelia burgdorferi*. *J. Biol. Chem.* 287, 16860–16868

Pal, U. *et al.* (2004) OspC facilitates *Borrelia burgdorferi* invasion of *Ixodes scapularis* salivary glands. *J. Clin. Invest.* 113, 220–230

Tilly, K. *et al.* (2006) *Borrelia burgdorferi* OspC protein required exclusively in a crucial early stage of mammalian infection. *Infect. Immun.* 74, 3554–3564

Xu, Q. *et al.* (2007) Identification of an *ospC* operator critical for immune evasion of *Borrelia burgdorferi*. *Mol. Microbiol.* 64, 220–231

Tilly, K. *et al.* (2013) Lipoprotein succession in *Borrelia burgdorferi*: Similar but distinct roles for OspC and VlsE at different stages of mammalian infection. *Mol. Microbiol.* 89, 216–227

Carrasco, S.E. *et al.* (2015) Outer surface protein OspC is an antiphagocytic factor that protects *Borrelia burgdorferi* from phagocytosis by macrophages. *Infect. Immun.* 83, 4848–4860

Qiu, W.G. *et al.* (1997) A population genetic study of *Borrelia burgdorferi* sensu stricto from eastern Long Island, New York, suggested frequency-dependent selection, gene flow and host adaptation. *Hereditas* 127, 203–216

Qiu, W.G. *et al.* (2002) Geographic uniformity of the Lyme disease spirochete (*Borrelia burgdorferi*) and its shared history with tick vector (*Ixodes scapularis*) in the northeastern United States. *Genetics* 160, 833–849

Wang, I.N. *et al.* (1999) Genetic diversity of *ospC* in a local population of *Borrelia burgdorferi* sensu stricto. *Genetics* 151, 15–30

Charlesworth, B. *et al.* (1997) The effects of local selection, balanced polymorphism and background selection on equilibrium patterns of genetic diversity in subdivided populations. *Genet. Res.* 70, 155–174

May, G. *et al.* (1999) The signature of balancing selection: Fungal mating compatibility gene evolution. *Proc. Natl. Acad. Sci.* 96, 9172–9177

Brisson, D. *et al.* (2012) Genetics of *Borrelia burgdorferi*. *Annu. Rev. Genet.* 46, 515–536

Haven, J. *et al.* (2011) Pervasive recombination and sympatric genome diversification driven by frequency-dependent selection in *Borrelia burgdorferi*, the Lyme disease bacterium. *Genetics*

189, 951–966

Seifert, S.N. *et al.* (2015) Evolution and population genomics of the Lyme borreliosis pathogen, *Borrelia burgdorferi. Trends Genet.* 31, 201–207

Gilmore, R.D. *et al.* (1996) Outer surface protein C (OspC), but not P39, is a protective immunogen against a tick-transmitted *Borrelia burgdorferi* challenge: Evidence for a conformational protective epitope in OspC. *Infect. Immun.* 64, 2234–2239

Probert, W.S. *et al.* (1997) Immunization with outer surface protein (Osp) A, but not OspC, provides cross-protection of mice challenged with North American isolates of *Borrelia burgdorferi. J. Infect. Dis.* 175, 400–405

Devevey, G. *et al.* (2015) First arrived takes all: inhibitory priority effects dominate competition between co-infecting *Borrelia burgdorferi* strains. *BMC Microbiol.* 15, 61

Vuong, H.B. *et al.* (2014) Occurrence and transmission efficiencies of *Borrelia burgdorferi ospC* types in avian and mammalian wildlife. *Infect. Genet. Evol.* 27, 594–600

States, S.L. *et al.* (2014) Lyme disease risk not amplified in a species-poor vertebrate community: Similar *Borrelia burgdorferi* tick infection prevalence and OspC genotype frequencies. *Infect. Genet. Evol.* 27, 566–575

Hanincova, K. *et al.* (2006) Epidemic spread of Lyme borreliosis, northeastern United States. *Emerg. Infect. Dis.* 12, 604–611

Ogden, N.H. *et al.* (2008) Role of migratory birds in introduction and range expansion of *Ixodes scapularis* ticks and of *Borrelia burgdorferi* and *Anaplasma phagocytophilum* in Canada. *Appl. Environ. Microbiol.* 74, 1780–1790

Mechai, S. *et al.* (2016) Evidence for host-genotype associations of *Borrelia burgdorferi* sensu stricto. *PLoS One* 11, e0149345

Seinost, G. *et al.* (1999) Four clones of *Borrelia burgdorferi* sensu stricto cause invasive infection in humans. *Infect. Immun.* 67, 3518–3524

Wormser, G.P. *et al.* (2008) *Borrelia burgdorferi* genotype predicts the capacity for hematogenous dissemination during early Lyme disease. *J. Infect. Dis.* 198, 1358–1364

Dykhuizen, D.E.D.E. *et al.* (2008) The propensity of different *Borrelia burgdorferi* sensu stricto genotypes to cause disseminated infections in humans. *Am. J. Trop. Med. Hyg.* 78, 806–810

Brisson, D. *et al.* (2011) Biodiversity of *Borrelia burgdorferi* strains in tissues of Lyme disease patients. *PLoS One* 6, e22926

Connell, J.H. and Orias, E. (1964) The ecological regulation of species diversity. *Am. Nat.* 98, 399–414

Schoener, T.W. (1974) Resource partitioning in ecological communities research on how similar species divide resources helps reveal the natural regulation of species diversity. *Science* 185, 27–39

Ricklefs, R.E. and Schluter, D. (1993) *Species diversity in ecological communities: historical and geographical perspectives*, University Of Chicago Press.

Wright, S.J. (2002) Plant diversity in tropical forests: a review of mechanisms of species coexistence. *Oecologia* 130, 1–14

Lubchenco, J. (1978) Plant species diversity in a marine intertidal community; imoprtance of herbivore food preference and algal competetive. *Am. Nat.* 112, 23–39

Williams, G.C. (1966) *Adaptation and Natural Selection*, Princeton University Press.

Thingstad, T.F. (2000) Elements of a theory for the mechanisms controlling abundance, diversity, and biogeochemical role of lytic bacterial viruses in aquatic systems. *Limnol. Oceanogr.* 45, 1320–1328

Thingstad, T.F. and Lignell, R. (1997) Theoretical models for the control of bacterial growth rate, abundance, diversity and carbon demand. *Aquat. Microb. Ecol.* 13, 19–27

Winter, C. *et al.* (2010) Trade-offs between competition and defense specialists among unicellular planktonic organisms: the "killing the winner" hypothesis revisited. *Microbiol. Mol. Biol. Rev.* 74, 42–57

Koskella, B. and Meaden, S. (2013) Understanding bacteriophage specificity in natural microbial communities. *Viruses* 5, 806–823

Suttle, C.A. (2007) Marine viruses-major players in the global ecosystem. *Nat. Rev. Microbiol.* 5, 801–812

Mitchell-Olds, T. *et al.* (2007) Which evolutionary processes influence natural genetic variation for phenotypic traits? *Nat. Rev. Genet.* 8, 845–856

Charlesworth, D. (2006) Balancing selection and its effects on sequences in nearby genome regions. *PLoS Genet.* 2, 379–384

Loisel, D.A. *et al.* (2006) Ancient polymorphism and functional variation in the primate MHC-DQA1 5' cis-regulatory region. *Proc. Natl. Acad. Sci.* 103, 16331–6

Kojima, K. and Tobari, Y. (1969) Selective modes associated with karyotypes in *Drosophila ananassae.* II. Heterosis and frequency-dependent selection. *Genetics* 63, 639–651

Mokkonen, M. *et al.* (2011) Negative frequency-dependent selection of sexually antagonistic alleles in *Myodes glareolus.* *Science* 334, 972–974

Koskella, B. and Lively, C.M. (2009) Evidence for negative frequency-dependent selection during experimental coevolution of a freshwater snail and a sterilizing trematode. *Evolution* 63, 2213–2221

Schmidt, P.S. (2001) The effects of diet and physiological stress on the evolutionary dynamics of an enzyme polymorphism. *Proc. R. Soc. B.* 268, 9–14

Roy, B.A. (1998) Differentiating the effects of origin and frequency in reciprocal transplant experiments used to test negative frequency-dependent selection hypotheses. *Oecologia* 115, 73–83

Siemens, D.H. and Roy, B.A. (2005) Tests for parasite-mediated frequency-dependent selection in natural populations of an asexual plant species. *Evol. Ecol.* 19, 321–338

Olendorf, R. *et al.* (2006) Frequency-dependent survival in natural guppy populations. *Nature* 441, 633–636

Thrall, P.H. *et al.* (2012) Rapid genetic change underpins antagonistic coevolution in a natural host-pathogen metapopulation. *Ecol. Lett.* 15, 425–435

Kershaw, K.A. (1963) Pattern in vegetation and its causality. *Ecology* 44, 377–388

McDonald, J.H. (1991) Contrasting amounts of geographical variation as evidence for direct selection: the *mpi* and *pgm* loci in eight crustacean species. *Heredity (Edinb).* 67, 215–219

Hoffmann, R.J. (1981) Evolutionary genetics of *Metridium senile.* I. Kinetic differences in phosphoglucose isomerase allozymes. *Biochem. Genet.* 19, 129–44

Schmidt, P.S. and Rand, D.M. (1999) Intertidal microhabitat and selection at Mpi: interlocus contrasts in the northern acorn barnacle, *Semibalanus glandula.* *Evolution* 53, 135–146

Schmidt, P.S. and Rand, D.M. (2001) Adaptive maintenance of genetic polymorphism in an intertidal barnacle: habitat- and life-stage-specific survivorship of Mpi genotypes. *Evolution* 55, 1336–1344

Figure 1 data generated in R

```
### 3 genotypes S1S2, S1S3, S2S3
### S1 pollen can only pollenate S2S3 plant, 50% of time get S1S2, 50% S1S3

##Outline ###
## Start with Plants at different freq
## A pollen grain is chosen at random (by percent each is found in the )
## pollen lands on 1 of the 100 plants (random number generator 1-100)
## Rejected if either allele of same plant is same as pollen
## if not rejected, makes baby plant with one of the 2 alleles (picked at random)
## repeat until 100 next generation plants
## repeat for 100 generations

#####  PARAMETERS ####
S1S2<- 500 ## Starting pop of variant S1S2
S1S3<- 450 ## Starting pop of variant S1S3
S2S3 <- 50  ## Starting pop of variant S2S3
PopSize<-S1S2+S1S3+S2S3
S1<- (S1S2+S1S3)/(2*(S1S2+S1S3+S2S3)) ## starting number of S1 pollen grains
S2<- (S1S2+S2S3)/(2*(S1S2+S1S3+S2S3)) ## starting number of S2 pollen grains
S3<- (S2S3+S1S3)/(2*(S1S2+S1S3+S2S3)) ## starting number of S3 pollen grains
S1S2ng<-0
S1S3ng<-0
S2S3ng<-0

#### HOUSEKEEPING STUFF #####
generations<-5
pollens<-1000000

S1vector <- {}
S2vector <- {}
S3vector <- {}

S1vector[1] <- (S1S2+S1S3)/(2*(S1S2+S1S3+S2S3))
S2vector[1] <- (S1S2+S2S3)/(2*(S1S2+S1S3+S2S3))
S3vector[1] <- (S1S3+S2S3)/(2*(S1S2+S1S3+S2S3))

#### START OF MODEL SIMULATION #####

for (gens in 2:generations) { ## number of generations loop
pS1S2<-S1S2/(S1S2+S1S3+S2S3)
pS1S3<-S1S3/(S1S2+S1S3+S2S3)
pS2S3<-S2S3/(S1S2+S1S3+S2S3)
```

```
        for (pol in 1:pollens) { ## mating loop
S1yes<-0
S2yes<-0
S3yes<-0
S1S2yes<-0
S2S3yes<-0
S1S3yes<-0

# choose pollen type

pollenRand <-runif(1)
if (pollenRand<=S1) {S1yes<-1}
if (pollenRand > S1+S2) {S3yes<-1}
if (S1yes==0 & S3yes==0) {S2yes<-1}

# choose plant type
PlantRand <-runif(1)
if (PlantRand<=pS1S2) {S1S2yes<-1}
if (PlantRand > pS1S2+pS1S3) {S2S3yes<-1}
if (S1S2yes==0 & S2S3yes==0) {S1S3yes<-1}

## next generation plants
if (S1yes == 1 & S2S3yes==1) {
        alleleSelect<-runif(1)
        if (alleleSelect<=.5){S1S2ng <- S1S2ng+1}
        else {S1S3ng <- S1S3ng+1}
}

if (S2yes == 1 & S1S3yes==1) {
        alleleSelect<-runif(1)
        if (alleleSelect<=.5){S1S2ng <- S1S2ng+1}
        else {S2S3ng <- S2S3ng+1}
}

if (S3yes == 1 & S1S2yes==1) {
        alleleSelect<-runif(1)
        if (alleleSelect<=.5){S1S3ng <- S1S3ng+1}
        else {S2S3ng <- S2S3ng+1}
}

#Stop when 100 babies
if (S1S2ng + S1S3ng + S2S3ng ==PopSize){break}

} ## mating loop

S1S2<-S1S2ng
```

```r
S1S3<-S1S3ng
S2S3<-S2S3ng
S1S2ng <-0
S1S3ng <-0
S2S3ng <-0

S1<- (S1S2+S1S3)/(2*(S1S2+S1S3+S2S3))
S2<- (S1S2+S2S3)/(2*(S1S2+S1S3+S2S3))
S3<- (S2S3+S1S3)/(2*(S1S2+S1S3+S2S3))

S1vector[gens] <- (S1S2+S1S3)/(2*(S1S2+S1S3+S2S3))
S2vector[gens] <- (S1S2+S2S3)/(2*(S1S2+S1S3+S2S3))
S3vector[gens] <- (S1S3+S2S3)/(2*(S1S2+S1S3+S2S3))

}## end generations

S1S2
S1S3
S2S3
S1vector
S2vector
S3vector

minVect<-{}
minVect[1]<-min(S1vector)
minVect[2]<-min(S2vector)
minVect[3]<-min(S3vector)

maxVect<-{}
maxVect[1]<-max(S1vector)
maxVect[2]<-max(S3vector)
maxVect[3]<-max(S3vector)

minTotal<-min(minVect)
maxTotal<-max(maxVect)

minTotal
maxTotal

plot(S1vector,type = "l", lwd=5, xlim=c(1, gens), ylim=c(minTotal-.01,maxTotal+.01), xaxt="n")
#axis(1, at = seq(1, 10, by = 1), las=2)
lines(S2vector,type = "l", lty=3, lwd=5)
lines(S3vector,type = "l", lty=4, lwd=5)

plot(S1vector,type = "l", lwd=5, xlim=c(1, gens), ylim=c(minTotal-.01,maxTotal+.01), xaxt="n")
axis(1, at = seq(1, 10, by = 1), las=2)
plot(S2vector,type = "l", lwd=5, xlim=c(1, gens), ylim=c(minTotal-.01,maxTotal+.01), xaxt="n")
axis(1, at = seq(1, 10, by = 1), las=2)
```

```r
plot(S3vector,type = "l", lwd=5, xlim=c(1, gens), ylim=c(minTotal-.01,maxTotal+.01), xaxt="n")
axis(1, at = seq(1, 10, by = 1), las=2)

#plot(S2vector,type = "l", lty=3, lwd=5, xlim=c(1, gens), ylim=c(0,1))
#plot(dtvectorN1freq,type = "l", lwd=5, xlim=c(0, gen), ylim=c(0,1))

#plot(dtvectorN1,type = "n", xlim=c(1, gen), ylim=c(0,Ka))
#lo <- loess(dtvectorN1~time)
#xl <- seq(min(time),max(time), (max(time) - min(time))/1000)
#lines(xl, predict(lo,xl), col='black', lwd=5)

##plot(dtvectorN2,type = "n", xlim=c(1, gen), ylim=c(0,Kb))
#lo <- loess(dtvectorN2~time)
#xl <- seq(min(time),max(time), (max(time) - min(time))/1000)
#lines(xl, predict(lo,xl), col='black', lwd=5)

#plot(dtvectorN1freq,type = "n", xlim=c(0, gen), ylim=c(0,1))
#lo <- loess(dtvectorN1freq~time)
#xl <- seq(min(time),max(time), (max(time) - min(time))/1000)
#lines(xl, predict(lo,xl), col='black', lwd=5)

#plot(dtvectorN2freq,type = "n", xlim=c(0, gen), ylim=c(0,1))
#lo <- loess(dtvectorN2freq~time)
#xl <- seq(min(time),max(time), (max(time) - min(time))/1000)
#lines(xl, predict(lo,xl), col='black', lwd=5)
```

Figure 3 data generated in R

Make a model with 2 niches (A and B) and two variants (1 and 2)
show how it is not frequency but abundance and carrying capacity that affect fitness

cycle through differential fitness values (home vs away) and migration values
show when fitness changes and when polymorphism maintained

PARAMETERS
N1aStart<- 7 ## Starting pop of variant 1 (all start in their home niche)
N2bStart<- 100 ## Starting pop of variant 2 (all start in their home niche)
N1bStart <- 5 ## Starting pop of variant 1 in niche b
N2aStart <- 2 ## Starting pop of variant 2 in niche a
N1m <-0 ## migrant pool
N2m <-0 ## migrant pool
Ka<- 10 ## Carrying capacity of niche a
Kb<- 10000 ## Carrying capacity of niche b
rh <- 1 ### growth rate in correct niche
##ra == growth rate in incorrect niche (cycle through this in for loop)

HOUSEKEEPING STUFF
i <- seq(0, .1, by=.05)
j <- seq(0, .1, by=.1)
k <- seq(0, 20, by=1)

dtvectorN1a<-{}
dtvectorN1am<-{}
dtvectorN1b<-{}
dtvectorN2a<-{}
dtvectorN2b<-{}
dtvectorN2am<-{}
dtvectorN1aT<-{}

START OF MODEL SIMULATION

for (ra in i) { ##Selection differential loop

for (m in j) { ## migration loop

N1a<- N1aStart
N2b<- N2bStart
N1b<- N1bStart
N2a<- N2aStart

for (gen in k){## generations loop

#ra=0
#m=0

```
#N1a<- N1aStart; N2b<- N2bStart; N1b<- N1bStart; N2a<- N2aStart

#N1a <- N1a * (1+rh* (1 - (N1a+N2a)/Ka))
N1a <- N1a * (1+rh* (1 - N1a/Ka))
N1b <- N1b * (1+ra* (1 - (N1b+N2b)/Kb))

N2a <- N2a * (1+ra* (1 - (N1a+N2a)/Ka))
N2b <- N2b * (1+rh* (1 - (N1b+N2b)/Kb))

N1aT<- N1a - N1a*m
N1bT<- N1b - N1b*m
N2aT<- N2a - N2a*m
N2bT<- N2b - N2b*m

if (N1aT < 0) {N1aT<-0}
if (N2aT < 0) {N2aT<-0}
if (N1bT < 0) {N1bT<-0}
if (N2bT < 0) {N2bT<-0}

N1am <- m * (N1a + N1b)/2
N1bm <- N1am
N2am <- m * (N2a + N2b)/2
N2bm<-N2am

NaSpace<- Ka-N1aT-N2aT ## migrants cannot displace the residents
NbSpace<- Kb-N1bT-N2bT## migrants cannot displace the residents
if(NaSpace<0){NaSpace<-0}
if(NbSpace<0){NbSpace<-0}

if ((N1am+N2am) > NaSpace){
N1a <- N1aT + NaSpace*N1am/(N1am+N2am)
N2a <- N2aT + NaSpace*N2am/(N1am+N2am)
} else {
N1a <- N1aT + N1am
N2a <- N2aT + N2am
}

if ((N1bm+N2bm) > NbSpace){
N1b <- N1bT + NbSpace*N1bm/(N1bm+N2bm)
N2b <- N2bT + NbSpace*N2bm/(N1bm+N2bm)
} else {
N1b <- N1bT + N1bm
N2b <- N2bT + N2bm
}
```

```
N1<-N1a+N1b
N2<-N2a+N2b

dtvectorN1a[gen]<-N1a
dtvectorN1aT[gen]<-N1aT
dtvectorN1b[gen]<-N1b
dtvectorN1am[gen]<-N1am
dtvectorN2a[gen]<-N2a
dtvectorN2b[gen]<-N2b
dtvectorN2am[gen]<-N2am

} ## end generations loop

} ##end migration loop

}## end selection loop

dtvectorN1a
dtvectorN1b
dtvectorN1am
dtvectorN2a
dtvectorN2am
dtvectorN2b
dtvectorN1aT
```