Title page

Myeloablation-associated deletion of ORF4 in a human coronavirus 229E infection

Alexander L. Greninger[1,2,#], Gregory Pepper[1], Ryan C. Shean[1], Anne Cent[1], Isabel Palileo[1],

Jane M. Kuypers[1], Joshua T. Schiffer[2,3], Keith R. Jerome[1,2]

[1]Department of Laboratory Medicine, University of Washington, Seattle, WA

[2]Vaccine and Infectious Disease Division, Fred Hutchinson Cancer Research Center, Seattle,

WA

[3]Department of Medicine, University of Washington, Seattle, WA

[#]corresponding author, agrening@uw.edu

Abstract

We describe metagenomic next-generation sequencing (mNGS) of a human coronavirus

229E from a patient with AML and persistent upper respiratory symptoms, who underwent

hematopoietic cell transplantation (HCT). mNGS revealed a 548-nucleotide deletion, which

comprised the near entirety of the ORF4 gene, and no minor allele variants were detected to

suggest a mixed infection. As part of her pre-HCT conditioning regimen, the patient

received myeloablative treatment with cyclophosphamide and 12 Gy total body irradiation.

Iterative sequencing and RT-PCR confirmation of 4 respiratory samples over the 4-week

peritransplant period revealed that the pre-conditioning strain contained an intact ORF4

gene, while the deletion strain appeared just after conditioning and persisted over a 2.5-

week period. This sequence represents one of the largest genomic deletions detected in a

human RNA virus and describes large-scale viral mutation associated with myeloablation

for HCT.

Keywords: coronavirus, 229E, human coronavirus 229E, myeloablation, genome evolution,

metagenomics, mNGS, irradiation, deletion, ORF4, ORF4a, ORF4b

Introduction

Coronaviruses are positive-stranded RNA viruses with large genomes, ranging in size from 25-35kb. Although much attention has been paid to the epidemic potential and high mortality rates for MERS and SARS coronaviruses, the most common human coronaviruses (HCoV) include 229E, OC43, HKU1, and NL63 viruses [1]. HCoV 229E and NL63 are alphacoronaviruses with genome lengths of 27-28kb, and HCoV HKU1 and OC43 are betacoronaviruses with genome lengths around 30-31kb [2]. Few studies have been conducted on the genome evolution of the common HCoV species [3,4].

HCoVs enact a major burden on hematopoietic cell transplant (HCT) patients with >10% cumulative incidence of HCoV infection among HCT by day 100 of transplant [5]. HCoV in HCT patients is associated with prolonged shedding, with a median duration of 3 weeks [5]. Conditioning regimens for HCT usually consist of chemotherapy and irradiation that may induce genomic alterations [6,7]. Treatment-related myeloid leukemias are well-known to be associated with radiation therapy or chemotherapeutic agents such as cyclophosphamide [8]. Here we describe the generation of a large deletion in an RNA virus that was temporally associated with the myeloablative conditioning regimen for HCT.

Materials and Methods

Cell culture and qRT-PCR

The HCoV 229E type strain ATCC VR-740 was obtained and used as a positive control for cell culture. 200ul of each patient nasal swab in viral transport media (50ul 229E control/150ul 2% MEM) was inoculated onto MRC-5 cells and incubated at 37C in stationary rack for 30 minutes. After incubation, 1.2mL of 2% MEM was added to each tube and incubated at 37C. Each day tubes were read for cytopathic effect and 50uL of supernatant was stored at -80C. 50uL culture supernatants were extracted on a Roche

MagNA Pure LC 2.0 and RNA was eluted in 100uL water. 10uL of RNA was used for qRT-

PCR with primers targeting the HCoV polymerase gene [9]. qRT-PCR on patient samples was

performed on 200uL of nasal swab in viral transport with the same extraction and PCR

conditions.

Metagenomic next-generation sequencing and confirmatory RT-PCR with deep sequencing

mNGS was performed as described previously [10]. Briefly, 20uL of extracted RNA

was treated with DNAse I and double stranded cDNA was created using random hexamers

with SuperScript III and Sequenase v2.0 (Thermo Fisher). mNGS libraries were created

using one-third volume NexteraXT reactions with 16 cycles of PCR amplification. mNGS

libraries from the deletion-containing specimens were prepared twice from RNA. Dual-

indexed libraries were sequenced on an Illumina MiSeq over several different types of runs

(1x190bp, 1x188bp, 2x94bp, 2x300bp) with consistent results. Sequencing data were

Nextera adapter and quality (Q30) trimmed using cutadapt, iteratively mapped to the HCoV

229E reference genome (NC_002645) using the Geneious v9.1 read aligner, de novo

assembled using SPAdes v3.9, and visualized using Geneious v9.1 and manually curated [11–

13]. The HCoV 229E reference genome and the strains sequenced in this study have an

average pairwise nucleotide identity of 98%. Majority consensus genomes from mNGS

were called using a minimum of 3X coverage.   Reads for each specimen were also mapped

to the day -12 consensus genome to look for minor variants that arose during treatment.

Due to low coverage, minor variants were called with a minimum allele frequency of 10% at

11X coverage, after trimming of inversions due to Nextera artifact, variants within 10

nucleotides of end of read, and PCR duplicates.

Confirmatory gap junction RT-PCR was performed on 1uL of RNA extracted as

above using 35 cycles following manufacturer's recommended conditions (Tm 55C) using

quarter-volumes of the Qiagen One-Step RT-PCR kit and primers HCoV_24034F (5′ –

TTGTTGTGAATCAACTAAACTTCC – 3′) and HCoV_24808R (5′ –

ACACACCAGAGTAGTACATTAAC – 3′). Deep sequencing of the day -9 and day -2 PCR

amplicons was performed using 1ng of PCR product purified with the Zymo DNA Clean and

Concentrator Kit-5 with 1/10th volumes of the the Kapa HyperPlus kit ligation of Y-shaped

adapters followed by 7 cycles of Truseq-adapter dual-indexed PCR amplification [14].

Sequencing was performed using a 315x310bp run on an Illumina MiSeq. From the day -9

amplicon 36,880 paired-end reads were recovered, while from the day -2 amplicon 231,990

paired-end reads were recovered. Truseq-adapter and quality-trimmed using bbduk and

aligned to the ORF4 intact and deleted alleles present in the day -12 and day 15 consensus

genomes using the Geneious read aligner with no gaps or mismatches allowed [15]. The

methods were performed in accordance with relevant guidelines and regulations and

approved by the University of Washington Institutional Review Board.

Results

The virus analyzed was from a female in her 40s with acute myelogeneous leukemia

presenting for hematopoetic cell transplant. The previous year she had undergone 4 cycles

of chemotherapy (G-CLAM, G-CLA, cytarabine, and decitabine-primed MEC). Her prior

month's bone marrow evaluation showed 16% blasts by flow cytometry with normal

cytogenetics. On day –12 prior to transplant, she was found to be infected by a coronavirus

229E with a cycle threshold (Ct) of 30 and persisted at the same Ct on day -9 (Figure 1A).

Based on the patient's risk of relapsing leukemia and relative rarity of HCoV pneumonitis,

the decision was made to continue with HCT despite ongoing viral shedding.   On days -7

and -6 she received cyclophosphamide 60mg/kg, and on days -4, -3, and -2 she received a

cumulative dose of twice-daily 2 Gy total body irradiation for a cumulative dose of 12 Gy

(Figure 1A). Other significant drugs the patient received during this time period included

acyclovir, granisetron, micafungin, voriconazole, and sulfamethoxazole-trimethoprim. A

mismatched unrelated donor peripheral blood stem-cell transplant was performed on day

0. Throughout the HCoV infection, the patient had mild nasal congestion but never required

supplemental oxygen. Her course was complicated by bacteremia on day 10, thought to be

due to grade 2 mucositis, and she was discharged on day 20.

Four nasal swab samples over a four-week period were available from the patient.

All qRT-PCR cycle thresholds (Ct) recovered over the four-week period were comparable,

ranging between 31.3-33.5 (Table). The percent of total reads mapping to HCoV 229E

ranged from 0.01-0.75%, consistent with the low concentrations of virus found by qRT-PCR

(Table). mNGS of the Day 15 post-transplant specimen revealed a large deletion that

encompassed 548 of the 660 nucleotides of the ORF 4 gene (Figure 1B) with average

coverage of 68X across the viral genome. Repeat library preparations of the same specimen

demonstrated the same deletion. The deletion was found whether mapping to the HCoV

229E reference genome or de novo assembly. No read was recovered within the area of the

deletion, suggesting the presence of a viral population entirely containing the genomic

deletion. mNGS of the day -12 and -9 specimens from the same patient revealed a

completely intact ORF4a/4b (Figure 1B). Other than the large deletion, no consensus

variants were observed between the HCoV 229E consensus genomes recovered from day -

12, day -9, and day 15 specimens. Mapping of reads from the day –12, day -9, day 15

samples to the day -12 consensus genome revealed two minor allele variants that reached

>10% allele frequency and had consistent allele frequencies between the day -12 and day -9

samples. The first was a c.12332A>G, p.K4111R change in the replicase protein that went

from undetectable in the day -12 and -9 specimens and was at 11.9% allele frequency in the

day 15 specimen. The second was a c.675_676insU that maintained at 10.6%, 10.8%, and

8.9% allele frequency in the day -12, day -9, and day 15 specimens.

No junctional reads across the 548-nucleotide deletion were found in either the

day -12 or day -9 specimen by mNGS. mNGS of the day -2 specimen also demonstrated the

548-nucleotide deletion with coverage across the genome consistent with RNA degradation

(Figure S1). No consensus variants were observed in the genomic regions covered in the

day -2 specimen. Confirmatory RT-PCR across the ORF 4a/4b gene demonstrated an intact

gene in the pre-myeloablation specimens and a smaller PCR product consistent with the

548-nucleotide deletion in the day -2 and day 15 post-myeloablation specimens (Figure 1C).

Deep sequencing of the day -9 and day -2 PCR amplicons revealed no junction reads from

the day -9 specimen mapping to the day 15 deletion strain locus with 11,279X coverage at

the same locus for the intact ORF4 gene in the day -12 strain (Figure S2). Conversely, only

one paired-end read from the day -2 specimen mapped to the deleted portion of the ORF4

gene locus in the day -12 strain with 158,938X coverage at the ORF4 locus in the day 15

deletion strain (Figure S2).

Attempts to culture each of the patient's specimens on MRC-5 cells proved

unsuccessful. No CPE was visualized for any of the patient's four clinical samples and HCoV

229E RNA levels in the culture supernatant were undetectable by qRT-PCR after

inoculation, while viral growth was apparent in the positive control HCoV 229E ATCC type

strain (Figure S3).

Discussion

We describe metagenomic detection of a large deletion in a HCoV 229E strain

temporally associated with a myeloablative regimen of cyclophosphamide and total body

irradiation in an HCT patient. The ORF4-deleted virus maintained the same viral load over

a 2.5 week period and was clonal by day 15 after transplant, suggesting that it was

replication competent in vivo. No other majority consensus variants in the HCoV genome

arose during the sampled period. One minor variant arose in the day 15 specimen that was

not present in the earlier specimens, while another minor variant maintained a similar

allele frequency throughout the sampled period. This rate of single nucleotide variant

evolution is consistent with the molecular clock of HCoV and with a study of that found no

HCoV genomic variants arose during the first month of infection [16]. The ability of the virus

to persist in vivo despite the near complete loss of ORF4 suggests that this accessory gene is

completely dispensable for HCoV 229E replication and may outcompete the wild-type virus

under certain conditions. Indeed, HCoV 229E type strain contains a 2-nucleotide deletion

that splits ORF4 into ORF4a/4b, and the alpaca alphacoronavirus contains a 1-nucleotide

insertion that also disrupts the reading frame [17,18]. Meanwhile, all other group1b

coronaviruses including all known HCoV 229E clinical strains contain an intact ORF4 [17,19].

This is the first detection of a non-intact ORF4 in HCoV 229E associated with human

infection.

The function of ORF4 in the alphacoronaviruses is unknown. The protein sequence

shows no significant alignment to any protein in Protein Data Bank by HHPred [20]. ORF4a of

the HCoV 229E type strain has been suggested to be a viroporin that possessed ion channel

activity [21]. Interestingly, replacement of ORF4 in the HCoV 229E type strain with Renilla

luciferase resulted in viral replication in vitro comparable to that of the wild-type strain,

consistent with our in vivo data that ORF4 is unlikely to be required for replication [22].

Taken together, these data suggest that ORF4 would make a poor antiviral target for the

human coronaviruses.

Many potential mechanisms for development of the deletion exist. The

chemotherapy and/or irradiation could have been directly toxic to the coronavirus genome.

Cyclophosphamide is a well-known clastogen and has recently been associated with

increased development of single nucleotide variations insertions and deletions [7,23].

Alternatively, the myeloablative regime could have removed immune pressure, allowing for

the loss of ORF4. It seems that in certain circumstances, deletion of ORF4 could prove a

selective advantage as it was the only allele present in the day 15 specimen.

This study has several limitations. We were unable to culture any of the patient's

HCoV specimens to demonstrate ability to replicate in vitro. This is not necessarily

surprising as HCoV 229E are not usually cultured in the clinical virology lab and the virus

was present at low concentrations in the sample. We are also limited in accurately calling

minor variants by the relatively low coverage of each viral genome. We did not attempt

reverse genetics of any alphacoronaviruses to definitively show ORF4 is dispensable for in

vitro replication, although the ability of the type strain to replicate despite a 2-nucleotide

deletion in ORF4 and the luciferase gene replacement experiment described above are

consistent with this possibility.

In summary, we describe a large deletion in HCoV 229E encompassing almost an

entire gene that was temporally associated with myeloblation. Little is known about the

genome evolution of HCoV 229E as only 20 genomes are currently available in NCBI

Genbank. Similarly sized (500-700 nucleotide) deletions have previously been found in

hepatitis D virus in patients receiving antiviral therapy [24]. Our study is also the first

demonstration of microbial evolution associated with cyclophosphamide treatment or

irradiation. Further research into how chemotherapy and irradiation affect the microbiome

is warranted. As mNGS is increasingly performed in the clinical lab [25], much intriguing

biology can be unlocked from clinical specimens, providing a continuing case for the

sequencing clinical microbial genomes [26-28].

Data Availability Statement

These sequences are deposited in NCBI Genbank under the strain names SC677 day 15

(KY369909), SC379 day -12 (KY621348), and SC399 day -9 (KY674914). The non-human

reads are deposited in the NCBI Short Read Archive under accessions SRR5809357-

SRR5809372, SRR5809395, SRR5816371-SRR5816372.

Author Contributions

ALG, GP, IP performed experiments; ALG, RCS, AC, JMK analyzed data; ALG, JTS, KRJ wrote

the paper.

Competing Interests

The authors declare they have no competing interests.

Supplementary information is available at the *Genomic Medicine* website.

References

223
224
1.   Lepiller, Q. *et al.* High incidence but low burden of coronaviruses and preferential

associations between respiratory viruses. *J. Clin. Microbiol.* **51,** 3039–3046 (2013).

2.   Forni, D., Cagliani, R., Clerici, M. & Sironi, M. Molecular Evolution of Human Coronavirus

Genomes. *Trends Microbiol.* **25,** 35–48 (2017).

3.   Dijkman, R. & Hoek, L. van der. Human Coronaviruses 229E and NL63: Close Yet Still So

Far. *J. Formos. Med. Assoc.* **108,** 270–279 (2009).

4.   Posthuma, C. C., Te Welthuis, A. J. W. & Snijder, E. J. Nidovirus RNA polymerases:

complex enzymes handling exceptional RNA genomes. *Virus Res.* (2017).

doi:10.1016/j.virusres.2017.01.023

5.   Milano, F. *et al.* Human rhinovirus and coronavirus detection among allogeneic

hematopoietic stem cell transplantation recipients. *Blood* **115,** 2088–2094 (2010).

6.   Gyurkocza, B. & Sandmaier, B. M. Conditioning regimens for hematopoietic cell

transplantation: one size does not fit all. *Blood* **124,** 344–353 (2014).

7.   Szikriszt, B. *et al.* A comprehensive survey of the mutagenic impact of common cancer

cytotoxics. *Genome Biol.* **17,** 99 (2016).

8.   Godley, L. A. & Larson, R. A. Therapy-related Myeloid Leukemia. *Semin. Oncol.* **35,** 418–

429 (2008).

9.   Kuypers, J. *et al.* Clinical disease in children associated with newly described

coronavirus subtypes. *Pediatrics* **119,** e70-76 (2007).

10.   Greninger, A. L. *et al.* Rapid Metagenomic Next-Generation Sequencing during an

Investigation of Hospital-Acquired Human Parainfluenza Virus 3 Infections. *J. Clin.*

Microbiol. **55,** 177–182 (2017).

11.   Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing

reads. *EMBnet.journal* **17,** 10–12 (2011).

12. Kearse, M. *et al.* Geneious Basic: an integrated and extendable desktop software

platform for the organization and analysis of sequence data. *Bioinforma. Oxf. Engl.* **28,**

1647–1649 (2012).

13. Bankevich, A. *et al.* SPAdes: A New Genome Assembly Algorithm and Its Applications to

Single-Cell Sequencing. *J. Comput. Biol.* **19,** 455–477 (2012).

14. Koelle, D. M. *et al.* Worldwide circulation of HSV-2 × HSV-1 recombinant strains. *Sci.*

Rep. **7,** 44084 (2017).

15. BBTools. *DOE Joint Genome Institute* Available at: http://jgi.doe.gov/data-and-

tools/bbtools/. (Accessed: 11th July 2017)

16. Ogimi, C. *et al.* Prolonged shedding of human coronavirus in hematopoietic cell

transplant recipients: risk factors and viral genome evolution. *J. Infect. Dis.*

doi:10.1093/infdis/jix264

17. Dijkman, R. *et al.* Human coronavirus 229E encodes a single ORF4 protein between the

spike and the envelope genes. *Virol. J.* **3,** 106 (2006).

18. Crossley, B. M., Mock, R. E., Callison, S. A. & Hietala, S. K. Identification and

Characterization of a Novel Alpaca Respiratory Coronavirus Most Closely Related to the

Human Coronavirus 229E. *Viruses* **4,** 3689–3700 (2012).

19. Farsani, S. M. J. *et al.* The first complete genome sequences of clinical isolates of human

coronavirus 229E. *Virus Genes* **45,** 433–439 (2012).

20. Söding, J., Biegert, A. & Lupas, A. N. The HHpred interactive server for protein homology

detection and structure prediction. *Nucleic Acids Res.* **33,** W244-248 (2005).

21. Zhang, R. *et al.* The ORF4a protein of human coronavirus 229E functions as a viroporin

that regulates viral production. *Biochim. Biophys. Acta BBA - Biomembr.* **1838,** 1088–

1095 (2014).

22. Worm, S. H. E. van den *et al.* Reverse Genetics of SARS-Related Coronavirus Using

Vaccinia Virus-Based Recombination. *PLOS ONE* **7**, e32857 (2012).

23. Povirk, L. F. & Shuker, D. E. DNA damage and mutagenesis induced by nitrogen

mustards. *Mutat. Res.* **318**, 205–226 (1994).

24. Hsu, C.-W. *et al.* Detection of hepatitis D virus RNA carrying large fragment deletions in

patients with severe hepatitis B/D receiving oral antiviral therapy. *J. Med. Virol.* **87**,

634–641 (2015).

25. Greninger, A. L. *et al.* A Metagenomic Analysis of Pandemic Influenza A (2009 H1N1)

Infection in Patients from North America. *PLoS ONE* **5**, e13381 (2010).

26. Greninger, A. L. *et al.* Copy number heterogeneity of JC virus standards. *J. Clin. Microbiol.*

(2016). doi:10.1128/JCM.02337-16

27. Greninger, A. L. *et al.* Clinical metagenomic identification of Balamuthia mandrillaris

encephalitis and assembly of the draft genome: the continuing case for reference

genome sequencing. *Genome Med.* **7**, 113 (2015).

28. Canuti, M. & van der Hoek, L. Virus discovery: are we scientists or genome collectors?

Trends Microbiol. **22**, 229–231 (2014).

Figure Legend

Figure – Detection of a 548-nucleotide deletion in a HCoV 229E strain associated with

myeloablation. A) Depiction of the case history surrounding myeloablation and

coronavirus infection. A woman in her 40s with AML was found to be infected with a HCoV

229E just before her HCT. Due to the patient's advanced disease, the decision was made to

continue with the transplant. Four successive HCoV-positive nasal swab specimens were

available from the patient; two from before treatment with myeloablative

cyclophosphamide and total body irradiation and two from after. All days are reported

relative to HCT. B) Coverage plots of mNGS reads mapped to the reference genome for

HCoV 229E (NC_002645) of three samples from the patient revealed the generation of a

548-nucleotide deletion in the ORF4a/4b gene. In addition to deleting 83% of the coding

region from ORF4a/4b, the deletion resulted in a frame shift that resulted in a premature

stop codon 10 amino acids before the reference genome stop codon. Primer binding sites

are denoted in blue for confirmatory junction RT-PCR. C) Confirmatory junction RT-PCR of

the four available HCoV-positive nasal swabs revealed the absence of the deletion before

myeloablation and the presence of the deletion in specimens taken after myeloablation.

NTC, no template control.

A

HCoV 229E intact strain
HCoV 229E deletion strain
Outpatient oncology
Inpatient oncology
2 Gy total body irradiation
Cyclophosphamide

Days relative to transplant

-16 -12 -8 -4 0 4 8 12 16 20 24

C

L   Day -12   Day -9   D

bp

B

Day -12   27X average

Day -9   16X average

Day 15   68X average

replicase polyprotein 1a
replicase polyprotein 1ab   replicase polyprotein 1ab   spike protein

MALGLFTLQFVS--DVYTCFVNTQLLE*

nucleotide 23,910   24,127   124X

24,676

spike protein   ORF 4a   ORF 4b

Table 1 – Specimens sequenced in this study. mNGS sequencing reads to HCoV 229E and total reads for each specimen are indicated, along with qRT-PCR cycle threshold.

Day	HCoV Ct	HCoV reads	Total reads	Strain name
-12	31.5	7,257	4,395,609	SC379
-9	33.5	3,733	30,166,065	SC399
-2	32.7	5,064	11,185,156	SC475
15	31.3	11,990	1,594,894	SC677