The Past Sure Is Tense: On Interpreting Phylogenetic Divergence Time Estimates

Joseph W. Brown[1,*], and Stephen A. Smith[1]

[1]*Department of Ecology & Evolutionary Biology, University of Michigan, Ann Arbor, MI 48109, USA*
[*]*Correspondence to be sent to: Department of Ecology & Evolutionary Biology, University of Michigan, Ann Arbor, MI 48109, USA;*
E-mail: josephwb@umich.edu

Abstract.—Divergence time estimation — the calibration of a phylogeny to geological time — is a integral first step in modelling the tempo of biological evolution (traits and lineages). However, despite increasingly sophisticated methods to infer divergence times from molecular genetic sequences, the estimated age of many nodes across the tree of life contrast significantly and consistently with timeframes conveyed by the fossil record. This is perhaps best exemplified by crown angiosperms, where molecular clock (Triassic) estimates predate the oldest (Early Cretaceous) undisputed angiosperm fossils by tens of millions of years or more. While the incompleteness of the fossil record is a common concern, issues of data limitation and model inadequacy are viable (if underexplored) alternative explanations. In this vein, Beaulieu *et al.* (2015) convincingly demonstrated how methods of divergence time inference can be misled by both (i) extreme state-dependent molecular substitution rate heterogeneity and (ii) biased sampling of representative major lineages. While these (essentially model-violation) results are robust (and probably common in empirical data sets), we note a further alternative: that the configuration of the statistical inference problem alone precluded the reconstruction of the paleontological timeframe for the crown age of angiosperms. We demonstrate, through sampling from the joint prior (formed by combining the tree (diversification) prior with the various calibration densities specified for fossil-calibrated nodes), that with no data present at all, an Early Cretaceous crown angiosperms is rejected (i.e., has essentially zero probability). More worrisome, however, is that for the 24 nodes calibrated by fossils, almost all have indistinguishable marginal prior and posterior age distributions, indicating an absence of relevant information in the data. Given that these calibrated nodes are strategically placed in disparate regions of the tree, they essentially anchor the tree scaffold, and so the posterior inference for the tree as a whole is largely determined by the pseudo-data present in the (often arbitrary) calibration densities. We recommend, as for any Bayesian analysis, that marginal prior and posterior distributions be carefully compared, especially for parameters of direct interest. Finally, we note that the results presented here do not refute the biological modelling concerns identified by Beaulieu *et al.* (2015). Both sets of issues remain apposite to the goals of accurate divergence time estimation, and only by considering them in tandem can we move forward more confidently. [Angiosperms; divergence time estimation; fossil record; marginal priors; information content; diptych; wild speculation.]

"Molecular clocks are not up to the job, but neither is the fossil record." Donoghue and Benton (2007)

Divergence time estimation from molecular genetic sequences is fraught with uncertainty. The errors involved in routine phylogenetic reconstruction (suboptimal alignments, inadequate substitution models, insufficient taxon/gene sampling, real gene tree/species tree conflict, etc.) are compounded by assumptions required to transform a phylogram (in units of expected number of substitutions per site) into a chronogram (in units of geological time): 1) an appropriate model of substitution rate heterogeneity among lineages and across time, and 2) temporal calibrations, generated from the fossil (or biogeographic) record, used to inform and constrain the extent of rate variation.

It is therefore not surprising that there are discrepancies between inferred molecular genetic and paleontological timescales. However, while many disagreements are minor and may innocuously be attributed to insufficient sampling (genes, taxa, or fossils), others are so severe and consistent that they cast serious doubt on the appropriateness of molecular clock models, the fossil record, or both. One prominent example concerns placental mammals, where molecular estimates (e.g., Meredith *et al.* 2011) of the crown age almost double those from the fossil record (e.g., O'Leary *et al.* 2013), obscuring the role of the K-Pg mass extinction on the evolutionary trajectory of this group. Another conspicuous example, also spanning the K-Pg boundary, is crown birds (Neornithes; Ksepka *et al.* 2014), where (re)analyses have repeatedly lead to incongruous inferred evolutionary timeframes (e.g., Ericson *et al.* 2006 vs. Brown *et al.* 2007; Jarvis *et al.* 2014 vs. Mitchell *et al.* 2015; Prum *et al.* 2015).

However, perhaps the best exemplified recalcitrant node in terms of absolute age is that of crown angiosperms, where molecular clocks pervasively infer a Triassic age (e.g., Bell *et al.* 2010; Smith *et al.* 2010; Zeng *et al.* 2014; Beaulieu *et al.* 2015; Foster *et al.* 2016; see a comprehensive review of estimates in Magallón *et al.* 2015) while the oldest undisputed fossil remains are restricted to the Early Cretaceous (136 Ma; Brenner 1996). Moreover, using a model of uniform random fossilization (Marshall 2008) applied to 137 fossils lead Magallón *et al.* (2015) to infer an upper bound on the origin of crown angiosperms of just ~140 Ma. The age of this one node, more than any other, has seriously called into question the utility of both molecular clock models and the fossil record.

Numerous reasons have been put forth to explain the disparity of molecular and paleontological timescales. On the one hand we have concerns with the fossil record. By their nature, fossils *must* postdate the origin of taxa, meaning that molecular estimates *should* predate those from the fossil record. Furthermore, it is clear that the fossil record is imperfect (more so for some groups than others), such that in some instances it may prove impossible to ever have a tight correlation with

molecular clock estimates. However, 'absence of evidence' vs. 'evidence of absence' is a complex matter, so a more productive avenue to pursue may be that of considering molecular approaches. While it has been speculated that molecular clocks might 'run fast' during radiations (thereby misleading clocks into inferring a long period of time has occurred; Benton 1999), this has no empirical support. However, it is known that substitution model mis-specification can mislead divergence time estimation (Phillips 2009; Schenk and Hufford 2010) and ultimately downstream analyses (Revell *et al.*, 2005). Likewise, mis-specification of relaxed clock models may also lead to inaccurate results (Dornburg *et al.* 2012; Worobey *et al.* 2014). Being only semi-identifiable, molecular clock methods require calibration from the fossil record, so appropriate calibration use is critically important (Inoue *et al.* 2010; Sauquet *et al.* 2011; Warnock *et al.* 2012; Magallón *et al.* 2013; Zhu *et al.* 2015). Finally, Beaulieu *et al.* (2015) recently demonstrated through simulation two further ways where molecular clocks might be mislead: (i) through extreme state-dependent molecular substitution rate heterogeneity, and (ii) biased sampling of representative major lineages (both of which are essentially instances of model violation). While these are all valid concerns that should always be kept in mind, we explore below a further non-biological possibility that may unknowingly be at play in many data analyses.

Diptych: A Metaphor For Data Analysis?

A diptych is a device commonly used in western art and literature. It consists of paired, complementary works, in the artistic tradition typically two images joined at a hinge (e.g., Fig. 1). The function of a diptych is to reciprocally illuminate the component images, ideally revealing some more holistic concept. It is this feature that suggests an association with Bayesian data analysis. It is *de rigueur* in any Bayesian analysis to carefully compare paired prior and posterior parameter distributions to gauge how information content (via the likelihood) drives the results, as well as to establish the sensitivity of inferences to prior specifications. A diptych is, we argue, therefore a useful metaphor for describing the process of changes in belief in parameter values from the prior (*before* data have been observed) to the posterior (*after* data have been observed). We note that the phylogenetic systematics community has been generally lax in this respect, despite available Bayesian software packages making such reflections straightforward. We argue that this is especially important in divergence time estimation analyses, as it is generally unappreciated that there are *three* sets of distributions to consider. In addition to the temporal fossil calibration specified by the investigator (the 'user prior') and the resulting marginal posterior distribution, there exists an intermediate distribution, the marginal prior (also called the 'effective' or 'joint' prior by some authors), which is formed by the interaction among user priors and the underlying 'tree

FIGURE 1. Diptych of Stilicho, ca. 395. Individual images can be viewed on their own, but the work as a whole is only coherent when considering the component pieces together. Photograph from Hayford Pierce and Royal Tyler, "L'art byzantin", Paris, 1932. Public domain.

prior' (for nodes not directly calibrated by a fossil; typically a birth-death prior). Here we turn our attention to these distributions to see what, if anything, we can glean about the age of crown angiosperms.

A Re-Reanalysis Of The Age Of Angiosperms

We reanalyzed the data set provided by Beaulieu *et al.* (2015). The molecular alignment consists of four genes (chloroplast: *atp*B, *psb*B, and *rbc*L; nuclear: 18S) for 124 taxa including 91 angiosperms representing all extant orders (data file provided in the Supplementary Material; see also data from the original paper available on Dryad at http://dx.doi.org/10.5061/dryad.629sc). Sampling was originally designed specifically for dating the origin of angiosperms by allowing the placement of 24 (15 angiosperm) fossil calibrations from across landplants. All dating analyses reported here, like those in the original paper, were performed using the uncorrelated lognormal (UCLN) clock model and birth-death tree prior in BEAST v1.8.2 (Drummond *et al.* 2006; Drummond and Rambaut 2007) using the CPU implementation of the BEAGLE v2.1.2 library (Ayres *et al.* 2012).

We regenerated the posterior results of Beaulieu *et al.* (2015) by employing the original lognormal user priors specified in Table 1 and the following analysis settings: 3 replicate analyses of 50 million generations, sampling every 1000 generations. As in the original Beaulieu *et al.* (2015) paper we fixed the tree topology (their Figure 1). These analyses were re-run without any data (i.e.,

TABLE 1. Lognormal fossil calibration parameters as originally defined in Beaulieu *et al.* 2015.

Name	Mean	St. Dev.	Offset	Clade
Anacostia	1.5	0.5	125.0	Angiosperms
Aquifoliaceae	1.5	0.5	65.0	Angiosperms
Araceae	1.5	0.5	112.0	Angiosperms
Archaefructus	1.5	0.5	125.0	Angiosperms
Arecales	1.5	0.5	85.5	Angiosperms
Cornales	1.5	0.5	85.8	Angiosperms
Endressinia	1.5	0.5	114.0	Angiosperms
Fagales	1.5	0.5	93.6	Angiosperms
Mauldinia	1.5	0.5	99.6	Angiosperms
Myrtales	1.5	0.5	85.8	Angiosperms
Sapindopsis	1.5	0.5	105.8	Angiosperms
Tricolpites	1.5	0.5	125.0	Angiosperms
Typhaceae	1.5	0.5	40.4	Angiosperms
Virginianthus	1.5	0.5	105.8	Angiosperms
Walkerpollis	1.5	0.5	125.0	Angiosperms
Araucarites	1.5	0.5	213.0	Gymnosperms
Cratonia	1.5	0.5	114.0	Gymnosperms
Emporia	2.0	0.5	299.0	Gymnosperms
Paleotaxus	1.5	0.5	197.0	Gymnosperms
Baragwanathia	2.0	0.5	423.0	Outgroups
Palaeosmunda	1.5	0.5	251.0	Outgroups
Pekinopteris	1.5	0.5	228.0	Outgroups
Rellimia	2.0	0.5	388.0	Outgroups
Stachypteris	1.5	0.5	168.0	Outgroups

Notes: 'Offset' denotes the age of the oldest undisputed fossil in Ma. Mean and standard deviation are given in log space. Distributions with a mean of 1.5 have 95% of the prior mass with 10.2 Ma of the fossil age, while those with a mean of 2.0 have 95% of the prior mass with 16.82 Ma of the fossil age.

sampling from the marginal prior). Because these latter analyses were not as computationally demanding, as well as to perform a thorough exploration of parameter space, analyses were run in duplicate for 1 billion generations each, sampling every 10 thousand generations. Finally, we replicated all analyses but replaced the lognormal user prior calibrations from Table 1 with 'extreme exponential' distributions with a mean of 1.0 (offset by fossil ages); such distributions lend the utmost credence to the fossil record, as 95% of the prior masses lie within 3 Ma of the relevant fossil ages. Importantly, all analyses (posterior and prior-only) were initialized with chronograms wherein crown angiosperms originated ∼140 Ma (i.e., consistent with the prescription of Magallón *et al.* 2015).

We focus here on the data (taxon, gene, and fossil sampling) and settings (model and priors) of Beaulieu *et al.* (2015) because it is representative of a standard dating analysis. However, to demonstrate that our own results are not restricted to this particular data set, we also reanalyze the data set of Magallón *et al.* (2015), albeit to a more limited degree because computational requirements. This data set consists of 5 genes (chloroplast: *atp*B, *rbc*L and *mat*K; nuclear: 18S and 26S) for 799 taxa (792 angiosperms) and 121 fossil calibrations (note that the original analysis included 137 fossil calibrations, but 16 non-essential calibrations were subsequently found to be missing from the file shared by the original authors). All analyses employed the UCLN model as above and a fixed tree topology (their Figure 3).

The posterior results of Magallón *et al.* (2015) were regenerated by running 3 replicate analyses of 100 million generations, sampling every 5000 generations. These analyses included the uniform calibration prior (139.5– 136 Ma) on the age of crown angiosperms. To assess the influence of this single prior, 4 replicate analyses of 50 million generations were performed without the prior. Finally, as above, analyses sampling from the marginal prior (i.e., without any data) were performed for both sets of fossil calibrations (each with 4 replicates of 50 million generations).

Prior to summarization, the MCMC log files from each set of replicated analyses were concatenated while removing a conservative 10% sample burnin using the `pxlog` program from the `phyx` package (Brown *et al.* 2017). All results were processed in R v3.3.2 (R Core Team, 2016) and were visualized using `ggplot2` v2.2.1 (Wickham, 2009) and code adapted from `phyloch` v1.5-3 (Heibl, 2008).

THE INACCESSIBILITY OF AN EARLY CRETACEOUS CROWN ANGIOSPERMS

As in the original Beaulieu *et al.* (2015) paper we were unable to recover a posterior age estimate for crown angiosperms that corresponded with the prevailing paleontological timeframe, even when employing overly precise exponential fossil calibration user priors. However, when considering the diptych interpretation by examining the joint marginal prior, it is clear that we need not invoke modelling complications (e.g., due to biased lineage sampling or structured excessive rate heterogeneity) to explain the results. Rather, when running the analysis without any data, we see that an Early Cretaceous crown angiosperms is precluded based on the configuration of the statistical problem alone (Fig. 2). From the trace plots (Fig. 3) we see that the parameter regarding the age of crown angiosperms departs immediately from ∼140 Ma to >200 Ma. In no instance did the MCMC samplers ever return to a 'young' age of angiosperms. The minimum post-burnin value for the prior and posterior analyses for the original lognormal calibration priors were 185.9 Ma and 192.0 Ma, respectively (181.1 Ma and 176.0 Ma for the exponential calibration priors). We note that these findings do not have to do with any peculiarity of the Beaulieu *et al.* (2015) data set as the results generated from the Magallón *et al.* (2015) data set without the uniform prior on crown angiosperms confirms the findings (minimum post-burnin ages for the prior and posterior analyses are 226.1 Ma and 197.2 Ma, respectively). In fact, the Magallón *et al.* (2015) data set, containing far more data (genes, taxa, and fossils) generated the oldest mean posterior estimate (249.7 Ma vs. 233.0 Ma for the original Beaulieu *et al.* (2015) priors vs. 213.3 Ma for the same data set using exponential calibration priors).

While rejecting an Early Cretaceous origin based on the marginal prior alone, both data sets *do* seem to

FIGURE 2. Diptychs comparing marginal prior (orange) and posterior (blue) distributions for the age of crown angiosperms. The top panel displays results using the original lognormal fossil calibration priors with the Beaulieu *et al.* (2015) data set, while the middle panel uses the exponential priors for the same data set. The bottom panel displays results for the Magallón *et al.* (2015) data set. Note that none of the analyses include a user prior for this node. For reference the uniform prior (139.5–136 Ma; pink) used by Magallón *et al.* (2015), reflecting the paleontological estimate, is shown.

FIGURE 3. Traces of the age of the crown angiosperm node across replicate analyses (burnin phase only) for Beaulieu *et al.* (2015) (top) and Magallón *et al.* (2015) (bottom) data sets. Top: prior (lognormal priors), orange ($n = 2$); posterior (lognormal priors), blue ($n = 3$); prior (exponential priors), red ($n = 2$); posterior (exponential priors), green ($n = 3$). Bottom: prior (no angiosperm constraint), orange ($n = 4$); posterior (no angiosperm constraint), blue ($n = 4$); prior (all constraints), green ($n = 4$); posterior (all constraints), red ($n = 3$; note that this is mostly obscured by the green prior trace). All analyses were initialized with the age of crown angiosperms set at ∼140 Ma.

contain signal relevant to the age of crown angiosperms,
as the marginal posterior estimates are shifted
significantly younger than the marginal prior (Fig. 2).
This raises the question: if true, what kind/amount of
data would be required to recover an Early Cretaceous
age for crown angiosperms? Ultimately this comes
down to quantifying phylogenetic 'information content'.
Intriguing possibilities to addressing this sort of question
therefore lie in Shannon information theory (Shannon
1948), a field which the systematics community has
largely ignored. Recently Lewis *et al.* (2016) made
substantial strides forward by applying this theory (in
the discrete case) towards assessing the information
content of prior and posterior tree topology distributions.
Applications to the continuous case, however, are much
more difficult, and no theory (let alone software)
currently exists to address the present problem (Paul
Lewis, personal communication).

PRIORS AND POSTERIORS: A DIPTYCH IN THREE PARTS

Above we introduced 'diptych' as a useful metaphor
for interpreting Bayesian analytical results. The paired
nature of a diptych mirrors the before (prior) and after
(posterior) reflection on what has been learned about
probable parameter values.
The metaphor is slightly more complicated for some
parameters involved in divergence time analyses. Nodes
not explicitly calibrated by fossil data (henceforth,
'uncalibrated' nodes) still require an age prior, which
is provided by a 'tree prior', typically a birth-death,
Yule, or coalescent prior. Such nodes thus have marginal
prior and posterior distributions, and are conducive to
the diptych metaphor. For those nodes that *are* directly
calibrated using fossil information, the interpretation of
the results of inference are more complicated. These
nodes have a 'user prior', a distribution constructed
in some way using information from the fossil record.

FIGURE 4.   Densities for fossil-calibrated nodes from Beaulieu *et al.* 2015 (see Table 1); green indicates the user-specified lognormal prior, orange indicates the marginal prior, and blue indicates the marginal posterior. It is clear, for the majority of calibrated nodes, that (from the nearly complete overlap of prior and posterior distributions) the data contain little information for the parameters of interest.

However, these nodes are also involved in the tree
prior. The resulting 'marginal prior' is a multiplicative
combination of the user and tree priors, and may
also be influenced by interactions with adjacent user-
calibrated nodes (e.g., ancestor and descendant nodes
which have overlapping user priors). The resulting
marginal prior thus does not necessarily reflect the
original user prior. This is, in our judgement, *far
too rarely assessed*. Finally, these nodes have marginal
posterior distributions. However, unlike the uncalibrated
nodes (which involve only two distributions, and thus
a simple interpretation), calibrated nodes involve three
distributions (two priors and one posterior), which
complicates interpretation. [We prefer to continue with

the diptych metaphor for these nodes, rather than the
obvious 'triptych', as the focus lies still on the change
in belief on parameter values before (prior) and after
(posterior) observing the data, even if the prior involves
two components.] The difference between the marginal
prior and marginal posterior, like the uncalibrated
nodes, reflects information in the data (that is, the
likelihood). However, the difference between the user and
marginal priors, if present, may be better described as
demonstrating the influence of 'pseudo-data' present in
the various user priors.

We now turn our attention to the fossil-calibrated
nodes. Ideally we would find that the user and marginal
priors are identical (that is, that the marginal priors

FIGURE 5.   Mean prior (grey) and posterior (black) age estimates using the original lognormal calibration priors for the Beaulieu *et al.* (2015) data set. Blue arrows indicate shifts in age estimates from the prior to the posterior (nodes without arrows have shifted less than 5 Ma from prior to posterior). Nodes that are calibrated by fossil prior distributions are indicated with red circles, except the Tricolpites node which is indicated with a yellow circle. Finally, the (uncalibrated) crown angiosperm node is indicated in green.

reflect the intentions of the researcher), and the marginal priors and posteriors to differ (indicating information present in the data relevant to the parameter of interest). We plot in Fig. 4 the three sets of distributions for the 24 fossil-calibrated nodes for the Beaulieu *et al.* 2015 data set. In general, user and marginal priors match quite closely. However, there is a stark exception involving the Tricolpites constraint: while the user prior specifies that 95% of the prior mass should lie between 135.2–125 Ma, the marginal prior has a 95% highest posterior density (HPD) range of 226.1–128.7 Ma. The marginal posterior

of this node age has a 95% HPD range of 170.7–146.6 Ma, which already surpasses the angiosperm paleontological estimate of ∼140 Ma despite being well nested within the clade. We note that Tricolpites is the oldest constraint within angiosperms used by Beaulieu *et al.* 2015 (see Table 1 and Fig. 5). However, it is clear that it is not this particular calibration which is forcing angiosperms to be 'too old'. Reanalysis without this specific constraint yielded even older posterior estimates for both this node (95% HPD: 181.2–154.3 Ma) and crown angiosperms (mean 241.0 Ma vs. 233.0 Ma with the constraint; data

not shown). We cannot currently identify the cause of the disruption of the Tricolpites user prior.

Finally, we consider differences between the marginal prior and posterior distributions of age estimates for these same fossil-calibrated nodes. As discussed above, shifts in these paired distributions (i.e., following the diptych metaphor) would indicate the presence/degree of relevant phylogenetic 'information' (although theory has not yet been worked out on how to quantify this). However, from Fig. 4 we note that, for the most part, these distributions are nearly indistinguishable. This pattern is even more clear in Fig. 5 where, while uncalibrated nodes show significant shifts between prior and posterior analyses, calibrated nodes show no movement. This lack of updating in belief from prior to posterior for the calibrated nodes suggests two non-mutually exclusive interpretations: (i) that the UCLN model, through allowing independent molecular substitution rates on all edges, may be overfitting the data, essentially allowing all calibrations to be mutually consistent (we do not expect, for example, that all of our calibration densities are equally 'good' - that is, that some constraints should conflict); (ii) that the data set considered here lacks relevant phylogenetic 'information', or at least insufficient information to overrule the pseudo-data present in the fossil calibration densities.

The finding of equivalent prior and posterior distributions may come as a surprise to some users, as unbounded (e.g., lognormal) temporal priors are typically used to 'let the data speak for themselves'. Certainly, divergence time inference is only semi-identifiable (dos Reis and Yang 2013), so we do expect some level of association. It is the *degree* of association that is worrying. We are unaware of any other type of Bayesian analysis in evolutionary biology where identical prior and posterior distributions would not cause concern. The present results go a long way to explaining why divergence time estimation shows such a strong sensitivity to the fossil calibrations used (Inoue *et al.* 2010; Sauquet *et al.* 2011; Warnock *et al.* 2012). The interpretation of fossil calibrations contributing pseudo-data (rather than, say, fossils setting simple minimum age constraints as they have been traditionally) suggests that we might benefit from rethinking lessons that have been learned in the early days of phylogenetic divergence time estimation. We briefly consider one now.

ARE MORE FOSSILS REALLY BETTER IN NODE DATING ANALYSES?

There is an adage in the divergence time estimation literature that as many fossils as possible should be used to calibrate nodes (Benton and Donoghue 2007). This makes sense, as of course we would like to include as much information as possible into a reconstruction. However, this advice largely came about when dating methods (e.g., r8s, Sanderson 2003; multidivtime, Thorne and Kishino 2002) employed constraints (e.g., boolean minimum ages for the age of the fossil) rather

than probabilistic distributions. As long as fossils were correctly placed within the phylogeny, the inclusion of more fossils should not produce misleading results. For instance, fossils that are 'too young' (that is, do not closely approximate in age the node they are calibrating) are either simply uninformative, or appropriately represent limitations of the fossil record. As an extreme example, a chicken bone found in a back alley gutter is a valid (if imprecise) minimum age constraint for *Gallus gallus* (recently estimated at 2.9 Ma; Stein *et al.* 2015).

However, from the results reported above (Figs. 4,5) we find several concerns with including as many fossils as possible in a probabilistic node-dating analysis. [We note that these concerns do not apply to the fossilized birth-death model (Heath *et al.* 2014) or tip-dating (Ronquist *et al.* 2012) approaches to divergence time estimation, which do not involve such calibrations.] First, as with the Tricolpites example above, calibrations can interact with each other and the tree prior in unpredictable ways to produce marginal priors that do not represent the originally intended user priors. While this is a recognized (though under-appreciated) issue, the available solutions work only for a small number of calibrations (Heled and Drummond 2012; Heled and Drummond 2015). Second, given that the marginal prior and posterior calibrated node ages are often indistinguishable (suggesting little relevant phylogenetic information content), it is worrisome that the act of employing temporal calibration priors can directly determine the resulting posterior patterns of rate heterogeneity across a tree. It is not inconceivable, for example, that the parametric use of the best available fossils from an incomplete fossil record could turn a clock-like data set unclock-like, needlessly increasing the model complexity (and therefore uncertainty) involved.

Our final concern with unrestrained parametric calibration use is the form of the calibrations themselves. A flexible assortment of distribution families are available (Ho and Phillips 2009; see also discussion in Brown and van Tuinen 2011), allowing essentially any prior belief to be employed. In addition, researchers can make use of the fossil calibration database (Ksepka *et al.* 2015), and prescribed 'best practices' (Parham *et al.* 2012) can help avoid naive errors when dealing with the fossil record. Nevertheless, the vast majority of user calibration priors employed in the literature are wholly idiosyncratic and arbitrary (we include ourselves here). This is not necessarily a result of molecular phylogeneticists lacking the appropriate paleontological expertise (and isn't that what collaboration is for?), but rather a property of data involved.

While methods exist to generate a distribution from a set of fossils (Marshall 2008; Nowak *et al.* 2013; Claramunt and Cracraft 2015), these require well sampled data. Scant data is an entirely different problem. How does one fit a distribution to a single (exceptionally old, and therefore exceptionally informative) fossil? Minimum bounds are simple (the age of the oldest

fossil), but as Parham *et al.* (2012) note, there exists no standard protocol for formulating maximum ages (let alone the shape of the distribution spanning the upper and lower bounds). Indeed, the process of constructing temporal priors is so nebulous that Lee and Skinner (2011) likened it to "educated guesswork". However, it is not the arbitrariness of the calibrations *per se* that is of concern, but rather that they act as a strong source of pseudo-data. Taken to a hyberbolic extreme, if calibration priors were applied to every node in a tree, then the results above would suggest that there would be no use in running the analysis at all. In other types of Bayesian analysis we expect that an increase in data can overrule poorly constructed priors and converge on an answer. It is presently unclear whether this is true when performing node-dating using the UCLN model, and if so, how much data (or 'information', whatever that turns out to be) would be required. Certainly it is known that the uncertainty in divergence time estimates cannot be reduced arbitrarily, even with infinite amounts of data (Rannala and Yang 2007; Yang and Rannala 2006; dos Reis and Yang 2013), but it is unclear how data can override the pseudo-data present in the node calibration priors.

WHERE TO GO FROM HERE?

The results presented here highlight several issues that should be considered as the field moves forward. In regard to angiosperms, is the amount of temporal 'information' present in empirical data (or, on the other hand, the adequacy of current relaxed clock models) insufficient to reconstruct such recalcitrant nodes as the age of crown angiosperms? If this is the case, the only way forward, given the methods of inference, may be to apply user priors that are *intended* to constitute pseudo-data. However, this fact needs to be more widely recognized. In some respects this is a defensible position as, if fossil calibrations are constructed with significant information about the fossil record, estimations will be constrained *to* existing fossil information. In this vein, the results of Magallón *et al.* (2015), which estimate nested angiosperm divergence times within a strict paleontologically-imposed age of the ancestral crown node, are reasonable in the context of the data available (Sanderson, 2015). Nevertheless, it should be clear when the molecular data, in this context, do not significantly alter the posterior distribution. If this is indeed the way forward, then care should be taken to assess both the validity of the fossils being used (Sanders and Lee 2007; Brown and Sorhannus 2010) and the form of the calibration priors (Inoue *et al.* 2010; Brown and Sorhannus 2010; Sauquet *et al.* 2011; Warnock *et al.* 2012; Foster *et al.* 2016). In addition, we strongly advocate the regular use of the diptych approach to data analysis by habitually comparing prior and posterior distributions: it is imperative to understand which parameters in our models are informed by the data present, and which simply recapitulate the prior. When hypothesis testing it is even more critically important to determine whether a hypothesis is rejected by the data or, as with the crown angiosperm age results above, are effectively precluded by the joint prior.

However, new methods of divergence time inference are emerging that largely bypass the concerns associated with node-dating (Heath and Moore, 2014). The fossilized birth-death model of Heath *et al.* (2014) incorporates extant and extinct (i.e., sampled fossils) lineages as evolving according to the same underlying diversification model. Alternatively, when morphological data are available for both extinct and extant taxa, divergence times can be estimated using the tip-dating approach of Ronquist *et al.* (2012). Both of these methods can take advantage of an arbitrary number of fossils within a lineage (rather than being reduced to a single distribution as in node-dating) and incorporate fossil temporal information directly without extrapolation. The excitement surrounding these methods might lead us to think it not unreasonable to suppose that in the near future node-dating will be regarded as a useful tool that was ultimately replaced by methods that more directly make use of the available data. However, both of these methods are relatively new, and it is unclear whether they will overthrow node-dating results for the most recalcitrant nodes (i.e., placental mammals, crown birds, crown angiosperms, etc.). These methods also raise new questions in regard to model adequacy, implied and explicit assumptions regarding both diversification and morphology models, and data availability and quality for extinct and extant lineages. Furthermore, the resulting divergence time estimates from these new methods may not differ as much as expected. For example, Eguchi and Tamura (2016) employed the fossilized birth-death model and found monocots arose 174.26–134.14 Ma, which does not conflict strongly with previous node-dating results. It thus appears far too premature to consider the tempo of angiosperm diversification solved.

FINAL THOUGHTS ON THE AGE OF ANGIOSPERMS

Finally, we note that the results presented above do not refute the concerns identified and demonstrated through simulation by Beaulieu *et al.* (2015) regarding violations of biological modelling. While among-lineage molecular substitution rate heterogeneity is regarded as ubiquitous, clade- and trait-specific correlations of rate variation explored by Beaulieu *et al.* (2015) are becoming increasingly recognized as important biological patterns of molecular evolution (Smith *et al.* 2010; Dornburg *et al.* 2012; Lartillot and Delsuc 2012; Worobey *et al.* 2014). Such processes must be correctly modelled if our divergence time estimates are to be accurate. In this vein, we note that the fit (Lepage *et al.*, 2007) and adequacy (Duchênene *et al.*, 2015) of alternative clock models are far too rarely assessed. In addition, the artefacts of lineage sampling identified by Beaulieu *et al.* (2015) casts doubt on the suitability of a homogeneous

birth-death model as a prior on node ages. This doubt is especially manifested with respect to dating the evolution of angiosperms, where it is known *a priori* that lineages exhibit an incredible breadth of diversification rates (Tank *et al.*, 2015), not to mention that the extant angiosperm diversity dwarfs other embryophyte clades. We thus regard our results above as complementary to those of Beaulieu *et al.* (2015), and only with both in mind can we confidently move forward.

Supplementary Material

Data available from the Dryad Data Repository:http://dx.doi.org/10.5061/dryad.XXXX.

Funding

J.W.B. and S.A.S. were both supported by the NSF AVATOL Grant 1207915.

Acknowledgements

We thank Jeremy Beaulieu and Brian O'Meara for sharing their thoughts on these issues, and Susana Magallón for graciously sharing her BEAST xml input file. We would also like to thank the following for helpful discussions: Ben Redelings, Joseph Walker, Ning Wang, Greg Stull, Aaron King, Michael Landis, Alex Taylor, Paul Lewis, and members of the Smith laboratory. J.W.B. thanks Don Van Vliet for help with the title, Emo Philips for help with broader dissemination, Mark Kozelek for encouragement, and Linda Slote for tracking down an obscure Testudines reference (that was unfortunately dropped from an earlier version of this paper).

REFERENCES

Ayres, D. L., Darling, A., Zwickl, D. J., Beerli, P., Holder, M. T., Lewis, P. O., Huelsenbeck, J. P., Ronquist, F., Swofford, D. L., Cummings, M. P., Rambaut, A., and Suchard, M. A. 2012. BEAGLE: an application programming interface and high-performance computing library for statistical phylogenetics. *Systematic Biology*, 61(1): 170–173.

Beaulieu, J. M., O'Meara, B. C., Crane, P., and Donoghue, M. J. 2015. Heterogeneous rates of molecular evolution and diversification could explain the Triassic age estimate for angiosperms. *Systematic Biology*, 64(5): 869–878.

Bell, C. D., Soltis, D. E., and Soltis, P. S. 2010. The age and diversification of the angiosperms re-revisited. *American Journal of Botany*, 97(8): 1296–1303.

Benton, M. J. 1999. Early origins of modern birds and mammals: molecules vs. morphology. *BioEssays*, 21(12): 1043–1051.

Benton, M. J. and Donoghue, P. C. J. 2007. Paleontological evidence to date the tree of life. *Molecular Biology and Evolution*, 24(1): 26–53.

Brenner, G. J. 1996. Evidence for the earliest stage of angiosperm pollen evolution: A paleoequatorial section from israel. In D. W. Taylor and L. J. Hickey, editors, *Flowering Plant Origin, Evolution & Phylogeny*, pages 91–115. Springer US, Boston, MA.

Brown, J. W. and Sorhannus, U. 2010. A molecular genetic timescale for the diversification of autotrophic stramenopiles (Ochrophyta): Substantive underestimation of putative fossil ages. *PLOS ONE*, 5(9): 1–11.

Brown, J. W. and van Tuinen, M. 2011. Evolving perceptions on the antiquity of the modern avian tree. In G. Dyke and G. Kaiser, editors, *Living Dinosaurs: The Evolutionary History of Modern Birds*, chapter 12, pages 306–324. John Wiley & Sons Ltd, Oxford.

Brown, J. W., Payne, R. B., and Mindell, D. P. 2007. Nuclear DNA does not reconcile 'rocks' and 'clocks' in Neoaves: a comment on Ericson et al. *Biology Letters*, 3(3): 257–260.

Brown, J. W., Walker, J. F., and Smith, S. A. 2017. phyx: Phylogenetic tools for unix. *Bioinformatics*. in press.

Claramunt, S. and Cracraft, J. 2015. A new time tree reveals Earth history's imprint on the evolution of modern birds. *Science Advances*, 1(11).

Donoghue, P. C. and Benton, M. J. 2007. Rocks and clocks: calibrating the tree of life using fossils and molecules. *Trends in Ecology & Evolution*, 22(8): 424–431.

Dornburg, A., Brandley, M. C., McGowen, M. R., and Near, T. J. 2012. Relaxed clocks and inferences of heterogeneous patterns of nucleotide substitution and divergence time estimates across whales and dolphins (Mammalia: Cetacea). *Molecular Biology and Evolution*, 29(2): 721–736.

dos Reis, M. and Yang, Z. 2013. The unbearable uncertainty of Bayesian divergence time estimation. *Journal of Systematics and Evolution*, 51(1): 30–43.

Drummond, A. J. and Rambaut, A. 2007. BEAST: Bayesian evolutionary analysis by sampling trees. *BMC Evolutionary Biology*, 7(1): 214.

Drummond, A. J., Ho, S. Y. W., Phillips, M. J., and Rambaut, A. 2006. Relaxed phylogenetics and dating with confidence. *PLOS Biology*, 4(5).

Duchênene, D. A., DuchDuchênene, S., Holmes, E. C., and Ho, S. Y. 2015. Evaluating the adequacy of molecular clock models using posterior predictive simulations. *Molecular Biology and Evolution*, 32(11): 2986.

Eguchi, S. and Tamura, M. N. 2016. Evolutionary timescale of monocots determined by the fossilized birth-death model using a large number of fossil records. *Evolution*, 70(5): 1136–1144.

Ericson, P. G., Anderson, C. L., Britton, T., Elzanowski, A., Johansson, U. S., Källersjö, M., Ohlson, J. I., Parsons, T. J., Zuccon, D., and Mayr, G. 2006. Diversification of Neoaves: integration of molecular sequence data and fossils. *Biology Letters*, 2(4): 543–547.

Foster, C. S. P., Sauquet, H., van der Merwe, M., McPherson, H., Rossetto, M., and Ho, S. Y. W. 2016. Evaluating the impact of genomic data and priors on Bayesian estimates of the angiosperm evolutionary timescale. *Systematic Biology*.

Heath, T. A. and Moore, B. R. 2014. Bayesian inference of species divergence times. In M.-H. Chen, L. Kuo, and P. O. Lewis, editors, *Bayesian Phylogenetics: Methods Algorithms, and Applications*, chapter 13, pages 277–318. CRC Press, Boca Raton, Florida.

Heath, T. A., Huelsenbeck, J. P., and Stadler, T. 2014. The fossilized birthdeath process for coherent calibration of divergence-time estimates. *Proceedings of the National Academy of Sciences*, 111(29): E2957–E2966.

Heibl, C. 2008. Phyloch: R language tree plotting tools and interfaces to diverse phylogenetic software packages.

Heled, J. and Drummond, A. J. 2012. Calibrated tree priors for relaxed phylogenetics and divergence time estimation. *Systematic Biology*, 61(1): 138–149.

Heled, J. and Drummond, A. J. 2015. Calibrated birthdeath phylogenetic time-tree priors for Bayesian inference. *Systematic Biology*, 64(3): 369–383.

Ho, S. Y. W. and Phillips, M. J. 2009. Accounting for calibration uncertainty in phylogenetic estimation of evolutionary divergence times. *Systematic Biology*, 58(3): 367–380.

Inoue, J., Donoghue, P. C. J., and Yang, Z. 2010. The impact of the representation of fossil calibrations on Bayesian estimation of species divergence times. *Systematic Biology*, 59(1): 74–89.

Jarvis, E. D., Mirarab, S., Aberer, A. J., Li, B., Houde, P., Li, C., Ho, S. Y. W., Faircloth, B. C., Nabholz, B., Howard, J. T., Suh, A., Weber, C. C., da Fonseca, R. R., Li, J., Zhang, F., Li, H., Zhou, L., Narula, N., Liu, L., Ganapathy, G., Boussau, B., Bayzid, M. S., Zavidovych, V., Subramanian, S., Gabaldón, T., Capella-Gutiérrez, S., Huerta-Cepas, J., Rekepalli, B., Munch, K., Schierup, M., Lindow, B., Warren, W. C., Ray, D., Green, R. E., Bruford, M. W., Zhan, X., Dixon, A., Li, S., Li, N., Huang, Y., Derryberry, E. P., Bertelsen, M. F., Sheldon, F. H., Brumfield, R. T., Mello, C. V., Lovell, P. V., Wirthlin, M., Schneider, M. P. C., Prosdocimi, F., Samaniego, J. A., Velazquez, A. M. V., Alfaro-Núñez, A., Campos, P. F., Petersen, B., Sicheritz-Ponten, T., Pas, A., Bailey, T., Scofield, P., Bunce, M., Lambert, D. M., Zhou, Q., Perelman, P., Driskell, A. C., Shapiro, B., Xiong, Z., Zeng, Y., Liu, S., Li, Z., Liu, B., Wu, K., Xiao, J., Yinqi, X., Zheng, Q., Zhang, Y., Yang, H., Wang, J., Smeds, L., Rheindt, F. E., Braun, M., Fjeldsa, J., Orlando, L., Barker, F. K., Jønsson, K. A., Johnson, W., Koepfli, K.-P., O'Brien, S., Haussler, D., Ryder, O. A., Rahbek, C., Willerslev, E., Graves, G. R., Glenn, T. C., McCormack, J., Burt, D., Ellegren, H., Alström, P., Edwards, S. V., Stamatakis, A., Mindell, D. P., Cracraft, J., Braun, E. L., Warnow, T., Jun, W., Gilbert, M. T. P., and Zhang, G. 2014. Whole-genome analyses resolve early branches in the tree of life of modern birds. *Science*, 346(6215): 1320–1331.

Ksepka, D. T., Ware, J. L., and Lamm, K. S. 2014. Flying rocks and flying clocks: disparity in fossil and molecular dates for birds. *Proceedings of the Royal Society of London B: Biological Sciences*, 281(1788).

Ksepka, D. T., Parham, J. F., Allman, J. F., Benton, M. J., Carrano, M. T., Cranston, K. A., Donoghue, P. C. J., Head, J. J., Hermsen, E. J., Irmis, R. B., Joyce, W. G., Kohli, M., Lamm, K. D., Leehr, D., Patan, J. L., Polly, P. D., Phillips, M. J., Smith, N. A., Smith, N. D., Van Tuinen, M., Ware, J. L., and Warnock, R. C. M. 2015. The fossil calibration database–a new resource for divergence dating. *Systematic Biology*, 64(5): 853–859.

Lartillot, N. and Delsuc, F. 2012. Joint reconstruction of divergence times and lifehistory evolution in placental mammals using a phylogenetic covariance modell. *Evolution*, 66(6): 1773–1787.

Lee, M. S. Y. and Skinner, A. 2011. Testing fossil calibrations for vertebrate molecular trees. *Zoologica Scripta*, 40(5): 538–543.

Lepage, T., Bryant, D., Philippe, H., and Lartillot, N. 2007. A general comparison of relaxed molecular clock models. *Molecular Biology and Evolution*, 24(12): 2669.

Lewis, P. O., Chen, M.-H., Kuo, L., Lewis, L. A., Fučíková, K., Neupane, S., Wang, Y.-B., and Shi, D. 2016. Estimating Bayesian hylogenetic information content. *Systematic Biology*, 65(6): 1009–1023.

Magallón, S., Hilu, K. W., and Quandt, D. 2013. Land plant evolutionary timeline: Gene effects are secondary to fossil constraints in relaxed clock estimation of age and substitution rates. *American Journal of Botany*, 100(3): 556–573.

Magallón, S., Gómez-Acevedo, S., Sánchez-Reyes, L. L., and Hernández-Hernández, T. 2015. A metacalibrated time-tree documents the early rise of flowering plant phylogenetic diversity. *New Phytologist*, 207(2): 437–453.

Marshall, C. R. 2008. A simple method for bracketing absolute divergence times on molecular phylogenies using multiple fossil calibration points. *The American Naturalist*, 171(6): 726–742.

Meredith, R. W., Janečka, J. E., Gatesy, J., Ryder, O. A., Fisher, C. A., Teeling, E. C., Goodbla, A., Eizirik, E., Simão, T. L. L., Stadler, T., Rabosky, D. L., Honeycutt, R. L., Flynn, J. J., Ingram, C. M., Steiner, C., Williams, T. L., Robinson, T. J., Burk-Herrick, A., Westerman, M., Ayoub, N. A., Springer, M. S., and Murphy, W. J. 2011. Impacts of the Cretaceous terrestrial revolution and KPg extinction on mammal diversification. *Science*, 334(6055): 521–524.

Mitchell, K. J., Cooper, A., and Phillips, M. J. 2015. Comment on "Whole-genome analyses resolve early branches in the tree of life of modern birds". *Science*, 349(6255): 1460–1460.

Nowak, M. D., Smith, A. B., Simpson, C., and Zwickl, D. J. 2013. A simple method for estimating informative node age priors for the fossil calibration of molecular divergence time analyses. *PLOS ONE*, 8(6): 1–13.

O'Leary, M. A., Bloch, J. I., Flynn, J. J., Gaudin, T. J., Giallombardo, A., Giannini, N. P., Goldberg, S. L., Kraatz, B. P., Luo, Z.-X., Meng, J., Ni, X., Novacek, M. J., Perini, F. A., Randall, Z. S., Rougier, G. W., Sargis, E. J., Silcox, M. T., Simmons, N. B., Spaulding, M., Velazco, P. M., Weksler, M., Wible, J. R., and Cirranello, A. L. 2013. The placental mammal ancestor and the post–K-Pg radiation of placentals. *Science*, 339(6120): 662–667.

Parham, J. F., Donoghue, P. C. J., Bell, C. J., Calway, T. D., Head, J. J., Holroyd, P. A., Inoue, J. G., Irmis, R. B., Joyce, W. G., Ksepka, D. T., Patané, J. S. L., Smith, N. D., Tarver, J. E., van Tuinen, M., Yang, Z., Angielczyk, K. D., Greenwood, J. M., Hipsley, C. A., Jacobs, L., Makovicky, P. J., Mller, J., Smith, K. T., Theodor, J. M., Warnock, R. C. M., and Benton, M. J. 2012. Best practices for justifying fossil calibrations. *Systematic Biology*, 61(2): 346–359.

Phillips, M. J. 2009. Branch-length estimation bias misleads molecular dating for a vertebrate mitochondrial phylogeny. *Gene*, 441(12): 132–140.

Prum, R. O., Berv, J. S., Dornburg, A., Field, D. J., Townsend, J. P., Lemmon, E. M., and Lemmon, A. R. 2015. A comprehensive phylogeny of birds (Aves) using targeted next-generation DNA sequencing. *Nature*, 526(7574): 569–573.

R Core Team 2016. *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria.

Rannala, B. and Yang, Z. 2007. Inferring speciation times under an episodic molecular clock. *Systematic Biology*, 56(3): 453–466.

Revell, L. J., Harmon, L. J., Glor, R. E., and Linder, P. 2005. Under-parameterized model of sequence evolution leads to bias in the estimation of diversification rates from molecular phylogenies. *Systematic Biology*, 54(6): 973–983.

Ronquist, F., Klopfstein, S., Vilhelmsen, L., Schulmeister, S., Murray, D. L., and Rasnitsyn, A. P. 2012. A total-evidence approach to dating with fossils, applied to the early radiation of the Hymenoptera. *Systematic Biology*, 61(6): 973–999.

Sanders, K. L. and Lee, M. S. 2007. Evaluating molecular clock calibrations using bayesian analyses with soft and hard bounds. *Biology Letters*, 3(3): 275–279.

Sanderson, M. J. 2003. r8s: inferring absolute rates of molecular evolution and divergence times in the absence of a molecular clock. *Bioinformatics*, 19(2): 301–302.

Sanderson, M. J. 2015. Back to the past: a new take on the timing of flowering plant diversification. *New Phytologist*, 207(2): 257–259.

Sauquet, H., Ho, S. Y. W., Gandolfo, M. A., Jordan, G. J., Wilf, P., Cantrill, D. J., Bayly, M. J., Bromham, L., Brown, G. K., Carpenter, R. J., Lee, D. M., Murphy, D. J., Sniderman, J. M. K., and Udovicic, F. 2011. Testing the impact of calibration on molecular divergence times using a fossil-rich group: The case of Nothofagus (Fagales). *Systematic Biology*, 61(2): 289–313.

Schenk, J. J. and Hufford, L. 2010. Effects of substitution models on divergence time estimates: Simulations and an empirical study of model uncertainty using cornales. *Systematic Botany*, 35(3): 578–592.

Shannon, C. E. 1948. Modeling compositional heterogeneity. *Bell System Technical Journal*, 27: 379–423, 623–656.

Smith, S. A., Beaulieu, J. M., and Donoghue, M. J. 2010. An uncorrelated relaxed-clock analysis suggests an earlier origin for flowering plants. *Proceedings of the National Academy of Sciences*, 107(13): 5897–5902.

Stein, R. W., Brown, J. W., and Mooers, A. Ø. 2015. A molecular genetic time scale demonstrates cretaceous origins and multiple diversification rate shifts within the order Galliformes (Aves). *Molecular Phylogenetics and Evolution*, 92: 155–164.

Tank, D. C., Eastman, J. M., Pennell, M. W., Soltis, P. S., Soltis, D. E., Hinchliff, C. E., Brown, J. W., Sessa, E. B., and Harmon, L. J. 2015. Nested radiations and the pulse of angiosperm diversification: increased diversification rates often follow whole genome duplications. *New Phytologist*, 207(2): 454–467. 2014-18202.

Thorne, J. L. and Kishino, H. 2002. Divergence time and evolutionary rate estimation with multilocus data. *Systematic Biology*, 51(5): 689.

Warnock, R. C. M., Yang, Z., and Donoghue, P. C. J. 2012. Exploring uncertainty in the calibration of the molecular clock. *Biology Letters*, 8(1): 156–159.

Wickham, H. 2009. *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York.

Worobey, M., Han, G.-Z., and Rambaut, A. 2014. A synchronized global sweep of the internal genes of modern avian influenza virus. *Nature*, 508(7495): 254–257.

Yang, Z. and Rannala, B. 2006. Bayesian estimation of species divergence times under a molecular clock using multiple fossil calibrations with soft bounds. *Molecular Biology and Evolution*, 23(1): 212–226.

Zeng, L., Zhang, Q., Sun, R., Kong, H., Zhang, N., and Ma, H. 2014. Resolution of deep angiosperm phylogeny using conserved nuclear genes and estimates of early divergence times. *Nature Communications*, 5: 4956.

Zhu, T., Dos Reis, M., and Yang, Z. 2015. Characterization of the uncertainty of divergence time estimation under relaxed molecular clock models using multiple loci. *Systematic Biology*, 64(2): 267–280.