1

No general relationship between mass and temperature in endothermic species

Kristina Riemer[1*], Robert P Guralnick[2], Ethan White[1,3]

Bergmann's rule is a widely-accepted biogeographic rule stating that individuals within a species are smaller in warmer environments. While there are many single-species studies and integrative reviews documenting this pattern, a data-intensive approach has not been used yet to determine the generality of this pattern. We assessed the strength and direction of the intraspecific relationship between temperature and individual mass for 952 bird and mammal species. For eighty-seven percent of species, temperature explained less than 10% of variation in mass, and for 79% of species the correlation was not statistically significant. These results suggest that Bergmann's rule is not general and temperature is not a dominant driver of biogeographic variation in mass. Further understanding of size variation will require integrating multiple processes that influence size. The lack of dominant temperature forcing weakens the justification for the hypothesis that global warming could result in widespread decreases in body size.

[1]Department of Wildlife Ecology and Conservation, 110 Newins-Ziegler Hall, PO Box 110430, University of Florida, Gainesville, FL 32611-0430, USA
[2]Department of Natural History, University of Florida, Gainesville, FL 32611
[3]Informatics Institute, 432 Newell Drive, PO Box 115585, University of Florida, Gainesville, FL 32611-8545

Correspondence and requests for materials should be addressed to K.R. (email: kristina.riemer@weecology.org)

INTRODUCTION

Bergmann's rule describes a negative relationship between body mass and temperature across space that is believed to be common in endothermic species (Bergmann, 1847; Brown and Lee, 1969; Kendeigh, 1969; Freckleton et al., 2003; Carotenuto et al., 2015). Many hypotheses have been proposed to explain this pattern (Blackburn et al., 1999; Ashton, 2002; Watt et al., 2010) including the heat loss hypothesis, which argues that the higher surface area to volume ratio of smaller individuals results in improved heat dissipation in hot environments (Bergmann, 1847). Though originally described for closely-related species (i.e., interspecific; Blackburn et al., 1999), the majority of studies have focused on the intraspecific form of Bergmann's rule (Rensch, 1938; Meiri, 2011) by assessing trends in individual size within a species (Langvatn and Albon, 1986; Gardner et al., 2009; Yom-Tov and Geffen, 2006). Bergmann's rule has been questioned both empirically and mechanistically (McNab, 1971; Geist, 1987; Huston and Wolverton, 2011; Teplitsky & Millien, 2014) but the common consensus from recent reviews is that the pattern is general (Ashton, 2002; Watt et al., 2011; Ashton et al., 2000; Meiri and Dayan, 2003).

It has recently been suggested that this negative relationship between mass and temperature could result in decreasing individual size across species in response to climate change (Sheridan and Bickford, 2011) and that this may be a "third universal response to warming" (Gardner et al., 2011). The resulting shifts in size distributions could significantly alter ecological communities (Brose et al., 2012), especially if the rate of size decrease varies among species (Sheridan and Bickford, 2011). While there is limited empirical research on body size responses to changes in temperature through time (but see Teplitsky and Millien, 2014; Smith et al., 1995; Caruso et al., 2014), the apparent generality of Bergmann's rule across space indicates the likelihood of a similar relationship in response to temperature change across time.

The generality of Bergmann's rule is based on many individual studies that analyze

empirical data on body size across an environmental gradient (e.g., Langvatn and Albon, 1986;

Barnett, 1977; Fuentes and Jaksic, 1979; Dayan et al., 1989; Sand et al., 1995) and reviews that

compile and evaluate the results from these studies (Ashton, 2002; Watt et al., 2010; Meiri and

Dayan 2003). Most individual studies of Bergmann's rule are limited by: 1) analyzing only one

or a few species (e.g., Langvatn and Albon, 1986); 2) using small numbers of observations (e.g.,

Fuentes and Jaksic, 1979); 3) only including data at the small scales typical of ecological studies

(e.g., Sand et al., 1995); 4) using latitude instead of directly assessing temperature (e.g., Barnett,

1977); and 5) focusing on statistical significance instead of the strength of the relationship (e.g.,

Dayan et al., 1989). The reviews tabulate the results of these individual studies and assess

patterns in the direction and significance of relationships across species. Such aggregation of

published results allows for a more general understanding of the pattern but, in addition to

limitations of the underlying studies, the conclusions may be influenced by publication bias and

selective reporting where studies or individual analyses that do not support Bergmann's rule are

published less frequently (Koricheva et al., 2013).

Previous analyses of publication bias in the context of Bergmann's rule have found no

evidence for selective publication, which supports the idea that it is a general rule (Ashton, 2002;

Meiri et al., 2004). However, two of the most extensive studies of Bergmann's rule, which both

used museum records to assess dozens of intraspecific Bergmann's rule relationships

simultaneously, found that the majority of species did not exhibit significant negative

relationships between latitude and size (McNab, 1971; Meiri et al., 2004). As a result,

understanding the generality of this ecophysiological rule and its potential implications for global

change requires more extensive analysis.

A data-intensive approach to analyzing Bergmann's rule, evaluating the pattern using

large amounts of broad scale data, has the potential to overcome existing limitations in the

literature and provides a new perspective on the generality of the intraspecific form of

Bergmann's rule. Understanding the generality of the temperature-mass relationship has

important implications for how size will respond to climate change. We use data from Vertnet

(Constable et al., 2010), a large compilation of digitized museum records that contains over

700,000 globally distributed individual-level size measures, to evaluate the intraspecific

relationship between temperature and mass for 952 mammal and bird species. The usable data

consist of 273,901 individuals with an average of 288 individuals per species, with individuals of

each species spanning an average of 75 years and 34 latitudinal degrees. This approach reduces

or removes many of the limitations to previous approaches and the results suggest that

Bergmann's rule is not a strong or general pattern.

RESULTS

Most of the species in this study showed weak non-significant relationships between

temperature and mass (Figures 1 and 2). The distribution of correlation coefficients was centered

near zero with a mean correlation coefficient of -0.05 across species (Figure 2A). Relationships

for 79% of species were not significantly different from zero, while 14% of species' relationships

were significant and negative and 7% were significant and positive (Figure 2A). Temperature

explained less than 10% of variation in mass (i.e., $-0.316 < r < 0.316$) for 87% of species,

indicating that temperature explained very little of the observed variation in mass for these

species (Figure 2A).

The weak, non-directional intraspecific relationships indicated by the distribution of

correlation coefficients are consistent across taxonomic groups and temporal lags. Mean

correlation coefficients for both endothermic classes are -0.006 and -0.065, for mammals and

birds respectively (Figure 2B). Similarly, correlation coefficient distributions were

approximately centered on zero for all of the 30 orders analyzed ($-0.2 < \bar{r} < 0.003$ for orders with

more than 10 species; Figure 3 and Figure 3—figure supplement 1), or for migrant or nonmigrant

bird species (Figure 2-figure supplement 1). Correlation coefficient distributions for temperature-

mass relationships using lagged temperatures were centered around zero like those using

temperature from the collection year (Figure 4 and Figure 4–figure supplement 1), indicating that

there was not a temporal lag effect on the response of species' masses to temperature. Correlation

coefficients did not vary systematically by sample size (Figure 5A), extent of variation in

temperature or mass (Figure 5B,C), species' average mass (Figure 5D), or species' average

latitude (Figure 5E). While temperature is considered the actual driver, some studies use latitude

as a proxy when evaluating variation in size (Bergmann, 1847; Stillwell, 2010) Using latitude,

the mean correlation coefficient was -0.05 with no statistically significant latitude-mass

relationship for 71% of species (Figure 2–figure supplement 2), while the respective values for

temperature were -0.05 and 79% (Figure 2A). Results were robust to a variety of decisions and

stringencies about how to filter the size (Figure 2–figure supplement 3 and 4) and species data

(Figure 2—figure supplement 5 and 6).

DISCUSSION

In contrast to conventional wisdom and several recent review papers, our analysis of 952

species shows little to no support for a negative intraspecific temperature-mass relationship that

is sufficiently strong or common to be considered a biogeographic rule. Three quarters of bird

and mammal species show no significant change in mass across a temperature gradient and

temperature explained less than 10% of intraspecific variation in mass for 87% of species (Figure

2A). This was true regardless of taxonomic group (Figure 2 and 3), temporal lag in temperature

(Figure 4), species' size, location, or sampling intensity or extent (Figure 5). These results are

consistent with two previous studies that examined museum specimen size measurements across

latitude. The first study showed that 22 out of 47 North American mammal species studied had

no relationship between latitude and length, and 10 of the 25 significant relationships were

opposite the expected direction (McNab, 1971). The second found a similar proportion of non-

significant results (42/87), but a lower proportion of significant relationships that opposed the

rule (9/45) for carnivorous mammals (Meiri et al., 2004). While more species had significant

negative relationships than positive in both our study and these two museum-based studies, in all

cases less than half of species had significant negative correlations (14 – 41%). In combination

with these two smaller studies, our results suggest that there is little evidence for a strong or

general Bergmann's rule when analyzing raw data instead of summarizing published results.

Our results are inconsistent with recent reviews, which have reported that the majority of

species conform to Bergmann's rule (Ashton, 2002; Watt et al., 2010; Meiri and Dayan, 2003).

While these reviews included results that were either non-significant or opposite of Bergmann's

rule, the proportion of significant results in support of Bergmann's rule was higher and therefore

resulted in conclusions that supported the generality of the temperature-mass relationship.

Generalizing from results in the published literature involves the common challenges of

publication bias and selective reporting (Koricheva et al., 2013). In addition, because the

underlying Bergmann's rule studies typically report minimal statistical information, often

providing only relationship significance or direction instead of p-values or correlation

coefficients (Meiri and Dayan, 2003), proper meta-analyses and associated assessments of

biological significance are not possible. While several reviews found no evidence for publication

bias using limited analyses (Ashton, 2002; Meiri et al., 2004), the notable differences between

the conclusions of our data-intensive approach and those from reviews suggests that publication

bias in papers about Bergmann's rule warrants further investigation. These differences also

demonstrate the value of data-intensive approaches in ecology for overcoming potential

weaknesses and biases in the published literature. Directly analyzing large quantities of data

from hundreds of species allows us to assess the generality of patterns originally reported in

smaller studies while avoiding the risk of publication bias. This approach also makes it easier to

integrate other factors that potentially influence size into future analyses. The new insight gained

from this data-intensive approach demonstrates the value of investing in large compilations of

ecologically-relevant data (Hampton et al., 2013) and the proper training required to work with

these datasets (Hampton et al., 2017).

Our analyses and conclusions are limited to the intraspecific form of Bergmann's rule.

This is the most commonly studied and well-defined form of the relationship and the one most

amenable to analyses using large compilations of museum data. Difficulty in interpreting

Bergmann's original formulation has resulted in an array of different ideas and implementations

of interspecific analyses (Blackburn et al., 1999; Meiri and Thomas, 2007; Watt et al., 2010;

Meiri, 2011). The most common forms of these interspecific analyses involve correlations

between various species-level size metrics and environmental measures and are conducted at

various taxonomic levels from genus to class (e.g., Clauss et al., 2013; Diniz-Filho, 2007; Boyer

et al., 2010; Blackburn and Gaston, 1996). Efforts to apply data-intensive approaches to the

interspecific form of this relationship will need to address the fact that occurrence records are not

evenly distributed across the geographic range of species and determine how the many

interpretations of interspecific Bergmann's rule are related to one another and the biological

expectations for interspecific responses to temperature.

The original formulation of Bergmann's rule, and the scope of our conclusions, apply

only to endotherms. However, negative temperature-mass relationships have also been

documented in ectotherms, with the pattern referred to as the size-temperature rule (Ray, 1960;

Angilletta and Dunham, 2003). In contrast to the hypotheses for Bergmann's rule, which are

based primarily on homeostasis (Gardner et al., 2011), the size-temperature rule in ectotherms is

thought to result from differences between growth and development rates (Forster et al., 2011).

The current version of Vertnet contained ectotherm size data for only seven species, which is not

sufficient to complete a comprehensive analysis of the ectotherm size-temperature rule. Future

work exploring the ectotherm size-temperature rule in natural systems using data-intensive

approaches is necessary for understanding the generality of this rule in ectotherms, and data may

be sought for this effort in the literature or via a coordinated effort by museums to continue

digitizing size measurements for specimens.

A number of mechanisms have been suggested to explain why higher temperatures

should result in lower body sizes, including heat loss, starvation, resource availability, migratory

ability, and phylogenetic constraints (Blackburn et al., 1999). Most of the proposed hypotheses

have not been tested sufficiently to allow for strong conclusions to be drawn about their potential

to produce Bergmann's rule (Blackburn et al., 1999; Watt et al., 2010; Teplitsky and Millien,

2014) and the widely studied heat loss hypothesis has been questioned for a variety of reasons

(Blackburn et al., 1999; Watt et al., 2010; McNab, 1971; James, 1970; McNamara et al., 2016).

While no existing hypotheses have been confirmed, it is possible that some processes are

producing negative relationships between size and temperature. The lack of a strong relationship

does not preclude processes that result in a negative temperature-mass relationship, but it does

suggest that these processes are have less influence relative to other factors that influence

intraspecific size.

The relative importance of the many factors besides temperature that can influence size

within a species is as yet unknown. Size is affected by abiotic factors such as humidity and

resource availability (Teplitsky and Millien, 2014), characteristics of individuals like clutch size

(Boyer et al., 2009), and community context, including if and what kinds of gaps there are in

size-related niches (Smith et al., 2010) and the trophic effects of primary productivity on

consumer size (Sheridan and Bickford, 2011). Temperature itself can have indirect effects on

size, such as via habitat changes in water flow or food availability, that result in size responses

opposite of Bergmann's rule (Gardner et al., 2011). Anthropogenic influences have been shown

to influence the effect of temperature on size (Faurby and Araùjo, 2016), and similar impacts of

dispersal, extinctions, and the varying scales of climate change have been proposed (Clauss et

al., 2013). Because our data primarily came from North America, further analyses focused on

species native to other continents could reveal differing temperature-mass relationships due to

varying temperature regimes. While our work shows that more species have negative significant

relationships between temperature and mass than positive, only 21% of species have statistically

significant relationships and it consequently appears that some combination of other factors more

strongly drives intraspecific size variation for most taxa.

The lack of evidence for temperature as a primary determinant of size variation in

endothermic species calls into question the hypothesis that decreases in organism size may

represent a third universal response to global warming. The potentially general decline in size

with warming was addressed by assessments that evaluated dynamic body size responses to

temperature using similar approaches to the Bergmann's rule reviews discussed above (Teplitsky

and Millien, 2014; Sheridan and Bickford, 2011; Gardner et al., 2011). These temporal reviews

had similar results to those for spatial relationships, but the conclusions of these studies clearly

noted the variability in body size responses and the need for future data-intensive work (Sheridan

and Bickford, 2011; Gardner et al., 2011) using broader temperature ranges (Teplitsky and

Millien, 2014) to fully assess the temperature-size relationship.

Our results in combination with those from other studies suggest that much of the

observed variation in size is not explained simply by temperature. While there is still potential

for the size of endotherms, and other aspects of organismal physiology and morphology, to

respond to both geographic gradients in temperature and climate change, these responses may

not be as easily explained solely by temperature as has been suggested (Sheridan & Bickford,

2011; Gardner et al., 2011). This requires that future attempts to explain variation in the size of

individuals across space or time use an integrative approach to include the influence of multiple

factors, and their potential interactions, on organism size. This will be facilitated by analyzing

spatiotemporal data similar to that used in this study, which includes wide ranges of time, space,

and environmental conditions for large numbers of species and individuals. This data-intensive

approach provides a unique perspective on the general responses of bird and mammal species to

temperature, and has potential to assist in further investigation of the complex combinations of

factors that determine biogeographic patterns of endotherm size and how species respond to

changes in climate.

METHODS

Data

Organismal data were obtained from Vertnet, a publicly available data platform for

digitized specimen records from museum collections primarily in North America, but also

includes global data (Constable et al., 2013). Body mass is routinely measured when organisms

are collected, with relatively high precision and consistent methods, by most field biologists,

whose intent is to use those organisms for research and preservation in natural history collections

(Winker, 2000; Hoffmann, 2010). These measurements are included on written labels and

ledgers associated with specimens, which are digitized and provided in standard formats, e.g.,

Darwin Core (Wieczorek, 2012). In addition to other trait information, mass has recently been

extracted from Darwin Core formatted records published in Vertnet and converted to a more

usable form (Guralnick et al., 2016). This crucial step reduces variation in how these

measurements are reported by standardizing the naming conventions and harmonizing all

measurement values to the same units (Guralnick et al., 2016). We downloaded the entire

datasets for Mammalia, Aves, Amphibia, and Reptilia available in September 2016 (Bloom et al.,

2016a; Bloom et al., 2016b; Bloom et al., 2016c; Bloom et al., 2016d) using the Data Retriever

(Morris and White, 2013) and filtered for those records that had mass measurements available.

Fossil specimen records with mass measurements were removed.

We only analyzed species with at least 30 georeferenced individuals whose collection

dates spanned at least 20 years and collection locations at least five degrees latitude, in order to

ensure sufficient sample size and spatiotemporal extent to accurately represent each species'

temperature-mass relationship. To determine if these thresholds were appropriate, we

additionally explored the impact of increasing and decreasing them (Figure 2–figure supplement

5 and 6). We selected individual records with geographic coordinates for collection location,

collection dates between 1900 and 2010, and species-level taxonomic identification, which were

evaluated to assure no issues with synonymy or clear taxon concept issues. To minimize

inclusion of records of non-adult specimens, we identified the smallest mass associated with an

identified adult life stage category for each species and removed all records with mass values

below this minimum adult size. Results were not qualitatively different due to either additional

filtering based on specimen lifestage (Figure 2–figure supplement 3) or removal of outliers

(Figure 2–figure supplement 4). Temperatures were obtained from the Udel_AirT_Precip global

terrestrial raster provided by NOAA from their website at http://www.esrl.noaa.gov/psd/, a 0.5

by 0.5 decimal degree grid of monthly mean temperatures from 1900 to 2010 (Willmott and

Matsuura, 2001). For each specimen, the mean annual temperature at its collection location was

extracted for the year of collection.

This resulted in a final dataset containing records for 273,901 individuals from 952 bird

and mammal species (MSB Mammal Collection (Arctos), 2015; Ornithology Collection

Passeriformes - Royal Ontario Museum, 2015; MVZ Mammal Collection (Arctos), 2015; MVZ

Bird Collection (Arctos), 2015; KUBI Mammalogy Collection, 2016; CAS Ornithology (ORN),

2015; DMNS Bird Collection (Arctos), 2015; UCLA Donald R, 2015; DMNS Mammal

Collection (Arctos), 2015; UAM Mammal Collection (Arctos), 2015; UWBM Mammalogy

Collection, 2015; UAM Bird Collection (Arctos), 2015; UMMZ Birds Collection, 2015; CUMV

Bird Collection (Arctos), 2015; CUMV Mammal Collection (Arctos), 2015; MLZ Bird

Collection (Arctos), 2015; LACM Vertebrate Collection, 2015; CHAS Mammalogy Collection

(Arctos), 2016; Ornithology Collection Non Passeriformes - Royal Ontario Museum, 2015;

KUBI Ornithology Collection, 2014; MSB Bird Collection (Arctos), 2015; Biodiversity

Research and Teaching Collections - TCWC Vertebrates, 2015; TTU Mammals Collection,

2015; CAS Mammalogy (MAM), 2015; Vertebrate Zoology Division - Ornithology, Yale

Peabody Museum, 2015; University of Alberta Mammalogy Collection (UAMZ), 2015; UAZ

Mammal Collection, 2016; Conner Museum, 2015; SBMNH Vertebrate Zoology, 2015; Cowan

Tetrapod Collection - Birds, 2015; Cowan Tetrapod Collection - Mammals, 2015; NMMNH

Mammal, 2015; Schmidt Museum of Natural History_Mammals, 2015; USAC Mammals

Collection, 2013; MLZ Mammal Collection (Arctos), 2015; Ohio State University Tetrapod

Division - Bird Collection (OSUM), 2015; AMNH Mammal Collections, 2015; DMNH Birds,

2015; CM Birds Collection, 2015; WNMU Mammal Collection (Arctos), 2015; UCM Mammals

Collection, 2015; UWYMV Bird Collection (Arctos), 2015; NCSM Mammals Collection, 2015;

Vertebrate Zoology Division - Mammalogy, Yale Peabody Museum, 2015; HSU Wildlife

Mammals, 2016; WNMU Bird Collection (Arctos), 2015; UWBM Ornithology Collection, 2015;

UCM Birds, 2015; University of Alberta Ornithology Collection (UAMZ), 2015; SDNHM Birds

Collection, 2015). The average number of individuals per species was 288, ranging from 30 to

15,415 individuals. The species in the dataset were diverse, including volant, non-volant,

placental, and marsupial mammals, and both migratory and non-migratory birds. There were

species from all continents except Antarctica, though the majority of the data were concentrated

in North America (Figure 1A). The distribution of the species' mean masses was strongly right-

skewed, as expected for broad scale size distributions (Brown and Nicoletto, 1991), with 74% of

species having average masses less than 100 g. Size ranged from very small (3.7 g desert shrew

Notiosorex crawfordi and 2.6 g calliope hummingbird *Stellula calliope*) to very large (63 kg

harbor seal *Phoca vitulina* and 5.8 kg wild turkey *Meleagris gallopavo*). These specimen data

and code have been deposited online in the Dryad Data Repository (http://datadryad.org/).

Analysis

We fit the intraspecific relationship between mean annual temperature and mass for each

species with ordinary least squares linear regression (e.g., Figure 1B,C,D and Figure 1–figure

supplement 1-12) using the statsmodels.formula.api module in Python (Seabold and Perktold,

2010). The strength of each species' relationship was characterized by the correlation coefficient

and its significant at alpha of 0.05. When assessing statistical significance with large numbers of

correlations it is important to consider the expected distribution of these correlations under the

null model that no correlation exists for any species.

We addressed this issue by using false discovery rate control (Benajmini and Hochberg,

1995) implemented with the stats package in R (R Core Team, 2016). This method determines

the expected distribution of values for p (or Z) in the case where no relationship exists for

individual correlation and adjusts observed values to control for excessive false positives.

Specifically, it maintains the Type I error rate (proportion of false positives) across all tests at the

chosen value of alpha and therefore gives an accurate estimate of the number of significant

relationships (Benajmini and Hochberg, 1995). This allows us to estimate the number of species

with true positive and negative correlations (i.e., those that have values that exceed those

expected from the null distribution). We then compared the number of species with positive and

negative correlation coefficients, and the proportion of those with statistically significant

adjusted p-values.

We investigated various potential correlates of the strength of Bergmann's rule. Because

it has been argued that Bergmann's rule is exhibited more strongly by some groups than others

(McNab, 1971), we examined correlation coefficient distributions within each class and order.

Additionally, distributions for migrant and nonmigrant bird species were compared due to

conflicting evidence about the impact of migration on temperature-mass relationships (Ashton,

2002). As a temporal lag in size response to temperature is likely due to individuals of a species

responding to temperatures prior to their collection year (e.g., Stacey and Fellowes, 2002), we

assessed species' temperature-mass relationships using temperatures from 1 to 110 years prior to

collection year. We also examined the relationship between species' correlation coefficients and

five variables to understand potential statistical and biological influences on the results. We did

so with the number of individuals, temperature range, and mass range to determine if the

relationship was stronger when more data points or more widely varying values were available.

Because it has been argued that Bergmann's rule is stronger in larger species (Steudel et al.,

1994) and at higher latitudes (Freckleton et al., 2003; Faurby and Araùjo, 2016), we examined

variability with both mean mass and mean latitude for each species. We also conducted all

analyses using latitude instead of mean annual temperature.

Acknowledgements

Thanks to all of the Vertnet data providers, Dan McGlinn for assistance with developing this

research, and Rafael LaFrance for his trait extraction work.

Competing financial interests

No competing financial interests

REFERENCES

AMNH Mammal Collections. American Museum of Natural History. Source:
http://ipt.vertnet.org:8080/ipt/resource.do?r=amnh_mammals (source published on 2015-
02-12, accessed on 2017-10-19)
Angilletta, Michael J, and Arthur E Dunham. "The Temperature-Size Rule in Ectotherms:
Simple Evolutionary Explanations May Not Be General." *American Naturalist* 162, no. 3
(2003): 332–42. doi:10.1086/377187.
Ashton, KG, MC Tracy, and Alan De Queiroz. "Is Bergmann's Rule Valid for Mammals?" *The*
American Naturalist 156, no. 4 (2000): 390–415.
Ashton, KG. "Patterns of Within-species Body Size Variation of Birds: Strong Evidence for
Bergmann's Rule." *Global Ecology and Biogeography* 11 (2002): 505–23.
Barnett, Raymond J. "Bergmann â   S Rule and Variation in Structures Related to Feeding in
the Gray Squirrel." *Evolution* 31, no. 3 (1977): 538–45.
Benajmini, Yoav, and Yosef Hochberg. "Controlling the False Discovery Rate : A Practical and
Powerful Approach to Multiple Testing Author (S): Yoav Benjamini and Yosef Hochberg
Source : Journal of the Royal Statistical Society . Series B (Methodological), Vol . 57 , No
. 1 Published by :" *J R Statist Soc B* 57, no. 1 (1995): 289–300.
Bergmann, C. "Uber Die Verhaltnisse Der Warmekonomie Der Tiere Zu Ihrer Grosse."
Gottinger Studien 3 (1847): 595–708.

Biodiversity Research and Teaching Collections - TCWC Vertebrates. Texas A&M University Biodiversity Research and Teaching Collections. Source: http://ipt.vertnet.org:8080/ipt/resource.do?r=tcwc_verts (source published on 2015-09-05, accessed on 2017-10-19)

Blackburn, TM, and KJ Gaston. "Spatial Patterns in the Body Sizes of Bird Species in the New World." *Oikos* 77 (1996): 436–46.

Blackburn, TM, KJ Gaston, and Natasha Loder. "Geographic Gradients in Body Size: A Clarification of Bergmann's Rule." *Diversity and Distributions*, 1999, 165–74.

Bloom, David, John Wieczorek, and Laura Russell. "Vertnet_Amphibia_Sep2016," 2016. http://datacommons.cyverse.org/browse/iplant/home/shared/commons_repo/curated/Vertnet _Amphibia_Sep2016.

Bloom, David, John Wieczorek, and Laura Russell. "Vertnet_Aves_Sep2016," 2016. http://datacommons.cyverse.org/browse/iplant/home/shared/commons_repo/curated/Vertnet _Aves_Sep2016.

Bloom, David, John Wieczorek, and Laura Russell. "Vertnet_Mammalia_Sep2016," 2016. http://datacommons.cyverse.org/browse/iplant/home/shared/commons_repo/curated/Vertnet _Mammalia_Sep2016.

Bloom, David, John Wieczorek, and Laura Russell. "Vertnet_Reptilia_Sep2016," 2016. http://datacommons.cyverse.org/browse/iplant/home/shared/commons_repo/curated/Vertnet _Reptilia_Sep2016.

Boyer, Alison G., Jean-Luc E. Cartron, and James H. Brown. "Interspecific Pairwise Relationships among Body Size, Clutch Size and Latitude: Deconstructing a Macroecological Triangle in Birds." *Journal of Biogeography* 37, no. 1 (August 2009): 47–56. doi:10.1111/j.1365-2699.2009.02175.x.

Brose, Ulrich, Jennifer A Dunne, Jose M Montoya, Owen L Petchey, Florian D Schneider, and Ute Jacob. "Climate Change in Size-Structured Ecosystems." *Philosophical Transactions of the Royal Society B: Biological Sciences* 367, no. 1605 (2012): 2903–12. doi:10.1098/rstb.2012.0232.

Brown, James H, and Paul F Nicoletto. "Spatial Scaling of Species Composition: Body Masses of North American Land Mammals." *The American Naturalist* 138, no. 6 (1991): 1478–1512. doi:10.1086/285297.

Brown, James H., and Anthony K. Lee. "Bergmann â ™ S Rule and Climatic Adaptation in Woodrats (Neotoma)." *Evolution* 23, no. 2 (1969): 329–38.

Carotenuto, Francesco, José Alexandre Felizola Diniz-Filho, and Pasquale Raia. "Space and Time: The Two Dimensions of Artiodactyla Body Mass Evolution." *Palaeogeography, Palaeoclimatology, Palaeoecology* 437 (2015): 18–25. doi:10.1016/j.palaeo.2015.07.013.

Caruso, Nicholas M., Michael W. Sears, Dean C. Adams, and Karen R. Lips. "Widespread Rapid Reductions in Body Size of Adult Salamanders in Response to Climate Change." *Global Change Biology* 20, no. 6 (2014): 1751–59. doi:10.1111/gcb.12550.

CAS Mammalogy (MAM). California Academy of Sciences. Source: http://ipt.calacademy.org:8080/ipt/resource.do?r=mam (source published on 2015-10-05, accessed on 2017-10-19)

CAS Ornithology (ORN). California Academy of Sciences. Source: http://ipt.calacademy.org:8080/ipt/resource.do?r=orn (source published on 2015-10-05, accessed on 2017-10-19)

CHAS Mammalogy Collection (Arctos). Chicago Academy of Sciences. Source: http://ipt.vertnet.org:8080/ipt/resource.do?r=chas_mammals (source published on 2016-06-

23, accessed on 2017-10-19)

Clauss, Marcus, Marie T. Dittmann, Dennis W H Müller, Carlo Meloro, and Daryl Codron. "Bergmann′s Rule in Mammals: A Cross-Species Interspecific Pattern." *Oikos* 122, no. 10 (2013): 1465–72. doi:10.1111/j.1600-0706.2013.00463.x.

CM Birds Collection. Carnegie Museum of Natural History. Source: http://ipt.vertnet.org:8080/ipt/resource.do?r=cm_birds (source published on 2015-10-11, accessed on 2017-10-19)

Conner Museum. Charles R. CRCM Vertebrate Collection. Source: http://ipt.vertnet.org:8080/ipt/resource.do?r=crcm_verts (source published on 2015-10-13, accessed on 2017-10-19)

Constable, Heather, Robert Guralnick, John Wieczorek, Carol Spencer, A. Townsend Peterson, Hank Bart, John Bates, et al. "VertNet: A New Model for Biodiversity Data Sharing." *PLoS Biology* 8, no. 2 (2010): 1–4. doi:10.1371/journal.pbio.1000309.

Cowan Tetrapod Collection - Birds. University of British Columbia Beaty Biodiversity Museum. Source: http://ipt.vertnet.org:8080/ipt/resource.do?r=ubc_bbm_ctc_birds (source published on 2015-04-07, accessed on 2017-10-19)

Cowan Tetrapod Collection - Mammals. University of British Columbia Beaty Biodiversity Museum. Source: http://ipt.vertnet.org:8080/ipt/resource.do?r=ubc_bbm_ctc_mammals (source published on 2015-04-07, accessed on 2017-10-19)

CUMV Bird Collection (Arctos). Cornell University Museum of Vertebrates. Source: http://ipt.vertnet.org:8080/ipt/resource.do?r=cumv_bird (source published on 2015-10-26, accessed on 2017-10-19)

CUMV Mammal Collection (Arctos). Cornell University Museum of Vertebrates. Source: http://ipt.vertnet.org:8080/ipt/resource.do?r=cumv_mamm (source published on 2015-10-26, accessed on 2017-10-19)

Dayan, Tamar, Eitan Tchernov, Yoram Yom-tov, and Daniel Simberloff. "Ecological Character Displacement in Saharo-Arabian Vulpes : Outfoxing Bergmann â S Rule." *Oikos* 55, no. 2 (1989): 263–72.

Diniz-Filho, José Alexandre F., Luis Mauricio Bini, Miguel Á. Rodríguez, Thiago Fernando L. V. B. Rangel, and Bradford a. Hawkins. "Seeing the Forest for the Trees: Partitioning Ecological and Phylogenetic Components of Bergmann's Rule in European Carnivora." *Ecography* 30, no. 4 (August 2007): 598–608. doi:10.1111/j.2007.0906-7590.04988.x.

DMNH Birds. Delaware Museum of Natural History. Source: http://ipt.vertnet.org:8080/ipt/resource.do?r=dmnh_birds (source published on 2015-10-20, accessed on 2017-10-19)

DMNS Bird Collection (Arctos). Denver Museum of Nature & Science. Source: http://ipt.vertnet.org:8080/ipt/resource.do?r=dmns_bird (source published on 2015-10-26, accessed on 2017-10-19)

DMNS Mammal Collection (Arctos). Denver Museum of Nature & Science. Source: http://ipt.vertnet.org:8080/ipt/resource.do?r=dmns_mamm (source published on 2015-10-26, accessed on 2017-10-19)

Faurby, Søren, and Miguel B. Araújo. "Anthropogenic Impacts Weaken Bergmann's Rule." *Ecography*, no. June (2016): 1–2. doi:10.1111/ecog.02287.

Forster, Jack, Andrew G Hirst, and Guy Woodward. "Growth and Development Rates Have Different Thermal Responses." *The American Naturalist* 178, no. 5 (November 2011): 668–78. doi:10.1086/662174.

Freckleton, RP, PH Harvey, and Mark Pagel. "Bergmann's Rule and Body Size in Mammals."

The American Naturalist 161, no. 5 (2003): 821–25.

Fuentes, Eduardo R, and Fabián M Jaksić. "Latitudinal Size Variation of Chilean Foxes: Tests of Alternative Hypotheses." *Ecology* 60, no. 1 (1979): 43–47. doi:10.2307/1936466.

Gardner, Janet L, Anne Peters, Michael R Kearney, Leo Joseph, and Robert Heinsohn. "Declining Body Size: A Third Universal Response to Warming?" *Trends in Ecology & Evolution* 26, no. 6 (June 2011): 285–91. doi:10.1016/j.tree.2011.03.005.

Gardner, Janet L, Robert Heinsohn, and Leo Joseph. "Shifting Latitudinal Clines in Avian Body Size Correlate with Global Warming in Australian Passerines." *Proceedings: Biological Sciences* 276, no. 1674 (2009): 3845–52. doi:10.2307/30245347.

Geist, V. "Bergmann's Rule Is Invalid." *Canadian Journal of Zoology*, no. 1984 (1987): 1984–87.

Guralnick, Robert P., Paula F. Zermoglio, John Wieczorek, Raphael LaFrance, David Bloom, and Laura Russell. "The Importance of Digitized Biocollections as a Source of Trait Data and a New VertNet Resource." *Database: The Journal of Biological Databases and Curation*, 2016, 1–13. doi:10.1093/database/baw158.

Hampton, S.E., M.B. Jones, L.A. Wasser, M.P. Schildhauer, S.R. Supp, J. Bruno, R.R. Hernandez, et al. "Skills and Knowledge for Data-Intensive Environmental Research." *BioScience* 67, no. 6 (2017): 546–57.

Hampton, Stephanie E., Carly a. Strasser, Joshua J. Tewksbury, Wendy K. Gram, Amber E. Budden, Archer L. Batcheller, Clifford S. Duke, and John H. Porter. "Big Data and the Future of Ecology." *Frontiers in Ecology and the Environment* 11, no. 3 (2013): 156–62. doi:10.1890/120103.

Hoffmann, A, J Decher, F Rovero, J Schaer, C Voigt, and G Wibbelt. "Field Methods and Techniques for Monitoring Mammals." *Manual on Field Recording Techniques and Protocols for All Taxa Biodiversity Inventories and Monitoring* 8 (2010): 482–529.

HSU Wildlife Mammals. Humboldt State University. Source: http://ipt.vertnet.org:8080/ipt/resource.do?r=hsu_wildlife_mammals (source published on 2016-03-14, accessed on 2017-10-19)

Huston, Michael A., and Steve Wolverton. "Regulation of Animal Size by eNPP, Bergmann's Rule, and Related Phenomena." *Ecological Monographs* 81, no. 3 (2011): 349–405. doi:10.1890/07-1861.1.

International, Birdlife. "Migratory Status of Bird Species of the World." Available from: http://datazone.birdlife.org/species/requestdis (accessed on 8/25/17) 2017.

James, Frances C. "Geographic Size Variation in Birds and Its Relationship to Climate." *Ecology* 51, no. 3 (1970): 365–90.

Kendeigh, S Charles. "Tolerance of Cold and Bergmann's Rule." *The Auk* 86, no. 1 (1969): 13–25.

Koricheva, Julia, Jessica Gurevitch, and Kerrie Mengersen. *Handbook of Meta-Analysis in Ecology and Evolution*, 2013. doi:10.2307/j.ctt24hq6n.

KUBI Mammalogy Collection. KU Biodiversity Institute. Source: http://ipt.nhm.ku.edu/ipt/resource.do?r=kubi_mammals (source published on 2016-02-03, accessed on 2017-10-19)

KUBI Ornithology Collection. KU Biodiversity Institute. Source: http://ipt.nhm.ku.edu/ipt/resource.do?r=kubi_ornithology (source published on 2014-12-17, accessed on 2017-10-19)

LACM Vertebrate Collection. Natural History Museum of Los Angeles County. Source: http://ipt.vertnet.org:8080/ipt/resource.do?r=lacm_verts (source published on 2015-04-17,

accessed on 2017-10-19)

Langvatn, R, and Steve D Albon. "Geographic Clines in Body-Weight of Norwegian Red Deer -
A Novel Explanation of Bergmann Rule." *Holarctic Ecology* 9, no. 4 (1986): 285–93.

McNab, Brian K. "On the Ecological Significance of Bergmann's Rule," 1971.

Mcnamara, John M., Andrew D. Higginson, and Simon Verhulst. "The Influence of the
Starvation-Predation Trade-off on the Relationship between Ambient Temperature and
Body Size among Endotherms." *Journal of Biogeography* 43, no. 4 (2016): 809–19.
doi:10.1111/jbi.12695.

Meiri, Shai, and Tamar Dayan. "On the Validity of Bergmann's Rule." *Journal of Biogeography*,
no. 1847 (2003): 331–51.

Meiri, Shai, Tamar Dayan, and Daniel Simberloff. "Carnivores, Biases and Bergmann' S Rule."
Biological Journal of the Linnean Society 81 (2004): 579–88. doi:10.1111/j.1095-
8312.2004.00310.x.

Meiri, Shai. "Bergmann's Rule-What's in a Name?" *Global Ecology and Biogeography* 20, no. 1
(2011): 203–7. doi:10.1111/j.1466-8238.2010.00577.x.

MLZ Bird Collection (Arctos). Moore Laboratory of Zoology. Source:
http://ipt.vertnet.org:8080/ipt/resource.do?r=mlz_bird (source published on 2015-10-26,
accessed on 2017-10-19)

MLZ Mammal Collection (Arctos). Moore Laboratory of Zoology. Source:
http://ipt.vertnet.org:8080/ipt/resource.do?r=mlz_mamm (source published on 2015-10-26,
accessed on 2017-10-19)

Morris, Benjamin D., and Ethan P. White. "The EcoData Retriever: Improving Access to
Existing Ecological Data." *PLoS ONE* 8, no. 6 (2013): 1–7.
doi:10.1371/journal.pone.0065848.

MSB Bird Collection (Arctos). Museum of Southwestern Biology. Source:
http://ipt.vertnet.org:8080/ipt/resource.do?r=msb_bird (source published on 2015-10-27,
accessed on 2017-10-19)

MSB Mammal Collection (Arctos). Museum of Southwestern Biology. Source:
http://ipt.vertnet.org:8080/ipt/resource.do?r=msb_mamm (source published on 2015-10-27,
accessed on 2017-10-19)

MVZ Bird Collection (Arctos). Museum of Vertebrate Zoology, UC Berkeley. Source:
http://ipt.vertnet.org:8080/ipt/resource.do?r=mvz_bird (source published on 2015-10-27,
accessed on 2017-10-19)

MVZ Mammal Collection (Arctos). Museum of Vertebrate Zoology, UC Berkeley. Source:
http://ipt.vertnet.org:8080/ipt/resource.do?r=mvz_mammal (source published on 2015-10-
27, accessed on 2017-10-19)

NCSM Mammals Collection. North Carolina Museum of Natural Sciences. Source:
http://ipt.vertnet.org:8080/ipt/resource.do?r=ncsm_mammals (source published on 2015-02-
20, accessed on 2017-10-19)

NMMNH Mammal. New Mexico Museum of Natural History and Science. Source:
http://ipt.vertnet.org:8080/ipt/resource.do?r=nmmnh_mammals (source published on 2015-
04-14, accessed on 2017-10-19)

Ohio State University Tetrapod Division - Bird Collection (OSUM). Ohio State University.
Source: http://hymfiles.biosci.ohio-state.edu:8080/ipt/resource.do?r=osum-birds (source
published on 2015-05-22, accessed on 2017-10-19)

Ornithology Collection Non Passeriformes - Royal Ontario Museum. Royal Ontario Museum.
Source: http://gbif.rom.on.ca:8180/ipt/resource.do?r=birdsnonpass (source published on

2015-06-02, accessed on 2017-10-19)

Ornithology Collection Passeriformes - Royal Ontario Museum. Royal Ontario Museum. Source: http://gbif.rom.on.ca:8180/ipt/resource.do?r=birdspass (source published on 2015-09-11, accessed on 2017-10-19)

R Core Team. "R: A Language and Environment for Statistical Computing," 2016. R Foundation for Statistical Computing, Vienna, Austria. https://www.r-project.org/.

Ray, Carleton. "The Application of Bergmann's and Allen's Rules to the Poikilotherms." *Journal of Morphology* 106, no. 1 (January 1960): 85–108. doi:10.1002/jmor.1051060104.

Rensch, Bernhard. "Some Problems of Geographical Variation and Species-Formation." *Proceedings of the Linnean Society of London* 150, no. 4 (1938): 275–78.

Sand, Håkan, Göran Cederlund, and Kjell Danell. "Geographical and Latitudinal Variation in Growth Patterns and Adult Body Size of Swedish Moose (Alces Alces)." *Oecologia* 102, no. 4 (1995): 433–42. doi:10.1007/BF00341355.

SBMNH Vertebrate Zoology. Santa Barbara Museum of Natural History. Source: http://ipt.vertnet.org:8080/ipt/resource.do?r=sbmnh_verts (source published on 2015-02-17, accessed on 2017-10-19)

Schmidt Museum of Natural History_Mammals. Schmidt Museum of Natural History, Emporia State University. Source: http://ipt.vertnet.org:8080/ipt/resource.do?r=kstc_schmidt_mammals (source published on 2015-02-12, accessed on 2017-10-19)

SDNHM Birds Collection. San Diego Natural History Museum. Source: http://ipt.vertnet.org:8080/ipt/resource.do?r=sdnhm_birds (source published on 2015-04-06, accessed on 2017-10-19)

Seabold, Skipper, and Josef Perktold. "Statsmodels: Econometric and Statistical Modeling with Python." *Proceedings of the 9th Python in Science Conference*, no. Scipy (2010): 57–61.

Sheridan, Jennifer A., and David Bickford. "Shrinking Body Size as an Ecological Response to Climate Change." *Nature Climate Change* 1, no. 8 (October 2011): 401–6. doi:10.1038/nclimate1259.

Smith, FA, JL Betancourt, and JH Brown. "Evolution of Body Size in the Woodrat over the Past 25,000 Years of Climate Change." *SCIENCE*, 1995. http://biology.unm.edu/fasmith/web_page_pdfs/smith_et_al_1995_science.pdf.

Smith, Felisa a, Alison G Boyer, James H Brown, Daniel P Costa, Tamar Dayan, S K Morgan Ernest, Alistair R Evans, et al. "The Evolution of Maximum Body Size of Terrestrial Mammals." *Science (New York, N.Y.)* 330, no. 6008 (2010): 1216–19. doi:10.1126/science.1194830.

Stacey, David A., and Mark D E Fellowes. "Temperature and the Development Rates of Thrips: Evidence for a Constraint on Local Adaptation?" *European Journal of Entomology* 99, no. 3 (2002): 399–404.

Steudel, K., W. P. Porter, and D. Sher. "The Biophysics of Bergmann's Rule: A Comparison of the Effects of Pelage and Body Size Variation on Metabolic Rate." *Canadian Journal of Zoology* 72, no. 1 (1994): 70–77. doi:10.1139/z94-010.

Stillwell, R. Craig. "Are Latitudinal Clines in Body Size Adaptive?" *Oikos* 119, no. 9 (2010): 1387–90. doi:10.1111/j.1600-0706.2010.18670.x.

Teplitsky, Celine, and Virginie Millien. "Climate Warming and Bergmann's Rule through Time: Is There Any Evidence?" *Evolutionary Applications* 7, no. 1 (January 2014): 156–68. doi:10.1111/eva.12129.

TTU Mammals Collection. Museum of Texas Tech University (TTU). Source:

http://ipt.vertnet.org:8080/ipt/resource.do?r=ttu_mammals (source published on 2015-02-26, accessed on 2017-10-19)

UAM Bird Collection (Arctos). University of Alaska Museum. Source: http://ipt.vertnet.org:8080/ipt/resource.do?r=uam_bird (source published on 2015-10-27, accessed on 2017-10-19)

UAM Mammal Collection (Arctos). University of Alaska Museum. Source: http://ipt.vertnet.org:8080/ipt/resource.do?r=uam_mamm (source published on 2015-10-27, accessed on 2017-10-19)

UAZ Mammal Collection. University of Arizona Museum of Natural History. Source: http://ipt.vertnet.org:8080/ipt/resource.do?r=uaz_mammals (source published on 2016-03-29, accessed on 2017-10-19)

UCLA Donald R. University of California, Los Angeles. Dickey Bird and Mammal Collection. Source: http://ipt.vertnet.org:8080/ipt/resource.do?r=ucla_birds_mammals (source published on 2015-03-02, accessed on 2017-10-19)

UCM Birds. University of Colorado Museum of Natural History. Source: http://ipt.vertnet.org:8080/ipt/resource.do?r=ucm_birds (source published on 2015-10-27, accessed on 2017-10-19)

UCM Mammals Collection. University of Colorado Museum of Natural History. Source: http://ipt.vertnet.org:8080/ipt/resource.do?r=ucm_mammals (source published on 2015-10-27, accessed on 2017-10-19)

UMMZ Birds Collection. University of Michigan Museum of Zoology. Source: http://ipt.vertnet.org:8080/ipt/resource.do?r=ummz_birds (source published on 2015-10-28, accessed on 2017-10-19)

University of Alberta Mammalogy Collection (UAMZ). University of Alberta Museums. Dataset DOI: doi:10.18165/brxhjf. Source: http://web.macs.ualberta.ca:8088/ipt/resource?r=uamz_mammalogy (source published on 2015-09-20, accessed on 2017-10-19)

University of Alberta Ornithology Collection (UAMZ). University of Alberta Museums. Dataset DOI: doi:10.18165/srgkci. Source: http://web.macs.ualberta.ca:8088/ipt/resource?r=uamz_ornithology (source published on 2015-09-24, accessed on 2017-10-19)

USAC Mammals Collection. Universidad de San Carlos de Guatemala. Source: http://ipt.vertnet.org:8080/ipt/resource.do?r=usac_mammals (source published on 2013-11-04, accessed on 2017-10-19)

UWBM Mammalogy Collection. University of Washington Burke Museum. Source: http://ipt.vertnet.org:8080/ipt/resource.do?r=uwbm_mammals (source published on 2015-02-23, accessed on 2017-10-19)

UWBM Ornithology Collection. University of Washington Burke Museum. Source: http://ipt.vertnet.org:8080/ipt/resource.do?r=uwbm_birds (source published on 2015-02-24, accessed on 2017-10-19)

UWYMV Bird Collection (Arctos). University of Wyoming Museum of Vertebrates. Source: http://ipt.vertnet.org:8080/ipt/resource.do?r=uwymv_bird (source published on 2015-10-27, accessed on 2017-10-19)

Vertebrate Zoology Division - Mammalogy, Yale Peabody Museum. Yale Peabody Museum. Source: http://ipt.peabody.yale.edu/ipt/resource.do?r=ipt_vz_mam (source published on 2015-10-09, accessed on 2017-10-19)

Vertebrate Zoology Division - Ornithology, Yale Peabody Museum. Yale Peabody Museum.

Source: http://ipt.peabody.yale.edu/ipt/resource.do?r=ipt_vz_orn (source published on
2015-10-09, accessed on 2017-10-19)
Watt, Cortney, and Volker Salewski. "Bergmann's Rule Encompasses Mechanism: A Reply to
Olalla-Tárraga (2011)." *Oikos* 120, no. 10 (October 2011): 1445–47. doi:10.1111/j.1600-
0706.2011.19968.x.
Watt, Cortney, Sean Mitchell, and Volker Salewski. "Bergmann's Rule; a Concept Cluster?"
Oikos 119, no. 1 (January 2010): 89–100. doi:10.1111/j.1600-0706.2009.17959.x.
Wieczorek, John, David Bloom, Robert Guralnick, Stan Blum, Markus Döring, Renato
Giovanni, Tim Robertson, and David Vieglais. "Darwin Core: An Evolving Community-
Developed Biodiversity Data Standard." *PLoS ONE* 7, no. 1 (2012): e29715–e29715.
doi:10.1371/journal.pone.0029715.
Willmott, C.J., and K. Matsuura. "Terrestrial Air Temperature and Precipitation: Monthly and
Annual Time Series (1950-1999)," 2001.
http://climate.geog.udel.edu/~climate/html_pages/README.ghcn_ts2.html.
Winker, Kevin. "Obtaining, Preserving, and Preparing Bird Specimens." *Journal of Field
Ornithology* 71, no. 2 (2000): 250–97. doi:10.1648/0273-8570-71.2.250.
WNMU Bird Collection (Arctos). Western New Mexico University. Source:
http://ipt.vertnet.org:8080/ipt/resource.do?r=wnmu_bird (source published on 2015-10-27,
accessed on 2017-10-19)
WNMU Mammal Collection (Arctos). Western New Mexico University. Source:
http://ipt.vertnet.org:8080/ipt/resource.do?r=wnmu_mamm (source published on 2015-10-
27, accessed on 2017-10-19)
Yom-Tov, Yoram, and Eli Geffen. "Geographic Variation in Body Size: The Effects of Ambient
Temperature and Precipitation." *Oecologia* 148, no. 2 (2006): 213–18. doi:10.1007/s00442-006-
0364-9.

701
702
703 **FIGURE LEGENDS**
704

705
706
Figure 1. Species spatial distributions and selected temperature-mass relationships. (A)
Spatial collection locations of all individual specimens. All species shown with black points
except three species, whose relationships between mean annual temperature and mass are shown
at bottom (B-D), are marked with colored points. These species were chosen as representative of
the variability in relationship strength and direction exhibited by the 952 species from the study:
Martes pennanti had a negative relationship with temperature explaining a substantial amount of
variation in mass (B; blue points); *Tamias quadrivittatus* had no directional relationship between
temperature and mass with temperature having little explanatory power (C; yellow points);
Synaptomys cooperi had a strong positive temperature-mass relationship with a correlation
coefficient (r) in the 99[th] percentile of all species' values (D; red points). Intraspecific
temperature-mass relationships are shown with black circles for all individuals and ordinary least
squares regression trends as blue lines. Linear regression correlation coefficients and p-values in
upper left hand corner of figure for each species. See figure supplement 1.
720

721
722
Figure 2. Species correlation coefficients by statistical significance and taxonomic class. (A)
Stacked histogram of correlation coefficients (r) for all species' intraspecific temperature-mass
relationships. Colored bars show species with statistically significant relationships, both negative
(purple) and positive (green), while white bars indicate species with relationship slopes that are
not significantly different from zero. Percentages shown are percentages of species in each
group. (B) Stacked histogram of all species' correlation coefficients with bar color corresponding
to taxonomic class. Dark vertical lines are correlation coefficients of zero. See figure
supplements 1-6.
731

732
733
734 **Figure 3. Species correlation coefficients for selected taxonomic orders.** Histograms of
735 correlation coefficients (r) from intraspecific temperature-mass relationships for each taxonomic
736 order represented by more than ten species, with order shown above histogram. Height of y-axis
737 varies depending on number of species. Bar color indicates taxonomic class. Dark vertical lines
738 are correlation coefficients of zero. For remaining orders, see figure supplement 1.
739

Figure 4. Species correlation coefficients with selected past year temperatures. Histograms of correlation coefficients (r) for all species' intraspecific temperature-mass relationships with mean annual temperature from (A) the year in which individuals were collected, (B) 25 years prior to collection year, and (C) 50 years prior to collection year. Dark vertical lines are correlation coefficients of zero. For all past year temperatures, see figure supplement 1.

748
749

Figure 5. Variability of species correlation coefficients across several variables. Variation in all species' correlation coefficients (r) across the following variables for each species: (A) number of individuals, (B) difference between hottest and coldest collection year temperatures, (C) mass range, (D) mean mass, and (E) absolute mean latitude. Horizontal lines are correlation coefficients of zero. The x-axes of some plots (A, C, D) are on a log scale to better show spread of values.