The Protein Phosphatase 4 complex promotes the Notch pathway and *wingless* transcription

Eric T. Hall[1,*], Tirthadipa Pradhan-Sundd[1,2,*], Faaria Samnani[1] and Esther M. Verheyen[1,#]

[1] Department of Molecular Biology and Biochemistry, Centre for Cell Biology, Development and Disease, Simon Fraser University, British Columbia, Canada

[2] Present address: Department of Pathology, University of Pittsburgh School of Medicine, Pittsburgh, Pennsylvania, USA

* These authors contributed equally to the study

#Corresponding author: everheye@sfu.ca, T: 1-778-782-4665, F: 1-778-782-5583

Running Title: PP4 regulates Notch signaling

Keywords: PP4, Flfl, Wingless Signaling, Notch Signaling

Summary statement: The protein phosphatase 4 complex promotes Notch signaling and target gene expression during Drosophila wing development.

Abstract:

The Wnt/Wingless (Wg) pathway controls cell fate specification, tissue differentiation and organ development across organisms. Using an *in vivo* RNAi screen to identify novel kinase and phosphatase regulators of the Wg pathway, we identified subunits of the serine threonine phosphatase Protein phosphatase 4 (PP4). Knockdown of the catalytic and the regulatory subunits of PP4 cause reductions in the Wg pathway targets Senseless and Distal-less. We find that PP4 regulates the Wg pathway by controlling Notch-driven *wg* transcription. Genetic interaction experiments identified that PP4 likely promotes Notch signaling within the nucleus of the Notch-receiving cell. Although the PP4 complex is implicated in various cellular processes, its role in the regulation of Wg and Notch pathways was previously uncharacterized. Our study identifies a novel role of PP4 in regulating Notch pathway, resulting in aberrations in Notch-mediated transcriptional regulation of the Wingless ligand. Furthermore, we show that PP4 regulates proliferation independent of its interaction with Notch.

Introduction

The progression from a fertilized egg into a multicellular organism is a complex process, requiring proliferation and intricate cell-cell communication between individual cells for the eventual formation of tissues and organs. Only a handful of evolutionarily conserved signal transduction pathways are used reiteratively, both spatially and temporally to control development. In metazoans, the Wnt signaling [Wingless (Wg) in Drosophila] pathway regulates growth and proliferation, cell-fate differentiation, stem-cell renewal and homeostasis (Clevers and Nusse, 2012; Swarup and Verheyen, 2012). Wnt signaling alone does not control all these processes; its activity is extensively regulated by other signaling pathways and cellular mechanisms (Collu et al., 2014; Itasaki and Hoppler, 2010; Kim and Jho, 2014; Zeller et al., 2013). Determining how these interactions occur is critical for understanding basic cellular function and disease progression, as the disruption of the Wnt pathway has been implicated in a variety of developmental disorders and cancer (Clevers and Nusse, 2012).

The Drosophila wing imaginal disc is a powerful tool for studying Wg signaling (Swarup and Verheyen, 2012). In the developing wing disc the Wg ligand is expressed throughout different stages of disc development. At the end of the larval third instar stage, Wg expression is

confined to the presumptive wing margin along the dorsal/ventral (D/V) boundary, which

controls patterning and fate specification (Couso et al., 1994; Williams et al., 1993). Wg

produced in this narrow band of cells is secreted and acts as a morphogen to induce the nested

expression of target genes including Senseless (Sens) and Distal-less (Dll), in the flanking non-

boundary cells (Neumann and Cohen, 1997; Zecca et al., 1996).

The directed expression of Wg at the D/V boundary requires the transmembrane receptor

Notch in these boundary cells. The Notch ligands Delta (Dl) and Serrate (Ser) signal from the

flanking non-boundary cells, inducing proteolytic cleavages of Notch to generate a free Notch

intracellular domain (N^{ICD}) (Bray, 2016; Fortini, 2009). N^{ICD} translocates to the nucleus where it

binds transcriptional co-activators and DNA binding proteins to initiate target gene transcription,

including *wg* and *cut* (de Celis et al., 1996; Rulifson and Blair, 1995). The absence of Notch

results in reduced *wg* transcription and therefore reduced Wnt pathway activation (Rulifson and

Blair, 1995).

Both Notch and Wg signaling act to regulate common developmental processes such as

tissue patterning, fate specification and growth of different Drosophila appendages (Hing et al.,

1994). These two pathways share a number of common regulators which affect the activity of

their signaling outcome. In an *in vivo* RNAi screen to identify novel kinase and phosphatase

modulators of the Wg pathway, we found that the components of the Protein Phosphatase 4

(PP4) complex appeared to promote Wg signaling (Swarup et al., 2015). The serine threonine

phosphatase PP4 belongs to the Protein Phosphatase 2A (PP2A) group of phosphatases (Cohen

et al., 2005). Similar to what is found with PP2A, PP4 forms a heterotrimeric complex, which in

Drosophila consists of a catalytic subunit, Protein Phosphatase 4-19C (PP4-19C), and two

regulatory subunits called Protein Phosphatase 4 Regulatory subunit 2-related protein (PPP4R2r)

and PP4R3/Falafel (Flfl) (Cohen et al., 2005; Gingras et al., 2005).

PP4 is a highly conserved phosphatase seen across metazoans, and has been implicated in

a wide range of cellular processes, including chemotaxis in slime molds (Mendoza et al., 2007),

developmental signaling pathways such as Hedgehog (Jia et al., 2009), JNK (Huang and Xue,

2015; Zhou et al., 2002), Insulin-like growth factor (Mihindukulasuriya et al., 2004), as well as

TOR (Raught et al., 2001). The major functional role of PP4 is as a key regulator in cell cycle

progression and regulation of cell division (Helps et al., 1998; Huang et al., 2016). No previous

studies have implicated PP4 in Notch or Wnt/Wg signaling.

87   In this study we demonstrate that our previous observations of reduced Wg signaling due

88 to knockdown of PP4 components are due to effects on *wg* transcriptional regulation by Notch.

89 Using genetic interaction studies and expression of mutant Flfl, we determine that PP4 promotes

90 the activity of nuclear Notch. We further elucidated that the function of PP4 in promoting Notch

91 signaling was independent of its previously described role in cell cycle progression and

92 proliferation. Taken together we have identified a novel role for PP4 in promoting Notch

93 signaling and expression of *wg* during Drosophila development.

95 **Results**

97 *PP4 promotes Wg signaling in the Drosophila wing imaginal disc*

99   In a screen for modifiers of Wg signaling in the Drosophila wing imaginal disc, three

100 components of PP4 were found to reduce Wg target genes following their knockdown through

101 RNAi (Swarup et al., 2015). An involvement of PP4 in Wg signaling has not been previously

102 identified, so we were curious to determine mechanistically how PP4 may be involved in

103 regulating the output of the Wg pathway. In the developing wing imaginal disc the Wg target

104 genes *Sens* and *Dll* are expressed in a distinct nested pattern along the dorsoventral (D/V)

105 boundary (Fig. 1A, A'). We utilized *dpp-GAL4* expressed in a narrow band of cells along the

106 anterior/posterior (A/P) boundary (marked by GFP; Fig. 1B', B") to express RNAi constructs to

107 knockdown expression of the individual PP4 components. The knockdown of the catalytic

108 subunit *PP4-19C* caused a strong reduction in Sens and Dll (Fig. 1C, C'). Reduction of the PP4

109 targeting subunits *ppp4R2* or *flfl* via RNAi caused reduced Dll protein levels (Fig. 1E', G'), but

110 did not noticeably affect Sens (Fig. 1E, G).

111   To confirm that the reduction of PP4 components did not critically affect cell viability as

112 loss of PP4 has been shown to promote JNK-dependant cell death (Huang and Xue, 2015), discs

113 were stained for the apoptotic marker, cleaved caspase-3 (C. Casp-3). Compared to control cells

114 expressing GFP (Fig. 1B, B"), reduction of any individual PP4 component did not increase levels

115 of apoptosis within the *dpp* domain of the imaginal disc (Fig. 1D, F, H). Following Wg pathway

116 activation, the key effector protein Arm is stabilized at the highest concentration in two bands of

117 cells flanking the Wg-producing cells of the D/V boundary (Fig. 1I, I') (Peifer et al., 1991).

Expression of *flfl-RNAi* in the posterior domain of the wing imaginal disc, using *engrailed (en)-*

Gal4 (Fig. 1I, J), caused a reduction of stabilized Arm (Fig. 1J, arrow in J'). In subsequent

experiments, we utilized *flfl-RNAi* to reduce PP4 activity, as it has been previously confirmed as

a functional indicator of the entire complex (Sousa-Nunes et al., 2009). Together, these data

suggest PP4 is required for promoting Wg pathway activation.

PP4 promotes Wg signaling through Notch pathway activation

As the reduction of Wg target genes and Arm was apparent upon knockdown of PP4

components, we next wanted to look at the Wg ligand and its transcription. In third instar wing

imaginal discs, *wg* is transcribed, translated, and undergoes post-translational modification,

which can affect its stability, before being secreted to activate the Wg pathway in neighbouring

cells (Franch-Marro et al., 2008; Tanaka et al., 2002). Utilizing Wg antibodies, and the *wg*

transcriptional reporter, *wg-lacZ*, we could identify any defects in the ligand's transcription,

processing, or stability. In a wild type wing imaginal disc, Wg and *wg-lacZ* expression are

refined along the D/V boundary in a narrow band 2-3 cells wide (Fig. 2A, B). Expression of *flfl-*

RNAi in the posterior domain of the wing disc, using *hedgehog* (*hh*)-*Gal4*, resulted in a reduction

of both total Wg protein levels and transcription (Fig. 2E, F) suggesting PP4 is involved in

regulation of *wg* transcription.

wg transcription is controlled along the D/V boundary of the wing disc by Notch

signaling (Rulifson and Blair, 1995). We next wished to determine if PP4 regulation of *wg*

transcription is mediated through the involvement of the Notch signaling pathway. *cut*, another

Notch target gene (de Celis et al., 1996), is expressed in a similar pattern as Wg along the D/V

boundary (Fig. 2C). The reduction of *flfl* in the posterior domain of the wing disc via RNAi

resulted in a strong loss of Cut expression, indicating an overall reduction in Notch signaling

(Fig. 2G). Looking at the Notch ligand Delta (Dl) which is enriched in the cells adjacent to the

D/V boundary (Fig. 2D), it was apparent that *flfl-RNAi* expressed in the posterior domain of the

disc, resulted in reduced Dl and a failure of its refinement (Fig. 2H). We could not discern if this

effect on Dl is from upstream regulation of *Dl* expression, or on Notch activation itself, as the

refinement of Dl involves a cis/trans feedback mechanism with N for pattern refinement, through

lateral inhibition of each gene (Axelrod, 2010). Together, these results demonstrate that PP4

normally appears to influence Notch signaling to promote multiple pathway targets including *wg*.

PP4 promotes Notch signaling in the Notch signal receiving cells.

Having identified that Flfl, and by extension PP4, is involved in promoting Notch

signaling we sought to further elucidate how. During wing imaginal disc development Notch and

its ligands Dl and Serrate (Ser) undergo refinement through lateral inhibition resulting in high

levels of active Notch (N^{ICD}) being expressed along the D/V boundary and suppressed in the

flanking cells (Fig. 3D arrow head), which conversely have high levels of Dl (Fig. 2D) and Ser

(de Celis et al., 1996). This feed forward loop of lateral inhibition creates Notch signal-sending

cells (cells flanking the D/V boundary), and signal-receiving cells (D/V boundary cells) with

active Notch signaling (Axelrod, 2010).

To further analyze the role of the PP4 complex in the complementary ligand-expressing

and Notch-expressing cells, we used *C5-Gal4* and *wg-Gal4* respectively, to express *flfl-RNAi* in

the wing imaginal disc. *C5-Gal4* is expressed in N ligand expressing cells flanking the D/V

boundary (Fig. 3E), while *wg-Gal4* is expressed along the D/V boundary, in the active N signal

receiving cells (Fig. 3J). By looking at Wg and Notch target genes, we could determine in which

cells Flfl and by extension PP4 is working to affect the Notch pathway. *C5>flfl-RNAi* appeared

to have no affect on *Dll* expression (Fig. 3A, F), Wg (Fig. 3B, G), or Cut (Fig. 3C, H). The

enrichment of the N^{ICD} along the D/V boundary also appeared to be unaffected (Fig. 3D, I

arrowheads). *wg>flfl-RNAi* gave very contrasting results. Knockdown of *flfl* in the Notch

receptor expressing cells resulted in a complete loss of *Dll-lacZ (*Fig. 3K), strong reduction of

Wg (Fig. 3L), loss of Cut (Fig. 3M), and a failure of enrichment of N^{ICD} (Fig. 3N). Knockdown

of the other components of PP4 using *wg-Gal4*, but not *C5-gal4*, also caused loss of Wg

expression (Fig. S1A-D). Taken together, these results suggest that PP4 functions within the

Notch-expressing cell to promote full pathway activation and target gene expression.

PP4 functions within the nucleus to promote Notch signaling.

To further refine where PP4 functions within the Notch signal receiving cell, we utilized

mutant transgenes of Flfl and N in the adult Drosophila wing. During pupal wing

metamorphosis, the activation and refinement of Dl and N are required to refine adult vein

formation (Huppert et al., 1997). Notch signaling is also critical for the development of sensory

bristles along the adult wing margin (reviewed in Posakony, 1994). Any developmental defects

from expression of the various Flfl transgenes in adult wings could provide insight into PP4's

role in the Notch pathway. Using *MS1096-Gal4*, which is expressed across the entire developing

wing pouch, to ectopically express wild type Flfl or a cytoplasmic form, $Flfl^{\Delta 3NLS+2NES}$ (Flfl-

cyto), had no effect on the adult wing compared to wild type (Fig. 4A-C). Endogenous Flfl is a

predominantly nuclear protein and the wild type Flfl transgene has been shown to function

similarly (Sousa-Nunes et al., 2009). Knockdown of *flfl* via RNAi induced ectopic and thicker

veins in the adult wing (Fig. 4D), a hallmark of reduced Notch activity (Huppert et al., 1997).

This phenotype could be suppressed by reintroduction of the wild type *flfl* transgene (Fig. 4E),

but not with expression of *flfl-cyto* (Fig. 4F). This suggests that Flfl functions within the nucleus,

rather than the cytoplasm, to promote Notch.

To confirm the hypothesis that Flfl likely acts in nuclear Notch signaling, we expressed a

construct encoding the intracellular domain of Notch that localizes to the nucleus (N^{nucl})(Rebay

et al., 1993). This activated nuclear Notch suppressed wing vein formation and the formation of

sensory bristles (Fig. 4G, G') compared to wild type (Fig. 4A). Notch-dependent activation of *wg*

expression is essential for Wg signaling to induce expression of proneural genes like *Sens* for the

specification of sensory organ precursor (SOP) cells (Nolo et al., 2000). SOPs then divide and

differentiate giving rise to the sensory bristles in the adult fly via Notch signaling (Guo et al.,

1996; Hartenstein and Posakony, 1990). Although ectopic Notch signaling increases the number

of SOPs in the wing disc via Wg, they do not differentiate correctly resulting in double socket

cells, and loss of bristle cells (Guo et al., 1996). Knockdown of *flfl* in the N^{nucl} expressing cells

was able to partially recover vein loss, but did not significantly rescue the bristle defect (Fig. 4H

arrow, H'). Conversely, the expression of Flfl or Flfl-cyto had no affect on the N^{nucl} phenotype

(Fig. 4I-J').We interpret the inability to specifically rescue the bristle defect as being due to the

combination of the strength of N^{nucl} as well as an incomplete knockdown of *flfl* via RNAi. To

confirm that this phenotype could be rescued in our assay, we tested whether loss of *wg* could

rescue the effect since up-regulated target genes expression causes the N^{nucl} phenotype.

Expression of a weak *wg-RNAi* transgene was able to induce sporadic sensory bristle loss (Fig.

4K, K' arrow), due to reduced SOPs (Nolo et al., 2000; Parker et al., 2002). When combined

with N^{nucl}, *wg-RNAi* can partially suppress the overactive Notch phenotype of inhibited bristle

formation (Fig. 4L, L' arrowhead).

Flfl is required for proliferation and maintenance of overall tissue size independent of Notch

signaling.

Adult flies with reduced *flfl* expression displayed smaller wing blades compared to

control flies (Fig. 4A, D). This was expected given PP4's known role in cycle progression and

growth (Helps et al., 1998; Huang et al., 2016; Martin-Granados et al., 2008; Zhuang et al.,

2014). Notch has been implicated in cell proliferation in the wing imaginal disc, but a direct

mechanism for its involvement is not fully understood (Baonza and Garcia-Bellido, 2000;

Giraldez and Cohen, 2003; Go et al., 1998). We quantified the area of the adult wings of the

different genotypes, in order to determine if PP4/Flfl's role in growth is mediated through Notch

signaling. Overexpression of Flfl and Flfl-cyto had no significant effect on wing size compared

to wild type (Fig. 5A, box plot a,b,c). Knockdown of *flfl* resulted in a ~28% reduction in wing

size, and could be fully rescued by the wild type *flfl* transgene (Fig. 5A, box plot d,e). Flfl-cyto

was able to slightly rescue the growth defect from *flfl-RNAi*, but not to a significant level (Fig.

5A, box plot f). The ability to rescue partially may be due to PP4's role in mitotic progression

after nuclear envelope break down at prometaphase, allowing for Flfl-cyto to perform its

function at this step in mitosis (Huang et al., 2016). Wings expressing N^{nucl} did not have a

significantly smaller area than wild type wings, and were unable to rescue the growth defect

from *flfl-RNAi* (Fig. 5A, box plot g,h). As N^{nucl} is unable to rescue the growth defect from

reduced Flfl levels, it indicates that PP4/Flfl's role in regulating growth is likely independent to

its function in propagating Notch signaling. An alternative interpretation could be that Flfl acts

downstream of N^{nucl} or that Flfl is required for proper Notch function in the proliferation.

To determine if the growth defects from loss of PP4 were due to decreased cell

proliferation or overall cell size, we looked at the developing wing imaginal disc. We utilized *en-*

Gal4 driving GFP to mark the posterior portion of the wing disc, representing roughly 50% of

the overall tissue (Fig. 5B). We compared the size of the GFP-positive region to the control

anterior side of the disc in different genotypes to determine the effects that changes in the levels

of Flfl have on tissue growth. In control discs expressing *UAS-lacZ*, the ratio between the

internal control area of the anterior, to the GFP positive posterior was equal (Fig. 5B, E).

Overexpression of Flfl in the posterior domain did not affect the posterior/anterior area ratio

(Fig. 5C, E). However, reduction of *flfl* via RNAi resulted in a significant decrease in size of the

posterior domain (Fig. 5D, E).

We simultaneously looked at the number of mitotic cells in these genotypes using a

phospho-Histone H3 (Ser10) (PH3) antibody. There was no significant difference between our

control discs and those overexpressing of Flfl (Fig. 5B', C', F). Surprisingly, we found that *flfl-*

RNAi tissue had significantly elevated levels of PH3 positive cells (Fig. 5D', F). This result was

perplexing considering the decreased tissue size, yet apparent increase in proliferation rate. As

PH3 marks condensed chromatin prior to chromosomal segregation in cells along the G2/M

transition (Hendzel et al., 1997), it is possible that *flfl-RNAi* cells were arrested during early

mitosis, and had not undergone mitotic exit. Previously, Sousa-Nunes et al. (2009) identified a

similar effect. Drosophila *flfl/+* brains exhibited much lower rates of Bromodeoxyuridine (BrdU)

incorporation and reduced proliferation, yet exhibited elevated levels of PH3 positive cells,

demonstrating *flfl* is important for mitotic progression. Similar results have been found in

multiple cases (Huang et al., 2016; Martin-Granados et al., 2008). Our results suggest a similar

function for PP4/Flfl in the wing disc, where PP4 is critical for cell cycle progression and mitosis

affecting proliferation rates and overall tissue size, yet this function is independent of its role in

promoting Notch signaling.

aPKC is not involved with PP4 and Notch signaling in the Drosophila wing imaginal disc

We examined whether PP4 mechanistically interacted with other known modulators of

the Notch pathway. One well-characterized regulator is atypical protein kinase C (aPKC). aPKC

is a protein kinase widely studied for its role in developmental processes, including asymmetric

cell division (ACD). The regulators of ACD (including aPKC, Bazooka, and Crumbs) act

upstream of Notch signaling and determine the identity of the Notch signal sending and signal

receiving cells (reviewed in Knoblich, 2008). aPKC also promotes the Notch pathway by

inhibiting Numb-mediated endocytosis of the Notch pathway components (Frise et al., 1996;

Smith et al., 2007; Wang et al., 2006). Previous studies have shown a role for PP4/Flfl in the

localization of the Miranda complex to promote neuroblast ACD in Drosophila by acting

downstream or parallel to aPKC (Sousa-Nunes et al., 2009). Since such a role has not been

identified in a proliferating epithelium like the wing disc we sought to investigate if this

mechanism was conserved during the development of the wing disc.

Using *MS1096-Gal4* to express *aPKC-RNAi* in the wing disc resulted in small, crumpled

adult wings, with only sporadic sensory bristles (Fig. S2B), indicating disrupted patterning and

growth. Overexpression of an aPKC transgene did not induce any visible phenotype (Fig. S2C),

and was able to rescue the *aPKC-RNAi* phenotype, confirming the phenotype seen from the

RNAi was directly a result of loss of *aPKC* (Fig. S2D). The knockdown of *flfl* via RNAi or over

expression of the wild type *flfl* transgene had no affect on the *aPKC-RNAi* phenotype (Fig. S2E,

F). Importantly, expression of N^{nucl} was unable to rescue the crumpled wing from *aPKC*

knockdown, indicating aPKC is most likely not a direct upstream regulator of Notch in the wing

imaginal disc (Fig. S2G). In addition to this, overexpression of *aPKC* had no effect on the *flfl-*

RNAi vein thickening compared to *flfl-RNAi* alone (Fig. 4D, S2H). Although aPKC is involved

with PP4 and Notch signaling in ACD (Sousa-Nunes et al., 2009; Zhang et al., 2016), it does not

appear to be directly involved in Notch signaling and pattering in the developing wing imaginal

disc.

Discussion

An *in vivo* RNAi screen initially identified three components of the PP4 enzyme complex

as modulators of endogenous Wg signaling during wing development (Swarup et al., 2015). The

PP4 complex in *Drosophila melanogaster* has been implicated in many signaling pathways,

cellular functions and developmental processes, yet its role in regulating Wg signaling was

previously uncharacterized. In this study, we revealed that the effect on Wg pathway was at the

level of expression of the Wg ligand, by promotion of Notch signaling. Notch signaling regulates

the precise expression of Wg in the cells of the D/V boundary of the wing imaginal disc. This

expression is required for specification of the wing margin and bristle structures (Couso et al.,

1994; de Celis et al., 1996; Neumann and Cohen, 1996).

A partial knockdown of Flfl, PP4-19C and PPP4R2 by RNAi in cells along the A/P
boundary of the wing imaginal discs was able to effectively reduce Wg target genes, yet did not
induce elevated levels of JNK-mediated cell death as previously reported (Huang and Xue,
2015). Cell death was inducible upon stronger expression of the RNAi using *act-Gal4* with heat-
shock inducible flip-out clones (data not shown). The knockdown of Flfl was further found to
reduce expression of the Wg ligand as well as other Notch pathway target genes, implicating PP4
in the Notch pathway. Reduction of PP4 proteins in the D/V boundary all resulted in reduced Wg
expression, while their knockdown in neighbouring Dl- and Ser-expressing cells had no effect.
This result highlights that the PP4 complex acts in boundary cells to regulate Notch-dependent
gene expression.

Previously PP4 has been indirectly associated with Notch signaling in Drosophila for its
involvement in asymmetrical cell division (ACD) of the developing neuroblasts (Sousa-Nunes et
al., 2009; Zhang et al., 2016). While PP4 acts in concert with aPKC and Notch signaling to drive
proper ACD, we were unable to identify a role for aPKC in Notch signaling in the epithelial cells
of the wing imaginal disc. However, further genetic interaction experiments also identified that
PP4's involvement in Notch signaling in the wing imaginal disc appears to be independent of
PP4's role in cell cycle progression and tissue growth. These results demonstrated that although
both PP4 and Notch are required for cell proliferation (Giraldez and Cohen, 2003; Go et al.,
1998; Huang et al., 2016), in the wing imaginal disc it is likely not through the same mechanism.

Subsequent genetic interaction studies revealed that most likely Flfl acts to promote
Notch through its role in the nucleus. A cytoplasmic form of Flfl could not recue the phenotypes
generated by *flfl-RNAi* in the adult wing, while expression of a wildtype transgene could. The
required function of Flfl in the nucleus was further bolstered by the fact that the wing phenotype
induced by activated nuclear N was partially suppressed by *flfl-RNAi*. As the N^{ICD} enters the
nucleus and binds to Suppressor of Hairless and Mastermind to initiate target gene transcription,
a multitude of cofactors must be recruited, while others must be removed, from the
transcriptional initiation site (reviewed in Bray, 2016). This includes the inhibition of histone
deacetylase (HDAC) co-repressor complexes (Kao et al., 1998). PP4 has been previously
identified to dephosphorylate and inhibit HDAC activity, while its depletion stimulates HDACs
(Zhang et al., 2005). Taken together a possible mechanism for PP4 to promote Notch signaling is
through the dephosphorylation of HDACs. This could allow for increased chromatin

remodelling, which is needed for the binding of other transcriptional co-factors to ensure full

transcriptional initiation of target genes (reviewed Bray, 2016). This is just one possibility, as

PP4 may be responsible for the dephosphorylation and modulation of any number of components

that cooperate with transcription factors, or regulate the activity of N^{ICD} leading to appropriate

target gene expression. Future studies will hopefully address the exact mechanism PP4 plays in

promoting nuclear Notch signaling for full expression of target genes like *wg*.

Methods and Materials

Fly Strains and Crosses

Fly strains and crosses were raised on standard medium at 25°C unless stated otherwise. *w^{1118}*

was utilized as wild type. In assays examining genetic interactions between two UAS-driven

transgenes, control crosses were performed with *UAS-lacZ* and *UAS-gfp* to eliminate effects

caused by the titration of Gal4. The following fly strains were used in this study: *UAS-GFP,*

UAS-lacZ/TM6B, *UAS-dicer, dpp-Gal4*, *Dll-lacZ*, *MS1096-Gal4*, *UAS-aPKC-GFP* (obtained

from the Bloomington Drosophila Stock Center)*, UAS-GFP, C5-Gal4 (*Hugo Bellen)*, UAS-*

flflΔ3NLS+2NES (flfl-cyto) (Gregory Somers), *UAS-flfl (*Zoltan Lipinszki)*, wg-lacZ/CyO and*

UAS-N[nucl] (Spyros Artavanis-Tsakonas), *UAS-flfl-RNAi* (VDRC 24143,103793), *UAS-PP4-*

19C-RNAi (VDRC 25317, 103317, 43250), *UAS-PPP4R2R-RNAi* (VDRC 25445, 105399), *UAS-*

*aPKC-RNAi (*VDRC 2907,105624*)* [obtained from the Vienna Drosophila Resource Center

(Dietzl et al., 2007)], *hh-Gal4/TM6B*, *UAS-wg-RNAi, wg-Gal4 (*ND382*)* and *en-Gal4,UAS-GFP*

(gifts from Konrad Basler, Institute of Molecular Life Sciences University of Zurich,

Switzerland).

Crosses involving *C5-Gal4* were performed at 29°C to induce maximal Gal4 expression in the

developing wing disc.

Immunofluorescence of wing imaginal discs

Third-instar larvae were dissected in phosphate-buffered saline (PBS). Wing imaginal discs were

fixed in 4% paraformaldehyde at room temperature for 20 minutes and then washed 3 times for 5

minutes in PBS. Discs were blocked [2% BSA diluted in PBS 0.1% Triton X-100 (PBST)] for 45

minutes at room temperature, followed by incubation with primary antibodies (diluted in block)

overnight at 4°C. Tissue was then washed 3 times in PBST and incubated with secondary

antibodies (diluted in block) at room temperature for 1.5 hours. Tissue was then washed a final

three times in PBST and mounted in a 70% glycerol solution. The following primary antibodies

and dilutions were used: mouse anti-β-galactosidase (1:2000 Promega), mouse anti-Wg (1:100

DSHB), mouse anti-Cut (1:50 DSHB), mouse anti-N^{ICD} (1:50 DSHB), mouse anti-Delta (1:50

DSHB), mouse anti-Arm (1:50 DSHB), rabbit anti-cleaved Caspase 3 (1:100 Cell Signaling),

rabbit anti-PH3 (1:200 Cell Signaling), guinea pig anti-Sens (1:500, a gift from Hugo Bellen,

Dept. of Molecular and Human Genetics, Baylor College of Medicine, USA), mouse anti-Dll

(1:300, a gift from Ian Duncan, Dept. of Biology, Washington University in St. Louis, USA).

Secondary antibodies (Jackson ImmunoResearch) were used at a 1:200 dilution.

Adult wing mounting

Adult wings were dissected in 95% ethanol followed by mounting in Aquatex (EMD Chemicals

Inc.) A minimum of 8 wings were mounted per genotype for analysis.

Imaging, Analysis and Quantification

Fluorescent images were taken with a Nikon A1R laser scanning confocal microscope and

processed using Adobe Photoshop CS6. Adult wings were imaged with an Axioplan 2

microscope. Adult wing and wing disc areas were quantified using ImageJ software. PH3 cell

counts were performed using the ImageJ plugin, "Cell Counter". To compare PH3 positive cell

counts per genotype, counts were converted as a ratio of PH3 cells/area, then analysed as

experimental condition over control tissue. Significance between groups was assessed by one-

way analysis of variance (ANOVA), and *p<0.01* was considered significant unless stated

otherwise.

Hall, Pradhan-Sundd et al.

Acknowledgements:

We are grateful to Jessica Blaquiere for technical assistance in the early stage of this study. We

also thank Zoltan Lipinszki, Gregory Somers, Spyros Artavanis-Tsakonas, Rita Sousa-Nunes,

Hugo Bellen, Juergen Knoblich, Konrad Basler, James Skeath, the Bloomington *Drosophila*

Stock Centre, the Vienna *Drosophila* RNAi Centre and the Developmental Studies Hybridoma

Bank for providing fly strains and antibodies.

Competing Interests

The authors declare that they have no competing interests.

Author contributions

E.T.H., T.P.-S. and E.M.V. designed the experiments, E.T.H., T.P.-S., F.S. and E.M.V.

performed the experiments, and E.T.H., T.P.-S. and E.M.V analyzed the data and wrote the

manuscript.

Funding

This work was supported by an operating grant from the Canadian Institutes of Health Research

(CIHR) [grant number FRN 133522].

References

Axelrod, J. D. (2010). Delivering the Lateral Inhibition Punchline: It's All About the Timing.

Sci. Signal. **3**, pe38-pe38.

Baonza, A. and Garcia-Bellido, A. (2000). Notch signaling directly controls cell proliferation

in the Drosophila wing disc. *Proc. Natl. Acad. Sci.* **97**, 2609–2614.

Bray, S. J. (2016). Notch signalling in context. *Nat. Rev. Mol. Cell Biol.* **17**, 722–735.

Clevers, H. and Nusse, R. (2012). Wnt/β-Catenin Signaling and Disease. *Cell* **149**, 1192–1205.

Cohen, P. T. W., Philp, A. and Vázquez-Martin, C. (2005). Protein phosphatase 4 – from

obscurity to vital functions. *FEBS Lett.* **579**, 3278–3286.

Collu, G. M., Hidalgo-Sastre, A. and Brennan, K. (2014). Wnt-Notch signalling crosstalk in
development and disease. *Cell. Mol. Life Sci.* **71**, 3553–67.

Couso, J. P., Bishop, S. A. and Martinez Arias, A. (1994). The wingless signalling pathway
and the patterning of the wing margin in Drosophila. *Development* **120**, 621–36.

de Celis, J. F., Garcia-Bellido, A. and Bray, S. J. (1996). Activation and function of Notch at
the dorsal-ventral boundary of the wing imaginal disc. *Development* **122**,.

Dietzl, G., Chen, D., Schnorrer, F., Su, K.-C., Barinova, Y., Fellner, M., Gasser, B., Kinsey,
K., Oppel, S., Scheiblauer, S., et al. (2007). A genome-wide transgenic RNAi library for
conditional gene inactivation in Drosophila. *Nature* **448**, 151–156.

Fortini, M. E. (2009). Notch Signaling: The Core Pathway and Its Posttranslational Regulation.
Dev. Cell **16**, 633–647.

Franch-Marro, X., Wendler, F., Griffith, J., Maurice, M. M. and Vincent, J.-P. (2008). In
vivo role of lipid adducts on Wingless. *J. Cell Sci.* **121**, 1587–1592.

Frise, E., Knoblich, J. A., Younger-Shepherd, S., Jan, L. Y. and Jan, Y. N. (1996). The
Drosophila Numb protein inhibits signaling of the Notch receptor during cell-cell
interaction in sensory organ lineage. *Proc. Natl. Acad. Sci. U. S. A.* **93**, 11925–32.

Gingras, A.-C., Caballero, M., Zarske, M., Sanchez, A., Hazbun, T. R., Fields, S.,
Sonenberg, N., Hafen, E., Raught, B. and Aebersold, R. (2005). A novel, evolutionarily
conserved protein phosphatase complex involved in cisplatin sensitivity. *Mol. Cell.*
Proteomics **4**, 1725–40.

Giraldez, A. J. and Cohen, S. M. (2003). Wingless and Notch signaling provide cell survival
cues and control cell proliferation during wing development. *Development* **130**, 6533–6543.

Go, M. J., Eastman, D. S. and Artavanis-Tsakonas, S. (1998). Cell proliferation control by
Notch signaling in Drosophila development. *Development* **125**, 2031–40.

Guo, M., Jan, L. Y. and Jan, Y. N. (1996). Control of daughter cell fates during asymmetric
division: interaction of Numb and Notch. *Neuron* **17**, 27–41.

Hartenstein, V. and Posakony, J. W. (1990). A dual function of the Notch gene in Drosophila
sensillum development. *Dev. Biol.* **142**, 13–30.

Helps, N. R., Brewis, N. D., Lineruth, K., Davis, T., Kaiser, K. and Cohen, P. T. (1998). Protein phosphatase 4 is an essential enzyme required for organisation of microtubules at centrosomes in Drosophila embryos. *J. Cell Sci.* 1331–40.

Hendzel, M. J., Wei, Y., Mancini, M. A., Van Hooser, A., Ranalli, T., Brinkley, B. R., Bazett-Jones, D. P. and Allis, C. D. (1997). Mitosis-specific phosphorylation of histone H3 initiates primarily within pericentromeric heterochromatin during G2 and spreads in an ordered fashion coincident with mitotic chromosome condensation. *Chromosoma* **106**, 348–60.

Hing, H. K., Sun, X. and Artavanis-Tsakonas, S. (1994). Modulation of wingless signaling by Notch in Drosophila. *Mech. Dev.* **47**, 261–268.

Huang, J. and Xue, L. (2015). Loss of flfl Triggers JNK-Dependent Cell Death in Drosophila. *Biomed Res. Int.* **2015**, 623573.

Huang, X., Liu, J., Shen, T., Meng, X., Dou, L., Lin, Y. and Li, J. (2016). Protein phosphatase 4 plays dual roles during cell proliferation. *Cell Prolif.* **49**, 219–235.

Huppert, S. S., Jacobsen, T. L. and Muskavitch, M. A. (1997). Feedback regulation is central to Delta-Notch signalling required for Drosophila wing vein morphogenesis. *Development* **124**, 3283–91.

Itasaki, N. and Hoppler, S. (2010). Crosstalk between Wnt and bone morphogenic protein signaling: a turbulent relationship. *Dev. Dyn.* **239**, 16–33.

Jia, H., Liu, Y., Yan, W. and Jia, J. (2009). PP4 and PP2A regulate Hedgehog signaling by controlling Smo and Ci phosphorylation. *Development* **136**, 307–316.

Kao, H. Y., Ordentlich, P., Koyano-Nakagawa, N., Tang, Z., Downes, M., Kintner, C. R., Evans, R. M. and Kadesch, T. (1998). A histone deacetylase corepressor complex regulates the Notch signal transduction pathway. *Genes Dev.* **12**, 2269–77.

Kim, M. and Jho, E.-H. (2014). Cross-talk between Wnt/β-catenin and Hippo signaling pathways: a brief review. *BMB Rep.* **47**, 540–5.

Knoblich, J. A. (2008). Mechanisms of asymmetric stem cell division. *Cell* **132**, 583–97.

Martin-Granados, C., Philp, A., Oxenham, S. K., Prescott, A. R. and Cohen, P. T. W.

479 (2008). Depletion of protein phosphatase 4 in human cells reveals essential roles in
480 centrosome maturation, cell migration and the regulation of Rho GTPases. *Int. J. Biochem.*
481 *Cell Biol.* **40**, 2315–2332.

482 **Mendoza, M. C., Booth, E. O., Shaulsky, G. and Firtel, R. A.** (2007). MEK1 and Protein
483 Phosphatase 4 Coordinate Dictyostelium Development and Chemotaxis. *Mol. Cell. Biol.* **27**,
484 3817–3827.

485 **Mihindukulasuriya, K. A., Zhou, G., Qin, J. and Tan, T.-H.** (2004). Protein Phosphatase 4
486 Interacts with and Down-regulates Insulin Receptor Substrate 4 following Tumor Necrosis
487 Factor- Stimulation. *J. Biol. Chem.* **279**, 46588–46594.

488 **Neumann, C. J. and Cohen, S. M.** (1996). A hierarchy of cross-regulation involving Notch,
489 wingless, vestigial and cut organizes the dorsal/ventral axis of the Drosophila wing.
490 *Development* **122**, 3477–85.

491 **Neumann, C. J. and Cohen, S. M.** (1997). Long-range action of Wingless organizes the dorsal-
492 ventral axis of the Drosophila wing. *Development* **124**, 871–80.

493 **Nolo, R., Abbott, L. A. and Bellen, H. J.** (2000). Senseless, a Zn finger transcription factor, is
494 necessary and sufficient for sensory organ development in Drosophila. *Cell* **102**, 349–62.

495 **Parker, D. S., Jemison, J. and Cadigan, K. M.** (2002). Pygopus, a nuclear PHD-finger protein
496 required for Wingless signaling in Drosophila. *Development* **129**,.

497 **Peifer, M., Rauskolb, C., Williams, M., Riggleman, B. and Wieschaus, E.** (1991). The
498 segment polarity gene armadillo interacts with the wingless signaling pathway in both
499 embryonic and adult pattern formation. *Development* **111**, 1029–43.

500 **Posakony, J. W.** (1994). Nature versus nurture: asymmetric cell divisions in Drosophila bristle
501 development. *Cell* **76**, 415–8.

502 **Raught, B., Gingras, A.-C. and Sonenberg, N.** (2001). The target of rapamycin (TOR)
503 proteins. *Proc. Natl. Acad. Sci.* **98**, 7037–7044.

504 **Rebay, I., Fehon, R. G. and Artavanis-Tsakonas, S.** (1993). Specific truncations of Drosophila
505 Notch define dominant activated and dominant negative forms of the receptor. *Cell* **74**,
506 319–29.

Rulifson, E. J. and Blair, S. S. (1995). Notch regulates wingless expression and is not required

for reception of the paracrine wingless signal during wing margin neurogenesis in

Drosophila. *Development* **121**, 2813–24.

Smith, C. A., Lau, K. M., Rahmani, Z., Dho, S. E., Brothers, G., She, Y. M., Berry, D. M.,

Bonneil, E., Thibault, P., Schweisguth, F., et al. (2007). aPKC-mediated phosphorylation

regulates asymmetric membrane localization of the cell fate determinant Numb. *EMBO J.*

26, 468–80.

Sousa-Nunes, R., Chia, W. and Somers, W. G. (2009). Protein Phosphatase 4 mediates

localization of the Miranda complex during Drosophila neuroblast asymmetric divisions.

Genes Dev. **23**, 359–372.

Swarup, S. and Verheyen, E. M. (2012). Wnt/Wingless signaling in Drosophila. *Cold Spring*

Harb. Perspect. Biol. **4**,.

Swarup, S., Pradhan-Sundd, T. and Verheyen, E. M. (2015). Genome-wide identification of

phospho-regulators of Wnt signaling in Drosophila. *Development* **142**, 1502–1515.

Tanaka, K., Kitagawa, Y. and Kadowaki, T. (2002). Drosophila Segment Polarity Gene

Product Porcupine Stimulates the Posttranslational N-Glycosylation of Wingless in the

Endoplasmic Reticulum. *J. Biol. Chem.* **277**, 12816–12823.

Wang, H., Somers, G. W., Bashirullah, A., Heberlein, U., Yu, F. and Chia, W. (2006).

Aurora-A acts as a tumor suppressor and regulates self-renewal of Drosophila neuroblasts.

Genes Dev. **20**, 3453–3463.

Williams, J. A., Paddock, S. W. and Carroll, S. B. (1993). Pattern formation in a secondary

field: a hierarchy of regulatory genes subdivides the developing Drosophila wing disc into

discrete subregions. *Development* **117**, 571–84.

Zecca, M., Basler, K. and Struhl, G. (1996). Direct and long-range action of a wingless

morphogen gradient. *Cell* **87**, 833–44.

Zeller, E., Hammer, K., Kirschnick, M. and Braeuning, A. (2013). Mechanisms of RAS/β-

catenin interactions. *Arch. Toxicol.* **87**, 611–32.

Zhang, X., Ozawa, Y., Lee, H., Wen, Y.-D., Tan, T.-H., Wadzinski, B. E. and Seto, E.

(2005). Histone deacetylase 3 (HDAC3) activity is regulated by interaction with protein

serine/threonine phosphatase 4. *Genes Dev.* **19**, 827–839.

Zhang, F., Huang, Z.-X., Bao, H., Cong, F., Wang, H., Chai, P. C., Xi, Y., Ge, W., Somers,
W. G., Yang, Y., et al. (2016). Phosphotyrosyl phosphatase activator facilitates localization
of Miranda through dephosphorylation in dividing neuroblasts. *Development* **143**, 35–44.

Zhou, G., Mihindukulasuriya, K. A., MacCorkle-Chosnek, R. A., Van Hooser, A., Hu, M.
C.-T., Brinkley, B. R. and Tan, T.-H. (2002). Protein Phosphatase 4 Is Involved in Tumor
Necrosis Factor- -induced Activation of c-Jun N-terminal Kinase. *J. Biol. Chem.* **277**, 6391–
6398.

Zhuang, X., Semenova, E., Maric, D. and Craigie, R. (2014). Dephosphorylation of Barrier-
to-autointegration Factor by Protein Phosphatase 4 and Its Role in Cell Mitosis. *J. Biol.*
Chem. **289**, 1119–1127.

Figure Legends:

Figure 1. Reduction of PP4 subunits inhibits Wg pathway activation without inducing cell death. (A-B'') Wild-type expression pattern of Wg target genes Sens (A), and Dll (A'), as well as cleaved caspase 3 (B) with normal expression pattern of *dpp-Gal4* (B', B'') along the anterior/posterior boundary in the developing wing disc. (C-D) The knockdown of PP4-19C with RNAi along the anterior/posterior boundary causes a loss of Sens (C, arrow head) and Dll (C', arrow head), but does not elevate C. Caspase 3 activity (D). (E-F) Knockdown of PPP4R2 was able to effectively reduce Dll (E', arrow head), but did not affect Sens (E) or C. Casp3 (F). (G-H) Flfl knockdown did not affect Sens (G) or C. Casp3 levels (H), but did reduces Dll (G', arrow head). Normal expression pattern of *en-Gal4* (I) in the posterior domain of the wing disc, shown with Arm (I, I'). (J-J') The knockdown of Flfl in the posterior domain (shown with GFP) induced a marked reduction in stabilized Arm (J'). Scale bar: 50 µm.

Figure 2. PP4 promotes Wg signaling through Notch pathway activation. (A-D) Wild-type

pattern of Wg protein (A), *wg* transcription (B), Cut protein (C), and Dl protein (D) in the

developing wing disc. (E-H) Using *hh-Gal4*, expressed in the posterior domain of the wing disc

(right of the dotted line), to express *flfl-RNAi*, caused a reduction in total Wg protein levels (E),

as well as *wg* transcription (F), and induced a loss of Cut (G), and Dl (H). Scale bar: 50 μm.

Figure 3. PP4 promotes Notch signaling in the Notch-signal receiving cells. (A-D) Wild-type

expression pattern of *Dll-lacZ* (A), Wg (B), Cut (C), and N^{ICD} (D) in the developing wing disc.

N^{ICD} is enriched along the dorsal/ventral (D/V) boundary (D, arrowhead) and suppressed in the

adjacent cells. (E) Expression pattern of *C5-Gal4* driving GFP in the D/V boundary flanking

cells of the wing pouch. (F-I) The knockdown of *flfl* with RNAi in the D/V boundary flanking

cells does not affect *Dll-lacZ* (F), Wg (G), Cut (H), or N^{ICD} (I). (J) Expression pattern of *wg-Gal4*

driving GFP along the D/V boundary. (K-N) Knockdown of *flfl* in the D/V boundary cells causes

a loss of *Dll-lacZ* (K), strong reduction of Wg (L), loss of Cut (M), and a failure of N^{ICD}

enrichment along the D/V boundary (N, arrow). Scale bar: 50 μm.

Figure 4. PP4 likely functions within the nucleus to promote Notch. (A-C) Adult wild-type

wing and margin (A, inset). Over expression of Flfl (B), or Flfl-cyto(C) throughout the entire

wing does not induce any noticeable phenotype. (D-E) Knockdown of Flfl induces ectopic veins

and thickening of veins, as well as a reduced wing size (D). This effect can be primarily rescued

by reintroduction of a full length Flfl transgene (E), but not by Flfl-cyto (F). (G-L') Over-

expression of N^{nucl} induces a loss of wing veins (G) and wing margin bristles (G'). Knockdown

of *flfl* induced a mild rescue the N^{nucl} loss of vein phenotype, and still reduces the overall wing

size (H, H'). The over expression of Flfl (I, I'), or Flfl-cyto (J, J') did not disrupt the N^{nucl}

phenotype. (K-L') Expression of *wg-RNAi* in the wing induced sporadic loss of margin bristles

(K, K' arrow). N^{nucl} with *wg-RNAi* is able to maintain several margin bristles (L, L' arrowhead).

Figure 5. Flfl is required for proliferation and overall tissue size independent of Notch

signaling. (A) Box plots representing total wing area of genotypes shown in Fig. 4 (n = 8-13).

Over expression of Flfl (b) or Flfl-cyto (c) did not affect wing size compared to wild type (a).

flfl-RNAi caused a significant reduction in wing size (d). The *flfl-RNAi* size defect could be fully

rescued by reintroduction of full length Flfl (e), but no effect was seen with Flfl-cyto (f). Wings

expressing N^{nucl} (g) are slightly smaller than wild type wings. N^{nucl} and *flfl-RNAi* wings (h) have

a significant size reduction compared to wild type (a), equivalent to that of *flfl-RNAi* alone (d).

Data are presented as box plot 25-75 percentile, whiskers 10-90 percentile, (—) median, (+)

mean, (•) outliers, letters above representing significance from corresponding genotypes,

(*p<0.01*) generated from one-way ANOVA. (B-F) The normal expression pattern of *en-Gal4*

marked by GFP (B) in the posterior domain of the developing wing disc, shown with mitotic cell

marker PH3 (B') and represented as a ratio of posterior domain vs. the anterior control (n = 8)(E,

F). Over expression of Flfl in the posterior domain had no effect on area (C, E) or proliferation

rate (n = 7)(C' F). The knockdown of Flfl with RNAi in the posterior domain induced a

significant reduction in area (D, E), and exhibited a significantly higher number of PH3

positively marked cells (n=9)(D' F). Data are presented as means ± s.d., (*) *p<0.05*, (**) *p<0.01*

generated from one-way ANOVA. Scale bar: 50 μm.

Hall, Pradhan-Sundd et al.

Hall, Pradhan-Sundd *et al.* - Supplemental Information

Supplemental Figure 1. PP4 subunits promote Notch signaling in the Notch receiving cells.

(A, B) *C5-Gal4* expressing *Pp4-19C-RNAi* or *PPP4R2-RNAi* along the D/V boundary flanking

cells of the wing pouch did not affect Wg protein levels or patterning. (C, D) *wg-Gal4* expression

of *Pp4-19C-RNAi* or *PPP4R2-RNAi* resulted in strong reduction of Wg in the wing disc. Scale

bar: 50 μm.

Supplemental Figure 2. aPKC does not promote Notch signaling in the Drosophila wing.(A)

Adult wild-type wing. (B) Knockdown of aPKC with RNAi throughout the entire wing induces

small crumpled, blistered, and malformed wings. (C-D) Over expression of an aPKC transgene

induces no visible phenotype (C), but can fully rescue the effects of *aPKC-RNAi* (D). (E-G) The

knockdown of Flfl (E), or its over expression (F) had no affect on the *aPKC-RNAi* phenotype.

N^{nucl} was also unable to rescue the *aPKC-RNAi* phenotype (G). (H) Over expression aPKC does

not affect the *flfl-RNAi* phenotype. Scale bar: 300 μm.