Adaptive pathways of coral populations on the Great Barrier Reef

Mikhail V. Matz[1*], Eric A. Treml[2], Galina V. Aglyamova[1], Madeleine J. H. van Oppen[2,3] and Line K. Bay[3]

[1] Department of Integrative Biology, University of Texas at Austin, 205 W 24th St. C0990, Austin, Texas 78712, USA
[2] School of BioSciences, University of Melbourne, Victoria 3010 Australia
[3] Australian Institute of Marine Science, QLD, Australia

* Author for correspondence, matz@utexas.edu

Abstract

There is an urgent need to understand factors influencing the potential of reef-building corals to adapt to future thermal regimes. Here we infer demographic parameters in five populations of the common coral *Acropora millepora* and use them to model future adaptation of the species on the Great Barrier Reef (GBR). Genetic analysis of samples collected 2002-2009 revealed that the loss of about half the coral cover in the preceding three decades did not yet result in detectable loss of genetic diversity or changes in gene flow. Both genetic and biophysical models indicated the prevalence of southward migration along the GBR, which would facilitate the spread of heat-tolerant alleles to higher latitudes as climate warms. Using a newly developed metapopulation adaptation model we find that standing genetic variation could be sufficient to fuel rapid adaptation of *A. millepora* to warming for the next 100-200 years. However, thermal anomalies such as ENSO cycles will drive increasingly severe coral mortality episodes. New mutations are unlikely to sustain genetic variation in the face of such strong selection, and so after the initial rapid adaptive response coral populations will go extinct, beginning with warmer locations.

Hot water coral bleaching, caused by global warming, is devastating coral reefs around
the world (*1*) but there is room for hope if corals can adapt to increasing temperatures (*2*). The
fact that current coral generation is suffering high mortality does not necessarily imply that the
next coral generation would not be better adapted. Many coral species have wide distributions
that span environments that differ dramatically in their thermal regimes, demonstrating that
efficient thermal adaptation has occurred in the past (*3*). But can coral adaptation keep up with
the unprecedentedly rapid current rate of global warming (*4*)? One way for corals to achieve rapid
thermal adaptation is through genetic rescue, involving the spread of existing heat tolerance
alleles from low-latitude, warm-adapted populations to higher-latitude, warming regions, via
larval migration (*5*, *6*). We have previously demonstrated the presence of genetic variants
conferring high thermal tolerance in a low-latitude *A. millepora* population (*5*). It can be
hypothesized that global warming will cause preferential survival of migrants from warmer to
cooler locations because they will be following their thermal optimum, whereas individuals
migrating in the opposite direction would find themselves in increasingly mismatched
environments (Fig. 1 A, B). Another likely population-level effect of recent declines in coral
cover (*7*) is a reduction in overall genetic diversity, potentially limiting both the scope and the
rate of adaptation. Here, we tested these predictions in *Acropora millepora*, a common reef-
building coral from the most ecologically prominent and diverse coral genus in the Indo-Pacific
(staghorn corals, *Acropora*), and used the obtained demographic estimates to model the future
adaptive potential of *A. millepora* on the Great Barrier Reef (GBR).

Results

Locations and genotyping

We used samples collected in 2002-2009 from five populations of *A. millepora* along the
latitudinal range of the GBR (Fig. 1 A). Environmental parameters (obtained from
http://eatlas.org.au/) varied widely among these locations (Fig. 1 C). Importantly, maximum
summer temperature (the major cause of bleaching-related mortality) followed the latitudinal
gradient with one notable exception: one of the near-shore populations from the central GBR
(Magnetic Island) experienced summers as hot as the lowest-latitude population (Wilkie Island
(Fig. 1 C).

We genotyped 18-28 individuals per population using 2bRAD (*8*) at >98% accuracy and
with a >95% genotyping rate. Analysis of population structure based on ~11,500 biallelic SNPs
separated by at least by 2,500 bases agreed with previous microsatellites results (*9*, *10*), and
revealed very low levels of genetic divergence, with only the Keppel Islands population being
potentially different from the others (Fig. 1 D and Fig. S1). Pairwise F_{ST} were small and did not
exceed 0.014 even between the southernmost and northernmost populations (Keppel and Wilkie).

Demographic subdivision and migration patterns

To more rigorously test for population subdivision and infer unidirectional migration rates among populations and population sizes, we used Diffusion Approximation for Demographic Inference (*dadi,* (*11*)). *dadi* is a method that optimizes parameters of a pre-specified demographic model to maximize the likelihood of generating the observed allele frequency spectrum (AFS). For two populations AFS is essentially a two-dimensional histogram of allele frequencies (Fig. S2). Being a likelihood-based method, *dadi* can be used to compare alternative models using likelihood ratio tests and Akaike Information Criterion (AIC). Most importantly, unlike previously used approaches (*10*, *12*, *13*) *dadi* does not rely on assumptions of genetic equilibrium (stability of population sizes and migration rates for thousands of generations) or equality of population sizes and therefore is potentially more realistic and sensitive for natural populations.

We used bootstrap-AIC approach to confirm that our populations are separate demographic units. For each pair of populations we generated 120 bootstrapped datasets by resampling genomic contigs and performed delta-AIC comparison of two demographic models, a split-with-migration model and a no-split model (Fig. S3 C). The split-with-migration model assumed two populations that split some time *T* in the past with potentially different sizes *N1* and *N2*, and exchange migrants at different rates (*m12* and *m21*) depending on direction. The no-split

model allowed for ancestral population size to change but not for a population split, so the
experimental data were modeled as two random samples from the same population of size *N*. The
majority of bootstrap replicates (53-100%) showed AIC advantage of the split-with-migration
model for all pairs of populations (Fig. S3). This indicates that *A. millepora* populations on the
GBR are in fact demographically distinct, despite typically non-significant F_{ST} reported by
previous studies based on allozymes (*12*, *13*) and microsatellite markers (*10*).
110

111
<u>Figure 2</u>. Demography of *A. millepora* populations on the GBR. (A) Arc-plot of migration rates among
populations reconstructed from population genetic data. Inset: *dadi* model used: ancestral population splits
into two populations of unequal sizes (N1 and N2) some time T in the past, these populations exchange
migrants at different rates depending on direction. (B) Migration rates according to the biophysical model.
On panels A and B, the arcs should be read clockwise to tell the direction of migration; line thickness is
proportional to the migration rate. (C) Correlation between log-transformed biophysical and genetic
migration rates (Mantel r = 0.58, *P* = 0.05). Grey symbols are migration rates to Magnetic Island, the high-
temperature location in the central GBR. (D) Box plot of effective population sizes inferred by the split-
with-migration model (panel A) across all population pairs and bootstrap replicates. (E) Historical changes
in effective population sizes inferred using a single-population model with two periods of exponential
growth (*T0* and *T1*, reaching sizes *N0* and *N1*, inset), averaged across bootstrap replicates.
123
AFS-based analysis allows rigorous estimation of unidirectional migration rates between
populations. The classical F_{ST} –based approach only allows estimating bi-directional migration
rate (*13*) and even that calculation has been highly criticized because its underlying assumptions
are rarely realistic (*14*). We determined unidirectional migration rates from the split-with-
migration model and estimated their confidence limits from bootstrap replicates. In theory,
migration rate can be confounded with population divergence time, since in the AFS higher
migration often looks similar to more recent divergence (*15*). To ascertain that the model with
ancient population divergence and migration is preferable to the model with very recent

divergence and no migration, we performed the delta-AIC bootstrap comparison between these
models and obtained overwhelming support for the model with ancient divergence and migration
(Fig. S4). Notably, for all pairwise analyses migration in southward direction exceeded northward
migration, and this difference was significant in seven out of nine cases (Fig. 2 A and Fig. S3A,
C). Linear mixed model analysis of direction dependent median migration rates with a random
effect of destination (to account for variation in total immigration rate) confirmed the overall
significance of this southward trend (P_{MCMC} <1e-4). Full listing of parameter estimates and their
bootstrap-derived 95% confidence limits is given in Table S1.
140
To investigate whether the southward migration bias was due to higher survival of warm-
adapted migrants, as predicted under global warming (Fig. 1 B), we developed a biophysical
model of coral larval dispersal on the Great Barrier Reef. This model quantified the per-
generation migration potential among coral reef habitat patches in the GBR based on ocean
currents and parameters of larval biology (16, 17). The biophysical model predicted very similar
migration rates as our genetic model (Mantel r =0.58, p = 0.05), recapitulating the southward bias
(Fig. 2 A-C). Importantly, the same southward bias was predicted for population pairs in which
southward migration corresponded to movement to the same-temperature or even to warmer
location, such as migrations to the Magnetic Island (Fig. 2 C, grey points). This indicates that
southward migration bias is predominantly driven by ocean currents and not by preferential
survival of warm-adapted coral genotypes migrating to cooler locations.
152
Migration estimates from our main genetic model (Fig. 2 A, inset) represented historical
averages since the populations split and did not resolve any potential recent migration changes.
To determine if there were any recent changes in southward migration, we evaluated an extended
split-with-migration model that allowed for a change in migration over the past 75-100 years. The
extended model suggested some recent migration changes, including southward migration
increases (Fig. S5) but once again, these changes did not correspond with an increase in migration
from warmer to cooler locations. We conclude that with the current data and analysis techniques
we cannot (yet) detect an effect of recent warming on preferential direction of coral migration on
the GBR.
162
163   *Genetic diversity trends*
164
The GBR has warmed considerably since the end of last century (18), which may have
already reduced genetic diversity in *A. millepora* populations. We used *dadi* to infer effective
population sizes, which is a measure of genetic diversity and one of the key parameters
determining the population's adaptive potential (19). The results of the split-with-migration
model (Fig. 2 A) were consistent for all population pairs and indicated that Keppel population
was about one-fifth the size of others (Fig. 2 D, E). This result was not surprising since the
Keppel population frequently suffers high mortality due to environmental disturbances and was
therefore is expected to show diminished long-term effective population size (9). We also used a
single-population *dadi* model that allowed for two consecutive growth/decline periods (Fig. 2 E,

inset) to reconstruct effective sizes of individual populations through time (Fig. 2 E and Fig S6).
All populations showed evidence of growth prior to the last glaciation, between 500 and 20
thousand years ago (Fig 2 E), which aligned well with the fossil record of rising dominance of
Acropora corals on Indo-Pacific reefs during this period (*20*). It was hypothesized that the fast
growth and early sexual maturation of *Acropora* corals gave them an advantage relative to most
other reef-building corals during dynamic changes in the reef-forming zone due to the sea level
changes accompanying glacial cycles (*20*). Our results suggest that *A. millepora* populations have
been in decline since sea level stabilized after the last deglaciation, roughly 10 thousand years ago
(Fig. S6). This decline was the most pronounced for the Keppel population (Fig. 2 E), but delta-
AIC bootstrap supported inclusion of additional growth/decline period for all populations except
Magnetic Island (Fig. S7). None of the populations showed evidence of accelerated decline in
effective population size over the past hundred years, despite recent GBR-wide decline in overall
coral cover (*7*). It must be noted that recent decline in population size is hard to detect using the
AFS method unless the sample size is very large, since it would predominantly affect the
frequencies of rare alleles (*15*, *21*). Despite this shortcoming, we can conclude that genetic
diversity in *A. millepora* has not yet been strongly affected by warming over the past century,
although the populations appear to have been in long-term decline GBR-wide for the past several
thousand years.
192
193   *Metapopulation adaptation model*
194
To evaluate whether standing genetic variation contributed by local thermal adaptation
could facilitate rapid adaptation of the *A. millepora* metapopulation in response to warming, we
developed an individual-based multigene model of metapopulation adaptation in the SLiM
software environment (*22*). The model's code is highly flexible and can simulate any number of
populations with any configuration of population sizes, migration rates, and environmental trends.
The model also allows varying the number and effect sizes of QTLs, heritability, and phenotypic
plasticity. Here, we used population sizes and migration rates inferred from the genetic analysis
(Fig. 2 A, D) and incorporated differences in midsummer monthly mean temperature among
populations (Fig. 1 A). The populations were allowed to adapt to local thermal conditions for
2,000 generations. Assuming a generation time of 5 years in *A. millepora* (*23*) this corresponded
to the period of stable temperature since the last deglaciation. After this pre-adaptation, the
temperature began to increase at a rate of 0.05^{o}C per generation in all populations, corresponding
to the projected 0.1^{o}C warming per decade (*24*). During warming period, a population declining
in fitness would shrink in size and stop contributing migrants to other populations. Throughout
the simulation the temperature was allowed to fluctuate randomly between generations to
approximate El Nino Southern Oscillation (ENSO).
211
We found that, with only ten thermal QTLs, under a broad range of settings for
heritability and plasticity the pre-adapted metapopulation was able to persist through the warming
for at least 20 - 50 generations (100 - 250 years) and, under some parameter combinations, much
longer (Fig. 3 and S8). Migration substantially contributed to this persistence (Fig. 3 E, F),

underscoring the importance of divergent local adaptation and genetic rescue (*5*). However, when
existing standing genetic variation was finally depleted, all populations inevitably went extinct
beginning with warmer locations. Allowing for new mutations could not avert this extinction
even when the mutation rate was extremely high - 1e-5 per QTL per gamete or higher while
assuming that half of all mutations were beneficial (Fig. 4).
221

222
Figure 3. Modeling coral metapopulation persistence under global warming based on standing genetic
variation (no new mutations). (A-D) Fitness of modeled populations depending on the magnitude of non-
heritable phenotypic component (*Esd,* standard deviation of normally distributed random value added to the
sum of QTL effects, in degrees C), phenotypic plasticity (σ, standard deviation of the Gaussian slope of
fitness decline away from the phenotypic optimum, in degrees C), and presence-absence of migration (E,
F). On panels A-F, y-axis is observed fitness relative to maximum attainable with perfect heritability at the
genetically determined optimum, averaged over all individuals in a population. Warm-adapted populations
(W and M) are shown as red-tint traces, populations from mild thermal regime (S and O) are green-tint
traces, and the cool-adapted population (K) is the blue trace. Note nearly complete overlap between traces
for pairs of populations pre-adapted to the same temperature (W, M and S, O). (G) Thermal tolerances of
evolving populations. Thin noisy lines are modeled temperatures at different locations. (H) Modeled
random temperature anomalies (grey line) and fluctuations in populations' fitness (colored lines: residuals
from loess regression over fitness traces on panel D; Wilkie: orange line, Keppel: blue line). Note the
inverse sign of temperature anomalies: this more clearly shows the correspondence between rise in
temperature and drop in fitness. As warming progresses, populations (especially originally warm-adapted
ones) become increasingly sensitive to random temperature fluctuations.
239
A notable tendency observed with all parameter settings was that during warming the
fitness (and hence the size) of adapting populations began to fluctuate following random thermal
anomalies, and the amplitude of these fitness fluctuations increased as the warming progressed

even though the amplitude of thermal anomalies did not change (Fig. 3 H). These fluctuations
correspond to severe mortality events induced by thermal extremes that can occur as a result of
ENSO and affected warm-adapted populations most, which very much resembles the situation
currently observed throughout the world (*1*).
247

248
Figure 4. New mutations cannot prevent extinction. Top row of graphs (A, C, E and G) show population
fitness; bottom row (B, D, F, H) – mean thermal tolerance. All cases share the settings of *Esd* = 1 and σ =
1. Mutation rate (*mu*) per QTL per gamete is listed above the graphs; effect sizes of new mutations were
drawn from a normal distribution with mean 0 and standard deviation 0.2. Note a few "evolutionary
rescue" events at high mutation rates, when a new adaptive mutation spreads through metapopulation
leading to temporary acceleration of phenotypic evolution (F, H).
255
256 *What helps corals adapt?*
257
Predictably, higher phenotypic plasticity promoted metapopulation persistence and
stability against random thermal anomalies, but we were rather surprised to observe a similar
positive effect of lower heritability (i.e., higher non-heritable component, Fig 3, S8 and S9). Low
heritability of thermal tolerance is expected for reef-building corals: much of natural variation in
this trait in corals is due to the type of algal symbionts (*Symbiodinium* spp. (*25*)). Photo-
symbionts are not transmitted from parent to offspring in the majority of coral species (*26*), and
although host genetics can have some effect on the choice of *Symbiodinium* in the next generation
(*27*) environment has stronger effect on this association (*25, 28*).
266
Longer metapopulation persistence under low heritability and high plasticity was most
likely due to their enhancing effect on standing genetic variation (Fig. 5). Higher plasticity
promoted both higher number of variants retained in populations and larger effect sizes of these
variants, whereas higher heritability also led to higher number of retained variants but notably
smaller effect sizes (Fig. 5 A-K). It can be said that under high heritability local adaptation was
based on many mutations of small effect, whereas low heritability promoted adaptation involving
fewer mutations of larger effect. Importantly, the cumulative absolute effect of QTL variants in a
population was consistently higher under the setting of low heritability, despite lower number of

variants (Fig. 5 K, L). During warming, this variation lasted longer as the source of adaptive
genetic variants, enabling up to 4°C increase in mean thermal tolerance (Fig. 3 G and S9).
Another important effect of higher plasticity was that it partially rescued the drop in average
population fitness due to low heritability (Fig. 3 B, D and S9). This drop happened because low
heritability prevented individuals from attaining maximum fitness even if their genetics was
perfectly matched to the environment.
281

282
<u>Figure 5</u>. Effects of non-heritable phenotypic component (*Esd*) and phenotypic plasticity (σ) on standing
genetic variation at thermal QTL loci. (A - I) Dependence of the number of individuals in each population
bearing a mutation on the mutation's effect size (change in thermal tolerance, in °C). Each mutation is
represented by up to five points colored according to the population in which it is found (see legend).
Lower heritability (higher *Esd*) and higher plasticity promote retention of mutations with larger effect sizes.
(A, D, G): *Esd*=0. (B, E, H): *Esd*=1. (C, F, I): *Esd*=2. (A-C): σ = 0.5. (D-F): σ = 1. (G-I): σ = 2. (K, L)
Genetic variation retained after 2000 generations of adaptation to benign local thermal conditions
(aggregating 12 simulation replicates for two populations, S and O). (K) Number of mutations at QTL loci.
(L) Cumulative effect size (sum of products of mutation's absolute effect size and the number of
individuals bearing the mutation). Lower heritability (higher *Esd*) results in retaining fewer mutations, but
these mutations amount to up to four-fold larger cumulative effect size, while higher plasticity promotes
both higher number of mutations and larger effect sizes.

295
296   *Potential pitfalls*
297
298   There are several uncertainties in our model associated with coral biology. Below we
299   argue that, while more research is certainly needed to resolve these uncertainties, our modeling
300   was conservative overall.
301

We assumed only ten QTLs, which is likely much fewer that the actual number of
thermal QTLs in acroporid corals (20). Higher number of QTLs and/or their larger effect sizes
would promote higher genetic variation and lead to longer population persistence. We also kept
the distribution of QTL effect sizes narrow: with the current settings and ten QTLs, at the start of
simulation only about 2% of corals deviated from the mean thermal tolerance by more than 1.5°C
in either direction. Such narrow variation makes adaptation to the thermal gradient of ~3°C along
the GBR non-trivial, but still, at present there is no experimental data to evaluate whether even
such narrow variation is realistic.
310
We used effective population sizes suggested by genetic analysis as census sizes. In
highly fecund marine organisms census sizes tend to substantially exceed effective population
sizes, sometimes by orders of magnitude (29), which would strongly promote higher genetic
diversity and population persistence. Moreover, we modeled only our five populations rather than
the whole GBR, which would have resulted in much higher standing genetic variation in the
metapopulation, promoting longer persistence.
317
As for phenotypic plasticity, in simulations shown on Figs. 3 $\sigma = 0.5$ and $\sigma = 1$
corresponded to 86% and 40% decline in fitness if the individual's phenotype mismatched the
environment by 1°C. The existing data on the issue of coral thermal plasticity are somewhat
conflicting. One study shows that acroporid corals can successfully acclimatize to environments
differing in maximum temperatures by as much as 2°C (30); however, another study found that
coral grew 52-80% more slowly when transplanted among locations differing by 1.5°C average
temperature, (31). Although it is not possible to directly interpret these results in terms of width
of the fitness function (as plasticity is encoded in our model) the former study likely supports the
higher plasticity setting ($\sigma = 1$) while the latter study supports $\sigma = 0.5$. It must also be noted that
both these studies involved *in situ* transplantations and hence the effect of temperature remains
confounded with other local fitness-affecting environmental parameters. Also, in adult corals
plasticity is likely lower that in larvae and recruits, which are expected to exhibit non-reversible
developmental plasticity associated with metamorphosis and establishment within a novel
environment (32). One particularly important event during this developmental transition is
establishment of association with local algal symbionts. Since symbionts also adapt to local
thermal conditions (28) this would elevate the fitness of the coral host despite possible mismatch
between its own genetically determined thermal optimum and local temperature, which in our
model implies broadening of the fitness function (i.e., higher plasticity). Future experiments that
expose multiple genetically distinct coral individuals to a range of temperatures under controlled
laboratory settings are required to rigorously quantify variation in thermal optima and plasticity in
natural populations.
339
Our demographic analysis and, by implication, our modeling results are contingent on the
accuracy of the mutation rate estimate. The mutation rate used here, 4e-9 per base per year, or 10
mutations per genome per generation (see Methods), is but a rough estimate based on
phylogenetic distances (33). Higher mutation rate would lead to smaller population sizes, higher

migration rates, and more recent time stamps for population splits or population size changes (see
Methods, Unit Conversion section). As a consequence, it would lead to lower standing genetic
variation in the metapopulation and therefore to diminished potential to persist under global
warming. Although the rate used here is within the ballpark of mutation rates measured in
multicellular animals (*34*) in the future a more precise estimate of mutation rate should be
obtained to sharpen the demographic estimates and parameterize the adaptation model more
realistically.
351
It may be argued that our samples are genetically out of date, not capturing the effects of
disturbances that happened on the GBR since the time of their collection (2002-2009). However,
as we mentioned above, very recent demographic events (in our case, 2-3 generations ago) are
undetectable at the level of neutral genetic variation unless the number of disturbance-surviving
individuals is similar to the study's sample size (i.e., when either the disturbance was truly
catastrophic or the sample size is very large). Thus, our samples can still be considered well
representative of major patterns of genetic diversity of our study species.
359
Finally, our model assumed that recovery from high mortality events would happen
without impediment, through reseeding by survivors and migrant influx from other coral
populations. However, ecological feedbacks such as shifts to an alternative ecological stable state
(*35*) might substantially decrease the rate of reseeding and recovery of affected reefs. In that case,
the increase in severity of bleaching-related mortality might lead to much faster coral extinction
than predicted by our model.
366
367   *Conclusions*
368
Our study provides a novel integrated empirical and modeling framework to evaluate the
risk of extinction in natural populations. We found that genetic diversity and migration patterns of
Acropora millepora were not yet strongly affected by climate change and were well positioned to
facilitate persistence of the GBR metapopulation for a century or more. Our results underscore
the pivotal role of standing genetic variation and genetic exchange in the future metapopulation
persistence. This implies that any intervention that would reduce this variation (for example,
captive breeding based on selection of only a few "winner" genotypes or clonally propagating a
small number of genotypes for reef restoration) is likely to have negative impact on corals'
adaptive potential. In contrast, efforts facilitating the spread of genetic variation, such as assisted
gene flow (*36*), could be much more helpful in the long term. With the estimated natural
migration rates on the order of 5-100 migrants per generation, human-assisted genotype exchange
could appreciably contribute to the genetic rescue without risking disruption of the natural local
adaptation patterns (*37*). However, despite good prospects for short-term adaptation, corals are
predicted to become increasingly more sensitive to random thermal anomalies, especially in the
originally warm-adapted populations. The 10-85% mortality in the Northern GBR as a result of
2016 bleaching event (*38*) could be a particularly sobering recent manifestation of this trend.
Finally, it is important to point out that adaptation based on genetic rescue will not save corals

from eventual extinction: it will only buy us some time to take action against global warming,
which hopefully can be stopped before corals run out of genetic variation.
388
389 **Methods**
390
391 *Genotyping*
392
This study relied predominantly on samples described by van Oppen et al (*10*), with
addition of several samples from Orpheus and Keppel islands that were used in the reciprocal
transplantation experiment described by Dixon et al (*39*). The samples were genotyped using
2bRAD (*8*) modified for Illumina sequencing platform; the latest laboratory and bioinformatics
protocols are available at https://github.com/z0on/2bRAD_GATK. BcgI restriction enzyme was
used and the samples retained for this analysis had 2.3-20.2 (median: 7.45) million reads after
trimming and quality filtering (no duplicate removal was yet implemented in this 2bRAD
version). The reads were mapped to the genome of the outgroup species, *Acropora digitifera* (*40*,
41), to polarize the allelic states into ancestral (as in *A. digitifera*) and derived, e.g., (*42*, *43*).
Genotypes were called using GATK pipeline (*44*).
403
Preliminary analysis of sample relatedness using vcftools (*45*) revealed that our samples
included several clones: four repeats of the same genotype from the Keppel Island (van Oppen et
al (*10*) samples K210, K212, K213 and K216), another duplicated genotype from Keppel
(samples K211 and K219), and one duplicated genotype from Magnetic Island (samples M16 and
M17). All other samples were unrelated. We took advantage of these clonal replicates to extract
SNPs that were genotyped with 100% reproducibility across replicates and, in addition, appeared
as heterozygotes in at least two replicate pairs (script replicatesMatch.pl with hetPairs=2 option).
These 7,904 SNPs were used as "true" SNP dataset to train the error model to recalibrate variant
quality scores at the last stage of the GATK pipeline. During recalibration, we used the transition-
transversion (Ts/Tv) ratio of 1.438 determined from the "true" SNPs to assess the number of false
positives at each filtering threshold (as it is expected that an increase of false positive calls would
decrease the Ts/Tv ratio towards unity). We chose the 95% tranche, with novel Ts/Tv = 1.451.
After quality filtering that restricted the calls to only bi-allelic polymorphic sites, retained only
loci genotyped in 95% or more of all individuals, and removed loci with the fraction of
heterozygotes exceeding 0.6 (possible lumped paralogs), we ended up with 25,090 SNPs. In total,
2bRAD tags interrogated 0.18% of the genome. The genotyping accuracy was assessed based on
the match between genotyped replicates using script repMatchStats.pl. Overall agreement
between replicates was 98.7% or better with the heterozygote discovery rate (fraction of matching
heterozygote calls among replicates) exceeding 96%.
423
424 *Genome-wide genetic divergence*
425
To begin to characterize genome-wide divergence between populations we used pairwise
genome-wide Weir and Cockerham's F_{ST} calculated by vcftools (*45*), principal component

analysis (PCA) using R package adegenet (*46*), and ADMIXTURE (*47*). For PCA and
ADMIXTURE, the data were thinned to keep SNPs separated by 5kb on average and by at least
2.5 kb, choosing SNPs with highest minor allele frequency (script thinner.pl with options
'interval=5000 criterion=maxAF'). The optimal K in ADMIXTURE analysis was determined
based on the cross-validation procedure incorporated within ADMIXTURE software; the lowest
standard error in cross-validation was observed at K=1.
434
435   *Demographic analysis and bootstrapping*
436
Prior to demographic analysis, Bayescan (*48*) was used to identify sites potentially under
selection among populations, and 73 sites with q-value <0.5 were removed. This aggressive
removal of potential non-neutral sites resulted in better agreement between bootstrap replicates
compared to an earlier of analysis where only 13 sites with q-value < 0.05 were removed.
Demographic models were fitted to 120 bootstrapped datasets, which were generated in two
stages. First, three alternatively thinned datasets were generated for which SNPs were randomly
drawn to be on average 5 kb apart and not closer than 2.5 kb. This time the SNPs were drawn at
random to avoid distorting the allele frequency spectrum (unlike thinning for PCA and
ADMIXTURE where the highest minor allele frequency SNPs were selected). Then, 40
bootstrapped replicates were generated for each thinned dataset by resampling contigs of the
reference genome with replacement (script dadiBoot.pl). The fitted model parameters were
summarized after excluding bootstrap replicates that fell into the lowest 15% likelihood quantile
and the ones where model fitting failed to converge, leading to some parameters being
undetermined or at infinity (less than 10% of total number of runs). Delta-AIC values were
calculated for each bootstrap replicate that passed these criteria for both compared models, and
summarized to obtain bootstrap support value, the percentage of replicates favoring the
alternative model. While fitting *dadi* models, the data for each population were projected to
sample sizes maximizing the number of segregating sites in the analysis, resulting in 7000-8172
segregating sites per population. Initially, our models included a parameter designed to account
for ancestral state misidentification rate when constructing the polarized AFS (e.g., (*49*)), but
since this parameter was consistently estimated to be on the order of 0.001 and had negligible
effect on the models' likelihood, we removed it from the final set of models.
459
460   *Unit conversion*
461
To convert *dadi*-reported coalescent parameter values (θ, T and M) into time in years (t),
effective population sizes in number of individuals (Ne) and migration rates as fraction of new
immigrants per generation (m), we estimated the mutation rate (μ) from the time-resolved
phylogeny of *Acorpora* genus based on *paxC* intron (*33*), at 4e-9 per base per year. Although *A.*
millepora can reproduce after 3 years (*23*) we assumed a generation time of 5 years reasoning that
it would better reflect the attainment of full reproductive potential as the colony grows. Assuming
a genome size of 5e+8 bases (*40*) the number of new mutations per genome per generation is 10.
Since the fraction 2bRAD-sequenced genome in our experiment was 1.8e-3, the mutation rate per

2bRAD-sequenced genome fraction per generation is $\mu = 0.018$. This value was used to obtain:
-   Ancestral effective population size: $Ne = \theta / 4\mu$
-   Migration rate: $m = M / 2Ne$
-   Time in years: $t = 2TNe \cdot 5$
474
475   *Biophysical model*
476
A spatially-explicit biophysical modeling framework (*16*, *50*) was used to quantify
migration between coral reef habitats of the broader region surrounding the Great Barrier Reef,
thereby revealing the location, strength, and structure of a species' potential population
connectivity. The model's spatial resolution of ca. 8 km coincides with hydrodynamic data for the
broader region (1/12.5 degree; HYCOM+NCODA Reanalysis and Analysis product; hycom.org).
Our biophysical dispersal model relies on geographic data describing the seascape environment
and biological parameters capturing coral-specific life-histories. Coral reef habitat data are
available from the UNEP World Conservation Monitoring Centre (UNEP-WCMC;
http://data.unep-wcmc.org/datasets/1) representing a globally-consistent and up-to-date
representation of coral reef habitat. To capture specific inter-annual variability, two decades of
hydrodynamic data were used from 1992 to 2013 (*51*).
488
Coral-specific biological parameters for *A. millipora* included relative adult density
(dependent on the habitat), reproductive output, larval spawning time and periodicity (e.g.,
Magnetic Island populations spawn a month earlier than the other GBR sites (*52*)), maximum
dispersal duration, pre-competency and competency periods, and larval mortality (*53*, *54*). The
spatially explicit dispersal simulations model the dispersal kernel (2-D surface) as a 'cloud' of
larvae, allowing it to be concentrated and/or dispersed as defined by the biophysical parameters.
An advection transport algorithm is used for moving larvae within the flow fields (*55*).
496
Simulations were carried out by releasing a cloud of larvae into the model seascape at all
individual coral reef habitat patches and allowing the larvae to be transported by the currents.
Ocean current velocities, turbulent diffusion, and larval behavior move the larvae through the
seascape at each time-step. Larval competency, behavior, density, and mortality determine when
and what proportion of larvae settle in habitat cells at each time step. When larvae encounter
habitat, the concentration of larvae settling with the habitat is recorded at that time-step. From the
dispersal data, we derived the coral migration matrix representing the proportion of settlers to
each destination patch that came from a source patch, which is analogous to the source
distribution matrix (*56*) and is equivalent to migration matrices derived from population genetic
analysis. It is important to note that migration matrices extracted for the field sites represent the
potential migration through all possible stepping-stones.
508
509   *Metapopulation adaptation model*
510
The model was implemented in SLiM (*22*), the forward evolutionary simulator, by

modifying the provided recipe "Quantitative genetics and phenotypically-based fitness". The
model simulates Fisher-Wright populations with discreet generations. At the start of the
simulation, populations are established at specified population sizes and pairwise migration rates
(genetic replacement rates), and all QTLs in all individuals are given a mutation with the effect
size drawn from a normal distribution with mean zero and specified standard deviation, to create
standing genetic variation. The phenotype of each individual is calculated as the sum of QTL
effects plus random noise of specified magnitude to simulate non-heritable phenotypic
component. Then, fitness of each individual is calculated based on the difference between the
individual's phenotype (thermal optimum), temperature of the environment, and the setting for
phenotypic plasticity, modeled as the width of the fitness curve: the standard deviation of the
Gaussian slope of fitness decline away from the phenotypic optimum. Then, parents are chosen
to produce the next generation according to their fitness; parents for immigrant individuals are
chosen from among individuals in the source population. New mutations at QTLs happen at the
specified rate when transitioning to the next generation and the effect of a new mutation adds to
the previous QTL effect. To better model population dynamics, we implemented linear scaling of
the population size and immigration rates with the population's mean fitness. In the model
described here this scaling was applied during warming period, so that a population declining in
fitness relative to its state at the end of pre-adaptation period shrinks in size and stops
contributing migrants to other populations.
531
532   Adjustable model parameters and their settings in this study:
533
-   Number of QTLs and the distribution of their effect sizes. To keep the model conservative,
we modeled only ten QTLs with normal distribution of effect sizes with a standard deviation
of $0.2^{o}C$. With ten QTLs, this setting implied that at the start of simulation only about 2% of
corals deviated from mean thermal tolerance by more than $1.5^{o}C$ in either direction. Since
thermal differences between our populations exceeded $3^{o}C$, this narrow variation made local
adaptation rather non-trivial.
-   Dominance of QTLs (set to 0.5 in our simulation).
-   Phenotypic plasticity: standard deviation of the Gaussian curve describing decline in fitness
away from phenotypic optimum. We modeled three plasticity settings, 0.5, 1 and 2, which
corresponded to 86%, 40% and 13% fitness drop when the environment mismatched
phenotypic optimum by $1^{o}C$.
-   Non-heritable phenotypic component: standard deviation of a normal distribution with mean
zero from which a random value is drawn to be added to the sum of QTL effects when
computing phenotype. Setting this parameter to zero corresponds to trait heritability of one.
Higher values of this parameter imply heritability less than one; however, the exact value of
heritability (the proportion of phenotypic variation explained by genetics) could still vary
depending on the extent of genetic variation.
-   Mutation rate. It was either set to zero to explore the role of standing genetic variation or
varied between 1e-6 and 1e-4 per QTL per gamete. This range covers and exceeds the range
of trait-level deleterious mutation rates observed in humans (57). Therefore, values at higher

end of the range most likely strongly over-estimate the rate of adaptive mutations, which was
done deliberately to show that no realistic mutation rate could help sustain genetic variation
in the face of strong selection by warming.
557
558   The model's code, available at https://github.com/z0on/Adaptive-pathways-of-coral-
559 populations-on-the-Great-Barrier-Reef, is designed for general modeling of multilocus adaptation
560 in metapopulations. It can read user-supplied files of environmental conditions, population sizes
561 and migration matrices for arbitrary number of populations.
562
Here, we modeled our five populations with effective population sizes and pairwise
migration rates inferred by *dadi*. We modeled identical thermal trends across populations with
population-specific offsets. During pre-adaptation period lasting 2000 generations, the
temperature was constant on average but experienced random fluctuations across generations
drawn from a normal distribution with a standard deviation of 0.25°C (to approximate ENSO
events). The temperature was offset by $+1.6^{\circ}$C in Wilkie and Magnetic populations and by -1.8°C
in the Keppel population, to model differences in midsummer monthly mean temperature among
populations (Fig. 1). After 2000 generations a linear increase at 0.05°C per generation was added
to simulate warming.
572
573   All combinations of parameter settings were run ten times to ensure consistency. We
574 found that with population sizes in thousands, such as in our case, the results were very
575 consistent among independent runs. We therefore did not aggregate results over many replicated
576 runs but show one randomly chosen run for each tested parameter combination.
577

578 **Acknowledgements**
579
We wish to thank Ryan Gutenkunst and Benjamin Haller for their continuous support of
dadi and SLiM users, respectively. The bioinformatics analysis was accomplished using
computational resources of the Texas Advanced Computer Center. This study has been supported
by NSF (DEB-1054766) grant to M.V.M, ARC (LP120200245) and University of Melbourne
ECR grants to E. A.T., a Coral Reef Alliance grant ("Coral Adaptation Challenge") to E.A.T and
M.V.M, Queensland Government funding to L.K.B and AIMS funding to L.K.B. and M.J.V.O

Data and code availability

The finalized genotyping dataset in VCF format, detailed bioinformatic walkthrough,
accessory formatting and plotting scripts, *dadi* scripts and the SLiM model code are available
from https://github.com/z0on/Adaptive-pathways-of-coral-populations-on-the-Great-Barrier-
Reef. Raw sequencing data has been deposited to National Center for Biotechnology
Information's Short Reads Archive (accession number pending).
592
593

References

1. A. C. Baker, P. W. Glynn, B. Riegl, Climate change and coral reef bleaching: An ecological assessment of long-term impacts, recovery trends and future outlook. *Estuar. Coast. Shelf Sci.* **80**, 435–471 (2008).

2. C. A. Logan, J. P. Dunne, C. M. Eakin, S. D. Donner, Incorporating adaptive responses into future projections of coral bleaching. *Glob. Chang. Biol.* **20**, 125–139 (2014).

3. T. P. Hughes *et al.*, Climate change, human impacts, and the resilience of coral reefs. *Science.* **301**, 929–933 (2003).

4. J. M. Pandolfi, S. R. Connolly, D. J. Marshall, A. L. Cohen, Projecting coral reef futures under global warming and ocean acidification. *Science.* **333**, 418–22 (2011).

5. G. B. B. Dixon *et al.*, Genomic determinants of coral heat tolerance across latitudes. *Science.* **348**, 1460–1462 (2015).

6. P. K. Ingvarsson, Restoration of genetic variation lost – the genetic rescue hypothesis. *Trends Ecol. Evol.* **16**, 62–63 (2001).

7. G. De'ath, K. E. Fabricius, H. Sweatman, M. Puotinen, The 27-year decline of coral cover on the Great Barrier Reef and its causes. *Proc. Natl. Acad. Sci. U. S. A.* **109**, 17995–9 (2012).

8. S. Wang, E. Meyer, J. K. McKay, M. V Matz, 2b-RAD: a simple and flexible method for genome-wide genotyping. *Nat. Methods.* **9**, 808–810 (2012).

9. M. J. H. van Oppen, V. Lukoschek, R. Berkelmans, L. M. Peplow, A. M. Jones, A population genetic assessment of coral recovery on highly disturbed reefs of the Keppel Island archipelago in the southern Great Barrier Reef. *PeerJ.* **3**, e1092 (2015).

10. M. J. H. Van Oppen, L. M. Peplow, S. Kininmonth, R. Berkelmans, Historical and contemporary factors shape the population genetic structure of the broadcast spawning coral, Acropora millepora, on the Great Barrier Reef. *Mol. Ecol.* **20**, 4899–4914 (2011).

11. R. N. Gutenkunst, R. D. Hernandez, S. H. Williamson, C. D. Bustamante, Inferring the joint demographic history of multiple populations from multidimensional SNP frequency data. *PLoS Genet.* **5**, e1000695 (2009).

12. D. J. Ayre, T. P. Hughes, Genotypic diversity and gene flow in brooding and spawning corals along the Great Barrier Reef, Australia. *Evolution (N. Y).* **54**, 1590–1605 (2000).

13. K. J. Miller, D. J. Ayre, The role of sexual and asexual reproduction in structuring high latitude populations of the reef coral Pocillopora damicornis. *Heredity (Edinb).* **92**, 557–568 (2004).

14. M. C. Whitlock, D. E. McCauley, Indirect measures of gene flow and migration: FST≠1/(4Nm+1). *Heredity (Edinb).* **82**, 117–125 (1999).

15. J. D. Robinson, A. J. Coffman, M. J. Hickerson, R. N. Gutenkunst, Sampling strategies for frequency spectrum-based population genomic inference. *Bmc Evol. Biol.* **14**, 254 (2014).

16. E. A. Treml, J. Roberts, P. N. Halpin, H. P. Possingham, C. Riginos, The emergent geography of biophysical dispersal barriers across the Indo-West Pacific. *Divers. Distrib.* **21**, 465–476 (2015).

17. E. A. Treml *et al.*, Reproductive Output and Duration of the Pelagic Larval Stage Determine Seascape-Wide Connectivity of Marine Populations. *Integr. Comp. Biol.* **52**, 525–537 (2012).

18. J. Lough, A. Hobday, Observed climate change in Australian marine and freshwater environments. *Mar. Freshw. Res.* **62**, 984–999 (2011).

19. B. Charlesworth, Fundamental concepts in genetics: Effective population size and patterns of molecular evolution and variation. *Nat. Rev. Genet.* **10**, 195–205 (2009).

20. W. Renema *et al.*, Are coral reefs victims of their own past success? *Sci. Adv.* **2**, e1500850 (2016).

21. A. Keinan, A. G. Clark, Recent Explosive Human Population Growth Has Resulted in an Excess of Rare Genetic Variants. *Science.* **336**, 740–743 (2012).

22. B. C. Haller, P. W. Messer, SLiM 2: Flexible, interactive forward genetic simulations. *Mol. Biol. Evol.* **34**, 230–240 (2017).

23. M. Vanessa, B. Baria, D. W. Dela Cruz, R. D. Villanueva, J. R. Guest, Spawning of three-year-old Acropora millepora corals reared from larvae in Northern Philippines. *Bull. Mar. Sci.* **88**, 61–62 (2012).

24. IPCC, *Climate Change 2007 - The Physical Science Basis. Contribution of Working Group I to the*

Fourth Assessment Report of the IPCC (Cambridge University Press, New York, NY, 2007).

25. R. Berkelmans, M. J. H. van Oppen, The role of zooxanthellae in the thermal tolerance of corals: a
"nugget of hope" for coral reefs in an era of climate change. *Proc. R. Soc. B-Biological Sci.* **273**,
2305–2312 (2006).

26. A. H. Baird, J. R. Guest, B. L. Willis, Systematic and biogeographical patterns in the reproductive
biology of scleractinian corals. *Annu. Rev. Ecol. Evol. Syst.* **40**, 551–571 (2009).

27. K. Quigley, B. Willis, L. Bay, Heritability of the Symbiodinium community in vertically- and
horizontally-transmitting broadcast spawning corals. *bioRxiv*. **doi: https** (2017).

28. E. J. Howells *et al.*, Coral thermal tolerance shaped by local adaptation of photosymbionts. *Nat.*
Clim. Chang. **2** (2011), pp. 116–120.

29. F. P. Palstra, D. J. Fraser, Effective/census population size ratio estimation: a compendium and
appraisal. *Ecol. Evol.* **2**, 2357–2365 (2012).

30. S. R. Palumbi, D. J. Barshis, N. Traylor-Knowles, R. A. Bay, Mechanisms of reef coral resistance
to future climate change. *Science*. **344**, 895–8 (2014).

31. E. J. Howells, R. Berkelmans, M. J. H. van Oppen, B. L. Willis, L. K. Bay, Historical thermal
regimes define limits to coral acclimatization. *Ecology*. **94**, 1078–1088 (2013).

32. D. Nettle, M. Bateson, Adaptive developmental plasticity: what is it, how can we recognize it and
when can it evolve? *Proc. R. Soc. B Biol. Sci.* **282**, 20151005 (2015).

33. Z. T. Richards, D. J. Miller, C. C. Wallace, Molecular phylogenetics of geographically restricted
Acropora species: Implications for threatened species conservation. *Mol. Phylogenet. Evol.* **69**,
837–851 (2013).

34. M. Lynch, Evolution of the mutation rate. *Trends Genet.* **26**, 345–52 (2010).

35. T. J. Done, Phase shifts in coral reef communities and their ecological significance. *Hydrobiologia*.
247, 121–132 (1992).

36. O. Hoegh-Guldberg *et al.*, Assisted colonization and rapid climate change. *Science*. **321**, 345–346
(2008).

37. S. N. Aitken, M. C. Whitlock, Assisted Gene Flow to Facilitate Local Adaptation to Climate
Change. *Annu. Rev. Ecol. Evol. Syst.* **44**, 367–388 (2013).

38. Great Barrier Reef Marine Park Authority, *Interim report on the environmental impacts of the 2016*
coral bleaching event.

39. G. B. Dixon, L. K. Bay, M. V. Matz, Bimodal signatures of germline methylation are linked with
gene expression plasticity in the coral Acropora millepora. *BMC Genomics*. **15**, 1109 (2014).

40. C. Shinzato *et al.*, Using the Acropora digitifera genome to understand coral responses to
environmental change. *Nature*. **476**, 320-U82 (2011).

41. M. J. H. van Oppen, B. J. McDonald, B. Willis, D. J. Miller, The Evolutionary History of the Coral
Genus Acropora (Scleractinia, Cnidaria) Based on a Mitochondrial and a Nuclear Marker:
Reticulation, Incomplete Lineage Sorting, or Morphological Convergence? *Mol. Biol. Evol.* **18**,
1315–1329 (2001).

42. B. F. Voight, S. Kudaravalli, X. Wen, J. K. Pritchard, A map of recent positive selection in the
human genome. *PLoS Biol.* **4**, e72 (2006).

43. I. K. Jordan *et al.*, A universal trend of amino acid gain and loss in protein evolution. *Nature*. **433**,
633–8 (2005).

44. A. Mckenna *et al.*, The Genome Analysis Toolkit: A MapReduce framework for analyzing next-
generation DNA sequencing data. *Genome Res.* **20**, 1297–1303 (2010).

45. P. Danecek *et al.*, The variant call format and VCFtools. *Bioinformatics*. **27**, 2156–8 (2011).

46. T. Jombart, adegenet: a R package for the multivariate analysis of genetic markers. *Bioinformatics*.
24, 1403–5 (2008).

47. D. H. Alexander, J. Novembre, K. Lange, Fast model-based estimation of ancestry in unrelated
individuals. *Genome Res.* **19**, 1655–64 (2009).

48. T. Günther, G. Coop, Robust identification of local adaptation from allele frequencies. *Genetics*.
195, 205–20 (2013).

49. M. Tine *et al.*, European sea bass genome and its variation provide insights into adaptation to
euryhalinity and speciation. *Nat. Commun.* **5**, 5770 (2014).

50.   E. A. Treml, P. N. Halpin, D. L. Urban, L. F. Pratson, Modeling population connectivity by ocean
currents, a graph-theoretic approach for marine conservation. *Landsc. Ecol.* **23**, 19–36 (2008).
51.   E. P. Chassignet *et al.*, The HYCOM (HYbrid Coordinate Ocean Model) data assimilative system.
J. Mar. Syst. **65**, 60–83 (2007).
52.   R. C. Babcock *et al.*, Synchronous Spawnings of 105 Scleractinian Coral Species on the Great-
Barrier-Reef. *Mar. Biol.* **90**, 379–394 (1986).
53.   S. W. Davies, E. A. Treml, C. D. Kenkel, M. V. Matz, Exploring the role of Micronesian islands in
the maintenance of coral genetic diversity in the Pacific Ocean. *Mol. Ecol.* **24**, 70–82 (2015).
54.   S. R. Connolly, A. H. Baird, Estimating dispersal potential for marine larvae: dynamic models
applied to scleractinian corals. *Ecology*. **91**, 3572–3583 (2010).
55.   P. K. Smolarkiewicz, J. Szmelter, An MPDATA-based solver for compressible flows. *Int. J.*
Numer. Methods Fluids. **56**, 1529–1534 (2008).
56.   R. K. Cowen, G. Gawarkiewicz, J. Pineda, S. Thorrold, F. Werner, Population Connectivity in
Marine Systems: An Overview. *Oceanography*. **20**, 14–21 (2007).
57.   M. R. Cummings, *Human heredity : principles and issues* (Brooks/Cole, Cengage Learning, 2014).
715

  Supplemental Figures

721 Figure S1. Principal component analysis of genetic diversity in sampled populations. (A) Map of sampled
722 locations with one-letter population identifiers. (B) Principal component analysis of genome-wide genetic
723 variation. On panel D, centroid labels are initial letters of population names as in panel A.

729 Figure S2. Example of two-population *dadi* model fit. (A) The model: ancestral population splits into two
730 populations of unequal sizes (N1 and N2) some time T in the past, which exchange migrants with different
731 rates depending on direction. (B) Observed allele frequency spectrum comparing Wilkie (W) and Keppel
732 (K) populations. (C) Allele frequency spectrum generated by the fitted model. (D, E) Map and histogram of
733 residuals (absolute scale).

Figure S3. Bootstrap analysis of migration rates, divergence times, and population subdivision. (A)

Migration among population pairs, with bootstrap-derived 95% confidence intervals. The pairs are

identified on the x-axis and sorted by increasing geographical distance. Black bars – southward migration,

grey bars – northward migration. (B) Boxplot of divergence times (in years, y-axis) between pairs of

populations (x-axis) across bootstrap replicates. (C) Models being compared: the split model (left) implies

populations' split into two different sizes (N1 and N2) at time T in the past, since when they exchanged

migrants at unequal rates depending on direction. No-split model (right) allows for population size change

at time T in the past but does not include population split: the two genotyped groups (p1 and p2) are

regarded as two samples from the same population. (D-M) Histograms of delta-AIC values comparing split

and no-split models (panel A) for bootstrap replicates (bootstrap was performed over genomic contigs of

the draft genome of *A. digitifera*). Positive numbers indicate support for the split model. The letters on top

of each panel identify compared populations, the number is the proportion of positive bootstrap replicates

750   (i.e., bootstrap support for the full model).

753
Figure S4. Delta-AIC bootstrap comparison of models with and without migration (A), to confirm that the
model with migration and ancient divergence is preferable to the model with no migration but very recent
divergence. (B-K) Histograms of delta-AIC values for bootstrap replicates comparing models with and
without migration. Positive numbers indicate support for the model with migration. The letters on top of
each panel identify compared populations, the number is the proportion of bootstrap replicates supporting
the model with migration and ancient split. For all pairs of populations the model of ancient split with
migration is strongly supported. (L, M) Example of residuals from the two models. Model without
migration under-estimates the number of shared low-frequency polymorphisms and over-estimates the
number of shared high-frequency polymorphisms.

763
764
765
766

767
Figure S5. Migration rates inferred by the *dadi* model allowing for the change in migration rates over the
last 0.01 T units (15-20 generations or 75-100 years, in our case). Box plots show historical (N, S) and
recent (Nr, Sr) migration rates inferred among pairs of population across bootstrap replicates. Numbers in
the top left corner are delta-AIC bootstrap support for the model with the recent change in migration
compared to the split-with-migration model with no recent change (Fig. 2A, inset). Only in three cases
(OW, MW, and KW) the model with recent migration change is favored, and in two of these cases (OW,
MW) southward migration shows pronounced recent increase. Southward migration increase is also
suggested for three more population pairs in the central GBR (OS, MS and MO): although in these cases
bootstrap support for the recent increase model is below 50%, the model is not strongly disfavored.

777
778
779
780

781
782   Figure S6. Population history. (A-E) Historical population sizes with bootstrap-derived 95% confidence
783   intervals, according to the two-growth model (Fig. S7 F). (F) Sea level with shaded area corresponding to
784   standard error (*41*).

785
786
787

788
Figure S7. Delta-AIC bootstrap analysis of single-population models. (A-E) Histograms of delta-AIC
values for 100 bootstrap replicates comparing two-growths and one-growth models (panel F). Positive
numbers indicate support for the two-growth model. The letter on top of each panel identify the population,
the number is the proportion of positive bootstrap replicates (i.e., bootstrap support for the two-growth
model). The two-growth model is well supported for populations W, O, and K (panels A, C and E), weakly
supported for population S (panel B), and not supported for population M (panel D). (F) Models compared.
The full model (left) includes two exponential growth periods (any of which could be growth or decline),
the reduced model (right) has only one growth period.

797
798

799
Figure S8. Higher plasticity and lower heritability facilitate metapopulation persistence during warming.
The graphs show fitness of populations (relative to maximal fitness at the genetically determined optimum)
after pre-adaptation period and under warming, depending on the magnitude of non-heritable component
(*Esd*, standard deviation of normally distributed random value added to the sum of QTL effects when
calculating individual's phenotype; higher *Esd* implies lower heritability) and phenotypic plasticity (σ,
standard deviation of the Gaussian slope of fitness decline away from the phenotypic optimum, in degrees
C). Higher plasticity confers stability against random thermal fluctuations (compare panels A, D and G)
and partially rescues the drop in fitness due to high *Esd* (i.e., lower heritability - compare pre-warming
generations, from -100 to 0, on panels B, E and H or C, F, and I).
809

811
<u>Figure S9</u>. Higher plasticity and lower heritability promote longer and more extensive evolution in response
to warming. The graphs show mean thermal tolerance of populations after pre-adaptation period and under
warming, depending on the magnitude of non-heritable component (*Esd*, standard deviation of normally
distributed random value added to the sum of QTL effects when calculating individual's phenotype; higher
Esd implies lower heritability) and phenotypic plasticity (σ, standard deviation of the Gaussian slope of
fitness decline away from the phenotypic optimum, in degrees C).
818

Table S1. Summary of demographic analysis based on split-with-migration model (Fig. 2 A). "lo" and "hi" values are bootstrap-based 95% confidence limits.

This table (S2M_results_summary) as well as full listing of raw parameter values obtained for all bootstrap replicates (s2m_rawParameters) are available at https://github.com/z0on/Adaptive-pathways-of-coral-populations-on-the-Great-Barrier-Reef

pop1	pop2	nu1.median	nu1.hi	nu1.lo	nu2.median	nu2.hi	nu2.lo	T.median	T.lo	T.hi	m12.median	m12.lo	m12.hi	m21.median	m21.lo	m21.hi
W	S	9800	5000	14000	8000	4200	14000	290000	110000	600000	0.0020	0.0008	0.0120	0.0060	0.0010	0.0130
W	O	12000	9600	14000	7600	4500	9900	300000	130000	700000	0.0017	0.0012	0.0021	0.0094	0.0061	0.0140
W	M	11000	7700	14000	8000	5500	12000	220000	110000	400000	0.0015	0.0009	0.0026	0.0065	0.0028	0.0110
W	K	14000	13000	16000	1900	1100	3100	130000	96000	180000	0.0005	0.0000	0.0008	0.0054	0.0031	0.0099
S	O	13000	8000	15000	11000	7400	14000	170000	96000	850000	0.0011	0.0003	0.0045	0.0096	0.0026	0.0150
S	M	14000	8800	15000	13000	7300	29000	140000	61000	720000	0.0009	0.0003	0.0025	0.0130	0.0053	0.0180
S	K	15000	13000	18000	1500	840	2700	150000	110000	300000	0.0004	0.0000	0.0018	0.0086	0.0043	0.0150
O	M	11000	2700	14000	11000	4900	16000	140000	60000	580000	0.0016	0.0008	0.0170	0.0090	0.0007	0.0130
O	K	15000	13000	19000	1800	750	2600	140000	94000	320000	0.0009	0.0005	0.0019	0.0068	0.0041	0.0170
M	K	18000	15000	21000	2800	830	3400	370000	120000	430000	0.0021	0.0012	0.0024	0.0041	0.0033	0.0170