CRISPR/Cas9 screening using unique molecular identifiers

Bernhard Schmierer[1,#], Sandeep K. Botla[1,#], Jilin Zhang[1], Mikko Turunen[2], Teemu Kivioja[2] and Jussi Taipale[1,2,*]

[1]Department of Medical Biochemistry and Biophysics, Karolinska Institutet, 171 77 Stockholm, Sweden.
[2]Genome-Scale Biology Research Program, Faculty of Medicine, University of Helsinki, PO Box 63 FI-00014 Helsinki, Finland.

* corresponding author
# equal contribution

Loss of function screening by CRISPR/Cas9 gene knockout with pooled, lentiviral guide libraries is a widely applicable method for systematic identification of genes contributing to diverse cellular phenotypes. Here, random sequence labels (RSLs) are incorporated into the guide library, which act as unique molecular identifiers (UMIs) to allow massively parallel lineage tracing and lineage dropout screening. RSLs greatly improve the reproducibility of results by increasing both the precision and the accuracy of screens. They reduce the number of cells and sequencing reads needed to reach a set statistical power, or allow a more robust screen using the same number of cells.

Pooled CRISPR/Cas9 loss of function screening is a powerful approach to identify genes contributing to a wide range of phenotypes[1]. A library of guide sequences is integrated lentivirally into Cas9-expressing cells, which are then subjected to a selection pressure. Relative guide frequencies in the population before and after selection are quantified by next generation sequencing (NGS) to determine depleted and enriched guides.

The approach has been applied successfully[2-5], but suffers from several shortcomings: First, the presence of a guide does not necessarily cause loss of the corresponding gene, and cells sharing the same guide have distinct genotypes and phenotypes. Second, identification of guides that are under negative selection can be confounded by random drift and undersampling. Third, growth characteristics of individual cells can vary substantially[6, 7] and the site of viral integration can affect the phenotype. For these reasons, each guide needs to be present in a large number of cells. In conventional screens, only the sum of all cells with a specific guide is measured, and no information regarding the distribution of cell behaviors can be obtained. Optimal identification of hit genes would require a method that individually tracks clonal lineages derived from single virus-transduced cells.

Here, we address these issues by incorporating an RSL into the guide-library plasmid (**Fig. 1a**) to allow tracing of thousands of individual virus-transduced cell lineages in a CRISPR screen. Depending on the kinetics of editing, these unique molecular identifiers (UMIs)[8] either trace single clones of identically edited cells, or small populations of sublineages with different editing outcomes at the same locus (**Supplementary Fig. 1a**). Such massively parallel lineage tracing enables both lineage dropout analysis (LDA), and the creation and analysis of internal replicates (IRA), while retaining the option of conventional, total read count analysis (TCA, **Fig. 1b**). Perhaps counterintuitively, analysis of hundreds to thousands of

RSL-labelled cell lineages per guide neither requires more cells per guide, nor deeper sequencing. The RSL
approach simply splits the total guide read count obtained to read counts representing individual
constituent cell lineages, thus increasing the amount of information that is obtained, and consequently
improving both precision and accuracy of the screen.

To demonstrate the power and flexibility of the approach, we screened the human colorectal
carcinoma cell line RKO for essential genes with an RSL-guide library targeting 2325 genes with 10 guides
per gene[5]. For experimental details see **Supplementary Information**. Briefly, Cas9 expressing RKO cells
were transduced with the guide library, and samples were taken at Day 4 and Day 28 after transduction
(control and treatment time points, respectively). Guide frequencies in the two samples were then
assessed by NGS. The experiments were run at far larger screen size (we define "screen size" as the number
of cells per guide sequence) and sequencing depth (reads per guide) than previous screens [4, 5]. Such
redundancy allows subsequent subsampling using RSL information, and robust testing of different
analytical methods at varying screen sizes (**Fig. 1c**).

The plasmid library input contained 78 million unique RSL-guide combinations, 93% of which were
also detected in the virus-transduced cell populations (**Supplementary Fig. 1b**). Based on the Poisson-
distribution, this indicates that about half of the RSL-guides were incorporated into one or two cell
lineages. Because only a subset of the cells can be harvested at each time-point, undersampling is
unavoidable, and some RSL-guides and thus cell lineages were present only in one of the time points
(**Supplementary Fig. 1b, right**). Such undersampling and loss of cell lineages occurs whether or not RSLs
are present, however goes undetected in their absence. With RSLs, the effect becomes apparent and can
be used in quality control of individual experiments as well as in filtering out inconsistently sampled
lineages prior to data analysis.

RSL-labelled, distinguishable guide sequences can be used to split the data into internal replicates,
which in turn allow the usage of classical statistical tools to test for significant differences. To demonstrate
the approach, RSL-guides were binned into 64 internal replicates per guide. The median effect size (**Fig.
1d**) as well as a median-based version of strictly standardized mean difference (SSMD)[9] were then used to
rank the guides (internal replicate analysis using SSMD, IRA/SSMD, **Supplementary Fig. 1c**). In addition,
RSL-labelled guides enable lineage dropout screening, where gene hits are called solely based on the
number of lost RSL-guide lineages (lineage dropout analysis, LDA, **Fig. 1e**).

To evaluate IRA/SSMD and LDA, and to compare them with conventional TCA performed with the
pipeline MAGeCK[10], we assessed the ranks of a set of known essential genes (accuracy), and the hit gene
overlap between experimental replicates (precision). In principle, RSL-based methods should outperform
TCA when the number of cells per guide is relatively low, and their benefit should progressively decrease
as the number of cells per guide approaches infinity. Thus, the comparisons were performed using the
complete dataset, and subsamples of the data that were similar in sample size to published screens [4, 5]
(**Fig. 2**).

Both IRA/SSMD and LDA were more accurate than TCA, as indicated by lower hit ranks of 20 known
essential, ribosomal proteins (**Fig. 2a, Supplementary Fig. 1d**). Both IRA/SSMD and LDA were also more
precise than TCA, with much improved replicate concordance between the top-ranked 5% of genes (**Fig.
2b**). Consistently with the theoretical considerations, our analysis revealed that the RSL-based methods

were far more robust at smaller screen sizes than TCA. Even when the screen size was approximately one
order of magnitude larger, TCA was still inferior to either RSL-based method. At smaller screen size, the
number of highly significant hit genes (FDR < 1%) was massively increased in lineage dropout analysis when
compared to total readcount analysis (**Fig. 2c**).

To summarize, RSLs dramatically improve accuracy, precision, and statistical power in CRISPR/Cas9
screening. The RSL strategy is not limited to CRISPR knockout screening, but can be applied in other
screening methods such as CRISPR-dependent inhibition or activation screens[2, 11]. We expect the RSL
method to become instrumental in the interrogation of small genomic features, e.g. exons, promoters,
and even individual transcription factor binding sites. In many of these cases there is just one possible
guide sequence, and the inclusion of RSLs is the only way to obtain the replicates that are required for hit
calling. In the absence of precise knowledge of both on- and off-target activity, inclusion of multiple guide
positions is however still important, and rescue experiments and/or analysis of the mutational spectrum
of the cutsite are necessary to establish that the mutation induced by the guide results in the observed
phenotype. Incorporation of RSLs is technically straightforward, and does not require a higher number of
cells or sequencing reads compared to conventional approaches. In contrast, RSLs give the same statistical
power at lower number of cells and/or sequence reads, improving the economy of CRISPR/Cas9 screens.
Conversely, RSLs improve accuracy and precision at a given number of cells per guide, which is particularly
advantageous in cases where cell numbers are limiting, such as in primary cells.

Figure Legends.

Figure 1.

a. **Library design. Top.** Guide plasmid. The plasmid library contains an i7 index read primer and an
inert, untranscribed RSL downstream of the tracrRNA. **Bottom.** Sequencing library. Sequencing
was done with a custom primer (Seq) placed directly upstream of the guide (gRNA). The sample
index and RSL were read as two index reads with illumina i5 and i7 index primers, respectively
(20+6+6 sequencing cycles).

b. **RSL guides allow additional methods of analysis.** In Total guide read Count Analysis (TCA, left) RSL
information is ignored and only the sum of readcounts for all RSL-guides is taken into account. In
internal replicate analysis (IRA, middle), readcounts of RSL-guides are binned such that internal
replicates are created for each guide. The example shown bins into four internal replicates. In
lineage dropout analysis (LDA, right) each RSL-guide is monitored separately.

c. **Screen size and sequencing depth.** The screens were performed at a very large screen size of
roughly 4500 cells per guide and sequenced to a depth of 30,000 reads per guide. Using RSL
information, the data from these oversized experiments were then subsampled bioinformatically

to approximately one quarter and one sixteenth, to test different analysis methods at different
screen sizes. The corresponding values for two published screens are indicated for comparison [4, 5]

d. **Internal replicate analysis (IRA).** RSL-guides were binned to create 64 internal replicates. Effect
sizes (log2 fold change in readcount between Day 4 and Day 28 after virus transduction) for each
bin are plotted in ascending order, 10 guides each for MYCN (**top left**) and MYC (**top right**), as well
ass 50 representative non-targeting guides (**bottom**, these non-cutters seem to have a small
fitness advantage). Red dots, median effect size (MES) of the 64 internal replicates; black line, MES
of all guides in the library. Hits for this type of data were called from MES and SSMD
(**Supplementary Figure 1b,** see Supplementary Information for details).

e. **Lineage dropout analysis (LDA).** The average fraction of RSL-guides lost from day 4 to day 28 in
each experimental replicate is plotted for each gene. Red, positive controls; blue non-targeting
controls; black line, linear regression. The number of virus-transduced cell lineages lost is the most
direct readout of the guide effect on cell viability.

Figure 2.

a. **RSLs increase accuracy of hit calling.** Ranks of known positive controls (20 ribosomal proteins out
of a total of 2,335 interrogated genes) in one experimental replicate for the full screen size (left),
as well as one quarter (middle) and 1/16 (right) of the full screen size. Red line, median rank. At all
screen sizes, IR/SSMD analysis and LDA assigned lower ranks to the positive controls than TCA.
The variance of the ranking increased substantially with decreasing screen size in TCA, but not in
the two RSL-based methods.

b. **RSLs increase the precision of gene ranking.** Average percent overlap of the top-ranked 5% of
genes (116 genes) between two experimental replicates. Error bars, standard deviation of four
subsamples. LDA is the most precise method, followed by IRA/SSMD. Again, both RSL-based
methods are superior to TCA and much more robust at smaller screen sizes.

c. **RSLs boost statistical power at small screen size.** Hit gene overlap between experimental
replicates (<1% false discovery rate) at full screen size, one quarter and one sixteenth of the full
screen size. Error bars, standard deviation for hit gene overlap between four subsamples in
experimental replicate 1 and four subsamples in experimental replicate 2 (16 comparisons in
total). Only at full screen size, TCA matches LDA. At more practical screen sizes, LDA has much
higher statistical power and identifies considerably more hit genes.

Accession codes

Read data: European Nucleotide Archive, PRJEB18436. Scripts will be made available on Github under
public license.

Acknowledgements

The authors would like to thank Drs. Inderpreet Kaur Sur, Jenna Persson and Minna Taipale for suggestions
on the manuscript. Part of this work was carried out at Karolinska High Throughput Center (KHTC) and the
High Throughput Genome Engineering Facility (HTGE) funded by SciLifeLab.

Authorship contributions.

B.S., S.K.B. and J.T. developed the approach, B.S., S.K.B. and M.T. performed the experiments, B.S., S.K.B,
J.Z. and T.K. analyzed the data, B.S. and J.T. wrote the manuscript.

References

1.   Shalem, O., Sanjana, N.E. & Zhang, F. High-throughput functional genomics using CRISPR-Cas9. *Nat*
Rev Genet **16**, 299-311 (2015).
2.   Gilbert, L.A. et al. Genome-Scale CRISPR-Mediated Control of Gene Repression and Activation. *Cell*
159, 647-661 (2014).
3.   Koike-Yusa, H., Li, Y., Tan, E.P., Velasco-Herrera Mdel, C. & Yusa, K. Genome-wide recessive genetic
screening in mammalian cells with a lentiviral CRISPR-guide RNA library. *Nat Biotechnol* **32**, 267-
273 (2014).
4.   Shalem, O. et al. Genome-scale CRISPR-Cas9 knockout screening in human cells. *Science* **343**, 84-
87 (2014).
5.   Wang, T. et al. Identification and characterization of essential genes in the human genome. *Science*
350, 1096-1101 (2015).
6.   Levy, S.F. et al. Quantitative evolutionary dynamics using high-resolution lineage tracking. *Nature*
519, 181-186 (2015).
7.   Sandler, O. et al. Lineage correlations of single cell division time as a probe of cell-cycle dynamics.
Nature **519**, 468-471 (2015).
8.   Kivioja, T. et al. Counting absolute numbers of molecules using unique molecular identifiers.
Nature methods **9**, 72-74 (2012).
9.   Zhang, X.D. A pair of new statistical parameters for quality control in RNA interference high-
throughput screening assays. *Genomics* **89**, 552-561 (2007).
10.   Li, W. et al. MAGeCK enables robust identification of essential genes from genome-scale
CRISPR/Cas9 knockout screens. *Genome Biol* **15**, 554 (2014).
11.   Konermann, S. et al. Genome-scale transcriptional activation by an engineered CRISPR-Cas9
complex. *Nature* **517**, 583-588 (2015).

Figure 1

a. Library design.

b. Levels of analysis

c. Screen size and sequencing depth

d. Internal replicate analysis (IRA)

e. Lineage dropout analysis (LDA)

Figure 2

a. RSLs increase accuracy - Ranks of positive controls

b. RSLs increase precision - gene rank overlap

c. RSLs increase statistical power at smaller screen size

Schmierer et al., Supplementary Figure 1

a. Lineage drop out versus lineage depletion

Lineage drop out

Example of lineage depletion

b. Library complexity and carry-through

c. IRA analysis by median effect sized and SSMD

Double flashlight plot

IRA/SSMD Ranking Score

d. RSLs increase accuracy - Ranks of positive controls, replicate 2

Full dataset

1/4th of the dataset

1/16th of the dataset

Supplementary Figure S1.

a. Lineage dropout versus lineage depletion. Depending on the kinetics of editing, single cell lineages harboring a single RSL-guide against an essential gene can either disappear (drop out) or decrease in their abundancy (depletion). **Left.** Dropout happens if the editing occurs early on, either before the cell can divide, or in several independent events at later time points (grey, dead cell; white, unedited cell). **Right**. In lineage depletion, editing occurs either after several cell divisions and/or with several different outcomes, some of which will retain gene function of the essential gene (blue and green edits). The traced lineage is then comprised of several sublineages.

b. **Number of distinct sequences carried through the experiment in one representative experimental replicate. Left.** Boxplot. An average of 3600 RSLs per guide in the plasmid library reduced to an average of 2800 RSLs per guide in the samples taken from the cell populations. Many of these RSLs have very low read counts, those were filtered out and not used for downstream data analysis. **Right.** Venn diagram. Both timepoints together covered 93% of input RSL-guides (78 million unique sequences in the cell population). The overlap between day 4 and day 28 was two thirds, with about one sixth of sequences found either only in Day 4 or only in Day 28. This is a consequence of unavoidable undersampling, which also occurs in the absence of RSLs, however in their presence becomes apparent and allows removal of inconsistently sampled lineages.

c. **Internal replicate analysis by strictly standardized mean difference (IRA/SSMD). Left.** Double flash-light plot. SSMDs for all >23,000 guide sets are plotted against their median effect sizes (MES). Green circles indicate non-targeting control guide sets, yellow circles indicate the 10 guide-sets targeting MYC. **Right.** Ranking score. A hit score was defined as the product of MES and SSMD. This score is negative for depleted guides and positive for enriched guides, and was used for guide ranking. The score was plotted against the median effect size for all guides. Green circles indicate non-targeting control guide sets, yellow circles indicate the 10 guide-sets targeting MYC.

d. **RSLs increase accuracy of hit calling.** As Fig. 2a, but for the second experimental replicate.

Supplementary Information

Oligo synthesis and library cloning. The guide library targeted 2325 genes and contains a total of
23,279 guides. The targeted gene set contains all human transcription factors [1], other genes of interest
as well as ribosomal proteins as positive controls and 101 non-targeting guides as negative controls.
All sgRNA sequences used in this library were taken from a previously published, genome-wide library[2]
(Supplementary file *RSL_guide_library.csv*). The 5' part of the library construct (blue + black, 122bp),
containing the sgRNA was synthesized by oligo array (CustomArray). The 3' part containing the RSL and
the Illumina i7 index primer sequence (green) was synthesized as a single 119 bp oligo
(black+green+red). These two oligos were annealed to each other at the overlapping part (black) and
double stranded by PCR (T_M = 64C) using outer primers (underlined).

GGCTTTATATATCTTGTGGAAAGGACGAAACACCGNNNNNNNNNNNNNNNNNNNNNNgtttAagagctag
aaatagcaagttTaaataaggctagtccgttatcaacttgaaaaagtggcaccgagtcggtgcTTTTT
TgatcggaagagcacacgtctgaactccagtcacNNNNNNaagcttggcgtaactagatcttgagaca
aa

The PCR product was cloned by Gibson assembly into the lentiviral vector pLenti-Puro-AU-flip-
3xBsMBI, which was created by modifying lentiGuide-Puro (a gift from Feng Zhang, Addgene #52963)
by replacing the sequence

gttttagagctagaaatagcaagttaaaataaggctagtccgttatcaacttgaaaaagtggcaccga
gtcggtgcTTTTTT

with

gtttAagagctagaaatagcaagttTaaataaggctagtccgttatcaacttgaaaaagtggcaccga
gtcggtgcTTTTTTcgtctct).

Gibson assembly, transformation and amplification of the library. 100 ng vector and 12 ng insert
where assembled in a total reaction volume of 100 µl (NEBuilder® HiFi DNA Assembly Master Mix,
NEB). The reaction was cleaned via a Minelute reaction cleanup column (Qiagen) and transformed into
6 x 50 µl electrocompetent *E. coli* (Endura™ ElectroCompetent Cells, Lucigen) using a 1.0 mm cuvette,
25µF, 400 Ohms, 1800 Volts. Bacteria were plated on several 24x24 cm agar plates and colonies were
grown overnight. Colonies were scraped into LB medium and the contained plasmids were isolated by
Maxiprep.

Library packaging. The library was packaged in HEK 293T cells by cotransfecting the library plasmid
and the two packaging plasmids psPAX2 (a gift from Didier Trono, Addgene #12260) and pCMV-VSV-G
(a gift from Bob Weinberg, Addgene # 8454) in equimolar ratios. After 48 hours, the virus-containing
supernatant was concentrated 40-fold using Lenti-X concentrator (Clontech), aliquoted for one time
use and stored at -140C.

Cell lines and cell culture. All the cells used in this study were purchased directly from ATCC. Cells were
regularly tested for mycoplasma using the Mycoalert detection kit (Lonza; cat# LT07-218).

Creating editing-proficient Cas9 cell lines. To rapidly generate editing-proficient cell lines, we
synthesized a lentiviral construct (pLenti-Cas9-sgHPRT1) that encodes a codon optimized WT-SpCas9
that is flanked by two nuclear localization signals (derived from lenti-dCAS-VP64_Blast, a gift from Feng
Zhang, Addgene #61425). In addition, the construct codes for blasticidin resistance, and carries an
sgRNA against HPRT1 (GATGTGATGAAGGAGATGGG). HPRT1 loss confers resistance to the

antimetabolite 6-thioguanine (6-TG). Lentivirally transduced cells were selected in 5 µg/ml Blasticidin
and after one week to 10 days additionally with 5µg/ml 6-TG until control cells had died. Only cells that
both express Cas9 and are editing proficient, as indicated by loss of HPRT1 function, will survive. The
method allows rapid establishment of a pool of editing proficient cells. Compared to single cell clones,
this method retains the genetic heterogeneity of the original cell line, avoids potential clonal effects
of the particular integration site of Cas9, and greatly accelerates cell line generation. These benefits
need to be weighed carefully against possible disadvantages, such as synthetic lethality with HPRT1
loss, or potential effects of the presence of a second guide in the cell.
56
57   **Library transduction.** Per replicate, 100 million RKO Cas9 cells were transduced with the library virus.
58   Cells were then selected for guide integration and expression by 1 µg/ml puromycin selection for 48
59   hours. A proportion of cells will contain more than one guide. Because of the vast number of RSL-
60   guides, any ineffective passenger guides will associate with effective guides randomly and will not be
61   significantly enriched or depleted in the population.
62
63   **Cell propagation and sample preparation.** Cells were kept in culture for a total of 28 days after
64   transduction by sub-culturing them every three to four days. 100 million cells were reseeded at each
65   split, and genomic DNA was prepared from 50 – 80 million cells at Days 4 and 28 after transduction.
66   Day 4 after transduction was considered the control time point.
67
Preparation of the sequencing library from genomic DNA. The sequencing library preparation consists
of 3 PCR steps, PCR1 amplifies the genomic region containing the guide sequence using the primers 1F
and 1R. PCR2 and PCR3 then incorporate the Illumina adaptors with primers 2F/2R and 3F/3R,
respectively. 3F contains the Illumina index for multiplexing, indicated by NNNNNN in the sequence
given.

Genomic DNA was isolated using Blood and Tissue Maxi Kit (Qiagen), and 200 µg, theoretically
corresponding to 30 million diploid cells, were used as PCR template in 40 parallel PCR reactions (5 µg
template DNA each) using KAPA HiFi HotStart polymerase (KAPA Biosystems). After 14 cycles, the
reactions were pooled. PCR2 used 5 µl of pooled PCR1 as template and was run for 19 cycles, PCR3
used 2µl of PCR2 as template and was run for 14 cycles. The resulting product of 288 bp was gel purified
and sequenced on an Illumina HiSeq 4000 instrument using single read 20 cycles plus two 6 bp index
reads, where index read 1 reads the RSL and index read 2 reads the Illumina sample index.
80
1fw   GGACTATCATATGCTTACCGTAACTTGAAAGTATTTCG
1ref   CTTTAGTTTGTATGTCTGTTGCTATTATGTCTACTATTCTTTCC
2fw   TCTTTCCCTACACGACGCTCTTCCGATCtcttgtggaaaggacgaaacac
2rev   AGAAGACGGCATACGAGATctgccatttgtctcaagatctagttac
3fw   AATGATACGGCGACCACCGAGATCTACAC NNNNNN TCTTTCCCTACACGACGCTCTTCCG
3rev   CAAGCagaagacggcatacgagatctgccatttg
87
88   The final library product was sequenced with a custom primer and the i5 and i7 index primers
89   (underlined) by running 20+6+6 cycles on the Illumina HiSeq4000.
90
91   <u>AATGATACGGCGACCACCGAGATCTACAC</u>[i5]**NNNNNN**TCTTTCCCTACACGACGCTCTTC<u>CGATCt</u>
92   <u>cttgtggaaaggacgaaacac</u>CG**NNNNNNNNNNNNNNNNNNNNNNNN**gtttAagagctagaaatagcaagtt
93   TaaataaGgctagtccgttatcaacttgaaaaagtggcaccgagtcggtgcTTTTTT<u>gatcggaagag</u>
94   <u>cacacgtctgaactccagtcac</u>[i7]**BBBBBB**aagcttggcgtaactagatcttgagacaaatggcag
95   ATCTCGTATGCCGTCTTCTGCTTG
96
97

Scripts used for counting RSL-guides and for binning. RSL-guides were counted in the original fastq
files with the Perl scripts *BatchRun-pub2.pl*, which requires the script *GuideUMI_pub2.pl*. Binning of
RSL-guide counts was done using the script *countTruncatedRSLs.pl*. Generally, sequences whose total
readcount in control and treatment was less than five were filtered out prior to data analysis.

SSMD analysis of read count data.

Normalization. In RNASeq, methods such as median normalization are commonly preferred to total
read-count normalization, mainly to compensate for the effect of a few very highly expressed genes
that can take up a significant proportion of the total read count. CRISPR/Cas9 screening data are
comparably well balanced and we thus chose the most basic normalisation method, total read count
normalisation, to compensate for different sequencing depths. c_{ij} and t_{ij} represent the raw read
counts for RSL-guide j in guide-set i for control (Day 4 after lentiviral transduction) and treatment (Day
28 after lentiviral transduction), respectively. The normalised read counts c'_{ij} and t'_{ij} are then
111

112   $$c'_{ij} = c_{ij} \frac{\sum_{ij}(c_{ij} + t_{ij})}{2 \sum_{ij} c_{ij}}$$

114   $$t'_{ij} = t_{ij} \frac{\sum_{ij}(c_{ij} + t_{ij})}{2 \sum_{ij} t_{ij}}$$

116   **Median effect size and variability of the guide-sets.** We defined the effect size ES_{ij} for each RSL-guide
117   j in guide-set i as the log2 of the fold change between treatment count and control count. To handle
118   total loss of an RSL-guide in the treatment sample, we added a pseudo-count of 1 to all counts:
119

120   $$ES_{ij} = log_2 \frac{t'_{ij} + 1}{c'_{ij} + 1}$$

122   Next, we calculated the median effect size for guide set *i*, MES_i, and the median of the absolute
123   deviations (MAD) of all RSL-guides j in guide-set i from MES_i
124

$$MES_i = \underset{j}{median}\, ES_{ij}$$

$$MAD_i = 1.4826\, \underset{j}{median}|ES_{ij} - MES_i|$$

The factor 1.4826 was chosen such that the MAD is approximately equal to the standard deviation
under the assumption of normal distribution[3].

Median effect size and variability of the control guide-sets. The RSL library contains 101 non-targeting
guide-sets. We calculate a single median effect size and MAD for this control set in the following way:

Median effect size of all non-targeting RSL-guides

$$MES_{CON} = \underset{ij}{median}\, ES_{ij}^{NONT}$$

Median absolute deviation of all non-targeting RSL-guides:

$$MAD_{CON} = 1.4826\, \underset{ij}{median}\, |ES_{ij}^{NONT} - MES_{CON}|$$

Strictly standardized mean difference (SSMD). SSMD is a measure for the significance of the difference in behaviour of sample *i* and the non-targeting controls. It takes into account both the effect size and the variability of the data.

$$SSMD_i = \frac{MES_i - MES_{CON}}{\sqrt{MAD_i^2 + MAD_{CON}^2}}$$

For samples with relatively small effect size, the SSMD can still become large if the spread is small. We thus introduce a score in which the effect size weighs more strongly, and which is used as a ranking parameter:

$$Score_i = MES_i |SSMD_i|$$

For hit calling, the average score and standard deviation were calculated for all non-targeting guide sets. The script used in these calculations is *SSMD.sh*, which calls the script R-script *SSMD.R*. Guide-sets were then ranked according to their score and the resulting ranked list was analysed with α-RRA, a robust rank aggregation algorithm as implemented in the "pathway" function of MAGeCK[4, 5] using Supplementary file *RSL_guide_library.gmt.*

Lineage dropout.

An RSL-guide was considered a dropout if it had less than two readcounts in the treatment time point. The numbers of RSLs per guide at Day 4 and Day 28 were then used as input in MAGeCK to obtain a ranked gene list and FDRs.

Subsampling.

For subsampling the full data set, RSL-guides were grouped according to their RSL-sequence. For medium screen size, the whole dataset was split into four groups (RSLs starting with A,C,G and T). For small screen size, the whole dataset was split into 16 groups, the first four of which (AA,AC,AG,AT) were used for analysis.

Supplementary References

1. Vaquerizas, J.M., Kummerfeld, S.K., Teichmann, S.A. & Luscombe, N.M. A census of human transcription factors: function, expression and evolution. *Nat Rev Genet* **10**, 252-263 (2009).
2. Wang, T., Wei, J.J., Sabatini, D.M. & Lander, E.S. Genetic screens in human cells using the CRISPR-Cas9 system. *Science* **343**, 80-84 (2014).
3. Zhang, X.D. Illustration of SSMD, z score, SSMD*, z* score, and t statistic for hit selection in RNAi high-throughput screens. *J Biomol Screen* **16**, 775-785 (2011).
4. Kolde, R., Laur, S., Adler, P. & Vilo, J. Robust rank aggregation for gene list integration and meta-analysis. *Bioinformatics* **28**, 573-580 (2012).
5. Li, W. et al. MAGeCK enables robust identification of essential genes from genome-scale CRISPR/Cas9 knockout screens. *Genome Biol* **15**, 554 (2014).