Alphaproteobacteria fix nitrogen in a *Sphagnum*-dominated peat bog using molybdenum-

dependent nitrogenase

Melissa J. Warren[1*], Xueju Lin[2**], John C. Gaby[2***], Cecilia B. Kretz[1****], Peter L. Morton[3],

Jennifer Pett-Ridge[4], David J. Weston[5], Christopher W. Schadt[5], Joel E. Kostka[1,2], Jennifer B.

Glass[1,2#]

[1]School of Earth and Atmospheric Sciences, Georgia Institute of Technology, Atlanta, GA, USA

[2]School of Biology, Georgia Institute of Technology, Atlanta, GA, USA

[3]Department of Earth, Ocean and Atmospheric Science, Florida State University, Tallahassee,

Florida, USA

[4]Physical and Life Sciences Directorate, Lawrence Livermore National Laboratory, Livermore,

CA, USA

[5]Biosciences Division, Oak Ridge National Laboratory, Oak Ridge, TN, USA

Now at: *CH2M, Atlanta, GA, USA; **InstantLabs, Baltimore, MD, USA; ***Faculty of

Chemistry, Biotechnology and Food Science, Norwegian University of Life Sciences, Oslo,

Norway; ****Division of Bacterial Diseases, National Center for Immunization and Respiratory

Diseases, Centers for Disease Control and Prevention, Atlanta, GA, USA

#Corresponding author: Jennifer B. Glass, 404-894-3942; jennifer.glass@eas.gatech.edu

20
Running Title: Molybdenum-based nitrogen fixation in a *Sphagnum* peat bog

Abstract

Microbial N_2 fixation (diazotrophy) represents an important nitrogen source to oligotrophic

peatland ecosystems, which are important sinks for atmospheric CO_2 and susceptible to changing

climate. The objective of this study was to investigate the pathways and controls of diazotrophy,

as well as the active microbial groups that mediate the process in *Sphagnum*-dominated peat

bogs. In an ombrotrophic peat bog (S1) in the Marcell Experimental Forest (Minnesota, USA),

low (μM) levels of inorganic nitrogen were observed, suggesting that diazotrophy could have a

significant influence on ecosystem properties. Despite higher dissolved vanadium (V) (11 nM)

than molybdenum (Mo) (3 nM) in surface peat, a combination of metagenomic, amplicon

sequencing and activity measurements indicated that Mo-containing nitrogenases dominate over

the V-containing form. Diazotrophy was only detected in surface peat exposed to light, with the

highest rates observed in peat collected from hollows with the highest water content. Rates were

suppressed by O_2, and unaffected by CH_4 and CO_2 amendments. Acetylene fully inhibited CH_4

consumption under oxic conditions, but only partially inhibited $^{15}N_2$ incorporation in degassed

incubations, and had a minimal effect on oxic $^{15}N_2$ incorporation. Through a close coupling of

process rate measurements with molecular analysis of the metabolically active microbial

communities, our findings suggest that diazotrophy in surface layers of the S1 bog is mediated

by *Alphaproteobacteria* (*Bradyrhizobiaceae* and *Beijerinckiaceae*) supported by photosynthate,

rather than methane, for carbon and/or energy.

Importance

Previous studies indicate that diazotrophy provides an important nitrogen source and is linked to

methanotrophy in *Sphagnum*-dominated peatlands. However, the environmental controls and

enzymatic pathways of diazotrophy, as well as the metabolically active microbial populations

that catalyze this process in peatlands, remain in question. Our findings indicate that oxygen

levels and photosynthetic activity override low nutrient availability in limiting diazotrophy, and

that primarily non-methanotrophic members of the *Alphaproteobacteria* (*Bradyrhizobiaceae* and

Beijerinckiaceae) catalyze this process primarily at the bog surface.

Introduction

High-latitude peatlands store approximately one-third of global soil carbon and may pose a climatic threat if rising global temperatures accelerate the release of this stored carbon in gaseous forms as either carbon dioxide or methane (32, 59, 94). Mineral-poor (ombrotrophic) peatlands receive most of their nutrient inputs from atmospheric deposition and contain *Sphagnum* moss as their primary plant cover (11, 59). The peatmoss *Sphagnum* is a keystone genus in these ecosystems with its biological function and abiotic environment responsible for much of the primary production and recalcitrant dead organic matter (12, 92). *Sphagnum* mosses also host complex microbiomes (8, 53, 69, 70), including N_2-fixers (diazotrophs) that are significant nitrogen sources to peatland ecosystems (4).

Despite decades of research, there is still much debate on the identity of the dominant diazotrophs in ombrotrophic peatlands. Early work implicated Cyanobacteria (1, 33, 34) or heterotrophic bacteria (78) based primarily on microscopic studies, while more recent molecular analyses argue for the importance of methanotrophic *Beijerinckiaceae* (18) as major diazotrophs in *Sphagnum* peat bogs (7, 21, 40, 89). Possible contributions from other potential diazotrophs, such as strictly anaerobic methanogenic *Euryarchaeaota*, remain unknown. However, it is quite possible that diverse diazotrophs exist within defined niches of peatland environments (55).

Diazotrophy is catalyzed by the nitrogenase metalloenzyme, a complex of thee subunits (H, D and K) that contains abundant iron as Fe-S clusters. This enzyme is extremely O_2 sensitive (93), and must be protected from exposure to O_2 for diazotrophy to occur (24). The most common form of nitrogenase, encoded by *nif* genes, contains molybdenum (Mo) as its cofactor. When Mo is scarce, some species of Bacteria and Archaea express nitrogenases containing vanadium (V; *vnf* genes) or iron (Fe; *anf* genes) in place of Mo, but these "alternative"

nitrogenases are less efficient than the Mo form (67, 90). Phylogenetic analyses of the most

conserved nitrogenase gene, *nifH*, revealed four clusters: aerobic bacteria (cluster I); alternative

nitrogenases (cluster II); anaerobic bacteria and archaea (cluster III); uncharacterized sequences

(cluster IV), and paralogs related to chlorophyll biosynthesis (cluster V; 76). Consistent with

higher concentrations of V than Mo in most rocks (91), microbes from diverse soils contain *vnf*

(cluster II) genes (3, 6, 16, 41, 65). Given the oligotrophic conditions that predominate in

peatlands, trace metals may limit diazotrophy. However, little is known about trace metal

availability and the role of alternative nitrogenase pathways in ombrotrophic peatlands.

Similarly and importantly, methane monooxygenase (MMO) (the enzyme that catalyzes

the first step of methane oxidation) occurs in particulate (copper (Cu)-containing pMMO) and

soluble (Fe-containing sMMO) forms. While pMMO has more specific substrate requirements,

pathways that employ sMMO can use a wider range of compounds (14). Both forms of MMO

are inhibited by acetylene (C_2H_2) (13, 73). In organisms with both sets of genes, pMMO is

expressed when Cu is abundant, whereas Cu limitation induces sMMO expression (80). The

dominant peatland methanotrophs in the *Alphaproteobacteria* and *Gammaproteobacteria* tend to

possess both MMOs (10, 23, 35, 47, 63), although *Methylocella* species containing solely sMMO

have been isolated from peat bogs (19, 20). While most studies have primarily targeted the *pmoA*

gene (23, 47), *mmoX* genes and transcripts have also been reported in peatlands (58, 63, 75),

raising questions about the relative importance of each form for peatland methane oxidation.

The acetylene reduction assay (ARA) is commonly used as a proxy for diazotroph

activity (37, 38). This assay is effective for capturing the potential activity of diazotrophic

bacteria that are not inhibited by C_2H_2, such as *Cyanobacteria* and non-methanotrophic

Proteobacteria (e.g. *Bradyrhizobiaceae*) (45). However, a number of functional guilds of

microorganisms (including methanotrophs, methanogens, sulfate reducers, and nitrifiers) are

inhibited by C_2H_2 (15, 44, 71, 72, 82, 84). If these or other C_2H_2-sensitive microbes (see

Fulweiler, et al. (25)) perform diazotrophy and/or provide substrates to other diazotrophs, ARA

may underestimate N_2 fixation activity in that system. Thus, recent studies have shifted to

tracking N_2 fixation by incorporation of the stable isotope tracer, $^{15}N_2$ (50, 55, 56, 89).

In this study of the S1 peat bog at the Marcell Experimental Forest in Minnesota, USA,

dissolved macro- (NH_4^+, NO_3^-, PO_4^{3-}) and micro- (Fe, Cu, V, Mo) nutrients were profiled along

with the community composition and abundance of diazotrophic microorganisms. We also

performed separate laboratory incubation experiments to measure potential rates of ARA and

$^{15}N_2$ incorporation to: (a) assess environmental controls (light, O_2, CH_4) on diazotrophy; (b)

quantify the effect of C_2H_2 on rates of diazotrophy and methanotrophy; and (c) search for

diagnostic markers for alternative nitrogenase activity such as a low conversion factor of ARA to

$^{15}N_2$-incorporation (3) and C_2H_2 reduction to ethane (22). Finally, we make recommendations on

universal *nifH* primers for amplicon sequencing and quantitative PCR based on our findings.

Materials and Methods

Site description and sample collection. Samples were collected from the S1 (black spruce-

Sphagnum spp.) peat bog at Marcell Experimental Forest (MEF; 47°30.476' N; 93°27.162' W),

the site of the DOE SPRUCE (Spruce and Peatland Responses Under Climatic and

Environmental Change) experiment in northern Minnesota, USA (36). The S1 bog is

ombrotrophic and acidic (average pH 3.5-4; 52, 79). Dissolved O_2 levels decrease to below

detection (~20 ppb) within the top 5 cm of the bog. Three locations were sampled along S1 bog

transect 3 (T3) at near, middle and far sites (see Lin, et al. (64) for further details). Surface (0-10

cm depth) peat was collected from hollows dominated by a mixture of *Sphagnum fallax* and *S.*

angustifolium, and hummocks dominated by *S. magellanicum*. Peat depth cores (0-200 cm) were

sampled from hollows where the water level reached the surface of the *Sphagnum* layer.

Macronutrients. Peat porewater was collected using piezometers from 0, 10, 25, 50, 75,

150 and 200 cm depth. Piezometers were recharged the same day as collection, and porewater

was pumped to the surface, filtered through sterile 0.2 μm polyethersulfone membrane filters,

and stored frozen until analysis. Nitrate (NO_3^-) and nitrite (NO_2^-) were analyzed using the

spectrophotometric assay as described by García-Robledo, et al. (28). Ammonium (NH_4^+)

concentrations were determined with the indophenol blue assay (85). Phosphate concentrations

were measured with the molybdate-antimony ascorbic acid colorimetric assay (68).

Micronutrients. Peat porewater was collected from two locations in the S1 bog from

cores at 0-30 cm, 30-50 cm, and 100-150 cm depths by filtration through 0.15 μm Rhizon soil

samplers (Rhizosphere Research Products). All plastics were HCl washed prior to sampling;

Rhizon soil samplers were cleaned by pumping 10 mL of 1 N HCl through them, followed by a

rinsing with ultrapure water until the pH returned to neutral (~100 mL/filter). After collection,

samples were acidified to 0.32 M HNO_3 (Fisher Optima) and analyzed using a Thermo

ELEMENT 2 HR-ICP-MS (National High Magnetic Field Laboratory, Florida State University).

Initial analyses resulted in frequent clogging of the nebulizer, likely due to abundant dissolved

organic carbon. Therefore, samples were diluted 1:10 to minimize interruptions from nebulizer

clogs. Concentrations were quantified with a 7-point external calibration using standards

prepared in 0.32 N HNO_3 from a multi-element standard mix (High Purity Standards). In order to

measure lower concentrations but avoiding sample dilution, subsequent samples were digested

before ICP-MS analysis. Briefly, 1 mL porewater samples were digested for 36 hours at 250°C

in a trace metal clean polypropylene exhausted laminar flow hood with 1 mL of 16 N NHO$_3$

(Ultrex II, JT Baker) and 100 µL of 30% H$_2$O$_2$ (Ultrex II, JT Baker) in acid-washed Teflon vials.

Samples were then evaporated to near dryness, resuspended in 0.32 N HNO$_3$, and analyzed by

ELEMENT2 ICP-MS along with parallel blank solutions.

Quantification and sequencing of gene and transcript amplicons. Peat was frozen on

dry ice at the field site in July 2013, or in liquid N$_2$ after 7 days at 25°C in the light under

degassed (80% N$_2$ + 20% CO$_2$) headspace with 1% C$_2$H$_2$, with or without 1% CH$_4$ for June 2014

incubations (see ARA section for more details). DNA and RNA extractions were performed with

MoBio PowerSoil DNA and total RNA extraction kits, respectively, as described in Lin, et al.

(63). cDNA was prepared by reverse transcription as in Esson, et al. (23). Primer pairs were

polF/polR for *nifH* (25), A189f/Mb661r for *pmoA* (51), mmoX-206f/mmoX-886r for *mmoX* (43),

Mlas/mcrA-rev for *mcrA* (83), nifD820F/nifD1389R followed by nifD820F/nifD1331R for *nifD*

(3)*,* and vnfD_anfD548F/vnfD_anfD1337R followed by vnfD_anfD548F/vnfD_anfD1291R for

vnfD/anfD (3). PCR conditions were based on (27), (23), (58), (83), (3) and (9) for *nifH, pmoA,*

mmoX, mcrA, nifD/vnfD/anfD and 16S rRNA, respectively.

Gene fragments for qPCR were amplified from genomic DNA and plasmid standards

were constructed according to Lin, et al. (61). To prepare cDNA standards, plasmid DNA with a

positive gene insert was linearized with NcoI restriction enzyme following the manufacturer's

protocol (Promega), and purified by MinElute PCR purification kit (Qiagen). RNA was

synthesized from the linearized plasmid DNA using the Riboprobe in vitro transcription system

according to the manufacturer's protocol (Promega). Then, cDNA standards and cDNA of

environmental RNA samples were synthesized using the GoScript Reverse Transcription system

following the manufacturer's protocol (Promega). Samples were run in duplicate against a

plasmid DNA standard curve on a StepOnePlus Real-Time PCR System (ABI) using Power

SYBR Green PCR master mix. Functional gene and transcript copy numbers were normalized to

dry weight of peat as well as 16S rRNA transcript copies for incubation samples. Amplicons

were sent to the University of Illinois at Chicago for DNA sequencing using a 454 platform. Raw

sequences were demultiplexed, trimmed, and quality filtered in CLCbio. The phylogeny of

vnfD/anfD sequences was inferred using the Maximum Likelihood method based on the Kimura

2-parameter model in MEGA5 (86).

Acetylene reduction and methane consumption rates. Samples of bulk peat (*Sphagnum*

spp. and surrounding soil) was collected from 0-10 and 10-30 cm depths in September 2013,

April 2014, June 2014, September 2014 and August 2015, and stored at 4°C until the start of

laboratory incubations. Samples from 0-10 cm depth were gently homogenized so as not to

rupture *Sphagnum* spp. tissues, while peat samples from 10-30 cm depth were fully

homogenized. For each sample, 5 g of bulk peat were placed in 70 mL glass serum bottles,

stoppered with black butyl stoppers (Geo-Microbial Technologies) and sealed with an aluminum

crimp seal. Headspaces were oxic (room air, 80% N_2 + 20% O_2) or degassed (100% N_2 or 80%

N_2 + 20% CO_2) with or without 1% C_2H_2 or 1% CH_4. Treatments were incubated for one week at

25°C in the light or dark. A gas chromatograph with a flame ionization detector (SRI

Instruments, Torrance, CA, USA) equipped with a HayeSep N column was used to quantify CH_4,

C_2H_2 and C_2H_4. Samples were measured for C_2H_4 production daily until C_2H_4 production was

linear (~7 days). Controls not amended with C_2H_2 did not produce ethylene (C_2H_4). Incubations

of hollow peat from June 2014 incubated in oxic headspace with and without 1% C_2H_2 were also

monitored for consumption of 1% CH_4. Statistical analysis was performed with JMP Pro (v.

12.1.0) using the Student's T-test.

^{15}N_2 incorporation rates. In September 2014, samples were quantified for N_2 fixation

rates by $^{15}N_2$ incorporation in parallel with ARA measurements. Incubations were set up as

described above and supplemented with 7 mL of 98% $^{15}N_2$ (Cambridge Isotope Laboratories,

Tewksbury, MA, USA). After 7 days, samples were dried at 80°C, homogenized into a fine

powder, and analyzed for N content and $\delta^{15}N$ by isotope-ratio mass spectrometry (IRMS) with a

MICRO cube elemental analyzer and IsoPrime100 IRMS (Elementar) at the University of

California, Berkeley, corrected relative to National Institute of Standards and Technology (NIST,

Gaithersburg, MD, USA) standards.

Metagenomic analyses. Metagenomes were generated in a previous study (62).

Diazotrophic and methanotrophic pathways were interrogated using the following bioinformatics

approaches. Briefly, Illumina reads were filtered by quality (Phred33 score threshold of Q25)

using Trim Galore (Babraham Bioinformatics) and a minimum sequence length cut off of 100

bp. The sequences were then queried using RAPSearch2 (95) against the NCBI-nr database of

non-redundant protein sequences as of November 2013. Sequences with a bit-score of 50 and

higher were retained to determine the total number of functional genes for normalization across

the different samples. The taxonomic composition of protein-coding sequences was determined

based on the taxonomic annotation of each gene according to the NCBI-nr taxonomy in

MEGAN5 (42) (min score: 50; max expected: 0.01; top percent: 10; min complexity: 0.3).

To classify sequences by nitrogenase cluster type, genes were analyzed using BLASTX

(e-value 0.1; bit-score 50) versus a custom *nifH* database that includes a phylogenetic tree to

distinguish the principal clusters (I, V, III) in the *nifH* phylogeny, as well as paralogous cluster

IV, *nifH*-like sequences (26). *nifH* genes from the four clusters were normalized to total protein-

coding genes from RAPSearch2 output sequences. The relative abundance of particulate vs.

soluble methane monooxygenase (*pmoA* vs. *mmoX*, respectively) was based on previous analyses

in Lin, et al. (63).

Nucleotide sequence accession numbers. Metagenomes were reported in a previous

study (62) and deposited in BioProject XXXXXXX. *pmoA* cDNA amplicons were reported in a

previous study (23) and deposited in BioProject PRJNA311735. *nifH, mcrA, nifD,* and

vnfD/anfD cDNA amplicons were deposited in BioProject XXXXXXX.

Results

Macro/micro-nutrients and functional genes. In S1 bog hollows, NH_4^+ remained at 2 μM from

the surface to 25 cm depth, and increased at greater depths (Fig. 1; Fig. S1a). Nitrate was <1 μM

in surface peat and decreased with depth (Fig. S1b). Phosphate was <0.1 μM from the surface to

25 cm depth, and then increased with depth (Fig. S1c). With the polF/polR primer pair, we

measured 1.2×10^7 copies g^{-1} *nifH* genes at 1 and 20 cm, and 0.2×10^7 copies g^{-1} at 30 and 75 cm;

nifH transcripts (12.2×10^7 copies g^{-1}) were only detected at 1 cm (a 10:1 transcript: gene ratio),

and not at deeper depths (Fig. 1; Fig. S2a). Sequencing of cDNA from surface peat amplified

with polF/polR (*nifH*) and nifD820F/nifD1331R (*nifD*) primers showed that the vast majority of

nitrogenase transcripts belonged to cluster I (*Alphaproteobacteria*), with *Bradyrhizobiaceae*

dominating *nifH* sequences and *Beijerinckiaceae* dominating *nifD* sequences (Fig. S2). *nifH*

transcripts from *Gammaproteobacteria*, *Cyanobacteria* (*Oscillatoriophycideae*) and *Nitrospira*

were also observed at lower abundance in cDNA amplicon libraries (data not shown). Like *nifH*,

pmoA and *mcrA* transcripts showed the highest abundance in surface peat (Fig. S2b,c). Surface

pmoA transcripts mapped to *Methylocystaceae* (75%) and *Methylococcaceae* (25%). Surface

mcrA transcripts mapped to *Methanosarcina* (58%), *Methanocella* (28%), and *Methanoregula*

(11%). Attempts to amplify *mmoX* from cDNA were unsuccessful (data not shown).

Vanadium (5-21 nM) was consistently more abundant than Mo (1-7 nM) at 0-30, 30-50

and 100-150 cm depth intervals (Fig. 2). In metagenomes, *nifH* genes were roughly equally

distributed between Mo-dependent clusters I and III, and cluster IV/V paralogs (Fig. 2; Table

S1). Sequences from Cluster II (alternative nitrogenases, *vnfH/anfH*) were scarce at all depths

(<5% overall); two *vnfD* genes from metagenomes showed phylogenetic similarity to those from

soil *Proteobacteria* (Fig. S3). Attempts to amplify *vnfD/anfD* from cDNA yielded few reads;

those recovered were most similar to *Alphaproteobacteria anfD* from *Rhodopseudomonas*

species (Fig. S4). Iron (7-35 μM) was three orders of magnitude higher than Cu (7-38 nM; Fig.

3). Trace nutrients Co (5-20 nM), Ni (10-80 nM), Zn (50-250 nM) and Mn (60-2220 nM) were

also quantified (data not shown). In metagenomes, genes for *pmoA* were dominant in surface

peat, whereas *mmoX* sequences became more abundant with depth (Fig. 3). In hollow surface

waters, other dissolved metals analyzed included Mn (~500 nM), Ni (~30 nM) and Co (~10 nM).

Rates of diazotrophy and methanotrophy. Potential rates of acetylene reduction were

measured for peat collected from S1 bog hollows and hummocks in April, June, August and

September 2013-2015 and incubated for 1 week at 25°C. Acetylene reduction to ethylene was

only detected in surface (0-10 cm) peat samples incubated in the light, and not in deep (10-30

cm) peat, nor in surface peat incubated in the dark; acetylene reduction to ethane was not

detected in any incubation (data not shown). *Sphagnum* in peat incubations exposed to light

became visibly greener over the course of the incubation (Fig. 4). In hollows, where surface peat

was dominantly covered by a mix of *S. fallax* and *S. angustifolium*, ARA rates were higher and

more variable in degassed in comparison to oxic incubations (0-163 vs. 2-23 μmol C_2H_4 g^{-1} h^{-1}),

and were unaffected by the presence or absence of 20% CO_2. *nifH* transcripts were also higher

and more variable in hollows than hummocks (Fig. S5). ARA in hollows incubated with

degassed headspace was positively correlated ($P < 0.0001$) with peat water content (93 to 96%).

In hummocks with lower water content (90-91%), surface peat was dominantly covered by *S.*

magellanicum, and oxic and degassed treatments had similarly low ARA rates (0-8 μmol C_2H_4 g^{-}

^{1} h^{-1}). In both hollows and hummocks, ARA rates were not affected by addition of 1% CH_4.

$^{15}N_2$ incorporation showed similar overall trends as ARA (e.g. >90% higher rates in

degassed vs. oxic conditions, and no significant CH_4 effect; Fig. 5). Using the four sites

measured with both methods, a conversion factor of 3.9 for $^{15}N_2$-to-ARA was calculated (Fig.

S6). In degassed treatments, 1% C_2H_2 inhibited $^{15}N_2$ incorporation by 55% but had no effect on

oxic treatments. In oxic treatments, CH_4 consumption rates were 100 times higher than $^{15}N_2$

incorporation rates, and 1% C_2H_2 addition suppressed CH_4 oxidation rates by 95% (Fig. 5). *nifH*

transcripts in surface peat from hollows incubated under degassed headspace with 1% C_2H_2, with

or without 1% CH_4, ranged from 10^4 - 10^7 copies g^{-1} (July 2014; data not shown), which was 1-4

orders of magnitude lower than field samples from the previous summer (10^8 copies g^{-1}; Fig. 1).

Discussion

Peat bog nutrient concentrations and metalloenzymes. By definition, the only source of

nutrients to ombrotrophic peat bogs is the atmosphere, and thus these ecosystems are well known

to be oligotrophic (60). Scarce macronutrients in S1 peat pore water and low rates of atmospheric

deposition in Minnesota peat bogs (39) are consistent with oligotrophic conditions (60, 79, 87).

Micronutrient concentrations (~15 μM Fe, ~1 μM Zn and nM-level Ni, Cu, V, Mo and Co)

reported for acidic peatlands in Northern Europe (48) were similar to those we measured in the

S1 bog. Molybdenum sorption to peat is enhanced at low pH (5), and Mo concentrations (<10

nM) here were within the same range as in other freshwaters worldwide (30). In laboratory

cultures, <10 nM Mo can either limit growth (31) or induce expression of alternative nitrogenase

(2). Although S1 peat bog waters contained higher V (11 nM) than Mo (3 nM), the majority of

nitrogenase H-subunit genes retrieved from metagenomes belonged to Mo-containing clusters I

and III. A significant number of sequences was also obtained from uncharacterized Cluster IV,

recently shown to contain a functional nitrogenase (96) that likely binds a Mo-Fe cofactor (66).

Cluster I (*Alphaproteobacteria*; *Rhizobiales*) showed the highest relative abundance in transcript

libraries, with *Bradyrhizobiaceae* and *Beijerinckiaceae* dominating *nifD* and *nifH* amplicons,

respectively, as seen in prior studies (58, 89). Overall, evidence for alternative nitrogenases was

scant: (a) very few cluster II *vnfD/anfD* genes were found in metagenomes; (b) minimal

vnfD/anfD amplification was obtained from peat cDNA; (c) ethane was undetectable in ARA

incubations; and (d) the $^{15}N_2$-to-ARA conversion factor (3.9) – within the same range (3-4) as

other peat bogs (37, 57, 89) – matched Mo-nitrogenase in pure culture experiments (3), as

opposed to the lower values measured for alternative nitrogenases. Thus, in spite of lower

abundance of Mo than V in S1 bog porewaters, our collective evidence strongly suggests N_2

fixation by the Mo-containing form of the nitrogenase enzyme.

Like diazotrophy, methanotrophy was apparently also mediated by the enzyme requiring

the scarcer metal; dissolved Fe was consistently orders of magnitude higher than Cu in peat

porewater, yet *pmoA* sequences were more abundant than *mmoX* sequences in surface peat in

which the highest CH_4 consumption was observed (23, 63). Overall, these findings suggest that

trace metal availability is not the overriding environmental pressure driving selection for

diazotrophs and methanotrophs in *Sphagnum*-dominated peat bogs. Rather, water potential and

oxygen appear to be more important. Low metal and phosphate concentrations add further

evidence to previous suggestions that these nutrients may limit *Sphagnum* productivity (55)

and/or complex mechanisms may exist for nutrient scavenging at ultra-low concentrations.

Additionally, the inherent nature of the peat matrix, with characteristically high levels of

particulate and dissolved organic matter, likely also affects metal bioavailability in complex

ways not addressed in our study (88).

Environmental controls on diazotrophy. Methanotrophs have the potential to serve as a

biofilter to suppress methane emissions, along with providing a nitrogen source for the dominant

primary producers in peatland ecosystems. Previous field and laboratory studies point to a close

coupling of diazotrophy and methanotrophy in *Sphagnum*-dominated peatlands (49, 55, 74, 89).

This work has led others to suggest that methanotrophs provide the unaccounted nitrogen input

resulting from an imbalance in atmospheric nitrogen deposition and accumulation in *Sphagnum*

mosses (40). However, while substantial evidence indicates that methanotrophs are a source of

nitrogen to peat mosses, the ecological relevance of diazotrophic methanotrophs and their

importance to nitrogen flow still remains unclear because the two Alphaproteobacteria families

that appear to be the dominant diazotrophs in *Sphagnum* peat bogs grow on complex organics

(*Bradyrhizobiaceae* (29)) as well as simple alkanes and C1 compounds (*Beijerinckiaceae* (17)).

Our results from laboratory incubations of native peats revealed that diazotrophy was

stimulated by light, suppressed by O_2, and minimally affected by CH_4 and CO_2. Consistent with

previous reports (77), ARA and *nifH* transcription were only detected in surface samples

incubated in the light, although it is possible that our methods excluded C_2H_2-sensitive

diazotrophs at deeper depths (50). *nifH* transcripts were 1-4 orders of magnitude lower in

laboratory incubations than in the field, suggesting either that the diazotrophs in our incubations

were stressed (e.g. nutrient-limited), and/or that a different assemblage of diazotrophs grew in

the incubations than in the field due to primer specificity (see next section).

Complete CH_4 consumption in air with 1% CH_4 showed that methanotrophs were active

in our peat incubations. If these aerobic methanotrophs were the dominant diazotrophs, as

previously proposed (55, 89), the presence of O_2 and CH_4 should have stimulated $^{15}N_2$

incorporation. Instead, our study and previous work (56) found the highest rates of diazotrophy

at low O_2 and independent of CH_4 content. Acetylene fully inhibited CH_4 consumption under

oxic conditions, but only partially inhibited $^{15}N_2$ incorporation under degassed conditions (as in

Kox, et al. (54)) and had a minimal effect on oxic $^{15}N_2$ incorporation. Assuming that these

incubations reflect field conditions, our data imply: (a) peatland diazotrophy is mediated by O_2-

sensitive microorganisms that rely on photosynthesis for carbon and/or energy sources; and (b) at

least part of the diazotrophic community is C_2H_2 sensitive. When we consider this in concert

with the molecular information from the native peats, our data indicate that heterotrophic

Bradyrhizobiaceae (C_2H_2 insensitive, O_2 sensitive (46)) and *Beijerinckiaceae* (C_2H_2 insensitive,

O_2 insensitive (81)) were the primary diazotrophs in our incubations. Lesser contributions likely

came from methanotrophic *Beijerinckiaceae* (C_2H_2 sensitive, O_2 insensitive), photosynthetic

cyanobacteria from the *Oscillatoriophycideae* (C_2H_2 insensitive, O_2 sensitive (24)), and C_2H_2-

sensitive, O_2-sensitive members of anaerobic cluster III. Future studies are warranted to

interrogate contrasts in the activity and gene expression of diazotrophs when comparing

laboratory microcosm experiments to the field, since microcosm experiments are most often used

as a proxy for *in situ* activity.

Molecular markers for diazotrophy. We end with a word of caution with regard to the

molecular detection of diazotrophs. The majority of studies in peatlands have employed PCR

amplification and sequencing of nitrogenase genes (nifH) for studying the dynamics of

diazotrophs in peatlands. A wide range of *nifH* primer sets exist, with varying universality (27).

Peat bog sequencing efforts have used polF/polR (this study; 58), F1/R6 (89),

FGPH19/polR+polF/AQER (56), and 19F/nifH3+nifH1(1)/nifH2(2) with nested PCR (7, 54).

We discourage use of polF/polR for peatland samples because *in silico* evaluation predicts that

this primer set will not amplify the majority of *Proteobacteria* and/or *Cyanobacteria* and Group

III *nifH* sequences (27). Of the *nifH* primer sets used previously, F1/R6, 19F/nifH3 and

nifH1(1)/nifH2 are predicted to have the highest coverage for soils (>80% predicted primer

binding for sequences from soil ecosystems). However, it is important to be aware that the F1/R6

primer set contains a number of mismatches with cluster III, including methanogens represented

in peatlands (Table S2). In order to maximize sequence coverage, we suggest the primer set

IGK(3)/DVV for future studies (27).

Acknowledgments

This research was funded by DOE support to J.E.K. under grant numbers DE-SC0007144 and

DE-SC0012088. We thank Loren Dean Williams for inspiration with visuals, and Heather Dang

(UC Berkeley) for IRMS analysis. Work at LLNL was conducted under the auspices of DOE

Contract DE-AC52-07NA27344, with funding provided by LDRD 14-ER-038.

Figure 1. Depth profiles of NH_4^+ concentrations (black circles) and *nifH* gene copies (white squares) and transcripts (white triangle) in units of copies per gram of dry weight for the S1 bog T3 mid site. Concentrations are means of measurements from May, June and September 2014. *nifH* copy numbers are for July 2013. Error bars are standard errors. *nifH* transcripts were not detected at 20, 30 and 75 cm depths. Surface (0-10 cm) and deep (10-30 cm) peat depth intervals used for rate measurements are designated by dashed lines.

Figure 2. Dissolved molybdenum (white) and vanadium (gray) concentrations in pore water from three depths in S1 peat hollows (mid and far sites along T3 transect). Pie charts show the relative abundance of genes encoding the five nitrogenase H-subunit clusters from metagenomes for each depth; clusters I and III encode Mo-Fe nitrogenases (*nifH)*; cluster II encodes alternative (*vnfH, anfH*) nitrogenases; cluster IV encode nitrogenase paralogs. Deepest metagenomes were from 75 cm; insufficient numbers of nitrogenase H-subunit sequences were recovered from the far site for cluster analysis.

382
Figure 3. Dissolved copper (white) and iron (shaded) concentrations in pore water from

three depths in S1 peat hollows (mid and far sites along T3 transect). Pie charts show the

relative abundance of *pmoA* (Cu-containing particulate methane monooxygenase) and *mmoX*

(Fe-containing soluble methane monooxygenase) sequences recovered from metagenomes from

T3 mid sites (0-10, 25-50, and 75-100 cm).

Figure 4. Acetylene reduction rates for hummocks (90-91% water content) and hollows (93-96% water content) at the S1 bog, T3 transect (0-10 cm depth incubated in the light at 25°C for 7 days). ARA units are μmol ethylene produced per gram of dry weight per hour. Photo insets show dominant *Sphagnum species* in (a) hummocks (*S. magellanicum*) and (b) hollows (*S. fallax* and *S. angustifolium*). Photo inset (c) shows *Sphagnum* greening after incubation of hollow samples in the light for 7 days at 25°C; bottle B2* (April 2014) received air headspace with 1% CH_4 and treatment D2 (Sept 2013) received air headspace without CH_4.

Figure 5. Effect of 1% C_2H_2 on $^{15}N_2$ incorporation and CH_4 consumption for S1 bog surface peat. Rates were measured for samples collected from the NW S1 bog transect in September 2014. $^{15}N_2$ incorporation treatment conditions were 80%N_2+20% CO_2±1% CH_4 (shaded bars) or 80%N_2+20% O_2±1% CH_4 (white bars), with (a) and without (b) 1% C_2H_2; units are nmol $^{15}N_2$ incorporated per gram of dry weight per hour. CH_4 consumption treatments were 80%N_2+20% CO_2+1% CH_4 (black circles) with (a) and without (b) 1% C_2H_2; units are nmol CH_4 consumed per gram of dry weight per hour. Error bars are standard errors.

References

1. **Basilier, K., U. Granhall,T.-A. Stenström.** 1978. Nitrogen fixation in wet minerotrophic moss communities of a subarctic mire. Oikos**:**236-246.
2. **Bellenger, J. P., T. Wichard, Y. Xu,A. M. L. Kraepiel.** 2011. Essential metals for nitrogen fixation in a free living N2 fixing bacterium: chelation, homeostasis and high use efficiency. Environ. Microbiol. **13:**1395–1411.
3. **Bellenger, J. P., Y. Xu, X. Zhang, F. M. M. Morel,K. A.M.L.** 2014. Possible contribution of alternative nitrogenases to nitrogen fixation by asymbiotic N2-fixing bacteria in soils. Soil Biol Biochem **69:**413-420.
4. **Berg, A., Å. Danielsson,B. H. Svensson.** 2013. Transfer of fixed-N from N2-fixing cyanobacteria associated with the moss *Sphagnum riparium* results in enhanced growth of the moss. Plant Soil **362:**271-278.
5. **Bertine, K.** 1972. The deposition of molybdenum in anoxic waters. Marine Chemistry **1:**43-53.
6. **Betancourt, D. A., T. M. Loveless, J. W. Brown,P. E. Bishop.** 2008. Characterization of diazotrophis containing Mo-independent nitrogenases, isolated from diverse natural environments. Appl Environ Microbiol **74:**3471-3480.
7. **Bragina, A., S. Maier, C. Berg, H. Müller, V. Chobot, F. Hadacek,G. Berg.** 2011. Similar diversity of alphaproteobacteria and nitrogenase gene amplicons on two related *Sphagnum* mosses. Front Microbiol **2:**275. doi: 210.3389/fmicb.2011.00275.
8. **Bragina, A., L. Oberauner ‐ Wappis, C. Zachow, B. Halwachs, G. G. Thallinger, H. Müller,G. Berg.** 2014. The *Sphagnum* microbiome supports bog ecosystem functioning under extreme conditions. Mol Ecol **23:**4498-4510.
9. **Caporaso, J. G., C. L. Lauber, W. A. Walters, D. Berg-Lyons, J. Huntley, N. Fierer, S. M. Owens, J. Betley, L. Fraser,M. Bauer.** 2012. Ultra-high-throughput microbial community analysis on the Illumina HiSeq and MiSeq platforms. ISME J **6:**1621-1624.
10. **Chen, Y., M. G. Dumont, J. D. Neufeld, L. Bodrossy, N. Stralis-Pavese, N. P. McNamara, N. Ostle, M. J. Briones,J. C. Murrell.** 2008. Revealing the uncultivated majority: combining DNA stable isotope probing, multiple displacement amplification and metagenomic analyses of uncultivated *Methylocystis* in acidic peatlands. Environ Microbiol **10:**2609-2622.
11. **Clymo, R.** 1984. The limits to peat bog growth. Philos T Roy Soc B **303:**605-654.
12. **Clymo, R.,P. Hayward.** 1982. The ecology of *Sphagnum*, p. 229-289, Bryophyte Ecology. Springer.
13. **Colby, J.,H. Dalton.** 1976. Some properties of a soluble methane mono-oxygenase from *Methylococcus capsulatus* strain Bath. Biochem J **157:**495-497.
14. **Crombie, A. T.,J. C. Murrell.** 2014. Trace-gas metabolic versatility of the facultative methanotroph *Methylocella silvestris*. Nature **510:**148–151.
15. **Dalton, H.,R. Whittenbury.** 1976. The acetylene reduction technique as an assay for nitrogenase activity in the methane oxidizing bacterium *Methylococcus capsulatus* strain Bath. Arch Microbiol **109:**147-151.
16. **Darnajoux, R., J. Constantin, J. Miadlikowska, F. Lutzoni,J. P. Bellenger.** 2014. Is vanadium a biometal for boreal cyanolichens? New Phytol **202:**765-771.

17.   **Dedysh, S.,P. Dunfield.** 2016. Beijerinckiaceae. Bergey's Manual of Systematics of
Archaea and Bacteria.
18.   **Dedysh, S. N.,P. F. Dunfield.** 2016. Beijerinckiaceae. Bergey's Manual of Systematics
of Archaea and Bacteria **1-4**.
19.   **Dedysh, S. N., W. Liesack, V. N. Khmelenina, N. E. Suzina, Y. A. Trotsenko, J. D.**
Semrau, A. M. Bares, N. S. Panikov,J. M. Tiedje. 2000. *Methylocella palustris* gen.
nov., sp. nov., a new methane-oxidizing acidophilic bacterium from peat bogs,
representing a novel subtype of serine-pathway methanotrophs. Int J Syst Evol Microbiol
50:955-969.
20.   **Dedysh, S. N., N. S. Panikov, W. Liesack, R. Großkopf, J. Zhou,J. M. Tiedje.** 1998.
Isolation of acidophilic methane-oxidizing bacteria from northern peat wetlands. Science
282:281-284.
21.   **Dedysh, S. N., P. Ricke,W. Liesack.** 2004. NifH and NifD phylogenies: an evolutionary
basis for understanding nitrogen fixation capabilities of methanotrophic bacteria.
Microbiology **150:**1301-1313.
22.   **Dilworth, M. J., R. R. Eady, R. L. Robson,R. W. Miller.** 1987. Ethane formation from
acetylene as a potential test for vanadium nitrogenase *in vivo*. Nature **327:**167-168.
23.   **Esson, K. C., X. Lin, D. Kumaresan, J. P. Chanton, J. C. Murrell,J. E. Kostka.** 2016.
Alpha-and gammaproteobacterial methanotrophs codominate the active methane-
oxidizing communities in an acidic boreal peat bog. Appl Environ Microbiol **82:**2363-
2371.
24.   **Fay, P.** 1992. Oxygen relations of nitrogen fixation in cyanobacteria. Microbio Mol Biol
Rev **56:**340-373.
25.   **Fulweiler, R. W., E. M. Heiss, M. K. Rogener, S. E. Newell, G. R. LeCleir, S. M.**
Kortebein,S. W. Wilhelm. 2015. Examining the impact of acetylene on N-fixation and
the active sediment microbial community. Front Microbiol **6**.
26.   **Gaby, J. C.,D. H. Buckley.** 2014. A comprehensive aligned nifH gene database: a
multipurpose tool for studies of nitrogen-fixing bacteria. Database **2014:**bau001.
27.   **Gaby, J. C.,D. H. Buckley.** 2012. A comprehensive evaluation of PCR primers to
amplify the *nifH* gene of nitrogenase. PLoS One **7:**e42149.
28.   **García-Robledo, E., A. Corzo,S. Papaspyrou.** 2014. A fast and direct
spectrophotometric method for the sequential determination of nitrate and nitrite at low
concentrations in small volumes. Marine Chemistry **162:**30-36.
29.   **Garrity, G., J. Bell,T. Lilburn.** 2015. Bradyrhizobiaceae fam. nov. Bergey's Manual of
Systematics of Archaea and Bacteria**:**1-4.
30.   **Glass, J. B., R. P. Axler, S. Chandra,C. R. Goldman.** 2012. Molybdenum limitation of
microbial nitrogen assimilation in aquatic ecosystems and pure cultures. Front Microbiol
3:331.
31.   **Glass, J. B., F. L. Wolfe-Simon, J. J. Elser,A. D. Anbar.** 2010. Molybdenum-nitrogen
co-limitation in freshwater and coastal heterocystous cyanobacteria. Limnol Oceanogr
55:667-676.
32.   **Gorham, E., C. Lehman, A. Dyke, D. Clymo,J. Janssens.** 2012. Long-term carbon
sequestration in North American peatlands. Quat Sci Rev **58:**77-82.
33.   **Granhall, U.,A. V. Hofsten.** 1976. Nitrogenase activity in relation to intracellular
organisms in Sphagnum mosses. Physiol Plantarum **36:**88-94.
34.   **Granhall, U.,H. Selander.** 1973. Nitrogen fixation in a subarctic mire. Oikos**:**8-15.

35.   **Gupta, V., K. A. Smemo, J. B. Yavitt,N. Basiliko.** 2012. Active methanotrophs in two
contrasting North American peatland ecosystems revealed using DNA-SIP. Microbial
Ecol **63:**438-445.
36.   **Hanson, P. J., J. S. Riggs, W. R. Nettles, J. R. Phillips, M. B. Krassovski, L. A.**
Hook, L. Gu, A. D. Richardson, D. M. Aubrecht,D. M. Ricciuto. 2017. Attaining
whole-ecosystem warming using air and deep-soil heating methods with an elevated CO2
atmosphere. Biogeosciences **14:**861-883.
37.   **Hardy, R., R. Burns,R. Holsten.** 1973. Applications of the acetylene-ethylene assay for
measurement of nitrogen fixation. Soil Biol Biochem **5:**47-81.
38.   **Hardy, R. W. F., R. D. Holsten, E. K. Jackson,R. C. Burns.** 1968. The acetylene-
ethylene assay for N2 fixation: laboratory and field evaluation. Plant Physiol **43:**1185-
1207.
39.   **Hill, B. H., T. M. Jicha, L. L. Lehto, C. M. Elonen, S. D. Sebestyen,R. K. Kolka.**
2016. Comparisons of soil nitrogen mass balances for an ombrotrophic bog and a
minerotrophic fen in northern Minnesota. Sci Total Environ **550:**880-892.
40.   **Ho, A.,P. L. Bodelier.** 2015. Diazotrophic methanotrophs in peatlands: the missing link?
Plant Soil **389:**419-423.
41.   **Hodkinson, B., J. Allen, L. Forrest, B. Goffinet, E. Seurasiaux, O. Andresson, V.**
Miao, J. P. Bellenger,F. Lutzoni. 2014. Lichen-symbiotic cyanobacteria associated with
Peltigera have an alternative vanadium-dependent nitrogen fixation system. Eur J Phycol
49:11-19.
42.   **Huson, D. H., S. Mitra, H.-J. Ruscheweyh, N. Weber,S. C. Schuster.** 2011. Integrative
analysis of environmental sequences using MEGAN4. Genome research **21:**1552-1560.
43.   **Hutchens, E., S. Radajewski, M. G. Dumont, I. R. McDonald,J. C. Murrell.** 2004.
Analysis of methanotrophic bacteria in Movile Cave by stable isotope probing. Environ
Microbiol **6:**111-120.
44.   **Hynes, R.,R. Knowles.** 1982. Effect of acetylene on autotrophic and heterotrophic
nitrification. Can J Microbiol **28:**334-340.
45.   **Keister, D.** 1975. Acetylene reduction by pure cultures of *Rhizobia*. J Bacteriol
123:1265-1268.
46.   **Keister, D. L.,W. R. Evans.** 1976. Oxygen requirement for acetylene reduction by pure
cultures of rhizobia. J Bacteriol **127:**149-153.
47.   **Kip, N., B. E. Dutilh, Y. Pan, L. Bodrossy, K. Neveling, M. P. Kwint, M. S. Jetten,H.**
J. Op den Camp. 2011. Ultra-deep pyrosequencing of *pmoA* amplicons confirms the
prevalence of *Methylomonas* and *Methylocystis* in *Sphagnum* mosses from a Dutch peat
bog. Environ Microbiol Rep **3:**667-673.
48.   **Kip, N., W. Ouyang, J. van Winden, A. Raghoebarsing, L. van Niftrik, A. Pol, Y.**
Pan, L. Bodrossy, E. G. van Donselaar,G.-J. Reichart. 2011. Detection, isolation, and
characterization of acidophilic methanotrophs from *Sphagnum* mosses. Appl Environ
Microbiol **77:**5643-5654.
49.   **Kip, N., J. F. van Winden, Y. Pan, L. Bodrossy, G.-J. Reichart, A. J. Smolders, M. S.**
Jetten, J. S. S. Damsté,H. J. O. den Camp. 2010. Global prevalence of methane
oxidation by symbiotic bacteria in peat-moss ecosystems. Nature Geosci **3:**617-621.
50.   **Knorr, K. H., M. A. Horn,W. Borken.** 2015. Significant nonsymbiotic nitrogen fixation
in Patagonian ombrotrophic bogs. Glob Change Biol **21:**2357-2365.

51.   **Kolb, S., C. Knief, S. Stubner,R. Conrad.** 2003. Quantitative detection of
methanotrophs in soil by novel *pmoA*-targeted real-time PCR assays. Appl Environ
Microbiol **69:**2423-2429.
52.   **Kolka, R., S. Sebestyen, E. S. Verry,K. Brooks.** 2011. Peatland biogeochemistry and
watershed hydrology at the Marcell experimental forest. CRC Press.
53.   **Kostka, J. E., D. J. Weston, J. B. Glass, E. A. Lilleskov, A. J. Shaw,M. R. Turetsky.**
2016. The *Sphagnum* microbiome: new insights from an ancient plant lineage. New
Phytol **211:**57-64.
54.   **Kox, M. A., C. Lüke, C. Fritz, E. van den Elzen, T. van Alen, H. J. O. den Camp, L.**
P. Lamers, M. S. Jetten,K. F. Ettwig. 2016. Effects of nitrogen fertilization on
diazotrophic activity of microorganisms associated with *Sphagnum magellanicum*. Plant
Soil **406:**83-100.
55.   **Larmola, T., S. M. Leppänen, E.-S. Tuittila, M. Aarva, P. Merilä, H. Fritze,M.**
Tiirola. 2014. Methanotrophy induces nitrogen fixation during peatland development.
Proc Natl Acad Sci **111:**734-739.
56.   **Leppänen, S. M., A. J. Rissanen,M. Tiirola.** 2014. Nitrogen fixation in *Sphagnum*
mosses is affected by moss species and water table level. Plant Soil**:**1-12.
57.   **Leppänen, S. M., M. Salemaa, A. Smolander, R. Mäkipää,M. Tiirola.** 2013. Nitrogen
fixation and methanotrophy in forest mosses along a N deposition gradient. Environ Exp
Bot **90:**62-69.
58.   **Liebner, S.,M. M. Svenning.** 2013. Environmental transcription of *mmoX* by methane-
oxidizing Proteobacteria in a subarctic palsa peatland. Appl Environ Microbiol **79:**701-
706.
59.   **Limpens, J., F. Berendse, C. Blodau, J. Canadell, C. Freeman, J. Holden, N. Roulet,**
H. Rydin,G. Schaepman-Strub. 2008. Peatlands and the carbon cycle: from local
processes to global implications: a synthesis. Biogeosciences **5:**1475-1491.
60.   **Limpens, J., M. M. Heijmans,F. Berendse.** 2006. The nitrogen cycle in boreal
peatlands, p. 195-230, Boreal peatland ecosystems. Springer.
61.   **Lin, X., S. Green, M. Tfaily, O. Prakash, K. Konstantinidis, J. Corbett, J. Chanton,**
W. Cooper,J. Kostka. 2012. Microbial community structure and activity linked to
contrasting biogeochemical gradients in bog and fen environments of the Glacial Lake
Agassiz Peatland. Appl Environ Microbiol **78:**7023-7031.
62.   **Lin, X., K. M. Handley, J. A. Gilbert,J. E. Kostka.** 2015. Metabolic potential of fatty
acid oxidation and anaerobic respiration by abundant members of Thaumarchaeota and
Thermoplasmata in deep anoxic peat. ISME J **9:**2740-2744.
63.   **Lin, X., M. M. Tfaily, S. Green, J. M. Steinweg, P. Chanton, A. Imvittaya, J. P.**
Chanton, W. Cooper, C. W. Schadt,J. E. Kostka. 2014. Microbial metabolic potential
for carbon degradation and nutrient (nitrogen and phosphorus) acquisition in an
ombrotrophic peatland. Appl Environ Microbiol **80:**3531-3540.
64.   **Lin, X., M. M. Tfaily, J. M. Steinweg, P. Chanton, K. Esson, Z. K. Yang, J. P.**
Chanton, W. Cooper, C. W. Schadt,J. E. Kostka. 2014. Microbial community
stratification linked to utilization of carbohydrates and phosphorus limitation in a boreal
peatland at Marcell Experimental Forest, Minnesota, USA. Appl Environ Microbiol
80:3518-3530.

65. **Loveless, T. M., J. R. Saah,P. E. Bishop.** 1999. Isolation of nitrogen-fixing bacteria
containing molybdenum-independent nitrogenases from natural environments. Appl
Environ Microbiol **65:**4223-4226.
66. **McGlynn, S. E., E. S. Boyd, J. W. Peters,V. J. Orphan.** 2013. Classifying the metal
dependence of uncharacterized nitrogenases. Front Microbiol **3:**419.
67. **Miller, R.,R. Eady.** 1988. Molybdenum and vanadium nitrogenases of *Azotobacter*
chroococcum. Low temperature favours N2 reduction by vanadium nitrogenase. Biochem
J **256:**429-432.
68. **Murphy, J.,J. Riley.** 1962. A modified single solution method for the determination of
phosphate in natural waters. Anal Chim Acta **27:**31-36.
69. **Opelt, K., C. Berg, S. Schonmann, L. Eberl,G. Berg.** 2007. High specificity but
contrasting biodiversity of *Sphagnum*-associated bacterial and plant communities in bog
ecosystems independent of the geographical region. The ISME Journal **1:**502-516.
70. **Opelt, K., V. Chobot, F. Hadacek, S. Schonmann, L. Eberl,G. Berg.** 2007.
Investigations of the structure and function of bacterial communities associated with
Sphagnum mosses. Environ Microbiol **9:**2795-2809.
71. **Oremland, R. S.,B. F. Taylor.** 1975. Inhibition of methanogenesis in marine sediments
by acetylene and ethylene: validity of the acetylene reduction assay for anaerobic
microcosms. Appl Microbiol **30:**707-709.
72. **Payne, W.,M. Grant.** 1982. Influence of acetylene on growth of sulfate-respiring
bacteria. Appl Environ Microbiol **43:**727-730.
73. **Prior, S. D.,H. Dalton.** 1985. Acetylene as a suicide substrate and active site probe for
methane monooxygenase from *Methylococcus capsulatus* (Bath). FEMS Microbiol Lett
29:105-109.
74. **Raghoebarsing, A. A., A. J. Smolders, M. C. Schmid, W. I. C. Rijpstra, M. Wolters-**
Arts, J. Derksen, M. S. Jetten, S. Schouten, J. S. S. Damsté, L. P. Lamers, J. G. M.
Roelofs, H. J. M. Op den Camp,M. Strous. 2005. Methanotrophic symbionts provide
carbon for photosynthesis in peat bogs. Nature **436:**1153-1156.
75. **Rahman, M. T., A. Crombie, Y. Chen, N. Stralis-Pavese, L. Bodrossy, P. Meir, N. P.**
McNamara,J. C. Murrell. 2011. Environmental distribution and abundance of the
facultative methanotroph Methylocella. The ISME journal **5:**1061-1066.
76. **Raymond, J., J. L. Siefert, C. R. Staples,R. E. Blankenship.** 2004. The natural history
of nitrogen fixation. Mol Biol Evol **21:**541-554.
77. **Rosswall, T.,U. Granhall.** 1980. Nitrogen cycling in a subarctic ombrotrophic mire.
Ecological Bulletins**:**209-234.
78. **Schwintzer, C. R.** 1983. Nonsymbiotic and symbiotic nitrogen fixation in a weakly
minerotrophic peatland. Am J Bot **70:**1071-1078.
79. **Sebestyen, S. D., C. Dorrance, D. M. Olson, E. S. Verry, R. K. Kolka, A. E. Elling,R.**
Kyllander. 2011. Long-term monitoring sites and trends at the Marcell Experimental
Forest. Peatland Biogeochemistry and Watershed Hydrology at the Marcell Experimental
Forest**:**93-134.
80. **Semrau, J. D., A. A. DiSpirito,S. Yoon.** 2010. Methanotrophs and copper. FEMS
Microbiol Rev **34:**496-531.
81. **Spiff, E.,C. Odu.** 1973. Acetylene reduction by Beijerinckia under various partial
pressures of oxygen and acetylene. Microbiology **78:**207-209.

82. **Sprott, G., K. Jarrell, K. Shaw,R. Knowles.** 1982. Acetylene as an inhibitor of
methanogenic bacteria. J Gen Microbiol **128:**2453-2462.
83. **Steinberg, L. M.,J. M. Regan.** 2008. Phylogenetic comparison of the methanogenic
communities from an acidic, oligotrophic fen and an anaerobic digester treating
municipal wastewater sludge. Appl Environ Microbiol **74:**6663-6671.
84. **Stirling, D. I.,H. Dalton.** 1977. Effect of metal-binding agents and other compounds on
methane oxidation by two strains of *Methylococcus capsulatus*. Arch Microbiol **114:**71-
76.
85. **Strickland, J.,T. Parsons.** 1972. A practical handbook of seawater analysis, vol. 167.
86. **Tamura, K., D. Peterson, N. Peterson, G. Stecher, M. Nei,S. Kumar.** 2011. MEGA5:
molecular evolutionary genetics analysis using maximum likelihood, evolutionary
distance, and maximum parsimony methods. Mol Biol Evol **28:**2731-2739.
87. **Urban, N.,S. Eisenreich.** 1988. Nitrogen cycling in a forested Minnesota bog. Can J Bot
66:435-449.
88. **Vaughan, D., D. Lumsdon,D. Linehan.** 1993. Influence of dissolved organic matter on
the bio-availability and toxicity of metals in soils and aquatic systems. Chem Ecol **8:**185-
201.
89. **Vile, M. A., R. K. Wieder, T. Živković, K. D. Scott, D. H. Vitt, J. A. Hartsock, C. L.**
Iosue, J. C. Quinn, M. Petix,H. M. Fillingim. 2014. N2-fixation by methanotrophs
sustains carbon and nitrogen accumulation in pristine peatlands. Biogeochemistry
121:317-328.
90. **Walmsley, J.,C. Kennedy.** 1991. Temperature-dependent regulation by molybdenum
and vanadium of expression of the structural genes encoding 3 nitrogenases in
Azotobacter vinelandii. Appl Environ Microbiol **57:**622-624.
91. **Wedepohl, K. H.** 1995. The composition of the continental crust. Geochem Cosmochim
Acta **59:**1217-1232.
92. **Weston, D. J., C. M. Timm, A. P. Walker, L. Gu, W. Muchero, J. Schmutz, A. J.**
Shaw, G. A. Tuskan, J. M. Warren,S. D. Wullschleger. 2015. *Sphagnum* physiology in
the context of changing climate: Emergent influences of genomics, modeling and host-
microbiome interactions on understanding ecosystem function. Plant Cell Environ
38:1737-1751.
93. **Wong, P. P.,R. Burris.** 1972. Nature of oxygen inhibition of nitrogenase from
Azotobacter vinelandii. Proc Natl Acad Sci **69:**672-675.
94. **Yu, Z.** 2012. Northern peatland carbon stocks and dynamics: a review. Biogeosciences
4071–4085:4071–4085.
95. **Zhao, Y., H. Tang,Y. Ye.** 2012. RAPSearch2: a fast and memory-efficient protein
similarity search tool for next-generation sequencing data. Bioinformatics **28:**125-126.
96. **Zheng, H., C. Dietrich, R. Radek,A. Brune.** 2016. *Endomicrobium proavitum*, the first
isolate of Endomicrobia class. nov.(phylum Elusimicrobia)–an ultramicrobacterium with
an unusual cell cycle that fixes nitrogen with a Group IV nitrogenase. Environ Microbiol
18:191-204.
671

Acetylene reduction rate (μmol g⁻¹ h⁻¹) vs **Water content (%)**

Legend:
- ■ CO₂
- ● CO₂/CH₄
- ● CH₄
- □ O₂
- ○ O₂/CH₄

Sphagnum fallax
Sphagnum angustifolium

Hollows

Sphagnum magellanicum

Hummocks

B2*   D2