Molybdenum-based diazotrophy in a *Sphagnum* peatland in northern Minnesota

Melissa J. Warren[1*], Xueju Lin[2**], John C. Gaby[2***], Cecilia B. Kretz[1****], Max Kolton[2], Peter L.

Morton[3], Jennifer Pett-Ridge[4], David J. Weston[5], Christopher W. Schadt[5], Joel E. Kostka[1,2],

Jennifer B. Glass[1,2#]

[1]School of Earth and Atmospheric Sciences, Georgia Institute of Technology, Atlanta, GA, USA

[2]School of Biology, Georgia Institute of Technology, Atlanta, GA, USA

[3]Department of Earth, Ocean and Atmospheric Science, Florida State University, Tallahassee,

Florida, USA

[4]Physical and Life Sciences Directorate, Lawrence Livermore National Laboratory, Livermore,

CA, USA

[5]Biosciences Division, Oak Ridge National Laboratory, Oak Ridge, TN, USA

Now at: *CH2M, Atlanta, GA, USA; **InstantLabs, Baltimore, MD, USA; ***Faculty of

Chemistry, Biotechnology and Food Science, Norwegian University of Life Sciences, Oslo,

Norway; ****Division of Bacterial Diseases, National Center for Immunization and Respiratory

Diseases, Centers for Disease Control and Prevention, Atlanta, GA, USA

#Corresponding author: Jennifer B. Glass, 404-894-3942; jennifer.glass@eas.gatech.edu

19
Running Title: Nitrogen fixation in a *Sphagnum* peat bog

Abstract

Microbial N_2 fixation (diazotrophy) represents an important nitrogen source to oligotrophic

peatland ecosystems, which are important sinks for atmospheric CO_2 and susceptible to changing

climate. The objectives of this study were: (i) to determine the active microbial group and type of

nitrogenase mediating diazotrophy in a ombrotrophic *Sphagnum*-dominated peat bog (the S1

peat bog, Marcell Experimental Forest, Minnesota, USA); and (ii) to determine the effect of

environmental parameters (light, O_2, CO_2, CH_4) on potential rates of diazotrophy measured by

acetylene (C_2H_2) reduction and $^{15}N_2$ incorporation. Molecular analysis of metabolically active

microbial communities suggested that diazotrophy in surface peat was primarily mediated by

Alphaproteobacteria (*Bradyrhizobiaceae* and *Beijerinckiaceae*). Despite higher dissolved

vanadium (V; 11 nM) than molybdenum (Mo; 3 nM) in surface peat, a combination of

metagenomic, amplicon sequencing and activity measurements indicated that Mo-containing

nitrogenases dominate over the V-containing form. Acetylene reduction was only detected in

surface peat exposed to light, with the highest rates observed in peat collected from hollows with

the highest water content. Incorporation of $^{15}N_2$ was suppressed 90% by O_2 and 55% by C_2H_2,

and was unaffected by CH_4 and CO_2 amendments. These results suggest that peatland

diazotrophy is mediated by a combination of C_2H_2-sensitive and C_2H_2-insensitive microbes that

are more active at low O_2 and show similar activity at high and low CH_4.

Importance

Previous studies indicate that diazotrophy provides an important nitrogen source and is linked to methanotrophy in *Sphagnum*-dominated peatlands. However, the environmental controls and enzymatic pathways of peatland diazotrophy, as well as the metabolically active microbial populations that catalyze this process remain in question. Our findings indicate that oxygen levels and photosynthetic activity override low nutrient availability in limiting diazotrophy, and that members of the *Alphaproteobacteria* (*Rhizobiales*) catalyze this process at the bog surface using the molybdenum-based form of the nitrogenase enzyme.

Introduction

High-latitude peatlands store approximately one-third of global soil carbon and may pose a climatic threat if rising global temperatures accelerate the release of this stored carbon in gaseous forms, as either carbon dioxide or methane (35, 64, 106). Mineral-poor (ombrotrophic) peatlands receive most of their nutrient inputs from atmospheric deposition and contain *Sphagnum* moss as their primary plant cover (12, 64). The peatmoss *Sphagnum* is a keystone genus in these ecosystems and is responsible for much of the primary production and recalcitrant dead organic matter (13, 103). *Sphagnum* mosses also host complex microbiomes (9, 58, 78, 79), including N_2-fixers (diazotrophs) that are significant nitrogen sources to peatland ecosystems (5).

Despite decades of research, there is still much debate on the identity of the dominant diazotrophs in ombrotrophic peatlands. Early work implicated *Cyanobacteria* (2, 36, 37) or heterotrophic bacteria (88) based primarily on microscopic studies, while more recent molecular analyses argue for the importance of methanotrophic *Beijerinckiaceae* (19) as major diazotrophs in *Sphagnum* peat bogs (8, 22, 44, 99). Possible contributions from other potential diazotrophs, such as strictly anaerobic methanogenic *Euryarchaeota*, remain unknown. However, it is quite possible that diverse diazotrophs exist within defined niches of peatland environments (60).

Diazotrophy is catalyzed by the nitrogenase metalloenzyme, a complex of thee subunits (H, D and K) that contains abundant iron as Fe-S clusters. This enzyme is extremely O_2 sensitive (105), and must be protected from exposure to O_2 for diazotrophy to occur (25). The most common form of nitrogenase, encoded by *nif* genes, contains molybdenum (Mo) as its cofactor. When Mo is scarce, some species of Bacteria and Archaea express nitrogenases containing vanadium (V; *vnf* genes) or iron (Fe; *anf* genes) in place of Mo, but these "alternative" nitrogenases are less efficient than the Mo form (74, 101). The most conserved nitrogenase gene,

nifH (77), has become the marker gene of choice for environmental diazotrophy (28, 29, 107).

Phylogenetic studies show five *nifH* clusters: aerobic bacteria (cluster I); alternative nitrogenases

(cluster II); anaerobic bacteria and archaea (cluster III); uncharacterized sequences (cluster IV),

and paralogs related to chlorophyll biosynthesis (cluster V; 85). Because *vnfH* and *anfH* genes in

cluster II cannot be differentiated by sequence alone, the D-subunit (*nifD*/*vnfD*/*anfD*) has

become the preferred marker gene for studies of alternative nitrogenases (4). Consistent with

higher concentrations of V than Mo in most rocks (102), microbes from diverse soils have been

shown to contain *vnfD* genes (4, 7, 18, 45, 70, 73). Given that oligotrophic conditions dominate

in peatlands, trace metals may limit diazotrophy. However, little is known about trace metal

availability and the role of alternative nitrogenase pathways in ombrotrophic peatlands.

Similarly, methane monooxygenase (MMO, the enzyme that catalyzes the first step of

methane oxidation) occurs in particulate (copper (Cu)-containing pMMO) and soluble (Fe-

containing sMMO) forms. While pMMO has more specific substrate requirements, pathways

that employ sMMO can use a wider range of compounds (15). Both forms of MMO are inhibited

by acetylene (C_2H_2) (14, 82). In organisms with both sets of genes, pMMO is expressed when Cu

is abundant, whereas Cu limitation induces sMMO expression (90). The dominant peatland

methanotrophs in *Alphaproteobacteria* and *Gammaproteobacteria* tend to possess both MMOs

(11, 24, 39, 52, 68), although *Methylocella* and *Methyloferula* species containing solely sMMO

have been isolated from peat bogs (20, 21, 100). While most studies have primarily targeted the

pmoA gene (24, 52), *mmoX* genes and transcripts have also been reported in peatlands (63, 68,

84), raising questions about the relative importance of each form for peatland methane oxidation.

The acetylene reduction assay (ARA) is commonly used as a proxy for diazotroph

activity (41, 42). This assay is effective for capturing the potential activity of diazotrophs that are

not inhibited by C_2H_2, such as *Cyanobacteria* and non-methanotrophic *Proteobacteria* (e.g.

Bradyrhizobiaceae) (50). However, a number of functional guilds of microorganisms, including

methanotrophs, methanogens, sulfate reducers, and nitrifiers, are inhibited by C_2H_2 (16, 49, 80,

81, 92, 94). If these or other C_2H_2-sensitive microbes perform diazotrophy and/or provide

substrates to other diazotrophs (see Fulweiler, et al. (27)), ARA may underestimate diazotrophy

in that system. Thus, recent studies have shifted to tracking diazotrophy by incorporation of the

stable isotope tracer, $^{15}N_2$ (55, 60, 61, 99).

In this study of the S1 peat bog at the Marcell Experimental Forest in northern

Minnesota, USA, dissolved macro- (NH_4^+, NO_3^-, PO_4^{3-}) and micro- (Fe, Cu, V, Mo) nutrients

were profiled along with the community composition and abundance of diazotrophic

microorganisms. We also performed separate laboratory incubation experiments to measure

potential rates of ARA and $^{15}N_2$ incorporation in order to: (a) assess environmental controls

(light, O_2, CH_4) on diazotrophy; (b) quantify the effect of C_2H_2 on rates of diazotrophy and

methanotrophy; and (c) search for diagnostic markers for alternative nitrogenase activity such as

a low conversion factor of ARA to $^{15}N_2$-incorporation (4) and C_2H_2 reduction to ethane (23).

Finally, we make recommendations on universal *nifH* primers for amplicon sequencing and

quantitative PCR based on our findings.

Materials and Methods

Site description and sample collection. Samples were collected from the S1 (black spruce-

Sphagnum spp.) peat bog at Marcell Experimental Forest (MEF; 47°30.476' N; 93°27.162' W),

the site of the DOE SPRUCE (Spruce and Peatland Responses Under Climatic and

Environmental Change) experiment in northern Minnesota, USA (40). The S1 bog is

ombrotrophic and acidic (average pH 3.5-4; 57, 89). Over the summer months, the water table is

±5 cm from the hollow surface (38, 89). Dissolved O_2 levels decrease to below detection (~20

ppb) within the top 5 cm of the bog. Three locations were sampled along S1 bog transect 3 (T3)

at near, middle and far sites (see Lin, et al. (69) for further details). Surface (0-10 cm depth) peat

was collected from hollows dominated by a mixture of *Sphagnum fallax* and *S. angustifolium*,

and hummocks dominated by *S. magellanicum*. Peat depth cores (0-200 cm) were sampled from

hollows where the water level reached the surface of the *Sphagnum* layer.

Macronutrients. Peat porewater was collected using piezometers from 0, 10, 25, 50, 75,

150 and 200 cm depth. Piezometers were recharged the same day as collection, and porewater

was pumped to the surface, filtered through sterile 0.2 µm polyethersulfone membrane filters,

and stored frozen until analysis. Nitrate (NO_3^-) and nitrite (NO_2^-) were analyzed using the

spectrophotometric assay as described by García-Robledo, et al. (31). Ammonium (NH_4^+)

concentrations were determined with the indophenol blue assay (95). Phosphate concentrations

were measured with the molybdate-antimony ascorbic acid colorimetric assay (75).

Micronutrients. Peat porewater was collected from two locations in the S1 bog from

cores at 0-30 cm, 30-50 cm, and 100-150 cm depths by filtration through 0.15 µm Rhizon soil

samplers (Rhizosphere Research Products). All plastics were washed with HCl prior to sampling;

Rhizon soil samplers were cleaned by pumping 10 mL of 1 N HCl through them, followed by a

rinsing with ultrapure water until the pH returned to neutral (~100 mL/filter). After collection,

samples were acidified to 0.32 M HNO_3 (Fisher Optima) and analyzed using a Thermo

ELEMENT 2 HR-ICP-MS (National High Magnetic Field Laboratory, Florida State University).

Initial analyses resulted in frequent clogging of the nebulizer, likely due to abundant dissolved

organic carbon. Therefore, samples were diluted 1:10 to minimize interruptions from nebulizer

clogs. Concentrations were quantified with a 7-point external calibration using standards

prepared in 0.32 N HNO$_3$ from a multi-element standard mix (High Purity Standards).

In order to generate an organic-free sample matrix suitable for ICP-MS analysis without

contaminating or diluting the sample, subsequent samples were digested as follows: 1 mL

aliquots of the porewater samples were heated in 15-mL Teflon beakers (Savillex) with 1 mL of

16 N HNO$_3$ (Ultrex II, JT Baker) and 100 μL of 30% H$_2$O$_2$ (Ultrex II, JT Baker) for 36 h at

230°C in a trace metal clean, polypropylene exhaust hood. The HNO$_3$/H$_2$O$_2$ mixture oxidizes any

DOM to CO$_2$, but the resulting matrix is too acidic for direct ICP-MS introduction. Therefore,

samples were evaporated to near dryness and resuspended in a 0.32 N HNO$_3$ matrix suitable for

ICP-MS analysis, and analyzed by ELEMENT2 ICP-MS along with parallel blank solutions.

Quantification and sequencing of gene and transcript amplicons. Peat was frozen on

dry ice at the field site in July 2013, or in liquid N$_2$ after 7 day incubations at 25°C in the light

under degassed (80% N$_2$ + 20% CO$_2$) headspace with 1% C$_2$H$_2$, with or without 1% CH$_4$ for June

2014 incubations (see ARA section). DNA and RNA were extracted with MoBio PowerSoil

DNA and total RNA extraction kits, respectively, as described in Lin, et al. (68). RNA was

cleaned with a TURBO DNA-free kit (Ambion). Nucleic acid purity was analyzed for 260/280

absorbance ratio (1.8-2.0) on a NanoDrop spectrophotometer. cDNA was synthesized using the

GoScript reverse transcription system (Promega) according to the manufacturer's protocol.

Plasmid standards for qPCR were constructed according to Lin, et al. (66). Primer pairs

are given in Table 1. The gene fragments of *nifH*, *pmoA* and *mcrA* for constructing plasmid

standards for qPCR were amplified from genomic DNA of *Rhodobacter sphaeroides*,

Methylococcus capsulatus Bath, and S1 peat bog peat soil, respectively. To prepare cDNA

standards, plasmid DNA with a positive gene insert was linearized with NcoI restriction enzyme

following the manufacturer's protocol (Promega), and purified by MinElute PCR purification kit

(Qiagen). RNA was synthesized from the linearized plasmid DNA using the Riboprobe *in vitro*

transcription system according to the manufacturer's protocol (Promega), followed by cDNA

synthesis using the GoScript reverse transcription system (Promega) according to the

manufacturer's protocol.

The abundance of functional gene transcripts were quantified in samples run in duplicate

on a StepOnePlus Real-Time PCR System (ABI) using Power SYBR Green PCR master mix.

Reaction mixtures were 20-µL reaction mixtures with 2-µL of template cDNA (10-100 ng/µL)

added to 10 µL of SYBR green master mix, 0.5-1.6 µL of each forward and reverse primer (0.3-

0.8 µM final concentration; Table 1), and 4.8-6.5 µL of PCR-grade water. Samples were run

against a cDNA standard curve (10^1 to 10^7 copies of plasmid gene fragment) on a StepOnePlus

qPCR instrument with 96 wells with an initial denaturation step of 2-5 min at 95°C and 40 cycles

of denaturation at 95°C for 15-30 s, annealing at 55-64°C for 30-45 s, extension at 72°C for 30-

45 s, and data acquisition at 83-86.5°C for 16-30 s. To minimize the effects of inhibitors in

assays, peat DNA was diluted to 1/40 of original concentrations, and duplicate 20-µL reaction

mixtures, each containing 2-µL of diluted DNA, were run for each sample. Functional gene and

transcript copy numbers were normalized to dry weight of peat, or 16S rRNA transcript copies

for incubation samples. Amplicons were sent to the University of Illinois at Chicago for DNA

sequencing using a 454 platform. Raw sequences were demultiplexed, trimmed, and quality

filtered in CLCbio. The phylogeny of *vnfD/anfD* sequences was inferred using the Maximum

Likelihood method based on the Kimura 2-parameter model in MEGA5 (96).

Acetylene reduction and methane consumption rates. Samples of bulk peat (*Sphagnum*

spp. and surrounding soil) were collected from 0-10 and 10-30 cm depths in September 2013,

April 2014, June 2014, September 2014 and August 2015, and stored at 4°C until the start of

laboratory incubations. Samples from 0-10 cm depth were gently homogenized so as not to

rupture *Sphagnum* spp. tissues, while peat samples from 10-30 cm depth were fully

homogenized. For each sample, 5 g of bulk peat were placed in 70 mL glass serum bottles,

stoppered with black bromobutyl stoppers (Geo-Microbial Technologies, pre-treated by boiling

3x in 0.1 M NaOH) and sealed with an aluminum crimp seal. Headspaces were oxic (room air,

80% N_2 + 20% O_2) or degassed (100% N_2 or 80% N_2 + 20% CO_2) with or without 1% C_2H_2 or

1% CH_4. Treatments were incubated for one week at 25°C in the light or dark. A gas

chromatograph with a flame ionization detector (SRI Instruments) equipped with a HayeSep N

column was used to quantify CH_4, C_2H_2 and C_2H_4. Samples were measured for C_2H_4 production

daily until C_2H_4 production was linear (~7 days). Controls not amended with C_2H_2 did not

produce ethylene (C_2H_4). Incubations of hollow peat from June 2014 incubated in oxic

headspace with and without 1% C_2H_2 were also monitored for consumption of 1% CH_4.

Statistical analysis was performed with JMP Pro (v. 12.1.0) using the Tukey-Kramer HSD

comparison of all means.

$^{15}N_2$ incorporation rates. In September 2014, samples were quantified for N_2 fixation

rates by $^{15}N_2$ incorporation in parallel with ARA measurements. Incubations were set up as

described above and supplemented with 7 mL of 98% $^{15}N_2$ (Cambridge Isotope Laboratories,

Tewksbury, MA, USA). After 7 days, samples were dried at 80°C, homogenized into a fine

powder, and analyzed for N content and $\delta^{15}N$ by isotope-ratio mass spectrometry (IRMS) with a

MICRO cube elemental analyzer and IsoPrime100 IRMS (Elementar) at the University of

California, Berkeley, corrected relative to National Institute of Standards and Technology (NIST,

Gaithersburg, MD, USA) standards.

Metagenomic analyses. Metagenomes were generated in a previous study (67).

Diazotrophic and methanotrophic pathways were investigated using the following bioinformatics

approaches. Briefly, Illumina reads were filtered by quality (Phred33 score threshold of Q25)

using Trim Galore (Babraham Bioinformatics) and a minimum sequence length cut off of 100

bp. The sequences were then queried using RAPSearch2 (109) against the NCBI-nr database of

non-redundant protein sequences as of November 2013. Sequences with a bit-score of 50 and

higher were retained to determine the total number of functional genes for normalization across

the different samples. The taxonomic composition of protein-coding sequences was determined

based on the taxonomic annotation of each gene according to the NCBI-nr taxonomy in

MEGAN5 (47) (min score: 50; max expected: 0.01; top percent: 10; min complexity: 0.3).

To classify sequences by nitrogenase cluster type, genes were analyzed using BLASTX

(e-value 0.1; bit-score 50) versus a custom *nifH* database that includes a phylogenetic tree to

distinguish the principal clusters (I, V, III) in the *nifH* phylogeny, as well as paralogous cluster

IV, *nifH*-like sequences (28). *nifH* genes from the four clusters were normalized to total protein-

coding genes from RAPSearch2 output sequences. The relative abundance of particulate (*pmoA*)

vs. soluble (*mmoX*) methane monooxygenase was based on previous analyses in Lin, et al. (68).

Nucleotide sequence accession numbers. Metagenomes were reported in a previous

study (67) and deposited in BioProject PRJNA382698 (SAMN06712535-06712540). *pmoA*

cDNA amplicons were reported in a previous study (24) and deposited in BioProject

PRJNA311735. *nifH*, *mcrA*, *nifD*, and *vnfD/anfD* cDNA amplicons were deposited in

BioProjects PRJNA382268, PRJNA382282, PRJNA382288 and PRJNA382295, respectively.

Results

Macro- and micro-nutrient profiles. In S1 bog hollows, NH_4^+ was ~2 µM from the surface to 25

cm depth, and increased at greater depths (Fig. 1; Fig. S1a). Nitrate was <1 µM in surface peat

and decreased with depth (Fig. S1b). Phosphate was <0.1 µM from the surface to 25 cm depth,

and then increased with depth (Fig. S1c). For the metal pairs of greatest interest to this study, V

(5-21 nM) was consistently more abundant than Mo (1-7 nM; Fig. 3), and Fe (7-35 µM) was

three orders of magnitude higher than Cu (7-38 nM; Fig. S3) at all three depth intervals (0-30,

30-50 and 100-150 cm), and at three sampling dates (September 2014, June 2015, and September

2015; data shown for June 2015 in Figs. 3 and S2). Other trace nutrients were in the nano- to

micro-molar range: Co (5-20 nM), Ni (10-80 nM), Zn (50-250 nM) and Mn (60-2220 nM; Table

S1). Macro- and micro-nutrient profiles were essentially identical in profiles from Zim bog,

another ombrotrophic bog ~80 km southeast of the S1 bog, sampled in September 2014, with the

exception of lower Fe at the surface (data not shown).

Nitrogenase expression and phylogeny. With the polF/polR primer pair, we measured

1.2×10^7 copies g^{-1} *nifH* genes at 1 and 20 cm, and 0.2×10^7 copies g^{-1} at 30 and 75 cm; *nifH*

transcripts (12.2×10^7 copies g^{-1}) were only detected at 1 cm (10:1 transcript: gene ratio), and not

at deeper depths (Fig. 1; Fig. S3a). Sequencing of cDNA from surface peat amplified with

polF/polR (*nifH*) and nifD820F/nifD1331R (*nifD*) primers showed that the majority of

nitrogenase transcripts belonged to cluster I (*Alphaproteobacteria*), with *Beijerinckiaceae*

dominating *nifH* sequences and *Bradyrhizobiaceae* dominating *nifD* sequences (Fig. 2).

Additional alphaproteobacterial *nif* transcripts matched to *Rhodospirillaceae, Rhizobiaceae,*

Rhodobacteraceae, Methylocystaceae and *Xanthobacteraceae* (Fig. 2). *Gammaproteobacteria*,

Cyanobacteria (*Oscillatoriophycideae*) and *Nitrospira* were also observed at lower abundance in

cDNA amplicon libraries (data not shown).

In metagenomes, *nifH* genes were roughly equally distributed between Mo-dependent

clusters I and III, and cluster IV/V paralogs (Fig. 2; Table S2). Sequences from Cluster II

(alternative nitrogenases, *vnfH*/*anfH*) were scarce at all depths (<5% overall); two *vnfD* genes

from metagenomes showed phylogenetic similarity to those from soil *Proteobacteria* (Fig. S4).

Attempts to amplify *vnfD/anfD* from cDNA yielded few reads; those recovered were most

similar to *Alphaproteobacteria anfD* from *Rhodopseudomonas* species (Fig. S4).

Methane-related gene expression and phylogeny. Like *nifH*, particulate methane

monooxygenase (*pmoA)* and methyl coenzyme M reductase (*mcrA*) transcripts showed the

highest abundance in surface peat (Fig. S3b,c). Surface *pmoA* transcripts mapped to

Methylocystaceae (75%) and *Methylococcaceae* (25%). Surface *mcrA* transcripts mapped to

Methanosarcina (58%), *Methanocella* (28%), and *Methanoregula* (11%). Attempts to amplify

mmoX from cDNA were unsuccessful (data not shown). In metagenomes, genes for *pmoA* were

dominant in surface peat, whereas the relative abundance of *mmoX* sequences increased with

depth (Fig. S2).

Rates of diazotrophy and methanotrophy. Potential rates of acetylene reduction were

measured for peat collected from S1 bog hollows and hummocks in April, June, August and

September 2013-2015 and incubated for 1 week at 25°C. Acetylene reduction to ethylene was

only detected in surface (0-10 cm) peat samples incubated in the light, and not in deep (10-30

cm) peat, nor in surface peat incubated in the dark. Acetylene reduction to ethane was not

detected in any incubation (data not shown). *Sphagnum* in peat incubations exposed to light

became visibly greener over the course of the incubation (Fig. 4).

In hollows, where surface peat was dominantly covered by a mix of *S. fallax* and *S.*

angustifolium, ARA rates were higher and more variable in degassed vs. oxic incubations (0-163

vs. 2-23 µmol C_2H_4 g^{-1} h^{-1}), and were unaffected by the presence or absence of 20% CO_2. ARA

in hollows incubated with degassed headspace was positively correlated ($P < 0.0001$) with peat

water content (93 to 96%). In both hollows and hummocks, ARA rates were not affected by

addition of 1% CH_4. In hummocks with lower water content (90-91%), surface peat was

dominantly covered by *S. magellanicum*, and oxic and degassed treatments had similarly low

ARA rates (0-8 µmol C_2H_4 g^{-1} h^{-1}). *nifH* transcripts in surface peat from hollows incubated under

degassed headspace with 1% C_2H_2, with or without 1% CH_4, ranged from 10^4 - 10^7 copies g^{-1}

(July 2014; data not shown), which was 1-4 orders of magnitude lower than field samples from

the previous summer (10^8 copies g^{-1}; Fig. 1), and higher and more variable in hollows than *nifH*

transcripts from hummock incubations (Fig. S5).

$^{15}N_2$ incorporation showed similar overall trends as ARA (e.g. 90% higher rates in

degassed vs. oxic conditions, and no significant CH_4 effect; Fig. 5). In degassed treatments, 1%

C_2H_2 inhibited $^{15}N_2$ incorporation by 55% but had no effect on oxic treatments. In oxic

treatments, CH_4 consumption rates were 100 times higher than $^{15}N_2$ incorporation rates, and 1%

C_2H_2 addition suppressed CH_4 oxidation rates by 95% (Fig. 5). Using the four sites measured

with both methods, a conversion factor of 3.9 for $^{15}N_2$-to-ARA was calculated (Fig. S6). In sum,

laboratory incubations of native peats revealed that diazotrophy was stimulated by light,

suppressed by O_2, and minimally affected by CH_4 and CO_2.

Discussion

Diazotrophs are active in surface peat. By definition, the only source of nutrients to

ombrotrophic peat bogs is the atmosphere (65). Scarce (low µM) dissolved nitrogen and

phosphorus in S1 peat pore water suggests oligotrophic conditions (65, 89, 97), consistent with

low rates of atmospheric deposition in Minnesota peat bogs (43). Low nutrient concentrations

add further evidence to previous suggestions that these nutrients may limit *Sphagnum*

productivity (60) and/or complex mechanisms may exist for nutrient scavenging at ultra-low

concentrations. Diazotroph activity (ARA and *nifH* transcription) was solely detected in surface

peat samples incubated in the light. The apparent absence of diazotrophy at greater depths is

consistent with previous reports (86), and may be due to light limitation and/or remineralization

of organic nitrogen to ammonium, which is preferentially used as a nitrogen source by microbes.

Alphaproteobacteria (Rhizobiales) dominate peatland nitrogenase sequences.

Diazotrophic methanotrophs have the potential to serve as a methane biofilter and a nitrogen

source for peatland ecosystems. Previous work showed that *Alphaproteobacteria (Rhizobiales),*

including Type II methanotrophs (63, 99), were the dominant diazotrophs in *Sphagnum*-

dominated peatlands and may provide the unaccounted nitrogen input resulting from an

imbalance in atmospheric nitrogen deposition and accumulation in *Sphagnum* mosses (44, 54,

60, 83, 99). However, the carbon metabolism of peatland diazotrophs remains unclear because

the two dominant *Rhizobiales* families grow on both complex organics (*Bradyrhizobiaceae* (32))

as well as simple alkanes and C1 compounds (*Beijerinckiaceae* (19)). Consistent with previous

studies, *Rhizobiales* showed the highest relative abundance in transcript libraries in this study.

Beijerinckiaceae and *Bradyrhizobiaceae* dominated *nifH* and *nifD* amplicons, respectively.

However, the complex taxonomic classification of *Rhizobiales nifH* and *nifD* genes (22)

prevented distinguishing methanotrophic *Beijerinckiaceae* from heterotrophic *Beijerinckia*

indica and *Bradyrhizobiaceae* solely on the basis of *nifH* and *nifD* phylogenies.

Diazotrophy is catalyzed by the molybdenum form of nitrogenase. Peatland conditions

such as low pH, nitrogen, and temperature would be expected to favor diazotrophy by alternative

nitrogenases. Molybdenum sorption to peat is enhanced at low pH (6), biological requirements

for Mo are higher when bacteria are fixing N_2 than growing on other nitrogen compounds (33),

and alternative nitrogenases have higher activity and expression at lower temperatures (74, 101).

Below 10 nM Mo, diazotrophy is limited in laboratory cultures (1, 26, 34, 87, 108) and

alternative nitrogenases, if present, are expressed (3). Transcription of alternative nitrogenase

genes has been reported for *Peltigera* cyanolichens (45), but to our knowledge has not previously

been investigated for *Sphagnum* peatlands.

Molybdenum concentrations in S1 bog porewaters were 1-7 nM, which is within the

same range as Mo in other oligotrophic freshwaters (33), and lower than V (5-21 nM) and Fe (7-

35 μM). Similar metal concentrations between: (a) sampling dates in September 2014, June

2015, and September 2015; (b) in another ombrotrophic Minnesota peatland, Zim bog, ~80 km

from the S1 bog; and (c) in acidic peatlands in Northern Europe (53) suggest that the values we

measured are spatially and temporally representative for diverse northern peatlands.

Intriguingly, despite the presence of conditions that would be expected to favor

alternative nitrogenase expression (low pH, long winters, and Fe>V>Mo), our collective

evidence suggests that diazotrophy at the S1 bog was catalyzed by the Mo-containing form of the

nitrogenase enzyme. The majority of *nifH* genes retrieved from metagenomes belonged to Mo-

containing clusters I and III. A significant number of sequences came from uncharacterized

Cluster IV, recently shown to contain a functional nitrogenase (110) that likely binds a Mo-Fe

cofactor (72). The metagenomes contained very few cluster II *vnfD/anfD* genes, and minimal

vnfD/anfD transcripts were amplified from peat cDNA. Ethane, a biomarker of alternative

nitrogenase, was undetectable in ARA incubations. Finally, the $^{15}N_2$-to-ARA conversion factor

(3.9) was within the same range (3-4) as other peat bogs (41, 62, 99), and matched Mo-

nitrogenase in pure culture experiments as opposed to the lower values measured for alternative

nitrogenases (4). The question of how peatland diazotrophs access scarce Mo remains uncertain;

it is possible that Mo bound to peat organic matter can be scavenged by diazotrophs as is the case

in forest soils (71, 104).

Methanotrophs are active in incubations and inhibited by acetylene. We performed

bottle experiments to test whether methanotrophs were active. Complete CH_4 consumption in air

amended with 1% CH_4 showed that methanotrophs were active in our incubations, and that the

black bromobutyl stoppers we used were not toxic to peatland methanotrophs, whereas non-

halogenated stoppers are toxic to aquatic aerobic methanotrophs (76). Acetylene fully inhibited

CH_4 consumption, demonstrating that the methanotrophs in our incubations were C_2H_2 sensitive,

similar to laboratory strains tested in previous studies (16, 94). Like diazotrophy, methanotrophy

was apparently also mediated by the enzyme requiring the scarcer metal; dissolved Fe was

consistently orders of magnitude higher than Cu in peat porewater, yet *pmoA* sequences were

more abundant than *mmoX* sequences in surface peat where the highest CH_4 consumption was

observed (24, 68). This finding is also consistent with higher transcription of *pmoA* vs. *mmoX* in

other acidic peatlands (10, 53, 63). In laboratory studies, the "copper switch" from sMMO to

pMMO supported growth occurs when Cu > 1 μM (90), which is several orders of magnitude

higher than Cu concentrations measured at the S1 bog (7-38 nM). It is possible that the "copper

switch" occurs at a lower threshold in peat bogs, or that there are other factors controlling the

type of MMO expressed, such as CH_4 or O_2 availability. Additionally, the inherent nature of the

peat matrix, with characteristically high levels of particulate and dissolved organic matter, likely

also affects metal bioavailability in complex ways (98) not addressed in our study.

Surface peatland diazotrophy is sensitive to oxygen and acetylene. If methanotrophs

were the dominant diazotrophs in peatlands, as previously proposed (60, 99), CH_4 addition

should have stimulated $^{15}N_2$ incorporation in our bottle experiments, and C_2H_2 should have

inhibited it. Instead, this and previous work (61) found that $^{15}N_2$ incorporation was not affected

by CH_4 addition. Acetylene partially inhibited $^{15}N_2$ incorporation under degassed conditions (as

in Kox, et al. (59)) and had minimal effect on oxic $^{15}N_2$ incorporation, suggesting that C_2H_2- and

O_2-sensitive microbial clades contributed approximately half of the diazotrophic activity in our

incubations. This finding highlights the importance of quantifying peatland diazotrophy by $^{15}N_2$

incorporation instead of, or in addition to, ARA.

Based on *nif, pmoA* and *mcrA* phylogeny from native peat, the O_2- and C_2H_2-sensitive

diazotrophs were likely methanotrophic *Beijerinckiaceae* that can only fix N_2 under micro-oxic

conditions, and/or strict anaerobes in cluster III, such as methanogenic *Euryarchaeota*. These

two families are the most active in surface peat based on transcript and amplicon sequences.

Since *nifH* transcripts were 1-4 orders of magnitude lower in our incubations than native peat, it

is likely that diazotrophs were stressed, possibly due to O_2 exposure during the sampling process.

Indeed, we observed highest ARA rates in hollows where the water table was typically at the bog

surface, limiting O_2 penetration into the peat. The other half of O_2-sensitive diazotrophic activity

was likely performed by C_2H_2-insensitive, heterotrophic *Bradyrhizobiaceae* and/or

Beijerinckiaceae (51, 91), or C_2H_2-insensitive, methanotrophic *Methylocystaceae*, which can

adapt to a wide range of CH_4 concentrations (17). Less O_2-sensitive microbes, including

heterotrophic *Beijerinckiaceae* (91) and/or photosynthetic *Oscillatoriophycideae* (25), likely

contributed to the minor amount of ARA activity in the presence of O_2.

Molecular markers for diazotrophy. We end with a word of caution with regard to the

molecular detection of diazotrophs. The majority of studies in peatlands have employed PCR

amplification and sequencing of the *nifH* marker gene for studying the dynamics of diazotrophs

in peatlands. A wide range of *nifH* primer sets exist, with varying universality (29). Peat bog

sequencing efforts have used polF/polR (this study; 63), F1/R6 (99), FGPH19/polR+polF/AQER

(61), and 19F/nifH3+nifH1(1)/nifH2(2) with nested PCR (8, 59). *In silico* evaluation predicts

that the polF/polR primer set will not amplify the majority of *Proteobacteria* and/or

Cyanobacteria and Group III *nifH* sequences (29), however this primer set yields the highest

efficiency for qPCR (30). Of the *nifH* primer sets used previously, F1/R6, 19F/nifH3 and

nifH1(1)/nifH2 are predicted to have the highest coverage for soils (>80% predicted primer

binding for sequences from soil ecosystems). However, it is important to be aware that the F1/R6

primer set contains a number of mismatches with cluster III in peatlands, including

methanogenic *Euryarchaeota* represented (Fig. S7). In order to maximize sequence coverage, we

suggest primer sets that can amplify *nifH* from cluster III, such as IGK(3)/DVV (29), for future

studies.

Conclusions. This study revealed that peatland diazotrophs preferentially transcribed the

Mo-based, rather than the V-based, form of the nitrogenase enzyme, despite the dominance of V

over Mo in the environment. It also highlighted the sensitivity of diazotrophic peatland

communities to O_2 exposure during sample collection, and quantified the inhibitory effect of

C_2H_2 addition on peatland diazotrophy. Under our experimental conditions in lab incubations, we

did not observe CH_4-stimulated diazotrophy. However, quantification of the relative

contributions of methanotrophic and heterotrophic diazotrophy *in situ* awaits further

investigation.

Acknowledgments

This research was funded by DOE support to J.E.K. under grant numbers DE-SC0007144 and

DE-SC0012088, and was supported by NASA Exobiology grant NNX14AJ87G to J.B.G. We

thank Loren Dean Williams for inspiration with visuals, Heather Dang (UC Berkeley) for IRMS

analysis, and Will Overholt for technical assistance. Work at LLNL was conducted under the

auspices of DOE Contract DE-AC52-07NA27344, with funding provided by LDRD 14-ER-038.

Table 1. qPCR and sequencing primers used in this study. PCR conditions were based on

refs. (24), (63), (93), (29), and (4) for *pmoA, mmoX, mcrA*, *nifH*, and *nifD/vnfD/anfD*,

respectively.

Gene	Primer set	Reference
mmoX	mmoX-206f/mmoX-886r	(48)
pmoA	A189f/Mb661r	(46)
mcrA	Mlas/mcrA-rev	(93)
nifH	polF/polR	(56)
nifD	1) nifD820F/nifD1389R 2) nifD820F/nifD1331R	(4)
vnfD/anfD	1) vnfD_anfD548F/vnfD_anfD1337R 2) vnfD_anfD548F/vnfD_anfD1291R	(4)

Figure Captions

Figure 1. Depth profiles of NH_4^+ concentrations (black circles) and *nifH* gene copies (white squares) and transcripts (white triangle) in units of copies per gram of dry weight for the S1 bog T3 mid site. Ammonium concentrations are means of measurements from May, June and September 2014. *nifH* copy numbers are from July 2013. Error bars are standard errors. *nifH* transcripts were not detected at 20, 30 and 75 cm depths. Surface (0-10 cm) and deep (10-30 cm) peat depth intervals used for rate measurements are designated by dashed lines. The maximum water table depth at the S1 site in July 2013 was 2 cm below the hollow surface (dotted line) (38).

Figure 2. Numbers of cDNA amplicon sequence reads for *nifH* and *nifD* alphaproteobacterial transcripts. Primer sets were polF/polR and nifD820F/nifD1331R for *nifH* and *nifD*, respectively.

Figure 3. Dissolved molybdenum (white) and vanadium (gray) concentrations in pore water from three depths in S1 peat hollows (mid and far sites along T3 transect) from June 2015. Pie charts show the relative abundance of genes encoding the five nitrogenase H-subunit clusters from metagenomes for each depth; clusters I and III encode Mo-Fe nitrogenases (*nifH)*; cluster II encodes alternative (*vnfH, anfH*) nitrogenases; cluster IV encode nitrogenase paralogs. Deepest metagenomes were from 75 cm; insufficient numbers of nitrogenase H-subunit sequences were recovered from the far site for cluster analysis.

Figure 4. Acetylene reduction rates for hummocks (90-91% water content) and hollows

(93-96% water content) at the S1 bog, T3 transect (0-10 cm depth incubated in the light at

25°C for 7 days). ARA units are μmol ethylene produced per gram of dry weight per hour.

Photo insets show dominant *Sphagnum species* in (a) hummocks (*S. magellanicum*) and (b)

hollows (*S. fallax* and *S. angustifolium*). Photo inset (c) shows *Sphagnum* greening after

incubation of hollow samples in the light for 7 days at 25°C; bottle B2* (April 2014) received air

headspace with 1% CH_4 and treatment D2 (Sept 2013) received air headspace without CH_4. The

vertical dotted line divides the hummock samples (dominated by S. *magellanicum*) from the

hollow samples (dominated by *S. fallax/angustifolium*) in terms of water content (92%).

Figure 5. Effect of 1% C_2H_2 on $^{15}N_2$ incorporation and CH_4 consumption for S1 bog surface

peat. Rates were measured for samples collected from the NW S1 bog transect in September

2014. $^{15}N_2$ incorporation treatment conditions were 80%N_2+20% CO_2±1% CH_4 (shaded bars) or

80%N_2+20% O_2±1% CH_4 (white bars), with (a) and without (b) 1% C_2H_2; units are nmol $^{15}N_2$

incorporated per gram of dry weight per hour. CH_4 consumption treatments were 80%N_2+20%

CO_2+1% CH_4 (black circles) with (a) and without (b) 1% C_2H_2; units are nmol CH_4 consumed

per gram of dry weight per hour. Error bars are standard errors. Different letters indicate

statistically different elemental contents ($p < 0.05$ based on Tukey-Kramer HSD test).

References

1. **Attridge, E. M.,P. Rowell.** 1997. Growth, heterocyst differentiation and nitrogenase activity in the cyanobacteria *Anabaena variabilis* and *Anabaena cylindrica* in response to molybdenum and vanadium. New Phytologist **135:**517-526.

2. **Basilier, K., U. Granhall,T.-A. Stenström.** 1978. Nitrogen fixation in wet minerotrophic moss communities of a subarctic mire. Oikos**:**236-246.

3. **Bellenger, J. P., T. Wichard, Y. Xu,A. M. L. Kraepiel.** 2011. Essential metals for nitrogen fixation in a free living N2 fixing bacterium: chelation, homeostasis and high use efficiency. Environ Microbiol **13:**1395–1411.

4. **Bellenger, J. P., Y. Xu, X. Zhang, F. M. M. Morel,K. A.M.L.** 2014. Possible contribution of alternative nitrogenases to nitrogen fixation by asymbiotic N2-fixing bacteria in soils. Soil Biol Biochem **69:**413-420.

5. **Berg, A., Å. Danielsson,B. H. Svensson.** 2013. Transfer of fixed-N from N2-fixing cyanobacteria associated with the moss *Sphagnum riparium* results in enhanced growth of the moss. Plant Soil **362:**271-278.

6. **Bertine, K.** 1972. The deposition of molybdenum in anoxic waters. Marine Chemistry **1:**43-53.

7. **Betancourt, D. A., T. M. Loveless, J. W. Brown,P. E. Bishop.** 2008. Characterization of diazotrophis containing Mo-independent nitrogenases, isolated from diverse natural environments. Appl Environ Microbiol **74:**3471-3480.

8. **Bragina, A., S. Maier, C. Berg, H. Müller, V. Chobot, F. Hadacek,G. Berg.** 2011. Similar diversity of alphaproteobacteria and nitrogenase gene amplicons on two related *Sphagnum* mosses. Front Microbiol **2:**275. doi: 210.3389/fmicb.2011.00275.

9. **Bragina, A., L. Oberauner ‐ Wappis, C. Zachow, B. Halwachs, G. G. Thallinger, H. Müller,G. Berg.** 2014. The *Sphagnum* microbiome supports bog ecosystem functioning under extreme conditions. Mol Ecol **23:**4498-4510.

10. **Chen, Y., M. G. Dumont, N. P. McNamara, P. M. Chamberlain, L. Bodrossy, N. Stralis-Pavese,J. C. Murrell.** 2008. Diversity of the active methanotrophic community in acidic peatlands as assessed by mRNA and SIP-PLFA analyses. Environ Microbiol **10:**446-459.

11. **Chen, Y., M. G. Dumont, J. D. Neufeld, L. Bodrossy, N. Stralis-Pavese, N. P. McNamara, N. Ostle, M. J. Briones,J. C. Murrell.** 2008. Revealing the uncultivated majority: combining DNA stable isotope probing, multiple displacement amplification and metagenomic analyses of uncultivated *Methylocystis* in acidic peatlands. Environ Microbiol **10:**2609-2622.

12. **Clymo, R.** 1984. The limits to peat bog growth. Philos T Roy Soc B **303:**605-654.

13. **Clymo, R.,P. Hayward.** 1982. The ecology of *Sphagnum*, p. 229-289, Bryophyte Ecology. Springer.

14. **Colby, J.,H. Dalton.** 1976. Some properties of a soluble methane mono-oxygenase from *Methylococcus capsulatus* strain Bath. Biochem J **157:**495-497.

15. **Crombie, A. T.,J. C. Murrell.** 2014. Trace-gas metabolic versatility of the facultative methanotroph *Methylocella silvestris*. Nature **510:**148–151.

507   16.   **Dalton, H.,R. Whittenbury.** 1976. The acetylene reduction technique as an assay for
508   nitrogenase activity in the methane oxidizing bacterium *Methylococcus capsulatus* strain
509   Bath. Arch Microbiol **109:**147-151.
510   17.   **Dam, B., S. Dam, J. Blom,W. Liesack.** 2013. Genome analysis coupled with
511   physiological studies reveals a diverse nitrogen metabolism in *Methylocystis* sp. strain
512   SC2. PLoS One **8:**e74767.
513   18.   **Darnajoux, R., J. Constantin, J. Miadlikowska, F. Lutzoni,J. P. Bellenger.** 2014. Is
514   vanadium a biometal for boreal cyanolichens? New Phytol **202:**765-771.
515   19.   **Dedysh, S.,P. Dunfield.** 2016. Beijerinckiaceae. Bergey's Manual of Systematics of
516   Archaea and Bacteria.
517   20.   **Dedysh, S. N., W. Liesack, V. N. Khmelenina, N. E. Suzina, Y. A. Trotsenko, J. D.**
518   **Semrau, A. M. Bares, N. S. Panikov,J. M. Tiedje.** 2000. *Methylocella palustris* gen.
519   nov., sp. nov., a new methane-oxidizing acidophilic bacterium from peat bogs,
520   representing a novel subtype of serine-pathway methanotrophs. Int J Syst Evol Microbiol
521   **50:**955-969.
522   21.   **Dedysh, S. N., N. S. Panikov, W. Liesack, R. Großkopf, J. Zhou,J. M. Tiedje.** 1998.
523   Isolation of acidophilic methane-oxidizing bacteria from northern peat wetlands. Science
524   **282:**281-284.
525   22.   **Dedysh, S. N., P. Ricke,W. Liesack.** 2004. NifH and NifD phylogenies: an evolutionary
526   basis for understanding nitrogen fixation capabilities of methanotrophic bacteria.
527   Microbiology **150:**1301-1313.
528   23.   **Dilworth, M. J., R. R. Eady, R. L. Robson,R. W. Miller.** 1987. Ethane formation from
529   acetylene as a potential test for vanadium nitrogenase *in vivo*. Nature **327:**167-168.
530   24.   **Esson, K. C., X. Lin, D. Kumaresan, J. P. Chanton, J. C. Murrell,J. E. Kostka.** 2016.
531   Alpha-and gammaproteobacterial methanotrophs codominate the active methane-
532   oxidizing communities in an acidic boreal peat bog. Appl Environ Microbiol **82:**2363-
533   2371.
534   25.   **Fay, P.** 1992. Oxygen relations of nitrogen fixation in cyanobacteria. Microbio Mol Biol
535   Rev **56:**340-373.
536   26.   **Fay, P.,L. Vasconcelos.** 1974. Nitrogen metabolism and ultrastructure in *Anabaena*
537   *cylindrica*. 2. Effect of molybdenum and vanadium. Archiv. Microbiol. **99:**221-230.
538   27.   **Fulweiler, R. W., E. M. Heiss, M. K. Rogener, S. E. Newell, G. R. LeCleir, S. M.**
539   **Kortebein,S. W. Wilhelm.** 2015. Examining the impact of acetylene on N-fixation and
540   the active sediment microbial community. Front Microbiol **6**.
541   28.   **Gaby, J. C.,D. H. Buckley.** 2014. A comprehensive aligned nifH gene database: a
542   multipurpose tool for studies of nitrogen-fixing bacteria. Database **2014:**bau001.
543   29.   **Gaby, J. C.,D. H. Buckley.** 2012. A comprehensive evaluation of PCR primers to
544   amplify the *nifH* gene of nitrogenase. PLoS One **7:**e42149.
545   30.   **Gaby, J. C.,D. H. Buckley.** 2017. The use of degenerate primers in qPCR analysis of
546   functional genes can cause dramatic quantification bias as revealed by investigation of
547   *nifH* primer performance. Microbial Ecol **doi:10.1007/s00248-017-0968-0**.
548   31.   **García-Robledo, E., A. Corzo,S. Papaspyrou.** 2014. A fast and direct
549   spectrophotometric method for the sequential determination of nitrate and nitrite at low
550   concentrations in small volumes. Marine Chemistry **162:**30-36.
551   32.   **Garrity, G., J. Bell,T. Lilburn.** 2015. Bradyrhizobiaceae fam. nov. Bergey's Manual of
552   Systematics of Archaea and Bacteria**:**1-4.

33.   **Glass, J. B., R. P. Axler, S. Chandra,C. R. Goldman.** 2012. Molybdenum limitation of
microbial nitrogen assimilation in aquatic ecosystems and pure cultures. Front Microbiol
3:331.
34.   **Glass, J. B., F. L. Wolfe-Simon, J. J. Elser,A. D. Anbar.** 2010. Molybdenum-nitrogen
co-limitation in freshwater and coastal heterocystous cyanobacteria. Limnol Oceanogr
55:667-676.
35.   **Gorham, E., C. Lehman, A. Dyke, D. Clymo,J. Janssens.** 2012. Long-term carbon
sequestration in North American peatlands. Quat Sci Rev **58:**77-82.
36.   **Granhall, U.,A. V. Hofsten.** 1976. Nitrogenase activity in relation to intracellular
organisms in Sphagnum mosses. Physiol Plantarum **36:**88-94.
37.   **Granhall, U.,H. Selander.** 1973. Nitrogen fixation in a subarctic mire. Oikos**:**8-15.
38.   **Griffiths, N. A.,S. D. Sebestyen.** 2016. Dynamic vertical profiles of peat porewater
chemistry in a northern peatland. Wetlands **36:**1119-1130.
39.   **Gupta, V., K. A. Smemo, J. B. Yavitt,N. Basiliko.** 2012. Active methanotrophs in two
contrasting North American peatland ecosystems revealed using DNA-SIP. Microbial
Ecol **63:**438-445.
40.   **Hanson, P. J., J. S. Riggs, W. R. Nettles, J. R. Phillips, M. B. Krassovski, L. A.**
Hook, L. Gu, A. D. Richardson, D. M. Aubrecht,D. M. Ricciuto. 2017. Attaining
whole-ecosystem warming using air and deep-soil heating methods with an elevated CO2
atmosphere. Biogeosciences **14:**861-883.
41.   **Hardy, R., R. Burns,R. Holsten.** 1973. Applications of the acetylene-ethylene assay for
measurement of nitrogen fixation. Soil Biol Biochem **5:**47-81.
42.   **Hardy, R. W. F., R. D. Holsten, E. K. Jackson,R. C. Burns.** 1968. The acetylene-
ethylene assay for N2 fixation: laboratory and field evaluation. Plant Physiol **43:**1185-
1207.
43.   **Hill, B. H., T. M. Jicha, L. L. Lehto, C. M. Elonen, S. D. Sebestyen,R. K. Kolka.**
2016. Comparisons of soil nitrogen mass balances for an ombrotrophic bog and a
minerotrophic fen in northern Minnesota. Sci Total Environ **550:**880-892.
44.   **Ho, A.,P. L. Bodelier.** 2015. Diazotrophic methanotrophs in peatlands: the missing link?
Plant Soil **389:**419-423.
45.   **Hodkinson, B., J. Allen, L. Forrest, B. Goffinet, E. Seurasiaux, O. Andresson, V.**
Miao, J. P. Bellenger,F. Lutzoni. 2014. Lichen-symbiotic cyanobacteria associated with
Peltigera have an alternative vanadium-dependent nitrogen fixation system. Eur J Phycol
49:11-19.
46.   **Holmes, A. J., A. Costello, M. E. Lidstrom,J. C. Murrell.** 1995. Evidence that
participate methane monooxygenase and ammonia monooxygenase may be
evolutionarily related. FEMS Microbiology Letters **132:**203-208.
47.   **Huson, D. H., S. Mitra, H.-J. Ruscheweyh, N. Weber,S. C. Schuster.** 2011. Integrative
analysis of environmental sequences using MEGAN4. Genome research **21:**1552-1560.
48.   **Hutchens, E., S. Radajewski, M. G. Dumont, I. R. McDonald,J. C. Murrell.** 2004.
Analysis of methanotrophic bacteria in Movile Cave by stable isotope probing. Environ
Microbiol **6:**111-120.
49.   **Hynes, R.,R. Knowles.** 1982. Effect of acetylene on autotrophic and heterotrophic
nitrification. Can J Microbiol **28:**334-340.
50.   **Keister, D.** 1975. Acetylene reduction by pure cultures of *Rhizobia*. J Bacteriol
123:1265-1268.

51.   **Keister, D. L.,W. R. Evans.** 1976. Oxygen requirement for acetylene reduction by pure
600   cultures of rhizobia. J Bacteriol **127:**149-153.
601   52.   **Kip, N., B. E. Dutilh, Y. Pan, L. Bodrossy, K. Neveling, M. P. Kwint, M. S. Jetten,H.**
602   **J. Op den Camp.** 2011. Ultra-deep pyrosequencing of *pmoA* amplicons confirms the
603   prevalence of *Methylomonas* and *Methylocystis* in *Sphagnum* mosses from a Dutch peat
604   bog. Environ Microbiol Rep **3:**667-673.
605   53.   **Kip, N., W. Ouyang, J. van Winden, A. Raghoebarsing, L. van Niftrik, A. Pol, Y.**
606   **Pan, L. Bodrossy, E. G. van Donselaar,G.-J. Reichart.** 2011. Detection, isolation, and
607   characterization of acidophilic methanotrophs from *Sphagnum* mosses. Appl Environ
608   Microbiol **77:**5643-5654.
609   54.   **Kip, N., J. F. van Winden, Y. Pan, L. Bodrossy, G.-J. Reichart, A. J. Smolders, M. S.**
610   **Jetten, J. S. S. Damsté,H. J. O. den Camp.** 2010. Global prevalence of methane
611   oxidation by symbiotic bacteria in peat-moss ecosystems. Nature Geosci **3:**617-621.
612   55.   **Knorr, K. H., M. A. Horn,W. Borken.** 2015. Significant nonsymbiotic nitrogen fixation
613   in Patagonian ombrotrophic bogs. Glob Change Biol **21:**2357-2365.
614   56.   **Kolb, S., C. Knief, S. Stubner,R. Conrad.** 2003. Quantitative detection of
615   methanotrophs in soil by novel *pmoA*-targeted real-time PCR assays. Appl Environ
616   Microbiol **69:**2423-2429.
617   57.   **Kolka, R., S. Sebestyen, E. S. Verry,K. Brooks.** 2011. Peatland biogeochemistry and
618   watershed hydrology at the Marcell experimental forest. CRC Press.
619   58.   **Kostka, J. E., D. J. Weston, J. B. Glass, E. A. Lilleskov, A. J. Shaw,M. R. Turetsky.**
620   2016. The *Sphagnum* microbiome: new insights from an ancient plant lineage. New
621   Phytol **211:**57-64.
622   59.   **Kox, M. A., C. Lüke, C. Fritz, E. van den Elzen, T. van Alen, H. J. O. den Camp, L.**
623   **P. Lamers, M. S. Jetten,K. F. Ettwig.** 2016. Effects of nitrogen fertilization on
624   diazotrophic activity of microorganisms associated with *Sphagnum magellanicum*. Plant
625   Soil **406:**83-100.
626   60.   **Larmola, T., S. M. Leppänen, E.-S. Tuittila, M. Aarva, P. Merilä, H. Fritze,M.**
627   **Tiirola.** 2014. Methanotrophy induces nitrogen fixation during peatland development.
628   Proc Natl Acad Sci **111:**734-739.
629   61.   **Leppänen, S. M., A. J. Rissanen,M. Tiirola.** 2014. Nitrogen fixation in *Sphagnum*
630   mosses is affected by moss species and water table level. Plant Soil**:**1-12.
631   62.   **Leppänen, S. M., M. Salemaa, A. Smolander, R. Mäkipää,M. Tiirola.** 2013. Nitrogen
632   fixation and methanotrophy in forest mosses along a N deposition gradient. Environ Exp
633   Bot **90:**62-69.
634   63.   **Liebner, S.,M. M. Svenning.** 2013. Environmental transcription of *mmoX* by methane-
635   oxidizing Proteobacteria in a subarctic palsa peatland. Appl Environ Microbiol **79:**701-
636   706.
637   64.   **Limpens, J., F. Berendse, C. Blodau, J. Canadell, C. Freeman, J. Holden, N. Roulet,**
638   **H. Rydin,G. Schaepman-Strub.** 2008. Peatlands and the carbon cycle: from local
639   processes to global implications: a synthesis. Biogeosciences **5:**1475-1491.
640   65.   **Limpens, J., M. M. Heijmans,F. Berendse.** 2006. The nitrogen cycle in boreal
641   peatlands, p. 195-230, Boreal peatland ecosystems. Springer.
642   66.   **Lin, X., S. Green, M. Tfaily, O. Prakash, K. Konstantinidis, J. Corbett, J. Chanton,**
643   **W. Cooper,J. Kostka.** 2012. Microbial community structure and activity linked to

contrasting biogeochemical gradients in bog and fen environments of the Glacial Lake
Agassiz Peatland. Appl Environ Microbiol **78:**7023-7031.

67.   **Lin, X., K. M. Handley, J. A. Gilbert,J. E. Kostka.** 2015. Metabolic potential of fatty
acid oxidation and anaerobic respiration by abundant members of Thaumarchaeota and
Thermoplasmata in deep anoxic peat. ISME J **9:**2740-2744.

68.   **Lin, X., M. M. Tfaily, S. Green, J. M. Steinweg, P. Chanton, A. Imvittaya, J. P.**
Chanton, W. Cooper, C. W. Schadt,J. E. Kostka. 2014. Microbial metabolic potential
for carbon degradation and nutrient (nitrogen and phosphorus) acquisition in an
ombrotrophic peatland. Appl Environ Microbiol **80:**3531-3540.

69.   **Lin, X., M. M. Tfaily, J. M. Steinweg, P. Chanton, K. Esson, Z. K. Yang, J. P.**
Chanton, W. Cooper, C. W. Schadt,J. E. Kostka. 2014. Microbial community
stratification linked to utilization of carbohydrates and phosphorus limitation in a boreal
peatland at Marcell Experimental Forest, Minnesota, USA. Appl Environ Microbiol
80:3518-3530.

70.   **Loveless, T. M., J. R. Saah,P. E. Bishop.** 1999. Isolation of nitrogen-fixing bacteria
containing molybdenum-independent nitrogenases from natural environments. Appl
Environ Microbiol **65:**4223-4226.

71.   **Marks, J., S. Perakis, E. King,J. Pett-Ridge.** 2015. Soil organic matter regulates
molybdenum storage and mobility in forests. Biogeochemistry.

72.   **McGlynn, S. E., E. S. Boyd, J. W. Peters,V. J. Orphan.** 2013. Classifying the metal
dependence of uncharacterized nitrogenases. Front Microbiol **3:**419.

73.   **McRose, D. L., X. Zhang, A. M. Kraepiel,F. M. Morel.** 2017. Diversity and activity of
alternative nitrogenases in sequenced genomes and coastal environments. Front
Microbiol **8:**267.

74.   **Miller, R.,R. Eady.** 1988. Molybdenum and vanadium nitrogenases of *Azotobacter*
chroococcum. Low temperature favours N2 reduction by vanadium nitrogenase. Biochem
J **256:**429-432.

75.   **Murphy, J.,J. Riley.** 1962. A modified single solution method for the determination of
phosphate in natural waters. Anal Chim Acta **27:**31-36.

76.   **Niemann, H., L. Steinle, J. Blees, I. Bussmann, T. Treude, S. Krause, M. Elvert,M.**
F. Lehmann. 2015. Toxic effects of lab‐grade butyl rubber stoppers on aerobic
methane oxidation. Limnol Oceangr: Methods **13:**40-52.

77.   **Normand, P., P. Simonet,R. Bardin.** 1988. Conservation of *nif* sequences in *Frankia*.
Mol Gen Genet **213:**238-246.

78.   **Opelt, K., C. Berg, S. Schonmann, L. Eberl,G. Berg.** 2007. High specificity but
contrasting biodiversity of *Sphagnum*-associated bacterial and plant communities in bog
ecosystems independent of the geographical region. The ISME Journal **1:**502-516.

79.   **Opelt, K., V. Chobot, F. Hadacek, S. Schonmann, L. Eberl,G. Berg.** 2007.
Investigations of the structure and function of bacterial communities associated with
Sphagnum mosses. Environ Microbiol **9:**2795-2809.

80.   **Oremland, R. S.,B. F. Taylor.** 1975. Inhibition of methanogenesis in marine sediments
by acetylene and ethylene: validity of the acetylene reduction assay for anaerobic
microcosms. Appl Microbiol **30:**707-709.

81.   **Payne, W.,M. Grant.** 1982. Influence of acetylene on growth of sulfate-respiring
bacteria. Appl Environ Microbiol **43:**727-730.

82.   **Prior, S. D.,H. Dalton.** 1985. Acetylene as a suicide substrate and active site probe for
690   methane monooxygenase from *Methylococcus capsulatus* (Bath). FEMS Microbiol Lett
691   **29:**105-109.
692   83.   **Raghoebarsing, A. A., A. J. Smolders, M. C. Schmid, W. I. C. Rijpstra, M. Wolters-**
693   **Arts, J. Derksen, M. S. Jetten, S. Schouten, J. S. S. Damsté, L. P. Lamers, J. G. M.**
694   **Roelofs, H. J. M. Op den Camp,M. Strous.** 2005. Methanotrophic symbionts provide
695   carbon for photosynthesis in peat bogs. Nature **436:**1153-1156.
696   84.   **Rahman, M. T., A. Crombie, Y. Chen, N. Stralis-Pavese, L. Bodrossy, P. Meir, N. P.**
697   **McNamara,J. C. Murrell.** 2011. Environmental distribution and abundance of the
698   facultative methanotroph Methylocella. The ISME journal **5:**1061-1066.
699   85.   **Raymond, J., J. L. Siefert, C. R. Staples,R. E. Blankenship.** 2004. The natural history
700   of nitrogen fixation. Mol Biol Evol **21:**541-554.
701   86.   **Rosswall, T.,U. Granhall.** 1980. Nitrogen cycling in a subarctic ombrotrophic mire.
702   Ecological Bulletins**:**209-234.
703   87.   **Scherer, P.** 1988. Vanadium and molybdenum requirement for the fixation of molecular
704   nitrogen by two *Methanosarcina* strains. Arch. Microbiol. **151:**44-48.
705   88.   **Schwintzer, C. R.** 1983. Nonsymbiotic and symbiotic nitrogen fixation in a weakly
706   minerotrophic peatland. Am J Bot **70:**1071-1078.
707   89.   **Sebestyen, S. D., C. Dorrance, D. M. Olson, E. S. Verry, R. K. Kolka, A. E. Elling,R.**
708   **Kyllander.** 2011. Long-term monitoring sites and trends at the Marcell Experimental
709   Forest. Peatland Biogeochemistry and Watershed Hydrology at the Marcell Experimental
710   Forest**:**93-134.
711   90.   **Semrau, J. D., A. A. DiSpirito,S. Yoon.** 2010. Methanotrophs and copper. FEMS
712   Microbiol Rev **34:**496-531.
713   91.   **Spiff, E.,C. Odu.** 1973. Acetylene reduction by Beijerinckia under various partial
714   pressures of oxygen and acetylene. Microbiology **78:**207-209.
715   92.   **Sprott, G., K. Jarrell, K. Shaw,R. Knowles.** 1982. Acetylene as an inhibitor of
716   methanogenic bacteria. J Gen Microbiol **128:**2453-2462.
717   93.   **Steinberg, L. M.,J. M. Regan.** 2008. Phylogenetic comparison of the methanogenic
718   communities from an acidic, oligotrophic fen and an anaerobic digester treating
719   municipal wastewater sludge. Appl Environ Microbiol **74:**6663-6671.
720   94.   **Stirling, D. I.,H. Dalton.** 1977. Effect of metal-binding agents and other compounds on
721   methane oxidation by two strains of *Methylococcus capsulatus*. Arch Microbiol **114:**71-
722   76.
723   95.   **Strickland, J.,T. Parsons.** 1972. A practical handbook of seawater analysis, vol. 167.
724   96.   **Tamura, K., D. Peterson, N. Peterson, G. Stecher, M. Nei,S. Kumar.** 2011. MEGA5:
725   molecular evolutionary genetics analysis using maximum likelihood, evolutionary
726   distance, and maximum parsimony methods. Mol Biol Evol **28:**2731-2739.
727   97.   **Urban, N.,S. Eisenreich.** 1988. Nitrogen cycling in a forested Minnesota bog. Can J Bot
728   **66:**435-449.
729   98.   **Vaughan, D., D. Lumsdon,D. Linehan.** 1993. Influence of dissolved organic matter on
730   the bio-availability and toxicity of metals in soils and aquatic systems. Chem Ecol **8:**185-
731   201.
732   99.   **Vile, M. A., R. K. Wieder, T. Živković, K. D. Scott, D. H. Vitt, J. A. Hartsock, C. L.**
733   **Iosue, J. C. Quinn, M. Petix,H. M. Fillingim.** 2014. N2-fixation by methanotrophs

sustains carbon and nitrogen accumulation in pristine peatlands. Biogeochemistry
121:317-328.

100.   **Vorobev, A. V., M. Baani, N. V. Doronina, A. L. Brady, W. Liesack, P. F.**
Dunfield,S. N. Dedysh. 2011. *Methyloferula stellata* gen. nov., sp. nov., an acidophilic,
obligately methanotrophic bacterium that possesses only a soluble methane
monooxygenase. Int J Syst Evol Microbiol **61:**2456-2463.

101.   **Walmsley, J.,C. Kennedy.** 1991. Temperature-dependent regulation by molybdenum
and vanadium of expression of the structural genes encoding 3 nitrogenases in
Azotobacter vinelandii. Appl Environ Microbiol **57:**622-624.

102.   **Wedepohl, K. H.** 1995. The composition of the continental crust. Geochem Cosmochim
Acta **59:**1217-1232.

103.   **Weston, D. J., C. M. Timm, A. P. Walker, L. Gu, W. Muchero, J. Schmutz, A. J.**
Shaw, G. A. Tuskan, J. M. Warren,S. D. Wullschleger. 2015. *Sphagnum* physiology in
the context of changing climate: Emergent influences of genomics, modeling and host-
microbiome interactions on understanding ecosystem function. Plant Cell Environ
38:1737-1751.

104.   **Wichard, T., B. Mishra, S. C. B. Myneni, J. P. Bellenger,A. M. L. Kraepiel.** 2009.
Storage and bioavailability of molybdenum in soils increased by organic matter
complexation. Nature Geosci. **2:**625-629.

105.   **Wong, P. P.,R. Burris.** 1972. Nature of oxygen inhibition of nitrogenase from
Azotobacter vinelandii. Proc Natl Acad Sci **69:**672-675.

106.   **Yu, Z.** 2012. Northern peatland carbon stocks and dynamics: a review. Biogeosciences
4071–4085:4071–4085.

107.   **Zehr, J. P., B. D. Jenkins, S. M. Short,G. F. Steward.** 2003. Nitrogenase gene diversity
and microbial community structure: a cross‐system comparison. Environ Microbiol
5:539-554.

108.   **Zerkle, A. L., C. H. House, R. P. Cox,D. E. Canfield.** 2006. Metal limitation of
cyanobacterial N_2 fixation and implications for the Precambrian nitrogen cycle.
Geobiology **4:**285-297.

109.   **Zhao, Y., H. Tang,Y. Ye.** 2012. RAPSearch2: a fast and memory-efficient protein
similarity search tool for next-generation sequencing data. Bioinformatics **28:**125-126.

110.   **Zheng, H., C. Dietrich, R. Radek,A. Brune.** 2016. *Endomicrobium proavitum*, the first
isolate of Endomicrobia class. nov.(phylum Elusimicrobia)–an ultramicrobacterium with
an unusual cell cycle that fixes nitrogen with a Group IV nitrogenase. Environ Microbiol
18:191-204.

NH_4^+ (μM)

water table max

surface

deep

Depth (cm)

□ genes △ transcripts

nifH (copies g^{-1})