1

A Comprehensive Assessment of the Genetic Determinants in *Salmonella* Typhimurium for

Resistance to Hydrogen Peroxide

Sardar Karash[1], Rohana Liyanage[2,3], Abdullah Qassab[1], Jackson O. Lay, Jr.[2,3], and

Young Min Kwon[1,4*]

[1]Cell and Molecular Biology Program, University of Arkansas, Fayetteville, AR 72701,

[2]Department of Chemistry, University of Arkansas, Fayetteville, AR 72701,

[3]Statewide Mass Spectrometry Facility, Fayetteville, AR 72701, and

[4]Department of Poultry Science, University of Arkansas, Fayetteville, AR 72701

*Corresponding author:

Department of Poultry Science

College of Agricultural, Food and Life Sciences

University of Arkansas

Fayetteville, AR 72701

Phone : (479) 575-4935

Email ykwon@uark.edu

ABSTRACT

Salmonella is a Gram-negative bacterium that infects a wide range of hosts. *Salmonella* Typhimurium causes gastroenteritis in human, and can survive and replicate in macrophages. An essential mechanism used by the macrophages to eradicate *Salmonella* is production of reactive oxygen species. Here, we used proteogenomic approaches to determine the candidate genes and proteins that have a role in resistance of *S*. Typhimurium to H_2O_2. For Tn-seq, a highly saturated Tn5 insertion library was grown *in vitro* in Luria-Bertani broth (LB) medium as well as LB containing either 2.5 (H_2O_2L) or 3.5 mM H_2O_2 (H_2O_2H). We identified two sets of overlapping genes that are required for resistance of *S*. Typhimurium to H_2O_2L and H_2O_2H, and the result was validated via phenotypic evaluation of 50 selected mutants. The enriched pathways for resistance to H_2O_2 included DNA repair, aromatic amino acid biosynthesis, Fe-S cluster biosynthesis, iron homeostasis, flagellar genes, H_2O_2 scavenging enzymes, and DNA adenine methylase. Particularly, we identified aromatic amino acid biosynthesis (*aroB*, and *aroK*) and putative iron transporter system (*ybbK*, *ybbL*, and *ybbM*) as novel mechanisms for resistance to H_2O_2. The proteomics revealed that the majority of essential proteins, including ribosomal proteins, were downregulated upon exposure to H_2O_2. A subset of the proteins identified by Tn-seq were analyzed by targeted proteomics, and 70 % of them were upregulated upon exposure to H_2O_2. The identified candidate genes will deepen our understanding on the mechanisms of *S*. Typhimurium survival in macrophages, and can be exploited to develop new antimicrobial drugs.

IMPORTANCE

Salmonella infection is frequently caused by consumption of contaminated food or water. The

infection may lead to gastroenteritis or typhoid fever, depending on the *Salmonella* serovars. Even

though the bacterium encounters the immune defense arsenals of the infected host, including

reactive oxygen species in phagocytes, the bacterium can survive and replicate. In this study,

proteogenomic approaches were used in order to identify the genes and proteins that have a role

in resistance to H_2O_2. In addition to the H_2O_2 scavenging and degrading enzymes, aromatic amino

acid biosynthesis and iron homeostasis were identified among the most important pathways for

H_2O_2 resistance. These findings will deepen our knowledge on the mechanisms of *Salmonella*

survival in phagocytes and other niches with oxidative stress, and also provides novel targets to

develop new antimicrobial therapeutics.

56
57

Introduction

Salmonella is a Gram-negative bacterium that infects humans and animals. *Salmonella enterica*

has numerous serovars, which include typhoidal and non-typhoidal strains. In contrast to the

typhoidal salmonellae which are human restricted pathogens, the non-typhoidal salmonellae

(NTS), serovar Enteritidis and Typhimurium, are able to infect a wide range of hosts, causing

gastroenteritis (1). The NTS strains, including *Salmonella enterica* serovar Typhimurium, account

for 11% (1.2 million cases) of the total foodborne illnesses caused by different pathogens in the

United States (2). It has been estimated that *Salmonella* is responsible for 93.8 million cases of

gastroenteritis, leading to 155,000 deaths worldwide annually (3). The pathogen remains a

continuous threat to the food safety, and public health.

To initiate an infection and survive inside the host, *Salmonella* needs to overcome a myriad of host

defense mechanisms. As *Salmonella* reaches the intestine and breaches the epithelial tissue, it

enters the macrophages and activates different virulence strategies in order to survive and replicate

in them (4). An essential mechanism used by the phagocytes to kill and eradicate *Salmonella* is

production of reactive oxygen species (ROS). Hydrogen peroxide (H_2O_2), superoxide anion (O_2^-

), and the hydroxyl radical ($HO^.$) are derivatives of ROS. The short-lived O_2^-, produced by the

NADPH-dependent phagocytic oxidase, quickly dismutates into H_2O_2, which diffuses across

semipermeable bacterial cell membranes. Eventually, H_2O_2 reduces to $HO^.$ by Fe^{2+} via Fenton

reaction (5-7). The ROS, including H_2O_2, can damage DNA, iron-sulfur cluster-containing

proteins, and other biological molecules in the bacterial cells (8-10).

Numerous genetic factors and proteins that are important for resistance of *S.* Typhimurium to H_2O_2

have been discovered and the underlying mechanisms have been explored (11,12). Various

approaches and techniques have been employed to study global response of *Salmonella* or related

bacteria to H_2O_2 *in vitro* as a model system to simulate the bacterium's response to ROS in

phagocytic cells: (i) Two-dimensional gel electrophoresis identified H_2O_2-induced proteins in

Salmonella (13), (ii) DNA microarray identified H_2O_2 induced genes in *E. coli* (14), and (iii) RNA-

seq identified H_2O_2 induced genes in *Salmonella* (15). However, the factors required for fitness

under the given condition cannot be identified with high confidence based on the analysis of

transcriptomics or proteomics data (16). Microarray-based tracking of random transposon

insertions was used to identify numerous genes in *Salmonella* that are required for survival in mice

and macrophages (17-18). However, the genetic factors responsible for resistance to ROS cannot

be sorted out among all of the genetic factors identified in the study that are required for fitness in

the presence of multiple host stressors.

To shed more insights into the underlying mechanisms of *Salmonella* resistance to H_2O_2, more

direct approach linking the gene-phenotype relationships in a genome-wide scale would be

necessary. Tn-seq is a powerful approach to allow direct and accurate assessment of the fitness

requirement of each gene on the entire genome of a prokaryotic organism (19). In Tn-seq method,

a saturated transposon insertion library (input) is exposed to a selective condition, and the mutant

population altered through the selection (output) is recovered. Then, the genomic junctions of the

transposon insertions are specifically amplified and sequenced from both input and output pools

by high-throughput sequencing. The gene fitness can be obtained by calculating the change in

relative abundance of the sequence reads corresponding to each gene in the entire genome between

the two pools. Tn-seq has been employed to assign gene functions to *Salmonella* genomes in

numerous studies: (i) Previously, our lab identified conditionally essential genes that are required

for growth in the presence of bile, limited nutrients, and high temperature (20), (ii) The genes

required for intestinal colonization were identified in chickens, pigs, and cattle (21), (iii) Candidate

essential genes and genes contributing toward bile resistance were identified (22), (iv) Core

conserved genes for growth in rich media were identified in serovars Typhi and Typhimurium

(23). In addition to Tn-seq, electrospray ionization liquid chromatography tandem mass

spectrometry (ESI-LC-MS/MS) is a powerful approach in identifying and quantifying proteins in

a large scale. The system-wide protein regulation can be determined using mass spectrometry

signal intensities of tryptic peptides obtained from two different culture conditions (24). The post-

translational modification in proteins can be revealed by using proteomic analysis (25). Many

studies took advantage of proteomic analysis of *Salmonella*. However, to the best of our

knowledge, this study is the first to investigate proteogenomics of a bacterium by combining Tn-

seq and proteome analysis simultaneously to the same stressor.

In this work, we used Tn-seq method and proteomic analysis in combination to determine system-

wide responses of *S.* Typhimurium to two different concentrations of H_2O_2 (H_2O_2L and H_2O_2H).

We obtained a comprehensive list of 137 genes that are putatively required for the resistance of *S.*

Typhimurium 14028 to H_2O_2. The role of 50 selected genes in resistance to H_2O_2 were determined

by phenotypic evaluation of the individual deletion mutants. Also, we identified a set of 246

proteins that are differentially expressed in response to H_2O_2, using data-dependent acquisition

(DDA) proteomics, which are largely overlapped with the genes identified by Tn-seq; targeted

proteomics showed 70% of the proteins identified by Tn-seq were upregulated by H_2O_2. In addition

to the genes of *S.* Typhimurium previously known to be important for resistance to H_2O_2, we

identified approximately 80 genes that have not been previously associated with resistance to

oxidative stress. The result of this study highlighted that the genes in aromatic amino acid

biosynthesis, *aroB* and *aroK*, and iron homeostasis, *ybbK*, *ybbL*, and *ybbM*, are crucially important

for growth fitness under H_2O_2 stress. The identified candidate genes will expand our understanding

on the molecular mechanisms of *Salmonella* survival in macrophages, and serve as new

antimicrobial drug targets.

Results and Discussion

The H_2O_2 concentrations and the selections of Tn5 library

First, we sought to determine the growth response of the wild type *S.* Typhimurium 14028 cells in

LB media containing varying concentrations of H_2O_2. The wild type cells were grown in LB media

that contain different concentrations of H_2O_2 in 96-well plates. After evaluating the growth rates

for the cultures, 2.5 and 3.5 mM H_2O_2 were chosen for Tn-seq selections in our study, and termed

H_2O_2L and H_2O_2H, respectively. In comparison to *Salmonella* grown in LB media with no H_2O_2,

H_2O_2L and H_2O_2H reduced the growth rates by 10% and 28%, respectively (Fig. 1A). The lag time

increased by a 5.7-fold (0.5 vs. 2.9 hr), and an 11-fold (0.5 vs. 5.6 hr) in H_2O_2L and H_2O_2H,

respectively. The maximum OD_{600} decreased by only 1% for the H_2O_2L and 2% for the H_2O_2H in

comparison to LB media (Fig. 1A).

For the selection of Tn5 library, 20 ml cultures in 300 ml Erlenmeyer flasks containing LB, H_2O_2L,

or H_2O_2H were inoculated with the same Tn5 library at the seeding CFUs of the library at 3.5×10^6.

This seeding level provided ~10 CFUs for each Tn5 insertion mutant in the library. The cultures

were grown until the mid-exponential phase, in which the CFUs reached 1.17×10^8 (SE 0.01×10^8).

It required 7.5 and 9.2 h to reach the cell density as measured by optical density for H_2O_2L and

H_2O_2H, respectively, in contrast to 5 h for LB medium (Fig. 1B). We observed some differences

in growth responses between the cultures in a 96-well plate and in a 300-ml Erlenmeyer flask. The

optical density readings by the plate reader was different in comparison to those by Bio-photometer

that we used to measure optical density of the culture in the flask. As a result, the growth curve in

Fig.1.A which was based on 96-well plate reader, dose not match exactly with the time required

for the Tn5 library to reach the target mid-exponential phase in the flask cultures. In addition, we

observed that the H_2O_2 is stable in LB media free of *Salmonella* during the window of time used

for the library selection process (Fig.S1), which was also supported by Bogomolnaya et al. (26).

Preparation of Tn5-seq amplicon library

The *Salmonella* mutants were generated by using the delivery plasmid pBAM1 via conjugation. A

total of 325,000 mutant colonies were recovered from 50 plates. Each mutant contained a single

random insertion of Tn5 transposon in the chromosome or plasmid according to DNA sequencing

of Tn5-junction sequences for a small set (n = 71) of randomly selected Tn5 mutants. We found a

significant portion (~20%) of the mutants in the library that were not genuine Tn5 insertions, but

the mutants generated as a result of pBAM1 integration into chromosome as determined by their

ability to grow in the presence of ampicillin. To prevent the Illumina sequencing reads from being

wasted on sequencing Tn5 junctions from these cointegrants, we digested genomic DNA of the

input and output libraries with PvuII, which digests immediately outside the inverted repeats on

both sides of Tn5. The digested DNA was then used to prepare Tn-seq amplicon library as

described in Materials and Methods. Our Tn-seq data analysis indicated that our strategy of

removing the DNA sequences originating from cointegrants was effective because only 0.55% of

the total HiSeq reads corresponding to Tn5-junctions matched to pBAM1. It should be possible to

remove them completely by ensuring complete digestion of genomic DNA with PvuII. The method

for Tn-seq amplicon library we developed and used in this study has multiple advantages over

other Tn-seq protocols, because our method requires only 100 ng of the genomic DNA, and the

whole process can be completed in a day (27). When the extension step in the protocol was

performed using a conventional 20 nucleotide primer, and the final products of exponential PCR

were separated on agarose gel electrophoresis, even the negative controls (the wild type genomic

DNA or mutant library genomic DNA without linear extension) showed smear patterns of

nonspecific background amplification. However, when dual priming oligonucleotide (DPO)

primer was used in place of the conventional primer for linear extension, non-specific background

amplification was completely disappeared. Therefore, we adopted the DPO primer in linear

extension step for all library samples in this study. Then, the single-stranded extension products

were C-tailed, and used as templates for the exponential PCR step using nested primer specific to

Tn5 and poly G primer that contain Illumina adapter sequences along with sample index sequences

(Fig. S2). The final PCR products were separated on an agarose gel, and the fragments within the

range of 325-625 bp were gel-purified. After pooling of multiple samples, the combined library

was sequenced on a HiSeq 3000.

Summary of Tn-seq DNA analysis

After de-multiplexing and C-tail trimming of all sequence reads, ~72 million reads of Tn5-

junctions with mean read length of 94 bp were obtained. The number of the reads mapped to the

complete genome of *S.* Typhimurium 14028 were ~25, 15, and 19 million for LB, H_2O_2L, and

H_2O_2H, respectively. The number of unique insertions on the chromosome were 125,449 in the

input library, excluding the plasmid (Table S1). On average, Tn5 was inserted in every 39 bp.

Number of raw reads per open reading frame (ORF) for H_2O_2L was plotted over the corresponding

number of H_2O_2H, which yielded an R^2 of 0.91, indicating the mutants in the input library

quantitatively responded in a similar way for both H_2O_2L and H_2O_2H as expected (Fig. S3). The

insertions were mapped to 5,428 genes or 8,022 genes/intergenic regions. Interestingly, the ORF

STM14_5121, which is 16.7 kbp long, had the highest number of insertions (~700 insertions) and

reads (0.25 M).

Comparison of various bioinformatics pipelines for Tn-seq data analysis

We used 3 different Tn-seq analysis tools to identify the genes and compare the results across the

methods with the goal of comprehensive identification of "all" genes required for resistance to

H_2O_2. The first tool, ARTIST (28), created small non-overlapping genomic windows of 100 bp

and the reads from each window were arbitrarily assigned into the middle of the window. The

default normalization script of the tool was used. Then, the relative proportions of insertion sites

in the output library versus the input were tabulated. Mann-Whiney *U* (MWU) test was used to

assess the essentiality of the locus. To consider a gene/intergenic region conditionally essential for

growth in the presence of H_2O_2, *p* value had to be ≤ 0.05 in 90 of the 100 conducted MWU tests.

Subsequently, 20 genes and 1 intergenic region were identified for H_2O_2L and 4 genes for H_2O_2H

(Data Set S1). We speculate the reason that more genes were identified for H_2O_2L in comparison

to H_2O_2H, was partially due to the lower number of total reads of H_2O_2L as compared to H_2O_2H,

even though the read numbers of H_2O_2L was normalized to those of the input.

The second tool, Tn-seq Explorer (29), counted insertions in overlapping windows of a fixed size.

Using a 550 bp window size, each annotated gene was assigned an essentiality index (EI) which

is determined mainly based on the insertion count in a window in this gene. The bimodal

distribution of insertion counts per window divided the essential genes to the left and the non-

essential genes to the right. To find conditional essential genes, the EI of the output was subtracted

from the EI of the input. The genes with negative ΔEI were ranked based on the change in read

fold change (Log2 (H_2O_2L or H_2O_2H/Input)). We found 114 consensus genes between H_2O_2L and

H_2O_2H that had at least four-fold reduction in H_2O_2H read counts as compared to the input. The

four-fold reduction (Log2FC = -2) threshold was chosen based on our validation study of Tn-seq

data by single mutant assays (Data Set S1).

The third tool, TRANSIT (30), determined read counts of genes in the input and output library.

The differences of total read counts between the input and outputs were obtained. The insertion

sites were permutated for a number that is specified by the user (we used 10,000 sample). This

sampling for each gene gave difference in read counts. The p value was calculated from the null

distribution of the difference in read counts. We identified 8 and 21 genes for the H_2O_2L and

H_2O_2H, respectively, using a p value ≤ 0.05 (Data Set S1). For normalization, trimmed total reads

(TTR) was chosen for this study

The combined list of the genes identified by the 3 Tn-seq analysis tools for both H_2O_2L and H_2O_2H

included 137 genes (Data Set S1). All of the genes on this list are expected to have a role in

conferring resistance to H_2O_2 and allow *Salmonella* to survive and replicate in the presence of

H_2O_2 *in vitro*. Of the 21 genes identified by TRANSIT, 19 of these genes were also identified by

Tn-seq Explorer, but only 3 out of this 21 were identified by ARTIST. The 19 genes were *hscA*,

rbsR, *fepD*, *efp*, *oxyR*, *polA*, *ybaD*, *aroD*, *ruvA*, *xthA*, *dps*, *aroB*, *uvrD*, *tonB*, *uvrA*, *aroK*, *ybbM*,

lon, and *proC*. Two genes, *fepD* and *xthA*, were identified by the all 3 methods and for both

conditions.

The 3 Tn-seq analysis tools are very valuable for Tn5 data analysis, but each tool has its own

advantages and disadvantages. For ARTIST, (i) the user must know how to run scripts in Matlab

software, (ii) the analysis is very slow on a personal computer with the HiSeq data, (iii) it has only

one method for normalization, but (iv) it can search for essentiality in the intergenic regions. For

Tn-seq Explorer, (i) there is no data normalization, and (ii) prediction on small genes is prone to

be inaccurate, but (iii) its very user-friendly and runs fast. For TRANSIT, (i) the user should have

some knowledge on running scripts on terminal, (ii) it may need some modification in its Python

script according to the way the library was prepared for sequencing, and (iii) a few software

packages should be installed on the computer as TRANSIT pre-requisites, but (iv) it does have 6

different methods for data normalization and it runs very fast on a personal computer. Although

ARTIST and Tn-seq Explorer are very useful tools for Tn-seq data analysis, we prefer using

TRANSIT in our future data analysis for conditionally essential genes. In the following sections,

we continued the downstream analysis mainly based on the 137 genes that include all of the genes

identified by all 3 methods.

The enriched pathways for resistance to H$_2$O$_2$

In order to categorize the identified genes that are required for *Salmonella* resistance to the H$_2$O$_2$,

the 137 genes were subjected to pathway enrichment analysis using DAVID Bioinformatics

Resources 6.7, NIAID/NIH (31). A total of 15 KEGG pathways (32) were recognized for 69 genes

on the list. The enriched pathways include homologous recombination (*ruvC*, *polA*, *ruvA*, *ruvB*,

priB, *recA*, *recR*, *holC*, *holD*, *recC*, *recG*), nucleotide excision repair (*uvrD*, *polA*, *uvrA*, *uvrC*),

mismatch repair (*dam*, *uvrD*, *holC*, *holD*), RNA degradation (*pnp*, *hfq*, *ygdP*), purine and

pyrimidine metabolism (*apaH*, *polA*, *pnp*, *arcC*, *spoT*, *holC*, *holD*, *cmk*, *dcd*, *pnp*), phenylalanine,

tyrosine and tryptophan biosynthesis (*aroD*, *aroB*, aroA, *aroK*, *aroE_2*), arginine and proline

metabolism (*proC*, *arcC*, *arcA*), starch and sucrose metabolism (*otsA*, *crr*, *pgm*), glycolysis and

gluconeogenesis (*crr*, *aceE*, *pgm*, *tpiA*), oxidative phosphorylation (*atpG*, *atpA*, *cydA*), DNA

replication (*polA*, *holC*, *holD*), flagellar assembly (*fliJ*, *fliD*, *flhD*, *fliC*), ABC transporter (*sapC*,

yejE, *yjgP*, *fepC*, *fepD*, *cbiM*, *cysP*), and two component system (*rcsB*, *rcsA*, *fliC*, *arcA*, *arcB*,

phoR, *barA*, *flhD*). Since KEGG was not able to recognize many genes on the list, we used

SP_PIR_Keywords of functional categories, which recognized majority of the genes and

categorized them into 55 functional categories (Table S2), excluding 15 uncharacterized genes

(ORFs). Among these categories were stress response (*rpoE*, *lon*, *dnaJ*, *hfq*, *yaiB*), iron (*dps*, *entD*,

iscA, *yjeB*, *yhgI*), and transcription regulation (*rcsA*, *oxyR*, *rpoE*, *yjeB*, *arcA*, *argR*, *rbsR*, *rpoS*,

fadR, *rcsB*, *furR*, *flhD*).

Validation of Tn-seq results using individual mutants

For the selected 50 genes among the 137 genes identified by Tn-seq, the growth phenotype was

determined using individual single deletion mutants in LB, H_2O_2L, and H_2O_2H. The genes were

considered to play a role in resistance to H_2O_2, if (i) lag phase time increased, (ii) growth rate

reduced or (iii) maximum OD_{600} decreased in the presence of H_2O_2 in comparison to the wild type

strain grown in the same conditions. Of the 50 single deletion mutants, 42 mutants were shown to

have a role in resistance to H_2O_2 (Fig. 2 and Data Set S2). One gene, *yhaD*, was identified by all

3 analysis tools, but it did not show the expected phenotype. The *fliD* was also identified by

ARTIST, but did not show any phenotype distinguishable from the wild type. The remaining 6

genes that did not show the phenotype was identified by Tn-seq Explorer. Based on the results of

the individual mutant assay, we conclude that 84% (42/50) of the genes identified by the Tn-seq

analysis and tested using single deletion mutants have a role for resistance to H_2O_2. These results

indicate that our Tn-seq analysis identified the genes in *S.* Typhimurium that are required for the

wild type level resistance to H_2O_2 with high accuracy.

Proteomics of H_2O_2 response

With ESI-LC-MS/MS in data-dependent acquisition (DDA) mode, the protein regulation was

determined using MS1 filtering technique that skyline software offers (33). It uses signal

intensities of tryptic peptides derived from the proteins of the wild type strain grown in the

presence of H_2O_2 in comparison to the control (LB). As described in Materials and Methods

section, trypsin digestion of the protein extracts under different conditions generates tryptic

peptides that are uniquely related to individual proteins. Tryptic peptides separated by liquid

chromatography from the complex samples were first subjected to simple mass measurement

(MS1) followed by intensity dependent fragmentation of these peptide ions to produce sequence

specific fragment ions by collision-induced dissociation (MS/MS). Tryptic peptides were then

identified using these sequence specific fragment ions via MASCOT database search software

(34), where the sequence specific fragment ions were matched to the proteins in *S.* Typhimurium

14028S reference proteome database (24, 35). This method of protein analysis is normally referred

to as data dependent analysis (DDA) method. At the beginning of data analysis, the H_2O_2L and

H_2O_2H data were compared to LB separately, however it turned out that comparison was not

sensitive enough to differentiate between H_2O_2L and H_2O_2H conditions. Hence, the data of H_2O_2L

and H_2O_2H were combined in the analysis in comparison to LB. We identified 1,104 proteins of

Salmonella for the 3 conditions (Data Set S3); of these, 246 proteins were differentially expressed

in response to H_2O_2 with *p* values ≤ 0.05 and 90% CI. The upregulated proteins were 121 and the

downregulated proteins were 125 in response to H_2O_2. Since Tn-seq revealed genetic requirements

for fitness under the selection conditions, the identified genes are expected to express their proteins

under the conditions to perform their cellular functions. Often the proteins required for fitness

under a given condition are overexpressed under the condition, but it may not be the case for some

proteins. In this study, we had a unique opportunity to comparatively analyze both Tn-seq and the

MS data to understand the relationship between genetic requirements and changes in expression

level under the condition of interest, which was H_2O_2 in this study. We also obtained the list of

essential genes based on our Tn-seq data, which could not tolerate insertions by definition, and if

we were not certain about essentiality of a gene from our Tn-seq data, the gene was searched for

essentiality in the previously reported list of *Salmonella* essential genes (22). The comprehensive

list of essential genes allowed us to study any correlation between the essentiality and the changes

in protein expression. Among the 246 proteins, there were 78 essential and 168 non-essential

proteins. Among the 78 essential proteins, 25 were upregulated whereas 53 were downregulated.

On the contrary, the majority (n = 96) of the detected non-essential proteins were upregulated,

while 72 non-essential proteins were downregulated. To further examine the quantitative

relationships closely, 64 genes/proteins identified by both methods (Data Set S3) were focused on.

Among the 64 genes/proteins, 57 genes showed negative Log2FC based on Tn-seq data, and 41

proteins among the 57 were upregulated at protein level. However, only 12 proteins had p values

of ≤ 0.05 (AhpC, ArcA, Crr, DksA, FliC, IcdA, OxyR, Pgm, RecA, RpoS, SlpA, and WecE).

Using KEGG pathway analysis, 150 proteins among the 246 were enriched in 21 pathways (Table

S3). Interestingly, of the all 59 30S and 50S ribosomal proteins in *S.* Typhimurium, 37 of these

proteins (63%) were downregulated in response to H_2O_2. Moreover, of the 8 identified proteins in

TCA cycle, 6 proteins were downregulated, including 2 essential proteins.

Although DDA method can be used to search for all proteins in a complex sample, it is prone to

miss the identification of important proteins due to the fact that fragmentation of tryptic peptides

from these proteins may not be triggered as a result of lower peptide ion intensities compared to

the threshold set. To quantify the proteins of the genes identified by Tn-seq more precisely and

accurately, we used targeted-proteomic approach by employing liquid chromatography coupled

with triple quadrupole mass spectrometry (LC-QQQ-ESI-MS). Here, tryptic peptides of the protein

were targeted for fragmentation (MS/MS) independent of their intensities, as described in

Materials and Methods, and the observed sequence specific fragment ion intensities from three

unique tryptic peptides were utilized for protein quantitation. Of the 137 Tn-seq identified genes,

we selected 33 genes to quantify their proteins in response to H_2O_2 by using targeted proteomics

(Dataset S3). Interestingly, 23 (70%) of the 33 tested proteins were upregulated in response to

H_2O_2. This shows a good agreement between the results of the Tn-seq and the targeted proteomics.

Aromatic amino acid biosynthesis and H_2O_2

Interestingly, our Tn-seq data revealed that the aromatic amino acid biosynthesis and metabolism

pathway play a role in conferring resistance in *Salmonella* to H_2O_2 (Fig. 3A and 3B). Five genes,

aroB, *aroD*, *aroE_2*, *aroK*, and *aroA* in the aromatic amino acid biosynthesis pathway were

identified by Tn-seq, and the fitness of the mutants were significantly reduced in the presence of

H_2O_2. To confirm this, 4 of these genes were evaluated using individual mutant assays. The

Salmonella aroK mutant showed the strongest phenotype, because it failed to grow in the presence

of H_2O_2L or H_2O_2H during 24 h incubation time. Also, the *aroB* mutant exhibited a strong

phenotype, significantly extending lag phase for both H_2O_2 conditions. The *aroE_2* mutant also

exhibited an extended lag time, but the *aroA* mutant did not show any difference in growth

phenotype in the presence of H_2O_2. In addition, targeted-proteomics also showed that all these 5

proteins were upregulated in response to H_2O_2. The most upregulated protein was *aroK* and this

was followed by the *aroE_2*, *aroA*, *aroB*, and *aroD* (Fig. 3C and Data Set S3).

The ROS damages a variety of biomolecules via Fenton reaction, which consequently lead to

metabolic defects, specifically auxotrophy for some aromatic amino acids (10). The *E. coli* mutants

that lack superoxide dismutase enzymes are unable to grow *in vitro* unless the medium are

supplemented with aromatic (Phe, Trp, Tyr), branched-chain (Ile, Leu, Val), and sulfur-containing

(Cys, Met) amino acids (36). We identified the genes in the aromatic amino acid biosynthesis

pathway that are critically important for resistance to H_2O_2. In this pathway, *aroK* catalyzes the

production of shikimate 3-phosphate from shikimate, which consequently leads to the production

of tryptophan, phenylalanine, tyrosine and some metabolites from the chorismate precursor in *E.*

coli. Further, the *aroK* mutant in *E*. coli displays increased susceptibility to protamine, a model

cationic antimicrobial peptide. It has been suggested that resistance to protamine is probably due

to the aromatic metabolites and product of *aroK* gene, which act as a signal molecule to simulate

the CpxR/CpxA system and Mar regulators (37). In our Tn-seq data, *cpxR*/*cpxA* and *marBCRT*

were in the list of non-required genes, but the proteomics data indicated that CpxR was

upregulated. Also, the *aroK* mutant in *E. coli* is resistance to mecillinam, a beta-lactam antibiotic

specific to penicillin-binding protein 2. It has been concluded that the AroK has a secondary

activity in addition to the aromatic amino acid biosynthesis, probably related to cell division (38).

In addition, the *aroK* gene presents a promising target to develop a non-toxic drug in

Mycobacterium tuberculosis because *aroK* is the only *in vitro* essential gene among the aromatic

amino acid pathway genes and blocking *aroK* kills the bacterium *in vivo* (39). Moreover, the *aroK*

gene plays a general role in *S.* Typhimurium persistence in pigs (40). The *aroB* is another gene in

the pathway that was identified by Tn-seq, which encodes 3-dehydroquinate synthase in the

Shikimate pathway, aromatic amino acid biosynthesis pathway. The *Salmonella* lacking *aroB*

showed a strong growth defect in the presence of H_2O_2. When this mutant was grown in the

presence of H_2O_2, it increased the lag phase time by a 114-fold for the H_2O_2L and a 347-fold for

the H_2O_2H as compared to the mutant grown in absence of H_2O_2. *S.* Typhimurium mutant lacking

the *aroB* gene is attenuated in BALB/c mice (41). In addition to *aroK* and *aroB*, *aroE_2* was also

shown to be important for resistance to H_2O_2, because deletion of the *aroE_2* reduced the growth

rate by 35% in the presence of H_2O_2 and increased the lag phase time, too. All these 3 genes in this

pathway are required for systemic infection of *Salmonella* in BALB/c mice in a more recent study

(18). We observed that there was a strong correlation between the fitness based on Tn-seq data,

growth rates measured by individual mutant assays, and upregulation of their proteins quantified

via targeted proteomics. This demonstrates the power of proteogenomic approach in discovering

and characterizing the genes that are required for growth under a specific condition.

The *ybbM*, *ybbK*, and *ybbL* have a role in H_2O_2 resistance

The mutants with single deletion in each of *ybbK*, *ybbL*, and *ybbM* genes on the same pathway

showed a strong phenotype against the activity of H_2O_2 in a dose-dependent manner. Based on Tn-

seq data, the fitness of *ybbM* was -1.16 and -1.79 for H_2O_2L and H_2O_2H, respectively (Fig.4A).

The *ybbM* mutant demonstrated decreased growth rate by 38% for H_2O_2L and 100% for the H_2O_2H

as compared to the mutant grown in the absence of H_2O_2. This mutant also increased the lag time

by a 126-fold and a 267-fold for H_2O_2L and H_2O_2H, respectively (Fig.4B). Also, the fitness score

of *ybbK* was -0.92 for H_2O_2L and -1.81 for H_2O_2H. The *ybbK* mutant showed decrease of growth

rate for both conditions H_2O_2L and H_2O_2H by 85% and 95%, respectively. The deletion increased

the lag phase by a 46-fold and a 114-fold in the presence of H_2O_2L and H_2O_2H, respectively

(Fig.4B). Moreover, the fitness of *ybbL* mutant was -1.05 and -1.73 for H_2O_2L and H_2O_2H,

respectively. Deleting the *ybbL* in *Salmonella* led to decrease in growth rate by 27% for H_2O_2L

and 92% for H_2O_2H. The lag phase time for this mutant increased by a 22-fold and a 33-fold for

H_2O_2L and H_2O_2H, respectively. In addition, YbbM, YbbL, and YbbK proteins were upregulated

in response to H_2O_2; YbbM was the most upregulated protein among the 3 proteins, followed by

YbbL, and YbbK (Fig. 4C and Data Set 3). The fitness scores of the Tn-seq of these 3 genes are

correlated strongly with the growth rate, lag time of their respective mutants, and upregulation of

their proteins. As the number of reads depletes after the selection for a mutant, (i) there was more

reduction in growth rate, (ii) the mutant stays longer in the lag phase, and (iii) the protein

expression elevates. These observations clearly point to their role in conferring resistance to the

H_2O_2-mediated stress. These genes were described in the *Salmonella* reference genome as follows:

ybbM, putative YbbM family transport protein, metal resistance protein; *ybbK*, putative inner

membrane proteins; *ybbL*, putative ABC transporter, ATP-binding protein YbbL. To the best of

our knowledge, there is only one published study on the *ybbM* and *ybbL* (42). Based on their

findings, YbbL and YbbM have a role in iron homeostasis in *E. coli* and are important for survival

when the bacterium was challenged with 10 mM H_2O_2 for 30 min; this putative ABC transporter

transports iron and lessens ROS species formation that generates via H_2O_2. In this study, we

identified an additional gene, *ybbK*, in the same pathway as the gene required for resistance to

H_2O_2, strongly establishing the role of these 3 genes in resistance to H_2O_2.

The H_2O_2 scavenging and degrading genes

Salmonella employs redundant enzymes to degrade or scavenge ROS. The *katE*, *katG*, and *katN*

genes encode catalases, which are involved in H_2O_2 degradation. The *ahpCF*, *tsaA*, and *tpx* genes

encode peroxidases, which scavenge H_2O_2. The *sodA*, *sodB*, *sodCI*, and *sodCII* genes encode

superoxide dismutases and these enzymes specifically scavenge O_2 (11, 12, 43 - 45). However,

none of these were present in the list of genes identified by Tn-Seq. Even though *katE*, *katG*, *ahpC*,

sodA, *sodCI*, and *sodCII* showed reduced fitness, they did not meet the statistical threshold.

However, the proteomics data indicated that AhpC, SodB, and TpX were upregulated in the

presence of H_2O_2 (Data Set S3) and KatG was also upregulated, but its *p* value was 0.054. This

reveals that these 4 proteins were the most important enzymes to scavenge H_2O_2 under our

experimental conditions. *Salmonella* containing an *ahpC* promoter-gfp fusion shows that

expression of the *ahpC* is regulated by ROS that is generated from macrophages or exogenous

H_2O_2 and the response to H_2O_2 is in a dose-dependent manner (46). *Salmonella* mutant that lacks

katE, *katG*, or *ahpCF* can degrade micromolar concentrations of H_2O_2. However, *Salmonella*

mutant that has deletions in the all 5 genes, *katE*, *katG*, *katN*, *ahpCF* and *tsaA* (HpxF), cannot

degrade H_2O_2, is unable to proliferate in macrophages, and showed reduced virulence in mice (11).

This data emphasizes that *ahpC*, *sodB*, and *tpx* may be the primary players in scavenging and

degrading H_2O_2 in our experiment. Why Tn-seq did not detect any of these genes, while proteomics

detected only these 3 proteins among others? It may reflect the functional redundancy in the

genetic network that prevented single deletions in one of the these genes from exhibiting fitness

defect. Alternatively, when these mutants were grown together with all other mutants in the library,

the functional protein lacking in one mutant due to Tn5 insertion could have been compensated by

the other mutants in the library.

In addition to these genes, *oxyR* was detected by Tn-seq (Fig. 1C) and DDA proteomics. The *oxyR*

was identified by all 3 analysis methods of Tn-seq data and it was on the top of the list, indicating

a severe fitness defect of the mutant. The *oxyR* gene encodes H_2O_2 sensor and transcription factor,

which mediates protection against ROS. The *katG* and *ahpCF* are regulated by OxyR, peroxide

response regulator (13, 14). Although *Salmonella* OxyR regulon is induced in the *Salmonella*-

containing vacuole in macrophage, the *oxyR* mutant was virulent in a BALB/c mouse and can grow

well in human neutrophils *in vitro* (47, 48). The fitness of *oxyR* mutant was reduced for both H_2O_2L

and H_2O_2H with the respective fitness score of -4.96 and -5.94. *Salmonella oxyR* mutant exhibited

the growth rate reduced by 24% and 40% for H_2O_2L and H_2O_2H, respectively. Comparing this

reduction in growth rate to the other mutants such as *rpoS* or *aroK*, we observed that the *oxyR*

mutant did not show severe phenotype and the mutant escaped from the lag phase easily.

Moreover, our targeted proteomics indicated that the OxyR was not upregulated significantly.

Further studies are needed to uncover the exact role of OxyR in response to ROS. However,

previous studies implied that OxyR plays an essential role in resistance to H_2O_2 by regulating other

proteins. OxyR induces Dps in *E. coli*, a ferritin-like protein that sequesters iron (49). Sequestering

of iron impairs the Fenton reaction, which consequently provides protection against ROS and

reduces the damage of biomolecules. The *dps* gene was identified by the Tn-seq and its fitness

score was -2.48. However, the Dps protein was downregulated based on the DDA proteomic

analysis. To confirm this unexpected finding, we conducted the proteomic assay twice and each

time with at least 4 technical replicates, but the Dps protein was significantly downregulated with

$p = 0.001$. Further, the targeted-proteomics gave the same result, pointing to the downregulation

of Dps in response to H_2O_2. This is contrary to the previously reported works on Dps in *Salmonella*

and the reason for the discrepancy is unclear.

DNA repair system and H_2O_2

The imposed exogenous H_2O_2 activates DNA repair system in *Salmonella* in order to repair or

eliminate the damage that occurred on the nucleotides. The *E. coli* RecA protein repairs double-

strand DNA lesions through recombination (50). In our Tn-seq analysis, the fitness score of this

mutant was -5.36 for both concentrations, and in proteomics, the RecA was upregulated.

Salmonella recA mutant decreased the maximum OD_{600} by 16% for H_2O_2L and 22% for H_2O_2H

as compared to the same mutant grown in LB. The *Salmonella recA* mutant was also sensitive to

exogenous H_2O_2 in aerated rich medium (26). Moreover, *recG*, recombination and DNA repair

gene (51), showed a stronger phenotype than *recA* mutant. The *regG* deletion in *Salmonella* caused

the growth rate reduction by 52% for H_2O_2L and 60% for H_2O_2H. This disruption in *recG* also

caused the cells to stay in lag phase for a longer time in the presence of H_2O_2 as compared to LB;

the lag time increased by a 62-fold and a 159-fold for H_2O_2L and H_2O_2H, respectively. In the blood

of patients with *Salmonella* Typhi bacteremia, the proteins encoded by *recA*, *recG*, and *xthA* genes

were detected, suggesting these proteins are actively expressed in the blood environment (52). The

XthA protein is another enzyme that participates in DNA repair that is induced by H_2O_2 and iron-

mediated Fenton reaction. The *xthA* encodes exonuclease III, which repairs the damaged DNA.

We found that the *xthA* gene was required based on the Tn-seq assay and its mutant had a reduced

fitness score of -3.06 for H_2O_2L and a -4.38 for the H_2O_2H. Further, *Salmonella* lacking the *xthA*

increased the lag time by 8-fold and a 12-fold for H_2O_2L and H_2O_2H, respectively. Targeted-

proteomics showed upregulation of XthA in response to H_2O_2. *Salmonella* enterica serovar

Enteritidis defective in *xthA* is susceptible to egg albumin (53). *E. coli xthA* mutant is

hypersensitive to H_2O_2 (54). The *xthA* is also required for *Mycobacterium tuberculosis* to infect

C57BL/6J mice (55). In addition to these genes that repair damaged DNA, *uvrA* encoding Holliday

junction DNA helicase motor protein, *uvrA* encoding exonuclease ABC subunit A, *uvrD* encoding

DNA-dependent helicase II, and *polA* encoding DNA polymerase I were among top of the list of

the genes identified by Tn-seq as required for resistance to H_2O_2. Collectively, DNA repair system

is crucial for the survival of the *Salmonella* in a niche that contains H_2O_2.

Flageller genes and H_2O_2

Some flageller genes, *fliC* and *fliB*, were shown to be important for resistance to H_2O_2. These two

genes were identified by Tn-seq and their proteins were shown to be upregulated based on

proteomics. *Salmonella* lacking either of these genes exhibited a strong phenotype in the presence

of H_2O_2. During 24 h of incubation, *fliC* and *fliB* mutants could not grow in both H_2O_2 conditions.

However, growth of *fliD* mutant was not affected by H_2O_2. The *fliC* was shown to have a role in

Salmonella Typhi interaction with human macrophages (56) and *Salmonella* Typhimurium *fliB*

mutant was defective in swarming motility (57). Currently it remains unclear how flagella genes

can be involved in the resistance of *Salmonella* to oxidative stress, which warrants future study

into this direction.

Fe-S cluster biogenesis system and H_2O_2

Salmonella requires the genes from Fe-S cluster biogenesis system in order to resist H_2O_2. Our

Tn-seq analysis identified 5 genes in this system as required for the resistance. In *isc* operon (Fe-

S cluster), *iscA*, *hscB*, and *hscA* were among the genes required to resist H_2O_2. Particularly, the

hscA is on the top of the gene list identified by Tn-seq. In *E. coli*, this operon is regulated by *iscR*,

iron sulfur cluster regulator (58); in *Salmonella* the gene *iscR* encoding this transcription regulator

is named *yfhP*. The HscB and HscA chaperones are believed to be involved in the maturation of

[2Fe–2S] proteins (59, 60). The second operon that is involved in Fe-S protein biogenesis is the

suf, sulfur mobilization operon. Tn-seq found that two genes in this operon were required for

Salmonella to resist H_2O_2; *sufS* and *sufC*. *Salmonella* mutant lacking *sufS* exhibited a strong

phenotype in the presence of H_2O_2 and could not grow during the 24 h of incubation as compared

to LB. The SufS with SufE in *E. coli* form a heterodimeric cysteine desulphurase and SufB, SufC,

and SufD form a pseudo-ABC-transporter that could act as a scaffold (60); this operon is regulated

by OxyR (14). The other known genes in these two operons that are present in *Salmonella* are *iscA*,

sufA, *sufB*, and *sufD*; they showed a reduced fitness, while their *p* values were greater than 0.05.

The damage of Fe-S clusters is not only problem for the defective proteins, but also it fuels the

Fenton reaction via the released iron and H_2O_2 (10). Thus, *Salmonella* uses Fe-S cluster repair

system as an arsenal to overcome the damage imposed by H_2O_2.

DNA adenine methylase

DNA adenine methylase genes, *dam* and *damX*, are involved in *Salmonella* resistance against

H_2O_2. Our Tn-seq data showed that fitness of *dam* and *damX* mutants were reduced in the presence

of H_2O_2. To confirm this, *Salmonella dam* mutant was grown in both conditions. Under H_2O_2L,

the growth rate was reduced by 42% as compared to the mutant in LB and the mutant could not

grow under H_2O_2H during the 24 h of incubation. In addition, the lag time of the *Salmonella dam*

mutant was extended by a 19-fold for H_2O_2L. While the *Salmonella damX* mutant displayed a

moderate phenotype as compared to the *dam* mutant, the *damX* mutant also showed the growth

rate decreased by 23% and 33% for H_2O_2L and H_2O_2H, respectively. The lag time was extended

for this mutant by a 25-fold and a 49-fold for H_2O_2L and H_2O_2H, respectively. The *dam* regulates

virulence gene expression in *S.* Typhimurium (61). The different levels of Dam affects virulence

gene expression, motility, flagellar synthesis, and bile resistance in the pathogenic *S.* Typhimurium

14028S (62). Dam methylation activates the gene that are involved in lipopolysaccharide synthesis

(63). Moreover, *Salmonella* defective in *damX* is very sensitive to bile (64). Collectively, our study

demonstrates the critical role of DNA adenine methylase in *Salmonella* resistance against H_2O_2.

Other genes for H_2O_2 resistance

Beside the important pathways described above, there were many additional genes also important

for resistance to H_2O_2. Among those, the 3 unrelated genes, *rpoS*, *pgm*, and *tonB*, are important

ones that deserve more attention. The *rpoS* mutant showed reduced fitness and its protein was

upregulated in the presence of H_2O_2. *Salmonella* mutant defective in *rpoS* showed a strong

phenotype when grown in the presence of H_2O_2. The *rpoS* encodes the alternative sigma factor σ^S,

subunit of RNA polymerase; it is the master regulator of stress response (65). This implies that

rpoS is an important component of the genetic regulatory network that *Salmonella* employs in

order to resist H_2O_2. Furthermore, the fitness of *pgm* mutant was reduced and its protein was

upregulated in the presence of H_2O_2. Knock out of *pgm* in *Salmonella* caused a decrease in growth

rate, increase the lag phase time, and reduce the maximum OD_{600} in the presence of H_2O_2. The

pgm encodes phosphoglucomutase required for catalysis of the interconversion of glucose 1-

phosphate and glucose 6-phosphate (66). This gene contributes to resistance against antimicrobial

peptides, is required for *in vivo* fitness in the mouse model, and participates in LPS biosynthesis

(67). Lastly, the fitness of *tonB* mutant was also reduced. *Salmonella* lacking *tonB* exhibited a

strong phenotype in the presence of H_2O_2 as compared to the mutant grown in LB. The gene

mediates iron uptake in the *Salmonella* (45). In addition, seven of the genes identified in our study

(*proC*, *arcA*, *barA*, *exbD*, *flhD*, *fliC*, and *fliD*) were previously shown to be important for

interaction of *Salmonella* Typhi with human macrophages (56). To see the full list of *S.*

Typhimurium genes required for H_2O_2 resistance, the supplementary information contains data of

Tn-seq, individual mutant assays, and proteomic analysis in Data Set S1, S2, and S3, respectively.

Conclusions

We applied Tn-seq and proteomic analysis to find the genes and proteins that are required in *S.*

Typhimurium to resist H_2O_2 *in vitro*. As the concentration of H_2O_2 increased, the growth rate

reduced, the lag time extended, the fitness of mutants decreased, and some proteins were

differentially expressed. Validation of Tn-seq results with individual mutant assays indicated the

accuracy of the identified genes in response to the two H_2O_2 concentrations. The targeted-

proteomics had a good agreement with Tn-seq. We found about 80 genes that have not been

associated to resistance to H_2O_2 previously. *Salmonella* employs multiple pathways to resist H_2O_2

and the most important ones are ROS detoxifying enzymes, amino acid biosynthesis (*aroK* and

aroB), putative iron transporters (*ybbK*, *ybbL*, *ybbM*), iron homeostasis, Fe-S cluster repair, DNA

repair, flagellar and DNA adenine methylase genes. The genes identified in this study will broaden

our understanding on the mechanisms used by *Salmonella* to survive and persist against ROS in

macrophages.

Our unbiased system-wide approach, Tn-seq, was successful in identifying novel genetic

determinants that have not been implicated previously in *Salmonella* resistance to oxidative stress.

Furthermore, the combined use of quantitative proteomic approach has provided additional

insights on the function or mode of action of the identified genetic determinants in resisting

oxidative stress. As expected, the majority of the proteins important for resistance to H_2O_2 were

upregulated in response to the same stressor. However, the expression level did not increase for

some proteins, in spite of their known roles in resistance to H_2O_2. Interestingly, the downregulation

of Dps and other proteins was counterintuitive to the common mode of protein regulation and

function, yet it may point to some unknown aspects of how *Salmonella* regulates the expression

of those proteins to better cope with the oxidative stress during infection in macrophage.

Materials and Methods

Construction of Tn5 mutant library

We mutagenized *Salmonella* enterica subsp. enterica serovar Typhimurium str. ATCC 14028S

(with spontaneous mutation conferring resistance to nalidixic acid (NA)), by biparental mating

using *Escherichia coli* SM10 λ*pir* carrying a transposon-delivery plasmid vector pBAM1

(Ampicillin (Amp) resistance) as a donor strain (68). The plasmid pBAM1 was generously

provided by Victor de Lorenzo (Systems and Synthetic Biology Program, Centro Nacional de

Biotecnología, Madrid, Spain). The donor strain*, E. coli* Sm10 λpir (pBAM1), was grown

overnight in LB with 50 µg/ml Amp and recipient strain was grown in LB with 50 µg/ml NA at

37 °C . Equal volumes (1 ml) of donor and recipient were mixed, centrifuged, washed in 10 mM

MgSO$_4$, and re-suspended in 2 ml PBS (pH 7.4). Then, the mating mixture was concentrated and

laid on a 0.45-µm nitrocellulose filters (Millipore). The filter was incubated at 37°C on the surface

of LB agar plate. After 5 h of conjugation, the cells on the filter was washed in 10 mM MgSO$_4$,

and resuspended in 1 ml MgSO4. The conjugation mixture was plated on LB agar containing 50

µg/ml NA and 50 µg/ml kanamycin (Km). After approximately 24 h at 37°C, we scraped the

colonies into LB broth containing 50 µg/ml Km and 7% DMSO. The yield was approximately

68,000 individual colonies from each conjugation. Five independent conjugations were conducted

to yield approximately 325,000 mutants. The library was stored at −80°C in aliquots. To determine

the frequency of the mutants that have been produced by integration of the entire delivery plasmid,

the colonies were picked randomly and streaked on LB plates (Km) and LB plates (Km and Amp).

It was shown that ~20% of the cells in the library were resistant to Amp, indicating a significant

portion of the Km-resistant colonies was not from authentic transposition events.

Measuring growth responses of *S.* Typhimurium to H_2O_2

To determine the effect of H_2O_2 concentrations on growth parameters, overnight culture of the

wild type *S.* Typhimurium 14028s was inoculated into fresh LB broth media (1:200 dilution) to

give a seeding concentration corresponding to OD_{600} of ~0.1. The LB broth contained freshly

prepared H_2O_2 to give the final concentrations ranging from 0.05 to 10 mM. The cultures were

directly added into 96-well microplate (200 µl/well). The microplate was incubated in a Tecan

Infinite 200 microplate reader at 37°C, with shaking duration 5 s, shaking amplitude 1.5 mm, and

reading OD_{600} every 10 min. The number of replicates were at least three. The lag time, growth

rate, and maximum OD_{600} were calculated using GrowthRates script (69). Growth Rate % decrease

was calculated as follows (70): Growth Rate % decrease $= ((\mu_{PC} - \mu_S)/\mu_{PC}) \times 100$; where $\mu =$ the

maximum slope (growth rate), $\mu_{PC} =$ growth rate of positive control (without H_2O_2), $\mu_S =$ growth

rate in the presence of H_2O_2.

Selection of the mutant library for Tn-seq analysis

The transposon library was thawed at room temperature and diluted 10^{-1} in fresh LB broth. To

activate the library, the diluted library was incubated at 37°C with shaking at 225 rpm for an hour.

Then, the culture was washed twice with PBS and resuspended in LB broth medium. The library

was inoculated to 20 ml LB broth and LB broth supplemented with either 2.5 or 3.5 mM H_2O_2

(H_2O_2L and H_2O_2H, respectively), seeding CFU was 3.5 x 10^6 per ml. Then, when the cultures

reached mid-exponential phase, OD_{600} of 2.7 (~1.17 x 10^8 CFU/ml), the incubation was stopped,

and the culture was immediately harvested by centrifugation, and stored at -20°C.

Preparation of Tn-seq amplicon libraries

Genomic DNA was extracted from the harvested cells using DNeasy Blood & Tissue kit (Qiagen),

and quantified using Qubit dsDNA RB Assay kit (Invitrogen). As described above, 20% of the

mutants in the library were the result of the integration of pBAM1 into chromosome. To remove

the Tn5-junction sequences originated from the plasmid in the Tn-seq amplicon libraries, genomic

DNA was digested with PvuII-HF (New England Biolabs), which digests immediately outside the

inverted repeats on both sides of Tn5 in pBAM1, and purified with DNA Clean & Concentrator-5

kit (Zymo Reaerch). Then, a linear PCR extension was performed using a Tn5-specific primer in

order to produce single stranded DNA corresponding to Tn5-junction sequences. To increase the

specificity in extending into Tn5-junction sequences, the linear PCR was conducted with a dual

priming oligonucleotide Tn5-DPO (5'-AAGCTTGCATGCCTGCAGGTIIIIICTAGAGGATC-3')

that is specific to Tn5 end (71). The PCR reaction contained 25 μl Go Taq Colorless Master Mix

(Promega), 20 μM Tn5-DPO primer, 100 ng gDNA, and 50 μl MQ-H_2O. The PCR cycle consisted

of the initial denaturation at 95°C for 2 min, followed by 50 cycles at 95°C for 30 sec, 62°C for 45

sec, and 72°C for 10 sec. The PCR product was purified with DNA Clean & Concentrator-5 kit

and eluted in 13 μl TE buffer. After that, C-tail was added to the 3' end of the single-stranded

DNA. The C-tailing reaction was consisted of 2 μl terminal transferase (TdT) buffer (New England

Biolabs), 2 μl $CoCl_2$, 2.4 μl 10 mM dCTP, 1 μl 1 mM ddCTP, 0.5 μl TdT and 13 μl purified linear

PCR product. The reaction was performed at 37°C for 1 h and the enzyme was inactivated by

incubation at 70°C for 10 min. The C-tailed product was purified with DNA Clean & Concentrator-

5 kit and eluted in 12 μl TE. Next, the exponential PCR was performed with forward primer, P5-

BRX-TN5-MEO,

AATGATACGGCGACCACCGAGATCT<u>ACACTCTTTCCCTACACGACGCTCTTCCGATC</u>

<u>T</u>NNNNAG-6 nt barcode-CCTAGGCGGCCTTAATTAAAGATGTGTATAAGAG and reverse

primer, P7-16G,

CAAGCAGAAGACGGCATACGAGCTCTTCCGATCTGGGGGGGGGGGGGGGGGG to attach

Illumina adapter sequences along with the sample barcodes. The PCR reaction contained 25 μl Go

Taq Green Master Mix, 10 μM P5-BRX-TN5-MEO primer, 10 μM P7-16G primer, 1 μl purified

C-tailed genomic junctions, and MQ-H$_2$O to 50 μl; the PCR condition started with initial

denaturation at 95°C for 2 min, followed by 36 cycles of 95°C for 30 sec, 60°C for 30 sec, and

72°C for 20 sec, with the final extension at 72°C for 5 min. Then, the size selection of the DNA

was performed using agarose gel electrophoresis. The 50 μl PCR products were incubated at 60°C

for 15 min and incubated on ice for 5 min, and immediately loaded on the 1% agarose gel in 0.5%

TAE buffer. After running the gel, the DNA fragment of size 325 – 625 bp was cut and put in a

microtube for each sample. The DNA was extracted from the gel using Zymoclean Gel DNA

Recovery kit (Zymo Reaerch). The prepared DNA libraries were quantified using Qubit dsDNA

RB Assay kit. Since each library has its own barcode, the libraries were combined and sequenced

on a flow cell of HiSeq 3000 using single end read and 151 cycles (Illumina) at the Center for

Genome Research & Biocomputing in Oregon State University.

Tn-seq data analysis

The preliminary data analysis was conducted by using a super computer in the High Performance

Computing Center (AHPCC) at the University of Arkansas. The libraries that were multiplexed

for sequencing were de-multiplexed using a custom Python script. The script searched for the six-

nucleotide barcode for each library for perfect matches. In order to extract the transposon genomic

junctions, we used Tn-Seq Pre-Processor (TPP) tool (30) with some modifications in the script.

The TPP searched for the 19 nucleotide inverted repeat (IR) sequence and identified five

nucleotides (GACAG) at the end of the IR sequence, allowing one nucleotide mismatch. The Tn5-

junctions that start immediately after GACAG were extracted and the C-tails at the end of junctions

were removed. Tn5-junction sequences less than 20 nucleotides were discarded and remaining

Tn5-junctions were mapped to the *Salmonella* enterica serovar Typhimurium 14028S genome and

plasmid using BWA-0.7.12 (72). To identify genes that are required for H_2O_2 resistance, the

following three Tn-seq analysis tools were used for comparative analysis: (i) ARTIST (28): the

genomic junctions were mapped to the reference genome using Bowtie 2.2.7 (73). The number of

insertions and reads were determined for genes and intergenic regions. The data were normalized

with default script in the ARTIST. Then, the relative abundance of Tn5 insertions in the output

library versus the input were calculated. Later, the *p* values were calculated from a 100 independent

Mann–Whitney U test (MWU) analysis that were carried out on input and output data for each

gene. Finally, the genes were considered conditionally essential if the *p* values were ≤ 0.05 in the

90 of the 100 MWU tests. (ii) Tn-seq Explorer (29): The output SAM files from the TTP were

used as input to the Tn-seq Explorer. The unique insertions with less than 20 reads were removed

from the input and outputs. Using the window size of 550 and excluding 5% of beginning of genes

and 20% of the end of genes, Essentiality Index (EI), number of unique insertions, and total number

of reads per gene were counted. The EI of more than 10 were removed from the input. Genes with

less than 300 nucleotides were removed. Deferential EI were calculated from input and outputs

(ΔIE = output EI–input EI) and genes with ΔIE more than -1 were removed. Log2 fold change of

reads were calculated from input and output (Log2FC = log2 (output reads/input reads)) and the

genes were ranked based on the Log2FC value from least value to highest. The genes with Log2FC

value of less than -2 from the H_2O_2H and present in H_2O_2L were considered conditionally essential.

(iii) TRANSIT (30): The output wig files from the TTP was used as input data file for TRANSIT.

The comparative analysis was conducted with Tn5 resampling option. The reads were normalized

with trimmed total reads (TTR). Insertions outside the 5% and 10% sequences from 5'- and 3'-

ends were removed, respectively. The genes were considered conditionally essential if p values $\leq$

0.05.

Phenotypic evaluation of individual deletion mutants

The mutants were obtained from *Salmonella enterica* subsp. *enterica*, 14028s (Serovar

Typhimurium) Single-Gene Deletion Mutant Library through BEI Resources

(www.beiresources.org). The overnight cultures of *S.* Typhimurium mutants were added into fresh

LB broth media containing freshly prepared H_2O_2 (2.5, or 3.5 mM/ml) (1:200 dilution) to give

seeding OD_{600} of 0.1. The cultures were directly added into 96-well microplates and incubated in

Tecan Infinite 200 at 37ºC for 24 h. The lag time, growth rate, and maximum OD_{600} were calculated

using GrowthRates (69).

Sample preparation for proteomics and mass spectrometry analysis

The overnight culture of the wild type *S.* Typhimurium 14028 was diluted 1:200 in 50 ml LB

medium, and LB containing either 2.5 or 3.5 mM H_2O_2 in a 300-ml flask. The cultures were grown

until mid-exponential phase (OD_{600} of 2.7), and the 50 ml volume of cultures were used for a total

protein extraction by using Qproteome Bacterial Protein Prep kit (Qiagen). The proteins were

denatured with mercaptoethanol and separated by SDS-PAGE gel electrophoresis. For each

condition, there were three lanes with approximately 300 μg of proteins. The gel portions of 3

lanes for each condition were cut out and chopped into small pieces, pooled together, washed twice

with 50 mM NH_4HCO_3, destained with NH_4HCO_3/ 50% Acetonitrile (ACN), and dried with pure

ACN. Then, the proteins were reduced using 10 mM Dithiothreitol in 50 mM NH_4CO_3 and the

alkylation was conducted with 10 mg/ml Iodoacetamide Acid in 50 mM NH_4CO_3. After that, the

proteins were washed with NH_4HCO_3, and dried with pure ACN. Mass spectrometry grade Trypsin

gold from Promega (~20 ng/μl in 50 mM NH_4HCO_3) was added to dried gels, and left it overnight

for efficient in-gel digestion of the proteins at 37ºC. During the digestion, tryptic peptides diffused

out into the solution. Later, these digests were analyzed by ESI-LC-MS/MS at State Wide Mass

Spectrometry Facility, University of Arkansas at Fayetteville. Data dependent analysis (DDA) for

the in-gel trypsin digested samples from each condition were performed by using an Agilent 1200

series micro flow HPLC in line with Bruker Amazon-SL quadrupole ion trap ESI mass

spectrometer (QIT-ESI-MS). All the ESI-MS analyses were performed in a positive ion mode

using Bruker captive electrospray source with a dry nitrogen gas temperature of 200°C, with

nitrogen flow rate of 3 L/minute. LC-MS/MS data were acquired in the Auto MS(n) mode with

optimized trapping condition for the ions at m/z 1000. MS scans were performed in the enhanced

scanning mode (8100 m/z/second), while the collision-induced dissociation or the MS/MS

fragmentation scans were performed automatically for top ten precursor ions with a set threshold

for one minute in the UltraScan mode (32,500 m/z/second). Tryptic peptides were separated by

reverse-phase high-performance liquid chromatography (RP-HPLC) using a Zorbax SB C18

column, (150 × 0.3 mm, 3.5 μm particle size, 300 Å pore size, Agilent Technologies), with a

solvent flow rate of 4 μL/minute, and a gradient of 5%–38% consisting of 0.1% FA (solvent A)

and ACN (solvent B) over a time period of 320 minutes. Tryptic peptides were then identified by

searching MS/MS data in *S.* Typhimurium 14028S reference proteome database (24, 35) by using

MASCOT database search software (34). MS1 intensities of the integrated areas of these identified

tryptic peptides were compiled and grouped in skyline software according to the

replicates/conditions to perform statistical analysis. Targeted protein work were performed using

Shimadzu UPLC-20A coupled to 8050 triple quadrupole ESI-MS with heated probe. Sequence

specific fragment ion intensities from at least three unique tryptic peptides from the protein of

interest were used in the protein quantitation. Multiple reaction monitoring (MRM) events

corresponding to sequence specific fragment ions derived from the precursor tryptic peptides were

targeted to operate at a certain specific retention time intervals predicted by in house retention time

library. This library was generated using the correlation of relative hydrophobicity of the tryptic

peptides with their retention times (RT) from highly common housekeeping proteins, for the UPLC

method used in this analysis as described below. While the RT were correlated well within 99%

confidence, sufficient number of sequence specific fragment ions were used as basis for

identification of the tryptic peptide by MS/MS alone. Specificity and the confidence was achieved

by incorporating RT prediction. In addition to the application of skyline in quantitation, skyline

software was also used in predicting RT and optimizing parameters such as collision energies and

voltages with the help of Shimadzu Labsolution software. Tryptic peptides were separated by

reverse-phase ultra-high-performance liquid chromatography (RP-UPLC) compatible Shimadzu

C18, 1.9-micron particle size, 50x2.1 mm column (SN # 16041880T), with a solvent flow rate of

0.3 mL/minute, and a gradient of 5%–90% consisting of 0.1% FA (solvent A) and 0.1% FA in

ACN (solvent B) over a time period of 10 minutes.

Accession numbers

Tn-seq sequencing data are available on NCBI Sequence Read Archive. LB: PRJNA352537;

H_2O_2L: PRJNA352862; H_2O_2H: PRJNA352865.

Funding

The bench fee of this work was came from a fund from Arkansas Biosciences Institute. The first

author was supported by his parents, Cell and Molecular Biology (CEMB) program at the

University of Arkansas, and Human Capacity Development Program-Kurdistan Regional

Government (HCDP-KRG).

Acknowledgment

We thank Jack Jiang, a high school student, for writing the demultiplexing Python script. We are

thankful for Dr. Thomas R. Ioerger and Dr. Michael A. DeJesus at Texas A&M University for the

help on the use of TRNASIT. We extend our special thanks to Dr. Jeff F. Pummill and Dr. Pawel

Wolinski at Arkansas High Performance Computing Center for their help to use the facility.

Author Contributions

Conceived and designed the experiments: YK SK. Performed the experiments, analyzed the data,

wrote the manuscript: SK. Proteomics work: RL SK AQ JL. Revised the manuscript: YK SK. All

authors read the final version of the manuscript.

Supplementary Information

Data Set S1

Full list of *Salmonella* Tn-seq genome for H_2O_2L and H_2O_2H analyzed by ARTIST, Tn-Seq

Explorer, and ARTIST. The list of 137 *Salmonella* genes for H_2O_2 resistance.

Data Set S2

Lag time, growth rate, and maximum $OD6_{00}$ of *Salmonella* grown in LB, H_2O_2L, H_2O_2H.

Data Set S3

Salmonella proteomic analysis in response to the H_2O_2, non-targeted and targeted proteomics.

Table S1

Salmonella Tn-seq sequencing in numbers for LB, H_2O_2L, and H_2O_2H

Table S2

List of functional categories required for *Salmonella* H_2O_2 resistance.

Table S3

List of deferentially expressed proteins of *Salmonella* in response to H_2O_2 and their pathways.

Fig S1

Stability of H_2O_2 in the medium and during the experiment time.

Fig S2

Tn-seq library preparation diagram for Illumina sequencing.

Fig S3

Correlation between reads of *Salmonella* Tn-seq conditions, H_2O_2L and H_2O_2H.

REFERENCES

1. **LaRock DL, Chaudhary A, Miller SI.** 2015. Salmonellae interactions with host
processes. Nat Rev Microbiol **13:**191-205.

2. **Scallan E, Hoekstra RM, Angulo FJ, Tauxe RV, Widdowson MA, Roy SL, Jones JL,**
Griffin PM. 2011. Foodborne illness acquired in the United States – major pathogens.
Emerg Infect Dis **17:**1-15.

3. **Majowicz SE, Musto J, Scallan E, Angulo FJ, Kirk M, O'Brien SJ, Jones TF, Fazil A,**
Hoekstra RM. 2010. The global burden of nontyphoidal *Salmonella* gastroenteritis. Clin
Infect Dis **50:**882-889.

4.   **Haraga A, Ohlson MB, Miller SI.** 2008. Salmonellae interplay with host cells. Nat Rev
Microbiol **6:**53-66.

5.   **Nauseef W.** 2004. Assembly of the phagocyte NADPH oxidase. Histochem Cell Biol
122:277-291.

6.   **Seaver LC, Imlay JA.** 2001. Hydrogen peroxide fluxes and compartmentalization inside
growing *Escherichia coli*. J Bacteriol **183:**7182-7189.

7.   **Park S, You X, Imlay JA.** 2005. Substantial DNA damage from submicromolar
intracellular hydrogen peroxide detected in Hpx- mutants of *Escherichia coli*. Proc Natl
Acad Sci U S A **102:**9317-9322.

8.   **Imlay JA, Linn S.** 1988. DNA damage and oxygen radical toxicity. Science **240:**1302-
1309.

9.   **Imlay JA.** 2006. Iron-sulphur clusters and the problem with oxygen. Mol Microbiol
59:1073-1082.

10.   **Slauch JM.** 2011. How does the oxidative burst of macrophages kill bacteria? Still an open
question. Mol Microbiol **80:**580-583.

11.   **Hébrard M, Julie P. M. Viala, Méresse S, Barras F, Aussel L.** 2009. Redundant
Hydrogen Peroxide Scavengers Contribute to *Salmonella* Virulence and Oxidative Stress
Resistance. J Bacteriol **191:**4605-4614.

12.   **Horst SA, Jaeger T, Denkel LA, Rouf SF, Rhen M, Bange F.** 2010. Thiol peroxidase
protects *Salmonella enterica* from hydrogen peroxide stress *in vitro* and facilitates
intracellular growth. J Bacteriol **192:**2929-2932.

13.   **Morgan RW, Christman MF, Jacobson FS, Storz G, Ames BN.** 1986. Hydrogen
Peroxide-Inducible Proteins in *Salmonella* typhimurium Overlap with Heat Shock and

Other Stress Proteins. Proc Natl Acad Sci U S A **83:**8059-8063.

14.   **Zheng M, Wang X, Templeton LJ, Smulski DR, LaRossa RA, Storz G.** 2001. DNA

microarray-mediated transcriptional profiling of the *Escherichia coli* response to hydrogen

peroxide. J Bacteriol **183:**4562-4570.

15.   **Kroger C, Colgan A, Srikumar S, Handler K, Sivasankaran S, Hammarlof D, Canals**

R, Grissom J, Conway T, Hokamp K, Hinton J. 2013. An infection-relevant

transcriptomic compendium for *Salmonella enterica* Serovar Typhimurium. Cell Host

Microbe **14:**683-695.

16.   **Turner KH, Wessel AK, Palmer GC, Murray JL, Whiteley M.** 2015. Essential genome

of *Pseudomonas aeruginosa* in cystic fibrosis sputum. Proc Natl Acad Sci U S A **112:**4110-

4115.

17.   **Chan K, Kim CC, Falkow S.** 2005. Microarray-based detection of *Salmonella enterica*

Serovar Typhimurium transposon mutants that cannot survive in macrophages and mice.

Infect Immun **73:**5438-5449.

18.   **Silva-Valenzuela C, Molina-Quiroz R, Desai P, Valenzuela C, Porwollik S, Zhao M,**

Hoffman R, Andrews-Polymenis H, Contreras I, Santiviago C, McClelland M. 2016.

Analysis of two complementary single-gene deletion mutant libraries of *Salmonella*

Typhimurium in intraperitoneal infection of BALB/c Mice. Front Microbiol **6:**1455.

19.   **van Opijnen T, Bodi KL, Camilli A.** 2009. Tn-seq: high-throughput parallel sequencing

for fitness and genetic interaction studies in microorganisms. Nat Methods **6:**767-772.

20.   **Khatiwara A, Jiang T, Sung S, Dawoud T, Kim JN, Bhattacharya D, Kim H, Ricke**

SC, Kwon YM. 2012. Genome scanning for conditionally essential genes in *Salmonella*

enterica Serotype Typhimurium. Appl Environ Microbiol **78:**3098-3107.

21.   **Chaudhuri R, Morgan E, Peters S, Pleasance S, Hudson D, Davies H, Wang J, van**

Diemen P, Buckley A, Bowen A, Pullinger G, Turner D, Langridge G, Turner A,

Parkhill J, Charles I, Maskell D, Stevens M. 2013. Comprehensive assignment of roles

for *Salmonella* Typhimurium genes in intestinal colonization of food-producing animals.

PLoS Genet **9:**e1003456.

22.   **Langridge GC, Phan M, Turner DJ, Perkins TT, Parts L, Haase J, Charles I, Maskell**

DJ, Peters SE, Dougan G, Wain J, Parkhill J, Turner AK. 2009. Simultaneous assay of

every *Salmonella* Typhi gene using one million transposon mutants. Genome Res **19:**2308-

2316

23.   **Barquist L, Langridge GC, Turner DJ, Phan M, Turner AK, Bateman A, Parkhill J,**

Wain J, Gardner PP. 2013. A comparison of dense transposon insertion libraries in the

Salmonella serovars Typhi and Typhimurium. Nucleic Acids Res **41:**4549-4564.

24.   **Domon B, Aebersold R.** 2010. Options and considerations when selecting a quantitative

proteomics strategy. Nat Biotechnol **28:**710-721.

25.   **Jensen ON, Mann M.** 2003. Proteomic analysis of post-translational modifications. Nat

Biotechnol **21:**255-261.

26.   **Bogomolnaya L, Andrews K, Talamantes M, Maple A, Ragoza Y, Vazquez-Torres A,**

Andrews-Polymenis H. 2013. The ABC-Type efflux pump MacAB protects *Salmonella*

enterica serovar Typhimurium from oxidative stress. Mbio **4:**e00630.

27.   **Kwon YM, Ricke SC, Mandal RK.** 2016. Transposon sequencing: methods and

expanding applications. Appl Microbiol Biotechnol **100:**31-43.

28.   **Pritchard JR, Chao MC, Abel S, Davis BM, Baranowski C, Zhang YJ, Rubin EJ,**

Waldor MK. 2014. ARTIST: high-resolution genome-wide assessment of fitness using

transposon-insertion sequencing. PLoS Genet **10:**e1004782-e1004782.

29.   **Solaimanpour S, Sarmiento F, Mrazek J.** 2015. Tn-Seq Explorer: A tool for analysis of

high-throughput sequencing data of transposon mutant libraries. Plos One **10:**e0126070.

30.   **DeJesus MA, Ambadipudi C, Baker R, Sassetti C, Ioerger TR.** 2015. TRANSIT--A

software tool for Himar1 TnSeq analysis. PLoS Comput Biol **11:**e1004401-e1004401.

31.   **Huang DW, Sherman BT, Lempicki RA.** 2009. Bioinformatics enrichment tools: paths

toward the comprehensive functional analysis of large gene lists. Nucleic Acids Res **37:**1-

13.

32.   **Kanehisa M, Goto S.** 2000. KEGG: Kyoto Encyclopedia of genes and genomes. Nucleic

Acids Res **28:**27-30.

33.   **MacLean B, Tomazela D, Shulman N, Chambers M, Finney G, Frewen B, Kern R,**

Tabb D, Liebler D, MacCoss M. 2010. Skyline: an open source document editor for

creating and analyzing targeted proteomics experiments. Bioinformatics **26:**966-968.

34.   **Perkins D, Pappin D, Creasy D, Cottrell J.** 1999. Probability-based protein identification

by searching sequence databases using mass spectrometry data. Electrophoresis **20:**3551-

3567.

35.   **Cravatt BF, Yates III JR, Simon GM.** 2007. The biological impact of mass-

spectrometry-based proteomics. Nature **450:**991-1000.

36.   **Carlioz A, Touati D.** 1986. Isolation of superoxide dismutase mutants in *Escherichia coli*:

is superoxide dismutase necessary for aerobic life? Embo j **5:**623-630.

37.   **Weatherspoon-Griffin N, Yang D, Kong W, Hua Z, Shi Y.** 2014. The CpxR/CpxA Two-

component Regulatory System Up-regulates the Multidrug Resistance Cascade to

Facilitate *Escherichia coli* Resistance to a Model Antimicrobial Peptide. J Biol Chem

289:32571-32582.

38.   **Vinella D, Gagny B, Joseleau-Petit D, D'Ari R, Cashel M.** 1996. Mecillinam resistance

899   in *Escherichia coli* is conferred by loss of a second activity of the AroK protein. J Bacteriol

900   **178:**3818-3828.

901   39.   **Kaur P, Datta S, Shandil RK, Kumar N, Robert N, Sokhi UK, Guptha S, Narayanan**

902   **S, Anbarasu A, Ramaiah S.** 2016. Unravelling the secrets of mycobacterial cidality

903   through the lens of antisense. PLoS One **11:**e0154513.

904   40.   **Van Parys A, Boyen F, Leyman B, Verbrugghe E, Haesebrouck F, Pasmans F.** 2011.

905   Tissue-specific *Salmonella* Typhimurium gene expression during persistence in Pigs. Plos

906   One **6:**e24120.

907   41.   **Günel-Özcan A, Brown KA, Allen AG, Maskell DJ.** 1997. *Salmonella* typhimurium

908   aroB mutants are attentuated in BALB/ c mice. Microb Pathog **23:**311-316.

909   42.   **Nicolaou SA, Fast AG, Nakamaru-Ogiso E, Papoutsakis ET.** 2013. Overexpression of

910   fetA (ybbL) and fetB (ybbM), encoding an Iron exporter, enhances resistance to oxidative

911   stress in *Escherichia coli*. Appl Environ Microbiol **79:**7210-7219.

912   43.   **Robbe-Saule V, Coynault C, Ibanez-Ruiz M, Hermant D, Norel F.** 2001. Identification

913   of a non-haem catalase in *Salmonella* and its regulation by RpoS (σS). Mol Microbiol

914   **39:**1533-1545.

915   44.   **Craig M, Slauch J.** 2009. Phagocytic superoxide specifically damages an

916   extracytoplasmic target to inhibit or kill *Salmonella*. Plos One **4:**e4975.

917   45.   **Tsolis RM, Bäumler AJ, Heffron F, Stojiljkovic I.** 1996. Contribution of TonB- and Feo-

918   mediated iron uptake to growth of *Salmonella* typhimurium in the mouse. Infect Immun

919   **64:**4549-4556.

46.   **Aussel L, Zhao W, Hebrard M, Guilhon AA, Viala JPM, Henri S, Chasson L, Gorvel**

JP, Barras F, Meresse S. 2011. *Salmonella* detoxifying enzymes are sufficient to cope

with the host oxidative burst. Mol Microbiol **80:**628-640.

47.   **Taylor PD, Inchley CJ, Gallagher MP.** 1998. The *Salmonella* typhimurium AhpC

polypeptide is not essential for virulence in BALB/c Mice but is recognized as an antigen

during infection. Infect Immun **66:**3208-3217.

48.   **Papp-Szabò E, Firtel M, Josephy PD.** 1994. Comparison of the sensitivities of

Salmonella typhimurium oxyR and katG mutants to killing by human neutrophils. Infect

Immun **62:**2662-2668.

49.   **Imlay J.** 2015. Diagnosing oxidative stress in bacteria: not as easy as you might think. Curr

Opin Microbiol **24:**124-131.

50.   **Carlsson J, Carpenter VS.** 1980. The recA+ gene product is more important than catalase

and superoxide dismutase in protecting *Escherichia coli* against hydrogen peroxide

toxicity. J Bacteriol **142:**319-321.

51.   **Ishioka K, Iwasaki H, Shinagawa H.** 1997. Roles of the recG gene product of *Escherichia*

coli in recombination repair: Effects of the Delta recG mutation on cell division and

chromosome partition. Genes Genet Syst **72:**91-99.

52.   **Sheikh A, Charles R, Sharmeen N, Rollins S, Harris J, Bhuiyan M, Arifuzzaman M,**

Khanam F, Bukka A, Kalsy A, Porwollik S, Leung D, Brooks W, LaRocque R,

Hohmann E, Cravioto A, Logvinenko T, Calderwood S, McClelland M, Graham J,

Qadri F, Ryan E. 2011. In Vivo Expression of *Salmonella enterica* Serotype Typhi Genes

in the Blood of Patients with Typhoid Fever in Bangladesh. PLoS Negl Trop Dis **5:**e1419.

53.   **Lu S, Killoran PB, Riley LW.** 2003. Association of *Salmonella enterica* Serovar

Enteritidis YafD with resistance to chicken egg albumen. Infect Immun **71:**6734-6741.

54.   **Demple B, Halbrook J, Linn S.** 1983. *Escherichia coli* xth mutants are hypersensitive to

hydrogen peroxide. J Bacteriol **153:**1079-1082.

55.   **Sassetti CM, Rubin EJ.** 2003. Genetic requirements for mycobacterial survival during

infection. Proc Natl Acad Sci U S A **100:**12989-12994.

56.   **Sabbagh S, Lepage C, McClelland M, Daigle F.** 2012. Selection of *Salmonella enterica*

Serovar Typhi genes involved during interaction with human macrophages by screening of

a transposon mutant library. Plos One **7:**e36643.

57.   **Bogomolnaya L, Aldrich L, Ragoza Y, Talamantes M, Andrews K, McClelland M,**

Andrews-Polymenis H. 2014. Identification of novel factors involved in modulating

motility of *Salmonella enterica* Serotype Typhimurium. Plos One **9:**e111513.

58.   **Schwartz CJ, Giel JL, Patschkowski T, Luther C, Ruzicka FJ, Beinert H, Kiley PJ.**

2001. IscR, An Fe-S cluster-containing transcription factor, represses expression of

Escherichia coli genes encoding Fe-S cluster assembly proteins. Proc Natl Acad Sci U S

A **98:**14895-14900.

59.   **Unciuleac M, Chandramouli K, Naik S, Mayer S, Huynh B, Johnson M, Dean D.** 2007.

Vitro activation of apo-aconitase using a [4Fe-4S] cluster-loaded form of the IscU [Fe-S]

cluster scaffolding protein. Biochemistry **46:**6812-6821.

60.   **Py B, Barras F.** 2010. Building Fe–S proteins: bacterial strategies. Nat Rev Microbiol

8:436-446.

61.   **Balbontín R, Rowley G, Pucciarelli MG, López-Garrido J, Wormstone Y, Lucchini**

S, Portillo FG, Jay C. D. Hinton, Casadesús J. 2006. DNA adenine methylation regulates

virulence gene expression in *Salmonella enterica* Serovar Typhimurium. J Bacteriol

188:8160-8168.

62.   **Badie G, Heithoff DM, Sinsheimer RL, Mahan MJ.** 2007. Altered levels of *Salmonella*

DNA adenine methylase are associated with defects in gene expression, motility, flagellar

synthesis, and bile resistance in the pathogenic strain 14028 but not in the laboratory strain

LT2. J Bacteriol **189:**1556-1564.

63.   **Sarnacki SH, Marolda CL, Llana MN, Giacomodonato MN, Valvano MA, Cerquetti**

MC. 2009. Dam methylation controls O-antigen chain length in *Salmonella enterica*

Serovar Enteritidis by regulating the expression of Wzz protein. J Bacteriol **191:**6694-

6700.

64.   **Lopez-Garrido J, Cheng N, Garcia-Quintanilla F, Garcia-del Portillo F, Casadesus J.**

2010. Identification of the *Salmonella enterica* damX gene product, an inner membrane

protein involved in bile resistance. J Bacteriol **192:**893-895.

65.   **Hengge-Aronis R.** 2002. Signal transduction and regulatory mechanisms involved in

control of the σS (RpoS) subunit of RNA polymerase. Microbiol Mol Biol Rev **66:**373-

395.

66.   **Lu M, Kleckner N.** 1994. Molecular cloning and characterization of the pgm gene

encoding phosphoglucomutase of *Escherichia coli*. J Bacteriol **176:**5847-5851.

67.   **Paterson GK, Cone DB, Peters SE, Maskell DJ.** 2009. The enzyme phosphoglucomutase

(Pgm) is required by *Salmonella enterica* serovar Typhimurium for O-antigen production,

resistance to antimicrobial peptides and in vivo fitness. Microbiology **155:**3403-3410.

68.   **Martínez-García E, Calles B, Arévalo-Rodríguez M, de Lorenzo V.** 2011. pBAM1: an

all-synthetic genetic tool for analysis and construction of complex bacterial phenotypes.

BMC Microbiol **11:**38-38.

69.   **Hall B, Acar H, Nandipati A, Barlow M.** 2014. Growth rates made easy. Mol Biol Evol

31:232-238.

70.   **Yao Q, Luo Z, Yuan X, Yu Y, Zhang C, Xie J, Lee JY.** 2014. Assembly of nanoions via

electrostatic interactions: ion-like behavior of charged noble metal nanoclusters. Sci Rep

4:3848-3848.

71.   **Chun J, Kim K, Hwang I, Kim Y, Lee D, Lee I, Kim J.** 2007. Dual priming

oligonucleotide system for the multiplex detection of respiratory viruses and SNP

genotyping of CYP2C19 gene. Nucleic Acids Res **35:**e40-e40.

72.   **Li H, Durbin R.** 2009. Fast and accurate short read alignment with Burrows-Wheeler

transform. Bioinformatics **25:**1754-1760.

73.   **Langmead B, Trapnell C, Pop M, Salzberg S.** 2009. Ultrafast and memory-efficient

alignment of short DNA sequences to the human genome. Genome Biol **10:**R25-R25.

1001

1002

1003

1004

1005

1006

1007

1008

1009

1010

1011

1012

1013

1014

1015

1016

1017

1018

FIG 1

The Tn-seq can identify the genes that are required H_2O_2 resistance. (A) The effect of H_2O_2 on the

growth rate of wild type *Salmonella* Typhimurium. An overnight culture of bacteria was diluted

1:200 in the LB medium contains either 2.5 mM H_2O_2 (H_2O_2L) or 3.5 mM H_2O_2 (H_2O_2H), LB

without H_2O_2 was used as control. The cultures were incubated at 37°C for 24 h in a 96-well plate.

The reduced growth rates for the H_2O_2 were in comparison to the control. In the all growth curve

figures in this work, the blue color represents just LB (no H_2O_2), the red is H_2O_2L, and green is

H_2O_2H. (B) Schematic representation of the Tn-seq study. The *Salmonella* transposon mutant

library was inoculated into LB and LB contains H_2O_2L or H_2O_2H. The three cultures were grown

until they reached mid-exponential phase. The DNA was extracted form each culture and subjected

to library preparation, sequencing, and data analysis. (C) The Tn-seq profile of the three

conditions. It shows 34 kb of the *Salmonella* genome which starts with *metF* gene and ends with

rrsB, horizontal axis. The height of vertical axis represents number of reads which is 1500 reads.

The highlighted genes in red are *katG*, catalase peroxidase, was tolerated insertions in the both

H_2O_2 conditions; *oxyR* was not tolerated insertions in presence of H_2O_2 and indicated that the gene

is required to H_2O_2 resistance; *murI*, glutamate racemase, was not tolerated any insertions at all

and considered as essential gene, its required for the biosynthesis of a component of cell wall

peptidoglycan.

1037

1038

1039

1040

1041

FIG 2

Growth curve of 50 mutants and a wild type *Salmonella* in the form of heat map. The lag phase

time, growth rate, and maximum OD$_{600}$ of the individual *Salmonella* Typhimurium mutants and

the wild type in the growth conditions of LB (no H_2O_2), H_2O_2L and H_2O_2H. The overnight culture

of the mutants and the wild type bacteria were diluted 1:200 in the LB medium, and the LB

contains either 2.5 mM H_2O_2 (H_2O_2L) or 3.5 mM H_2O_2 (H_2O_2H). The cultures were incubated at

37°C for 24 h in 96-well plate and the OD_{600} was recorded every 10 min. The lag phase time,

growth rate and maximum OD_{600} were calculated and shown here as a graphical representation.

The pale pink color indicates short lag phase time, high growth rate, and high OD_{600}. The red color

indicates that the bacteria was stayed in lag phase, growth rate was close to zero, and the OD_{600} of

the culture was not raised in the 24 h time of the assays.

1073

1074

1075

1076

1077

1078

1079

1080

1081

1082

1083

1084

1085

1086

1087

1088

FIG 3

The role of aromatic amino acid biosynthesis genes in resistance to the H_2O_2. (A) Schematic

representation of aromatic amino acid biosynthesis, adapted from the KEGG pathway database.

The genes in red color were identified by the Tn-seq for H_2O_2 resistance in *Salmonella*. The red

bold color genes were identified by the Tn-seq and the phenotypes were validated by the individual

mutant assays. (B) The overnight cultures of the individual mutants were diluted 1:200 in the LB

(no H_2O_2) and the LB contains either 2.5 mM H_2O_2 (H_2O_2L) or 3.5 mM H_2O_2 (H_2O_2H). The

cultures were incubated at 37°C for 24 h in 96-well plate. The color of growth curve figures is blue

for the LB, red is H_2O_2L, and green is H_2O_2H. In the $\varDelta$ *aroK* growth curve, the red color is under

the green color. (C) Differential expression of *Salmonella* proteins in response to the H_2O_2L

compared to the LB. Wild type *Salmonella* was grown in LB, H_2O_2L, and H_2O_2H until mid-

exponential phase. Targeted-proteomics was quantified AroB, AroE_2, AroK, and AroA protein

expressions in response to H_2O_2L. The shown peaks represent a unique peptide of the protein.

1102

1103

1104

1105

1106

1107

1108

1109

1110

1111

1112

1113

1114

1115

1116

1117

1118

FIG 4

The *ybbK*, *ybbL*, and *ybbM* have a role in resistance to H_2O_2. (A) The Tn-seq profile of the LB (no

H_2O_2), H_2O_2L (2.5 mM), and H_2O_2H (3.5 mM). It shows ~6 kb of *Salmonella* Typhimurium

genome starts with *ybbK* and ends with *ybbO*, horizontal axis. The read scale for the conditions

are 4000, vertical axis. (B) The growth curve of *ΔyybK*, *ΔyybL*, and *ΔyybM*. The overnight cultures

of these three mutants were diluted 1:200 in LB, H_2O_2L, and H_2O_2H. The cultures were incubated

at 37°C for 24 h in 96-well plate. The growth curve colors are blue which represents just LB or no

H_2O_2, the red is H_2O_2L, and green is H_2O_2H. (C) Wild type *Salmonella* was grown in LB, H_2O_2L,

and H_2O_2H until mid-exponential phase. Targeted-proteomics was quantified YbbK, YbbL, and

YbbM protein expressions in response to H_2O_2L. The shown peaks represent a unique peptide of

the protein.

51