1

A Comprehensive Assessment of the Genetic Determinants in *Salmonella* Typhimurium for

Resistance to Hydrogen Peroxide Using Proteogenomics

Sardar Karash[1], Rohana Liyanage[2,3], Abdullah Qassab[1], Jackson O. Lay, Jr.[2,3], and

Young Min Kwon[1,4*]

[1]Cell and Molecular Biology Program, University of Arkansas, Fayetteville, AR 72701,

[2]Department of Chemistry, University of Arkansas, Fayetteville, AR 72701,

[3]Statewide Mass Spectrometry Facility, Fayetteville, AR 72701, and

[4]Department of Poultry Science, University of Arkansas, Fayetteville, AR 72701

*Corresponding author:

Department of Poultry Science

College of Agricultural, Food and Life Sciences

University of Arkansas

Fayetteville, AR 72701

Phone : (479) 575-4935

Email ykwon@uark.edu

Abstract

Salmonella is an intracellular pathogen that infects a wide range of hosts and can survive in macrophages. An essential mechanism uses by the macrophages to eradicate *Salmonella* is production of reactive oxygen species. Here, we used proteogenomics to determine the candidate genes and proteins that have a role in resistance of *S*. Typhimurium to H_2O_2. For Tn-seq, a highly saturated Tn5 insertion library was grown *in vitro* under either 2.5 (H_2O_2L) or 3.5 mM H_2O_2 (H_2O_2H). We identified two sets of overlapping genes that are required for resistance of *S*. Typhimurium to H_2O_2L and H_2O_2H, and the results were validated via phenotypic evaluation of 50 selected mutants. The enriched pathways for resistance to H_2O_2 included DNA repair, aromatic amino acid biosynthesis (*aroBK*), Fe-S cluster biosynthesis, iron homeostasis and a putative iron transporter system (*ybbKLM*), flagellar genes (*fliBC*), H_2O_2 scavenging enzymes, and DNA adenine methylase. Proteomics revealed that the majority of essential proteins, including ribosomal proteins, were downregulated upon exposure to H_2O_2. A subset of proteins identified by Tn-seq were analyzed by targeted proteomics, and 70 % of them were upregulated upon exposure to H_2O_2. The identified candidate genes will deepen our understanding about mechanisms of *S*. Typhimurium survival in macrophages, and can be exploited to develop new antimicrobial drugs.

Introduction

Salmonella is a Gram-negative bacterium that infects humans and animals. *Salmonella enterica* has numerous serovars, which include typhoidal and non-typhoidal strains. In contrast to the typhoidal salmonellae which are human restricted pathogens, the non-typhoidal salmonellae (NTS), serovar Enteritidis and Typhimurium, are able to infect a wide range of hosts, causing gastroenteritis[1]. The NTS strains, including *Salmonella enterica* serovar Typhimurium, account for 11% (1.2 million cases) of the total foodborne illnesses caused by different pathogens in the United States[2]. It has been estimated that *Salmonella* is responsible for 93.8 million cases of gastroenteritis, leading to 155,000 deaths worldwide annually[3]. The pathogen remains a continuous threat to the food safety, and public health.

To initiate an infection and survive inside the host, *Salmonella* needs to overcome a myriad of host defense mechanisms. As *Salmonella* reaches the intestine and breaches the epithelial tissue, it enters the macrophages and activates different virulence strategies in order to survive and replicate in them[4]. An essential mechanism uses by the phagocytes to kill and eradicate *Salmonella* is production of reactive oxygen species (ROS). Hydrogen peroxide (H_2O_2), superoxide anion ($O_2^{.-}$), and the hydroxyl radical ($HO^{.}$) are derivatives of ROS. The short-lived $O_2^{.-}$, produced by the NADPH-dependent phagocytic oxidase, quickly dismutates into H_2O_2, which diffuses across semipermeable bacterial cell membranes. Eventually, Fe^{2+} reduces H_2O_2 to $HO^{.}$ via the so called Fenton Reaction[5-7]. The ROS, including H_2O_2, can damage DNA, iron-sulfur cluster-containing proteins, and other biological molecules in the bacterial cells[8-10].

Numerous genetic factors and proteins that are important for resistance of *S*. Typhimurium to H_2O_2 have been discovered and the underlying mechanisms have been explored[11, 12]. A various approaches and techniques have been employed to study global response of *Salmonella* or

related bacteria to H_2O_2 *in vitro* as a model system to simulate the bacterium's response to ROS

in phagocytic cells: (i) Two-dimensional gel electrophoresis identified H_2O_2-induced proteins in

Salmonella[13], (ii) DNA microarray identified H_2O_2 induced genes in *E. coli*[14], and (iii) RNA-seq

identified H_2O_2 induced genes in *Salmonella*[15]. Yet, the factors required for fitness under the

given condition cannot be identified with high confidence based on the analysis of

transcriptomics or proteomics data[16]. Microarray-based tracking of random transposon insertions

was used to identify numerous genes in *Salmonella* that are required for survival in mice and

macrophages[17-18]. However, the genetic factors responsible for resistance to ROS cannot be

sorted out among all of the genetic factors identified in the study that are required for fitness in

the presence of multiple host stressors.

To shed more insights into the underlying mechanisms of *Salmonella* resistance to H_2O_2, more

direct approach linking the gene-phenotype relationships in a genome-wide scale would be

necessary. Tn-seq is a powerful approach to allow direct and accurate assessment of the fitness

requirement of each gene on the entire genome of a prokaryotic organism[19]. In Tn-seq method, a

saturated transposon insertion library (input) is exposed to a selective condition, and the mutant

population altered through the selection (output) is recovered. Then, the genomic junctions of the

transposon insertions are specifically amplified and sequenced from both input and output pools

by high-throughput sequencing. The gene fitness can be obtained by calculating the change in

relative abundance of the sequence reads corresponding to each gene in the entire genome

between the two pools. Tn-seq has been employed to assign gene functions to *Salmonella*

genomes in numerous studies: (i) Previously, our lab identified conditionally essential genes that

are required for growth in the presence of bile, limited nutrients, and high temperature[20], (ii) The

genes required for intestinal colonization were identified in chickens, pigs, and cattle[21], (iii)

Candidate essential genes and genes contributing toward bile resistance were identified[22], (iv)

Core conserved genes for growth in rich media were identified in serovars Typhi and

Typhimurium[23]. In addition to Tn-seq, electrospray ionization liquid chromatography tandem

mass spectrometry (ESI-LC-MS/MS) is a powerful approach for identifying and quantifying

proteins in a large scale. The system-wide protein regulation can be determined using mass

spectrometry signal intensities of tryptic peptides obtained from two different culture

conditions[24]. The post-translational modification in proteins can be revealed by using proteomic

analysis[25]. Many studies took advantage of proteomic analysis of *Salmonella*. However, to the

best of our knowledge, this study is the first to investigate proteogenomics of a bacterium by

combining Tn-seq and proteome analysis simultaneously to the same stressor.

In this work, we used Tn-seq method and proteomic analysis in combination to determine

system-wide responses of *S.* Typhimurium to two different concentrations of H_2O_2 (H_2O_2L and

H_2O_2H). We obtained a comprehensive list of 137 genes that are putatively required for the

resistance of *S.* Typhimurium 14028 to H_2O_2. The role of 50 selected genes in resistance to H_2O_2

were determined by phenotypic evaluation of the individual deletion mutants. Also, we identified

a set of 246 proteins that are differentially expressed in response to H_2O_2, using data-dependent

acquisition (DDA) proteomics, which are largely overlapped with the genes identified by Tn-seq;

targeted proteomics showed 70% of the proteins identified by Tn-seq were upregulated by H_2O_2.

In addition to the genes of *S.* Typhimurium previously known to be important for resistance to

H_2O_2, we identified approximately 80 genes that have not been previously associated with

resistance to oxidative stress. The results of this study highlighted that the genes in aromatic

amino acid biosynthesis, *aroB* and *aroK*, and iron homeostasis, *ybbK*, *ybbL*, and *ybbM*, are

crucially important for growth fitness under H_2O_2 stress. The identified candidate genes will

expand our understanding on the molecular mechanisms of *Salmonella* survival in macrophages,

and serve as new antimicrobial drug targets.

Results and Discussion

The H_2O_2 concentrations and the selections of Tn5 library

First, we sought to determine the growth response of wild type *S*. Typhimurium 14028 cells in

LB media containing varying concentrations of H_2O_2. The wild type cells were grown in LB

media that contain different concentrations of H_2O_2 in 96-well plates. After evaluating the

growth rates for the cultures, 2.5 and 3.5 mM H_2O_2 were chosen for Tn-seq selections in our

study, and termed H_2O_2L and H_2O_2H, respectively. In comparison to *Salmonella* grown in LB

media with no H_2O_2, H_2O_2L and H_2O_2H reduced the growth rates by 10% and 28%, respectively

(Fig 1A). The lag time increased by a 5.7-fold (0.5 vs. 2.9 hr), and an 11-fold (0.5 vs. 5.6 hr) in

H_2O_2L and H_2O_2H, respectively. The maximum OD_{600} decreased by only 1% for the H_2O_2L and

2% for the H_2O_2H in comparison to LB media (Fig 1A).

For the selection of Tn5 library, 20 ml cultures in 300 ml Erlenmeyer flasks containing LB,

H_2O_2L, or H_2O_2H were inoculated with the same Tn5 library at the seeding CFUs of the library

at 3.5×10^6. This seeding level provided ~10 CFUs for each Tn5 insertion mutant in the library.

The cultures were grown until the mid-exponential phase, in which the CFUs reached 1.17×10^8

(SE 0.01×10^8). It required 7.5 and 9.2 h to reach the cell density as measured by optical density

for H_2O_2L and H_2O_2H, respectively, in contrast to 5 h for LB medium (Fig 1B). We observed

some differences in growth responses between the cultures in a 96-well plate and in a 300-ml

Erlenmeyer flask. The optical density readings by the plate reader was different in comparison to

those by Bio-photometer that we used to measure optical density of the culture in the flask. As a

result, the growth curve in Fig 1A which was based on 96-well plate reader, dose not match

exactly with the time required for the Tn5 library to reach the target mid-exponential phase in the

flask cultures. In addition, we observed that the H_2O_2 is stable in LB media free of *Salmonella*

during the window of time used for the library selection process (Fig. S1), which was also

supported by Bogomolnaya et al.[26].

Preparation of Tn5-seq amplicon library

The *Salmonella* mutants were generated by using the delivery plasmid pBAM1 via conjugation.

A total of 325,000 mutant colonies were recovered from 50 plates. Each mutant contained a

single random insertion of Tn5 transposon in the chromosome or plasmid according to DNA

sequencing of Tn5-junction sequences for a small set (n = 71) of randomly selected Tn5 mutants.

We found a significant portion (~20%) of the mutants in the library that were not genuine Tn5

insertions, but the mutants generated as a result of pBAM1 integration into chromosome as

determined by their ability to grow in the presence of ampicillin. To prevent the Illumina

sequencing reads of being wasted on sequencing Tn5 junctions from these cointegrants, we

digested genomic DNA of the input and output libraries with PvuII, which digests immediately

outside the inverted repeats on both sides of Tn5. The digested DNA was then used to prepare

Tn-seq amplicon library as described in Materials and Methods. Our Tn-seq data analysis

indicated that our strategy of removing the DNA sequences originating from cointegrants was

effective because only 0.55% of the total HiSeq reads corresponding to Tn5-junctions matched to

pBAM1. It should be possible to remove them completely by ensuring complete digestion of

genomic DNA with PvuII. The method for Tn-seq amplicon library we developed and used in

this study has multiple advantages over other Tn-seq protocols, because our method requires

only 100 ng of the genomic DNA, and the whole process can be completed in a day[27]. When the

extension step in the protocol was performed using a conventional 20 nucleotide primer, and the

final products of exponential PCR were separated on agarose gel electrophoresis, even the

negative controls (the wild type genomic DNA or mutant library genomic DNA without linear

extension) showed smear patterns of nonspecific background amplification. However, when dual

priming oligonucleotide (DPO) primer was used in place of the conventional primer for linear

extension, non-specific background amplification was completely disappeared. Therefore, we

adopted the DPO primer in linear extension step for all library samples in this study. Then, the

single-stranded extension products were C-tailed, and used as templates for the exponential PCR

step using nested primer specific to Tn5 and poly G primer that contain Illumina adapter

sequences along with sample index sequences (Fig. S2). The final PCR products were separated

on an agarose gel, and the fragments within the range of 325-625 bp were gel-purified. After

pooling of multiple samples, the combined library was sequenced on a HiSeq 3000.

Summary of Tn-seq DNA analysis

175
After de-multiplexing and C-tail trimming of all sequence reads, ~72 million reads of Tn5-

junctions with mean read length of 94 bp were obtained. The number of the reads mapped to the

complete genome of *S*. Typhimurium 14028 were ~25, 15, and 19 million for LB, H_2O_2L, and

H_2O_2H, respectively. The number of unique insertions on the chromosome were 125,449 in the

input library, excluding the plasmid (Table S1). On average, Tn5 was inserted in every 39 bp.

Number of raw reads per open reading frame (ORF) for H_2O_2L was plotted over the

corresponding number of H_2O_2H, which yielded an R^2 of 0.91, indicating the mutants in the

input library quantitatively responded in a similar way for both H_2O_2L and H_2O_2H as expected

(Fig. S3). The insertions were mapped to 5,428 genes or 8,022 genes/intergenic regions.

Interestingly, the ORF STM14_5121, which is 16.7 kbp long, had the highest number of

insertions (~700 insertions) and reads (0.25 M).

Comparison of various bioinformatics pipelines for Tn-seq data analysis

We used 3 different Tn-seq analysis tools to identify the genes and compare the results across the

methods with the goal of comprehensive identification of "all" genes required for resistance to

H_2O_2. The first tool, ARTIST[28], created small non-overlapping genomic windows of 100 bp and

the reads from each window were arbitrarily assigned into the middle of the window. The default

normalization script of the tool was used. Then, the relative proportions of insertion sites in the

output library versus the input were tabulated. Mann-Whiney U (MWU) test was used to assess

the essentiality of the locus. To consider a gene/intergenic region conditionally essential for

growth in the presence of H_2O_2, p value had to be $\leq$ 0.05 in 90 of the 100 conducted MWU tests.

Subsequently, 20 genes and 1 intergenic region were identified for H_2O_2L and 4 genes for

H_2O_2H (Table S2). We speculate the reason that more genes were identified for H_2O_2L in

comparison to H_2O_2H, was partially due to the lower number of total reads of H_2O_2L as

compared to H_2O_2H, even though the read numbers of H_2O_2L was normalized to those of the

input.

The second tool, Tn-seq Explorer[29], counted insertions in overlapping windows of a fixed size.

Using a 550 bp window size, each annotated gene was assigned an essentiality index (EI) which

is determined mainly based on the insertion count in a window in this gene. The bimodal

distribution of insertion counts per window divided the essential genes to the left and the non-

essential genes to the right. To find conditional essential genes, the EI of the output was

subtracted from the EI of the input. The genes with negative ΔEI were ranked based on the

change in read fold change (Log2 (H_2O_2L or H_2O_2H/Input)). We found 114 consensus genes

between H_2O_2L and H_2O_2H that had at least four-fold reduction in H_2O_2H read counts as

compared to the input. The four-fold reduction (Log2FC = -2) threshold was chosen based on our

validation study of Tn-seq data by single mutant assays (Table S2).

The third tool, TRANSIT[30], determined read counts of genes in the input and output library. The

differences of total read counts between the input and outputs were obtained. The insertion sites

were permutated for a number that is specified by the user (we used 10,000 sample). This

sampling for each gene gave difference in read counts. The p value was calculated from the null

distribution of the difference in read counts. We identified 8 and 21 genes for the H_2O_2L and

H_2O_2H, respectively, using a p value ≤ 0.05 (Table S2).

The combined list of the genes identified by the 3 Tn-seq analysis tools for both H_2O_2L and

H_2O_2H included 137 genes (Table S2). All of the genes on this list are expected to have a role in

conferring resistance to H_2O_2 and allow *Salmonella* to survive and replicate in the presence of

H_2O_2 *in vitro*. Of the 21 genes identified by TRANSIT, 19 of these genes were also identified by

Tn-seq Explorer, but only 3 out of this 21 were identified by ARTIST. The 19 genes were *hscA*,

rbsR, *fepD*, *efp*, *oxyR*, *polA*, *ybaD*, *aroD*, *ruvA*, *xthA*, *dps*, *aroB*, *uvrD*, *tonB*, *uvrA*, *aroK*, *ybbM*,

lon, and *proC*. Two genes, *fepD* and *xthA*, were identified by the all 3 methods and for both

conditions.

The 3 Tn-seq analysis tools are very valuable for Tn5 data analysis, but each tool has its own

advantages and disadvantages. For ARTIST, (i) the user must know how to run scripts in Matlab

software, (ii) the analysis is very slow on a personal computer with the HiSeq data, (iii) it has

only one method for normalization, but (iv) it can search for essentiality in the intergenic

regions. For Tn-seq Explorer, (i) there is no data normalization, and (ii) prediction on small

genes is prone to be inaccurate, but (iii) its very user-friendly and runs fast. For TRANSIT, (i)

the user should have some knowledge on running scripts on terminal, (ii) it may need some

modification in its Python script according to the way the library was prepared for sequencing,

and (iii) a few software packages should be installed on the computer as TRANSIT pre-

requisites, but (iv) it does have 6 different methods for data normalization and it runs very fast on

a personal computer. Although ARTIST and Tn-seq Explorer are very useful tools for Tn-seq

data analysis, we prefer using TRANSIT in our future data analysis for conditionally essential

genes. In the following sections, we continued the downstream analysis mainly based on the 137

genes that include all of the genes identified by all 3 methods.

The enriched pathways for resistance to H_2O_2
241
In order to categorize the identified genes that are required for *Salmonella* resistance to the H_2O_2,

the 137 genes were subjected to pathway enrichment analysis using DAVID Bioinformatics

Resources 6.7, NIAID/NIH[31]. A total of 15 KEGG pathways[32] were recognized for 69 genes on

the list. The enriched pathways include homologous recombination (*ruvC*, *polA*, *ruvA*, *ruvB*,

priB, *recA*, *recR*, *holC*, *holD*, *recC*, *recG*), nucleotide excision repair (*uvrD*, *polA*, *uvrA*, *uvrC*),

mismatch repair (*dam*, *uvrD*, *holC*, *holD*), RNA degradation (*pnp*, *hfq*, *ygdP*), purine and

pyrimidine metabolism (*apaH*, *polA*, *pnp*, *arcC*, *spoT*, *holC*, *holD*, *cmk*, *dcd*, *pnp*),

phenylalanine, tyrosine and tryptophan biosynthesis (*aroD*, *aroB*, aroA, *aroK*, *aroE_2*), arginine

and proline metabolism (*proC*, *arcC*), glycolysis and gluconeogenesis (*crr*, *pgm*, *tpiA*), oxidative

phosphorylation (*atpG*, *atpA*, *cydA*), DNA replication (*polA*, *holC*, *holD*), and flagellar assembly

(*fliJ*, *fliD*, *flhD*, *fliC*). Since KEGG was not able to recognize many genes on the list, we used

SP_PIR_Keywords of functional categories, which recognized majority of the genes and

categorized them into 55 functional categories (Table S3), excluding 15 uncharacterized genes

(ORFs). Among these categories were stress response (*rpoE*, *lon*, *dnaJ*, *hfq*, *yaiB*), iron (*dps*,

entD, *iscA*, *yjeB*, *yhgI*), and transcription regulation (*rcsA*, *oxyR*, *rpoE*, *yjeB*, *arcA*, *argR*, *rbsR*,

rpoS, *fadR*, *rcsB*, *furR*, *flhD*).

Validation of Tn-seq results using individual mutants

For the selected 50 genes among the 137 genes identified by Tn-seq, the growth phenotype was

determined using individual single deletion mutants in LB, H_2O_2L, and H_2O_2H. The genes were

considered to play a role in resistance to H_2O_2, if (i) lag phase time increased, (ii) growth rate

reduced or (iii) maximum OD_{600} decreased in the presence of H_2O_2 in comparison to the wild

type strain grown in the same conditions. Of the 50 single deletion mutants, 42 mutants were

shown to have a role in resistance to H_2O_2 (Fig. 2 and Table S4). One gene, *yhaD*, was identified

by all 3 analysis tools, but it did not show the expected phenotype. The *fliD* was also identified

by ARTIST, but did not show any phenotype distinguishable from the wild type. The remaining

6 genes that did not show the phenotype was identified by Tn-seq Explorer. Based on the results

of the individual mutant assay, we conclude that 84% (42/50) of the genes identified by the Tn-

seq analysis and tested using single deletion mutants have a role for resistance to H_2O_2. These

results indicate that our Tn-seq analysis identified the genes in *S.* Typhimurium that are required

for the wild type level resistance to H_2O_2 with high accuracy.

Proteomics of H_2O_2 response

With ESI-LC-MS/MS in data-dependent acquisition (DDA) mode, the protein regulation was

determined using MS1 filtering technique that skyline software offers[33]. It uses signal intensities

of tryptic peptides derived from the proteins of wild type strain grown in the presence of H_2O_2 in

comparison to the control (LB). As described in Materials and Methods section, trypsin digestion

of the protein extracts under different conditions generates tryptic peptides that are uniquely

related to individual proteins. Tryptic peptides separated by liquid chromatography from the

complex samples were first subjected to simple mass measurement (MS1) followed by intensity

dependent fragmentation of these peptide ions to produce sequence specific fragment ions by

collision-induced dissociation (MS/MS). Tryptic peptides were then identified using these

sequence specific fragment ions via MASCOT database search software[34], where the sequence

specific fragment ions were matched to the proteins in *S*. Typhimurium 14028S reference

proteome database[24, 35]. This method of protein analysis is normally referred to as data dependent

analysis (DDA). At the beginning of data analysis, the H_2O_2L and H_2O_2H data were compared to

LB separately, however it turned out that comparison was not sensitive enough to differentiate

between H_2O_2L and H_2O_2H conditions. Hence, the data of H_2O_2L and H_2O_2H were combined for

analysis in comparison to LB. We identified 1,104 proteins of *Salmonella* for the 3 conditions

(Table S5); of these, 246 proteins were differentially expressed in response to H_2O_2 with *p* values

≤ 0.05 and 90% CI. Proteomics analysis showed that 121 and 125 proteins were upregulated and

downregulated in response to stress by H_2O_2, respectively. Since Tn-seq revealed genetic

requirements for fitness under the selection conditions, the identified genes are expected to

express corresponding proteins under the conditions to perform their cellular functions. Often the

proteins required for fitness under a given condition are overexpressed under the condition, but it

may not be the case for some proteins. In this study, we had a unique opportunity to

comparatively analyze both Tn-seq and the MS data to understand the relationship between

genetic requirements and changes in expression level under the condition of interest, which was

H_2O_2 in this study. We also obtained the list of essential genes based on our Tn-seq data, which

could not tolerate insertions by definition, and if we were not certain about essentiality of a gene

from our Tn-seq data, the gene was searched for essentiality in the previously reported list of

Salmonella essential genes 22. The comprehensive list of essential genes allowed us to study any

correlation between the essentiality and the changes in protein expression. Among the 246

proteins, there were 78 essential and 168 non-essential proteins. Among the 78 essential proteins,

25 were upregulated whereas 53 were downregulated. On the contrary, the majority (n = 96) of

the detected non-essential proteins were upregulated, while 72 non-essential proteins were

downregulated. To further examine the quantitative relationships closely, 64 genes/proteins

identified by both methods (Table S5) were focused on. Among the 64 genes/proteins, 57 genes

showed negative Log2FC based on Tn-seq data, and 41 proteins among the 57 were upregulated

at protein level. However, only 12 proteins had p values of ≤ 0.05 (AhpC, ArcA, Crr, DksA,

FliC, IcdA, OxyR, Pgm, RecA, RpoS, SlpA, and WecE).

Using KEGG pathway analysis, 150 proteins among the 246 were enriched in 21 pathways

(Table S6). Interestingly, of the all 59 30S and 50S ribosomal proteins in *S*. Typhimurium, 37 of

these proteins (63%) were downregulated in response to H_2O_2. Moreover, of the 8 identified

proteins in TCA cycle, 6 proteins were downregulated, including 2 essential proteins.

Although DDA method can be used to search for all proteins in a complex sample, it is prone to

miss identification of important proteins due to the fact that fragmentation of tryptic peptides

from these proteins may not be triggered as a result of lower peptide ion intensities compared to

the threshold set. To quantify proteins expressed for the genes identified by Tn-seq more

precisely and accurately, we used targeted-proteomic approach by employing liquid

chromatography coupled with triple quadrupole mass spectrometry (LC-QQQ-ESI-MS). Here,

tryptic peptides of the protein were targeted for fragmentation (MS/MS) independent of their

intensities, as described in Materials and Methods, and the observed sequence specific fragment

ion intensities from three unique tryptic peptides were utilized for protein quantitation. Of the

137 Tn-seq identified genes, we selected 33 genes to quantify their proteins in response to H_2O_2

by using targeted proteomics (Table S5). Interestingly, 23 (70%) of the 33 tested proteins were

upregulated in response to H_2O_2. This shows a good agreement between the results of the Tn-seq

and the targeted proteomics.

Aromatic amino acid biosynthesis genes are required for H_2O_2 resistance
331
Interestingly, our Tn-seq data revealed that the aromatic amino acid biosynthesis and metabolism

pathway play a role in conferring resistance in *Salmonella* to H_2O_2 (Fig. 3A and 3B). Five genes,

aroB, *aroD*, *aroE_2*, *aroK*, and *aroA* in the aromatic amino acid biosynthesis pathway were

identified by Tn-seq, and the fitness of the mutants were significantly reduced in the presence of

H_2O_2. To confirm this, 4 of these genes were evaluated using individual mutant assays. The

Salmonella aroK mutant showed the strongest phenotype, because it failed to grow in the

presence of H_2O_2L or H_2O_2H during 24 h incubation time. Also, the *aroB* mutant exhibited a

strong phenotype, significantly extending lag phase for both H_2O_2 conditions. The *aroE_2*

mutant also exhibited an extended lag time, but the *aroA* mutant did not show any difference in

growth phenotype in the presence of H_2O_2. In addition, targeted-proteomics also showed that all

these 5 proteins were upregulated in response to H_2O_2 (Fig. 3C and Table S5).

ROS damages a variety of biomolecules via Fenton reaction, which consequently lead to

metabolic defects, specifically auxotrophy for some aromatic amino acids 10. *E. coli* mutants

that lack superoxide dismutase enzymes are unable to grow *in vitro* unless the medium are

supplemented with aromatic (Phe, Trp, Tyr), branched-chain (Ile, Leu, Val), and sulfur-

containing (Cys, Met) amino acids[36]. We identified the genes in the aromatic amino acid

biosynthesis pathway that are critically important for resistance to H_2O_2. In this pathway, *aroK*

catalyzes the production of shikimate 3-phosphate from shikimate, which consequently leads to

the production of tryptophan, phenylalanine, tyrosine and some metabolites from the chorismate

precursor in *E. coli*. Further, *aroK* mutant in *E*. coli displays increased susceptibility to

protamine, a model cationic antimicrobial peptide. It has been suggested that resistance to

protamine is probably due to the aromatic metabolites and product of *aroK* gene, which act as a

signal molecule to simulate the CpxR/CpxA system and Mar regulators[37]. In our Tn-seq data,

cpxR/*cpxA* and *marBCRT* were in the list of non-required genes, but the proteomics data

indicated that CpxR was upregulated. Also, *aroK* mutant in *E. coli* is resistance to mecillinam, a

beta-lactam antibiotic specific to penicillin-binding protein 2. It has been concluded that the

AroK has a secondary activity in addition to the aromatic amino acid biosynthesis, probably

related to cell division[38]. In addition, *aroK* gene presents a promising target to develop a non-

toxic drug in *Mycobacterium tuberculosis* because *aroK* is the only *in vitro* essential gene among

the aromatic amino acid pathway genes and blocking *aroK* kills the bacterium *in vivo*[39].

Moreover, *aroK* gene plays a general role in *S.* Typhimurium persistence in pigs[40]. The *aroB* is

another gene in the pathway that was identified by Tn-seq, which encodes 3-dehydroquinate

synthase in the Shikimate pathway, aromatic amino acid biosynthesis pathway. *Salmonella*

lacking *aroB* showed a strong growth defect in the presence of H_2O_2. When this mutant grown in

the presence of H_2O_2, it increased the lag phase time by a 114-fold for the H_2O_2L and a 347-fold

for the H_2O_2H as compared to the mutant grown in absence of H_2O_2. *S.* Typhimurium mutant

lacking the *aroB* gene is attenuated in BALB/c mice[41]. In addition to *aroK* and *aroB*, *aroE_2*

was also shown to be important for resistance to H_2O_2, because deletion of the *aroE_2* reduced

the growth rate by 35% in the presence of H_2O_2 and increased the lag phase time, too. All these 3

genes in this pathway are required for systemic infection of *Salmonella* in BALB/c mice in a

more recent study[18]. We observed that there was a strong correlation between the fitness based

on Tn-seq data, growth rates measured by individual mutant assays, and upregulation of their

proteins quantified via targeted proteomics. This demonstrates the power of proteogenomic

approach in discovering and characterizing the genes that are required for growth under a

specific condition.

The *ybbM*, *ybbK*, and *ybbL* have a role in H_2O_2 resistance

The mutants with single deletion in each of *ybbK*, *ybbL*, and *ybbM* genes on the same pathway

showed a strong phenotype against the activity of H_2O_2 in a dose-dependent manner. Based on

Tn-seq data, the fitness of *ybbM* was -1.16 and -1.79 for H_2O_2L and H_2O_2H, respectively (Fig.

4A). The *ybbM* mutant demonstrated decreased growth rate by 38% for H_2O_2L and 100% for the

H_2O_2H as compared to the mutant grown in the absence of H_2O_2. This mutant also increased the

lag time by a 126-fold and a 267-fold for H_2O_2L and H_2O_2H, respectively (Fig. 4B). Also, the

fitness score of *ybbK* was -0.92 for H_2O_2L and -1.81 for H_2O_2H. The *ybbK* mutant showed

decrease of growth rate for H_2O_2L and H_2O_2H by 85% and 95%, respectively. The deletion

increased the lag phase by a 46-fold and a 114-fold in the presence of H_2O_2L and H_2O_2H,

respectively (Fig. 4B). Moreover, the fitness of *ybbL* mutant was -1.05 and -1.73 for H_2O_2L and

H_2O_2H, respectively. Deleting the *ybbL* in *Salmonella* led to decrease in growth rate by 27% for

H_2O_2L and 92% for H_2O_2H. The lag phase time for this mutant increased by a 22-fold and a 33-

fold for H_2O_2L and H_2O_2H, respectively. In addition, YbbM, YbbL, and YbbK proteins were

upregulated in response to H_2O_2; YbbM was the most upregulated protein among the 3 proteins

(1.46-fold), followed by YbbL (1.29-fold), and YbbK (1.25-fold) (Fig. 4C and Table S5). The

fitness scores of the Tn-seq of these 3 genes are correlated strongly with the growth rate, lag time

of their respective mutants, and upregulation of their proteins. As the number of reads depletes

after the selection for a mutant, (i) there was more reduction in growth rate, (ii) the mutant stays

longer in the lag phase, and (iii) the protein expression elevates. These observations clearly point

to their role in conferring resistance to the H_2O_2-mediated stress. These genes were described in

the *Salmonella* reference genome as follows: *ybbM*, putative YbbM family transport protein,

metal resistance protein; *ybbK*, putative inner membrane proteins; *ybbL*, putative ABC

transporter, ATP-binding protein YbbL. To the best of our knowledge, there is only one

published study on the *ybbM* and *ybbL*[42]. Based on their findings, YbbL and YbbM have a role

in iron homeostasis in *E. coli* and are important for survival when the bacterium was challenged

with 10 mM H_2O_2 for 30 min; this putative ABC transporter transports iron and lessens ROS

species formation that generates via H_2O_2. In this study, we identified an additional gene, *ybbK*,

in the same pathway as the gene required for resistance to H_2O_2, strongly establishing the role of

these 3 genes in resistance to H_2O_2.

The H_2O_2 scavenging and degrading genes
409
Salmonella employs redundant enzymes to degrade or scavenge ROS. The *katE*, *katG*, and *katN*

genes encode catalases, which are involved in H_2O_2 degradation. The *ahpCF*, *tsaA*, and *tpx* genes

encode peroxidases, which scavenge H_2O_2. The *sodA*, *sodB*, *sodCI*, and *sodCII* genes encode

superoxide dismutases and these enzymes specifically scavenge O_2[11, 12, 43-45]. However, none of

these were present in the list of genes identified by Tn-Seq. Even though *katE*, *katG*, *ahpC*,

sodA, *sodCI*, and *sodCII* showed reduced fitness, they did not meet the statistical threshold.

However, the proteomics data indicated that AhpC (1.48-fold), SodB (1.46-fold), and TpX (1.39-

fold) were upregulated in the presence of H_2O_2 (Table S5) and KatG was also upregulated, but its

p value was 0.054. This reveals that these 4 proteins were the most important enzymes for H_2O_2

resistance under our experimental conditions. *Salmonella* containing an *ahpC* promoter-gfp

fusion shows that expression of the *ahpC* is regulated by ROS that is generated from

macrophages or exogenous H_2O_2 and the response to H_2O_2 is in a dose-dependent manner[46].

Salmonella mutant that lacks *katE*, *katG*, or *ahpCF* can degrade micromolar concentrations of

H_2O_2. However, *Salmonella* mutant that has deletions in the all 5 genes, *katE*, *katG*, *katN*, *ahpCF*

and *tsaA* (HpxF), cannot degrade H_2O_2, is unable to proliferate in macrophages, and show

reduced virulence in mice[11]. This emphasizes that *ahpC*, *sodB*, and *tpx* may be the primary

players in scavenging and degrading H_2O_2 in our experiment. Why Tn-seq did not detect any of

these genes, while proteomics detected only these 3 proteins among others? It may reflect the

functional redundancy in the genetic network that prevented single deletions in one of these

genes from exhibiting fitness defect. Alternatively, when these mutants were grown together

with all other mutants in the library, the functional protein lacking in one mutant due to Tn5

insertion could have been compensated by the other mutants in the library.

In addition to these genes, *oxyR* was detected by Tn-seq (Fig. 1C) and DDA proteomics. The

oxyR was identified by all 3 analysis methods of Tn-seq data and it was on the top of the list,

indicating a severe fitness defect of the mutant. The *oxyR* gene encodes H_2O_2 sensor and

transcription factor, which mediates protection against ROS. The *katG* and *ahpCF* are regulated

by OxyR, peroxide response regulator[13, 14]. Although *Salmonella* OxyR regulon is induced in the

Salmonella-containing vacuole in macrophage, the *oxyR* mutant was virulent in a BALB/c mouse

and can grow well in human neutrophils *in vitro*[47, 48]. The fitness of *oxyR* mutant was reduced for

both H_2O_2L and H_2O_2H with the respective fitness score of -4.96 and -5.94. *Salmonella oxyR*

mutant exhibited a growth rate reduction by 24% and 40% for H_2O_2L and H_2O_2H, respectively.

Comparing this reduction in growth rate to the other mutants such as *rpoS* or *aroK*, we observed

that the *oxyR* mutant did not show severe phenotype and the mutant escaped from the lag phase

easily. Moreover, our targeted proteomics indicated that the OxyR was not upregulated

significantly. Further studies are needed to uncover the exact role of OxyR in response to ROS.

However, previous studies implied that OxyR plays an essential role in resistance to H_2O_2 by

regulating other proteins. OxyR induces Dps in *E. coli*, a ferritin-like protein that sequesters

iron[49]. Sequestering of iron impairs the Fenton reaction, which consequently provides protection

against ROS and reduces the damage of biomolecules. The *dps* gene was identified by the Tn-seq

and its fitness score was -2.48. However, the Dps protein was downregulated based on the DDA

proteomic analysis. To confirm this unexpected finding, we conducted the proteomic assay twice

and each time with at least 4 technical replicates, but the Dps protein was significantly

downregulated with $p = 0.001$. Further, the targeted-proteomics demonstrated the same result,

pointing to the downregulation of Dps in response to H_2O_2. This is contrary to the previously

reported works on Dps in *Salmonella* and the reason for the discrepancy is unclear.

DNA repair system is important for H_2O_2 resistance

The imposed exogenous H_2O_2 activates DNA repair system in *Salmonella* in order to repair or

eliminate the damage that occurred on the nucleotides. The *E. coli* RecA protein repairs double-

strand DNA lesions through recombination[50]. In our Tn-seq analysis, the fitness score of this

mutant was -5.36 for both concentrations, and in proteomics, the RecA was upregulated (1.79-

fold). *Salmonella recA* mutant decreased the maximum OD_{600} by 16% for H_2O_2L and 22% for

H_2O_2H as compared to the same mutant grown in LB. *Salmonella recA* mutant was also sensitive

to exogenous H_2O_2 in aerated rich medium[26]. Moreover, *recG*, recombination and DNA repair

gene[51], showed a stronger phenotype than *recA* mutant. The *regG* deletion in *Salmonella* caused

the growth rate reduction by 52% for H_2O_2L and 60% for H_2O_2H. This disruption in *recG* also

caused the cells to stay in lag phase for a longer time in the presence of H_2O_2 as compared to LB;

the lag time increased by a 62-fold and a 159-fold for H_2O_2L and H_2O_2H, respectively. In the

blood of patients with *Salmonella* Typhi bacteremia, the proteins encoded by *recA*, *recG*, and

xthA genes were detected, suggesting these proteins are actively expressed in the blood

environment[52]. The XthA protein is another enzyme that participates in DNA repair mechanism

induced by H_2O_2 and iron-mediated Fenton reaction. The *xthA* encodes exonuclease III, which

repairs the damaged DNA. We found that the *xthA* gene was required based on the Tn-seq assay

and its mutant had a reduced fitness score of -3.06 for H_2O_2L and a -4.38 for the H_2O_2H. Further,

Salmonella lacking the *xthA* increased the lag time by 8-fold and a 12-fold for H_2O_2L and

H_2O_2H, respectively. Targeted-proteomics showed upregulation of XthA (1.64-fold) in response

to H_2O_2. *Salmonella* enterica serovar Enteritidis defective in *xthA* is susceptible to egg albumin[53].

E. coli xthA mutant is hypersensitive to H_2O_2[54]. The *xthA* is also required for *Mycobacterium*

tuberculosis to infect C57BL/6J mice[55]. In addition to the aforementioned genes involved in

DNA repair system, *uvrA* encoding Holliday junction DNA helicase motor protein, *uvrC*

encoding exonuclease ABC subunit A, *uvrD* encoding DNA-dependent helicase II, and *polA*

encoding DNA polymerase I were among top of the list of the genes identified by Tn-seq as

required for resistance to H_2O_2. Collectively, DNA repair system is crucial for the survival of the

Salmonella in a niche that contains H_2O_2.

Flagellar genes have a role for H_2O_2 resistance
485
Some flagellar genes, *fliC* and *fliB*, were shown to be important for resistance to H_2O_2. These

two genes were identified by Tn-seq and their proteins were shown to be upregulated.

Salmonella lacking either of these genes exhibited a strong phenotype in the presence of H_2O_2.

During 24 h of incubation, *fliC* and *fliB* mutants could not grow in both H_2O_2 conditions.

However, growth of *fliD* mutant was not affected by H_2O_2. The *fliC* was shown to have a role in

Salmonella Typhi interaction with human macrophages and *Salmonella* Typhimurium *fliB*

mutant was defective in swarming motility[56, 57]. Currently it remains unclear how flagella genes

can be involved in the resistance of *Salmonella* to oxidative stress, which warrants future study

into this direction.

Fe-S cluster biogenesis system is required for H_2O_2 resistance

496
Salmonella requires the genes from Fe-S cluster biogenesis system in order to resist H_2O_2. Our

Tn-seq analysis identified 5 genes in this system that are required for the resistance. In *isc* operon

(Fe-S cluster), *iscA*, *hscB*, and *hscA* were among the genes required to resist H_2O_2. Particularly,

the *hscA* is on the top of the gene list identified by Tn-seq. In *E. coli*, this operon is regulated by

iscR, iron sulfur cluster regulator[58]; in *Salmonella* the gene *iscR* encoding this transcription

regulator is named *yfhP*. The HscB and HscA chaperones are believed to be involved in the

maturation of Fe–S proteins[59, 60]. The second operon that is involved in Fe-S protein biogenesis

is the *suf*, sulfur mobilization operon. Tn-seq found that two genes in this operon were required

for *Salmonella* to resist H_2O_2; *sufS* and *sufC*. *Salmonella* mutant lacking *sufS* exhibited a strong

phenotype in the presence of H_2O_2 and could not grow during the 24 h of incubation as compared

to LB. The SufS with SufE in *E. coli* form a heterodimeric cysteine desulphurase and SufB,

SufC, and SufD form a pseudo-ABC-transporter that could act as a scaffold[60]; this operon is

regulated by OxyR[14]. The other known genes in these two operons that are present in *Salmonella*

are *iscA*, *sufA*, *sufB*, and *sufD*; they showed a reduced fitness, while their *p* values were > 0.05.

The damage of Fe-S clusters is not only problem for the defective proteins, but also it fuels the

Fenton reaction via the released iron and H_2O_2[10]. Thus, *Salmonella* uses Fe-S cluster repair

system as an arsenal to overcome the damage imposed by H_2O_2.

DNA adenine methylase is important for H_2O_2 resistance

515
DNA adenine methylase genes, *dam* and *damX*, are involved in *Salmonella* resistance against

H_2O_2. Our Tn-seq data showed that fitness of *dam* and *damX* mutants were reduced in the

presence of H_2O_2. To confirm this, *Salmonella dam* mutant was grown in both conditions. Under

H_2O_2L, the growth rate was reduced by 42% as compared to the mutant in LB and the mutant

could not grow under H_2O_2H during the 24 h of incubation. In addition, the lag time of the

Salmonella dam mutant was extended by a 19-fold for H_2O_2L. While the *Salmonella damX*

mutant displayed a moderate phenotype as compared to the *dam* mutant, the *damX* mutant also

showed that the growth rate decreased by 23% and 33% for H_2O_2L and H_2O_2H, respectively. The

lag time was extended for this mutant by a 25-fold and a 49-fold for H_2O_2L and H_2O_2H,

respectively. The *dam* regulates virulence gene expression in *S*. Typhimurium[61]. The different

levels of Dam affects virulence gene expression, motility, flagellar synthesis, and bile resistance

in the pathogenic *S*. Typhimurium 14028S[62]. Dam methylation activates the gene that are

involved in lipopolysaccharide synthesis[63]. Moreover, *Salmonella* defective in *damX* is very

sensitive to bile[64]. Collectively, our study demonstrates the critical role of DNA adenine

methylase in *Salmonella* resistance against H_2O_2.

Other genes for H_2O_2 resistance

Beside the important pathways described above, there were many additional genes also

important for resistance to H_2O_2. Among those, the 3 unrelated genes, *rpoS*, *pgm*, and *tonB*, are

important ones that deserve more attention. The *rpoS* mutant showed reduced fitness and its

protein was upregulated in the presence of H_2O_2. *Salmonella* mutant defective in *rpoS* showed a

strong phenotype when grown in the presence of H_2O_2. The *rpoS* encodes the alternative sigma

factor σ^S, subunit of RNA polymerase; it is the master regulator of stress response[65]. This implies

that *rpoS* is an important component of the genetic regulatory network that *Salmonella* employs

in order to resist H_2O_2. Furthermore, the fitness of *pgm* mutant was reduced and its protein was

upregulated in the presence of H_2O_2. Knock out of *pgm* in *Salmonella* caused a decrease in

growth rate, increase the lag phase time, and reduce the maximum OD_{600} in the presence of

H_2O_2. The *pgm* encodes phosphoglucomutase which required for catalysis of the interconversion

of glucose 1-phosphate and glucose 6-phosphate[66]. This gene contributes to resistance against

antimicrobial peptides, is required for *in vivo* fitness in the mouse model, and participates in LPS

biosynthesis[67]. Lastly, the fitness of *tonB* mutant was also reduced. *Salmonella* lacking *tonB*

exhibited a strong phenotype in the presence of H_2O_2 as compared to the mutant grown in LB.

The gene mediates iron uptake in the *Salmonella*[45]. In addition, seven of the genes identified in

our study (*proC*, *arcA*, *barA*, *exbD*, *flhD*, *fliC*, and *fliD*) were previously shown to be important

for interaction of *Salmonella* Typhi with human macrophages[56].

In summary, we applied Tn-seq and proteomic analysis to find the genes and proteins that are

required in *S*. Typhimurium to resist H_2O_2 *in vitro*. As the concentration of H_2O_2 increased, the

growth rate reduced, the lag time extended, the fitness of mutants decreased, and some proteins

were differentially expressed. Validation of Tn-seq results with individual mutant assays

indicated the accuracy of the identified genes in response to the two H_2O_2 concentrations. The

targeted-proteomics had a good agreement with Tn-seq. We found many genes that have not

been associated to resistance to H_2O_2 previously and these genes will be focus of our future

research. *Salmonella* employs multiple pathways to resist H_2O_2 and the most important ones are

ROS detoxifying enzymes, amino acid biosynthesis (*aroK* and *aroB*), putative iron transporters

(*ybbK*, *ybbL*, *ybbM*), iron homeostasis, Fe-S cluster repair, DNA repair, flagellar and DNA

adenine methylase genes. The genes identified in this study will broaden our understanding on

the mechanisms used by *Salmonella* to survive and persist against ROS in macrophages.

Our unbiased system-wide approach, Tn-seq, was successful in identifying novel genetic

determinants that have not been implicated previously in *Salmonella* resistance to oxidative

stress. Furthermore, the combined use of quantitative proteomic approach has provided

additional insights on the function or mode of action of the identified genetic determinants in

resisting oxidative stress. As expected, the majority of the proteins important for resistance to

H_2O_2 were upregulated in response to the same stressor. However, the expression level did not

increase for some proteins, in spite of their known roles in resistance to H_2O_2. Interestingly, the

downregulation of Dps and other proteins was counterintuitive to the common mode of protein

regulation and function, yet it may point to some unknown aspects of how *Salmonella* regulates

the expression of those proteins to better cope with the oxidative stress during infection in

macrophage. The genes identified in this study will broaden our understanding on the

mechanisms used by *Salmonella* to survive and persist against ROS in macrophages.

Methods

Construction of Tn5 mutant library

We mutagenized *Salmonella* enterica subsp. enterica serovar Typhimurium str. ATCC 14028S

(with spontaneous mutation conferring resistance to nalidixic acid (NA)), by biparental mating

using *Escherichia coli* SM10 λ*pir* carrying a transposon-delivery plasmid vector pBAM1

(Ampicillin (Amp) resistance) as a donor strain[68]. The plasmid pBAM1 was generously provided

by Victor de Lorenzo (Systems and Synthetic Biology Program, Centro Nacional de

Biotecnología, Madrid, Spain). The donor strain*, E. coli* Sm10 λpir (pBAM1), was grown

overnight in LB with 50 µg/ml Amp and recipient strain was grown in LB with 50 µg/ml NA at

37°C. Equal volumes (1 ml) of donor and recipient were mixed, centrifuged, washed in 10 mM

MgSO₄, and re-suspended in 2 ml PBS (pH 7.4). Then, the mating mixture was concentrated and

laid on a 0.45-µm nitrocellulose filters (Millipore). The filter was incubated at 37°C on the

surface of LB agar plate. After 5 h of conjugation, the cells on the filter was washed in 10 mM

MgSO$_4$, and resuspended in 1 ml MgSO4. The conjugation mixture was plated on LB agar

containing 50 µg/ml NA and 50 µg/ml kanamycin (Km). After approximately 24 h at 37°C, we

scraped the colonies into LB broth containing 50 µg/ml Km and 7% DMSO. The yield was

approximately 68,000 individual colonies from each conjugation. Five independent conjugations

were conducted to yield approximately 325,000 mutants. The library was stored at −80°C in

aliquots. To determine the frequency of the mutants that have been produced by integration of

the entire delivery plasmid, the colonies were picked randomly and streaked on LB plates (Km)

and LB plates (Km and Amp). It was shown that ~20% of the cells in the library were resistant to

Amp, indicating a significant portion of the Km-resistant colonies was not from authentic

transposition events.

Measuring growth responses of *S*. Typhimurium to H$_2$O$_2$
601
To determine the effect of H$_2$O$_2$ concentrations on growth parameters, overnight culture of the

wild type *S*. Typhimurium 14028s was inoculated into fresh LB broth media (1:200 dilution) to

give a seeding concentration corresponding to OD$_{600}$ of ~0.1. The LB broth contained freshly

prepared H$_2$O$_2$ to give the final concentrations ranging from 0.05 to 10 mM. The cultures were

directly added into 96-well microplate (200 µl/well). The microplate was incubated in a Tecan

Infinite 200 microplate reader at 37°C, with shaking duration 5 s, shaking amplitude 1.5 mm, and

reading OD$_{600}$ every 10 min. The number of replicates were at least three. The lag time, growth

rate, and maximum OD$_{600}$ were calculated using GrowthRates script[69]. Growth Rate % decrease

was calculated as follows[70]: Growth Rate % decrease = $((\mu_{PC} - \mu_S)/\mu_{PC})$ x 100; where μ = the

maximum slope (growth rate), μ_{PC} = growth rate of positive control (without H$_2$O$_2$), μ_S = growth

rate in the presence of H$_2$O$_2$.

Selection of the mutant library for Tn-seq analysis
614

The transposon library was thawed at room temperature and diluted 10^{-1} in fresh LB broth. To

activate the library, the diluted library was incubated at 37°C with shaking at 225 rpm for an

hour. Then, the culture was washed twice with PBS and resuspended in LB broth medium. The

library was inoculated to 20 ml LB broth and LB broth supplemented with either 2.5 or 3.5 mM

H_2O_2 (H_2O_2L and H_2O_2H, respectively), seeding CFU was 3.5 x 10^6 per ml. Then, when the

cultures reached mid-exponential phase, OD_{600} of 2.7 (~1.17 x 10^8 CFU/ml), the incubation was

stopped, and the culture was immediately harvested by centrifugation, and stored at -20°C.

Preparation of Tn-seq amplicon libraries

Genomic DNA was extracted from the harvested cells using DNeasy Blood & Tissue kit

(Qiagen), and quantified using Qubit dsDNA RB Assay kit (Invitrogen). As described above,

20% of the mutants in the library were the result of the integration of pBAM1 into chromosome.

To remove the Tn5-junction sequences originated from the plasmid in the Tn-seq amplicon

libraries, genomic DNA was digested with PvuII-HF (New England Biolabs), which digests

immediately outside the inverted repeats on both sides of Tn5 in pBAM1, and purified with

DNA Clean & Concentrator-5 kit (Zymo Reaerch). Then, a linear PCR extension was performed

using a Tn5-specific primer in order to produce single stranded DNA corresponding to Tn5-

junction sequences. To increase the specificity in extending into Tn5-junction sequences, the

linear PCR was conducted with a dual priming oligonucleotide Tn5-DPO (5'-

AAGCTTGCATGCCTGCAGGTIIIIICTAGAGGATC-3') that is specific to Tn5 end[71]. The PCR

reaction contained 25 μl Go Taq Colorless Master Mix (Promega), 20 μM Tn5-DPO primer, 100

ng gDNA, and 50 μl MQ-H_2O. The PCR cycle consisted of the initial denaturation at 95°C for 2

min, followed by 50 cycles at 95°C for 30 sec, 62°C for 45 sec, and 72°C for 10 sec. The PCR

product was purified with DNA Clean & Concentrator-5 kit and eluted in 13 μl TE buffer. After

that, C-tail was added to the 3' end of the single-stranded DNA. The C-tailing reaction was

consisted of 2 μl terminal transferase (TdT) buffer (New England Biolabs), 2 μl $CoCl_2$, 2.4 μl 10

mM dCTP, 1 μl 1 mM ddCTP, 0.5 μl TdT and 13 μl purified linear PCR product. The reaction

was performed at 37°C for 1 h and the enzyme was inactivated by incubation at 70°C for 10 min.

The C-tailed product was purified with DNA Clean & Concentrator-5 kit and eluted in 12 μl TE.

Next, the exponential PCR was performed with forward primer, P5-BRX-TN5-MEO, 5'-

AATGATACGGCGACCACCGAGATCTACACTCTTTCCCTACACGACGCTCTTCCGATC

TNNNNAG-6 nt barcode-CCTAGGCGGCCTTAATTAAAGATGTGTATAAGAG-3' and

reverse primer, P7-16G, 5'-

CAAGCAGAAGACGGCATACGAGCTCTTCCGATCTGGGGGGGGGGGGGGGGG-3' to

attach Illumina adapter sequences along with the sample barcodes. The PCR reaction contained

25 μl Go Taq Green Master Mix, 10 μM P5-BRX-TN5-MEO primer, 10 μM P7-16G primer, 1

μl purified C-tailed genomic junctions, and MQ-H_2O to 50 μl; the PCR condition started with

initial denaturation at 95°C for 2 min, followed by 36 cycles of 95°C for 30 sec, 60°C for 30 sec,

and 72°C for 20 sec, with the final extension at 72°C for 5 min. Then, the size selection of the

DNA was performed using agarose gel electrophoresis. The 50 μl PCR products were incubated

at 60°C for 15 min and incubated on ice for 5 min, and immediately loaded on the 1% agarose

gel in 0.5% TAE buffer. After running the gel, the DNA fragment of size 325-625 bp was cut

and put in a microtube for each sample. The DNA was extracted from the gel using Zymoclean

Gel DNA Recovery kit (Zymo Reaerch). The prepared DNA libraries were quantified using

Qubit dsDNA RB Assay kit. Since each library has its own barcode, the libraries were combined

and sequenced on a flow cell of HiSeq 3000 using single end read and 151 cycles (Illumina) at

the Center for Genome Research & Biocomputing in Oregon State University.

Tn-seq data analysis
663
The preliminary data analysis was conducted by using a super computer in the High Performance

Computing Center (AHPCC) at the University of Arkansas. The libraries that were multiplexed

for sequencing were de-multiplexed using a custom Python script. The script searched for the

six-nucleotide barcode for each library for perfect matches. In order to extract the transposon

genomic junctions, we used Tn-Seq Pre-Processor (TPP) tool[30] with some modifications in the

script. The TPP searched for the 19 nucleotide inverted repeat (IR) sequence and identified five

nucleotides (GACAG) at the end of the IR sequence, allowing one nucleotide mismatch. The

Tn5-junctions that start immediately after GACAG were extracted and the C-tails at the end of

junctions were removed. Tn5-junction sequences less than 20 nucleotides were discarded and

remaining Tn5-junctions were mapped to the *Salmonella* enterica serovar Typhimurium 14028S

genome and plasmid using BWA-0.7.12[72]. To identify genes that are required for H_2O_2

resistance, the following three Tn-seq analysis tools were used for comparative analysis: (i)

ARTIST[28]: the genomic junctions were mapped to the reference genome using Bowtie 2.2.7[73].

The number of insertions and reads were determined for genes and intergenic regions. The data

were normalized with default script in the ARTIST. Then, the relative abundance of Tn5

insertions in the output library versus the input were calculated. Later, the *p* values were

calculated from a 100 independent Mann–Whitney U test (MWU) analysis that were carried out

on input and output data for each gene. Finally, the genes were considered conditionally essential

if the *p* values were ≤ 0.05 in the 90 of the 100 MWU tests. (ii) Tn-seq Explorer[29]: The output

SAM files from the TTP were used as input to the Tn-seq Explorer. The unique insertions with

less than 20 reads were removed from the input and outputs. Using the window size of 550 and

excluding 5% of beginning of genes and 20% of the end of genes, Essentiality Index (EI),

number of unique insertions, and total number of reads per gene were counted. The EI of more

than 10 were removed from the input. Genes with less than 300 nucleotides were removed.

Deferential EI were calculated from input and outputs (ΔIE = output EI–input EI) and genes with

ΔIE more than -1 were removed. Log2 fold change of reads were calculated from input and

output (Log2FC = log2 (output reads/input reads)) and the genes were ranked based on the

Log2FC value from least value to highest. The genes with Log2FC value of less than -2 from the

H_2O_2H and present in H_2O_2L were considered conditionally essential. (iii) TRANSIT[30]: The

output wig files from the TTP was used as input data file for TRANSIT. The comparative

analysis was conducted with Tn5 resampling option. The reads were normalized with trimmed

total reads (TTR). Insertions outside the 5% and 10% sequences from 5'- and 3'- ends were

removed, respectively. The genes were considered conditionally essential if p values ≤ 0.05.

Phenotypic evaluation of individual deletion mutants
698
The mutants were obtained from *Salmonella enterica* subsp. *enterica*, 14028s (Serovar

Typhimurium) Single-Gene Deletion Mutant Library through BEI Resources

(www.beiresources.org). The overnight cultures of *S.* Typhimurium mutants were added into

fresh LB broth media containing freshly prepared H_2O_2 (2.5, or 3.5 mM/ml) (1:200 dilution) to

give seeding OD_{600} of 0.1. The cultures were directly added into 96-well microplates and

incubated in Tecan Infinite 200 at 37ºC for 24 h. The lag time, growth rate, and maximum OD_{600}

were calculated using GrowthRates[69].

Sample preparation for proteomics and mass spectrometry analysis
707
The overnight culture of the wild type *S.* Typhimurium 14028 was diluted 1:200 in 50 ml LB

medium, and LB containing either 2.5 or 3.5 mM H_2O_2 in a 300-ml flask. The cultures were

grown until mid-exponential phase (OD_{600} of 2.7), and the 50 ml volume of cultures were used

for a total protein extraction by using Qproteome Bacterial Protein Prep kit (Qiagen). Disulfide

bonds within proteins were reduced with 2-Mercaptoethanol and separated by SDS-PAGE gel

electrophoresis. For each condition, there were three lanes with approximately 300 μg of

proteins. The gel then was stained via coomassie blue dye. The gel portions of 3 lanes for each

condition were cut out and chopped into small pieces, pooled together, washed twice with 50

mM NH_4HCO_3, destained with NH_4HCO_3/ 50% Acetonitrile (ACN), and dried with pure ACN.

Then, the proteins were reduced using 10 mM Dithiothreitol in 50 mM NH_4CO_3 and the

alkylation was conducted with 10 mg/ml Iodoacetamide Acid in 50 mM NH_4CO_3. After that, the

proteins were washed with NH_4HCO_3, and dried with pure ACN. Mass spectrometry grade

Trypsin gold from Promega (~20 ng/μl in 50 mM NH_4HCO_3) was added to dried gels, and left it

overnight for efficient in-gel digestion of the proteins at 37ºC. During the digestion, tryptic

peptides diffused out into the solution. Gel pieces then were extracted three times by 50%

CAN/0.1% TFA solution and incubated at 37°C for 15 minutes. Later, these digests were

analyzed by ESI-LC-MS/MS at State Wide Mass Spectrometry Facility, University of Arkansas

at Fayetteville. Data dependent analysis (DDA) for the in-gel trypsin digested samples from each

condition were performed by using an Agilent 1200 series micro flow HPLC in line with Bruker

Amazon-SL quadrupole ion trap ESI mass spectrometer (QIT-ESI-MS). All the ESI-MS analyses

were performed in a positive ion mode using Bruker captive electrospray source with a dry

nitrogen gas temperature of 200°C, with nitrogen flow rate of 3 L/minute. LC-MS/MS data were

acquired in the Auto MS(n) mode with optimized trapping condition for the ions at m/z 1000.

MS scans were performed in the enhanced scanning mode (8100 m/z/second), while the

collision-induced dissociation or the MS/MS fragmentation scans were performed automatically

for top ten precursor ions with a set threshold for one minute in the UltraScan mode (32,500

m/z/second). Tryptic peptides were separated by reverse-phase high-performance liquid

chromatography (RP-HPLC) using a Zorbax SB C18 column, (150 × 0.3 mm, 3.5 µm particle

size, 300 Å pore size, Agilent Technologies), with a solvent flow rate of 4 µL/minute, and a

gradient of 5%–38% consisting of 0.1% FA (solvent A) and ACN (solvent B) over a time period

of 320 minutes. Tryptic peptides were then identified by searching MS/MS data in *S.*

Typhimurium 14028S reference proteome database[24, 35] by using MASCOT database search

software[34]. MS1 intensities of the integrated areas of these identified tryptic peptides were

compiled and grouped in skyline software according to the replicates/conditions to perform

statistical analysis. Targeted protein work were performed using Shimadzu UPLC-20A coupled

to 8050 triple quadrupole ESI-MS with heated probe. Sequence specific fragment ion intensities

from at least three unique tryptic peptides from the protein of interest were used in the protein

quantitation. Multiple reaction monitoring (MRM) events corresponding to sequence specific

fragment ions derived from the precursor tryptic peptides were targeted to operate at a certain

specific retention time intervals predicted by in house retention time library. This library was

generated using the correlation of relative hydrophobicity of the tryptic peptides with their

retention times (RT) from highly common housekeeping proteins, for the UPLC method used in

this analysis as described below. While the RT were correlated well within 99% confidence,

sufficient number of sequence specific fragment ions were used as basis for identification of the

tryptic peptide by MS/MS alone. Specificity and the confidence was achieved by incorporating

RT prediction. In addition to the application of skyline in quantitation, skyline software was also

used in predicting RT and optimizing parameters such as collision energies and voltages with the

help of Shimadzu Labsolution software. Tryptic peptides were separated by reverse-phase ultra-

high-performance liquid chromatography (RP-UPLC) compatible Shimadzu C18, 1.9-micron

particle size, 50x2.1 mm column (SN # 16041880T), with a solvent flow rate of 0.3 mL/minute,

and a gradient of 5%–90% consisting of 0.1% FA (solvent A) and 0.1% FA in ACN (solvent B)

over a time period of 10 minutes.

Accession numbers

Tn-seq sequencing data are available on NCBI Sequence Read Archive. LB: PRJNA352537;

H_2O_2L: PRJNA352862; H_2O_2H: PRJNA352865.

References

1. LaRock, D. L., Chaudhary, A. & Miller, S. I. Salmonellae interactions with host

processes. *Nat. Rev. Microbiol.* **13**, 191-205 (2015).

2. Scallan, E. *et al*. Foodborne illness acquired in the United States – major

pathogens. *Emerg. Infect. Dis.***17** (2011).

3. Majowicz, S. E. *et al*. The global burden of nontyphoidal *Salmonella* gastroenteritis. *Clin.*

Infect. Dis. **50**, 882-889 (2010).

4. Haraga, A., Ohlson, M. B. & Miller, S. I. Salmonellae interplay with host cells. *Nat. Rev.*

Microbiol. **6**, 53-66 (2008).

5. Nauseef, W. Assembly of the phagocyte NADPH oxidase. *Histochem. Cell Biol.* **122**,

277-291 (2004).

6. Seaver, L. C. & Imlay, J. A. Hydrogen peroxide fluxes and compartmentalization inside

growing *Escherichia coli*. *J. Bacteriol.* **183**, 7182-7189 (2001).

7. Park, S., You, X. & Imlay, J. A. Substantial DNA damage from submicromolar

intracellular hydrogen peroxide detected in Hpx- mutants of *Escherichia coli*. *Proc. Natl.*

Acad. Sci. U. S. A. **102**, 9317-9322 (2005).

8. Imlay, J. A. & Linn, S. DNA damage and oxygen radical toxicity. *Science* **240**, 1302-
1309 (1988).

9. Imlay, J. A. Iron-sulphur clusters and the problem with oxygen. *Mol. Microbiol.* **59**,
1073-1082 (2006).

10. Slauch, J. M. How does the oxidative burst of macrophages kill bacteria? Still an open
question. *Mol. Microbiol.* **80**, 580-583 (2011).

11. Hébrard, M., Julie P. M. Viala, Méresse, S., Barras, F. & Aussel, L. Redundant Hydrogen
Peroxide Scavengers Contribute to *Salmonella* Virulence and Oxidative Stress
Resistance. *J. Bacteriol.* **191**, 4605-4614 (2009).

12. Horst, S. A. *et al*. Thiol Peroxidase Protects *Salmonella* enterica from Hydrogen Peroxide
Stress *In Vitro* and Facilitates Intracellular Growth. *J. Bacteriol.* **192**, 2929-2932 (2010).

13. Morgan, R. W., Christman, M. F., Jacobson, F. S., Storz, G. & Ames, B. N. Hydrogen
Peroxide-Inducible Proteins in *Salmonella* Typhimurium Overlap with Heat Shock and
Other Stress Proteins. *Proc. Natl. Acad. Sci. U. S. A.* **83**, 8059-8063 (1986).

14. Zheng, M. *et al*. DNA Microarray-Mediated Transcriptional Profiling of the *Escherichia*
coli Response to Hydrogen Peroxide. *J. Bacteriol.* **183**, 4562-4570 (2001).

15. Kroger, C. *et al*. An Infection-Relevant Transcriptomic Compendium for *Salmonella*
enterica Serovar Typhimurium. *Cell Host Microbe* **14**, 683-695 (2013).

16. Turner, K. H., Wessel, A. K., Palmer, G. C., Murray, J. L. & Whiteley, M. Essential
genome of Pseudomonas aeruginosa in cystic fibrosis sputum. *Proc. Natl. Acad. Sci. U.*
S. A. **112**, 4110 (2015).

17. Chan, K., Kim, C. C. & Falkow, S. Microarray-Based Detection of *Salmonella* enterica

Serovar Typhimurium Transposon Mutants That Cannot Survive in Macrophages and

Mice. *Infect. Immun.* **73**, 5438-5449 (2005).

18. Silva-Valenzuela, C. *et al*. Analysis of Two Complementary Single-Gene Deletion

Mutant Libraries of *Salmonella* Typhimurium in Intraperitoneal Infection of BALB/c

Mice. *Front. Microbiol.* **6** (2016).

19. van Opijnen, T., Bodi, K. L., & Camilli, A. Tn-seq: high-throughput parallel sequencing

for fitness and genetic interaction studies in microorganisms. *Nat. Methods* **6**, 767-772

(2009).

20. Khatiwara, A. *et al*. Genome Scanning for Conditionally Essential Genes in *Salmonella*

enterica Serotype Typhimurium. *Appl. Environ. Microbiol.* **78**, 3098-3107 (2012).

21. Chaudhuri, R. *et al*. Comprehensive Assignment of Roles for *Salmonella* Typhimurium

Genes in Intestinal Colonization of Food-Producing Animals. *PLoS Genet.* **9**, e1003456

(2013).

22. Langridge, G. C. *et al*. Simultaneous assay of every *Salmonella* Typhi gene using one

million transposon mutants. *Genome Res.* **19**, 2308-2316 (2009).

23. Barquist, L. *et al*. A comparison of dense transposon insertion libraries in the *Salmonella*

serovars Typhi and Typhimurium. *Nucleic Acids Res.* **41**, 4549-4564 (2013).

24. Domon, B. & Aebersold, R. Options and considerations when selecting a quantitative

proteomics strategy. *Nat. Biotechnol.* **28**, 710-721 (2010).

25. Jensen, O. N. & Mann, M. Proteomic analysis of post-translational modifications. *Nat.*

Biotechnol. **21**, 255-261 (2003).

26. Bogomolnaya, L. *et al*. The ABC-Type Efflux Pump MacAB Protects *Salmonella*

enterica serovar Typhimurium from Oxidative Stress. *Mbio* **4**, e00630 (2013).

27. Kwon, Y. M., Ricke, S. C. & Mandal, R. K. Transposon sequencing: methods and

expanding applications. *Appl. Microbiol. Biotechnol.* **100**, 31-43 (2016).

28. Pritchard, J. R. *et al*. ARTIST: high-resolution genome-wide assessment of fitness using

transposon-insertion sequencing. *PLoS Genet.* **10**, e1004782-e1004782 (2014).

29. Solaimanpour, S., Sarmiento, F. & Mrazek, J. Tn-Seq Explorer: A Tool for Analysis of

High-Throughput Sequencing Data of Transposon Mutant Libraries. *PLoS One* **10**,

e0126070 (2015).

30. DeJesus, M. A., Ambadipudi, C., Baker, R., Sassetti, C. & Ioerger, T. R. TRANSIT--A

Software Tool for Himar1 TnSeq Analysis. *PLoS Comput. Biol.* **11**, e1004401-e1004401

(2015).

31. Huang, D. W., Sherman, B. T. & Lempicki, R. A. Bioinformatics enrichment tools: paths

toward the comprehensive functional analysis of large gene lists. *Nucleic Acids Res.* **37**,

1-13 (2009).

32. Kanehisa, M. & Goto, S. KEGG: Kyoto Encyclopedia of Genes and Genomes. *Nucleic*

Acids Res. **28**, 27-30 (2000).

33. MacLean, B. *et al*. Skyline: an open source document editor for creating and analyzing

targeted proteomics experiments. *Bioinformatics* **26**, 966-968 (2010).

34. Perkins, D., Pappin, D., Creasy, D. & Cottrell, J. Probability-based protein identification

by searching sequence databases using mass spectrometry data. *Electrophoresis* **20**,

3551-3567 (1999).

35. Cravatt, B. F., Yates III, J. R. & Simon, G. M. The biological impact of mass-
spectrometry-based proteomics. *Nature* **450**, 991-1000 (2007).

36. Carlioz, A. & Touati, D. Isolation of superoxide dismutase mutants in *Escherichia coli*: is
superoxide dismutase necessary for aerobic life? *EMBO J.* **5**, 623-630 (1986).

37. Weatherspoon-Griffin, N., Yang, D., Kong, W., Hua, Z. & Shi, Y. The CpxR/CpxA Two-
component Regulatory System Up-regulates the Multidrug Resistance Cascade to
Facilitate *Escherichia coli* Resistance to a Model Antimicrobial Peptide. *J. Biol.*
Chem. **289**, 32571-32582 (2014).

38. Vinella, D., Gagny, B., Joseleau-Petit, D., D'Ari, R. & Cashel, M. Mecillinam resistance
in *Escherichia coli* is conferred by loss of a second activity of the AroK protein. *J.*
Bacteriol. **178**, 3818-3828 (1996).

39. Kaur, P. *et al*. Unravelling the Secrets of Mycobacterial Cidality through the Lens of
Antisense. *PLoS One* **11**, e0154513 (2016).

40. Van Parys, A. *et al*. Tissue-Specific *Salmonella* Typhimurium Gene Expression during
Persistence in Pigs. *PLoS One* **6**, e24120 (2011).

41. Günel-Özcan, A., Brown, K. A., Allen, A. G. & Maskell, D. J. *Salmonella* Typhimurium
aroB mutants are attentuated in BALB/ c mice. *Microb. Pathog.* **23**, 311-316 (1997).

42. Nicolaou, S. A., Fast, A. G., Nakamaru-Ogiso, E. & Papoutsakis, E. T. Overexpression of
fetA (*ybbL*) and *fetB* (*ybbM*), Encoding an Iron Exporter, Enhances Resistance to
Oxidative Stress in Escherichia coli. *Appl. Environ. Microbiol.* **79**, 7210-7219 (2013).

43. Robbe-Saule, V., Coynault, C., Ibanez-Ruiz, M., Hermant, D. & Norel, F. Identification
of a non-haem catalase in *Salmonella* and its regulation by RpoS (σS). *Mol.*
Microbiol. **39**, 1533-1545 (2001).

44. Craig, M. & Slauch, J. Phagocytic Superoxide Specifically Damages an Extracytoplasmic

Target to Inhibit or Kill *Salmonella*. *PLoS One* **4**, e4975 (2009).

45. Tsolis, R. M., Bäumler, A. J., Heffron, F. & Stojiljkovic, I. Contribution of TonB- and

Feo-mediated iron uptake to growth of *Salmonella* Typhimurium in the mouse. *Infect.*

Immun. **64**, 4549-4556 (1996).

46. Aussel, L. *et al.* *Salmonella* detoxifying enzymes are sufficient to cope with the host

oxidative burst. *Mol. Microbiol.* **80**, 628-640 (2011).

47. Taylor, P. D., Inchley, C. J. & Gallagher, M. P. The *Salmonella* Typhimurium AhpC

Polypeptide Is Not Essential for Virulence in BALB/c Mice but Is Recognized as an

Antigen during Infection. *Infect. Immun.* **66**, 3208-3217 (1998).

48. Papp-Szabò, E., Firtel, M. & Josephy, P. D. Comparison of the sensitivities of *Salmonella*

Typhimurium *oxyR* and *katG* mutants to killing by human neutrophils. *Infect. Immun.* **62**,

2662-2668 (1994).

49. Imlay, J. Diagnosing oxidative stress in bacteria: not as easy as you might think. *Curr.*

Opin. Microbiol. **24**, 124-131 (2015).

50. Carlsson, J. & Carpenter, V. S. The recA+ gene product is more important than catalase

and superoxide dismutase in protecting *Escherichia coli* against hydrogen peroxide

toxicity. *J. Bacteriol.* **142**, 319-321 (1980).

51. Ishioka, K., Iwasaki, H. & Shinagawa, H. Roles of the *recG* gene product of *Escherichia*

coli in recombination repair: Effects of the Delta *recG* mutation on cell division and

chromosome partition. *Genes Genet. Syst.* **72**, 91-99 (1997).

52. Sheikh, A. *et al*. In Vivo Expression of *Salmonella* enterica Serotype Typhi Genes in the

Blood of Patients with Typhoid Fever in Bangladesh. *PLoS Negl. Trop. Dis.* **5**, e1419

(2011).

53. Lu, S., Killoran, P. B. & Riley, L. W. Association of *Salmonella* enterica Serovar

Enteritidis YafD with Resistance to Chicken Egg Albumen. *Infect. Immun.* **71**, 6734-

6741 (2003).

54. Demple, B., Halbrook, J. & Linn, S. *Escherichia coli xth* mutants are hypersensitive to

hydrogen peroxide. *J. Bacteriol.* **153**, 1079-1082 (1983).

55. Sassetti, C. M. & Rubin, E. J. Genetic Requirements for Mycobacterial Survival during

Infection. *Proc. Natl. Acad. Sci. U. S. A.* **100**, 12989-12994 (2003).

56. Sabbagh, S., Lepage, C., McClelland, M. & Daigle, F. Selection of *Salmonella* enterica

Serovar Typhi Genes Involved during Interaction with Human Macrophages by

Screening of a Transposon Mutant Library. *PLoS One* **7**, e36643 (2012).

57. Bogomolnaya, L. *et al*. Identification of Novel Factors Involved in Modulating Motility

of *Salmonella* enterica Serotype Typhimurium. *PLoS One* **9**, e111513 (2014).

58. Schwartz, C. J. *et al*. IscR, An Fe-S Cluster-Containing Transcription Factor, Represses

Expression of *Escherichia coli* Genes Encoding Fe-S Cluster Assembly Proteins. *Proc.*

*Natl. Acad. Sci. U. S. A.***98**, 14895-14900 (2001).

59. Unciuleac, M. *et al*. Vitro activation of apo-aconitase using a [4Fe-4S] cluster-loaded

form of the IscU [Fe-S] cluster scaffolding protein. *Biochemistry (N. Y.)* **46**, 6812-6821

(2007).

60. Py, B. & Barras, F. Building Fe–S proteins: bacterial strategies. *Nat. Rev. Microbiol.* **8**,

436-446 (2010).

61. Balbontín, R. *et al*. DNA Adenine Methylation Regulates Virulence Gene Expression in

Salmonella enterica Serovar Typhimurium. *J. Bacteriol.* **188**, 8160-8168 (2006).

62. Badie, G., Heithoff, D. M., Sinsheimer, R. L. & Mahan, M. J. Altered Levels of

Salmonella DNA Adenine Methylase Are Associated with Defects in Gene Expression,

Motility, Flagellar Synthesis, and Bile Resistance in the Pathogenic Strain 14028 but Not

in the Laboratory Strain LT2. *J. Bacteriol.* **189**, 1556-1564 (2007).

63. Sarnacki, S. H. *et al*. Dam Methylation Controls O-Antigen Chain Length in *Salmonella*

enterica Serovar Enteritidis by Regulating the Expression of Wzz Protein. *J.*

Bacteriol. **191**, 6694-6700 (2009).

64. Lopez-Garrido, J., Cheng, N., Garcia-Quintanilla, F., Garcia-del Portillo, F. & Casadesus,

J. Identification of the *Salmonella* enterica *damX* Gene Product, an Inner Membrane

Protein Involved in Bile Resistance. *J. Bacteriol.* **192**, 893-895 (2010).

65. Hengge-Aronis, R. Signal Transduction and Regulatory Mechanisms Involved in Control

of the σS (RpoS) Subunit of RNA Polymerase. *Microbiol. Mol. Biol. Rev.* **66**, 373-395

(2002).

66. Lu, M. & Kleckner, N. Molecular cloning and characterization of the *pgm* gene encoding

phosphoglucomutase of *Escherichia coli*. *J. Bacteriol.* **176**, 5847-5851 (1994).

67. Paterson, G. K., Cone, D. B., Peters, S. E. & Maskell, D. J. The enzyme

phosphoglucomutase (*Pgm*) is required by *Salmonella* enterica serovar Typhimurium for

O-antigen production, resistance to antimicrobial peptides and in vivo

fitness. *Microbiology* **155**, 3403-3410 (2009).

68. Martínez-García, E., Calles, B., Arévalo-Rodríguez, M. & de Lorenzo, V. pBAM1: an

all-synthetic genetic tool for analysis and construction of complex bacterial

phenotypes. *BMC Microbiol.* **11**, 38-38 (2011).

69. Hall, B., Acar, H., Nandipati, A. & Barlow, M. Growth Rates Made Easy. *Mol. Biol.*

Evol. **31**, 232-238 (2014).

70. Yao, Q. *et al.* Assembly of nanoions via electrostatic interactions: ion-like behavior of

charged noble metal nanoclusters. *Sci. Rep.* **4**, 3848-3848 (2014).

71. Chun, J. *et al.* Dual priming oligonucleotide system for the multiplex detection of

respiratory viruses and SNP genotyping of CYP2C19 gene. *Nucleic Acids Res.* **35**, e40-

e40 (2007).

72. Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler

transform. *Bioinformatics* **25**, 1754-1760 (2009).

73. Langmead, B., Trapnell, C., Pop, M. & Salzberg, S. Ultrafast and memory-efficient

alignment of short DNA sequences to the human genome. *Genome Biol.* **10**, R25-R25

(2009).

Acknowledgment
950
951   We thank Jack Jiang, a high school student, for writing the demultiplexing Python script. We are

952   thankful for Dr. Thomas R. Ioerger and Dr. Michael A. DeJesus at Texas A&M University for

953   the help on the use of TRNASIT. We extend our special thanks to Dr. Jeff F. Pummill and Dr.

954   Pawel Wolinski at Arkansas High Performance Computing Center for their help to use the

955   facility.

Funding

This work was partially supported by Arkansas Biosciences Institute. The first author was supported by his parents, Cell and Molecular Biology (CEMB) program at the University of Arkansas, and Human Capacity Development Program-Kurdistan Regional Government (HCDP-KRG).

Author Contributions

Conceived and designed the experiments: YK SK. Performed the experiments, analyzed the data, wrote the manuscript: SK. Proteomics work: RL SK AQ JL. Revised the manuscript: YK SK. All authors read the final version of the manuscript.

Competing financial interests

 The authors declare no competing financial interests.

Supplementary information

Figure S1. Stability of H$_2$O$_2$ in LB medium during the experiments. LB broth media supplemented with freshly diluted 3.5 mM H$_2$O$_2$ at each of indicated time points. At 0 h, immediately after adding H$_2$O$_2$ to LB broth, the media inoculated with *Salmonella* Typhimurium. At 24 h, 24 hours before the inoculation media with bacteria H$_2$O$_2$ added to media, and at 11 d, 11 days before the inoculation media with bacteria H$_2$O$_2$ added. The media supplemented with H$_2$O$_2$ left at room temperature. LB was free of H$_2$O$_2$. The cultures were incubated at 37°C for 24 h in a 96-well plate with OD$_{600}$ reading every 10 minutes.

Figure S2. Tn-seq library preparation diagram for Illumina sequencing. The genomic DNA extracted from the selected library and subjected to two PCR amplifications. First PCR was linear and specific forward primer used to capture and amplify Tn5 junctions. Second PCR was

exponential and Illumina adaptors with a barcode added. The PCR product gel purified and

sequenced on an Illumina platform.

Figure S3. The reproducibility of Tn-seq. Correlation between reads per ORFs of *Salmonella*

Typhimurium Tn-seq conditions, H_2O_2L (2.5 mM) and H_2O_2H (3.5 mM). Two ORFs excluded in

this correlation, STM14_2422 and STM14_2428.

Table S1. *Salmonella* Typhimurium Tn-seq sequencing in numbers. Number of extracted

reads, mapped reads, and unique insertions for LB (H_2O_2 free), H_2O_2L (2.5 mM), and H_2O_2H

(3.5 mM) presented. Mean length of mapped genomic junctions and hits per nucleotide shown.

Table S2. Full list of *Salmonella* Typhimurium Tn-seq genome of the study. The list of 137

Salmonella required genes for H_2O_2 resistance. The full data set of Tn-seq genome analysis with

three tools, ARTIST, Tn-Seq Explorer, and TRANSIT for H_2O_2L (2.5 mM) and H_2O_2H (3.5

mM).

Table S3. List of functional categories required for *Salmonella* Typhimurium H_2O_2

resistance. SP_PIR_Keywords used with default options for functional categories analysis of the

137 genes that were required for H2O2 resistance in *S*. Typhimurium. The gene recognition by

the analysis tool based on official gene symbols.

Table S4. Lag time, growth rate, and maximum OD$_{600}$ of *Salmonella* Typhimurium

mutants. 50 mutants and wild-type of *S*. Typhimurium grown in LB (H_2O_2 free), H_2O_2L (2.5

mM), H_2O_2H (3.5 mM). The cultures were incubated at 37°C for 24 h in a 96-well plate with

OD$_{600}$ reading every 10 minutes. Lag time, growth rate, and maximum OD600 calculated for

each mutant and compared to the wild-type.

Table S5. Full list of *Salmonella* Typhimurium proteomic analysis in response to H_2O_2. *S*.

Typhimurium strain 14028S grown in LB (H_2O_2 free), H_2O_2L (2.5 mM), H_2O_2H (3.5 mM) till

mid-log phase. Proteome profiles analyzed by utilizing ESI-LC-MS/MS in data-dependent

acquisition (DDA) mode and LC-QQQ-ESI-MS for targeted proteomics.

Table S6. Deferentially expressed proteins of *Salmonella* Typhimurium in response to H_2O_2

and their pathways. *S*. Typhimurium strain 14028S grown in LB (H_2O_2 free), H_2O_2L (2.5 mM),

H_2O_2H (3.5 mM) till mid-log phase. KEGG pathway analysis used to categorize deferentially

expressed proteins ($p < 0.05$). Blue for downregulated proteins, red for upregulated proteins and

bold represents essential proteins.

1012

1013

Figure 1. Study design and identification of the genes required for H_2O_2 resistance using

Tn-seq. (A) The effect of H_2O_2 on the growth rate of wild type *Salmonella* Typhimurium. An

overnight culture of bacteria was diluted 1:200 in the LB medium contains either 2.5 mM H_2O_2

(H_2O_2L), 3.5 mM H_2O_2 (H_2O_2H), or LB without H_2O_2 was used as control. The cultures were

incubated at 37°C for 24 h in a 96-well plate. The reduced growth rates for the H_2O_2 were in

comparison to the control. In the all growth curve figures in this work, the blue color represents

LB (no H_2O_2), the red is H_2O_2L, and green is H_2O_2H. (B) Schematic representation of the Tn-seq

study. The *Salmonella* transposon mutant library was inoculated into LB and LB contains H_2O_2L

or H_2O_2H. The three cultures were grown until they reached mid-exponential phase. The DNA

was extracted from each culture and subjected to library preparation, sequencing, and data

analysis. (C) The Tn-seq profile of the three conditions. It shows 34 kb of the *Salmonella*

genome which starts with *metF* gene and ends with *rrsB*, horizontal axis. The height of vertical

axis represents number of reads which is 1500 sequencing reads. The highlighted genes in red

are *katG*, catalase peroxidase, was tolerated insertions in the both H_2O_2 conditions; *oxyR* was not

tolerated insertions in presence of H_2O_2 and indicated that the gene is required to H_2O_2

resistance; *murI*, glutamate racemase, was not tolerated any insertions at all and it was

considered an essential gene, *murI* is required for the biosynthesis of a component of cell wall

peptidoglycan.

1032

Figure 2. Growth curve of 50 mutants and a wild type *Salmonella* in LB, H_2O_2L and

H_2O_2H. The lag phase time, growth rate, and maximum OD_{600} of the individual *Salmonella*

Typhimurium mutants and the wild type in the growth conditions of LB (no H_2O_2), H_2O_2L and

H_2O_2H. The overnight cultures of the mutants and wild type *Salmonella* were diluted 1:200 in

the LB medium, and the LB contains either 2.5 mM H_2O_2 (H_2O_2L) or 3.5 mM H_2O_2 (H_2O_2H).

The cultures were incubated at 37°C for 24 h in a 96-well plate and the OD_{600} was recorded

every 10 min. The lag phase time, growth rate, and maximum OD_{600} were calculated and shown

here as a graphical representation. The pale pink color indicates a short lag phase time, a high

growth rate, and a high OD_{600}. The red color indicates that the bacteria was stayed in a lag

phase, growth rate was close to a zero, and the OD_{600} of the culture was not raised in the 24 h

time of assays. The data of this figure can be found in Table S4.

Figure 3. The role of aromatic amino acid biosynthesis genes in resistance to the H_2O_2. (A)

Schematic representation of aromatic amino acid biosynthesis, adapted from the KEGG pathway

database. The genes in red color were identified by the Tn-seq for H_2O_2 resistance in *Salmonella*.

The red bold color genes were identified by the Tn-seq and the phenotypes were validated by the

individual mutant assays. (B) The overnight cultures of the individual mutants were diluted

1:200 in the LB (no H_2O_2) and the LB contains either 2.5 mM H_2O_2 (H_2O_2L) or 3.5 mM H_2O_2

(H_2O_2H). The cultures were incubated at 37°C for 24 h in a 96-well plate. The colors of growth

curve figures are blue for LB, red for H_2O_2L, and green for H_2O_2H. In the $\Delta aroK$ growth curve,

the red color is under the green color. (C) Differential expression of *Salmonella* proteins in

response to the H_2O_2L compared to the LB. Wild type *Salmonella* was grown in LB, H_2O_2L, and

H_2O_2H until mid-exponential phase. Targeted-proteomics was quantified AroB, AroE_2, AroK,

and AroA protein expressions in response to H_2O_2L. The shown peaks represent a unique peptide

of the three peptides that were used of protein expression analysis.

1057

Figure 4. The *ybbK*, *ybbL*, and *ybbM* have a role in resistance to H_2O_2. (A) The Tn-seq

profile of the LB (no H_2O_2), H_2O_2L (2.5 mM), and H_2O_2H (3.5 mM). It shows ~6 kb of

Salmonella Typhimurium genome starts with *ybbK* and ends with *ybbO*, horizontal axis. The

read scale for the conditions are 4000, vertical axis. (B) The growth curve of $\Delta yybK$, $\Delta yybL$, and

$\Delta yybM$. The overnight cultures of these three mutants were diluted 1:200 in LB, H_2O_2L, and

H_2O_2H. The cultures were incubated at 37°C for 24 h in a 96-well plate. The growth curve colors

are blue which represents LB, red is H_2O_2L, and green is H_2O_2H. (C) Wild type *Salmonella* was

grown in LB, H_2O_2L, and H_2O_2H until mid-exponential phase. Targeted-proteomics was

quantified YbbK, YbbL, and YbbM protein expressions in response to H_2O_2L. The shown peaks

represent a unique peptide of the three peptides that were used of protein expression analysis.

1068