Title: Dissecting the genetic architecture of shoot growth in carrot (*Daucus carota* L.) using a

diallel mating design

Authors: Sarah D. Turner[*], Paul L. Maurizio[†,‡], William Valdar[‡], Brian S. Yandell[*,§], and

Philipp W. Simon[*,**]

Author Affiliations:

[*]Department of Horticulture, University of Wisconsin-Madison, 1575 Linden Drive, Madison,

WI 53706, USA

[†]Curriculum in Bioinformatics and Computational Biology, University of North Carolina, Chapel

Hill, NC 27599, USA

[‡]Department of Genetics, University of North Carolina, Chapel Hill, NC 27599

[§]Department of Statistics, University of Wisconsin-Madison, 1300 University Avenue, Madison,

WI 53706, USA

[**]USDA-ARS Vegetable Crops Research Unit, 1575 Linden Drive, Madison, WI 53706, USA

Title: Diallel analysis of carrot shoot growth

Keywords: Diallel, Bayesian mixed model, genetic architecture, multiple imputation, heterosis

Corresponding Author:

Philipp W. Simon

Department of Horticulture, University of Wisconsin-Madison

1575 Linden Drive

Madison, WI 53706

Ph: (608) 262-1248

philipp.simon@ars.usda.gov

Article Summary

Breeding for improved competitive ability is a priority in carrot, which suffers yield losses under

weed pressure. However, improvement and in-depth genetic studies for these traits relies on

knowledge of the underlying genetic architecture. This study estimated heritable and non-

heritable components of carrot shoot growth from a diallel mating design using a Bayesian

mixed model. Results directly contribute to improvement efforts by providing estimates of

combining ability, identifying a useful tester line, and characterizing the genetic and non-genetic

influences on traits for improved competitive ability in carrot.

ABSTRACT

Crop establishment in carrot (*Daucus carota* L.) is limited by slow seedling growth and

delayed canopy closure, resulting in high management costs for weed control. Varieties with

improved growth habit (i.e. larger canopy and increased shoot biomass) may help mitigate weed

control, but the underlying genetics of these traits in carrot is unknown. This project used a

diallel mating design coupled with recent Bayesian analytical methods to determine the genetic

basis of carrot shoot growth. Six diverse carrot inbred lines with variable shoot size were crossed

in WI in 2014. F1 hybrids, reciprocal crosses, and parental selfs were grown in a randomized

complete block design (RCBD) with two blocks in CA (2014, 2015) and in WI (2015).

Measurements included canopy height, canopy width, shoot biomass, and root biomass. General

and specific combining abilities were estimated using Griffing's Model I. In parallel, additive,

inbreeding, epistatic, and maternal effects were estimated from a Bayesian linear mixed model,

which is more robust to dealing with missing data, outliers, and theoretical constraints than

traditional biometric methods. Both additive and non-additive effects significantly influenced

shoot traits, with non-additive effects playing a larger role early in the growing season, when

weed control is most critical. Results suggest that early season canopy growth and root size

express hybrid vigor and can be improved through reciprocal recurrent selection.

Author Contribution Statement

SDT and PWS conceived and designed this study. SDT performed the crosses, collected phenotypic data, ran analyses, and wrote the manuscript. PLM and WV contributed to descriptions of BayesDiallel and assisted with corresponding analyses and interpretations. BSY helped perform imputation and contributed to the presentation and interpretation of results. PLM, WV, BSY, and PWS provided comments on the manuscript prior to submission.

Acknowledgements

We are grateful to Dr. Shelby Ellison, Guillaume Ramstein, Joe Gage, and Robert Corty for helpful discussion regarding the content of this manuscript and to Rob Kane for technical support. This research was supported by the USDA National Institute of Food and Agriculture under award number 2011-51300-30903 of the Organic Agriculture and Research Extension Initiative.

INTRODUCTION

Carrots are the 7[th] most consumed fresh vegetable in the United States, with an annual

per capita consumption of 3.9 kg (USDA-ERS 2016) and a production value of $762 million

USD in 2015 (USDA-NASS 2016). In the US, the high alpha- and beta-carotene content in

carrots is the leading source of dietary provitamin A (Block 1994; Simon et al. 2009), which is

essential for healthy immune function, vision, reproduction, and cellular communication

(Institute of Medicine, Food and Nutrition Board 2001; Johnson and Russel 2004; Solomons

2012). Despite the economic and dietary importance of carrots, crop establishment and

productivity remains limited by erratic germination, slow seedling growth, and delayed canopy

closure (Rubatzky et al. 1999). This growth habit, coupled with thin, highly segmented leaf

laminae, competes ineffectively with weeds for nutrients, water, and light, resulting in yield

losses caused by reductions in root size and marketability (Bellinder et al. 1997; Bell et al. 2000).

Furthermore, in a survey of weed competitiveness in 25 crops, carrot had the largest reduction in

yield under weed pressure (van Heemst 1985).

To limit yield loss, carrots have an extended critical weed-free period ranging from three

to six weeks, during which chemical and hand weeding are necessary (Swanton et al. 2010).

Hand weeding, while effective, is disruptive to plant growth and requires intensive labor, with

estimated costs exceeding $4000 USD/ha (Bell et al. 2000). For organic systems, which

constitute 14.4% of carrot acreage in the US (USDA-ERS 2016), hand weeding is typically the

primary method of weed management. Even in conventional systems, few herbicides are labeled

for carrots and can only be applied when plants reach a threshold height (e.g. linuron) or have

five to six true leaves (e.g. metribuzin), by which point weeds have exceeded control stages

(Bellinder et al. 1997).

Cultivars with increased weed competitiveness offer a low cost, nonchemical, and

sustainable addition to an integrated weed management program (Pérez de Vida et al. 2006;

McDonald and Gill 2009). Improved competitive ability has been linked to traits that increase

resource allocation, such as height and biomass accumulation, in other densely planted crops

such as maize (Mohammadi 2007; Zystro et al. 2012), rice (Ni et al. 2000; Fischer et al. 2001;

Pérez de Vida et al. 2006), wheat (Lemerle et al. 1996; Murphy et al. 2008), and soybean

(Jannink et al. 2000; da Silva et al. 2013). While improvement of these traits offers a potential

solution for weed management, it is unknown how these traits are inherited in carrot or how they

influence marketability (e.g. root biomass accumulation).

The diallel mating design, which consists of pairwise combinations among a group of

inbred parents, is a classical breeding approach to identify informative testers, select desirable

hybrid combinations, and to determine the primary genetic control for complex traits (Hayman

1954a; Hayman 1954b; Gardner and Eberhart 1966). First introduced to plant breeding by

Sprague and Tatum in 1942, the relationships generated in a diallel crossing scheme allow

estimation of general (GCA) and specific combining abilities (SCA), which correspond to the

proportion of additive and non-additive genetic variation, respectively (Sprague and Tatum 1942;

Hayman 1954a; Hayman 1954b; Griffing 1956a; Griffing 1956b). However, the diallel mating

design is underutilized in genetic studies due to resource constraints, the complexity of the

analysis, and untenable assumptions (e.g. no epistasis; independent distribution of genes among

parents) (Baker 1978). These challenges often lead to misinterpretation of the estimates obtained

from diallel analysis and have been the subject of much debate in the literature (Baker 1978;

Hallauer and Miranda 1981; Christie and Shattuck 1992).

Several methods of diallel analysis have been described, of which the methodology

proposed by Griffing (1956a) is one of the most common. This method estimates the significance

of GCA, SCA, and reciprocal cross effects using a general linear model approach and can be

modified to test interactions of main effects across environments (Lin et al. 1977; Zhang and

Kang 1997). However, these traditional methods are not robust in addressing common issues

encountered in field experiments, such as missing data, imbalance, and outliers. Missing data for

a single cross is often addressed by list- or pair-wise deletion, thereby substantially reducing the

number of observations and power of the analysis.

The challenges of missing data and theoretical ambiguity have been addressed by the

development of Bayesian methods for diallel analysis, which improve biological interpretability

and expand the types of questions that can be addressed (Greenberg et al. 2010; Lenarcic et al.

2012). In this study, we used the methodologies developed by Griffing (1956a) and by Lenarcic

et al. (2012) to elucidate the relative importance of genetic parameters for shoot growth in

carrots. The primary goals of this work were (1) to estimate the inheritance of shoot growth in

carrots as a resource to inform selection strategies, identify useful testers, and assess hybrid

performance and (2) to present an applied framework for the analysis of multiple environment

data using advanced multiple imputation and Bayesian methods.

129   **MATERIALS AND METHODS**

Plant material and measurements

Six inbred lines, with canopy heights ranging from short (31.4 cm) to tall (52.8 cm), were

selected from the USDA-VCRU carrot breeding program and included L6038, L7550, P0159,

Nbh2189, P6139, and B7262 (**Table 1, Figure 1, Figures S1-S2**). Inbred parents were combined

in all pairwise combinations for a total of 36 combinations. The resulting F1 progenies,

reciprocals, and parental selfs were grown in a randomized complete block design (RCBD) with

two blocks. Field sites included the Hancock Agricultural Research Station (Hancock, WI, USA;

2015) and the University of California Imperial County Cooperative Extension Station

(Holtville, CA, USA; 2015 & 2016). Carrots were grown on 1.5 meter (m) plots with 1 m

between-row spacing.

Measurements of each trait were taken for three biological replicates per plot and are

summarized in **Table 1**. Canopy height (cm) and width (cm) were measured at midseason, 80

days after planting (DAP), and at harvest, 130 DAP. At harvest, fresh and dry biomass (g) were

recorded separately for both shoot and root tissue. For dry biomass, samples were dried at 60°C

in a forced-draft oven until reaching a constant weight. A natural log transformation, ln(x), was

applied to biomass measurements to make the data distribution symmetric and stabilize the

variance.

Statistical Analyses

Diallel data for each phenotype was analyzed using two complementary approaches: a

traditional analysis after Griffing (1956a), which, owing to its requirement that data is complete

and balanced, was combined here with a multiple imputation procedure; and the recent Bayesian

mixed model decomposition of Lenarcic et al. (2012), performed on the raw (unimputed) data.

These are described in detail below. All analyses were performed in R. 3.2.2 (R Core Team

2016). Data and source code are available at https://github.com/mishaploid/carrot-diallel.

Multiple imputation of missing data for Griffing's analysis: To compensate for imbalance,

missing data (**Figure S3**) was imputed using the Multivariate Imputation by Chained Equations

package (R package mice; R/mice) (van Buuren and Groothuis-Oudshoorn 2011), and

specifically using that package's predictive mean matching method (PMM), which is a general

purpose, stochastic regression technique that is suitable for numeric data (Little 1988). The

predictors used for PMM were chosen based on recommendations in the R/mice documentation,

and included male parent, female parent, cross, year, replication, planting density, and numeric

measurements with complete data. The values imputed by the PMM were generated by running

its associated Markov chain Monte Carlo (MCMC) sampler until it reached a stationary

distribution (usually at around 40 iterations; **Figure S4**), and then recording sampled values from

a later iteration (e.g. iteration 70). This was repeated $m = 50$ times to generate m imputed data

sets.

Griffing's analysis: Each of the m imputed data sets was analyzed using Griffing's Method I,

Model I (1956a), which treats genotypes and blocks as fixed effects and has the base model:

$$Y_{ijk} = \mu + g_j + g_k + s_{jk} + r_{jk} + \varepsilon_{ijk}$$

where μ is the population mean, g_j and g_k are the GCA effects for the jth and kth parents,

respectively, s_{jk} is the SCA effect for the cross of the jth and kth parents ($s_{jk} = s_{kj}$), r_{jk} is the

reciprocal effect for the cross of the jth and kth parents ($r_{jk} = -r_{kj}$), and ε_{ijk} is the

environmental effect for the ijkth observation. Analysis was run using the diallele1 function in the

R package plantbreeding (Rosyara 2014), which we modified to include environmental effects

and genotype x environment interactions (GxE). The proportion of additive to non-additive

genetic variation was estimated from the fixed model by comparing the ratio of mean squares for

GCA ($\frac{MS_{GCA} - MS_{Error}}{2 \times n}$) and SCA ($MS_{SCA} - MS_{Error}$), where n is the number of parental lines

(Baker 1978). Values close to unity suggest higher predictability based solely on GCA. Mean

squares and approximate F-tests were pooled following the method proposed by Raghunathan

and Dong (2011, unpublished manuscript) (**Table S1**). Estimates for GCA, SCA, and reciprocal

effects were combined according to Rubin's rules (Rubin 1987) and as implemented in R/mice

(van Buuren and Groothuis-Oudshoorn 2011).

Bayesian mixed model for diallel analysis: Raw data for each phenotype (y_i), measured for

individuals $i \in \{1, \dots, n\}$ with female parent j and male parent k, were decomposed into additive

(a), inbreeding (b), maternal (m), symmetric cross-specific (v), and asymmetric cross-specific

effects (w), as described by Lenarcic et al. (2012) and Crowley et al. (2014) and implemented as

a Gibbs sampler in the R package BayesDiallel. Specifically, we used BayesDiallel's full

unsexed model ('fullu'):

195
$$y_i = \mu + \underbrace{x_i^T \beta}_{\text{user fixed}} + \underbrace{\sum_{r=1}^{R} u_i^{(r)}}_{\text{user random}} + \underbrace{a_{j[i]} + a_{k[i]}}_{\text{additive}} + \underbrace{I_{\{j[i]=k[i]\}}(\beta_{\text{inbred}} + b_{j[i]})}_{\text{inbreeding}} + \underbrace{m_{j[i]} - m_{k[i]}}_{\text{maternal}}$$

196
$$+ \underbrace{I_{\{j[i] \neq k[i]\}} v_{(jk)[i]}}_{\text{symmetric}} + \underbrace{I_{\{j[i] < k[i]\}} w_{(jk)[i]}}_{\text{asymmetric}} + \varepsilon_i \,,$$

where $j[i]$, $k[i]$ and $(jk)[i]$ respectively denote the female, male and female-male combination

relevant to individual i, each group of effects parameters is modeled from its own random effects

distribution, e.g. $a_j \sim N(0, \tau_a^2)$, fixed covariates x_i, and R additional random-effect components

are included as $u_i^{(r)} \sim N(0, \tau_r^2), \forall\, r \in \{1, \dots, R\}$. Planting density (1-3 scale, 1=low, 3=high) was

included as a fixed covariate to capture linear trends and as a random effect to estimate

deviations from linearity. Year (CA2015, WI2015, CA2016) was also included as a random

effect. This model was fitted in BayesDiallel using a MCMC Gibbs sampler with 5 chains,

10000 iterations, and a burn-in of 1000.

Although convention is to model heterosis as a dominance effect, this method instead

models inbred-specific deviations from heterozygote-based predictions; that is, homozygotes (i.e.

parental selfs), which are a minority in the diallel, are treated as a special class. Thus, additive

effects are a dosage effect of parent j in combination with another sampled parent. Epistasis,

modeled as deviations from additivity, is represented by symmetric and asymmetric effects.

Symmetric effects model differences specific to a given cross, regardless of parental inheritance

and independent of reciprocal effects (i.e. crosses *jk* and *kj* have the same effect). Asymmetric

effects model deviations from symmetric effects due to differences between reciprocal crosses

(i.e. *jk* and *kj* have different effects).

Diallel variance projection as a heritability-like measure: In order to report the relative

contribution of each diallel inheritance class to a given phenotype, Crowley et al. (2014)

proposed the diallel variance projection (VarP). This approach uses the posterior predictive

distribution of effects from BayesDiallel to simulate future, complete, perfectly balanced diallels

of the same parental lines. In each simulated dataset, the contribution of each inheritance class

(additive, inbreeding, etc.) is then calculated as its sum of squares (SS) divided by the total

phenotype SS. The resulting proportion, the VarP, is similar to the traditional heritability of

Mather and Jinks (1982) and Lynch and Walsh (1998) but with two important differences: 1) it is

explicitly prospective, in that it seeks to describe how much additive effects, say, would impact a

future experiment; and 2) its estimation is more precise, since it is calculated as a function

primarily of the effects parameters (e.g. $a_1, a_2, ..., a_6$), which are well informed by the data,

rather than of the variance components (τ_a^2, τ_m^2, etc.), which are typically not. VarPs are

calculated from multiple posterior draws leading to a complete posterior distribution of the VarP

for each inheritance class, summarized here as highest posterior credibility intervals. Credibility

intervals that include zero are interpreted as not contributing positive, nonzero information to the

prediction of y_i, whereas credibility intervals excluding zero provide strong evidence that an

effect is important to the model.

234   **RESULTS**

Imputation of missing data

There was a high incidence of missing data due to variation in seed production and

disease pressure (**Table 1, Figure S3**). A large proportion of missing data occurred in the

WI2015 environment, which was subject to severe infestation by Alternaria leaf blight, a fungal

pathogen that causes leaf necrosis and plant death in carrots (Pryor and Strandberg 2001).

Distributions of imputed data matched those expected from observed data when accounting for

environmental variation (**Figures S5-6**).

Additive and non-additive gene action contributed to observed phenotypes

Most phenotypes were positively correlated and significant at α=0.001, with the

exception of the ratio for shoot:root biomass with both canopy height and width at 80 DAP

(**Figure 2**). Griffing's analysis revealed significant genotypic differences for all phenotypes

(**Table 2**), which are reflected in posterior predicted means from BayesDiallel (**Figure 3 a, b;**

Figures S7-12). For all traits, both GCA and SCA contributed significantly to the observed

genotypic variation, suggesting that both additive and non-additive effects are important. The

ratio of GCA to SCA varied for each trait but was less than one for all phenotypes except

shoot:root ratio, indicating a more prominent role of non-additive gene action for most traits

(**Table 2**). This is reaffirmed by the results from BayesDiallel, in which the highest posterior

density intervals for inbreeding effects were greater and more dispersed than for additive effects

(**Figure 3 c, d; Figures S13-19**).

Inbreeding effects differed across genetic backgrounds

Results from Griffing's analysis indicated that the observed phenotypes, with the

exception of shoot:root ratio, were largely under non-additive genetic control (**Table 2**), which is

also reflected in the posterior predicted means and highest posterior density intervals from

BayesDiallel (**Figures S7-19**). These effects are illustrated by canopy height at 130 DAP, for

which inbred lines were an average of 8.4 cm shorter than their hybrid counterparts (overall

inbreeding, β_{inbred}, in **Figure 3**). Additionally, the intensity of inbreeding also varied across

genetic background (**Figure 3**, **Figures S13-19**). Relative to heterozygotes, line L6038 had a

minimal net reduction in canopy height of 3.0 cm ($\beta_{inbred} + b_{j[1]}$), while line B7262 had a net

22 cm reduction in canopy height ($\beta_{inbred} + b_{j[5]}$) (**Figure 3**).

Identification of superior parents, hybrids, and testers for applied breeding

GCA estimates were compared to determine the relative performance of each parent

(**Table 3**). Parent L6038 had negative and significant GCA for all traits except canopy height and

width at 80 DAP. Low and significant GCA was also observed in parent L7550 for height (130

DAP), shoot biomass, and the ratio of shoot:root biomass. For canopy height, parents with

positive and significant GCA included Nbh2189 (130 DAP), P6139 (80 and 130 DAP), and

B7262 (130 DAP). Parent Nbh2189 was the only inbred with significant and positive GCA for

canopy width (130 DAP). Parents P0159 and B7262 had high and significant GCA for both

shoot biomass and the ratio of shoot:root biomass. Positive and significant GCA for root biomass

was only observed for parent P0159.

GCA estimates largely agree with the results from BayesDiallel, which provided similar

rankings based on posterior predicted means (**Figure 3 a, b; Figures S7-12**) and HPD intervals

(**Fig. 3 c, d; Figures S13-19**). For canopy height (130 DAP), hybrids with parents L6038 and

L7750 were, on average, about 6.5 cm shorter, while hybrids with parents P6139 and B7262

were an average of 5.0 cm taller. (**Figure 3**). The observed and posterior predicted means for

canopy height (130 DAP) also demonstrate relatively higher values for hybrids with parents

P6139 and B7262, as well as lower values for hybrids containing parents L6038 and L7550

(**Figure 3 a, b**).

SCA effects were identified as crosses that performed better or worse than expected

based on the GCA values of the contributing parents (**Table 4**). Hybrid Nbh2189 x P6139 had

high SCA for both height and width (80 and 130 DAP). For shoot biomass, the largest SCA was

observed in hybrid Nbh2189 x B7262. Hybrids with high SCA for root biomass included L7550

x B7262, P0159 x Nbh2189, and Nbh2189 x B7262. No significant positive effects were

observed for shoot:root ratio. Line Nbh2189 was the most discriminating parent in hybrid

combination for all phenotypes except root biomass, suggesting it can serve as a valuable tester

for shoot growth in applied breeding.

Non-genetic factors and interactions influenced observed variation

Highly significant reciprocal effects were detected for all traits in Griffing's analysis

(**Table 2**), suggesting parent-of-origin influences phenotypic expression. For increasing height

and width (80 and 130 DAP), lines L6038 and P0159 tended to perform best as female parents

and lines L7550 and B7262 tended to perform best as male parents. Significant increases were

also observed for shoot biomass, root biomass, and shoot:root ratio when line L7550 was used as

a female parent and when lines P0159 and Nbh2189 were used as male parents.

Genotype by environment interaction (GxE) was significant for canopy height (80 and

130 DAP), shoot biomass (fresh and dry), and shoot:root ratio (**Table 2**). For corresponding

traits with significant GCAxE, SCAxE, and ReciprocalxE interactions, estimates for each year

are provided in **Tables S2-4**. Significant GCAxE interactions were observed for canopy height

(80 and 130 DAP), shoot biomass, and shoot:root ratio. For canopy height (130 DAP), GCA

ranked consistently negative across environments for parents L6038 and L7550 (**Figure 4**).

Parent Nbh2189 had positive GCA in all environments, but effects were only significant for the

CA2015 and CA2016 locations (**Table S2**). In general, relative rankings for parents remained

consistent across environments, but the ability to distinguish significant effects was diminished

in the WI2015 environment. The performance of parents P0159 and B7262 was notably

inconsistent and fluctuated between negative and positive values of GCA (**Figure 4**). SCAxE

interactions were significant for height (80 DAP), shoot biomass, and shoot:root ratio, but it was

still possible to identify consistently high performing hybrids across environments (**Table S3**).

Similarly, significant ReciprocalxE interactions were observed for canopy height (130 DAP) and

fresh shoot biomass (**Table S4**). Differences across replications within a year were not

significant.

Additional effects estimated in BayesDiallel included planting density (fixed and

random) and year (random) (**Figure S20**). On average, planting density increased plant height in

a mostly linear fashion, with a greater effect at 80 DAP (5.39 cm) compared to 130 DAP (3.51

cm) (**Figure 5**). Similarly, year had a greater influence at 80 DAP than at 130 DAP, with the

highest mean in the WI2015 season and the lowest mean in the CA2016 season (**Figure 5**).

Genetic architecture varied across traits and over developmental time

Although most correlations among phenotypes ranged from moderate (r>0.3) to high

(r>0.5) (**Figure 2**), genetic architecture varied substantially by trait and across developmental

time (**Figure 6**). For canopy height and width, the GCA/SCA ratio increased approximately

three-fold from 80 to 130 DAP, demonstrating a higher proportion of non-additive variation

early in the growing season (**Table 2**). Results from BayesDiallel exhibit a similar relationship,

with overall inbreeding (β_{inbred}) explaining more variation than additive effects for height and

width at 80 DAP (**Table 6, Figure 6**).

As described by Crowley et al. (2014), the variance projection of the additive diallel

inheritance class, VarP[a], can be likened to narrow sense heritability (h^2). Traits with significant

additive effects included canopy height at 80 DAP (VarP[a] = 0.22), canopy height at 130 DAP

(0.43), canopy width at 130 DAP (0.23), shoot biomass (0.31), root biomass (0.12), and

shoot:root ratio (0.29) (**Table 6, Figure 6**).

The influence of non-additive variation was largely due to overall inbreeding, which

contributed significantly to canopy height at 80 DAP (VarP[B] = 0.20), canopy width at 80 DAP

(0.21) and at 130 DAP (0.14), shoot biomass (0.09), and root biomass (0.27) (**Table 6, Figure**

6). However, parent-specific inbreeding effects, symmetric epistasis, and asymmetric epistasis

did not contribute significantly to the predicted phenotypes (**Table 6, Figure 6**). While parent-

of-origin is not a genetic effect, it did explain variation for canopy height at 80 DAP (Var[m] =

0.07) and 130 DAP (0.04), canopy width at 80 DAP (0.07) and 130 DAP (0.02), and shoot

biomass (0.05) (**Table 6, Figure 6**).

344   **DISCUSSION**

Primary gene action

In this study, we estimated genetic, parent-of-origin, and environmental effects on carrot

growth traits for six carrot inbred lines and their combinations in a 6x6 diallel framework.

Significant genetic variation contributed to all carrot growth attributes, suggesting that there is

potential to improve these traits in carrot.

Apart from canopy width at 80 DAP, all phenotypes had some level of predictability

based on inbred performance, as evidenced by the presence of non-zero additive effects. Traits

with high additivity included canopy height (130 DAP) and shoot biomass, both of which are

well documented as highly heritable polygenic traits that play a fundamental role in plant fitness

and adaptation (Khush 2001; Meyer et al. 2004; Peiffer et al. 2014).

For the parental lines in this study, we observed varying sensitivity to inbreeding, which

could be due to genetic divergence and/or differing levels of prior inbreeding (East 1936;

Birchler et al. 2003). This matches expectations based on the biological constraints of

outcrossing in carrot, which has putative susceptibility to inbreeding depression (Simon 2000).

Consequently, hybrid vigor was evident for canopy height (80 DAP), canopy width (80 DAP),

and root biomass, which had high proportions of non-additive genetic variation and significant

estimates of inbreeding, suggesting that causative loci may have heterozygote advantage

(Charlesworth and Willis 2009). This result coincides with widespread evidence of heterosis in

plants, whereby hybrids demonstrate increased developmental speed and greater biomass relative

to their inbred parents (East 1936; Birchler et al. 2003; Meyer et al. 2004), and agrees with

previous observations of hybrid vigor for root weight in carrot (Simon et al. 1982).

Breeding strategies

With discovery of cytoplasmic male sterility (CMS) in carrot (Welch and Grimball

1947), breeding strategies transitioned from selection in open-pollinated populations to an

inbred-hybrid system, thereby improving crop uniformity and vigor (Peterson and Simon 1986).

We expect that traits with a significant contribution of inbreeding, such as canopy height (80

DAP), canopy width (80 DAP), and root biomass, will be responsive to commonly used hybrid

breeding strategies in carrot, such as reciprocal recurrent selection. Alternatively, selection for

traits with high additivity, such as canopy height (130 DAP), canopy width (130 DAP), and

shoot biomass, may allow more rapid genetic gain while indirectly selecting for positively

correlated traits under non-additive control. For all phenotypes in this study, we identified

promising parental lines and hybrids for use in applied breeding. Additionally, line Nbh2189 was

notable as a favorable tester for future breeding efforts on shoot growth.

Accounting for genotype by environment interaction (GxE) is especially important in

biennial crops like carrot, as breeding programs rely on winter nurseries to achieve an annual

breeding cycle (Simon and Goldman 2007). We conducted trials in CA and WI, which are two of

the leading carrot production regions and representative of common, but contrasting, breeding

environments. In general, significant GxE interactions did not affect the ability to identify high

and low rankings among parents and hybrids. Thus, we anticipate that environmental differences

are important, but should have a minimal impact on selection efforts.

Source-sink relationships

Biomass partitioning between the shoot and root is a major consideration in carrot

breeding and has been extensively studied, both in a breeding and an ecological context (Barnes

1979; Currah and Thomas 1979; Hole et al. 1983; Thomas et al. 1983; Hole and Dearman 1991).

The ideotype for carrot shoot growth is rapid initial growth that plateaus following canopy

closure, simultaneously reducing the critical weed free period and promoting taproot

development. Equally important is avoiding growth habits with large, dense canopies, which

foster a microclimate that is conducive to the development of foliar diseases (Simon et al. 2008).

Consistent with findings by Hole et al (1983), we found a strong linear relationship

between the log transforms of shoot and root biomass ($r = 0.69$, $P<0.001$). However, the ratio of

shoot:root biomass had a wide range across parents (0.19-0.72), providing evidence that high

shoot biomass is not necessary to produce roots with high biomass, and vice versa.

Previous work has demonstrated rapid and early acquisition of dry matter in carrot

storage roots, with the taproot constituting 42% of the plant dry weight at 67 days after sowing

(Benjamin and Wren 1978). Interestingly, our results demonstrate that canopy height and width

at 80 DAP were negatively correlated with the ratio of shoot:root biomass ($r = -0.26$ and -0.29,

respectively, $P< 0.001$) and positively correlated with root biomass at harvest ($r = 0.58$ and 0.55,

respectively, *P*<0.001). Conversely, canopy height at harvest (130 DAP) was positively

correlated with shoot:root ratio (r = 0.39, *P*<0.001) and not significantly correlated with root

biomass (r = 0.10, *P*=0.18). This suggests that early shoot growth is important for root biomass

accumulation and agrees with previous conclusions that these traits are subject to hybrid vigor.

Method of analysis

When applied appropriately, traditional diallel analysis provides valuable information on

combining abilities for parental lines and the underlying gene action for complex traits (Griffing

1956a). However, the benefits of diallel mating designs are often overshadowed by practical and

theoretical constraints of the analysis. The requirement for complete, balanced data is especially

restrictive, particularly in crops with poor seed set and limited availability of inbred lines. In this

study, we performed classical evaluation using Griffing's Method I and a modern analysis using

a general Bayesian approach (BayesDiallel).   Both methods provided similar conclusions

regarding primary gene action, parental rankings, and hybrid performance.

Although the underlying model is more complex, BayesDiallel provided numerous

practical advantages over Griffing's analysis, which are thoroughly discussed by Lenarcic et al

(2012). Of relevance in this experiment was the robustness of BayesDiallel to missing data,

which was a substantial challenge when applying Griffing's model and is pervasive in field

experiments. For the latter, we chose to address missing data using multiple imputation, which

produces a set of plausible values to replace missing information (Rubin 1987). However, there

are several caveats and compromises regarding multiple imputation in that there are inadequate

or vague diagnostics and, although simple in principle, methods to pool multi-factor ANOVA

results are often vague or are not widely accessible (van Ginkel and Kroonenberg 2014; Grund et

al. 2016). Furthermore, the ability of BayesDiallel to analyze data sets in the presence of missing

data allows breeders to leverage data from partial diallels, which are often generated by proxy in

a breeding program.

A notable advantage of BayesDiallel was the option to add covariates for year and

planting density as random and fixed effects. Posterior distributions for year matched

expectations based on observed data, with higher means observed in WI2015 compared to

CA2015 and CA2016. The inclusion of planting density was especially advantageous and

matched expectations from previous studies, which demonstrated significant effects of planting

density on canopy height and biomass partitioning in carrot (Bleasdale 1967; Benjamin and

Sutherland 1992; Traka-Mavrona 1996; Li et al. 1996)

Precise estimates of heritability are useful when determining which breeding strategy will

result in the most genetic gain. While it is possible to estimate heritability through variance

component decomposition using Griffing's model II, this method requires that the parents are

selected from a random mating population (at least according to its traditional interpretation; see

discussion in Lenarcic et al. 2012), and is not accompanied by a measure of estimation

uncertainty. As an alternative, the diallel variance projection (VarP) in BayesDiallel has benefit

for applied breeding by (1) describing how several inheritance classes will influence future

experiments composed of the same parents, and (2) providing a 95% credibility interval as a

measure of uncertainty, which affords more flexibility when designing future experiments and

estimating potential for genetic gain.

Conclusions

The rise of sustainable agriculture has tasked breeders to develop cultivars with improved

weed competitive ability. Using traditional and modern approaches, we analyzed diallel data to

describe the quantitative inheritance of previously uncharacterized traits in carrot, which have

been demonstrated to confer improved competitive ability in other crops. However, future

trialing for weed competitive ability in carrot will be essential to validate the utility of these

traits, to determine the underlying mechanism of competitive ability (i.e. tolerance or

suppression), and to assess relative fitness costs (e.g. trade-offs with yield).

Results from this study support applied breeding efforts for carrot shoot growth in

numerous ways, most notably through the quantification of inbred performance, the

identification of a useful tester line (Nbh2189), and the assessment of potential hybrid

combinations. Furthermore, the detailed characterization of the inbred parents in this study

provides a foundation for the development of a multi-parental advanced generation intercross

(MAGIC) population in carrot, which will facilitate future in-depth genetic studies (Huang et al.

2015).

Lastly, we demonstrate the utility of the BayesDiallel framework for modeling heritable

and non-heritable components of carrot phenotypes. This example will make BayesDiallel more

accessible as a resource in the plant breeding community to maximize the potential of diallel

experiments, especially in under-resourced crops.

LITERATURE CITED

Baker RJ (1978) Issues in diallel analysis. Crop Sci 18:533. doi:
470   10.2135/cropsci1978.0011183X001800040001x

471   Barnes A (1979) Vegetable plant part relationships. II. a quantitative hypothesis for shoot/storage
472   root development. Ann Bot 43:487–499.

473   Bell CE, Boutwell BE, Ogbuchiekwe EJ, McGiffen ME (2000) Weed control in carrots: the
474   efficacy and economic value of linuron. HortScience 35:1089–1091.

475   Bellinder R, Kirkwyland J, Wallace R (1997) Carrot (*Daucus carota*) and weed response to
476   linuron and metribuzin applied at different crop stages. Weed Technol 11:235–240.

477   Benjamin LR, Sutherland RA (1992) Control of mean root weight in carrots (*Daucus carota*) by
478   varying within- and between-row spacing. J Agric Sci 119:59–70. doi:
479   10.1017/S0021859600071549

480   Benjamin LR, Wren MJ (1978) Root development and source-sink relations in carrot, *Daucus*
481   *carota* L. J. Exp. Bot. 29:425-433

482   Birchler JA, Auger DL, Riddle NC (2003) In search of the molecular basis of heterosis. Plant
483   Cell 15:2236–2239. doi: 10.1105/tpc.151030

484   Bleasdale JK (1967) The relationship between the weight of a plant part and total weight as
485   affected by plant density. J Hortic Sci 42:51–8.

486   Block G (1994) Nutrient sources of provitamin A carotenoids in the American diet. Am J
487   Epidemiol 139:290–293.

488   Charlesworth D, Willis JH (2009) The genetics of inbreeding depression. Nat Rev Genet
489   10:783–796. doi: 10.1038/nrg2664

490   Christie BR, Shattuck VI (1992) The diallel cross: design, analysis, and use for plant breeders.
491   In: Janick J (ed) Plant breeding reviews. John Wiley & Sons, Inc., Hoboken, NJ, pp 9–36

492   Crowley JJ, Kim Y, Lenarcic AB, et al (2014) Genetics of adverse reactions to haloperidol in a
493   mouse diallel: a drug-placebo experiment and Bayesian causal analysis. Genetics
494   196:321–347. doi: 10.1534/genetics.113.156901

495   Currah IE, Thomas TH (1979) Vegetable plant part relationships. III. modification of carrot
496   (*Daucus carota* L.) root and shoot weights by gibberellic acid and daminozide. Ann Bot
497   43:501–511.

498   da Silva AF, Galon L, Aspiazú I, et al (2013) Weed management in the soybean crop. In: El-
499   Shemy HA (ed) Soybean - pest resistance. InTechOpen. doi: 10.5772/54596

East EM (1936) Heterosis. Genetics 21:375–397.

Fischer AJ, Ramirez HV, Gibson KD, Da Silveira Pinheiro B (2001) Competitiveness of
semidwarf upland rice cultivars against palisadegrass (*Brachiara brizantha*) and
signalgrass (*B. decumbens*). Agron J 93:967–973.

Gardner CO, Eberhart AS (1966) Analysis and interpretation of the variety cross diallel and
related populations. Biometrics 22:439–452.

Greenberg AJ, Hackett SR, Harshman LG, Clark AG (2010) A hierarchical Bayesian model for a
novel sparse partial diallel crossing design. Genetics 185:361–373. doi:
10.1534/genetics.110.115055

Griffing B (1956a) Concept of general and specific combining ability in relation to diallel
crossing systems. Aust J Biol Sci 9:463–493.

Griffing B (1956b) A generalised treatment of the use of diallel crosses in quantitative
inheritance. Heredity 10:31–50.

Grund S, Lüdtke O, Robitzsch A (2016) Pooling ANOVA results from multiply imputed
datasets: a simulation study. Methodology 12:75–88. doi: 10.1027/1614-2241/a000111

Hallauer AR, Miranda JB (1981) Hereditary variance: mating designs. In: Quantitative genetics
in maize breeding, 2nd edn. Springer New York, New York, NY, pp 45–83

Hayman BI (1954a) The analysis of variance of diallel tables. Biometrics 10:235. doi:
10.2307/3001877

Hayman BI (1954b) The theory and analysis of diallel crosses. Genetics 39:789–809.

Hole CC, Barnes A, Thomas TH, et al (1983) Dry matter distribution between the shoot and
storage root of carrot (*Daucus carota* L.) I. comparison of varieties. Ann Bot 51:175–
187.

Hole CC, Dearman J (1991) Carbon economy of carrots during initiation of the storage root in
cultivars contrasting in shoot:root ratio at maturity. Ann Bot 68:427–434.

Huang BE, Verbyla KL, Verbyla AP, et al (2015) MAGIC populations in crops: current status
and future prospects. Theor Appl Genet 128:999–1017. doi: 10.1007/s00122-015-2506-0

Institute of Medicine, Food and Nutrition Board (2001) Dietary reference intakes for vitamin A,
vitamin K, arsenic, boron, chromium, copper, iodine, iron, manganese, molybdenum,
nickel, silicon, vanadium, and zinc. National Academy Press, Washington, DC

Jannink J-L, Orf JH, Jordan NR, Shaw RG (2000) Index selection for weed suppressive ability in
soybean. Crop Sci 40:1087–1094.

Johnson E, Russel R (2004) Beta-carotene. In: Coates P, Blackman M, Cragg G, et al. (eds)
Encyclopedia of dietary supplements, 2nd edn. Marcel Dekker, New York, NY, pp 81–87

Khush GS (2001) Green revolution: the way forward. Nat Rev Genet 2:815–822. doi:
10.1038/35093585

Lemerle D, Verbeek B, Cousens RD, Coombes NE (1996) The potential for selecting wheat
varieties strongly competitive against weeds. Weed Res 36:505–513. doi:
10.1111/j.1365-3180.1996.tb01679.x

Lenarcic AB, Svenson KL, Churchill GA, Valdar W (2012) A general Bayesian approach to
analyzing diallel crosses of inbred strains. Genetics 190:413–435. doi:
10.1534/genetics.111.132563

Li B, Watkinson AR, Hara T (1996) Dynamics of competition in populations of carrot (*Daucus*
carota). Ann Bot 78:203–214.

Lin C, Binns M, Thompson B (1977) The use of regression methods to study genotype-
environment interactions. Heredity 38:309–319.

Little RJA (1988) Missing-data adjustments in large surveys. J Bus Econ Stat 6:287–296. doi:
10.2307/1391878

Lynch M, Walsh B (1998) Genetics and analysis of quantitative traits. Sinauer, Sunderland, MA

Mather K, Jinks J (1982) Biometrical genetics: The study of continuous variation, 3rd edn.
Chapman Hall, London UK

McDonald GK, Gill GS (2009) Improving crop competitiveness with weeds: adaptations and
trade-offs. In: Crop physiology. Academic Press, San Diego, pp 449–488

Meyer RC, Törjék O, Becher M, Altmann T (2004) Heterosis of biomass production in
Arabidopsis. Establishment during early development. Plant Physiol 134:1813–1823. doi:
10.1104/pp.103.033001

Mohammadi GR (2007) Growth parameters enhancing the competitive ability of corn (*Zea mays*
L.) against weeds. Weed Biol Manag 7:232–236. doi: 10.1111/j.1445-6664.2007.00261.x

Murphy KM, Dawson JC, Jones SS (2008) Relationship among phenotypic growth traits, yield
and weed suppression in spring wheat landraces and modern cultivars. Field Crops Res
105:107–115. doi: 10.1016/j.fcr.2007.08.004

Ni H, Moody K, Robles RP, et al (2000) *Oryza sativa* plant traits conferring competitive ability
against weeds. Weed Sci 48:200–204. doi: 10.1614/0043-
1745(2000)048[0200:OSPTCC]2.0.CO;2

Peiffer JA, Romay MC, Gore MA, et al (2014) The genetic architecture of maize height.
Genetics 196:1337. doi: 10.1534/genetics.113.159152

Pérez de Vida FB, Laca EA, Mackill DJ, et al (2006) Relating rice traits to weed competitiveness
and yield: a path analysis. Weed Sci 54:1122–1131. doi: 10.1614/WS-06-042R.1

Peterson C, Simon PW (1986) Carrot breeding. In: Bassett MJ (ed) Breeding vegetable crops.
AVI Publishing, Westport, CT,

Pryor BM, Strandberg JO (2001) Alternaria leaf blight of carrot. In: Davis RM, Raid RN (eds)
Compedium of Umbelliferous crop diseases. American Phytopathological Society Press,
St. Paul, MN,

R Core Team (2016) R: A Language and Environment for Statistical Computing. R Foundation
for Statistical Computing, Vienna, Austria

Rosyara U (2014) plantbreeding: Analysis and visualization of data from plant breeding and
genetics experiments. http://R-Forge.R-project.org/projects/plantbreeding/

Rubatzky VE, Quiros CF, Simon PW (1999) Carrots and related vegetable Umbelliferae. CABI
Publishing, Wallingford, Oxon, UK: New York

Rubin DB (1987) Multiple imputation for nonresponse in surveys. Wiley, New York

Simon PW (2000) Domestication, historical development, and modern breeding of carrot. In:
Janick J (ed) Plant breeding reviews. John Wiley & Sons, Inc., Hoboken, NJ, pp 157–190

Simon PW, Freeman RE, Vieira JV, et al (2008) Carrot. In: Prohens J, Nuez F (eds) Vegetables
II: Fabaceae, Liliaceae, Solanaceae, and Umbelliferae. Springer, New York, NY, pp 327–
357

Simon PW, Goldman IL (2007) Carrot. In: Singh RJ (ed) Genetic resources, chromosome
engineering, and crop improvement. CRC Press, pp 497–516

Simon PW, Peterson CE, Lindsay RC (1982) Genotype, soil, and climate effects on sensory and
objective components of carrot flavor. J Amer Soc Hort Sci 107:644–648.

Simon PW, Pollak L, Clevidence B, et al (2009) Plant breeding for human nutritional quality. In:
Janick J (ed) Plant breeding reviews. John Wiley & Sons, Inc., Hoboken, NJ, pp 325–392

Solomons N (2012) Vitamin A. In: Erdman JW, Macdonald IA, Zeisel SH (eds) Present
knowledge in nutrition, 10th edn. Wiley-Blackwell, Oxford, UK, pp 157–83

Sprague GF, Tatum LA (1942) General vs. specific combining ability in single crosses of corn.
Agron J 34:923–932. doi: 10.2134/agronj1942.00021962003400100008x

Swanton CJ, O'Sullivan J, Robinson DE (2010) The critical weed-free period in carrot. Weed
Sci 58:229–233. doi: 10.1614/WS-09-098.1

Thomas TH, Barnes A, Rankin WEF, Hole CC (1983) Dry matter distribution between the shoot
and storage root of carrot (*Daucus carota* L.) II. effect of foliar application of gibberellic
acid. Ann Bot 51:189–199.

Traka-Mavrona E (1996) Effects of competition on phenotypic expression and differentiation of
five quality traits of carrot (*Daucus carota* L.) and their implications in breeding. Sci
Hortic 65:335–340. doi: 10.1016/0304-4238(96)00884-9

USDA Economic Research Service (ERS) Food availability (per capita) data system. In: Food
Availab. Capita Data Syst. http://www.ers.usda.gov/data-products/food-availability-(per-
capita)-data-system/faqs.aspx.

USDA National Agricultural Statistics Service (NASS) (2016) Quick stats database - statistics by
subject. https://www.nass.usda.gov/Statistics_by_Subject/result.php?9AF46B77-6FC6-
3021-8AB4-
12312EEF4AF7§or=CROPS&group=VEGETABLES&comm=CARROTS.

van Buuren S, Groothuis-Oudshoorn K (2011) mice: multivariate imputation by chained
equations in R. J Stat Softw 45:1–67.

van Ginkel JR, Kroonenberg PM (2014) Analysis of variance of multiply imputed data. Multivar
Behav Res 49:78–91. doi: 10.1080/00273171.2013.855890

van Heemst HDJ (1985) The influence of weed competition on crop yield. Agric Syst 18:81–93.
doi: 10.1016/0308-521X(85)90047-2

Welch JE, Grimball EL (1947) Male sterility in the carrot. Science 106:594. doi:
10.1126/science.106.2763.594

Zhang Y, Kang MS (1997) DIALLEL-SAS: a SAS program for Griffing's diallel analyses.
Agron J 89:176–182. doi: 10.2134/agronj1997.00021962008900020005x

Zystro JP, de Leon N, Tracy WF (2012) Analysis of traits related to weed competitiveness in
sweet corn (*Zea mays* L.). Sustainability 4:543–560. doi: 10.3390/su4040543

Figure 1. Variation in means and 95% confidence intervals for canopy height (130 DAP) among

carrot inbred lines used in this study.

Figure 2. Pearson's correlations (lower diagonal) and significance (upper diagonal) among

carrot growth traits measured in this study. Significance codes: *** $P \leq 0.001$, ** $P \leq 0.01$, * $P \leq$

0.05, NS not significant.

Figure 3. Diallel effects for carrot canopy height (cm) at 130 days after planting. **a** and **b** show observed and predicted means, respectively. Shading indicates height on a scale from 21.4 cm (lighter) to 63.1 cm (darker). Posterior predicted means in **b** are the result of fitting the model in BayesDiallel to the data in **a**. The highest posterior density (HPD) intervals for additive effects are shown in **c** and the HPD intervals for inbreeding effects are shown in **d**. For each effect, thin and thick horizontal lines show the 95% and 50% HPD intervals, respectively, and the vertical break displays the posterior mean. Key: 1 = L6038, 2 = L7550, 3 = P0159, 4 = Nbh2189, 5 = P6139, 6 = B7262.

642   **Figure 4.** Interaction of general combining ability (GCA) and year (WI2015, CA2015, CA2016)

643   for canopy height at 130 DAP.

Figure 5. Highest posterior density (HPD) intervals of year and planting density (stand; 1-3

scale, 1 = low, 3 = high) for canopy height at 80 DAP (black) and 130 DAP (red). For each

effect, thin and thick horizontal lines show the 95% and 50% HPD intervals, respectively, and

the vertical break displays the posterior mean.

Figure 6. Diallel variance projections characterizing the genetic architecture for each trait,

including additive (a), maternal (m), overall inbreeding penalty (B), parent-specific inbreeding

(b), symmetric (v), and asymmetric (w) effect classes. **Left:** mean and 95% credibility intervals

of effect classes for each trait. **Right:** Mean values showing overall genetic architecture.

654   **Table 1.** Carrot traits evaluated, their range among parents, and number of complete observations for each environment in this study.

Phenotype	Measurement	Unit	Parental range	Data Transformation	Number of complete observations[a]		
					CA2015	WI2015	CA2016
Canopy height	Three points within the plot	Centimeters (cm)	29.9 – 52.8	None	72	50	69
Canopy width	Three points within the plot	Centimeters (cm)	41.5 – 61.3	None	72	50	69
Shoot biomass	Fresh and dry	Grams (g)	6.43 – 21.3 (fresh) 1.02 – 3.39 (dry)	$\ln(x)$	72	49	68
Root biomass	Fresh and dry	Grams (g)	29.0 – 64.9 (fresh) 4.22 – 8.64 (dry)	$\ln(x)$	72	49	68
Shoot to root ratio	Shoot biomass/root biomass (dry)	Grams (g)	0.23 – 0.64	$\ln(x)$	72	49	68

[a] 72 observations possible per environment (36 entries x 2 replications)

Table 2. Griffing's Method I, Model I ANOVA mean square values for carrot growth traits, including: canopy height and width (cm), shoot biomass (g), root biomass (g), and the ratio of shoot:root biomass. The ratio of general combining ability (GCA) to specific combining ability (SCA) variance is also reported.

Source	df	Height 80 DAP	Height 130 DAP	Width 80 DAP	Width 130 DAP	Shoot Biomass[a] Fresh	Shoot Biomass[a] Dry	Root Biomass[a] Fresh	Root Biomass[a] Dry	Shoot:root ratio[a]
Genotype (G)	35	331.34 ***	489.3 ***	353.57 ***	433.77 ***	1.8 ***	2.13 ***	0.76 ***	0.91 ***	0.24 ***
GCA	5	516.06 ***	1488.95 ***	363.35 ***	1024.37 ***	7.27 ***	8.55 ***	1.13 ***	1.74 ***	1.27 ***
SCA	15	393.69 ***	376.33 ***	362.43 ***	435.66 ***	0.95 ***	1.21 ***	1.11 ***	1.3 ***	0.06 ***
Reciprocal	15	207.42 ***	269.07 ***	341.46 ***	235.02 ***	0.83 ***	0.91 ***	0.28 ***	0.24 **	0.07 ***
Year (E)	2	10794.6 ***	268.7 **	11668.22 ***	944.08 ***	0.11	2.06 ***	17.8 ***	19.71 ***	0.27 ***
G x E	70	25.91 *	68.48 ***	55.6	64.74	0.26 ***	0.28 **	0.12	0.12	0.04 ***
GCA x E	10	39.72 *	134.49 ***	55.63	110.75 *	0.45 ***	0.4 **	0.21 *	0.2	0.08 ***
SCA x E	30	25.26 *	43.07	36.82	50.95	0.26 **	0.31 **	0.14	0.13	0.04 ***
Reciprocal x E	30	21.96	71.88 **	74.37	63.2	0.19 *	0.2	0.09	0.09	0.02
rep(E)	3	24	12.49	56.44	106.03	0.1	0.12	0.14	0.15	0.01
Error	105	15.38	32.14	45.39	53.17	0.12	0.14	0.09	0.1	0.01
GCA/SCA		0.11	0.35	0.08	0.21	0.72	0.65	0.09	0.11	2.35

*** $P \leq 0.001$, ** $P \leq 0.01$, * $P \leq 0.05$

[a] Natural log transformation; measured at harvest (130 DAP)

660   **Table 3.** Pooled estimates of general combining ability (GCA) for carrot growth traits combined across all growing environments.

Parent	Height		Width		Biomass[a]		
	80 DAP	130 DAP	80 DAP	130 DAP	Shoot	Root	Shoot:root ratio
L6038	-1.39	-6.5***	0.09	-3.5**	-0.53***	-0.19*	-0.19***
L7550	-2.83	-4.94***	-0.63	-2.23	-0.15	-0.03	-0.08**
P0159	0.85	1.43	-1.56	-1.41	0.39***	0.27***	0.1***
Nbh2189	2.43	3.62**	3.15	7.13***	0.11	0.06	0.03
P6139	3.52*	3.87**	1.86	0.53	-0.15	-0.07	-0.03
B7262	-2.58	2.52*	-2.92	-0.53	0.33***	-0.03	0.18***
Grand Mean	32.27	45.77	43.57	53.74	1.55	2.55	0.6

*** $P \leq 0.001$, ** $P \leq 0.01$, * $P \leq 0.05$
[a] Natural log transformation; dry weight as measured at harvest (130 DAP)

662 **Table 4.** Pooled estimates of specific combining ability (SCA) for carrot growth traits combined

663 across all growing environments.

F1 hybrids	Height		Width		Biomass[a]		
	80 DAP	130 DAP	80 DAP	130 DAP	Shoot	Root	Shoot:root ratio
L6038 x L7550	-0.53	-0.75	-1.4	-2.27	0.1	0	0.06
L6038 x P0159	1.05	1.83	1.82	2.86	0.12	0.17	0.05
L6038 x Nbh2189	1.01	-2.03	0.45	-2.62	-0.24	-0.06	-0.1*
L6038 x P6139	1.66	0.96	-0.8	2.43	0.21	0.15	0.06
L6038 x B7262	2.05	0.95	4.62	2.67	-0.06	0.09	-0.04
L7550 x P0159	-1.08	-1.75	-1.91	0.61	0.1	0.19	-0.01
L7550 x Nbh2189	0.48	-0.63	-1.97	-0.28	0.28	0.04	0.09
L7550 x P6139	0.34	0.68	0.01	-0.76	-0.05	-0.03	0.01
L7550 x B7262	-3.43	-2.92	0.96	-1.45	0.33*	0.39***	0
P0159 x Nbh2189	1.46	1.77	1.38	0.23	0.09	0.28**	-0.05
P0159 x P6139	4.04*	4.99*	3.95	1.24	-0.14	-0.04	-0.05
P0159 x B7262	4.39**	2.93	4.28	4.79*	-0.12	-0.09	-0.08
Nbh2189 x P6139	5.57***	6.26**	7.01**	7.2**	0.17	0.12	0.05
Nbh2189 x B7262	-5.45**	0.6	-1.66	1.7	0.54***	0.3**	0.08
P6139 x B7262	2.32	3.17	2.73	5*	-0.22	-0.04	-0.09*
Parental selfs							
L6038	-5.24*	-1.97	-4.58	-3.73	-0.16	-0.33*	-0.03
L7550	2.35	4.31	1.74	3.14	-0.44*	-0.38**	-0.08
P0159	-13.12***	-12.06***	-10.72**	-11.63**	-0.43*	-0.83***	0.1
Nbh2189	-6.40**	-4.48	-9.11**	-8.38*	-0.47*	-0.52***	0.01
P6139	-18.85***	-19.93***	-16.83***	-19.55***	-0.15	-0.34*	-0.01
B7262	-3.42	-6.22	-8.6*	-10.37**	-0.81***	-0.88***	0.05

*** $P \le 0.001$, ** $P \le 0.01$, * $P \le 0.05$

[a] Natural log transformation; dry weight as measured at harvest (130 DAP)

666   **Table 5.** Pooled estimates of reciprocal cross effects for carrot growth traits over all growing

667   environments.

F1 hybrids	Height		Width		Biomass[a]		
	80 DAP	130 DAP	80 DAP	130 DAP	Shoot	Root	Shoot:root ratio
L6038 x L7550	4***	4.35**	4.51*	5.25**	-0.1	-0.08	-0.02
L6038 x P0159	-0.91	-2.21	-3.37	-2.62	0.19	0.12	0.05
L6038 x Nbh2189	2.97**	2.43	2.22	3.74*	0.12	0.05	0.03
L6038 x P6139	2.96**	2.76	3.01	3.61*	-0.01	0.1	-0.03
L6038 x B7262	7.18***	9.93***	8.75***	7.14***	-0.15	-0.06	-0.06
L7550 x P0159	-2.89**	-1.61	-3.44	0.32	0.58***	0.28***	0.14***
L7550 x Nbh2189	-7.17***	-4.3	-9.45***	-5.42*	0.77***	0.36***	0.22***
L7550 x P6139	-4.58***	-1.5	-7.17***	-4.13*	0.03	0.04	0.01
L7550 x B7262	-0.24	2.46	1.4	1.04	0.11	0.15	-0.02
P0159 x Nbh2189	-1.5	-6.63***	-3.29	-2.81	0.04	0.04	0.01
P0159 x P6139	-4.72***	-2.96	-6.78***	-4.24*	-0.01	-0.01	0.01
P0159 x B7262	3.18**	6.74***	1.31	4.09*	-0.07	-0.03	0.01
Nbh2189 x P6139	0.81	-1.92	-0.97	-0.82	0	0.07	-0.02
Nbh2189 x B7262	6.14***	7.35**	7.15**	8.75***	-0.28*	-0.09	-0.08*
P6139 x B7262	3.96***	2.58	5.39**	1.71	-0.01	-0.06	0

*** $P \leq 0.001$, ** $P \leq 0.01$, * $P \leq 0.05$
[a] Natural log transformation; dry weight as measured at harvest (130 DAP)

669 **Table 6.** Diallel variance projection (VarP) for carrot growth traits. Under each trait (column) is listed the predicted percentage and

670 95% credibility intervals of variance that would be attributable to each class of effect. Predictions are conditional on a future complete

671 diallel composed of the same parental lines.

Diallel inheritance class	Height		Width		Biomass[a]		
	80 DAP	130 DAP	80 DAP	130 DAP	Shoot	Root	Shoot:root ratio
Additive (a)	22.07 (0.49,37.72)	43.49 (31.24,53.98)	4.56 (-0.04,11.83)	22.75 (8.96,34.53)	31.43 (15.47,47.83)	11.84 (1.55,23.9)	29.21 (5.18,53.73)
Parent of origin (m)	7.24 (2.15,13.23)	4.48 (0.49,10.19)	6.67 (1.12,14.21)	2.22 (0.14,6.66)	5.37 (0.14,11.93)	3.95 (-0.98,10.31)	4.55 (-3.68,14.49)
Overall inbreeding (B)	19.8 (3.43,36.9)	9.1 (-0.57,20.49)	20.87 (8.54,34.36)	13.98 (2.85,26.76)	9.44 (0.82,19.86)	27.48 (7.3,49.06)	3.51 (-6.33,15.18)
Inbreeding (b)	3.79 (-11.2,21.29)	-0.08 (-9.57,11.05)	3.74 (-5.23,16.75)	3.37 (-6.07,16.45)	-0.91 (-9.46,7.08)	2.17 (-16.12,21.62)	7.34 (-6.41,22.94)
Symmetric epistasis (v)	10.82 (-0.93,36.13)	1.44 (-1.49,8.87)	4.09 (-0.24,13.07)	4.36 (-0.95,18.33)	10.6 (-0.34,23.6)	8.73 (-1.67,21.04)	11.12 (-4.57,27.55)
Asymmetric epistasis (w)	1.32 (-0.46,5.57)	1.73 (-0.2,7.15)	1.92 (-0.38,8.46)	1 (-0.1,4.46)	3.68 (-1.19,9.57)	3.27 (-1.8,9.05)	5.08 (-3.58,14.62)
Total variance explained	65.03 (56.69,72.02)	60.15 (51.04,67.74)	41.85 (29.6,52.9)	47.69 (36.41,57.67)	59.62 (50.38,67.66)	57.43 (48.6,65.16)	60.81 (51.58,68.61)
Unexplained variance	34.97 (27.98,43.31)	39.85 (32.26,48.96)	58.15 (47.1,70.4)	52.31 (42.33,63.59)	40.38 (32.34,49.62)	42.57 (34.84,51.4)	39.19 (31.39,48.42)

[a] Natural log transformation; dry weight as measured at harvest (130 DAP)