3   Title: Site and gene-wise likelihoods unmask influential outliers in phylogenomic
4   analyses
5

6   Joseph F. Walker[1*], Joseph W. Brown[1], and Stephen A. Smith[1*]
7
8   [1]Dept. Ecology and Evolutionary Biology, University of Michigan, Ann Arbor,
9   Michigan, 48109, USA
10 *Corresponding authors
11
12 Corresponding author emails: jfwalker@umich.edu, eebsmith@umich.edu
13

14   ABSTRACT

Despite the wealth of evolutionary information available from sequence data,

recalcitrant nodes in phylogenomic studies remain. A recent study of vertebrate

transcriptomes by Brown and Thomson (2016) revealed that less than one percent of

genes can have strong enough phylogenetic signal to alter the species tree. While

identifying these outliers is important, the use of Bayes factors, advocated by Brown and

Thomson (2016), is a heavy computational burden for increasingly large and growing

datasets. We do not find fault with the Brown and Thomson (2016) study, but instead

hope to build on their suggestions and offer some alternatives. Here we suggest that site-

and gene-wise likelihoods may be used to idenitfy discordant genes and nodes. We

demonstrate this in the vertebrate dataset analyzed by Brown and Thomson (2016) as

well as a dataset of carnivorous Caryophyllales (Eudicots: Superasterids). In both datasets,

we identify genes that strongly influence species tree inference, and can overrule the

signal present in all remaining genes altering the species tree topology. By using a less

computationally demanding approach, we can more rapidly examine competing

hypotheses, providing a more thorough assessment of overall conflict. For example, our

analyses highlight that the debated vertebrate relationship of Alligatoridae sister to turtles,

only has six genes with complete coverage for all species of Alligatoridae, birds and

turtles. We also find that two genes (~0.0016%) from the 1237 gene dataset of

carnivorous Caryophyllales drive the topological estimate and, when removed, the

species tree topology supports an alternative hypothesis supported by the remaining

genes. Additionally, while the genes highlighted by Brown and Thomson (2016) were

revealed to be the result of errors, we suggest that the topology produced by the outlier

genes in the carnivorous Caryophyllales may not be the result of methodological error.

Close examination of these genes revealed no obvious biases (i.e. no evidence of

misidentified orthology, alignment error, or model violations such as significant

compositional heterogeneity) suggesting the potential that these genes represent genuine,

but exceptional, products of the evolutionary process. Bayes factors have been

demonstrated to be helpful in addressing questions of conflict, but require significant

computational effort. We suggest that maximum likelihood can also address these

questions without the extensive computational burden. Furthermore, we recommend

more thorough dataset exploration as this may expose limitations in a dataset to address

primary hypotheses. While a dataset may contain hundreds or thousands of genes, only a

small subset may be informative for the primary biological question.

49   INTRODUCTION

The wealth of information in phylogenomic datasets offers the potential to resolve

the most difficult nodes in the tree of life. However, different datasets within the same

study or published by different authors that comprise equivalent taxonomic sampling

often infer competing hypotheses with high support (i.e., nonparametric bootstrap support

of 100% or posterior probability of 1.0). Prominent examples of recalcitrant nodes within

well-studied groups include many charismatic lineages such as the root of placental

mammals (Morgan et al. 2013; Romiguier et al. 2013), early branching in Neoaves

(Jarvis et al. 2014; Prum et al. 2015), and the earliest diverging lineage of angiosperms

(Wickett et al. 2014; Xi et al. 2014). Understanding the evolution of these clades relies on

being able to resolve these nodes. Finding the underlying causes of uncertainty in

phylogenomic datasets is an essential step toward resolving problematic nodes. Recently,

authors have developed means of exploring conflict between gene trees and species trees

specifically for phylogenomic datasets (Smith et al. 2015; Kobert et al. 2016; Pease et al.

2016), aiding in identification of regions of species trees with considerable uncertainty

despite strong statistical support from traditional support measures.

Brown and Thomson (2016) used Bayes factors as a means of uncovering genes

that have disproportionate influence on the reconstruction of species relationships

inferred from concatenated phylogenomic data. Using a previously published vertebrate

transcriptome dataset (Chiari et al. 2012) they found that two genes were capable of

altering the topology of the concatenated species tree with high support values (posterior

node probabilities of 1.0). Identification of these genes allowed for further analysis to

determine whether they were the result of errors in orthology detection or real biological

phenomena. While these analyses demonstrated the potential influence of strongly

conflicting genes on species tree construction, the reliance on a Bayes-factor approach

imposes an enormous computational burden on already computationally intense analyses

using large datasets.

Here we discuss an alternative to Bayes factors for identifying genes that have a

disproportionate contribution to the resolution of the species tree topology. We explore

gene-tree conflict by examining both site- and gene-specific log-likelihoods. Site-wise

log-likelihood analyses have been employed in phylogenomic datasets previously (Castoe

et al. 2009; Smith et al. 2011), primarily to compare two alternative topologies. Here, we

also examine per-gene log-likelihood differences and gene-wise conflict as in Smith et al.

(2015). We conducted these analyses on both the vertebrate dataset of Brown and

Thomson (2016) as well as a carnivorous Caryophyllales (Eudicots: Superasterids)

dataset. While the two genes that were discovered by Brown and Thomson (2016) in the

vertebrate dataset were identified to be errors of dataset construction, we discuss the

possibility that the outlier genes found in the carnivorous dataset not be a methodological

error. We hope to build upon the important conclusion drawn from Brown and Thomson

(2016) by suggesting a fast and computationally feasible means of finding outlier genes

in a phylogenomic dataset, discussing the importance of examining gene conflict and

signal, and illustrating the possibility that some outlier genes may be the result of

biological phenomena and not errors.

93   METHODS

94   *Data collection*

We obtained the 248 genes that were codon aligned analyzed by Brown and

Thomson (2016) from Dryad deposit (http://dx.doi.org/10.5061/dryad.8gm85) of the

original study (Chiari et al. 2012), focused on resolving relationships of amniotes. The

coding DNA sequences of the 1237 one-to-one orthologs were downloaded from

(XXXX) and used by Walker et. al (in review) to infer the relationships among

carnivorous Caryophyllales (Eudicots: Superasterids). All programs and commands used

in this analysis may be found at https://bitbucket.org/jfwalker/siteloglikelihood.

103   *Species trees*

Brown and Thomson (2016) used Bayesian analyses to obtain the topologies from

the Chiari et. al (2012) data set. As our study focused on the use of likelihood for

detecting overly influential genes, we ensured that maximum likelihood would

recapitulate the previous species-tree results. To construct a species tree for the vertebrate

dataset, the 248 individual genes vertebrate genes used in Brown and Thomson (2016) for

inference of highly influential genes were concatenated pxcat (Brown, Walker and Smith,

in press). The species tree was inferred with maximum likelihood as implemented in

RAxML v8.2.3 (Stamatakis 2014) using the GTR+CAT model of evolution with 200

rapid bootstrap replicates performed. The use of CAT in the species tree analysis was

performed to save computational time as final inference is still conducted under

GAMMA. The species tree for the vertebrate dataset was inferred both with all genes

present, and again inferred in the same means with the previously identified two most

highly informative genes (8916 and 11434) removed (see below). The species tree

inferred through maximum likelihood and containing all data from the carnivory dataset

was downloaded from (XXXX). Another species tree was inferred through maximum

likelihood after removing the two highly informative genes (cluster575 and cluster3300;

see below) from the supermatrix.

122   *Gene tree construction and analysis of conflict*

123   Individual gene trees were inferred using maximum likelihood with the

GTR+CAT model of evolution as implemented in RAxML. A SH-Like test (Anisimova

et al. 2011), as implemented in RAxML, was performed to analyze the gene tree support.

As this test examines alternative topologies by NNI, it is possible that during the test a

topology with a higher likelihood is found. If a better topology was found during the test

performed for this study, that topology was used in downstream analyses. All gene trees

were rooted on the outgroup (*Protopterus* for the vertebrate dataset and *Beta vulgaris* and

Spinacia oleraceae for the carnivory dataset) and any gene trees not containing the

outgroup were left out of the conflict analysis. Conflict was assessed by examining taxon

bipartitions as implemented in phyparts (Smith et al. 2015) with SH-Like support of < 80

treated as uninformative. The conflict was mapped on the species tree using the script

phypartspiecharts.py (available from

https://github.com/mossmatters/MJPythonNotebooks).

137   *Gene log-likelihood Analysis*

138   The alternate topologies obtained for the placement of turtles were used along

with the Chiari et. al (2012) concatenated dataset for a site-wise log-likelihood analysis as

implemented in RAxML using the GTR+GAMMA model of evolution. The difference in

site-wise log-likelihoods between the two topologies, as well as the gene-wise log-

likelihood differences (sum of gene-specific site log-likelihoods extracted from the

overall matrix), were calculated using R scripts (available from

https://bitbucket.org/jfwalker/siteloglikelihood).

146   *Testing for paralogy in carnivory dataset*

147   The homolog trees created from amino acid data in the study by Walker et. al (in

148   review) were downloaded from (XXX). We examined the maximum inclusion (Yang and

149   Smith 2014) homologs of the amino acid data and compared the clusters containing the

150   outlier genes to those nucleotide clusters containing the outlier genes. This allowed us to

151 examine the possibility that the nucleotide cluster contained homology errors that would

152 be exposed by the slower evolving amino acid dataset.

154   RESULTS

155   *Likelihood based species tree inferences*

Brown and Thomson (2016) inferred a species tree with Bayesian analyses. While

we do not have a specific criticism of this choice, we re-analyzed the dataset using

maximum likelihood analyses to ensure the results were recapitulated as our study

focused on the use of likelihood. From the full dataset, we recovered the same topology

of Brown and Thomson (2016), with turtles positioned as sister to the crocodilians

(genera *Alligator* and *Caiman*). The edge supporting the relationship of turtles was shown

to have a large amount of conflict and the dominant alternative position placed turtles

sister to the bird clade (Fig. 1). The overall difference in log-likelihood between the two

topologies for the vertebrate dataset was 15.83. The removal of the vertebrate genes 8916

and 11434, as shown by Brown and Thomson (2016), placed turtles sister to Aves, albeit

with low bootstrap support (BS = 12; Supplementary Fig. 1). In the carnivorous

Caryophyllales, the inferred species tree contained two edges with many conflicting gene

trees and one dominant alternative topology (Fig. 1). The edge supporting *Ancistrocladus*

and *Drosophyllum* sister to the rest of carnivorous plants received no bootstrap support

(BS = 0); however, when reanalyzed with cluster575 and cluster3300 removed the

position of *Ancistrocladus* and *Drosophyllum* changed and the relationship gained

support (BS = 100; Supplementary Fig. 1). The log-likelihood difference between the

Caryophyllales topologies was 74.94.

Gene tree conflict and log-likelihood analysis shows genes of disproportionate influence

For the vertebrate dataset, we limited conflict analysis to genes that contained

sequences for the outgroup *Protopterus*. This resulted in 93 (of 248) usable gene trees for

conflict analysis. Many genes were missing one or more taxa, with only five genes

containing information for all ingroup taxa (Table 1). Throughout the vertebrate tree, we

found conflict at many of the deeper nodes (Fig. 1). Also, the node representing the

controversial placement of turtles as sister to crocodilians had only seven genes with high

SH support (>80). Nine genes recovered, with high SH support, supported the dominant

alternative relationship of turtle's sister to a clade comprising of Alligatoridae and birds

with high SH support.

The site-wise log-likelihood analyses did not clearly identify major biases (Fig.

2A and Fig. 2C). The gene-wise log-likelihood comparison of the two dominant

topologies showed that two genes (ENSGALG00000008916 and

ENSGALG00000011434) exhibit a disproportionate influence on the overall likelihood

of the supermatrix (Fig. 2B). The genes identified using the likelihood approach

presented here were the same genes identified by Brown and Thomson (2016) using

Bayes factors. The genes had a difference in log-likelihood scores of 72.91 and 41.71,

respectively, and support the hypothesis of turtles sister to crocodilians with an average

difference in log-likelihood of any gene in the supermatrix being 3.28.

We performed this same comparison of log-likelihoods between the dominant

topologies on the carnivory dataset. We found two genes (cluster575 and cluster3300)

that contribute disproportionately to the overall likelihood and that individually have a

difference in log-likelihood scores of 33.06 and 16.63, respectively, with the average

difference of log-likelihood of a gene in the supermatrix to either topology being 2.882.

Disproportionate information may potentially be a biological reality

For the carnivorous Caryophyllales dataset, we explored the possibility that the

strongly conflicting genes cluster575 and cluster3300 reflected some methodological

error in the assembly pipeline, as is the case for the genes identified by Brown and

Thomson (2016). However, both the alignment and inferred phylogram for each gene

revealed no obvious problems or potential sources of systematic error (sparse alignment,

abnormally long branch lengths etc…). We also explored whether compositional

heterogeneity could explain the strongly conflicting results (i.e., that the relationships

were not truly conflicting, but instead incorrectly modeled). However, both RY-coding in

RAxML and explicit modeling of multiple equilibrium frequencies (2, 3, or 4

composition regimes) across the tree in p4 v1.0 (Foster 2004) failed to overturn the

inferred relationships. We further explored the possibility of misidentified orthology. By

examining the homolog tree produced from amino acid data, we identified the ortholog

from the nucleotide data to be complete (i.e., an ortholog within the homolog amino acid

tree). We found that with the slower amino acid data the sequences in the nucleotide

cluster575 were inferred as a single monophyletic ortholog within a duplicated homolog

(Supplementary Figure 2). The discrepancies that appeared between the amino acid

dataset and the CDS dataset were found to be either different in-paralogs/splice sites

maintained during the dataset cleaning procedure or short sequences that were not

identified as homologs in the CDS dataset (Supplementary table and Supplementary

Figure 2).

221   DISCUSSION

We found few genes that strongly supported the deeper relationships in the

vertebrate dataset (Fig. 1). Biological processes including substitution saturation,

hybridization, horizontal gene transfer, and incomplete lineage sorting can contribute to

conflicting signal and may explain both the conflict and lack of information. However,

limitations in the dataset might also be a factor, as few gene regions contained sequence

data for every species (Table 1). In fact, only six genes have complete sampling of all

species (birds, turtles and Alligatoridae) involved in examining the position of turtles in

relation to the crocodilian clade. Thirty-six genes had the species sampling necessary to

address the alternate hypothesis, with turtle's sister to birds. Surprisingly, only five of the

genes in the analysis contained information for all ingroup taxa (Table 1). Despite the

size of many of these phylogenomic datasets, the available data to address specific

questions may be significantly smaller and should be analyzed as it has been shown that

taxon sampling influences even large phylogenomic datasets (Walker et. al in review).

As has been noted by several authors, gene tree conflict and concordance should

be examined within phylogenomic datasets (Salichos et al. 2014; Smith et al. 2015;

Kobert et al. 2016). High support values can mask significant underlying conflict (Ryan

et al. 2013; Salichos et al. 2014; Wickett et al. 2014; Smith et al. 2015; Yang et al. 2015;

Kobert et al. 2016). This is clearly the case for the vertebrates (Chiari et al. 2012;

Crawford et al. 2015; Brown and Thomson 2016). Although the vertebrate dataset

contained gene tree conflict, significant missing data, and small likelihood differences

among alternate topologies, high posterior probabilities were reported at every node (PP

= 1.00) (Brown and Thomson 2016). The change in topology of the carnivory dataset is

also remarkable, as all 1237 genes are represented in each species and the removal of two

(0.0016%) of the genes resulted in a different topology with high support. Some authors

have noted discrepancies between coalescent and supermatrix results in these datasets

(Wickett et al. 2014; Xi et al. 2014;Walker et. al in review). The observation that a small

number of genes, in the context of supermatrix analyses, can influence the resulting

topology may help explain this phenomenon. The results from both datasets discussed

here emphasize that even with large, high coverage datasets, supermatrix analyses may be

sensitive to a small number of influential genes.

We found relatively small differences in overall log-likelihood scores between the

alternate competing hypotheses for both the carnivore and vertebrate datasets. Through a

site- and gene-wise log-likelihood comparison, we demonstrated the presence of genes

that disproportionally contribute to the species-tree inference and examine the strong

conflicting signal. While Bayes factors provide another means of finding these genes,

they are computationally expensive, which is a major concern given the growing size of

phylogenomic datasets. Using the site- and gene-wise likelihood approach, we identified

the genes that have a disproportionate effect on the likelihood in ~400 seconds using two

processors on a laptop. Identifying these genes is important for understanding potential

errors, biological processes (hybridization, horizontal gene transfer), and to avoid

violating model assumptions where strong conflicting signal cannot be incorporated.

Identifying these genes *quickly* allows for more thorough examination of the entire

dataset in addition to the outliers (not to mention the CPU years and carbon savings).

The two outlier genes in the vertebrate dataset were demonstrated to be

misidentified orthologs (Brown and Thomson 2016). Unfortunately, the genomic

resources are not available to fully examine the carnivorous outlier genes (e.g., we do not

have synteny or information on gene loss). However, we used tools and data such as

alignment analysis, compositional heterogeneity tests, and homolog analysis to examine

the two carnivorous genes to the best of our ability. Our analyses did not detect any

problems with the alignment or composition. Our homolog analyses identified one gene,

cluster 575, to be an ortholog of a gene that experienced a duplication. While we cannot

rule out every possible source of error, we also cannot identify a source of

methodological error, suggesting the possibility that the conflicting topology is the result

of real (albeit unknown) biological processes.

277   CONCLUSION

Brown and Thomson (2016) used Bayes factors to identify the phylogenetic

signal in genes and discovered that two genes supporting a topology conflicting with the

dominant topology can dominate species-tree inference. Although Bayes factors are a

powerful method of identifying support for topological relationships in a Bayesian

context, they are computationally expensive. We show that likelihood analyses, which are

significantly less computationally intensive, can also identify these genes in

phylogenomic datasets. This lower computational burden frees resources for more

thorough analyses of conflict of the identified outlier genes. We show that, despite the

size of many of these datasets, very few genes can address key topological questions due

to missing data and/or saturation. Additionally, the genes identified by Brown and

Thomson (2016) were suggested to be the result of error in orthology detection. In the

carnivory dataset, we also identify genes with strong conflicting signal, but this might be

the result of biological processes and not methodological error. The paper by Brown and

Thomson (2016) lays an exciting framework for exploring data used in phylogenomic

analyses, and we further highlight the importance of their finding. We show that for a

dataset of 1237 genes, removing two genes (0.0016%) alters the topology and provides

high support. Collectively these finding show that the potential impact of a small number

of genes on the estimation of species trees is a critical topic for further examination.

297   FUNDING

298 JFW is supported through a fellowship provided by University of Michigan department

299 of Ecology and Evolutionary Biology. JFW and SAS were supported by NFS 1354048

300 and JWB and SAS were supported by NSF 1207915.

302   ACKNOWLEDGEMENTS

303 We would like to thank Oscar Vargas, Greg Stull and Ning Wang for discussion of the

304 manuscript and methods.

306   REFERENCES

Anisimova M., Gil M., Dufayard J.F., Dessimoz C., Gascuel O. 2011. Survey of branch

support methods demonstrates accuracy, power, and robustness of fast likelihood-

based approximation schemes. Syst. Biol. 60:685–699.

Brown J.M., Thomson R.C. 2016. Bayes factors unmask highly variable information

content, bias, and extreme influence in phylogenomic analyses. Syst. Biol. syw101.

Brown J.W., Walker J.F., Smith S.A. 2017. phyx: Phylogenetic tools for Unix.

Bioinformatics. accepted.

Castoe T.A., de Koning A.P.J., Kim H.-M., Gu W., Noonan B.P., Naylor G., Jiang Z.J.,

Parkinson C.L., Pollock D.D. 2009. Evidence for an ancient adaptive episode of

convergent molecular evolution. Proc. Natl. Acad. Sci. 106:8986–8991.

Chiari Y., Cahais V., Galtier N., Delsuc F. 2012. Phylogenomic analyses support the

position of turtles as the sister group of birds and crocodiles (Archosauria). BMC

Biol. 10:65.

Crawford N.G., Parham J.F., Sellas A.B., Faircloth B.C., Glenn T.C., Papenfuss T.J.,

Henderson J.B., Hansen M.H., Simison W.B. 2015. A phylogenomic analysis of

turtles. Mol. Phylogenet. Evol. 83:250–257.

Foster P.G. 2004. Modeling compositional heterogeneity. Syst Biol. 53:485–495.

Jarvis E.D., Mirarab S., Aberer A.J., Li B., Houde P., Li C., Ho S.Y.W., Faircloth B.C.,

Nabholz B., Howard J.T., Suh A., Weber C.C., da Fonseca R.R., Li J., Zhang F., Li

H., Zhou L., Narula N., Liu L., Ganapathy G., Boussau B., Bayzid M.S.,

Zavidovych V., Subramanian S., Gabaldon T., Capella-Gutierrez S., Huerta-Cepas J.,

Rekepalli B., Munch K., Schierup M., Lindow B., Warren W.C., Ray D., Green R.E.,

Bruford M.W., Zhan X., Dixon A., Li S., Li N., Huang Y., Derryberry E.P.,

Bertelsen M.F., Sheldon F.H., Brumfield R.T., Mello C. V., Lovell P. V., Wirthlin

M., Schneider M.P.C., Prosdocimi F., Samaniego J.A., Velazquez A.M. V., Alfaro-

Nunez A., Campos P.F., Petersen B., Sicheritz-Ponten T., Pas A., Bailey T., Scofield

334 P., Bunce M., Lambert D.M., Zhou Q., Perelman P., Driskell A.C., Shapiro B.,

335 Xiong Z., Zeng Y., Liu S., Li Z., Liu B., Wu K., Xiao J., Yinqi X., Zheng Q., Zhang

336 Y., Yang H., Wang J., Smeds L., Rheindt F.E., Braun M., Fjeldsa J., Orlando L.,

337 Barker F.K., Jonsson K.A., Johnson W., Koepfli K.-P., O'Brien S., Haussler D.,

338 Ryder O.A., Rahbek C., Willerslev E., Graves G.R., Glenn T.C., McCormack J.,

339 Burt D., Ellegren H., Alstrom P., Edwards S. V., Stamatakis A., Mindell D.P.,

340 Cracraft J., Braun E.L., Warnow T., Jun W., Gilbert M.T.P., Zhang G. 2014. Whole-

341 genome analyses resolve early branches in the tree of life of modern birds. Science

342 (80-.). 346:1320–1331.

343 Kobert K., Salichos L., Rokas A., Stamatakis A. 2016. Computing the internode certainty

344 and related measures from partial gene trees. Mol. Biol. Evol. Advance Ac:1–17.

345 Morgan C.C., Foster P.G., Webb A.E., Pisani D., McInerney J.O., O'Connell M.J. 2013.

346 Heterogeneous models place the root of the placental mammal phylogeny. Mol. Biol.

347 Evol. 30:2145–56.

348 Pease J.B., Haak D.C., Hahn M.W., Moyle L.C. 2016. Phylogenomics Reveals Three

349 Sources of Adaptive Variation during a Rapid Radiation. PLoS Biol. 14:1–24.

350 Prum R.O., Berv J.S., Dornburg A., Field D.J., Townsend J.P., Moriarty Lemmon E.,

351 Lemmon A.R. 2015. A comprehensive phylogeny of birds (Aves) using targeted

352 next-generation DNA sequencing. Nature. 526:569–573.

353 Romiguier J., Ranwez V., Delsuc F., Galtier N., Douzery E.J.P. 2013. Less is more in

354 mammalian phylogenomics: AT-rich genes minimize tree conflicts and unravel the

355 root of placental mammals. Mol. Biol. Evol. 30:2134–44.

356 Ryan J.F., Pang K., Schnitzler C.E., Nguyen A.D., Moreland R.T., Simmons D.K., Koch

B.J., Francis W.R., Havlak P., Smith S.A., Putnam N.H., Haddock S.H., Dunn C.W.,

Wolfsberg T.G., Mullikin J.C., Martindale M.Q., Baxevanis A.D. 2013. The genome

of the ctenophore Mnemiopsis leidyi and its implications for cell type evolution.

Science (80-.). 342:1242592.

Salichos L., Stamatakis A., Rokas A. 2014. Novel information theory-based measures for

quantifying incongruence among phylogenetic trees. Mol. Biol. Evol. 31:1261–1271.

Smith S.A., Moore M.J., Brown J.W., Yang Y. 2015. Analysis of phylogenomic datasets

reveals conflict, concordance, and gene duplications with examples from animals

and plants. BMC Evol. Biol. 15:150.

Smith S.A., Wilson N.G., Goetz F.E., Feehery C., Andrade S.C.S., Rouse G.W., Giribet

G., Dunn C.W. 2011. Resolving the evolutionary relationships of molluscs with

phylogenomic tools. Nature. 480:364–367.

Stamatakis A. 2014. RAxML version 8: A tool for phylogenetic analysis and post-

analysis of large phylogenies. Bioinformatics. 30:1312–1313.

Walker, J.F., Yang, Y. Moore, M.J., Mikenas, J., Brockington, S.F., Timoneda, A., Smith

S.A. 2017. Widespread paleopolyploidy, gene tree conflict and recalcitrant

relationships among the carnivorous caryophyllales. in review.

Wickett N.J., Mirarab S., Nguyen N., Warnow T., Carpenter E., Matasci N.,

Ayyampalayam S., Barker M.S., Burleigh J.G., Gitzendanner M.A., Ruhfel B.R.,

Wafula E., Der J.P., Graham S.W., Mathews S., Melkonian M., Soltis D.E., Soltis

P.S., Miles N.W., Rothfels C.J., Pokorny L., Shaw A.J., DeGironimo L., Stevenson

D.W., Surek B., Villarreal J.C., Roure B., Philippe H., DePamphilis C.W., Chen T.,

Deyholos M.K., Baucom R.S., Kutchan T.M., Augustin M.M., Wang J., Zhang Y.,

Tian Z., Yan Z., Wu X., Sun X., Wong G.K.-S., Leebens-Mack J. 2014.

Phylotranscriptomic analysis of the origin and early diversification of land plants.

Proc. Natl. Acad. Sci. 111:E4859–E4868.

Xi Z., Liu L., Rest J.S., Davis C.C. 2014. Coalescent versus Concatenation Methods and

the Placement of Amborella as Sister to Water Lilies. Syst. Biol. 63:919–932.

Yang Y., Moore M.J., Brockington S.F., Soltis D.E., Wong G.K., Carpenter E.J., Zhang

Y., Chen L., Yan Z., Xie Y., Sage R.F., Covshoff S., Hibberd J.M., Nelson M.N.,

Smith S.A. 2015. Dissecting molecular evolution in the highly diverse plant clade

Caryophyllales using transcriptome sequencing. Mol. Biol. Evol.:msv081-.

Figure 1. Maximum likelihood trees inferred by RAxML for the two datasets.

Conflict analysis for the vertebrates (A) and carnivorous Caryophyllales (B). The

vertebrate analysis includes the 93 genes that contained the outgroup (*Protopterus*), and

the carnivorous Caryophyllales includes 1237 genes all of which had the outgroups

(*Spinacia oleraceae* and *Beta vulgaris*). Grey represents uninformative genes (SH-Like <

80 or no taxon representation for the bipartition), blue represents gene trees that are

concordant with the relationship, green represents the dominant alternate topology and

red represents all other conflict. Numbers on edges represent concordance/conflict.

Numbers at the nodes of the vertebrate dataset correspond to bipartition numbers in Table

1.

Figure 2. Site-specific log-likelihood analysis between competing hypotheses. Plots A

and C show the difference in log-likelihood for each site and plots B and D show the

difference in log-likelihood for each gene. Positive values in plots A and B indicate

support for turtles sister to crocodilians and negative values indicate support for turtles

sister to a clade consisting of birds and Alligatoridae. Positive values in plots C and D

indicate support for *Drosophyllum lusitanicum* and *Ancistrocladus robertsonorium* sister

to other carnivorous plants and negative values indicate support for *D. lusitanicum* and *A.*

robertsonorium forming a clade with the *Nepenthes* samples.

437   **Table 1. Number of gene trees in which all the species for a given bipartition are**

438   **present. Bipartitions correspond to node labels on figure 1.**

Bipartition number	Gene trees containing all species in the bipartition
0	5
1	5
2	246
3	248
4	5
5 (All turtles, Alligatoridae and birds)	6
6	248
7	6
8	23
9	36
10	45
11	69
12	51
13	94
Bipartition of turtles sister to a clade of birds and Alligatoridae	36

Supplementary Figure 1. Species trees inferred using maximum likelihood from the

different supermatrices. Support at each node was obtained from 200 rapid bootstrap

replicates. A) Carnivorous Caryophyllales species tree inferred from all 1237 genes. B)

Carnivorous Caryophyllales species tree inferred with cluster575 and cluster3300

removed from the supermatrix. C) Species tree for vertebrate dataset inferred with all 248

genes included in the supermatrix. D) Species tree for the vertebrate dataset inferred with

8916 and 11434 removed from the supermatrix.

Supplementary Figure 2. Homolog tree for Amino Acid clustered (726) and CDS

clustered (575) highly influential gene in the carnivorous Caryophyllales dataset.

Different genes identified in the ortholog clusters are circled on cluster 726. Genes

circled in red represent ones that are shorter and were not identified as orthologous in the

CDS dataset and genes circled in blue represent alternate paralogs or introsplice sites

used between the two clustering analyses.

468   **Supplementary Table. Sources of discrepancy between the orthologs detected in**

469   **highly influential nucleotide cluster575 and in matching amino acid homolog**

470   **cluster726.**

Ortholog in 575	Ortholog in 726	Seq length of 575 (Nuc)	Seq length of 726 (Nuc)	Reason for misidentification
Dino@67443 (*Dionaea*)	Dino@67450	2793	2991	Different copy of the in-paralog or intron splice site was retained
Dino@67443 (*Dionaea*)	Dino@9980	2793	510	Not identified as homologs in blast
RuprSFB@17320 (*Ruprechtia*)	RuprSFB@17330	2787	2787	Different copy of the in-paralog or intron splice site was retained
MJM3360@61692 (*Plumbago*)	MJM3360@44226	2211	2403	Different copy of the in-paralog or intron splice site was retained
Retr@34176 (*Reaumuria*)	Retr@1791	1044	546	Not identified as homologs in blast