1   Running head: LIKELIHOOD AND OUTLIERS IN PHYLOGENOMICS
2
3   Title: Analyzing contentious relationships and outlier genes in phylogenomics
4

5   Joseph F. Walker[1*], Joseph W. Brown[1], and Stephen A. Smith[1*]
6
7   [1]Deptartment Ecology and Evolutionary Biology, University of Michigan, Ann Arbor,
8   Michigan, 48109, USA
9   *Corresponding authors
10
11 Corresponding author emails: jfwalker@umich.edu, eebsmith@umich.edu

19   ABSTRACT

Despite the wealth of evolutionary information available from genomic and

transcriptomic data, recalcitrant relationships in phylogenomic studies remain throughout

the tree of life. Recent studies have demonstrated that conflict is common among gene

trees, and less than one percent of genes may ultimately drive species tree inference. In

this study, we examined a plant and a vertebrate dataset where supermatrix and

coalescent-based species trees conflict. Using a two-topology site-specific log-likelihood

test, we identified two highly influential genes in each dataset. While the outlier genes in

the vertebrate dataset have been shown to be the result of errors in orthology detection,

we demonstrate that the outlier genes from the plant dataset may be the result of

biological processes rather than model or methodological errors. When the outlier genes

were removed from each supermatrix, the inferred trees matched the topologies obtained

from coalescent analyses. While most tests of this nature limit the comparison to two

trees, gene tree topologies generated under processes such as incomplete lineage sorting

are unlikely to precisely match either topology in a two-topology test. We therefore

examined bipartitions across a set of trees and recover more support for the resolution

favored by coalescent analyses. These results suggest that by expanding beyond two-

topology comparisons, we can dramatically improve our understanding of the underlying

signal in phylogenomic datasets by asking more targeted edge-based questions.

                                                                                 INTRODUCTION

Many recent studies have highlighted that small changes to a dataset can yield conflicting

hypotheses at particular recalcitrant relationships with high support (i.e., 100% support

from nonparametric bootstrap (BS) or posterior probability (PP) values). Prominent

examples of this include many charismatic lineages such as the root of placental

mammals (Morgan et al. 2013; Romiguier et al. 2013), early branching in Neoaves

(Jarvis et al. 2014; Prum et al. 2015), and the earliest diverging lineage of extant

angiosperms (Zanis et al. 2002; Wickett et al. 2014; Xi et al. 2014). The resolution of

these relationships is critical to understanding the evolutionary history of their respective

clades (e.g., patterns of biochemical, morphological, and life history evolution).

49   Finding the underlying causes of uncertainty in phylogenomic datasets is an

essential step toward resolving problematic relationships. Recently, authors have

developed means of exploring conflict between gene trees and species trees specifically

for phylogenomic datasets (Salichos et al. 2014; Smith et al. 2015; Kobert et al. 2016),

aiding in the identification of regions of species trees with considerable uncertainty

despite strong statistical support from traditional support measures. Two studies have

shown that the disproportionate influence of just one or two genes on a supermatrix

analysis is capable of altering tree topology inference (Brown and Thomson 2017; Shen

et al. 2017). Brown and Thomson (2017) reanalyzed a series of datasets using Bayes

factors and found that the transcriptome data from Chiari et al. (2012) contained outlier

genes. When the outlier genes were included in phylogenetic reconstruction, a clade of

turtles+crocodilians was inferred to be sister to birds with 100% PP. The same topology

was previously inferred using ML in the original study by Chiari et al. (2012), but was

dismissed in favor of a coalescent reconstruction that placed turtles sister to

birds+crocodilians. When Brown and Thomson (2017) removed the two outlier genes,

the reduced supermatrix infers the same topology as the coalescent reconstruction with

100% PP. Another recently published study used a two-topology gene-wise likelihood

approach to examine contentious relationships across the tree of life and found

disproportionate influence of genes at all relationships examined (Shen et al. 2017).

Given the prevalence of outlier genes in phylogenomic datasets, and the continued

focus on contentious relationships in the tree of life, it is imperative that we develop

methods for analyzing conflict and selecting among alternative resolutions for recalcitrant

relationships. We build upon the discussions of Brown and Thomson (2017) and Shen et

al. (2017) by addressing whether these outlier genes violate models of evolution and

expanding beyond a comparison of only two-topologies by examining bipartitions (i.e.,

edges defining reciprocally monophyletic clades), across sites and genes. Typically, site-

wise and gene-wise log-likelihood analyses of phylogenomic datasets compare two

alternate topologies (e.g., Castoe et al. 2009; Smith et al. 2011; Shen et al. 2017).

However, given widespread gene tree discordance (e.g., due to incomplete lineage sorting,

ILS), it may be more realistic to assume that many alternative topologies are supported

within larger genomic datasets (e.g., Smith et al. 2015; Pease et al. 2016; Walker et al.

2017). Additionally, when the research question involves a single relationship and not the

entirety of the tree, it may be more appropriate to examine bipartitions instead completely

resolved topologies. This allows for any processes that may be causing conflict in the

non-focal parts of the tree to be accommodated without influencing the relationships of

interest. Here, we compare results from two-topology gene-wise log-likelihood analyses,

maximum gene-wise bipartition (MGWB) analyses, and maximum site-wise bipartition

(MSWB) analyses. We applied these measures to vertebrate (Chiari et al. 2012; Brown

and Thomson 2017) and carnivorous Caryophyllales datasets (Walker et al. 2017) (the

latter hereafter referred to as the carnivory dataset). Both datasets contain contentious

relationships, outlier genes, and, in their respective original studies, the authors dismissed

the supermatrix topology for the topology inferred using a coalescent method.

METHODS

Data collection

We obtained the 248 genes that were codon-aligned and analyzed by Brown and

Thomson (2017) from Dryad deposit (http://dx.doi.org/10.5061/dryad.8gm85) of the

original study (Chiari et al. 2012) that focused on resolving the placement of turtles

among amniotes. The coding DNA sequences of the 1237 one-to-one orthologs from

Walker et al. (2017) to infer the relationships among carnivorous Caryophyllales

(Eudicots: Superasterids) are available from Dryad

(http://datadryad.org/resource/doi:10.5061/dryad.vn730). All programs used in this

analysis may be found at https://bitbucket.org/jfwalker/maximizelikelihoods.

Species trees

Brown and Thomson (2017) used Bayesian analyses to obtain the topologies from the

Chiari et al. (2012) data set. As our study focused on the use of maximum likelihood

(ML) for detecting overly influential genes, we ensured that ML phylogenetic

reconstruction would recapitulate the previous species-tree results. To construct a

supermatrix tree for the vertebrate dataset, the 248 individual vertebrate genes used in

Brown and Thomson (2017) were concatenated using the Phyx program pxcat (Brown et

al. 2017). The species tree was inferred in RAxML v8.2.3 (Stamatakis 2014) using the

GTR+Γ model of evolution, and edge support was assessed from 200 rapid bootstrap

replicates. Supermatrix trees for the vertebrate dataset were inferred both with all genes

present, and again with the previously identified two most highly informative genes

(8916 and 11434) removed (see below). The ML inferred from all the data from the

carnivory dataset was downloaded from (http://dx.doi.org/10.5061/dryad.33m48) while a

novel ML tree was inferred from a reduced supermatrix that excluded two highly

informative genes (cluster575 and cluster3300; see below).

119   *Gene tree construction and analysis of conflict*

Individual gene trees for both datasets were inferred using ML with the GTR+ Γ model of

evolution as implemented in RAxML. A SH-like test (Anisimova et al. 2011), as

implemented in RAxML, was performed to assess gene tree edge support. As this test

examines alternative topologies by nearest-neighbor interchange (NNI), it is possible that

during the test a topology with a higher likelihood is found (i.e., an 'NNI-optimal'

topology). When a better topology was found during the test performed for this study,

that topology was used in downstream analyses. We used the pxrr program in the Phyx

package (Brown et al. 2017) to root all gene trees on the outgroup (*Protopterus* for the

vertebrate dataset, and *Beta vulgaris* and *Spinacia oleraceae* for the carnivory dataset)

and we excluded gene trees where an outgroup was not present. We mapped conflict onto

the supermatrix tree using phyparts (Smith et al. 2015) with SH-like support of < 80

treated as uninformative. Gene tree conflict was visualized using the script

phypartspiecharts.py (available from

https://github.com/mossmatters/MJPythonNotebooks. We conducted more detailed

conflict analyses used for bipartition comparisons discussed below using pxbp as part of

the Phyx package (Brown et al. 2017).

Calculating two-topology gene-wise log-likelihoods

The alternate topologies (supermatrix and coalescent) and data matrices for the vertebrate

and carnivory datasets were obtained from the original studies, Chiari et al. (2012) and

Walker et al. (2017), respectively. We calculated site-wise log-likelihood scores for the

two topologies in RAxML using the GTR+Γ model of evolution, with the data

partitioned by gene. The differences in site-wise log-likelihoods between the candidate

topologies were then calculated using scripts available from

https://bitbucket.org/jfwalker/maximizelikelihoods.

Calculating maximum gene-wise and site-wise bipartitions

We expanded beyond the fixed two-topology comparison by examining gene- and site-

wise log likelihoods over a set of trees and estimating the maximum gene-wise bipartition

(MGWB) and maximum site-wise bipartition (MSWB). The candidate set of trees

consisted of all unique gene trees as well as trees inferred from ML supermatrix and

coalescent analyses. For each tree in the candidate set, we calculated gene- and site-wise

likelihoods using RAxML with the GTR+Γ model of evolution. We then determined the

set of bipartitions in each dataset that conflicted with the supermatrix reconstruction of

the focal edge and identified which trees from the tree set resolved to which bipartition.

We calculated which tree representing each bipartition had the highest likelihood for each

gene and site. The MGWB was then identified as the bipartition containing the tree with

the highest log-likelihood. Because multiple trees can resolve the same bipartition for the

relationship of interest, the same tree need not have the highest likelihood for each gene

in order for the same bipartition to be the MGWB for each gene (e.g., ((A,B),C),(D,E))

and ((A,C),B),(D,E)) both have the bipartition separating A,B,C from D,E). The MSWBs

were found in the same way as the MGWBs but by site-wise log-likelihood instead of

gene-wise. We then recorded the frequency that each bipartition was identified as a

MGWB and MSWB. While we record the maximum log-likelihood for each site for each

bipartition, these are not comparable among bipartitions directly as the count of trees

displaying each bipartition varies and hence they differ in the number of estimated

parameters involved. However, as a coarse comparison we summed the gene-wise log-

likelihoods for the three bipartitions most frequently recognized as MGWBs and then

calculated AIC scores for each bipartition, accommodating for the increase in model

complexity (i.e., the number of topologies used for each bipartition).

171   *Testing for paralogy in carnivory dataset*

The homolog trees created from amino acid data in the study by Walker et al. (2017)

were downloaded from Dryad (http://datadryad.org/resource/doi:10.5061/dryad.vn730).

We examined the homologs clustered using amino acids as opposed to nucleotides. This

allowed us to examine whether a nucleotide cluster contained homology errors that may

be exposed by the slower evolving amino acid dataset. The code needed to perform this

analysis may be found at (https://bitbucket.org/jfwalker/maximizelikelihoods).

179   RESULTS

Gene tree conflict and log-likelihood analysis reveals genes of disproportionate influence

Our analysis of the vertebrate dataset recovered the same supermatrix topology (Fig. 1) as

found with ML by Chiari et al. (2012) and Bayesian inference by Brown and Thomson

(2017). The difference in log-likelihood between the supermatrix and coalescent

topologies for the vertebrate dataset was 15.83. Ninety-three of 248 gene trees could be

rooted on the outgroup *Protopterus* and only five of these had all taxa represented

(Supplementary Table 1). We found low support for relationships within gene trees (SH

<80) and significant gene tree conflict (Fig. 1). Of the gene trees with high support (SH

>80), seven resolved turtles+crocodilians as sister to birds (hereafter referred to as the

vertebrate supermatrix topology) and nine resolved crocodilians+birds sister to turtles

(hereafter referred to as the vertebrate coalescent topology).

The two-topology gene-wise log-likelihood comparison showed that 107 genes

had a higher likelihood score for the vertebrate supermatrix topology while 141 supported

the vertebrate coalescent topology (Figs. 2A, 3A). Two genes (ENSGALG00000008916

and ENSGALG00000011434, referred to here as 8916 and 11434, respectively) exhibited

a disproportionate influence on the overall likelihood of the supermatrix (Fig. 2A). These

two genes identified with maximum likelihood analyses matched those previously

identified as outliers using Bayes factors (Brown and Thomson 2017). These two genes

both supported the vertebrate supermatrix topology with log-likelihood scores of 72.91

and 41.71 greater than the alternative coalescent tree topology, respectively. The

difference in log-likelihood between the two topologies of the non-outlier genes ranged

from |0.015| to |20.756| with an average of |3.28| for all genes in the analysis. The

removal of the vertebrate genes 8916 and 11434, as shown by Brown and Thomson

(2017), recovered the coalescent topology, albeit with low bootstrap support (BS = 12;

Supplementary Fig. 1).

Previous work on the carnivory dataset, demonstrated that the placement of the

Ancistrocladus+Drosophyllum clade (Fig. 1) contained significant conflict and is

strongly influenced by species sampling (Walker et al. 2017). The log-likelihood

difference between the supermatrix and coalescent topologies was 74.94. The two-

topology log-likelihood comparison between the dominant topologies on the carnivory

dataset (Fig. 2C) showed that 623 genes supported *Ancistrocladus+Drosophyllum* sister

to all other carnivorous plants (hereafter referred to as carnivory supermatrix topology)

while 614 genes supported *Ancistrocladus+Drosophyllum* sister to *Nepenthes*

alata+Nepenthes ampullaria (hereafter referred to as carnivory coalescent topology; Fig.

2C&3D). Two genes (cluster575 and cluster3300) contributed disproportionately to the

overall likelihood. Individually these two genes have a difference in log-likelihood scores

between the two topologies of 33.06 and 16.63, respectively, and support the carnivory

supermatrix topology. When we reanalyzed the supermatrix with cluster575 and

cluster3300 removed, the carnivory coalescent topology was recovered, with 100% BS

support (Supplementary Fig. 1). The difference between the two topologies in log-

likelihood of the non-outlier genes ranged from |0.005| to |12.296| with an average of

|2.88| for all genes in the analysis.

Maximum Gene-wise and Site-wise bipartitions change supported topology

We compared MGWB and MSWB involving the contentious bird, crocodilian, and turtle

relationships in the vertebrate dataset (Figs. 2B, 3B,C). We found seven unique

topologies with the necessary species coverage to conduct the analyses: five gene tree

topologies from Chiari et al. (2012) and the two dominant species tree topologies. The set

of seven trees included three major conflicting bipartitions for the relationship in

question: the two resolutions found in the supermatrix and coalescent trees and birds

sister to crocodilian+mammals+turtles. 88 genes supported the vertebrate supermatrix

bipartition, 150 genes supported the vertebrate coalescent bipartition, and 10 genes

supported the third conflicting bipartition (Figs. 2B, 3B). The AIC scores for the summed

likelihoods of the supermatrix (three source trees), the coalescent (three source trees), and

the third bipartition (one source tree) were highest for the coalescent bipartition (Table 1).

The MSWB showed 76,792 sites supported the supermatrix resolution, 90,659 sites

supported the coalescent resolution, and 19,575 sites supported the third bipartition (Fig.

3C). When examining only the sites where the MSWB was 2 log-likelihood units better

than the second best, 144 sites supported the supermatrix resolution, no sites supported

the coalescent resolution, and 16 sites supported the third bipartition.

For the carnivory dataset, we found 168 unique tree topologies gene, supermatrix,

and coalescent trees. The 168 tree topologies contained 41 conflicting bipartitions for the

relationship in question with 3 dominant bipartitions. The MGWB analyses found 500

genes supported the supermatrix bipartition, 498 genes supported the coalescent

bipartition, and 239 genes supported 16 additional bipartitions (Figs. 2D, 3E). The AIC

scores for the supermatrix (44 source trees), the coalescent (56 source trees), and for the

third bipartition (24 source trees) were highest for the supermatrix bipartition with genes

and the coalescent bipartition with sites (Table 1). The MSWB showed that 272,137 sites

supported the supermatrix bipartition, 615,418 sites supported the coalescent bipartition,

and 1,230,028 sites supported 18 additional bipartitions (Fig. 3F). When examining only

sites where the MSWB was 2 log-likelihood units better than the second best, 270 sites

supported the supermatrix topology, no sites supported the coalescent topology, and 34

sites supported two other topologies.

254   *Outlier gene examination*

For the carnivory dataset, we explored the possibility that the strongly conflicting genes

cluster575 and cluster3300 reflected methodological error in the assembly pipeline, as is

the case for the genes identified by Brown and Thomson (2017). However, both the

alignment and inferred phylogram for each gene revealed no obvious problems or

potential sources of systematic error (sparse alignment, abnormally long branch lengths,

etc.). We also explored whether compositional heterogeneity could explain the strongly

conflicting results (i.e., that the relationships were not truly conflicting, but instead

incorrectly modeled). However, both RY-coding in RAxML and explicit modeling of

multiple equilibrium frequencies (2, 3, or 4 composition regimes) across the tree in p4

v1.0 (Foster 2004) failed to overturn the inferred relationships. We further explored the

possibility of misidentified orthology. By examining the homolog tree produced from

amino acid data, we identified the ortholog from the nucleotide data to be complete (i.e.,

an ortholog within the homolog amino acid tree). We found that with the slower amino

acid data the sequences in the nucleotide cluster575 were inferred as a single

monophyletic ortholog within a duplicated homolog (Supplementary Fig. 2). The

discrepancies that appeared between the amino acid dataset and the CDS dataset were

found to be either different in-paralogs/splice sites maintained during the dataset cleaning

procedure or short sequences that were not identified as homologs in the CDS dataset

(Supplementary Table 2 and Supplementary Fig. 2).

275   DISCUSSION

Biological processes including substitution saturation, hybridization, horizontal gene

transfer, and incomplete lineage sorting can contribute to conflicting signal and may

explain both conflict and lack of support widely found in phylogenomic datasets

(Salichos et al. 2014; Smith et al. 2015; Kobert et al. 2016). In addition to these processes,

limited taxonomic coverage for each gene is also a factor. While the carnivory dataset

had extensive data overlap, the vertebrate dataset only had five gene regions that

contained sequence data for every species (Supplementary Table 1). Another challenge

for phylogenomic datasets is that high support values can mask significant underlying

conflict (Ryan et al. 2013; Salichos et al. 2014; Smith et al. 2015; Kobert et al. 2016;

Pease et al. 2017). Both datasets examined here recovered high support for two different

topologies depending on the inclusion or exclusion of two genes with disproportionate

influence on the likelihood (Brown and Thomson 2017; Walker et al. 2017). In the case

of the carnivory dataset, the inferred topology changes with the inclusion or exclusion of

just 0.0016% of the genes.

Outlying genes may be the result of biological processes or methodological errors.

The outlier genes in the vertebrate dataset were found to be the result of errors in

orthology detection and not biological processes (Brown and Thomson 2017). Here, we

explored the potential sources of error for the carnivorous outlier genes. While the

genomic resources are not available to fully examine the carnivorous outlier genes (e.g.,

we do not yet have synteny or information on gene loss), our analyses did not detect any

obvious problems with alignment, compositional heterogeneity, or homology. We found

one gene, cluster 575, to be an ortholog of a gene that experienced a duplication prior to

the divergence of both ingroup and outgroup taxa (Supplementary Fig. 2). While we

cannot rule out every possible source of error, we also cannot identify a source of

methodological error, suggesting the possibility that the conflicting topology is the result

of real (albeit unknown) biological processes.

The two-topology examination explored by most authors (Castoe et al. 2009;

Smith et al. 2011; Brown and Thomson 2017; Shen et al. 2017), whether performed with

Bayes factors or maximum likelihood, can be very informative and has been shown to

help identify these outlier genes. However, when questions involve a specific relationship,

the two-topology examination may not be optimal. The two-topology analyses require

fully bifurcating trees, while the research question may surround the resolution of one

relationship. Conflict among gene trees is a reality and expected from processes such as

incomplete lineage sorting. For instance, Jarvis et al. (2014) reported that no gene trees

from a genomic data set of 48 species of birds matched the inferred species tree. The

results of a two-topology analysis may be driven by the resolution of a part of the

phylogeny other than the area of interest, as two topology analyses condition on fully

bifurcating trees that necessarily resolve conflict in the entire tree.

To overcome these limitations, instead of full topologies, we examined

bipartitions across a set of candidate topologies, as defined by the set of gene trees and

the two tree hypotheses in question. By examining bipartitions, we accommodate for

uncertainty across the rest of the tree, regardless of the process generating that

uncertainty. For both datasets, both the MGWB and MSWB analyses demonstrate

increased support for the coalescent resolution as well as support for alternative

bipartitions. The vertebrate dataset contained three alternative bipartitions for the

relationship of interest while the carnivory dataset contained 41 different bipartitions

representing 168 topologies. The AIC analysis of the vertebrate dataset MGWBs and

MSWBs also suggested better fit of the coalescent bipartition than the supermatrix

bipartition. However, while the AIC analysis of MSWBs for the carnivorous dataset

showed better fit for the coalescent resolution, the AIC analysis of the MGWBs

supported the supermatrix resolution. This may be due to a higher penalty for parameters

for the coalescent bipartition as this included 56 gene trees as compared to the

supermatrix bipartition with 44 gene trees, but could also be the result of some within

gene processes driving conflicting signal.

The MSWB analyses comparing bipartitions that were 2 log likelihood better

than the second best provide additional information on the underlying data. In these

analyses, and for both datasets, the supermatrix resolution was supported. This is in

contrast to the results when considering all sites, not just those with 2 log likelihood

better likelihoods, and the AIC results for MSWBs that both support the coalescent

resolution. This suggests that in addition to outlier genes, there may be outlier sites that

drive the resolution of these topologies in supermatrix analyses.

The results presented here contribute to a growing body of literature that address

the question of how phylogenomic analyses should proceed in the presence of highly

influential outlier genes, conflicting topologies, and ever expanding datasets (Wickett et

al. 2014; Pease et al. 2016; Brown and Thomson 2017; Shen et al. 2017; Yang et al.

2017). For example, some authors have noted, and it is the case here, that supermatrix

analyses may be more susceptible to the problem of strong outliers (Shen et al. 2017;

Walker et al. 2017). In these cases, often the resolutions inferred using a coalescent

method were favored. When the dominant process generating gene tree conflict is

incomplete lineage sorting, coalescent methods should perform better (i.e., when gene

tree diversity if modeled correctly). Some coalescent methods that weigh all gene tree

equally (e.g., Mirarab and Warnow 2015), may overcome the problem of outlier genes

even if incomplete lineage sorting is not the dominant source of conflict simply by

eliminating the disproportionate influence of one or two outlying genes. Here, we

demonstrate with two empirical examples that the coalescent resolution had higher

support when examining bipartitions without using an explicit coalescent method. While

we continue to learn about the patterns and processes that generate conflict and signal

within phylogenetic datasets, new approaches that explore that data in different ways will

continue to provide new insights. Further investigation into the behavior of each of these

methods under different evolutionary processes is warranted to better understand their

performance to the expected heterogeneity within transcriptomes or genomes.

358   FUNDING

359   JFW was supported through a fellowship provided by University of Michigan department

360   of Ecology and Evolutionary Biology. JFW and SAS were supported by NFS 1354048

361   and JWB and SAS were supported by NSF 1207915.

363   ACKNOWLEDGEMENTS

364   We would like to thank Nathanael Walker-Hale, Javier Igea, Oscar Vargas, Greg Stull,

365   and Ning Wang for discussion of the manuscript and methods.

367   REFERENCES

Anisimova M., Gil M., Dufayard J.F., Dessimoz C., Gascuel O. 2011. Survey of branch

support methods demonstrates accuracy, power, and robustness of fast likelihood-

based approximation schemes. Syst. Biol. 60:685–699.

Brown J.M., Thomson R.C. 2017. Bayes Factors Unmask Highly Variable Information

Content , Bias , and Extreme Influence in Phylogenomic Analyses. Syst. Biol.

66:517–530.

Brown J.W., Walker J.F., Smith S.A. 2017. Phyx: phylogenetic tools for unix.

Bioinformatics. 33:1886–1888.

Castoe T.A., de Koning A.P.J., Kim H.-M., Gu W., Noonan B.P., Naylor G., Jiang Z.J.,

Parkinson C.L., Pollock D.D. 2009. Evidence for an ancient adaptive episode of

convergent molecular evolution. Proc. Natl. Acad. Sci. 106:8986–8991.

Chiari Y., Cahais V., Galtier N., Delsuc F. 2012. Phylogenomic analyses support the

position of turtles as the sister group of birds and crocodiles (Archosauria). BMC

Biol. 10:65.

Foster P.G. 2004. Modeling compositional heterogeneity. Syst Biol. 53:485–495.

Jarvis E.D., Mirarab S., Aberer A.J., Li B., Houde P., Li C., Ho S.Y.W., Faircloth B.C.,

Nabholz B., Howard J.T., Suh A., Weber C.C., da Fonseca R.R., Li J., Zhang F., Li

H., Zhou L., Narula N., Liu L., Ganapathy G., Boussau B., Bayzid M.S.,

Zavidovych V., Subramanian S., Gabaldon T., Capella-Gutierrez S., Huerta-Cepas J.,

Rekepalli B., Munch K., Schierup M., Lindow B., Warren W.C., Ray D., Green R.E.,

Bruford M.W., Zhan X., Dixon A., Li S., Li N., Huang Y., Derryberry E.P.,

Bertelsen M.F., Sheldon F.H., Brumfield R.T., Mello C. V., Lovell P. V., Wirthlin

M., Schneider M.P.C., Prosdocimi F., Samaniego J.A., Velazquez A.M. V., Alfaro-

Nunez A., Campos P.F., Petersen B., Sicheritz-Ponten T., Pas A., Bailey T., Scofield

P., Bunce M., Lambert D.M., Zhou Q., Perelman P., Driskell A.C., Shapiro B.,

Xiong Z., Zeng Y., Liu S., Li Z., Liu B., Wu K., Xiao J., Yinqi X., Zheng Q., Zhang

Y., Yang H., Wang J., Smeds L., Rheindt F.E., Braun M., Fjeldsa J., Orlando L.,

Barker F.K., Jonsson K.A., Johnson W., Koepfli K.-P., O'Brien S., Haussler D.,

Ryder O.A., Rahbek C., Willerslev E., Graves G.R., Glenn T.C., McCormack J.,

Burt D., Ellegren H., Alstrom P., Edwards S. V., Stamatakis A., Mindell D.P.,

Cracraft J., Braun E.L., Warnow T., Jun W., Gilbert M.T.P., Zhang G. 2014. Whole-

genome analyses resolve early branches in the tree of life of modern birds. Science

(80-.). 346:1320–1331.

Kobert K., Salichos L., Rokas A., Stamatakis A. 2016. Computing the internode certainty

and related measures from partial gene trees. Mol. Biol. Evol. Advance Ac:1–17.

Mirarab S., Warnow T. 2015. ASTRAL-II: Coalescent-based species tree estimation with

many hundreds of taxa and thousands of genes. Bioinformatics. 31:i44–i52.

Morgan C.C., Foster P.G., Webb A.E., Pisani D., McInerney J.O., O'Connell M.J. 2013.

Heterogeneous models place the root of the placental mammal phylogeny. Mol. Biol.

Evol. 30:2145–56.

Pease J.B., Brown J.W., Walker J.F., Hinchliff C.E., Smith S.A. 2017. Quartet Sampling

distinguishes lack of support from conflicting support in the plant tree of life.

BioRxiv.

Pease J.B., Haak D.C., Hahn M.W., Moyle L.C. 2016. Phylogenomics Reveals Three

Sources of Adaptive Variation during a Rapid Radiation. PLoS Biol. 14:1–24.

Prum R.O., Berv J.S., Dornburg A., Field D.J., Townsend J.P., Moriarty Lemmon E.,

Lemmon A.R. 2015. A comprehensive phylogeny of birds (Aves) using targeted

next-generation DNA sequencing. Nature. 526:569–573.

Romiguier J., Ranwez V., Delsuc F., Galtier N., Douzery E.J.P. 2013. Less is more in

mammalian phylogenomics: AT-rich genes minimize tree conflicts and unravel the

root of placental mammals. Mol. Biol. Evol. 30:2134–44.

Ryan J.F., Pang K., Schnitzler C.E., Nguyen A.D., Moreland R.T., Simmons D.K., Koch

B.J., Francis W.R., Havlak P., Smith S.A., Putnam N.H., Haddock S.H., Dunn C.W.,

Wolfsberg T.G., Mullikin J.C., Martindale M.Q., Baxevanis A.D. 2013. The genome

of the ctenophore Mnemiopsis leidyi and its implications for cell type evolution.

Science (80-.). 342:1242592.

Salichos L., Stamatakis A., Rokas A. 2014. Novel information theory-based measures for

quantifying incongruence among phylogenetic trees. Mol. Biol. Evol. 31:1261–1271.

Shen X., Hittinger C.T., Rokas A. 2017. Contentious relationships in phylogenomic

studies can be driven by a handful of genes. Nat. Ecol. Evol. 1:1–10.

Smith S.A., Moore M.J., Brown J.W., Yang Y. 2015. Analysis of phylogenomic datasets

reveals conflict, concordance, and gene duplications with examples from animals

and plants. BMC Evol. Biol. 15:150.

Smith S.A., Wilson N.G., Goetz F.E., Feehery C., Andrade S.C.S., Rouse G.W., Giribet

G., Dunn C.W. 2011. Resolving the evolutionary relationships of molluscs with

phylogenomic tools. Nature. 480:364–367.

Stamatakis A. 2014. RAxML version 8: A tool for phylogenetic analysis and post-

analysis of large phylogenies. Bioinformatics. 30:1312–1313.

Walker J.F., Yang Y., Moore M.J., Mikenas J., Timoneda A., Brockington S.F., Smith

S.A. 2017. Widespread paleopolyploidy , gene tree conflict , and recalcitrant

relationships among the. Am. J. Bot. 104:858–867.

Wickett N.J., Mirarab S., Nguyen N., Warnow T., Carpenter E., Matasci N.,

Ayyampalayam S., Barker M.S., Burleigh J.G., Gitzendanner M.A., Ruhfel B.R.,

Wafula E., Der J.P., Graham S.W., Mathews S., Melkonian M., Soltis D.E., Soltis

P.S., Miles N.W., Rothfels C.J., Pokorny L., Shaw A.J., DeGironimo L., Stevenson

D.W., Surek B., Villarreal J.C., Roure B., Philippe H., DePamphilis C.W., Chen T.,

Deyholos M.K., Baucom R.S., Kutchan T.M., Augustin M.M., Wang J., Zhang Y.,

Tian Z., Yan Z., Wu X., Sun X., Wong G.K.-S., Leebens-Mack J. 2014.

Phylotranscriptomic analysis of the origin and early diversification of land plants.

Proc. Natl. Acad. Sci. 111:E4859–E4868.

Xi Z., Liu L., Rest J.S., Davis C.C. 2014. Coalescent versus Concatenation Methods and

the Placement of Amborella as Sister to Water Lilies. Syst. Biol. 63:919–932.

Yang Y., Moore M.J., Brockington S.F., Mikenas J., Olivieri J., Walker J.F., Smith S.A.

2017. Improved transcriptome sampling pinpoints 26 paleopolyploidy events in

Caryophyllales, including two paleo-allopolyploidy events. bioRxiv.

Zanis M.J., Soltis D.E., Soltis P.S., Mathews S., Donoghue M.J. 2002. The root of the

angiosperms revisited. Proc. Natl. Acad. Sci. U. S. A. 99:6848–53.

Figure 1. Maximum likelihood trees inferred by RAxML for the Chiari et al. 2012

(vertebrate) and Walker et al. 2017 (carnivorous Caryophyllales) datasets. Conflict

analysis for the vertebrate (A) and carnivory (B) datasets. The vertebrate analysis

includes the 93 genes that contained the outgroup (*Protopterus*), and the carnivory

analysis includes 1237 genes all of which had the outgroups (*Spinacia oleraceae* and

Beta vulgaris). Blue represents gene trees that are concordant with the relationship, grey

represents uninformative genes (SH-like < 80 or no taxon representation for the

bipartition), green represents the dominant alternate topology, and red represents all other

conflict. Numbers on edges represent concordance/conflict. Bold numbers at the nodes of

the vertebrate dataset correspond to bipartition numbers in Supplementary Table 1.

Figure 2. Comparison of two-topology gene-wise likelihood test and MGWB. A&C)

Show the results of the two-topology test on the vertebrate and carnivory dataset,

respectively, using the coalescent and supermatrix topologies as the comparison. B&D)

Show the results of the MGWB test on the vertebrate and carnivory dataset, respectively,

using the bipartitons found in the supermatrix topology, the coalescent topology, and

unique gene trees with full species coverage.

Figure 3. Bar plot representing the two-topology, MGWB, and MSWB methods for

identifying support at a node. The letters S, C, and DA represent supermatrix,

coalescent, and the dominant alternative bipartitions, respectively. Panels A-C show the

results of the different methods applied to the vertebrate dataset with the most common

dominant alternative shown in B&C having a clade consisting solely of

mammals+crocodilians+turtles. Panels D-F show the results of the different methods

applied to the carnivory dataset with the most common dominant alternative shown in

E&F having a clade consisting of the two species in the genus *Nepenthes* sister to all

other carnivorous Caryophyllales.

495 **Table 1. Results of model testing the various topologies.**

Comparison	Biparitition/Topology	AIC	ΔAIC	AIC$_w$
Vertebrate Dataset				
Two Topology	Supermatrix	2094518	31	1.85e-07
	Coalescent	2094487	0.0	9.99e-01
MGWB	Supermatrix	2093604.12	622.46	0.0
	Coalescent	2092981.66	0.0	1.0
	Dominant Alternative	2098006.78	5025.12	0.0
MSWB	Supermatrix	2076738.66	2753.57	0.0
	Coalescent	2073985.09	0.0	1.0
	Dominant Alternative	2102131.54	28146.45	0.0
Carnivory Dataset				
Two Topology	Supermatrix	26625000	0.0	1.0
	Coalescent	26625150	150	0.0
MGWB	Supermatrix	26611048.76	0.0	1.0
	Coalescent	26611339.93	291.17	0.0
	Dominant Alternative	26613649.05	2600.29	0.0
MSWB	Supermatrix	26032491.44	181865.6	0.0
	Coalescent	25850625.84	0.0	1.0
	Dominant Alternative	26101950.85	251325.01	0.0

APPENDICES

Supplementary Figure 1. Species trees inferred using maximum likelihood from the

different supermatrices. Support at each node was obtained from 200 rapid bootstrap

replicates. A) carnivorous Caryophyllales species tree inferred from all 1237 genes. B)

carnivorous Caryophyllales species tree inferred with cluster575 and cluster3300

removed from the supermatrix. C) Species tree for vertebrate dataset inferred with all 248

genes included in the supermatrix. D) Species tree for the vertebrate dataset inferred with

8916 and 11434 removed from the supermatrix.

Supplementary Figure 2. Homolog tree for Amino Acid clustered (726) and CDS

clustered (575) highly influential gene in the carnivorous Caryophyllales dataset.

Different genes identified in the ortholog clusters are circled on cluster 726. Genes

circled in red represent ones that are shorter and were not identified as orthologous in the

CDS dataset and genes circled in blue represent alternate paralogs or introsplice sites

used between the two clustering analyses.

519   **Supplementary Table 1.** Number of gene trees in which all the species for a given

520   bipartition are present. Bipartitions correspond to node labels on Fig. 1.

Bipartition number	Genes containing all species for the bipartition
0	5
1	5
2	246
3	248
4	5
5 (All turtle, crocodilians, and birds)	6
6	248
7	6
8	23
9	36
10	45
11	69
12	51
13	94
Bipartition of turtles sister to birds+crocodilians	36

523 **Supplementary Table 2. Sources of discrepancy between the orthologs detected in**

524 **highly influential nucleotide cluster575 and in matching amino acid homolog**

525 **cluster726.**

Ortholog in 575	Ortholog in 726	Seq length of 575 (Nuc)	Seq length of 726 (Nuc)	Reason for misidentification
Dino@67443 (*Dionaea*)	Dino@67450	2793	2991	Different copy of the in-paralog or intron splice site was retained
Dino@67443 (*Dionaea*)	Dino@9980	2793	510	Not identified as homologs in blast
RuprSFB@17320 (*Ruprechtia*)	RuprSFB@17330	2787	2787	Different copy of the in-paralog or intron splice site was retained
MJM3360@61692 (*Plumbago*)	MJM3360@44226	2211	2403	Different copy of the in-paralog or intron splice site was retained
Retr@34176 (*Reaumuria*)	Retr@1791	1044	546	Not identified as homologs in blast