Running head: LIKELIHOOD AND OUTLIERS IN PHYLOGENOMICS

Title: Analyzing contentious relationships and outlier genes in phylogenomics

Joseph F. Walker[1*], Joseph W. Brown[1], and Stephen A. Smith[1*]

[1]Deptartment Ecology and Evolutionary Biology, University of Michigan, Ann Arbor, Michigan, 48109, USA
*Corresponding authors

Corresponding author emails: jfwalker@umich.edu, eebsmith@umich.edu

18
ABSTRACT

Despite the wealth of evolutionary information available from genomic and

transcriptomic data, recalcitrant relationships in phylogenomic studies remain throughout

the tree of life. Recent studies have demonstrated that conflict is common among gene

trees, and less than one percent of genes may ultimately drive species tree inference in

supermatrix analyses. In this study, we examined plant and vertebrate datasets where

supermatrix and coalescent-based species trees conflict. Using a two-topology site-

specific log-likelihood test, we identified two highly influential genes in each dataset.

While the outlier genes in the vertebrate dataset have been shown to be the result of

errors in orthology detection, we demonstrate that the outlier genes from the plant dataset

may be the result of biological processes rather than model or methodological errors.

When the outlier genes were removed from each supermatrix, the inferred trees matched

the topologies obtained from coalescent analyses. While most tests of this nature limit the

comparison to a small number of fixed topologies, often two topologies, gene tree

topologies generated under processes such as incomplete lineage sorting are unlikely to

precisely match these topologies. We therefore examined edges across a set of trees and

recover more support for the resolution favored by coalescent analyses. These results

suggest that by expanding beyond fixed-topology comparisons, we can dramatically

improve our understanding of the underlying signal in phylogenomic datasets by asking

more targeted edge-based questions.

 INTRODUCTION

Recent studies have highlighted that small changes to a dataset can yield conflicting

hypotheses at particular recalcitrant relationships with high support (i.e., 100% support

from nonparametric bootstrap (BS) or posterior probability (PP) values). Prominent

examples of this include many charismatic lineages such as the root of placental

mammals (Morgan et al. 2013; Romiguier et al. 2013), early branching within Neoaves

(Jarvis et al. 2014; Prum et al. 2015), and the earliest diverging lineage of extant

angiosperms (Zanis et al. 2002; Wickett et al. 2014; Xi et al. 2014). The resolution of

these relationships is critical to understanding the evolutionary history of their respective

clades (e.g., patterns of biochemical, morphological, and life history evolution).

Finding the underlying causes of uncertainty in phylogenomic datasets is an

essential step toward resolving problematic relationships. Recently, authors have

developed means of exploring conflict between gene trees and species trees specifically

for phylogenomic datasets (Salichos et al. 2014; Smith et al. 2015; Kobert et al. 2016),

aiding in the identification of regions of species trees with considerable uncertainty

despite strong statistical support from traditional support measures. Two studies have

shown that the disproportionate influence of just one or two genes "outlier genes" on a

supermatrix analysis is capable of altering tree topology inference (Brown and Thomson

2017; Shen et al. 2017)(Brown and Thomson 2017; Shen et al. 2017)(Brown and

Thomson 2017; Shen et al. 2017). Using a Bayes factor approach Brown and Thomson

(2017) reanalyzed a series of published datasets and found that the transcriptome data

from Chiari et al. (2012) contained outlier genes. When the outlier genes were included in

phylogenetic reconstruction, a clade of turtles+crocodilians was inferred to be sister to

birds with 100% PP. The same topology was previously inferred using ML with

nucleotide data in the original study by Chiari et al. (2012), but was dismissed in favor of

a coalescent reconstruction that placed turtles sister to birds+crocodilians. When Brown

and Thomson (2017) removed the outlier genes, the reduced supermatrix infers the same

topology as the coalescent reconstruction with 100% PP. Another recently published

study compared gene-wise likelihoods across multiple topologies to examine contentious

relationships across the tree of life and found disproportionate influence of genes at all

contentious relationships examined (Shen et al. 2017).

Given the prevalence of outlier genes in phylogenomic datasets, and the continued

focus on contentious relationships in the tree of life, it is imperative that we develop

methods for analyzing conflict and selecting among alternative resolutions for recalcitrant

relationships. We build upon the discussions of Brown and Thomson (2017) and Shen et

al. (2017) by addressing whether these outlier genes violate models of evolution.

Furthermore, we present a method that expands on topology comparisons to instead

pursue edge-based questions. Typically, site-wise and gene-wise log-likelihood analyses

of phylogenomic datasets are performed in a pairwise manner on two or more fixed

alternate topologies (e.g., Castoe et al. 2009; Smith et al. 2011; Shen et al. 2017).

However, given widespread gene tree discordance (e.g., due to incomplete lineage

sorting), it may be more realistic to assume that many alternative topologies are

supported within larger genomic datasets (e.g., Smith et al. 2015; Pease et al. 2016;

Walker et al. 2017). Additionally, when the research question involves a single

relationship and not the entirety of the tree, it may be more appropriate to examine

targeted edges instead of resolved topologies (Lee and Hugall 2003). This allows for any

processes that may be causing conflict in the non-focal parts of the tree to be

accommodated without influencing the relationships of interest. Here, we compare results

from two-topology gene-wise log-likelihood analyses and a novel approach of gene-wise

edge (MGWE) analysis (see Methods below). We examine vertebrate (Chiari et al. 2012;

Brown and Thomson 2017) and carnivorous Caryophyllales datasets (Walker et al. 2017)

(the latter hereafter referred to as the carnivory dataset). Both datasets contain

contentious relationships, outlier genes, and, in their respective original studies, the

authors dismissed the supermatrix topology for the topology inferred using a coalescent

method. In both cases we find that the use of an edge based approach results in stronger

support for the topology hypothesized to be correct by researchers in the original study.

97   METHODS

98   *Data collection*

We obtained the 248 genes that were codon-aligned and analyzed by Brown and

Thomson (2017) from the Dryad deposit (http://dx.doi.org/10.5061/dryad.8gm85) of the

original study (Chiari et al. 2012) that focused on resolving the placement of turtles

among amniotes. The coding DNA sequences of the 1237 one-to-one orthologs from

Walker et al. (2017) to infer the relationships among carnivorous Caryophyllales

(Eudicots: Superasterids) are available from Dryad

(http://datadryad.org/resource/doi:10.5061/dryad.vn730). All programs used in this

analysis may be found at https://bitbucket.org/jfwalker/maximizelikelihoods and the code

to conduct the MGWE analysis may be found at

https://github.com/jfwalker/SiteSpecificLogLikelihood.

Species trees

Brown and Thomson (2017) used Bayesian analyses to obtain the topologies from the

Chiari et al. (2012) data set. As our study focused on the use of maximum likelihood

(ML) for detecting overly influential genes, we ensured that ML phylogenetic

reconstruction would recapitulate the previous species tree results. To construct a

supermatrix tree for the vertebrate dataset, the 248 individual vertebrate genes used in

Brown and Thomson (2017) were concatenated using the Phyx program pxcat (Brown et

al. 2017). The species tree was inferred in RAxML v8.2.1 (Stamatakis 2014) using the

GTR+Γ model of evolution, and edge support was assessed from 200 rapid bootstrap

replicates. Supermatrix trees for the vertebrate dataset were inferred both with all genes

present, and again with the previously identified two outlier genes (8916 and 11434)

removed (see below). The ML tree inferred from all the data from the carnivory dataset

was downloaded from (http://dx.doi.org/10.5061/dryad.33m48) while a novel ML tree

was inferred from a reduced supermatrix that excluded two highly informative genes

(cluster575 and cluster3300; see below).

Gene tree construction and analysis of conflict

Individual gene trees for both datasets were inferred using ML with the GTR+Γ model of

evolution as implemented in RAxML. A SH-like test (Anisimova et al. 2011), as

implemented in RAxML, was performed to assess gene tree edge support. As this test

examines alternative topologies by nearest-neighbor interchange (NNI), it is possible that

during the test a topology with a higher likelihood is found (i.e., an 'NNI-optimal'

topology). When a better topology was found during the test performed for this study,

that topology was used in downstream analyses. We used the pxrr program in the Phyx

package (Brown et al. 2017) to root all gene trees on the outgroup (*Protopterus* for the

vertebrate dataset, and *Beta vulgaris* and *Spinacia oleraceae* for the carnivory dataset)

and we excluded gene trees where an outgroup was not present. We mapped conflict onto

the supermatrix tree using phyparts (Smith et al. 2015) with SH-like support of < 80

treated as uninformative. We chose 80 as a support cutoff due to the traditional cutoff of

(95) being shown as overly conservative with this test (Guindon et al. 2010). Gene tree

conflict was visualized using the script phypartspiecharts.py (available from

https://github.com/mossmatters/MJPythonNotebooks). We conducted more detailed

conflict analyses used for edge comparisons discussed below using pxbp as part of the

Phyx package (Brown et al. 2017).

145   *Calculating two-topology gene-wise log-likelihoods*

The alternate topologies (supermatrix and coalescent) and data matrices for the vertebrate

and carnivory datasets were obtained from the original studies, Chiari et al. (2012) and

Walker et al. (2017), respectively. We calculated site-wise log-likelihood scores for the

two topologies in RAxML using the GTR+ Γ model of evolution, with the data

partitioned by gene. The differences in site-wise log-likelihoods between the candidate

topologies were then calculated using scripts available from

https://bitbucket.org/jfwalker/maximizelikelihoods and

https://github.com/jfwalker/SiteSpecificLogLikelihood.

                    Maximum gene-wise edge calculations

In addition to pairwise topological comparisons, we also examined the maximum

gene-wise edges (MGWE) (Fig 1.). For a single gene and a single focal edge, the MGWE

is the likelihood of a gene tree with the highest likelihood that also displays the edge of

interest. When calculating the MGWE for a focal edge across multiple genes, this

approach does not require each gene to have the same topology, just that the likelihood

comes from a tree that displays the edge of interest. This contrasts with a standard fixed

topology comparison where the topology for each gene would be required to be the same

(e.g., supermatrix vs. coalescent topology). Unlike the fixed topology approach the

MGWE allows for genes to have conflicting relationships outside of the edge of interest.

Here, we are interested in comparing the MGWE for sets of alternative and conflicting

edges in order to determine if, by relaxing the requirement for each gene to share the

topology, we gain insight into the signal for conflicting relationships. One could calculate

the MGWE on any number of edges, and we consider the dominant alternative edges as

identified in the literature.

While there are several ways that MGWEs could be calculated, we restricted the

tree space under under consideration by circumscribing a set of empirically-supported

topologies (TREESET) consisting of the supermatrix-inferred topology, coalescent

inferred topology, and individual gene trees that contained all taxa. We then identified the

conflicting trees and pooled trees based upon shared conflicting relationships for the

edges of interest (EDGE). We then calculated the maximum likelihood for each gene and

for each topology.

For the edges of interest, we calculated the MGWEs by retaining the likelihood

for the tree with the highest likelihood that displayed the focal EDGE. This became the

representative likelihood for that EDGE. We then summed the representative likelihoods

together. That value, however, is not comparable between edges because a different

number of trees may be compared (Theobald 2010). Therefore, we calculated AIC scores

(-2ln(L) + 2k) for each EDGE. This, effectively, allowed for comparisons between more

parameter rich models and parameter poor models. The parameters, k, were calculated

based on the number of taxa, n, and the number of genes in the analysis, g. The branch

length parameters equal $2 \times n - 3$ and the GTR $+ \Gamma$ model of evolution $= 6$ (where base

frequencies were empirical and not estimated). The supermatrix ML analyses that

assumed a single set of branch lengths on one topology and model parameters to be

unlinked across genes consisted of $2 \times n - 3 + 6 \times g$ parameters. For each EDGE, because

branch lengths were calculated for each gene tree, the parameters consisted of the sum of

the number of parameters used for each gene: g x (2 x n – 3 + 6). In addition to

calculating AICs for the coalescent and supermatrix topologies with a single set of branch

lengths across the gene set, we also calculated AICs allowing the branch lengths to vary

across genes. This calculation results in the same number of parameters as the EDGE

calculations. Here, we are focused on addressing conflicting signal between edges of

interest and so the increase in the number of parameters is acceptable considering our

examination of gene trees. However, future work could attempt to limit the expansion of

the number of parameters for each EDGE by sharing branch length estimates or model

parameters across genes.

 Testing for paralogy in carnivory dataset

The homolog trees created from amino acid data in the study by Walker et al. (2017)

were downloaded from Dryad (http://datadryad.org/resource/doi:10.5061/dryad.vn730).

We matched the sequences from the outlier genes to their corresponding sequence in the

amino acid homolog trees. This allowed us to examine whether a nucleotide cluster

contained homology errors that may be exposed by the slower evolving amino acid

dataset.

208   RESULTS

Gene tree conflict and log-likelihood analysis reveals genes of disproportionate influence

Our ML analysis of the vertebrate dataset recovered the same supermatrix topology (Fig.

2) as found with ML by Chiari et al. (2012) and Bayesian inference by Brown and

Thomson (2017). The difference in log-likelihood between the supermatrix and

coalescent topologies for the vertebrate dataset was 4.01. Ninety-three of 248 gene trees

could be rooted on the outgroup *Protopterus* and only five of these had all taxa

represented (Supplementary Table 1). We found low support for relationships within

gene trees (SH <80) and significant gene tree conflict (Fig. 2). Of the gene trees with high

support (SH >80), seven resolved turtles+crocodilians as sister to birds (hereafter referred

to as the vertebrate supermatrix topology) and nine resolved crocodilians+birds sister to

turtles (hereafter referred to as the vertebrate coalescent topology).

The two-topology gene-wise log-likelihood comparison showed that 105 genes

had a higher likelihood score for the vertebrate supermatrix topology while 143 supported

the vertebrate coalescent topology (Figs. 3A, 4A). Two genes (ENSGALG00000008916

and ENSGALG00000011434, referred to here as 8916 and 11434, respectively),

appeared as outliers, exhibiting a disproportionate influence on the overall likelihood of

the supermatrix (Fig. 3A). The outlier genes identified with maximum likelihood

analyses matched those previously identified as outliers using Bayes factors (Brown and

Thomson 2017). These two genes both supported the vertebrate supermatrix topology

with log-likelihood scores of 79.55 and 46.01 greater than the alternative coalescent tree

topology, respectively. The difference in log-likelihood between the two topologies of the

non-outlier genes ranged from |0.006| to |19.891| with an average of 3.31 for all genes in

the analysis. The removal of the vertebrate genes 8916 and 11434, as shown by Brown

and Thomson (2017), recovered the coalescent topology, albeit with low bootstrap

support (BS = 12; Supplementary Fig. 1).

Previous work on the carnivory dataset demonstrated that the placement of the

Ancistrocladus+Drosophyllum clade (Fig. 2) contained significant conflict and is

strongly influenced by species sampling (Walker et al. 2017). The log-likelihood

difference between the supermatrix and coalescent topologies was 74.94 in favor of the

former. The two-topology log-likelihood comparison between the dominant topologies on

the carnivory dataset (Fig. 3B) showed that 623 genes supported

Ancistrocladus+Drosophyllum sister to all other carnivorous plants (hereafter referred to

as carnivory supermatrix topology) while 614 genes supported

Ancistrocladus+Drosophyllum sister to *Nepenthes alata+Nepenthes ampullaria*

(hereafter referred to as carnivory coalescent topology; Figs. 3A & 4D). Two genes

(cluster575 and cluster3300) contributed disproportionately to the overall likelihood.

Individually these two genes have a difference in log-likelihood scores between the two

topologies of 33.06 and 16.63, respectively, and support the carnivory supermatrix

topology. When we reanalyzed the supermatrix with cluster575 and cluster3300 removed,

the carnivory coalescent topology was recovered, with 100% BS support (Supplementary

Fig. 1). The difference between the two topologies in log-likelihood of the non-outlier

genes ranged from |0.001| to |12.82| with an average of 2.82 for all genes in the analysis.

252   *Edge based analysis changes supported topology*

We compared MGWE and two topology gene-wise likelihoods involving the contentious

bird, crocodilian, and turtle relationships in the vertebrate dataset (Fig. 4B). We found

seven unique topologies with the necessary species coverage to conduct the analyses: five

gene tree topologies from Chiari et al. (2012) and the two dominant species tree

topologies. The set of seven trees included three major conflicting edges for the

relationship in question: the two resolutions found in the supermatrix and coalescent trees,

and birds sister to crocodilian+mammals+turtles. 91 genes supported the vertebrate

supermatrix edge, 144 genes supported the vertebrate coalescent edge, and 13 genes

supported the third conflicting edge (Fig. 4B). When comparing the supermatrix analysis

with a single set of branch lengths, to that where it was treated as a sum of gene tree

likelihoods, we found a superior AIC score for the sum of gene tree likelihoods (Table 1).

The MGWE AIC scores for the summed likelihoods of the supermatrix (three source

trees), the coalescent (three source trees), and the third conflicting edge (one source tree)

were highest for the coalescent edge and out of all tested models the coalescent edge was

inferred to be the best (Table 1).

For the carnivory dataset, we found 168 unique tree topologies to include in the

tree set. The 168 tree topologies contained 41 conflicting edges for the relationship in

question with 3 dominant edges. The MGWE analyses found 499 genes supported the

supermatrix edge, 466 genes supported the coalescent edge, and 272 genes supported 15

additional edges (Figs. 2D, 3E). When we further compared the MGWE AIC scores for

the supermatrix (44 source trees), the coalescent (56 source trees), and for the third edge

(24 source trees) we found the coalescent edge to have the best AIC score out of all tested

models (Table 1).

277   *Outlier gene examination*

For the carnivory dataset, we explored the possibility that the strongly conflicting genes

cluster575 and cluster3300 reflected methodological error in the assembly pipeline, as is

the case for the genes identified by Brown and Thomson (2017) for the vertebrate dataset.

However, both the alignment and inferred phylogram for each gene revealed no obvious

problems or potential sources of systematic error (sparse alignment, abnormally long

branch lengths, etc.). We also explored whether compositional heterogeneity could

explain the strongly conflicting results (i.e., that the relationships were not truly

conflicting, but instead incorrectly modeled). However, both RY-coding in RAxML and

explicit modeling of multiple equilibrium frequencies (2, 3, or 4 composition regimes)

across the tree in p4 v1.0 (Foster 2004) failed to overturn the inferred relationships. We

further explored the possibility of misidentified orthology. By examining the homolog

tree produced from amino acid data, we identified the ortholog from the nucleotide data

to be complete (i.e., an ortholog within the homolog amino acid tree). We found that with

the slower amino acid data the sequences in the nucleotide cluster575 were inferred as a

single monophyletic ortholog within a duplicated homolog (Supplementary Fig. 2). The

discrepancies that appeared between the amino acid dataset and the CDS dataset were

found to be either different in-paralogs/splice sites maintained during the dataset cleaning

procedure or short sequences that were not identified as homologs in the coding DNA

sequence (CDS) dataset (Supplementary Table 2 and Supplementary Fig. 2).

298   DISCUSSION

Biological processes including substitution saturation, hybridization, horizontal gene

transfer, and incomplete lineage sorting can contribute to conflicting signal and may

explain both conflict and lack of support widely found in phylogenomic datasets

(Salichos et al. 2014; Smith et al. 2015; Kobert et al. 2016). In addition to these

biological processes, other data set assembly issues such as limited taxonomic coverage

for each gene may also contribute to conflict and low support in these data sets. For

example, while the carnivory dataset had extensive data overlap, the vertebrate dataset

only had five gene regions that contained sequence data for every species (Supplementary

Table 1). To further complicate the challenges facing phylogenomic analyses, high

support values, especially from concatenated runs, can mask significant underlying

conflict (Lee and Hugall, 2003; Ryan et al. 2013; Salichos et al. 2014; Smith et al. 2015;

Kobert et al. 2016; Pease et al. 2017). Both datasets examined here recovered high

support for two different topologies depending on the inclusion or exclusion of two genes

with disproportionate influence on the likelihood (Brown and Thomson 2017; Walker et

al. 2017). In the case of the carnivory dataset, the inferred topology changes with the

inclusion or exclusion of just 0.0016% of the genes.

To address these challenges, several approaches have been outlined in the

literature. Recently, the discovery of outlier genes has resulted in the necessity to closely

examine gene tree topologies and likelihoods (Brown and Thomson 2017; Shen et al.

2017). Outlier genes may be the result of biological processes or methodological errors,

and due to their high influence of species tree inference should be thoroughly examined.

Previously, the outlier genes in a vertebrate dataset were found to be the result of errors

in orthology detection and not biological processes (Brown and Thomson 2017). We

explored, in this study, the potential sources of error for the outlier genes in a dataset of

carnivorous plants. While the genomic resources are not available to fully examine the

carnivorous outlier genes (e.g., we do not yet have synteny or information on gene loss),

our analyses did not detect any obvious problems with alignment, compositional

heterogeneity, or homology. We found one gene, cluster575, to be an ortholog of a gene

that experienced a duplication event prior to the divergence of both ingroup and outgroup

taxa (Supplementary Fig. 3). While we cannot rule out every possible source of error, we

also cannot identify a source of methodological error, suggesting the possibility that the

conflicting topology is the result of real (albeit unknown) biological processes.

Fixed topological and pairwise examinations explored by most authors (Castoe et

al. 2009; Smith et al. 2011; Shen et al. 2017), have been very informative for the

identification of not only outlier genes, but also for phylogenetic signal for and against

conflicting phylogenetic relationships. However, for many reasons, these fixed

topological examinations, where a single topology is assumed to underlie all genes, may

not be optimal. Conflict among gene trees is common and expected from processes such

as incomplete lineage sorting, hybridization, and other processes. For instance, Jarvis et

al. (2014) reported that no gene trees from a genomic data set of 48 species of birds

matched the inferred species tree. Furthermore, such a result becomes increasingly likely

as sampling breadth (both taxa within a clade as well as the age of the clade itself)

increases. The results of a fixed-topology analysis may be driven by the resolution of a

part of the phylogeny other than the area of interest, as fixed-topology analyses condition

on fully bifurcating trees that necessarily resolve conflict in the entire tree.

To overcome these limitations, instead of fixed singular topologies, we examined

edges across a set of empirically supported candidate topologies, as defined by the set of

inferred gene trees and the two tree hypotheses in question. By examining edges, we

accommodate for uncertainty across the rest of the tree, regardless of the process

generating that uncertainty. We examined this with both a vertebrate dataset and

carnivorous plants dataset discussed above. The vertebrate dataset contained three

alternative edges for the relationship of interest while the carnivory dataset contained 41

different edges representing 168 topologies. The MGWE analysis and AIC scores of both

the vertebrate dataset and the carnivory dataset both suggested a better fit of the

coalescent edge than the supermatrix edge (Table 1). Also, in both cases, we found that

the AIC score supported the higher parameterized model, as opposed to a single fixed

topology and branch lengths. While we do not suggest that this is the best fit model and

only the best of the ones analyzed here, this indicates that future studies may benefit from

allowing more heterogeneity than is typically involved in a concatenation analysis. This

will require careful examination of some of the complexity involved in these large

phylogenomic analyses. For example, there is the issue of how missing data is handled in

these calculations (e.g.,Stamatakis and Alachiotis 2010). Furthermore, the models

explored could potentially have significantly reduced parameters by sharing topologies

and branch lengths across some gene regions, including potentially scaling branch lengths

proportionally (e.g., as is possible with the -spp option in the program iqtree).

Nevertheless, the exploratory analyses presented here provide additional evidence that a

simple concatenation approach with these large datasets masks important heterogeneity

that can be analyzed further to help inform phylogenetic resolution.

367   The results presented here contribute to a growing body of literature that address

the question of how phylogenomic analyses should proceed in the presence of highly

influential outlier genes, conflicting topologies, and ever expanding datasets (Wickett et

al. 2014; Pease et al. 2016; Brown and Thomson 2017; Shen et al. 2017; Yang et al.

2017). For example, some authors have noted, and it is the case here, that supermatrix

analyses may be more susceptible to the problem of strong outliers (Shen et al. 2017;

Walker et al. 2017). In these studies, the resolutions inferred using a coalescent method

were generally favored. When the dominant process generating gene tree conflict is

incomplete lineage sorting, coalescent methods should perform better (i.e., when gene

tree diversity is modeled correctly). Some coalescent methods that weigh all gene tree

equally (e.g., Mirarab and Warnow 2015), may overcome the problem of outlier genes

even if incomplete lineage sorting is not the dominant source of conflict simply by

eliminating the disproportionate influence of one or two outlying genes. Here, we

demonstrate with two empirical examples that the coalescent resolution had higher

support when examining edges without using an explicit coalescent method.

While we continue to uncover the patterns and processes that generate conflicting

signal within phylogenomic datasets, it is imperative that we explore new methods that

accommodate conflict. Phylogenomic studies often focus sampling efforts around

particularly recalcitrant nodes, and it is important we develop methods designed with the

same purpose. Here we focus on conflicting edges and explore the MGWE method as a

means of analyzing these conflicting edges while allowing for topological heterogeneity

outside of the relationships of interest. This approach helps accommodate the biological

realities of heterogeneity among lineages, conflicting signal both for in and outside the

relationship of interest, and evolutionary processes that violate assumptions by

supermatrix and coalescent models. This approach, however, is just a start and future

research should examine how to better incorporate the underlying heterogeneity that has

emerged from our large data sets over the last few years. We believe further investigation

into edge based testing is warranted to better understand how we may incorporate the

process based conflict of phylogenomics into our analyses.

399   FUNDING

400   JFW was supported through a fellowship provided by University of Michigan department

401   of Ecology and Evolutionary Biology. JFW and SAS were supported by NFS 1354048

402   and JWB and SAS were supported by NSF 1207915.

404   ACKNOWLEDGEMENTS

We would like to thank Alexis Stamatakis for advice on RAxML and Nathanael Walker-

Hale, Javier Igea, Oscar Vargas, Greg Stull, and Ning Wang for discussion of the

manuscript and methods.

REFERENCES

Anisimova M., Gil M., Dufayard J.F., Dessimoz C., Gascuel O. 2011. Survey of branch

support methods demonstrates accuracy, power, and robustness of fast likelihood-

based approximation schemes. Syst. Biol. 60:685–699.

Brown J.M., Thomson R.C. 2017. Bayes Factors Unmask Highly Variable Information

Content , Bias , and Extreme Influence in Phylogenomic Analyses. Syst. Biol.

66:517–530.

Brown J.W., Walker J.F., Smith S.A. 2017. Phyx: phylogenetic tools for unix.

Bioinformatics. 33:1886–1888.

Castoe T.A., de Koning A.P.J., Kim H.-M., Gu W., Noonan B.P., Naylor G., Jiang Z.J.,

Parkinson C.L., Pollock D.D. 2009. Evidence for an ancient adaptive episode of

convergent molecular evolution. Proc. Natl. Acad. Sci. 106:8986–8991.

Chiari Y., Cahais V., Galtier N., Delsuc F. 2012. Phylogenomic analyses support the

position of turtles as the sister group of birds and crocodiles (Archosauria). BMC

Biol. 10:65.

Foster P.G. 2004. Modeling compositional heterogeneity. Syst Biol. 53:485–495.

Guindon S., Dufayard J.F., Lefort V., Anisimova M., Hordijk W., Gascuel O. 2010. New

algorithms and methods to estimate maximum-likelihood phylogenies: Assessing the

performance of PhyML 3.0. Syst. Biol. 59:307–321.

Jarvis E.D., Mirarab S., Aberer A.J., Li B., Houde P., Li C., Ho S.Y.W., Faircloth B.C.,

Nabholz B., Howard J.T., Suh A., Weber C.C., da Fonseca R.R., Li J., Zhang F., Li

H., Zhou L., Narula N., Liu L., Ganapathy G., Boussau B., Bayzid M.S.,

Zavidovych V., Subramanian S., Gabaldon T., Capella-Gutierrez S., Huerta-Cepas J.,

Rekepalli B., Munch K., Schierup M., Lindow B., Warren W.C., Ray D., Green R.E.,

Bruford M.W., Zhan X., Dixon A., Li S., Li N., Huang Y., Derryberry E.P.,

Bertelsen M.F., Sheldon F.H., Brumfield R.T., Mello C. V., Lovell P. V., Wirthlin

M., Schneider M.P.C., Prosdocimi F., Samaniego J.A., Velazquez A.M. V., Alfaro-

Nunez A., Campos P.F., Petersen B., Sicheritz-Ponten T., Pas A., Bailey T., Scofield

P., Bunce M., Lambert D.M., Zhou Q., Perelman P., Driskell A.C., Shapiro B.,

Xiong Z., Zeng Y., Liu S., Li Z., Liu B., Wu K., Xiao J., Yinqi X., Zheng Q., Zhang

Y., Yang H., Wang J., Smeds L., Rheindt F.E., Braun M., Fjeldsa J., Orlando L.,

Barker F.K., Jonsson K.A., Johnson W., Koepfli K.-P., O'Brien S., Haussler D.,

Ryder O.A., Rahbek C., Willerslev E., Graves G.R., Glenn T.C., McCormack J.,

Burt D., Ellegren H., Alstrom P., Edwards S. V., Stamatakis A., Mindell D.P.,

Cracraft J., Braun E.L., Warnow T., Jun W., Gilbert M.T.P., Zhang G. 2014. Whole-

genome analyses resolve early branches in the tree of life of modern birds. Science

(80-.). 346:1320–1331.

Kobert K., Salichos L., Rokas A., Stamatakis A. 2016. Computing the internode certainty

and related measures from partial gene trees. Mol. Biol. Evol. Advance Ac:1–17.

Lee M.S.Y., Hugall A.F. 2003. Partitioned Likelihood Support and the Evaluation of

Data Set Conflict. Syst. Biol. 52:15–22.

Mirarab S., Warnow T. 2015. ASTRAL-II: Coalescent-based species tree estimation with

many hundreds of taxa and thousands of genes. Bioinformatics. 31:i44–i52.

Morgan C.C., Foster P.G., Webb A.E., Pisani D., McInerney J.O., O'Connell M.J. 2013.

Heterogeneous models place the root of the placental mammal phylogeny. Mol. Biol.

Evol. 30:2145–56.

Pease J.B., Brown J.W., Walker J.F., Hinchliff C.E., Smith S.A. 2017. Quartet Sampling

distinguishes lack of support from conflicting support in the plant tree of life.

BioRxiv.

Pease J.B., Haak D.C., Hahn M.W., Moyle L.C. 2016. Phylogenomics Reveals Three

Sources of Adaptive Variation during a Rapid Radiation. PLoS Biol. 14:1–24.

Prum R.O., Berv J.S., Dornburg A., Field D.J., Townsend J.P., Moriarty Lemmon E.,

Lemmon A.R. 2015. A comprehensive phylogeny of birds (Aves) using targeted

next-generation DNA sequencing. Nature. 526:569–573.

Romiguier J., Ranwez V., Delsuc F., Galtier N., Douzery E.J.P. 2013. Less is more in

mammalian phylogenomics: AT-rich genes minimize tree conflicts and unravel the

root of placental mammals. Mol. Biol. Evol. 30:2134–44.

Ryan J.F., Pang K., Schnitzler C.E., Nguyen A.D., Moreland R.T., Simmons D.K., Koch

B.J., Francis W.R., Havlak P., Smith S.A., Putnam N.H., Haddock S.H., Dunn C.W.,

Wolfsberg T.G., Mullikin J.C., Martindale M.Q., Baxevanis A.D. 2013. The genome

of the ctenophore Mnemiopsis leidyi and its implications for cell type evolution.

Science (80-.). 342:1242592.

Salichos L., Stamatakis A., Rokas A. 2014. Novel information theory-based measures for

quantifying incongruence among phylogenetic trees. Mol. Biol. Evol. 31:1261–1271.

Shen X., Hittinger C.T., Rokas A. 2017. Contentious relationships in phylogenomic

studies can be driven by a handful of genes. Nat. Ecol. Evol. 1:1–10.

Smith S.A., Moore M.J., Brown J.W., Yang Y. 2015. Analysis of phylogenomic datasets

reveals conflict, concordance, and gene duplications with examples from animals

and plants. BMC Evol. Biol. 15:150.

Smith S.A., Wilson N.G., Goetz F.E., Feehery C., Andrade S.C.S., Rouse G.W., Giribet

G., Dunn C.W. 2011. Resolving the evolutionary relationships of molluscs with

phylogenomic tools. Nature. 480:364–367.

Stamatakis A. 2014. RAxML version 8: A tool for phylogenetic analysis and post-

analysis of large phylogenies. Bioinformatics. 30:1312–1313.

Stamatakis A., Alachiotis N. 2010. Time and memory efficient likelihood-based tree

searches on phylogenomic alignments with missing data. Bioinformatics. 26:132–

139.

Theobald D.L. 2010. A formal test of the theory of universal common ancestry. Nature.

465:219–222.

Walker J.F., Yang Y., Moore M.J., Mikenas J., Timoneda A., Brockington S.F., Smith

S.A. 2017. Widespread paleopolyploidy , gene tree conflict , and recalcitrant

relationships among the. Am. J. Bot. 104:858–867.

Wickett N.J., Mirarab S., Nguyen N., Warnow T., Carpenter E., Matasci N.,

Ayyampalayam S., Barker M.S., Burleigh J.G., Gitzendanner M.A., Ruhfel B.R.,

Wafula E., Der J.P., Graham S.W., Mathews S., Melkonian M., Soltis D.E., Soltis

P.S., Miles N.W., Rothfels C.J., Pokorny L., Shaw A.J., DeGironimo L., Stevenson

D.W., Surek B., Villarreal J.C., Roure B., Philippe H., DePamphilis C.W., Chen T.,

497 Deyholos M.K., Baucom R.S., Kutchan T.M., Augustin M.M., Wang J., Zhang Y.,

498 Tian Z., Yan Z., Wu X., Sun X., Wong G.K.-S., Leebens-Mack J. 2014.

499 Phylotranscriptomic analysis of the origin and early diversification of land plants.

500 Proc. Natl. Acad. Sci. 111:E4859–E4868.

501 Xi Z., Liu L., Rest J.S., Davis C.C. 2014. Coalescent versus Concatenation Methods and

502 the Placement of Amborella as Sister to Water Lilies. Syst. Biol. 63:919–932.

503 Yang Y., Moore M.J., Brockington S.F., Mikenas J., Olivieri J., Walker J.F., Smith S.A.

504 2017. Improved transcriptome sampling pinpoints 26 paleopolyploidy events in

505 Caryophyllales, including two paleo-allopolyploidy events. bioRxiv.

506 Zanis M.J., Soltis D.E., Soltis P.S., Mathews S., Donoghue M.J. 2002. The root of the

507 angiosperms revisited. Proc. Natl. Acad. Sci. U. S. A. 99:6848–53.

Tree set

MGWE

Edge	Gene 1	Gene 2	Gene 3	Gene 4	Gene 5
ABC | DE	-20.00 (t1)	-10.00 (t4)	-11.00 (t2)	-14.00 (t1)	-9.00 (t1)
ADC | BE	-90.00 (t3)	-5.00 (t5)	-4.00 (t5)	-50.00 (t3)	-20.00 (t5)
AEC | BD	-15.00 (t6)	-11.00 (t7)	-7.00 (t6)	-8.00 (t7)	-10.00 (t6)

514 **Figure 1. Outline for the MGWE procedure.** The inferred tree set is depicted at the top,

515 with the tree number in the top right hand corner of each box, and the edge representing

516 the relationship of interest in the bottom left hand corner. The MGWE shows the best

517 likelihood for each edge at each gene, with the tree from which that likelihood was

518 obtained in the box in parentheses next to the likelihood score.

Figure 2. Maximum likelihood trees inferred by RAxML for the Chiari et al. 2012

(vertebrate) and Walker et al. 2017 (carnivorous Caryophyllales) datasets. Conflict

analysis for the vertebrate (A) and carnivory (B) datasets. The vertebrate analysis

includes the 93 genes that contained the outgroup (*Protopterus*), and the carnivory

analysis includes 1237 genes all of which had the outgroups (*Spinacia oleraceae* and

Beta vulgaris). Blue represents gene trees that are concordant with the relationship, grey

represents uninformative genes (SH-like < 80 or no taxon representation for the edge),

532 green represents the dominant alternate topology, and red represents all other conflict.

533 Numbers on edges represent concordance/conflict. Bold numbers at the nodes of the

534 vertebrate dataset correspond to edge numbers in Supplementary Table 1.

542 **Figure 3. Identification of outlier genes using gene-wise likelihood test.** A&B) Show

543 the results of the two-topology test on the vertebrate and carnivory dataset, respectively,

544 using the coalescent (negative values) and supermatrix (positive values) topologies as the

545 comparison. The genes identified as outliers from the analysis are marked with an X.

Figure 4. Bar plot representing gene counts for the two-topology and MGWE methods. A&C) represent counts of genes that support the supermatrix inferred maximum likelihood (ML) topology and the maximum quartet support species tree (MQSST), for the vertebrate and carnivory datasets respectively. B&D) Show the results of the MGWB analysis for support of the edge found in the ML analysis, the conflicting edge from the MQSST analysis, and the sum of all genes supporting an alternative conflict from an edge in the TREE SET.

Table 1. Results of model testing the various topologies and edges.

Vertebrate	Supermatrix	Topology	-1,047,406.05	1517	2,097,846.11	22374.01
		As Gene Trees	-1,031,489.81	6442	2,075,863.63	391.53
		Edge	-1,031,423.65	6442	2,075,731.31	259.20
	Coalescent	Topology	-1,047,410.07	1517	2,097,854.15	22382.04
		As Gene Trees	-1,031,450.71	6442	2,075,785.43	313.32
		Edge	**-1,031,294.05**	**6442**	**2,075,472.10**	**0**
	Dominant Alternative	Edge	-1,033,773.81	6442	2,080,431.62	4959.52
Carnivory	Supermatrix	Topology	-13,305,055.20	7445	26,625,000.40	36618.41
		As Gene Trees	-13,205,130.14	35873	26,595,640.58	7258.59
		Edge	-13,258,387.61	35873	26,588,521.23	139.24
	Coalescent	Topology	-13,305,130.14	7445	26,625,150.28	36768.28
		As Gene Trees	-13,262,019.55	35873	26,595,785.10	7403.10
		Edge	**-13,258,317.99**	**35873**	**26,588,381.99**	**0**
	Dominant Alternative	Edge	-13,260,106.83	35873	26,591,959.66	3577.67

564 *In the type column, "Topology" represents the supermatrix or coalescent topology with a single set
565 of branch lengths, "As Gene Trees" is the supermatrix or coalescent topology with branch lengths
566 varying among genes, and "Edge" is the MGWE analysis. The top AIC score is bolded.
567
568

Supplementary Figure 1. Species trees inferred using maximum likelihood from the different supermatrices. Support at each node was obtained from 200 rapid bootstrap replicates. A) Species tree for vertebrate dataset inferred with all 248 genes included in the supermatrix. B) Species tree for the vertebrate dataset inferred with 8916 and 11434 removed from the supermatrix. C) carnivorous Caryophyllales species tree inferred from

579   all 1237 genes. D) carnivorous Caryophyllales species tree inferred with cluster575 and

580   cluster3300 removed from the supermatrix.

Supplementary Figure 2. Homolog tree for Amino Acid clustered (726) and CDS

clustered (575) highly influential gene in the carnivorous Caryophyllales dataset.

Different genes identified in the ortholog clusters are circled on cluster 726. Genes

circled in red represent ones that are shorter and were not identified as orthologous in the

CDS dataset and genes circled in blue represent alternate paralogs or introsplice sites

used between the two clustering analyses.

Supplementary Table 1. Number of gene trees in which all the species for a given edges

are present. edges correspond to node labels on Fig. 1.

Edge number	Genes containing all species for the edge
0	5
1	5
2	246
3	248
4	5
5 (All turtle, crocodilians, and birds)	6
6	248
7	6
8	23
9	36
10	45
11	69
12	51
13	94
edge of turtles sister to birds+crocodilians	36

597 **Supplementary Table 2. Sources of discrepancy between the orthologs detected in**

598 **highly influential nucleotide cluster575 and in matching amino acid homolog**

599 **cluster726.**

Ortholog in 575	Ortholog in 726	Seq length of 575 (Nuc)	Seq length of 726 (Nuc)	Reason for misidentification
Dino@67443 (*Dionaea*)	Dino@67450	2793	2991	Different copy of the in-paralog or intron splice site was retained
Dino@67443 (*Dionaea*)	Dino@9980	2793	510	Not identified as homologs in blast
RuprSFB@17320 (*Ruprechtia*)	RuprSFB@17330	2787	2787	Different copy of the in-paralog or intron splice site was retained
MJM3360@61692 (*Plumbago*)	MJM3360@44226	2211	2403	Different copy of the in-paralog or intron splice site was retained
Retr@34176 (*Reaumuria*)	Retr@1791	1044	546	Not identified as homologs in blast