1 **Deconvolution of DNA methylation identifies differentially methylated gene**
2 **regions on 1p36 across breast cancer subtypes**
3

4 Alexander J. Titus*[a,d], Gregory P. Way*[b], Kevin C. Johnson[c,d], and Brock C.
5 Christensen^[d,e,f]
6
*Authors contributed equally
[a] Program in Quantitative Biomedical Sciences, Geisel School of Medicine at Dartmouth,
Hanover, NH 03755
[b] Genomics and Computational Biology Graduate Program, University of Pennsylvania,
Philadelphia, PA 19104
[c] The Jackson Laboratory for Genomic Medicine, Farmington, CT 06032, USA
[d] Department of Epidemiology,
[e] Department of Molecular and Systems Biology
[f] Department of Community and Family Medicine, Geisel School of Medicine at
Dartmouth, Hanover, NH 03755
17
18 CORRESPONDING AUTHOR:
19 ^ 1 Medical Center Drive, Williamson Level 6, HB7650, Lebanon, NH 03766; Phone:
20 603-650-1828; Fax: 603-650-1840 Brock.Christensen@Dartmouth.edu
21
AUTHOR EMAILS:
AT: Alexander.J.Titus.gr@dartmouth.edu
GW: GregWay@upenn.edu
KJ: Kevin.C.Johnson@jax.org
BC: Brock.Christensen@dartmouth.edu

ABSTRACT:

Breast cancer is a complex disease and studying DNA methylation (DNAm) in tumors is

complicated by disease heterogeneity. We compared DNAm in breast tumors with

normal-adjacent breast samples from The Cancer Genome Atlas (TCGA). We

constructed models stratified by tumor stage and PAM50 molecular subtype and

performed cell-type reference-free deconvolution on each model. We identified nineteen

differentially methylated gene regions (DMGRs) in early stage tumors across eleven

genes (*AGRN, C1orf170, FAM41C, FLJ39609, HES4, ISG15, KLHL17, NOC2L,*

PLEKHN1, SAMD11, WASH5P). These regions were consistently differentially

methylated in every subtype and all implicated genes are localized on chromosome

1p36.3. We also validated seventeen DMGRs in an independent data set. Identification

and validation of shared DNAm alterations across tumor subtypes in early stage tumors

advances our understanding of common biology underlying breast carcinogenesis and

may contribute to biomarker development. We also provide evidence on the importance

and potential function of 1p36 in cancer.

INTRODUCTION:

Invasive breast cancer is a complex disease characterized by diverse etiologic factors[1]. Key genetic and epigenetic alterations are recognized to drive tumorigenesis and serve as gate-keeping events for disease progression[2]. Early DNA methylation (DNAm) events have been shown to contribute to breast cancer development[3]. Importantly, DNAm alterations have been implicated in the transition from normal tissue to neoplasia[4,5] and from neoplasia to metastasis[6]. Furthermore, patterns of DNAm are known to differ across molecular subtypes of breast cancer[7] - Luminal A (LumA), Luminal B (LumB), Her2-enriched and Basal-like – identified based on the prediction analysis of microarray 50 (PAM50) classification[8]. However, while DNAm differences across breast cancer subtypes have been explored, similarities across subtypes are less clear[9]. Such similarities found in early stage tumors can inform shared biology underpinning breast carcinogenesis and – as similarities would be agnostic to subtype – potentially contribute to biomarkers for early detection.

Studying DNAm in bulk tumors is complicated by disease heterogeneity. Heterogeneity is driven by many aspects of cancer biology including variable cell-type proportions found in the substrate used for molecular profiling[10]. Different proportions of stromal, tumor, and infiltrating immune cells may confound molecular profile classification when comparing samples[11] because cell types have distinct DNAm patterns[12–14]. The potential for cell–type confounding prompted the development of statistical methods to adjust for variation in cell-type proportions in blood[15] and solid tissue[16]. One such method, *RefFreeEWAS*, is a reference-free deconvolution method and does not require a reference population of cells with known methylation patterns and is agnostic to genomic location when performing deconvolution[17]. Instead, the unsupervised

method infers underlying cell-specific methylation profiles through constrained non-

negative matrix factorization (NMF) to separate cell-specific methylation differences

from actual aberrant methylation profiles observed in disease states. This method has

previously been shown to effectively determine the cell of origin in breast tumor

phenotypes[18].

We applied *RefFreeEWAS* to The Cancer Genome Atlas (TCGA) breast cancer

DNAm data and estimated cell proportions across the set. We compared tumor DNAm

with adjacent normal tissue stratified by tumor subtype[9] and identified common early

methylation alterations across molecular subtypes that are independent of cell type

composition. We identified a specific chromosomal location, 1p36.3, that harbors all 19

of the differentially methylated regions that are in common to early stage breast cancer

subtypes. 1p36 is a well-studied and important region in many different cancer types, but

there remain questions about how it may impact carcinogenesis and disease

progression[19]. Our study provides evidence that methylation in this region may provide

important clues about early events in breast cancer. We also performed *RefFreeEWAS* on

an independent validation set (GSE61805) and confirmed these results[20].

RESULTS:

DNA methylation deconvolution

Subject age and tumor characteristic data stratified by PAM50 subtype and stage

is provided in Table 1 for the 523 TCGA tumors analyzed. TCGA breast tumor sample

purity, estimated by pathologists from histological slides, was consistent across PAM50

subtypes and stages indicating that observed methylation differences are not

predominantly a result of large differences in tumor purity (Supplementary Fig. S1). To

correct for cell-proportion differences across tumor samples, we estimated the number of

cellular methylation profiles contributing to the mixture differences by applying NMF to

the matrix of beta values, which resulted in model specific dimensionality estimates

indicating diverse cellular methylation profiles (Supplementary Table S1). The reference-

free deconvolution altered the number and extent of significant differentially methylated

CpGs across all models that compared breast tumor methylation with adjacent normal

samples (Supplementary Fig. S2).

Table 1. Sample information stratified by PAM50 subtype

	Basal-like	Her2	Luminal A	Luminal B	Total with Assignment
TCGA tumors	86	31	279	127	523
Age, mean (SD)	56.8 (12.8)	60 (12.8)	58 (13.5)	57.1 (12.6)	57.8 (13.1)
Stage*, n (%)	--	--	--	--	--
Early (I/II)	70 (81%)	20 (65%)	207 (74%)	84 (66%)	381 (73%)
Late (III/IV)	14 (16%)	10 (32%)	69 (25%)	42 (33%)	135 (26%)
Missing	2 (2%)	1 (3%)	3 (1%)	1 (1%)	7 (1%)

*AJCC characterized stage, provided by TCGA

98   *Subtype specific methylation patterns*

99   In early stage tumors, we identified a set of nineteen DMGRs shared among

100   Luminal A, Luminal B, Her2, and Basal-like subtypes (DMGRs $Q < 0.01$, Figure 1A). In

101   the late stage tumors, we identified 31,931 DMGRs in common across subtypes (Figure

102   1B).

Figure 1. Numbers of overlapping differentially methylated gene regions in (A) early

stage tumors (n = 76,847) and (B) late stage tumors (n = 70,759) stratified by Basal-like,

Her2, Luminal A, and Luminal B PAM50 subtypes with a Q-value cutoff of 0.01.

Subtype specific methylation patterns in early stage tumors were most divergent for

Basal-like tumors versus other types, while in late stage tumors methylation alterations in

Luminal B tumors were most divergent (Supplementary Table S2). To test if collapsing

by genomic region had an appreciable effect on detecting DMGRs, we compared DMGR

results to results derived from regions defined by CpG island status (i.e. CpG island,

Shore, Shelf, Open Sea). Using CpG island context designations indicated similar results

(Supplementary Fig S3), though a lower number of common DMGRs were observed.

Therefore, downstream analyses used DMGRs identified based on probe position in

relation to TSS.

117   We identified nineteen DMGRs with common methylation alterations among

118   tumor subtypes in comparison with normal tissues that were annotated to eleven genes:

119   *AGRN, C1orf170, FAM41C, FLJ39609, HES4, ISG15, KLHL17, NOC2L, PLEKNH1,*

120   *SAMD11,* and *WASH5P* (Supplementary Table S3).

121   Dependent upon tumor subtype, some gene regions had a different directional

change in tumor methylation compared to normal tissue (e.g. *C1orf170*, *HES4*, *and*

ISG15). Additionally, of the eleven genes identified, we observed differential methylation

in different regions including gene body, promoter (TSS1500, and TSS200), and 3'UTR

(Table 2 and Supplementary Table S3). All nineteen DMGRs also had differential

methylation in at least one late stage tumor subtype, and thirteen of the nineteen DMGRs

were significantly differentially methylated across all tumor subtypes in late stage tumors

(Table 2 and Supplementary Table S4). A heatmap of the unadjusted beta values for

individual CpGs from the nineteen DMGRs demonstrated grouping of most of the Basal-

like tumors separate from a group of mixed Luminal and Her2 tumors (Figure 2).

Table 2. Nineteen differentially methylated gene regions in common to early stage tumors.

DMGR	Alternate Gene Name	Basal Med Q	Her2 Med Q	Lum A Med Q	Lum B Med Q	*Any late stage	*All late stage	Present in validation	Validation Median Q
AGRN Body	AGNR	2.44E-06	1.68E-04	1.82E-07	1.29E-06	Y	--	Y	7.80E-21
C1orf170 Body	PERM1	4.03E-11	1.68E-05	5.46E-09	9.69E-04	Y	Y	Y	1.31E-08
C1orf170 TSS1500	PERM1	5.40E-04	6.52E-03	7.82E-06	6.82E-05	Y	--	Y	9.23E-03
FAM41C Body	FAM41C	4.13E-03	4.20E-08	1.18E-20	3.43E-03	Y	Y	Y	8.25E-10
FAM41C TSS1500	FAM41C	3.27E-04	1.11E-04	8.38E-05	1.04E-34	Y	Y	Y	1.75E-24
FLJ39609 TSS200	LOC100130417	1.30E-04	6.02E-05	2.92E-06	3.67E-04	Y	Y	Y	5.24E-06
HES4 TSS1500	HES4	3.06E-03	5.15E-04	7.84E-05	2.20E-04	Y	--	Y	5.06E-04
ISG15 Body	ISG15	3.14E-07	2.40E-04	1.18E-05	3.58E-04	Y	Y	--	1.03E-01
KLHL17 3'UTR	KLHL17	3.14E-05	5.51E-07	3.83E-16	2.27E-03	Y	Y	Y	3.99E-08
KLHL17 Body	KLHL17	5.90E-06	1.10E-04	7.85E-04	7.24E-05	Y	--	Y	1.60E-06
NOC2L Body	NOC2L	3.15E-04	6.15E-04	6.56E-05	2.40E-06	Y	Y	Y	4.90E-11
PLEKHN1 3'UTR	PLEKHN1	5.17E-16	4.73E-06	3.10E-07	7.74E-06	Y	--	Y	9.83E-09
PLEKHN1 Body	PLEKHN1	8.94E-10	2.71E-09	7.58E-29	1.73E-30	Y	Y	Y	5.87E-18
PLEKHN1 TSS1500	PLEKHN1	3.14E-05	5.51E-07	2.59E-06	3.63E-07	Y	Y	Y	3.99E-08
PLEKHN1 TSS200	PLEKHN1	1.56E-18	5.77E-10	1.42E-03	1.22E-03	Y	Y	Y	2.93E-10
SAMD11 5'UTR	SAMD11	3.58E-03	7.23E-12	1.01E-09	2.21E-08	Y	Y	Y	4.59E-11
SAMD11 Body	SAMD11	7.13E-08	2.47E-08	8.49E-06	2.04E-04	Y	Y	Y	3.26E-23
SAMD11 TSS1500	SAMD11	2.38E-03	6.14E-04	8.56E-04	1.02E-03	Y	Y	Y	2.02E-05
WASH5P Body	WASH7P	2.93E-03	9.84E-03	1.64E-03	1.25E-05	Y	--	--	7.01E-02

*Reference to any or all breast cancer subtypes in late stage tumors

Figure 2. Raw beta value (unadjusted for cellular composition) heatmap of the

significantly differentially methylated CpG sites mapping to the common early stage

differentially methylated gene regions. The genomic context is given in the vertical color

bar and the PAM50 subtype and tumor information (stage and subtype) are given in the

horizontal bars. Yellow indicates low methylation and blue indicates high methylation

beta values.

DMGRs cluster on chromosome 1p36 and on gene bodies

Of the nineteen DMGRs identified, all of them are in eleven genes located on the

p36.3 cytoband of chromosome 1 (Supplementary Figure S4). Chromosome 1p36.3 is the

start section of chromosome 1 and of the eleven genes identified, one (*WASH5P*) is

located near the very start of the chromosome (chr1:14,362 - 29,370) and the other ten

genes are located end-to-end between chr1:868,071 - 1,056,116 (Supplementary Figure

S4).

Most of the DMGRs tracked to gene body regions: *AGRN, C1orf170, FAM41C,*

ISG15, KLHL17, NOC2L, PLEKHN1, SAMD11, and *WASH5P* all had gene body

methylation differences. Gene body regions were enriched among early stage tumor

DMGRs compared to all other regions (Fisher's Exact Test OR = 4.15, 95% CI = 1.04 –

23.83, *P* = 0.04). All differentially methylated CpG probe IDs are given in

Supplementary Table S5. DAVID pathway analysis applied to the top 400 most

aberrantly methylated genes in common to the four PAM50 subtypes identified the GO

term for the regulation of hormone levels to be significantly enriched (GO:0010817, *FDR*

= 0.035, Supplementary Table S6).

Breast cancer copy number alterations in 1p36

Among these 523 tumors, the prevalence of 1p36.3 copy number alteration was

only 1.2% (n=6), all were amplifications that affected ten of the eleven genes most distal

to the chromosome end. Among the six tumors with 1p36.3 amplification three were

Basal-like, two were Her2-enriched, and one was Luminal A. Exclusive of tumors with

copy number alterations, there was one tumor (Her2-enriched), with a truncating

mutation in *KLHL17,* and one tumor with a missense mutation in *PLEKHN1* (Basal-like).

DMGRs impact gene expression

We identified CpG sites with significant correlation of methylation with gene

expression for five genes (*AGRN, PLEKHN1, KLHL17, SAMD11,* and *FAM41C*),

associated with eight DMGRs (Supplementary Table S7 and Supplementary Figures S6-

9).

Validating DMGR hits in an independent dataset

We validated our findings in an independent 450K methylation data set from 186

tumors and 46 normal tissues described in Fleischer *et al.* (GSE60185). Seventeen of

nineteen DMGRs were significantly differentially methylated between tumor and normal

tissues in the replication set (all DMGRs at $Q < 0.01$; Table 2), and CpGs in these

DMGRs had similar patterns of beta value distributions (Supplemental Figure S10). The

remaining two gene regions were also highly ranked in the *q* value distribution (*WASH5P*

body: $Q = 0.07$; *ISG15* Body: $Q = 0.10$).

Reproducibility

All TCGA and validation data is publicly available. We also provide software

under an open source license for analysis reproducibility and to build upon our work[21].

DISCUSSION:

We were interested in identifying common biology underlying breast cancer

independent of molecular subtype and cell-type proportion. After applying a reference-

free deconvolution algorithm, we observed that early stage tumors harbor differentially

methylated gene regions localized entirely to a small region on 1p36.3 shared across four

major subtypes. Although DNA methylation alterations are widespread in early stage

tumors and prior work has demonstrated alterations that differ among breast tumor

subtypes[9,22] we observed only 19 DMGRs that overlapped molecular subtypes. All

DMGRs tracking to the same region on 1p36.3 suggests that altered regulation of this

region contributes to breast carcinogenesis irrespective of disease subtype.

Previously, alterations on chromosome 1 have been observed in breast cancer cell

lines and tumors[23]. Additionally, copy number deletions in this region have been shown

to be an important precursor in DCIS tumors [24] and in follicular lymphomas [25]. However,

the most prevalent copy number alterations on chromosome 1 are gains on the *q* arm and

losses on the *p* arm that do not typically fully encompass our implicated genes on

1p36.3[23,26,27]. The region is also well-studied and significantly altered in neuroblastoma –

the most common solid tissue tumor of childhood[28–31]. The biological underpinnings of

this region remain elusive[19,32] but a systematic understanding of how these specific

DMGRs may impact early cancer development may be important for other cancer types

and not just breast cancer.

Of the nineteen DMGRs identified, eighteen of them replicated in either one or

both late stage and independent validation sets. The one DMGR that did not replicate was

the *WASH5P body*. This region is located more than 830,000 base pairs (bp) away from

the much tighter region spanned by the remaining eighteen DMGRs (~188,000 bp),

suggesting a loose association between *WASH5P* and the other ten genes.

There is also additional evidence implicating the potential importance of the

identified genes assigned to the differentially methylated regions. For example, in a study

of mutational profiles in metastatic breast cancers, *AGRN* was more frequently mutated in

metastatic cancers compared with early breast cancers[33]. Similarly, expression of the

HES4 Notch gene is known to be significantly correlated with the presence of activating

mutations in multiple breast cancer cell lines, and is associated with poor patient

outcomes[34]. In addition, *ISG15* has been implicated as a key player in breast

carcinogenesis[35], though there is conflicting evidence[36]. However, the conflicting

evidence to date may be related to our observation of *ISG15* hypomethylation in Basal-

Like, Her2, and LumB tumors, and hypermethylation in LumA tumors (Supplementary

Table S3). Opposing methylation states among tumor subtypes relative to normal tissue

may contribute to subtype-specific roles of *ISG15* dysregulation in breast carcinogenesis.

Additionally, the *NOC2L* gene has been identified as a member of a group of prognostic

genes derived from an integrated microarray of breast cancer studies[37]. We also identified

three DMGRs – TSS1500, Body, & 5'UTR – in the *SAMD11* gene, which has

significantly reduced expression in breast cancer cells compared to normal tissues[38],

consistent with our findings of *SAMD11* hypermethylation across all four breast cancer

subtypes. As DNAm changes were observed consistently and robustly across subtypes, it

is likely that several of the other identified genes are cancer initiation factors that require

additional study.

231   Importantly, we validated the identified DMGRs in an independent set of invasive

breast tumors and normal tissues. Our validation is strengthened by the lack of molecular

subtype assignments in the validation set. The validation of DMGRs in a setting agnostic

to intrinsic subtype indicates that differential magnitude or direction of methylation

alterations that may be present in different subtypes did not limit our ability to identify

significant alterations. A limitation of the validation set is a lack of gene expression data

to further investigate relationships between expression and methylation for each gene

region. Nevertheless, additional targeted studies on this set of validated genes and gene

regions can enhance the understanding of methylation alterations at these DMGRs in

breast carcinogenesis.

241   Caution should be exercised in interpreting the results of the adjusted beta

coefficients from the reference-free algorithm. It is unclear if specific disease states are a

result of aberrant methylation profiles in specific cell types which then cause changes to

cell mixtures, or if the disease state is a result of cell-type proportion differences.

Additionally, the unsupervised clustering heatmaps plot unadjusted methylation beta

values and do not account for cell type adjustment. Lastly, the DMGR analysis drops

CpGs that do not track to gene regions, which may reduce detection of non-genic regions

related with breast carcinogenesis.

249   We identified and validated DMGRs in early stage breast tumors across PAM50

subtypes that are located on chromosome 1p36.3. The observed differential methylation

suggests that this region may contribute to the initiation or progression to invasive breast

cancer. Additional work is needed to investigate the scope of necessary and sufficient

alterations to 1p36.3 for transformation and to more clearly understand the implications

of 1p36.3 methylation alterations to gene regulation. Further investigation of DNAm

changes to 1p36.3 may identify opportunities for early identification of breast cancer or

risk assessment. Lastly, the reference-free approach we used could be applied to

methylation datasets from other tumor types to identify potential drivers of

carcinogenesis common across histologic or intrinsic molecular subtypes.

PATIENTS & METHODS:

Data Processing

We accessed breast invasive carcinoma Level 1 Illumina HumanMethylation450 (450K) DNAm data (n = 870) from the TCGA data access portal and downloaded all sample intensity data (IDAT) files. We processed the IDAT files with the R package *minfi* using the "Funnorm" normalization method on the full dataset [39]. We filtered CpGs with a detection *P*-value > 1.0E-05 in more than 25% of samples, CpGs with high frequency SNP(s) in the probe, probes previously described to be potentially cross-hybridizing, and sex-specific probes [40,41]. We filtered samples that did not have full covariate data (PAM50 subtype, pathologic stage[42,12]) and full demographic data (age and sex). All tumor adjacent normal samples were included regardless of missing data (n = 97, Table 1).

From an original set of 485,512 measured CpG sites on the Illumina 450K array, our filtering steps removed 2,932 probes exceeding the detection *P*-value limit, and 93,801 probes that were SNP-associated, cross-hybridizing, or sex-specific resulting in a final analytic set of 388,779 CpGs. From 870 TCGA breast tumors, we restricted to primary tumors with available PAM50 intrinsic subtype assignments of Basal-like (n = 86), Her2 (n = 31), Luminal A (n = 279), and Luminal B (n = 127), excluding Normal-like tumors due to limited sample size (n = 18). Lastly, we restricted the final total tumor set to only those with stage assignments resulting in a final analytic sample size of n = 523.

Reference-free cell type adjustment modeling

We stratified samples by PAM50 subtype (Basal-like, Luminal A, Luminal B,

Her2) and then by tumor stage dichotomizing as early (stage I and II tumors) and late

(stage III and IV tumors)[42], resulting in eight distinct models. To analyze DNAm

differences between tumor and normal tissue and to adjust for effects of cellular

heterogeneity across samples, we applied the reference-free deconvolution algorithm

from the *RefFreeEWAS* R package to each model adjusting for age[16]. The method

estimates the number of underlying tissue-specific cell methylation states contributing to

methylation heterogeneity through a constrained variant of NMF[43]. Briefly, the method

assumes the sample methylome is composed of a linear combination of the constituent

methylomes. It decomposes the matrix of sample methylation values (Y) into two

matrices ($Y = M\Omega^{T}$), where M is an $m \; x \; K$ matrix of m CpG-specific methylations states

for K cell types and Ω is a $n \; x \; K$ matrix of subject-specific cell-types. K is selected via

bootstrapping $K = 2\ldots10$ and choosing the optimal K that minimizes the bootstrapped

deviance. To correct for multiple comparisons, we converted all extracted P-values to Q-

values using the R package *qvalue*[44].

Identifying differentially methylated gene regions

To understand the genomic regions with common DNAm alterations we grouped

CpGs by gene and region relative to genomic location (transcription start site 1500

(TSS1500), TSS200, 3' untranslated region (3'UTR), 5'UTR, 1st exon, and gene body).

We used this gene-region taxonomy to collapse differentially methylated CpGs, as

defined by our Q-value cutoff, into specific differentially methylated gene regions

(DMGRs). This extended the Illumina 450K CpG annotation file to allow for a given

CpG to be associated with up to two genes depending on the proximity of the CpG site to

neighboring genes (Figure 3).

Figure 3. *Diagram of CpG sites relative to gene regions (Transcription start sites*

(TSS1500 & TSS200), Untranslated regions (5'UTR & 3'UTR), and the gene body). Dark

circles indicate methylated sites and empty circles indicate unmethylated sites.

We defined a differentially methylated CpG as one with a Q-value < 0.01 following cell-

type adjustment in a specific subtype model compared to normal tissue. To identify

DMGR sets for each stage and subtype, we analyzed all eight models independently.

Pathway Analysis

We performed a DAVID (the database for annotation, visualization and integrated

discovery) analysis[45,46] for the 400 genes with the lowest median CpG Q-values that are

in common to all early stage tumors regardless of PAM50 subtype, and extracted

enriched Gene Ontology (GO)[47] and Kyoto Encyclopedia of Genes and Genomes

(KEGG)[48] terms. We selected the top 400 genes based on recommended gene list sizes[49].

Copy number, gene expression, and genomic location

We downloaded TCGA Breast Invasive Carcinoma CNV data[9] and normalized

RNAseq using cBioPortal[50]. For the DMGRs we identified, we analyzed the prevalence

of copy number alterations and mutations in each gene across all samples, stratified by

molecular subtype. Similarly, to determine whether these DMGRs affect gene expression

of their target gene, we calculated Spearman correlations of DNAm beta values in

significant CpGs ($Q < 0.01$) to matched sample Illumina HiSeq gene expression data. We

used a Bonferroni correction to determine significant expression differences, resulting in

an acceptance alpha value of 9.36E-5.

Validation

To confirm the identified early stage DMGRs in common among intrinsic

molecular subtypes we applied the analysis workflow to TCGA late stage tumors and an

independent validation set (GSE60185)[20]. The validation set includes samples of ductal

carcinoma *in situ* (DCIS), mixed, invasive, and normal histology collected from Akershus

University Hospital and from the Norwegian Radium Hospital. We analyzed only the

invasive samples compared to normal samples using the same bioinformatics pipeline of

quality control CpG filtering steps and normalization procedures. However, we did not

have complete age information or intrinsic subtype assignments for the validation set and

the models are not adjusted for age or stratified by subtype. This resulted in a single

model comparing 186 invasive tumors with 46 normal controls measured across 390,253

CpGs.

ACKNOWLEDGEMENTS:

Funding was provided by P20GM104416 and R01DE02277 (BCC), by the Quantitative

Biomedical Sciences graduate program, and through a BD2K Fellowship to AJT

(T32LM012204).

COMPETING INTERESTS:

The authors declare that they have no competing interests

REFERENCES:

1. Perou CM, Sørlie T, Eisen MB, van de Rijn M, Jeffrey SS, Rees CA, Pollack JR, Ross
DT, Johnsen H, Akslen LA, Fluge O, Pergamenschikov A, Williams C, Zhu SX,
Lønning PE, Børresen-Dale AL, Brown PO, Botstein D. Molecular portraits of
human breast tumours. Nature. 2000 Aug 17;406(6797):747–752. PMID:
10963602

2. Vogelstein B, Papadopoulos N, Velculescu VE, Zhou S, Diaz LA, Kinzler KW.
Cancer genome landscapes. Science. 2013 Mar 29;339(6127):1546–1558.
PMCID: PMC3749880

3. Jones PA, Baylin SB. The fundamental role of epigenetic events in cancer. Nat
Rev Genet. 2002 Jun;3(6):415–428. PMID: 12042769

4. Yang X, Yan L, Davidson NE. DNA methylation in breast cancer. Endocr Relat
Cancer. 2001 Jun;8(2):115–127. PMID: 11446343

5. Baylin SB, Esteller M, Rountree MR, Bachman KE, Schuebel K, Herman JG.
Aberrant patterns of DNA methylation, chromatin formation and gene
expression in cancer. Hum Mol Genet. 2001 Apr;10(7):687–692. PMID:
11257100

6. Fang F, Turcan S, Rimner A, Kaufman A, Giri D, Morris LGT, Shen R, Seshan V,
Mo Q, Heguy A, Baylin SB, Ahuja N, Viale A, Massague J, Norton L, Vahdat LT,
Moynahan ME, Chan TA. Breast cancer methylomes establish an epigenomic
foundation for metastasis. Sci Transl Med. 2011 Mar 23;3(75):75ra25. PMCID:
PMC3146366

7. Kamalakaran S, Varadan V, Giercksky Russnes HE, Levy D, Kendall J, Janevski A,
Riggs M, Banerjee N, Synnestvedt M, Schlichting E, Karesen R, Shama Prasada K,
Rotti H, Rao R, Rao L, Eric Tang M-H, Satyamoorthy K, Lucito R, Wigler M,
Dimitrova N, Naume B, Borresen-Dale A-L, Hicks JB. DNA methylation patterns
in luminal breast cancers differ from non-luminal subtypes and can identify
relapse risk independent of other clinical variables. Mol Oncol. 2011
Feb;5(1):77–92. PMID: 21169070

8. Sørlie T, Perou CM, Tibshirani R, Aas T, Geisler S, Johnsen H, Hastie T, Eisen MB,
van de Rijn M, Jeffrey SS, Thorsen T, Quist H, Matese JC, Brown PO, Botstein D,
Lønning PE, Børresen-Dale A-L. Gene expression patterns of breast carcinomas
distinguish tumor subclasses with clinical implications. Proc Natl Acad Sci.
2001 Sep 11;98(19):10869–10874. PMID: 11553815

9. Cancer Genome Atlas Network. Comprehensive molecular portraits of human
breast tumours. Nature. 2012 Oct 4;490(7418):61–70. PMCID: PMC3465532

10.   Beca F, Polyak K. Intratumor Heterogeneity in Breast Cancer. Adv Exp Med Biol.
2016;882:169–189. PMID: 26987535

11.   Yoshihara K, Shahmoradgoli M, Martinez E, Vegesna R, Kim H, Torres-Garcia W,
Trevino V, Shen H, Laird PW, Levine DA, Carter SL, Getz G, Stemke-Hale K, Mills
GB, Verhaak RGW. Inferring tumour purity and stromal and immune cell
admixture from expression data. Nat Commun. 2013;4:2612. PMCID:
PMC3826632

12.   Bloushtain-Qimron N, Yao J, Snyder EL, Shipitsin M, Campbell LL, Mani SA, Hu
M, Chen H, Ustyansky V, Antosiewicz JE, Argani P, Halushka MK, Thomson JA,
Pharoah P, Porgador A, Sukumar S, Parsons R, Richardson AL, Stampfer MR,
Gelman RS, Nikolskaya T, Nikolsky Y, Polyak K. Cell type-specific DNA
methylation patterns in the human breast. Proc Natl Acad Sci U S A. 2008 Sep
16;105(37):14076–14081. PMCID: PMC2532972

13.   Christensen BC, Houseman EA, Marsit CJ, Zheng S, Wrensch MR, Wiemels JL,
Nelson HH, Karagas MR, Padbury JF, Bueno R, Sugarbaker DJ, Yeh R-F, Wiencke
JK, Kelsey KT. Aging and environmental exposures alter tissue-specific DNA
methylation dependent upon CpG island context. PLoS Genet. 2009
Aug;5(8):e1000602. PMCID: PMC2718614

14.   Santagata S, Thakkar A, Ergonul A, Wang B, Woo T, Hu R, Harrell JC, McNamara
G, Schwede M, Culhane AC, Kindelberger D, Rodig S, Richardson A, Schnitt SJ,
Tamimi RM, Ince TA. Taxonomy of breast cancer based on normal cell
phenotype predicts outcome. J Clin Invest. 2014 Feb;124(2):859–870. PMCID:
PMC3904619

15.   Koestler DC, Christensen B, Karagas MR, Marsit CJ, Langevin SM, Kelsey KT,
Wiencke JK, Houseman EA. Blood-based profiles of DNA methylation predict
the underlying distribution of cell types: a validation analysis. Epigenetics Off J
DNA Methylation Soc. 2013 Aug;8(8):816–826. PMCID: PMC3883785

16.   Houseman EA, Kile ML, Christiani DC, Ince TA, Kelsey KT, Marsit CJ. Reference-
free deconvolution of DNA methylation data and mediation by cell composition
effects. BMC Bioinformatics. 2016;17(1):259. PMID: 27358049

17.   Houseman EA, Kelsey KT, Wiencke JK, Marsit CJ. Cell-composition effects in the
analysis of DNA methylation array data: a mathematical perspective. BMC
Bioinformatics. 2015 Mar 21;16:95. PMCID: PMC4392865

18.   Houseman EA, Ince TA. Normal cell-type epigenetics and breast cancer
classification: a case study of cell mixture-adjusted analysis of DNA methylation
data from tumors. Cancer Inform. 2014;13(Suppl 4):53–64. PMCID:
PMC4264613

19. Bagchi A, Mills AA. The Quest for the 1p36 Tumor Suppressor. Cancer Res.
2008 Apr 15;68(8):2551–2556. PMID: 18413720

20. Fleischer T, Frigessi A, Johnson KC, Edvardsen H, Touleimat N, Klajic J, Riis ML,
Haakensen VD, Wärnberg F, Naume B, Helland A, Børresen-Dale A-L, Tost J,
Christensen BC, Kristensen VN. Genome-wide DNA methylation profiles in
progression to in situ and invasive carcinoma of the breast with impact on gene
transcription and prognosis. Genome Biol. 2014;15(8):435. PMCID:
PMC4165906

21. Titus AJ, Way GP, Johnson KC, Christensen BC. Analytical code for "Reference-
free deconvolution of DNA methylation signatures identifies common
differentially methylated gene regions on 1p36 across breast cancer subtypes."
2017 Mar 10; Available from: https://zenodo.org/badge/latestdoi/45754471

22. Fang F, Turcan S, Rimner A, Kaufman A, Giri D, Morris LGT, Shen R, Seshan V,
Mo Q, Heguy A, Baylin SB, Ahuja N, Viale A, Massague J, Norton L, Vahdat LT,
Moynahan ME, Chan TA. Breast Cancer Methylomes Establish an Epigenomic
Foundation for Metastasis. Sci Transl Med. 2011 Mar 23;3(75):75ra25. PMID:
21430268

23. Orsetti B, Nugoli M, Cervera N, Lasorsa L, Chuchana P, Rouge C, Ursule L,
Nguyen C, Bibeau F, Rodriguez C, Theillet C. Genetic profiling of chromosome 1
in breast cancer: mapping of regions of gains and losses and identification of
candidate genes on 1q. Br J Cancer. 2006 Nov 20;95(10):1439–1447. PMCID:
PMC2360604

24. Munn KE, Walker RA, Varley JM. Frequent alterations of chromosome 1 in
ductal carcinoma in situ of the breast. Oncogene. 1995 Apr 20;10(8):1653–
1657. PMID: 7731721

25. Mamessier E, Song JY, Eberle FC, Pack S, Drevet C, Chetaille B, Abdullaev Z,
Adelaïde J, Birnbaum D, Chaffanet M, Pittaluga S, Roulland S, Chott A, Jaffe ES,
Nadel B. Early lesions of follicular lymphoma: a genetic perspective.
Haematologica. 2014 Mar;99(3):481–488. PMCID: PMC3943311

26. Bieche I, Champeme MH, Lidereau R. Loss and gain of distinct regions of
chromosome 1q in primary breast cancer. Clin Cancer Res Off J Am Assoc
Cancer Res. 1995 Jan;1(1):123–127. PMID: 9815894

27. Curtis C, Shah SP, Chin S-F, Turashvili G, Rueda OM, Dunning MJ, Speed D, Lynch
AG, Samarajiwa S, Yuan Y, Graf S, Ha G, Haffari G, Bashashati A, Russell R,
McKinney S, Langerod A, Green A, Provenzano E, Wishart G, Pinder S, Watson P,
Markowetz F, Murphy L, Ellis I, Purushotham A, Borresen-Dale A-L, Brenton JD,
Tavare S, Caldas C, Aparicio S. The genomic and transcriptomic architecture of
2,000 breast tumours reveals novel subgroups. Nature. 2012 Apr
18;486(7403):346–352. PMCID: PMC3440846

28. White PS, Thompson PM, Gotoh T, Okawa ER, Igarashi J, Kok M, Winter C,
Gregory SG, Hogarty MD, Maris JM, Brodeur GM. Definition and
characterization of a region of 1p36.3 consistently deleted in neuroblastoma.
Oncogene. 2005 Apr 14;24(16):2684–2694. PMID: 15829979

29. Attiyeh EF, London WB, Mossé YP, Wang Q, Winter C, Khazi D, McGrady PW,
Seeger RC, Look AT, Shimada H, Brodeur GM, Cohn SL, Matthay KK, Maris JM.
Chromosome 1p and 11q Deletions and Outcome in Neuroblastoma. N Engl J
Med. 2005 Nov 24;353(21):2243–2253. PMID: 16306521

30. Caren H, Ejeskar K, Fransson S, Hesson L, Latif F, Sjoberg R-M, Krona C,
Martinsson T. A cluster of genes located in 1p36 are down-regulated in
neuroblastomas with poor prognosis, but not due to CpG island methylation.
Mol Cancer. 2005 Mar 1;4(1):10. PMCID: PMC554762

31. Carén H, Fransson S, Ejeskär K, Kogner P, Martinsson T. Genetic and epigenetic
changes in the common 1p36 deletion in neuroblastoma tumours. Br J Cancer.
2007 Nov 19;97(10):1416–1424. PMCID: PMC2360241

32. Henrich K-O, Schwab M, Westermann F. 1p36 tumor suppression--a matter of
dosage? Cancer Res. 2012 Dec 1;72(23):6079–6088. PMID: 23172308

33. Lefebvre C, Bachelot T, Filleron T, Pedrero M, Campone M, Soria J-C, Massard C,
Levy C, Arnedos M, Lacroix-Triki M, Garrabey J, Boursin Y, Deloger M, Fu Y,
Commo F, Scott V, Lacroix L, Dieci MV, Kamal M, Dieras V, Goncalves A, Ferrero
J-M, Romieu G, Vanlemmens L, Mouret Reynier M-A, Thery J-C, Le Du F, Guiu S,
Dalenc F, Clapisson G, Bonnefoi H, Jimenez M, Le Tourneau C, Andre F.
Mutational Profile of Metastatic Breast Cancers: A Retrospective Analysis. PLoS
Med. 2016 Dec;13(12):e1002201. PMCID: PMC5189935

34. Stoeck A, Lejnine S, Truong A, Pan L, Wang H, Zang C, Yuan J, Ware C, MacLean J,
Garrett-Engele PW, Kluk M, Laskey J, Haines BB, Moskaluk C, Zawel L, Fawell S,
Gilliland G, Zhang T, Kremer BE, Knoechel B, Bernstein BE, Pear WS, Liu XS,
Aster JC, Sathyanarayanan S. Discovery of biomarkers predictive of GSI
response in triple-negative breast cancer and adenoid cystic carcinoma. Cancer
Discov. 2014 Oct;4(10):1154–1167. PMCID: PMC4184927

35. Burks J, Reed RE, Desai SD. Free ISG15 triggers an antitumor immune response
against breast cancer: a new perspective. Oncotarget. 2015 Mar 30;6(9):7221–
7231. PMCID: PMC4466680

36. Andersen JB, Hassel BA. The interferon regulated ubiquitin-like protein, ISG15,
in tumorigenesis: friend or foe? Cytokine Growth Factor Rev. 2006
Dec;17(6):411–421. PMID: 17097911

37. Xu L, Tan AC, Winslow RL, Geman D. Merging microarray data from separate
breast cancer studies provides a robust prognostic test. BMC Bioinformatics.
2008 Feb 27;9:125. PMCID: PMC2409450

38. Rodriguez-Martinez A, Alarmo E-L, Saarinen L, Ketolainen J, Nousiainen K,
Hautaniemi S, Kallioniemi A. Analysis of BMP4 and BMP7 signaling in breast
cancer cells unveils time-dependent transcription patterns and highlights a
common synexpression group of genes. BMC Med Genomics. 2011 Nov 25;4:80.
PMCID: PMC3229454

39. Hansen KD, Fortin JP. Minfi tutorial. BioC2014. 2014;

40. Chen Y, Lemire M, Choufani S, Butcher DT, Grafodatskaya D, Zanke BW,
Gallinger S, Hudson TJ, Weksberg R. Discovery of cross-reactive probes and
polymorphic CpGs in the Illumina Infinium HumanMethylation450 microarray.
Epigenetics Off J DNA Methylation Soc. 2013 Feb;8(2):203–209. PMCID:
PMC3592906

41. Wilhelm-Benartzi CS, Koestler DC, Karagas MR, Flanagan JM, Christensen BC,
Kelsey KT, Marsit CJ, Houseman EA, Brown R. Review of processing and
analysis methods for DNA methylation array data. Br J Cancer. 2013 Sep
17;109(6):1394–1402. PMCID: PMC3777004

42. Edge S, Byrd D, Compton C, Fritz A, Greene F, Trotti A, editors. AJCC cancer
staging manual. 7th ed. New York, NY: Springer; 2010.

43. Brunet J-P, Tamayo P, Golub TR, Mesirov JP. Metagenes and molecular pattern
discovery using matrix factorization. Proc Natl Acad Sci U S A. 2004 Mar
23;101(12):4164–4169. PMCID: PMC384712

44. Dabney A, Storey J. qvalue: Q-value estimation for false discovery rate control.
R Package Version 1430.

45. Huang DW, Sherman BT, Lempicki RA. Bioinformatics enrichment tools: paths
toward the comprehensive functional analysis of large gene lists. Nucleic Acids
Res. 2009 Jan;37(1):1–13. PMCID: PMC2615629

46. Huang DW, Sherman BT, Lempicki RA. Systematic and integrative analysis of
large gene lists using DAVID bioinformatics resources. Nat Protoc.
2009;4(1):44–57. PMID: 19131956

47. Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, Davis AP,
Dolinski K, Dwight SS, Eppig JT, Harris MA, Hill DP, Issel-Tarver L, Kasarskis A,
Lewis S, Matese JC, Richardson JE, Ringwald M, Rubin GM, Sherlock G. Gene
ontology: tool for the unification of biology. The Gene Ontology Consortium.
Nat Genet. 2000 May;25(1):25–29. PMCID: PMC3037419

48.   Kanehisa M, Goto S. KEGG: kyoto encyclopedia of genes and genomes. Nucleic
Acids Res. 2000 Jan 1;28(1):27–30. PMCID: PMC102409

49.   Huang DW, Sherman BT, Lempicki RA. Systematic and integrative analysis of
large gene lists using DAVID bioinformatics resources. Nat Protoc.
2009;4(1):44–57. PMID: 19131956

50.   Gao J, Aksoy BA, Dogrusoz U, Dresdner G, Gross B, Sumer SO, Sun Y, Jacobsen A,
Sinha R, Larsson E, Cerami E, Sander C, Schultz N. Integrative analysis of
complex cancer genomics and clinical profiles using the cBioPortal. Sci Signal.
2013 Apr 2;6(269):pl1. PMCID: PMC4160307

SUPPLAMENTAL FIGURES:

Supplementary Figure S1. *Box plots show the distribution of tumor purity across all*

551   *subtypes for both early and late stages of the TCGA dataset. The measurements estimated*

552   *by TCGA are based on histology slides and indicate the estimated distribution of the*

553   *number of tumor cells, stromal cells, and normal cells in each sample. See the NCI CDE*

554   *Browser for more details.*

Supplementary Figure S2. *Volcano plots from the eight models. The left most panel in*

each model indicates unadjusted P values and the right panel indicates RefFree adjusted

P values. Each point represents a CpG considered in the model and the color of the

points represents the change in the beta coefficient following adjustment (delta value).

The red lines indicate a Q value cutoff of 0.01 and the black lines indicate a Q value

cutoff of 0.05.

Supplementary Figure S3. *Venn diagram depicting overlapping Illumina annotation file*

UCSC regions between (A) early and (B) late stage tumors stratified by subtype. The

regions consist of mappings relative to CpG island definitions (e.g. <Gene Name>

N_Shore).

Supplementary Figure S4. *Diagram of chromosome 1. (A) The entire chromosome 1*

with regions annotated. (B) A zoomed in view of chromosome 1p36.3 with each identified

gene annotated on a track and highlighted in red boxes indicating a gene cluster between

base pairs 868,071 - 1,056,116. (C) The negative log of the median Q-value for all CpG

sites within each DMGR, stratified by PAM50 subtype and arranged along the x-axis

according to genomic position reflected in panel B. (D) The negative log of the median

Q-value for all CpG sites within each DMGR in the ten gene cluster (without WASH5P),

stratified by PAM50 subtype and arranged along the x-axis according to genomic

position reflected in panel B.

Supplementary Figure S5. *The relationship between differentially methylated CpG sites*

and FAM41C gene expression in early stage tumors and normal tissue with matched

RNAseq samples stratified by PAM50 subtype. All tumors (orange), all normal tissue

(black), Luminal A (blue), Luminal B (green), Her2 (purple), and Basal-like (red) are

given in the different facets of the figure.

Supplementary Figure S6. *The relationship between differentially methylated CpG sites*

and AGRN gene expression in early stage tumors and normal tissue with matched

RNAseq samples stratified by PAM50 subtype. All tumors (orange), all normal tissue

(black), Luminal A (blue), Luminal B (green), Her2 (purple), and Basal-like (red) are

given in the different facets of the figure.

Supplementary Figure S7. *The relationship between differentially methylated CpG sites and PLEKHN1 gene expression in early stage tumors and normal tissue with matched RNAseq samples stratified by PAM50 subtype. All tumors (orange), all normal tissue (black), Luminal A (blue), Luminal B (green), Her2 (purple), and Basal-like (red) are given in the different facets of the figure.*

604 ***Supplementary Figure S8.*** *The relationship between differentially methylated CpG sites*

605 *and KLHL17 gene expression in early stage tumors and normal tissue with matched*

606 *RNAseq samples stratified by PAM50 subtype. All tumors (orange), all normal tissue*

607 *(black), Luminal A (blue), Luminal B (green), Her2 (purple), and Basal-like (red) are*

608 *given in the different facets of the figure.*

Supplementary Figure S9. The relationship between differentially methylated CpG sites

and SAMD11 gene expression in early stage tumors and normal tissue with matched

RNAseq samples stratified by PAM50 subtype. All tumors (orange), all normal tissue

(black), Luminal A (blue), Luminal B (green), Her2 (purple), and Basal-like (red) are

given in the different facets of the figure.

Supplementary Figure S10. *Results from the validation set (Fleischer et al 2014;*

GSE60185). Validation set raw (unadjusted) beta value heatmap of the significantly

differentially methylated CpG sites in the common early stage differentially methylated

gene regions (DMGRs) identified in the initial analysis. The genomic context is given in

the vertical color bar (blue = gene body, dark pink = TSS200, light pink = TSS1500) and

tumor vs. normal status is given in the horizontal color bar (black = tumor, white =

normal tissue). In the heatmap, yellow indicates low methylation and blue indicates high

methylation.

627   SUPPLAMENTAL TABLES:

628   Due to size limitations of this document and the size of the supplemental tables available

629   for this manuscript, supplemental tables may be found at the following DOI link:

630   DOI: 10.5281/zenodo.400247