Effect size and statistical power in the rodent fear conditioning literature – a systematic review

Short title: Effect sizes and statistical power in fear conditioning

Authors: Clarissa F. D. Carneiro[1*], Thiago C. Moulin[1*], Malcolm R. Macleod[2], Olavo B. Amaral[1]

[1]Institute of Medical Biochemistry Leopoldo de Meis, Federal University of Rio de Janeiro, Rio de Janeiro, Brazil and [2]Division of Clinical Neurosciences, University of Edinburgh, Edinburgh, UK

* Both authors contributed equally to the work.

Corresponding author:

Olavo B. Amaral, M.D., PhD.

Instituto de Bioquímica Médica Leopoldo de Meis
Av. Carlos Chagas Filho 373, E-38
Cidade Universitária
Rio de Janeiro, RJ, Brazil
CEP 21941-902
Phone: +55-21-39386762
E-mail: olavo@bioqmed.ufrj.br

Abstract

Proposals to increase research reproducibility frequently call for focusing on effect sizes instead of p values, as well as for increasing the statistical power of experiments. However, it is unclear to what extent these two concepts are indeed taken into account in basic biomedical science. To study this in a real-case scenario, we performed a systematic review of effect sizes and statistical power in studies on learning of rodent fear conditioning, a widely used behavioral task to evaluate memory. Our search criteria yielded 410 experiments comparing control and treated groups in 122 articles. Interventions had a mean effect size of 29.5%, and amnesia caused by memory-impairing interventions was nearly always partial. Mean statistical power to detect the average effect size observed in well-powered experiments with significant differences (37.2%) was 65%, and was lower among studies with non-significant results. Only one article reported a sample size calculation, and our estimated sample size to achieve 80% power considering typical effect sizes and variances (15 animals per group) was reached in only 12.2% of experiments. Actual effect sizes correlated with effect size inferences made by readers on the basis of textual descriptions of results only when findings were non-significant, and neither effect size nor power correlated with study quality indicators, number of citations or impact factor of the publishing journal. In summary, effect sizes and statistical power have a wide distribution in the rodent fear conditioning literature, but do not seem to have a large influence on how results are described or cited. Failure to take these concepts into consideration might limit attempts to improve reproducibility in this field of science.

Introduction

Biomedical research over the last decades has relied heavily on the concept of statistical significance – i.e. the probability that an effect equal to or larger than that observed experimentally would occur by chance under the null hypothesis – and classifying results as "significant" or "non-significant" on the basis of an arbitrary threshold (usually set at $p < 0.05$) has become standard practice in most fields. This approach, however, has well-described limitations that can lead to erroneous conclusions when researchers rely on p values alone to judge results [1–6]. First of all, p values do not measure the magnitude of an effect, and thus cannot be used by themselves to evaluate its biological significance [7]. Moreover, the predictive value of a significance test is heavily influenced by factors such as the prior probability of the tested hypothesis, the number of tests performed and their statistical power [8]; thus, similar p values can lead to very different conclusions in distinct scenarios [1].

Recent calls for improving research reproducibility have focused on reporting effect sizes and confidence intervals alongside or instead of p values [6–9] and for the use of both informal Bayesian inference [10] and formal data synthesis methods [11] when aggregating data from multiple studies. The concepts of effect size and statistical power are central for such approaches, as how much a given experiment will change a conclusion or an effect estimate will depend on both. However, it is unclear whether they receive much attention from authors in basic science publications. Discussion of effect sizes seems to be scarce, and recent data has shown that sample size and power calculations are very rare in the preclinical literature [12,13]. The potential impact of these omissions is large, as reliance on the results of significance tests without consideration of statistical power can decrease the reliability of study conclusions [14].

Another issue is that, if effect size is not taken into account, it is difficult to

adequately assess the biological significance of a given finding. As p values will be low

even for small effect sizes if sample size is large, biologically trivial effects can be

found to be statistically significant. In preclinical studies, overlooking effect sizes will

thus lead to inadequate assessment of therapeutic potential, whereas in basic research it

will cause difficulties in dissecting essential biological mechanisms from peripheral

modulatory influences [15]. The wealth of findings in the literature will thus translate

poorly into better comprehension of phenomena, and the abundance of statistically

significant findings with small effect sizes can eventually do more harm than good. This

problem is made much worse when many of these studies have low positive predictive

values due to insufficient power, leading a large fraction of them to be false positives

[8,14,16–18].

To analyze how effect sizes and statistical power are taken into account in the

description and publication of findings in a real-case scenario of basic biomedical

science, we chose to perform a systematic review of articles on learning of rodent fear

conditioning, probably the most widely used behavioral task to study memory in

animals [19]. Focusing on this task provides a major advantage in the fact that the vast

majority of articles use the same measure to describe results (i.e. percentage of time

spent in freezing behavior during a test session). As effect sizes are comparable across

studies, studying their distribution allows one to estimate the statistical power of

individual experiments to detect typical differences.

Our first objective in this study is to analyze the distribution of effect sizes and

statistical power in a large sample of articles using different interventions, showing how

they are related to the outcome of statistical significance tests. Next, we will study

whether these two measures are correlated, in order to look for evidence of publication

bias and effect size inflation. We will also correlate effect sizes and variances with

different aspects of experimental design, such as species, sex and type of conditioning,

as well as with indicators of risk of bias. To inquire whether effect size and power are

taken into consideration by authors when interpreting findings, we will evaluate

whether they correlate with effect size inferences made by readers based on textual

descriptions of results in the articles. Finally, we will analyze whether mean effect size

and power correlate with article-level metrics, such as number of citations and impact

factor of the publishing journal, to explore how they influence the publication of results.

115   **Results**

Article search and inclusion

As previously described in a protocol published in advance of full data

collection [20], we performed a PubMed search for fear conditioning articles published

online in 2013. The search process (**Fig. 1**) yielded 400 search hits, of which 386 were

original articles that were included if they fulfilled pre-established criteria (see

Methods). Two investigators examined all included articles, and agreement for

exclusions measured on a double-screened sample of 40 articles was 95%. This led to a

final sample of 122 articles and 410 experiments, used to build the database provided as

Supplementary Data.

Figure 1. Study flow diagram. Our PubMed search yielded 400 results, of which 14

were excluded based on initial screening of titles and abstracts and 386 were selected

for full-text analysis. This led to the inclusion of 122 articles, containing a total of 410

comparisons (i.e. individual experiments). The main reasons for exclusion are listed in

the figure, in compliance with the PRISMA statement [21].

Distribution of effect sizes among experiments

For each experiment, we initially calculated effect size as the relative difference

(i.e. percentage of change) in the freezing levels of treated groups when compared to

controls. As shown in **Fig. 2A**, this leads interventions that enhance memory acquisition

(i.e. those in which freezing is significantly higher in the treated group) to have larger

effect sizes than those that impair it (i.e. those in which freezing is significantly lower in

the treated group) due to an asymmetry that is inherent to ratios. To account for this and

make effect sizes comparable between both types of interventions, we used a

normalized effect size, with difference expressed as a percentage of the highest freezing

value between groups (**Fig. 2B**) [11].

Use of absolute differences in freezing instead of relative ones led to similar, but

more constrained distributions (**S1 Fig.**) due to mathematical limits on absolute

differences. Freezing levels in the reference group correlated negatively with relative

effect size and pooled coefficient of variation (i.e. the ratio between the sample size-

weighted pooled SD and the pooled mean); however, normalization by the highest-

freezing group reduced this effect (**S2 Fig. A-C**). Absolute effect size, on the contrary,

showed a positive correlation with freezing levels in the control or highest-freezing

group (**S2 Fig. D-F**). We also calculated effect sizes as standardized mean differences

(i.e Cohen's d, **S3 Fig.**), but chose to use relative percentages throughout the study, as

they are more closely related to the way results are expressed in articles.

Figure 2. Distribution of effect sizes. (A) Distribution of effect sizes, calculated as %

of control group freezing. Interventions were divided into memory-impairing (-48.6 ±

18.1%, n=146), memory-enhancing (71.6 ± 53.2%, n=53) or non-effective (-1.8 ±

26.2%, n=211) for visualization purposes, according to the statistical significance of the

comparison performed in the article. Additionally, the whole sample of experiments is

shown in grey (-9.0 ± 47.5% [-13.6 to -4.4], n=410). Values are expressed as mean ±

SD [95% confidence interval]. Lines and whiskers in the inset express median and

interquartile interval. (B) Distribution of normalized effect sizes, calculated as % of the

group with the highest mean (i.e. control group for memory-impairing interventions, or

treated group for memory-enhancing interventions).

All 410 experiments combined had a mean normalized effect size of 29.5 ±

22.4% (mean ± SD; 95% CI [27.4 to 31.7]). When results were divided according to the

statistical comparison originally performed in the article, mean normalized effect size

was 48.6 ± 18.1% for memory-impairing interventions, 37.6 ± 14.2% for memory-

enhancing interventions and 14.4 ± 14.2% for non-effective interventions – i.e. those in

which a significant difference between groups was not found. This does not imply that

data in each of these groups represents effects coming from a different distribution, or

that significant and non-significant results correspond to true positive or true negative

effects. On the contrary, each group likely represents a mixture of heterogeneous effect

size distributions, as sampling error and lack of statistical power can directly impact the

chances of a result achieving statistical significance. Distribution of mean effect sizes at

the article level showed similar results to those found at the level of experiments (**S4**

Fig.).

The distribution of effect sizes shows that the vast majority of memory-

impairing interventions cause partial reductions in learning, leaving the treated group

with residual freezing levels that are higher than those of a non-conditioned animal. In

fact, in all 35 memory-impairing experiments in which pre-conditioning freezing levels

were shown for the treated group, these were lower than those observed in the test

session – with p values below 0.05 in 25 (78%) out of the 32 cases in which there was

enough information for us to perform an unpaired *t* test between sessions **(S5 Fig.)**. It is

also worth noting that 26.5% of non-significant experiments had an effect size greater

than 20%, suggesting that these experiments might have been underpowered. With this

in mind, we went on to evaluate the distribution of statistical power among studies.

Distribution of statistical power among experiments

For analyzing statistical power, we first sought to evaluate the distribution of

sample sizes and coefficients of variation (both of which are determinants of power). As

shown in **Fig. 3A**, most experiments had mean sample sizes between 8 and 12

animals/group, and this distribution did not vary between enhancing, impairing and non-

effective interventions. On the other hand, higher coefficients of variation were more

frequent among non-effective interventions (**Fig. 3B**). This difference was partly

explained by freezing levels in the reference group – which correlated negatively with

coefficients of variation (**S2 Fig. G-I**) and were lower on average for non-significant

experiments (49.3% vs. 52.9% in memory-impairing and 61.3% in memory-enhancing

experiments).

Figure 3. Distribution of sample size, variation and statistical power. (A)

Distribution of mean sample size between groups. Gray bars show the distribution of

the whole sample, while colored lines show distributions of impairing (n=120),

enhancing (n=39) and non-significant (n=177) experiments separately for visualization

purposes. (B) Distribution of coefficients of variation (pooled standard deviation/pooled

mean) for each type of experiment. (C) Distribution of statistical power for memory-

impairing interventions: based on each experiment's variance and sample size, power

varies according to the difference to be detected for α=0.05. Dashed lines show the three

effect sizes used for point estimates of power in F, G and H. (D) Distribution of

statistical power for memory-enhancing interventions. (E) Distribution of statistical

power for non-effective interventions. (F) Distribution of statistical power to detect the

upper-bound effect size of 45.6% (right dashed line on C, D and E) for impairing (red),

enhancing (blue), non-significant (yellow) and all (grey) experiments. Lines and

whiskers express median and interquartile interval. (G) Distribution of statistical power

to detect the intermediate effect size of 37.2% (middle dashed line on C, D and E). (H)

Distribution of statistical power to detect the lower-bound effect size of 29.5% (left

dashed line on C, D and E). (I) Sample size vs. statistical power to detect the upper-

bound effect size of 45.6%. Continuous lines use the 50th percentile of coefficients of

variation for calculations, while dotted lines use the 25th and 75th percentiles. (J) Sample

size vs. statistical power to detect the intermediate effect size of 37.2%. (K) Sample size

vs. statistical power to detect the lower-bound effect size of 29.5%.

Based on each experiment's variance and sample size, we built power curves to

show how power varies according to the difference to be detected at α=0.05 for each

individual experiment (**Fig. 3C-E**). To detect the mean effect size of 45.6% found for

nominally effective interventions (i.e. those leading to statistically significant differences between groups), mean statistical power in our sample was 0.75 ± 0.26; 95% CI [0.72 - 0.78] (**Fig. 3F**). This estimate, however, is an optimistic, upper-bound calculation of the typical effect size of biologically effective interventions (from here on referred to as "upper-bound ES"): as only large effects will be detected by underpowered studies, basing calculations only on significant results leads to effect size inflation [14]. A more realistic estimate of effect size was obtained based only on experiments that achieved statistical power above 0.95 (n=60) in the first analysis (and are thus less subject to effect size inflation), leading to a mean effect size of 37.2%. Predictably, mean statistical power to detect this difference ("intermediate ES", **Fig. 3G**) fell to 0.65 ± 0.28 [0.62 - 0.68]. Using the mean effect size of all experiments ("lower-bound ES", 29.5%) led to an even lower power of 0.52 ± 0.29 [0.49 - 0.56] (**Fig. 3H**), although this estimate of a typical effect size is likely pessimistic, as it probably includes many true negative effects.

Interestingly, using mean absolute differences instead of relative ones to calculate statistical power led to a smaller number of experiments with very low power (**S6 Fig.**). This suggests that some of the underpowered experiments in the first analysis had low freezing levels in the reference group, as in this case even large relative differences will still be small when expressed in absolute terms for statistical analysis. Also of note is that, if one uses Cohen's traditional definitions of small (d=0.2), medium (d=0.5) and large (d=0.8) effect sizes [22] as the basis for calculations, mean power is 0.07 ± 0.01, 0.21 ± 0.07 and 0.44 ± 0.13, respectively (**S7 Fig.**). These much lower estimates reflect the fact that effect sizes are typically much larger in rodent fear conditioning than in psychology experiments, for which this arbitrary classification was originally devised, and suggests that it might not be applicable to other fields of science.

A practical application of these power curves is that we were able to calculate

the necessary sample size to achieve desired power for each effect size estimate,

considering the median coefficient of variation (as well as the 25th and 75th quartiles) of

experiments in our sample (**Fig. 3I-K**). Thus, for an experiment with typical variation,

around 15 animals per group are needed to achieve 80% power to detect our

'intermediate effect size' of 37.2%, which we consider our more realistic estimate for a

typical effect size in the field. Nevertheless, only 12.2% of comparisons in our sample

had a sample size of 15 or above in each experimental group, suggesting that such

calculations are seldom performed.

We also analyzed the distributions of statistical power at the level of articles

instead of individual experiments. Results for these analyses are shown in **S8 Fig.**, and

are generally similar to those obtained for the experiment-level analysis, except that the

long tail of non-significant experiments with large coefficients of variation is not

observed. This suggests that experiments with large variation and low power are

frequently found alongside others with adequate power within the same articles. It is

unclear, however, whether this means that the low power of some experiments is a

consequence of random fluctuations of experimental variance, or if these experiments

use protocols that lead to larger coefficients of variation – for example, by generating

lower mean levels of freezing (see **S2 Fig.**).

Correlation between effect sizes and statistical power/sample size

We next sought to correlate normalized effect size with sample size and

statistical power for each experiment. The presence of a negative correlation between

these variables has been considered an indirect measure of publication bias [23], as

articles with low power or sample size will be subject to effect size inflation caused by

selective reporting of significant findings [24]. In our analysis, no correlation was found

between effect size and sample size (**Fig. 4A**, r=0.0007, p=0.99); on the other hand, a

positive correlation between effect size and coefficient of variation was observed (**Fig.**

4B, r=0.37, p<0.0001). Part of this correlation was mediated by the association of both

variables with freezing levels (**S2 Fig.**), but the correlation remained significant after

adjustment for this variable (r=0.32, p<0.001).

Because of this, negative correlations between effect size and power were

observed for the three effect size estimates used (**Figs. 4C-E**), although they were larger

for the lower-bound estimate (**Fig. 4E**, r=-0.21, p<0.0001) than for the intermediate

(**Fig. 4D**, r=-0.16, p=0.003) and upper-bound (**Fig. 4C**, r=-0.12, p=0.03) ones due to a

ceiling effect on power. This negative correlation is observed even when power is

calculated based on absolute differences (**S9 Fig.**), for which the correlation between

coefficients of variation and reference freezing levels is in the opposite direction of that

observed with relative differences (see **S2 Fig.**). This strongly suggests that the

correlation represents a real phenomenon related to publication bias and/or effect size

inflation, and is not merely due to the correlation of both variables with freezing levels.

A correlation between effect size and power is also observed when both are calculated

on the basis of standardized mean differences (i.e. Cohen's d) (**S10 Fig.**). In this case,

the line separating significant and non-significant results for a given sample size is

clearer, as significance is more directly related to standardized mean differences.

Expressing effect sizes in Cohen's d also makes effect size inflation in experiments with

low sample size and power more obvious.

Figure 4. Correlations between effect size, variation and statistical power. (A)

Correlation between normalized effect size and mean sample size. No correlation is

found (r=0.0007, p=0.99; r=-0.26, p=0.64 after adjustment), although sample size

variation is limited. (B) Correlation between normalized effect size and coefficient of

variation. Correlation of the whole sample of experiments yields r=0.37, p<0.0001*

(n=336; r=0.32, p<0.001 after adjustment for freezing levels). (C) Correlation between

normalized effect size and statistical power based on upper-bound effect size of 45.6%.

Correlation of the whole sample of experiments yields r=-0.12, p=0.03 (r=0.11, p=0.84

after adjustment for freezing levels), but distribution is skewed due to a ceiling effect on

power. (D) Correlation between normalized effect size and statistical power based on

intermediate effect size of 37.2%; r=-0.16, p=0.003* (r=-0.16, p=0.48 after adjustment).

(E) Correlation between normalized effect size and statistical power based on lower-

bound effect size of 29.5%; r=-0.21, p<0.0001* (r=-0.1, p=0.06 after adjustment).

Asterisks indicate significant results according to Holm-Sidak correction for 28

experiment-level correlations.

Interestingly, the correlation between effect size and power was driven by a

scarcity of experiments with large effect size and high power. This raises the possibility

that truly large effects are unusual in fear conditioning, and that some of the large effect

sizes among low-powered experiments in our sample are inflated. On the other hand, a

pattern classically suggesting publication bias – i.e. a scarcity of low-powered

experiments with small effects [23] – is not observed. It should be noted, however, that

our analysis focused on individual experiments within articles, meaning that non-

significant results were usually presented alongside other experiments with significant

differences; thus, this analysis does not allow us to assess publication bias at the level of

articles.

Effects of methodological variables on the distributions of effect sizes and

coefficients of variation.

We next examined whether the distributions of effect sizes and coefficients of

variation were influenced by type of conditioning, species or sex of the animals (**Fig. 5**).

Mean normalized effect size was slightly larger in contextual than in cued fear

conditioning (33.2% vs. 24.4%, Student's t test $p < 0.0001$) and markedly larger in males

than in females (30.3% vs. 18.9% vs. 34.2% for experiments using both, one-way

ANOVA, $p = 0.004$), but roughly equivalent between mice and rats (29.8% vs. 29.1%,

$p = 0.76$). Coefficients of variation were higher in contextual conditioning (0.51 vs. 0.41,

Student's t test $p = 0.001$), in experiments using animals of both sexes (0.62 vs. 0.44 in

males and 0.41 in females, one-way ANOVA, $p < 0.0001$), and in those using mice (0.50

vs. 0.42, Student's t test, $p = 0.008$), although the latter difference was not statistically

significant after correction for multiple comparisons. All of these associations should be

considered correlational and not causal, as specific types of conditioning or animals of a

particular species or sex might be more frequently used for testing interventions with

particularly high or low effect sizes. Also of note is the fact that experiments on males were 7.7 times more common than those on females in our sample (277 vs. 36), indicating a strong preference of researchers for using male animals.

Figure 5. Effect sizes and coefficients of variation across different protocols, species and sexes. Colors indicate memory-enhancing (red), memory-impairing (blue) or non-effective (yellow) experiments, all of which are pooled for analysis. Lines and whiskers express median and interquartile interval. (A) Distribution of effect sizes across cued (n=171) and contextual (n=239) conditioning protocols. Student's t test,

p<0.0001*. (B) Coefficients of variation across cued (n=145) and contextual (n=191)

conditioning protocols. Student's t test, p=0.001*. (C) Distribution of effect sizes across

experiments using mice (n=237) or rats (n=173). Student's t test, p=0.76. (D)

Coefficients of variation across experiments using mice (n=193) or rats (n=143).

Student's t test, p=0.008. (E) Distribution of effect sizes across experiments using male

(n=277), female (n=36) or both (n=67) sexes. One-way ANOVA, p=0.004*; Tukey's

post-hoc test, male vs. female p=0.01, male vs. both p=0.40, female vs. both p=0.003.

30 experiments were excluded from this analysis for not stating the sex of animals. (F)

Coefficients of variation across experiments using male (n=233), female (n=28) or both

(n=60) sexes. One-way ANOVA, p<0.0001*; Tukey's test, male vs. female p=0.85,

male vs. both p<0.0001, female vs. both p=0.0006. For coefficient of variation analyses,

74 experiments were excluded due to lack of information on sample size for individual

groups. Asterisks indicate significant results according to Holm-Sidak correction for 14

experiment-level comparisons.

We also examined whether effect sizes and coefficients of variation differed

systematically according to the type, timing or anatomical site of intervention (**S11**

Fig.). Effect sizes did not differ significantly between surgical, pharmacological, genetic

and behavioral interventions (38.7% vs. 28.1% vs. 30.5% vs. 25.8% one-way ANOVA,

p=0.12), although there was a trend for greater effects with surgical interventions

(which were uncommon in our sample). No differences were found between the mean

effect sizes of systemic and intracerebral interventions (28.7% vs. 30.3%, Student's t

test, p=0.45) or between those of pre- and post-training interventions (30.5% vs. 25.4%,

Student's t test, p=0.07), although pre-training interventions had slightly higher

coefficients of variation (0.49 vs 0.37, Student's t test p=0.0015). Coefficients of

variation did not differ significantly between surgical, pharmacological, genetic and

behavioral interventions (0.41 vs. 0.43 vs. 0.50 vs. 0.50, one-way ANOVA p=0.08) or

between systemic and intracerebral interventions (0.49 vs. 0.45, Student's t test p=0.15).

Once again, these differences can only be considered correlational and not causal.

Risk of bias indicators and their relationship with effect size and power

As previous studies have shown that measures to reduce risk of bias are not

widely reported in animal research [12,13], we investigated the prevalence of these

measures in our sample of fear conditioning articles, and evaluated whether they were

correlated with effect sizes or power. **Table 1** shows the percentage of articles reporting

7 items thought to reduce risk of bias in animal studies, adapted and expanded from the

CAMARADES checklist [25]. Although some items were reported in most articles

(statement of compliance with animal regulations, adequate description of sample size,

blinding), others were virtually inexistent, such as the presence of a sample size

calculation (1 article) and compliance with the ARRIVE guidelines [26] (0 articles).

Contrary to previous reports in other areas [27–30], however, no significant association

was found between reporting of these indicators and either the percentage of significant

experiments, the mean effect size of effective interventions or the mean statistical power

of experiments in our sample (**S12 Fig.**). The region of origin of the article also had no

correlation with either of these variables (**S13 Fig.**). Nevertheless, it should be noted

that this analysis used only experiments on fear conditioning acquisition or

consolidation, which were not necessarily the only results or the main findings

presented in these articles. Thus, it is possible that other results in the article might have

shown higher correlation with risk of bias indicators.

Quality assessment	Randomization of allocation	Blinded or automated	Sample size calculation	Exact sample size description	Statement of compliance with regulatory	Statement on conflict	Statement of compliance

item	assessment				requirements	of interest	with ARRIVE
Number of articles (%)	18/77 (23.4%)	92/122 (75.4%)	1/122 (0.8%)	98/122 (80.3%)	118/122 (96.7%)	66/122 (54.1%)	0/122 (0%)

Table 1. Number of articles including quality assessment items. Percentages were calculated using all 122 articles, except in the case of randomization, which was calculated based on 77 articles, as it is not applicable to genetic interventions. In the case of blinding, 68 articles used automated analysis and 24 used blinded observers, totaling 92 articles scored for this item.

Correlation between effect sizes/statistical power and description of results

Given the wide distribution of effect sizes and statistical power in the literature on fear conditioning learning, we tried to determine whether these were taken into account by authors when describing results in the text. For each included comparison, we extracted the words or phrases describing the results of that experiment in the text or figure legends, and asked 14 behavioral neuroscience researchers to classify them according to the implicit information they contained about effect size. For comparisons with significant differences, terms were to be classified as implying strong (i.e. large effect size) or weak (i.e. small effect size) effects, or as neutral terms (i.e. those from which effect size could not be deduced). For non-significant differences, terms were to be classified as implying similarity between groups, as suggesting a trend towards difference, or as neutral terms (i.e. those from which the presence or absence of a trend could not be deduced). From the average of these classifications, we defined a score for each term (**S1 and S2 Tables**) and correlated these scores with the actual effect size and statistical power of experiments.

Agreement between researchers over classification was low, especially for terms

describing significant differences: single measures intraclass correlation coefficients

(reflecting the reliability of individual researchers when compared to the whole sample)

were 0.234 for significant interventions and 0.597 for non-significant ones, while

average measures coefficients (reflecting the aggregated reliability of the sample) were

0.839 and 0.962, respectively. This, along with a trend for the use of terms with little

effect size information ("increase", "decrease", "significantly more", "significantly

less", etc.), led most terms describing effective interventions to receive intermediate

scores approaching 1 (i.e. neutral). For these interventions, no correlations were

observed between this score and either effect size (r=-0.05, p=0.48) or statistical power

(r=0.03, p=0.73) (**Fig. 6A and 6B**). For non-effective interventions, a significant

correlation between description score and effect size was observed (**Fig 6C**, r=0.28,

p=0.0002), as larger effect sizes were associated with terms indicating a trend for

difference. Still, no correlation was observed between textual descriptions of results and

power (**Fig 6D**, r=0.03, p=0.74). Moreover, statistical power was rarely mentioned in

the textual description of results – the term "power" was used in this context in only 4

articles– suggesting that it is largely ignored when discussing findings, as shown in

other areas of research [31].

Figure 6. Correlation between description of results and effect size/statistical power. Description scores refer to the mean score given by 14 neuroscience researchers who rated terms as "weak" (0), "neutral" (1) or "strong" (2) in the case of those describing significant differences, or as "similar" (0), "neutral" (1) or "trend" (2) in the case of those describing non-significant ones. (A) Correlation between normalized effect size and description score for significant results. r=-0.05, p=0.48 (n=195). (B) Correlation between statistical power and description score for significant results. r=0.03, p=0.73 (n=155). (C) Correlation between normalized effect size and description score for non-significant results. r=0.28, p=0.0002* (n=174). (D) Correlation between upper-bound estimate of statistical power and description score for non-significant results. r=0.03, p=0.74 (n=146). Asterisk indicates significant result according to Holm-Sidak correction for 28 experiment-level correlations.

 Correlations of effect size, power and study quality with article citations

Finally, we investigated whether the percentage of significant experiments

reported in each article, mean effect size for effective interventions, mean statistical

power or a composite study quality score (aggregating the 7 risk of bias indicators

described in **Table 1**) correlated with article impact, as measured by the number of

citations (**Fig. 7**) and the impact factor of the publishing journal (**S14 Fig.**). None of the

correlations was significant after adjustment for multiple comparisons, although a weak

positive correlation was observed between study quality score and impact factor

(r=0.22, p=0.01), driven by associations of higher impact factors with blinding

(Student's t test with Welch's correction, p=0.0001), conflict of interest reporting

(Student's t test with Welch's correction, p=0.03) and exact sample size description

(Student's t test, p=0.03). It should be noted that the distribution of impact factors and

citations is heavily skewed, limiting the use of linear correlations as planned in the

original protocol – nevertheless, exploratory non-parametric analysis of the data

confirmed the lack of significance of correlations. Once again, our data refers only to

experiments on fear conditioning acquisition or consolidation – therefore, other data in

the articles could feasibly account for the variation in impact factor and citations.

Figure 7. Correlation between citations and percentage of significant experiments, effect size and statistical power. Citations were obtained for all articles on August 26th, 2016. (A) Correlation between % of significant results per article and citations. r=-0.03, p=0.75 (n=121). (B) Correlation between mean normalized effect size of effective interventions and citations. r=0.097, p=0.34 (n=98). (C) Correlation between mean statistical power (upper-bound estimate) and citations. r=-0.08, p=0.40 (n=104). (D) Correlation between study quality score and citations. r=0.09, p=0.31 (n=121). According to Holm-Sidak correction for 8 article-level correlations, none is significant.

Discussion

In light of the low reproducibility of published studies in various fields of biomedical research [32–34] which is thought by many to be a consequence of low statistical power and excessive reliance on significance tests [8,16] calls have been

made to report effect sizes and confidence intervals alongside or in place of p values

[4,6,7,9] and to increase statistical power [14,31,35]. However, it is unclear whether

these proposals have had much impact on most fields of basic science. We have taken

one particular memory task in rodents, in which outcomes and effect sizes are described

in a standardized way and are thus comparable across studies, in order to analyze how

these two concepts are dealt with in the study of fear learning.

Our first main finding is that most amnestic interventions targeting fear

acquisition or consolidation cause partial effects, with residual freezing remaining

significantly above pre-conditioning levels in 78% of the experiments with available

data. Moreover, most of the large effect sizes in our sample were found in

underpowered studies, suggesting that they could represent inflated estimates [24]. This

is not necessarily unexpected: as fear memories depend on a well distributed network,

both anatomically and molecularly [19], it seems natural that most interventions

directed at a specific site or pharmacological target will modulate learning rather than

fully block it. This creates a problem, however, when effect sizes are not considered in

the analysis of experiments, as it is not possible to differentiate essential mechanisms of

memory formation from modulatory influences on the basis of statistical significance

alone. This can lead to a situation in which accumulating evidence, even if correct, can

confuse rather than advance understanding, as has been suggested to occur in fields such

as long-term potentiation [15] and apoptosis [36].

Matters are complicated further by the possibility that many of these findings are

false positives and/or false negatives. The prevalence of both in relation to true positives

and negatives depends crucially on statistical power, which in turn depends on sample

size. Calculating the actual power of published experiments is challenging, as the

difference used for the calculations should not be based on the observed results – which

leads power analysis to become circular [18,37]. Thus, statistical power depend on expected effect sizes, which are arbitrary by nature – although they can sometimes be estimated from meta-analyses [14], which were not performed in this study due to the large variety of heterogeneous interventions. However, by considering the mean effect size for well-powered experiments in our sample, we arrived at an estimate of around 37.2% that might be considered "typical" for a published experiment with an intervention affecting fear conditioning acquisition or consolidation. Using the sample size and variation for each experiment, we found mean statistical power to detect this effect size to be 65% in our sample.

As sample size calculations are exceedingly rare, and insufficient power seems to be the norm in other fields of neuroscience as well [14,18], it is quite possible that classically used sample sizes in behavioral neuroscience (and perhaps in other fields of basic science) might thus be insufficient. Considering median variances and our intermediate effect size estimate, the ideal sample size to achieve 80% power would be around 15 animals per group. This number, however, was reached in only 12.2% of cases in our sample, as most experiments had sample sizes of 8 to 12, informally considered to be standard in the field. This seems to confirm recent models suggesting that current incentives in science favor the publication of underpowered studies [16,38], although they could also be due to restrictions on animal use imposed by ethical regulations. That said, average power in our sample for typical effect sizes was higher than those described in other areas of neuroscience by Button et al. [14]; however, this could reflect the fact that effect sizes in their study were calculated by meta-analysis, and might be smaller than those derived by our method of estimation, or underestimated due to the inclusion of negative results [18]. One should also note that the abovementioned power estimates were found to vary widely across subfields of

neuroscience [18] – in this sense, the power distribution of fear conditioning studies

seems to resemble those found for psychology and neurochemistry studies, in which a

reasonable number of well-powered studies coexist with underpowered ones.

On the other hand, our statistical power to detect Cohen's definitions of small,

medium and large effects [22] were even lower than those recently reported in cognitive

neuroscience studies by Szucs and Ioannidis (2017). That said, our data provides a

strong cautionary note against the use of these arbitrary definitions, originally devised

for psychology studies, in calculations of statistical power, as 88.7% of statistically

significant experiments (or 48.2% of the whole sample) fell into the "large" category of

Cohen's original proposal. This suggests that laboratory studies in rodents have larger

effects than those found in human psychology (an unsurprising finding, given the

greater invasiveness of the interventions), as has also been found in meta-analyses

studying similar treatments in laboratory animals and humans [39], demonstrating that

what constitutes a small or large effect can vary between different fields of science.

An old-established truism in the behavioral neuroscience field – as well as in

other fields of basic science – is that experiments in females tend to yield more variable

results due to estrous cycle variations [40]. However, at least in our analysis,

coefficients of variation were similar between experiments in males and females (and

predictably higher in experiments using both), as has been found in other areas of

science [41,42] suggesting that this belief is false. Nevertheless, adherence to it likely

accounts for the vast preponderance of experiments on male animals, which were nearly

8 times more common than those in females in our sample – a sex bias greater than

those described for most fields [43] although smaller than that recently reported for

rodent models of anxiety [44]. Previous work in clinical [45] and preclinical [40,43]

data has pointed out the drawbacks of concentrating experiments in male populations.

However, despite calls for increasing the number of studies on females [46] this problem remains strikingly present in the fear learning field.

Concerning risk of bias indicators, the prevalence found in our sample was roughly similar to previous reports on animal studies for randomization and conflict of interest reporting [13] but were distinctly higher for blinded assessment of outcome, largely because 59% of articles used automated software to measure freezing, which we considered to be equivalent to blinded assessment. If one considers only articles reporting manual scoring of freezing, however, blinding was reported in 57% of cases, which is still higher than most areas of preclinical science [13]. As described previously in many fields [12,13,31] sample size calculations were almost non-existent, which helps to explain why many experiments are underpowered. Interestingly, although we analyzed a sample of papers published 3 years after the ARRIVE guidelines they were not mentioned in any of the articles, suggesting that their impact, at least in the field of behavioral neuroscience, was still rather limited at this time.

Contrary to previous studies, however, we did not detect an impact of these risk of bias indicators on article-level measures such as percentage of fear conditioning learning experiments with significant results, mean effect size of significant experiments and mean statistical power. This could mean that, compared to preclinical studies, bias towards positive results is lower in studies on fear learning. However, it seems more likely that, as we selected particular experiments within papers containing other results, we were not as likely to detect effects of bias on article-level measures. As basic science articles typically contain numerous results, it is perhaps less likely that all comparisons will be subject to bias towards positive findings. Moreover, the experiments in our sample probably included negative controls for other findings, which might have been expected to yield non-significant results. Thus, although our results do

not indicate an impact of bias on article-level results, they should not be taken as

evidence that this does not occur.

The same reasoning applies for the evaluation of publication bias, in which the

experiments we analyzed could have been published along with positive ones.

Nevertheless, we were still able to detect a negative correlation between effect size and

statistical power, suggesting effect size inflation due to low statistical power to be

present in studies on fear conditioning learning. Although the pattern we detected was

less suggestive of actual publication bias, our capability to detect it was likely smaller

due to the choice to use experiments within articles. Other methods to detect publication

bias, such as the Ioannidis excess significance test [47] and the use of p-value

distributions [48–50] were also considered, but found to be inappropriate for use with

our methodology (in the first case due to the absence of a meta-analytic effect estimate,

and in the second because exact p values were infrequently provided in articles).

One of the most interesting findings of our article was the lack of correlation of

effect sizes inferred from textual description of results with the actual effect sizes of

significant experiments, as well as with statistical power. Although this suggests that

these measures are not usually considered in the interpretation of results, there are

caveats to this data. First of all, agreement between what words describe a "strong" or

"weak" effect between researchers evaluating them was strikingly low, suggesting that

written language is a poor descriptor for quantitative data. Moreover, the fact that most

terms used to describe differences were neutral to effect sizes (e.g. "significantly

higher", "significantly lower", etc.) limited our ability to detect a correlation. That said,

the high prevalence of neutral terms by itself is evidence that effect sizes are not usually

taken into account when reporting results, as differences tend to be described in the text

by their statistical significance only.

This point is especially important to consider in the light of recent calls for basic

science to use data synthesis tools such as meta-analysis [11] and formal or informal

Bayesian inference [2,8,10,51]. In both of these cases, the incremental effect of each

new experiment on researchers' beliefs on the veracity of a finding is dependent both on

the effect size of the result and on its statistical significance. However, even exact p

values were uncommonly reported in our sample, with the majority of articles

describing p as being above or below a threshold value. This seems to suggest that

researchers in the field indeed tend to consider statistical significance as a binary

outcome, and might not be quite ready or willing to move towards Bayesian logic,

which would require a major paradigm shift in the way results are reported and

discussed.

An interesting question is that, if researches in the field indeed were to move

away from null-significance hypothesis testing, the concept of statistical power as it is

defined today would largely lose its meaning (as it is intrinsically linked to the idea of a

significance threshold). Nevertheless, the necessity of adequate sample size for

statistical robustness would remain – in this case, not in order to detect significant

differences and prevent false-negatives and false-positives, but to estimate effect sizes

with adequate precision. The current notion of statistical power to detect a given

difference could thus be replaced with a desired confidence interval for the obtained

result when performing sample size calculations – a formulation that might be useful in

terms of differentiating biologically significant results from irrelevant ones.

Concerning article impact metrics, our results are in line with previous work

showing that journal impact factor does not correlate with statistical power [14] or with

most risk of bias indicators [13]. Furthermore, we showed that, in articles on fear

conditioning, this lack of correlation also occurs for the percentage of significant

experiments and the mean effect size for significant differences, and that it extends to

citations measured over 2 subsequent years. That said, our article-level analysis was

limited by the fact that, for many articles, the included experiments represented a

minority of the findings. Moreover, most articles tend to cluster around intermediate

impact factors (i.e. between 3 and 6) and relatively low (< 20) citation numbers. Thus,

our methodology might not have been adequate to detect correlations between these

metrics with article-wide effect size and power estimates.

The choice to focus on a particular type of experiment – in this case,

interventions directed at rodent fear conditioning acquisition or consolidation – is both

one of the main strengths and the major limitation of our findings. On one hand, it

allows us to look at effect sizes that are truly on the same scale, as fear conditioning

protocols tend to be reasonably similar across laboratories, and all included experiments

described their results using the same metric. Thus, the studied effect sizes are not

abstract and have real-life meaning. On the other hand, this decision limits our

conclusions to this specific field of science, and also weakens our article-level

conclusions, as most articles had only a fraction of their experiments analyzed.

Dealing with multiple experiments using different outcomes presents a major

challenge for meta-research in basic science, and all alternatives present limitations. A

radically opposite approach of converting all effect sizes in a field to a single metric

(e.g. Pearson's r, Cohen's d, etc.) has been used by other researchers investigating

similar topics in neuroscience and psychology [17,23,31,35]. Although normalizing

effect sizes allows one to obtain results from a wider field, it also leads them to be

abstract and not as readily understandable by experimental researchers. Moreover, this

approach can lead to the aggregation of results from disparate types of experiments for

which effect sizes are not in the same scale, leading to important distortions in

calculating power for individual experiments. Finally, recent evidence indicates that,

even within neuroscience, features such as statistical power have very different

distributions across subfields [18], suggesting that surveys of individual areas are likely

to be more reliable for studying them.

In our case, studying the concrete scenario of a specific methodology leads to

more readily applicable suggestions for experimental researchers, such as the rule-of-

thumb recommendation that the average number of animals per group in a fear

conditioning experiments to achieve 80% power would be around 15 for typical effect

sizes and variances. Our approach also allowed us to detect correlations between results

and specific methodological factors (e.g. context vs. cued conditioning, female vs. male

animals) that would not be apparent if multiple types of experiments were pooled

together. Still, to provide more solid conclusions on the causal influence of these factors

on experimental results, even our methodology has too wide a focus, as analyzing

multiple interventions limits our possibilities to perform meta-analysis and meta-

regression to control for confounding variables. Follow-up studies with more specific

aims (i.e. meta-analyses of specific interventions in fear conditioning) are thus

warranted to understand the variation between results in the field.

Finally, it is important to note that, while our study has led to some illuminating

conclusions, they are inherently limited to the methodology under study. Thus,

extrapolating our findings to other types of behavioral studies, not to mention other

fields of science, requires data to be collected for each specific subfield. While this

might appear herculean at first glance, it is easily achievable if scientists working within

specific domains start to design and perform their own systematic reviews. Only

through this dissemination of meta-research across different areas of science will we be

able to develop solutions that, by respecting the particularities of individual subfields,

will be accepted enough to have an impact on research reproducibility.

Materials and Methods

The full protocol of data selection, extraction and analysis was initially planned

on the basis of a pilot analysis of 30 papers, and was registered, reviewed and published

ahead of full data extraction [20]. In brief, we searched PubMed for the term "fear

conditioning" AND ("learning" OR "consolidation" OR "acquisition") AND ("mouse"

OR "mice" OR "rat" OR "rats")" to obtain all articles published online in 2013. Titles

and abstracts were first scanned for articles presenting original results involving fear

conditioning in rodents that were written in English. Selected articles underwent full-

text screening for selection of experiments that (a) described the effects of a single

intervention on fear conditioning acquisition or consolidation, (b) had a clearly defined

control group to which the experimental group is compared to, (c) used freezing

behavior as a measure of conditioned fear in a test session and (d) had available data on

mean freezing, SD or SEM, as well as on the significance of the comparison. Articles

were screened by one of two investigators (C.F.D.C. or T.C.M.) for relevant data and

were analyzed by the other – thus, all included experiments were dual-reviewed.

Only experiments analyzing the effect of interventions performed before or up to

6 hours after the training session (i.e. those affecting fear conditioning acquisition or its

immediate consolidation) were included. Data on mean freezing and SD or SEM were

obtained for each group from the text when available; otherwise, it was extracted using

Gsys 2.4.6 software (Hokkaido University Nuclear Reaction Data Centre). When exact

sample size for each group was available, the experiment was used for the analysis of

effect size and statistical power – otherwise, only effect size was obtained, and the

experiment was excluded from power analysis. For individual experiments, study

design characteristics were also obtained, including species and sex of the animals, type

of conditioning protocol, type, timing and site of intervention.

From each comparison, we also obtained the description term used by the

authors in the results session of the paper. Classification of the terms used to describe

effects (**S1 and S2 Tables**) was based on a blinded assessment of words or phrases by a

pool of 14 researchers who were fluent or native speakers of English and had current or

past experience in the field of behavioral neuroscience. Categories were given a score

from 0 to 2 in order of magnitude (i.e. 0 = weak, 1 = neutral, 2 = strong for significant

results; 0 = similar, 1 = neutral, 2 = trend for non-significant results), and the average

results for all researchers was used as a continuous variable for analysis.

Apart from experiment-level variables, we also extracted article-level data such

as impact factor of the journal in which it was published (based on the 2013 Journal

Citations Report), number of citations (obtained for all articles on August 26[th] 2016),

country of origin (defined by the corresponding author's affiliation) and the 7 risk of

bias indicators described on **Table 1**. For article-level correlations, we compiled these

measures into a normalized score.

After completion of data extraction, all calculations and analyses were

performed according to the previously specified protocol. Specific details of

calculations (as well as the raw data used) are presented as **Supplementary Data**. After

this, the following additional analyses were performed in an exploratory fashion:

(a) To confirm that residual freezing levels after memory-impairing

interventions were indeed above training values, demonstrating that most amnestic

intervention have partial effects, we extracted pre-conditioning freezing levels from

training sessions when these were available. These levels were obtained for pre-shock

periods only, and separated as baseline values for contextual (i.e. freezing in the absence

of tone) or tone conditioning (i.e. freezing in the presence of a tone, but before shock),

as displayed in **S5 Fig**. These were compared to the corresponding test session values

for treated groups in memory-impairing interventions by an unpaired *t* test based on the

extracted means, SD or SEM and sample size when these were available.

(b) In the original protocol, only the mean of all effective interventions (i.e.

upper-bound effect size) was planned as a point estimate to be used for power

calculations, although we acknowledged this to be optimistic [20]. We later decided to

perform power calculations based on the mean effect size of the experiments achieving

power above 0.95 on the first analysis (i.e. intermediate effect size) to avoid effect size

inflation, as we reached the conclusion that this would provide a more realistic estimate.

Additionally, we calculated power based on the mean effect size of the whole sample of

experiments as a lower-bound estimate, and presented all three estimates in the results

section and figures.

(c) In order to evaluate whether the distribution of effect sizes and statistical

power varied if effect sizes were defined as absolute differences in freezing levels

instead of relative ones, we repeated the analyses in **Figs. 2, 3** and **4** using absolute

differences in **S1 Fig., S6 Fig.** and **S9 Fig.**. This proved to be particularly important to

demonstrate that correlations between effect sizes and power were not the consequence

of a confounding association of both variables with coefficients of variation. We also

repeated power and correlation analyses using effect sizes as standardized mean

differences (e.g. Cohen's d) in **S7 Fig.** and **S10 Fig.**)

(d) To further evaluate the possible impact of the negative correlation between

coefficients of variation and freezing levels on our results, we decided to use freezing

levels as a covariate in the correlations shown in **Fig. 4**. We also checked whether

adding freezing levels as a covariate influenced the statistical analyses in **Fig. 5**, **Fig. 6**

and **S11 Fig.**, but as this did not have a significant impact on the results in these figures,

we only reported the originally planned analyses.

(e) All of our planned analyses were parametric; after extraction, however, it

was clear that some of the data deviated from a normal distribution (especially in the

case of power estimates, citation counts and impact factor). Because of this, we

performed additional non-parametric analyses for the correlations of citations and

impact factor with percentage of significant results, mean normalized effect size,

statistical power and study quality score.

(f) In the protocol, we had planned to test correlations between normalized effect

sizes and statistical power, mean sample size and absolute freezing levels (using the

group with the highest freezing). After analyzing the results, we also decided to

correlate normalized effect sizes with coefficients of variation (as this, rather than

sample size, seemed to explain the lower power of non-significant results), additional

power estimates (as using our original estimate led to a ceiling effect) and different

estimates of freezing based on the control group or on the mean freezing of both groups

(to compare these forms of normalization with the one we chose).

767 (g) Due to the correlation of study quality assessment with journal impact

768 factor, we performed an exploratory analysis of the correlation of this metric with each

769 of the individual quality assessment items by performing a Student's t test (corrected for

770 unequal variances by Welch's correction) between the impact factors of studies with

771 and without each item.

772 (h) Because of the additional analyses above, we adjusted the number of

773 comparisons/correlations used as the basis of the Holm-Sidak correction for multiple

774 comparisons. The total numbers used for each correction were 14 for experiment-level

775 comparisons, 17 for article-level comparisons, 28 for experiment-level correlations and

776 8 for article-level correlations, leading to significance thresholds between 0.003 and

777 0.05.

778        **References**

779 1. Nuzzo R. Statistical errors. Nature. 2014;506: 150–152.

780 2. Colquhoun D. An investigation of the false discovery rate and the

781 misinterpretation of p-values. R Soc Open Sci. 2014;1: 140216–140216.

782 3. Altman N, Krzywinski M. Points of significance: P values and the search for

783 significance. Nat Methods. 2016;14: 3–4.

784 4. Halsey LG, Curran-Everett D, Vowler SL, Drummond GB. The fickle P value

785 generates irreproducible results. Nat Methods. 2015;12: 179–185.

786 5. Wasserstein RL, Lazar NA. The ASA's statement on p-values : context, process,

787 and purpose. 2016;70: 129-133.

788 6. Greenland S, Senn SJ, Rothman KJ, Carlin JB, Poole C, Goodman SN, et al.

789 Statistical tests, P values, confidence intervals, and power: a guide to misinterpretations.

790 Eur J Epidemiol. 2016;31: 337-350.

7.   Nakagawa S, Cuthill IC. Effect size, confidence interval and statistical

significance: a practical guide for biologists. Biol Rev. 2007;82: 591–605.

8.   Ioannidis JPA. Why most published research findings are false. PLoS Med.

2005;2: 696–701.

9.   Trafimow D, Marks M. Editorial. Basic Appl Soc Psych. 2015;37: 1–2.

10.   Goodman SN, Fanelli D, Ioannidis JPA. What does research reproducibility

mean? Sci Transl Med. 2016;8: 1–6.

11.   Vesterinen HM, Sena ES, Egan KJ, Hirst TC, Churolov L, Currie GL, et al.

Meta-analysis of data from animal studies: A practical guide. J Neurosci Methods.

2014;221: 92–102.

12.   Kilkenny C, Parsons N, Kadyszewski E, Festing MFW, Cuthill IC, Fry D, et al.

Survey of the quality of experimental design, statistical analysis and reporting of

research using animals. PLoS One. 2009;4: e7824.

13.   Macleod MR, Lawson McLean A, Kyriakopoulou A, Serghiou S, de Wilde A,

Sherratt N, et al. Risk of bias in reports of in vivo research: a focus for improvement.

PLoS Biol. 2015;13: e1002273.

14.   Button KS, Ioannidis JP, Mokrysz C, Nosek B, Flint J, Robinson ESJ, et al.

Power failure: why small sample size undermines the reliability of neuroscience. Nat

Rev Neurosci. 2013;14: 365–76.

15.   Sanes JR, Lichtman JW. Can molecules explain long-term potentiation? Nat

Neurosci. 1999;2: 597–604.

16.   Higginson AD, Munafò MR. Current incentives for scientists lead to

underpowered studies with erroneous conclusions. PLoS Biol. 2016;14: e2000995.

17.   Szucs D, Ioannidis JPA. Empirical assessment of published effect sizes and

power in the recent cognitive neuroscience and psychology literature. PLoS Biol.

2017;15: e2000797.

18.   Nord CL, Valton V, Wood J, Roiser JP. Power-up: a reanalysis of "power

failure" in neuroscience using mixture modelling. J Neurosci. 2017;37: 3592–16.

19.   Maren S. Neurobiology of Pavlovian fear conditioning. Annu Rev Neurosci.

2001;24: 897–931.

20.   Moulin TC, Carneiro CFD, Macleod MR, Amaral OB. Protocol for a systematic

review of effect sizes and statistical power in the rodent fear conditioning literature.

Evid Based Preclin Med. 2016;3: 24–32.

21.   Moher D, Liberati A, Tetzlaff J, Altman DG, Altman D, Antes G, et al. Preferred

reporting items for systematic reviews and meta-analyses: The PRISMA statement.

PLoS Med. 2009;6: e1000097.

22.   Cohen J. Statistical power analysis for the behavioral sciences (2nd ed.). New

York: Academic Press; 1977.

23.   Kuhberger A, Fritz A, Scherndl T. Publication bias in psychology: A diagnosis

based on the correlation between effect size and sample size. PLoS One. 2014;9:

e105825.

24.   Ioannidis JPA. Why most discovered true associations are inflated.

Epidemiology. 2008;19: 640–648.

25.   Sena E, van der Worp HB, Howells D, Macleod M. How can we improve the

pre-clinical development of drugs for stroke? Trends Neurosci. 2007;30: 433–9.

26.   Kilkenny C, Browne WJ, Cuthill IC, Emerson M, Altman DG. Improving

bioscience research reporting: the ARRIVE guidelines for reporting animal research.

PLoS Biol. 2010;8: e1000412.

27.   Macleod MR, van der Worp HB, Sena ES, Howells DW, Dirnagl U, Donnan

GA. Evidence for the efficacy of NXY-059 in experimental focal cerebral ischaemia is

confounded by study quality. Stroke. 2008;39: 2824–9.

28.   Vesterinen HM, Sena ES, Ffrench-Constant C, Williams A, Chandran S,

Macleod MR. Improving the translational hit of experimental treatments in multiple

sclerosis. Mult Scler. 2010;16: 1044–55.

29.   Rooke EDM, Vesterinen HM, Sena ES, Egan KJ, Macleod MR. Dopamine

agonists in animal models of Parkinson's disease: A systematic review and meta-

analysis. Parkinsonism Relat Disord. 2011;17: 313–320.

30.   Currie GL, Delaney A, Bennett MI, Dickenson AH, Egan KJ, Vesterinen HM, et

al. Animal models of bone cancer pain: Systematic review and meta-analyses. Pain.

2013;154: 917–926.

31.   Sedlmeier P, Gigerenzer G. Do studies of statistical power have an effect on the

power of studies? 1989;105: 309–316.

32.   Scott S, Kranz JE, Cole J, Lincecum JM, Thompson K, Kelly N, et al. Design,

power, and interpretation of studies in the standard murine model of ALS. Amyotroph

Lateral Scler. 2008;9: 4–15.

33.   Prinz F, Schlange T, Asadullah K. Believe it or not: how much can we rely on

published data on potential drug targets? Nat Rev Drug Discov. 2011;10: 712.

34.   Begley CG, Ellis LM. Drug development: Raise standards for preclinical cancer

research. Nature. 2012;483: 531–3.

35.   Cohen J. The statistical power of abnormal-social psychological research: a

review. J Abnorm Soc Psychol. 1962;65: 145–53.

36.   Lazebnik Y. Can a biologist fix a radio?--Or, what I learned while studying

apoptosis. Cancer Cell. 2002;2: 179–82.

37.   Goodman SN, Berlin JA. The use of predicted confidence intervals when

planning experiments and the misuse of power when interpreting results. Ann Intern

Med. 1994; 121: 200-206.

38.   Smaldino PE, McElreath R. The natural selection of bad science. R Soc Open

Sci. 2016;3: 160384.

39.   Norberg MM, Krystal JH, Tolin DF. A meta-analysis of D-cycloserine and the

facilitation of fear extinction and exposure therapy. Biol Psychiatry. 2008;63: 1118–

1126.

40.   Wald C, Wu C. Of mice and women: The bias in animal models. Science.

2010;327: 1571–1572.

41.   Mogil JS, Chanda ML. The case for the inclusion of female subjects in basic

science studies of pain. Pain. 2005;117: 1–5.

42.   Prendergast BJ, Onishi KG, Zucker I. Female mice liberated for inclusion in

neuroscience and biomedical research. Neurosci Biobehav Rev. 2014;40: 1–5.

43.   Beery AK, Zucker I. Sex bias in neuroscience and biomedical research. Neurosci

Biobehav Rev. 2011;35: 565–572.

44.   Mohammad F, Ho J, Woo JH, Lim CL, Poon DJJ, Lamba B, et al. Concordance

and incongruence in preclinical anxiety models: Systematic review and meta-analyses.

Neurosci Biobehav Rev. 2016;68: 504–529.

45.   Wizemann TM. Sex-specific reporting of scientific research. Washington:

National Academies Press. 2012.

46.   Clayton JA, Collins FS. Policy: NIH to balance sex in cell and animal studies.

Nature. 2014;509: 282–3.

47.   Ioannidis JP, Trikalinos TA. An exploratory test for an excess of significant

findings. Clin Trials. 2007;4: 245–253.

48.   Ridley J, Kolm N, Freckelton RP, Gage MJG. An unexpected influence of

widely used significance thresholds on the distribution of reported P-values. J Evol

Biol. 2007;20: 1082–1089.

49.   Simonsohn U, Nelson LD, Simmons JP. *p* -Curve and Effect Size. Perspect

Psychol Sci. 2014;9: 666–681.

50.   de Winter JC, Dodou D. A surge of *p*-values between 0.041 and 0.049 in recent

decades (but negative results are increasing rapidly too). PeerJ. 2015;3: e733.

51.   Wagenmakers EJ. A practical solution to the pervasive problems of p values.

Psychon Bull Rev. 2007;14: 779–804.